A systems based framework to computationally predict putative transcription factors and signaling pathways regulating glycan biosynthesis

Theodore Groth[1] and Sriram Neelamegham[1,2,3,*]

[1]Chemical and Biological Engineering, [2]Biomedical Engineering and [3]Medicine
University at Buffalo, State University of New York, Buffalo, NY 14260, USA

Running title: Systems Glycobiology

* Correspondence: Sriram Neelamegham, 906 Furnas Hall, Buffalo, NY, 14260,
neel@buffalo.edu, Ph: 716-645-1200; Fax: 716-645-3822

Abstract

Glycosylation is a common post-translational modification, and glycan biosynthesis is

regulated by a set of 'glycogenes'. The role of transcription factors (TFs) in regulating the

glycogenes and related glycosylation pathways is largely unknown. This manuscript presents a

multi-omics data-mining framework to computationally predict putative, tissue-specific TF

regulators of glycosylation. It combines existing ChIP-Seq (Chromatin Immunoprecipitation

Sequencing) and RNA-Seq data to suggest 22,519 potentially significant TF-glycogene

relationships. This includes interactions involving 524 unique TFs and 341 glycogenes that span

29 TCGA (The Cancer Genome Atlas) cancer types. Here, TF-glycogene interactions appeared

in clusters or 'communities', suggesting that changes in single TF expression during both health

and disease may affect multiple carbohydrate structures. Upon applying the Fisher's exact test

along with glycogene pathway classification, we identify TFs that may specifically regulate the

biosynthesis of individual glycan types. Integration with knowledge from the Reactome database

provided an avenue to relate cell-signaling pathways to TFs and cellular glycosylation state.

Whereas analysis results are presented for all 29 cancer types, specific focus is placed on

human luminal and basal breast cancer disease progression. Overall, the computational

predictions in this manuscript present a starting point for systems-wide validation of TF-

glycogene relationships.

Keywords: Glycoinformatics, transcription factor, glycosylation, ChIP-Seq, TCGA

Introduction

The glycan signatures of cells and tissue is controlled by the expression pattern of 300-350 glycosylating genes that are together termed 'glycogenes' [1,2]. These glycogenes include the glycosyltransferases, glycosidases, sulfotransferases, transporters etc. The expression of these glycogenes is in turn driven by the action of a class of proteins called transcription factors (TFs). These TFs regulate gene expression by binding proximal to the promoter regions of genes, facilitating the binding of RNA polymerases. They may homotropically or heterotropically associate with additional TFs in order to directly or indirectly control messenger RNA (mRNA) expression. Among the TFs, some 'pioneer factors' can pervasively regulate gene regulatory circuits, and access chromatin despite it being in a condensed state [3]. These TFs act as 'master regulators', promoting the expression of several genes across many signaling pathways, such as differentiation, apoptosis and cell proliferation. The precise targets of the TFs is controlled by their tissue-specific expression, DNA binding domains and nucleosome interaction sequences [3]. Additional factors regulating transcriptional activity include: i. cofactors and small molecules that enable TF-DNA recognition and RNA polymerase recruitment [3]; ii. chomatin modifications, such as acetylation, methylation and phosphorylation, which alter TF access; and iii. methylation of CpG islands in promoter regions which inhibit gene expression [4,5].

There are currently several isolated studies of TF-glycogene interactions, but to our best knowledge a systematic "systems level analysis" is absent. Many of these previous studies are based on discrete glycogene promoter region analysis and reporter assays. These studies have established some notable TF-glycogene relationships, though they are limited to distinct cell types. Examples include the regulation of MGAT5 by ETS2 in NIH3T3 fibroblasts [6], control of the α2-6 sialyltransferases ST6Gal-I/II by Hypoxic Nuclear Factor 1-α (HNF1- α) in HepG2 cells [7], c-JUN-B3GNT8 regulatory relationships in gastric carcinoma cell lines [8], and SP1-B4GALT1 relations in lung cancer A549 cells [9]. A recent study also used computational

predictions and wet-lab experiments to determine that ZNF263 is a potential heparin sulfate

master regulator [10]. This TF regulates two sulfotransferases, HS3ST1 and HS3ST3A1. The

above approaches have limitations in that: i. they do not consider the cellular epigenetic state

that could impact TF binding; and ii. proximal regulators are studied but enhancers present

several kilobases away from the Transcription State Site (TSS) away are neglected.

In the current manuscript, we propose that more global and higher-throughput TF-

glycogene relationships under biologically relevant conditions may be discovered using multi-

omics data-mining. To this end, we sought to utilize multi-omics experimental datasets and

curated pathway databases (DB) to relate cell-specific signaling processes to TFs, TFs to

glycogenes, and glycogenes to glycosylation pathways (**Fig. 1A**). These connections were

made using data available from Cistrome Cancer DB [11], Reactome DB [12] and by the manual

curation of various human glycogenes into pathways at GlycoEnzDB

(www.virtualglycome.org/GlycoEnzDB, **Fig. 1B**). Here, the Cistrome Cancer DB uses TF-gene

binding data from previously published ChIP-Seq studies for various cell systems and cancer

tissue RNA-seq data from The Cancer Genome Atlas (TCGA) [13]□. It curates putative TF-

gene relationships for 29 TCGA cancer types provided they satisfy three inclusive criteria: i. TFs

should be expressed at high levels in a given tissue; ii. Changes in TF gene expression should

correlate with RNA changes in target genes; iii. Chromatin Immunoprecipitation sequencing

(ChIP-Seq) data must support the TF-gene binding proximal to the transcription start site (TSS).

Next, knowledge curated in the Reactome DB [12]□ is used to establish links between TFs and

signaling pathways. In the final step, manually curated glycogene classifications were utilized to

determine TFs that disproportionately regulate individual glycosylation pathways. The findings

from this study

Figure 1. A systems glycobiology framework to link multi-OMICs data: **a.** Cell signaling proceeds to trigger transcription factor (TF) activity. The binding of TFs to sites proximal to the transcriptional start site triggers glycogene expression. A complex set of reaction pathways then results in the synthesis of various carbohydrate types, many of which are either secreted or expressed on the cell surface. **b.** Data available at various resources can establish the link between cell signaling and glycan biosynthesis. The Reactome DB contains vast cell signaling knowledge. Chip-Seq and RNA-Seq data available at the Cistrome Cancer DB describe the link between the TFs and glycogenes. Pathway curation at the GlycoEnzDB establishes the link between glycogenes and glycan structures. Cell illustration created using BioRender.com.

Figure 2. Analysis workflow: ChIP-seq provides evidence of TF binding to promoter regions with regulatory potential ($0 \leq RP \leq 1$) quantifying the likelihood that this is functionally important. RNA-Seq quantifies Spearman's correlation (ρ) between TF and gene expression. Filtering these data establishes potential TF-glycogene interactions in specific cancer types. TFs disproportionately regulating specific glycosylation pathways were identified using the above TF-glycogene relationships, and biochemical knowledge available at GlycoEnzDB (green region). ReactomeDB analysis helped establish cell signaling-TF-glycosylation pathway connectivity that are visualized using Alluvial plots. Independently, cytoscape maps enabled visualization of TF-glycogene relationships in different cancers (orange region).

1 Clusters in the resulting interactomes were related to pathway maps and signaling processes, thus
2 developing TF-community signaling pathway relationships.
3
4
Results

TF-Glycogene interaction map and relation to cell signaling pathways: The manuscript

follows a workflow shown in **Figure 2**. It infers TF-glycosylation pathway relationships, as well

as TF-glycogene communities by using publicly available ChIP-seq data and RNA-Seq results.

These data were obtained from the Cistrome Cancer DB [14], which curated TF-gene

relationships by integrating ChIP-seq data from Cistrome DB and RNA-seq data from The

Cancer Genome Atlas (TCGA). The Cistrome Cancer DB uses three filtering criteria to

determine putative TF-gene relationships: 1. The TF should be active in a cancer type, i.e. the

RPKM (Reads Per Kilobase Million) value in a cancer type must be greater than the TF's

median RPKM expression across all 29 different cancer types. 2. The RNA expression of TF

and target gene should be correlated. To determine this, Cistrome first compares the selected

TF-gene correlation with a null distribution computed by randomly selecting 1 million TF-gene

pairs. Linear regression and statistical analysis are then performed on the top 5-percentile hits

(positive and negative coefficients) to establish TF-gene correlations. This analysis accounts for

target gene copy number, tumor purity, and promoter methylation extent. 3. TF-gene

relationships must be supported by Chromatin Immunoprecipitation sequencing (ChIP-Seq)

evidence. Here, a non-linear weighted sum called "regulatory potential (RP)" quantifies the

strength of TF-gene interactions based on the proximity of TF binding site to the gene

transcriptional start site, and also the number of TF-gene binding interactions based

experimentally detected ChIP-peaks [15,16].

In the current manuscript, we passed the TF-gene relationships established in Cistrome

Cancer DB, to identify TFs potentially interacting with 341 glycogenes (**Supplemental Table**

S1). The two metrics for this select were $RP \geq 0.5$ and TF-glycogene expression correlation

coefficient $\rho \geq 0.4$. Such analysis was performed for 29 cancer types listed in **Supplemental**

Table S2. Based on our selected thresholding, the analysis revealed 22,519 potential TF-

glycogene interactions. The above data were used for two types of analysis described below.

First, the Fisher's exact test was used to infer TF-glycogene interactions that may

regulate individual glycosylation pathways. This analysis was based on pathway classifications

from GlycoEnzDB (**Supplemental Table S3**) that grouped 208 glycogenes into 20 glycosylation

pathways/groups. TFs having disproportionately larger number of relationships with individual

glycosylation pathways were determined, with respect to all TF-glycogene relationships.

Reactome DB was then used to associate these TFs to potential signaling pathways. This

method, thus, resulted in a relationship between cell-signaling, TF activity regulation and glycan

structure changes (**Supplemental Table S4, S5**). The data are presented as Alluvial plots for

the 29 cancer types (**Supplemental Fig. S1**). Here, the TFs were linked to glycosylation

pathways by colored bands if they were found to regulate a disproportionately high fraction of

glycogenes belonging to that pathway. Likewise, biological pathways were linked with TFs if

that TF was found to be enriched in the biological pathway.   Reading these alluvial plots left to

right, one can deduce which biological pathways may be potentially involved in regulating TFs,

and how these TFs could regulate glycosylation.

Second, we visualized TF-glycogene interactions using Cytoscape maps for each of the

cancers individually (**Supplemental File S2**). This analysis revealed clusters of TFs regulating

common sets of glycogenes. We sought to understand which glycosylation pathways were

overrepresented in each of these cluster using the Fisher's exact test. This analysis revealed

406 glycopathway enrichments in the TF-glycogene communities across the 29 cancer types

(**Supplemental Table S6**). Next, we determined, using the Reactome DB overrepresentation

API, if the TFs identified in these clusters could be related to specific cell signaling pathways.

Here we noted 901 pathway enrichments across the different cancer types (**Supplemental**

Table S7). Common TFs that we observed across all TF-glycogene communities include the

TCF and LEF families, FOXO and FOXP, RUNX family, and IRF family TFs, which were found to

regulate diverse glycosylation pathways, such as sialylation pathways, complex N-linked glycan

synthesis, chondroitin and dermatan sulfate synthesis.

Overall, the above analysis revealed the existence of communities of TF-glycogene

relationships, that could be linked to both cell signaling processes and specific glycosylation

pathways. Future, quantitative computational modeling and mass spectrometry studies may

establish the biological significance of these predictions.

TF-pathway relationships in breast cancer: As a usage example of the above framework, we

provide more detailed investigation using breast cancer as an example. This disorder appears in

5 unique molecular subtypes based on the PAM50 classification [17]. These include: i. normal-

like, ii-iii. luminal A and luminal B which overexpress estrogen receptor ESR1, iv. Her2+ tumors

that overexpress the epidermal growth factor receptor (ERBB), and v. basal (triple negative) that

express neither ESR1 nor ERBB. Each of these subtypes has unique signaling mechanisms

that may contribute to different glycan signatures.

In our analysis, TF-glycogene relationships for breast cancer derived by filtering

Cistrome Cancer were enriched for the glycosylation pathways. **Fig. 3** summarizes these cancer

related TF-glycosylation pathway relationships for luminal (type A and B together) and basal

breast cancers. The analysis suggests that TF transformations accompanying cancer

progression may impact all four major classes of glycans: O- and N-glycans found on

glycoproteins, glycosaminoglycans and glycolipids. Thus, multiple glycan changes may

accompany oncological transformation.

**Figure 3. Summary of TFs enriched to glycosylation pathways for luminal and basal breast
cancer:** The TFs found to be enriched to glycosylation pathways and the glycogenes they regulate are
shown in pink for luminal and orange for basal breast cancer. Note that some of the TFs shown above do
not appear in the alluvial plots in figures 4 and 5 because they were not enriched to a signaling pathway
in Reactome. The glycans synthesized by the enriched glycogenes are shown in SNFG format [18]. All
figures were generated using DrawGlycan-SNFG [19].

TF-glycogene communities in luminal and basal breast cancer (cytoscape plots):

Cytoscape plots were generated for luminal breast cancer (**Fig. 4a**). Here, using the bipartite

graph community detection methods [20], we inferred three large communities of TF-glycogene

interactions. The largest community detected in this analysis had TFs enriched for RUNX3

signaling, IL-21 signaling, MECP2, and PTEN regulation. Over-representation glycosylation

pathway analysis performed on the TFs in this community suggests that these TFs may regulate

pathways related to sialylation, hyaluronan synthesis, and chondroitin and dermatan sulfate

elongation. Here, STAT1, 4, and 5 proteins were enriched in the IL-21 signaling pathway.

Luminal breast cancers are known to express STAT1, 3 and STATs 2 and 4 are known to be

expressed in luminal breast cancer cell lines. STAT5 is known to be constitutively active in

luminal breast cancer and confers anti-apoptotic characteristics to cells [21]. The other two

communities detected consisted primarily of chromatin-modifying enzymes. Complex N-linked

glycan synthesis and the dolichol pathway were significantly enriched in the second community.

  In the third community, O-linked mannose and LacdiNAc synthesis were disproportionately

  regulated. Overall, the pathway maps suggest that chromatin remodeling enzymes could

  potentially play roles in regulating glycan synthesis in luminal breast cancer.

Figure 4. Luminal breast cancer signaling pathway enrichment and glycogene connections: a. *TF-to-glycogene communities in luminal breast cancer.* Three large TF-to-glycogene communities were discovered in the luminal breast subnetwork. Community 1 was enriched for pathways involving RUNX3, RUNX1, IL-21, and PTEN. Communities 2 and 3 consist primarily of chromatin modifying enzymes. **b.** *Signaling pathway enrichment analysis for luminal breast cancer.* Connections between signaling pathways and transcription factors found to be statistically significant for luminal breast cancer. Some pathways enriched to TFs were condensed to conserve space. More TF-to-glycogene relationships exist in luminal breast cancer and these can be viewed in the cytoscape figures (**Supplemental Figure S2**).

Like luminal, basal breast cancer TF-glycogene relationships were also clustered into

three communities. Here, the first community was enriched for chromatin modifying enzymes,

with complex N-linked glycan synthesis being the primary glycosylation pathway being affected

(**Fig. 5a**). The second community was enriched for interferon α/β/γ signaling pathways, with

interferon regulatory factor (IRF) TFs being enriched. In this regard, the TFs IRF-1 and IRF-5

have been shown to act as tumor suppressors in breast cancer [22,23]. Their loss-of-function in

1 breast cancer could potentially downregulate O-linked fucosylation. The third community did not

2 exhibit any specific TF pathway enrichments.

3
4
Figure 5. Basal breast cancer signaling pathway enrichments and glycogene connections: a. *TF-*
to-glycogene communities in basal breast cancer: Three large TF-to-glycogene communities were
discovered in the basal breast subnetwork. Community 1 has TFs enriched to chromatin modifying
enzymes, and community 2 has TFs enriched to interferon α/β/γ signaling. Community 3 did not have
any signaling pathways enriched. **b.** *Signaling pathway enrichment analysis for basal breast cancer*:
Connections between signaling pathways and TFs found to be statistically significant for basal breast
cancer. TFs displayed have been enriched to the displayed glycosylation pathways using the Fisher's
exact test.
13

Linking cell signaling to TF and glycogenes for luminal breast cancer (alluvial plot): The

links between biological signaling pathways, TFs, and glycosylation pathways are shown in

alluvial plots for luminal (**Fig. 4b**) and basal breast cancers (**Fig. 5b**), with additional plots

provided for additional cancer types in Supplemental Material. In the case of luminal breast

cancer:

CREB3L4 and PRDM1 disproportionately affect the Type I & II LacNAc pathway in luminal

breast cancer: Our analysis suggests that CREB3L4 (enrichment p-value = 0.036) and PRDM1

(enrichment p-value = 0.039) may regulate the Type 1 and 2 LacNAc pathways. CREB3L4 is

known to primarily be expressed in the prostate and some breast cancer cell lines, and has

been linked to diverse roles involving chromatin organization in spermiogenesis, adipocyte

regulation, and dysregulation in prostate cancer[24,25]. It has been found to be upregulated in

breast cancer with respect to normal. PRDM1, also known as Blimp-1, is a transcriptional

repressor, and its upregulation in cancer is known to dysregulate other proteins [26].The

increase Poly LacNAc structures have been shown to play roles in cancer metastasis [27].

CREB3L4 was found to regulate B4GALT3 glycogene (ρ=0.56, RP=0.94) which adds galactose

in a β1-4 linkage, and PRDM1 was found to regulate B3GNT5 which is critical for lacto-

/neolactoseries glycolipids (ρ=0.60, R.P.=0.84).

MEF2C disproportionately regulates Glycosaminoglycan synthesis pathways: MEF2C was

found to regulate several genes in the chondroitin and dermatan sulfate synthesis pathways

(p=0.008). This TF plays roles in development, particularly with the development of neurons

and hematopoetic cell differentiation towards myeloid lineages. It is known that MEF2C is

directly impacted by TGF-β signaling, thus increasing the metastatic potential of cancer [28].

MEF2C was found to be inhibited by MECP2 based on Reactome pathway enrichment. Since

the glycosaminoglycan elongation pathways positively correlate to MEFC2 expression, and

MEFC2 is amplified in cancer, it is possible that MECP2 may not sufficiently expressed to

repress MEFC2 in call cancer cells. MEF2C was found to regulate CSGALNACT1 (ρ=0.66,

RP=0.71), CHST3 (ρ=0.50, RP=0.74), CHST11 (ρ=0.47, RP=0.84), DSEL (ρ=0.40, RP=0.81),

and UST (ρ=0.42, RP=0.95). Here, CSGALNACT1 is responsible for the addition of GalNAc to

glucuronic acid to increase chondroitin polymer length, CHST3, CHST11, and UST are involved

in the sulfation of GalNAc and iduronic acid, and DSEL is the epimerase which converts

glucuronic acid to iduronic acid in CS/DS chains.

MECP2 disproportionately regulated heparan sulfate chain elongation: The MECP2 (enrichment

p-value=0.037) was found to positively regulate heparan sulfate elongation. MECP2 regulates

gene expression by binding to methylated promoters, and then by recruiting chromatin

remodeling proteins to condense DNA and repress gene expression [29,30]. MECP2 was

found to regulate sulfotransferase NDST1 (ρ=0.41, RP=0.67).

Linking cell signaling to TF and glycogenes for basal breast cancer (alluvial plot): Fewer

transcription factors were found to be enriched to signaling pathways in basal breast cancer

compared to luminal cancer (**Fig. 4b**). Despite this, there are many other TF-glycosylation

pathway enrichments for basal breast cancer available for analysis in Supplementary Material.

The roles of the enriched TFs and their relation to glycogenes and cancer is elaborated below.

RUNX3 and Fucosylation: The terminal fucosyltransferase FUT7 (ρ=0.49, RP=0.89) was found

to be positively regulated by the RUNX3 TF (enrichment p-value = 0.033). The RUNX family of

transcription factors (including RUNX1-3), are involved in several developmental processes,

including hematopoiesis, immune cell activation, and skeletal development. It was discovered

that RUNX3 acts as a tumor suppressor gene in breast cancer. Upon cancer development, the

RUNX3 promoter is hypermethylated, leading to reduced TF activity and loss of tumor

suppression activity [31]. Our data suggest that this may be associated with a reduction of FUT7

activity thus impacting the expression of the sialyl Lewis-X antigens in basal tumors. Sialyl

Lewis-X is considered to be an important regulator of cancer metastasis as it binds the selectins

on various vascular and blood cell types.

Regulation of GalNAc-type O-linked glycans by SMAD2: SMAD2 was found to significantly

affect core 1 & 2 O-linked glycan structures (enrichment p-value=0.035). SMAD proteins are

activated by TGF-β signaling and bind to DNA to act as cofactors to recruit TFs. SMAD2 has

been shown to act as a tumor metastasis suppressor in cell lines [32,33]. This TF was found to

regulate GALNT1 (ρ=0.54, RP=1.00), which adds GalNAc to serine or threonine residues to

being core 1 and 2 O-linked glycan synthesis. Thus, SMAD2 may play a key role in regulating

Tn-antigen expression in proteins like MUC-1 that are associated with breast cancer

progression.

Discussion

In the current analysis, we sought to identify strategies to enhance systems glycobiology

knowledge by leveraging existing, publicly-available, high-throughput ChIP-Seq and RNA-Seq

data. This analysis suggests putative TFs regulating glycogenes and glycosylation processes

across 29 different cancer types. Approximately three glycogenes were regulated by a given TF

based on our filtering criteria, with this number ranging from 1-10. These findings are tissue-

specific, as TF and glycogene expression vary widely among the different cell types. The

analysis also suggests putative TF-glycogene interactions that disproportionately impact specific

glycosylation pathways. Knowing which TF regulates which glycogene and pathway in a

context-dependent manner can provide insight as to how signaling pathways contribute to

altered glycan structures in diseases such as diabetes and cancer. Thus, this work represents a

rich starting point for wet lab validation and glycoinformatics database construction.

Visualizing TF-glycogene interaction networks revealed communities of glycogenes in

each cancer type. The presence of chromatin-modifying enzymes in large regulatory

communities in both luminal and basal breast cancer suggests a role of epigenetics in

glycogene regulation. To date, a systems-level investigation evaluating the epigenetic states of

cell systems on the resulting glycome has not been performed. Our results suggest that

complex N-linked branching and glycosylation may be sensitive to these processes. The

signaling pathways enriched in the largest community in luminal breast cancer were reflected in

our pathway enrichment findings. RUNX3, interleukin signaling, and the involvement of MECP2

regulation were all found to disproportionately regulate sialic acid and GAG synthesis pathways.

Several of the TFs enriched to glycosylation pathways were either regulated by or involved in TGF-β signaling and Wnt β-catenin signaling. These TFs primarily affected glycosaminoglycan synthesis pathways, sialylation and Type-2 LacNAc synthesis. Cell cycle and metabolic regulatory TFs were shown to regulate some glycogenes involved in the dolichol pathway. The crosstalk between cell cycle and glycosylation is not well explored, and may potentially be important for understanding N-linked glycosylation flux in cancer. Some TFs were found to interact with methyl CpG binding TFs when regulating glycosaminoglycan proteins, implicating methylation as a possible modulator of glycosylation in cancer.

The framework described in this manuscript represents a starting point for experimentally investigating TFs regulating glycosylation, however much more is needed to validate and establish these relationships. First, selected values of RP and ρ used to filter TF-glycogene relationships from the Cistrome Cancer database were selected to initiate a query into the available ChIP-seq data. Further studies are needed in order to determine how the selected thresholds affect the discovered relationships. Second, the glycogenes in individual pathways in this manuscript were classified using current knowledge of glycobiology. Different classification methods may result in slightly different TF-glycogene relationships [34]. Third, while Cistrome Cancer DB systematically filters TF-gene relationships based on ChIP-seq and RNA-seq evidence, the database has some biases. In one aspect, only TFs that were considered to be sufficiently expressed were considered in this analysis. Lower expressed TFs that may also be functional, are excluded from the analysis. Additionally, while RNA-Seq relationships in Cistrome Cancer are selected based on specific tissue type, supporting ChIP-seq evidence is not cell type specific. The tissue-dependent nature of TF-glycogene binding thus requires further validation through either surveying other datasets or performing experimental validation. Many databases curating ChIP-seq and other omic information could aid such TF-glycogene relationship validation. Some examples include: i. ChIP-Seq databases, such as the Gene Transcription Regulatory Database (GTRD) [35] □ that have analyzed vast

publicly available ChIP-Seq data to systematically cataloged TF-gene relationships across

several organisms and cellular contexts using multiple ChIP-seq peak calling algorithms, ii. the

Regulatory Circuits database [36]□ that quantifies the activity of promoter and enhancer

regions through cap analysis of gene expression (CAGE) and expression Quantitative Trait Loci

(eQTL), respectively, and iii. integration of TF binding motifs, protein-protein interactions, and

co-expression networks, as was done using data from GTEx and a method called PANDAS [37].

Such cross-platform analysis may reduce the number of candidate TF-glycogene interactions

presented in this manuscript. This would also aid more focused wet-lab validation using either

CRISRP-Cas9 knockouts and single cell RNA-Seq, or using structural methods that involve

glycomics and glycoproteomics based mass spectrometry.

11
12
Conclusion

A majority of current studies in the Glycoscience field use experimental data and

curations related to glycans only. Fewer investigations examine the links between the glycans,

glycogenes and glycosylation pathways, and other non-glyco datasets. We set out to establish

these relationships using bioinformatics and data-mining. Using this, we describe putative

regulatory relationships between TFs and glycogenes across 29 cancer types. Some TFs

appear to regulate glycogenes in communities, indicating potential cross-talk across pathways

in regulating glycosylation. The communities varied with cancer type, even in a single tissue,

suggesting that these TF-glycogene interactions are dynamic in nature. Groups of TFs enriched

to glycosylation pathways were also associated with signaling pathways, thus establishing a

framework to discover how cell signaling processes may regulate cellular glycosylation by

modifying TF activity. Overall, the putative TF-glycosylation pathway enrichments found here

represent the starting point for wet-lab validation. Such studies could enhance our fundamental

understanding of glycosylation pathway regulation, and lead to novel ways to control the

glycogenes and glycan structures during disease.

EXPERIMENTAL METHODS

Glycogene-pathway classification: A list of 208 unique glycogenes involved in 20 different

glycosylation pathways were used in this work (**Supplemental Table S3**). These data are

collated from GlycoEnzDB (virtualglycome.org/GlycoEnzDB), with original data coming from

various sources in literature [38,39]. The following is a summary of the pathways studied and

the enzymes involved:

1. Glycolipid core: The enzymes in this group are involved in the biosynthesis of the glucosyl-

ceramide (GlcCer) and galactosyl (GalCer)-ceramide lipid core. Here, the GlcCer core is formed

by the UDP-glucose:ceramide glucosyltransferase (UGCG) which transfers the first glucose.

Following this, lactosylceramide is formed by the action of the β1,4GalT activity of B4GalT5 (and

possibly also B4GalT3, 4 and 6). The GalCer core is typically structurally small and is made by

UDP-Gal:ceramide galactosyltransferase (UGT8). These structures can be further sulfated by

GAL3ST1 or sialylated by ST3GAL5.

2. P1-Pk Blood Group: The Pk, P1 and P antigens are synthesized on lactosyl-ceramide

glycolipid core. The activity of α1-4GalT (A4GALT) on this core results in the Pk antigen,

followed by β1-3GalNAcT (B3GALNT1) to form the P antigen. The P1 antigen, on the other

hand, is formed by the sequential action of β1-3GlcNAcT (B3GNT5), β1-4GalT (B4GALT1-6)

and α1-4GalT (A4GALT) on the glycolipid core.

3. Gangliosides: This pathway encompasses all glycogenes responsible for synthesizing a/b/c

gangliosides. UGCG is included to consider the addition of glucose to ceramide. ST3GAL5,

and ST8SIA enzymes are added to take the core ganglioside structures to the a,b and c levels.

B4GALTs and B4GALNT1 are included to account for ganglioside elongation. Decoration of the

gangliosides with sialic acid occurs using ST6GALNAC3-6 and also ST8SIA1/3/5.

4. Dolichol Pathway: This results in the formation of the dolichol-linked 14-monosaccharide

precursor oligosaccharide. This glycan is co-translationally transferred en bloc onto Asn-X-

Ser/Thr sites of the newly synthesized protein as it enters the endoplasmic reticulum. The

enzymes involved is such synthesis include the ALG (Asparagine-linked N-glycosylation)

enzymes, and additional proteins (part of OSTA and OSTB) involved in the transfer of the glycan

to the nascent protein.

5. Complex N-glycans: This pathway includes glycogenes responsible for processing the N-

linked precursor structure emerging from the dolichol pathway into complex structures.

Enzymes involved include mannosidases, glucosidases, some enzymes facilitating protein

folding and also enzymes that direct acid hydolases to the lysosome.

6. N-glycan branching: These glycogenes are responsible for the addition of GlcNAc to

processed N-linked glycan structures. These include all the MGAT enzymes.

7. GalNAc-type O-glycans: O-linked glycans are attached to serine or threonine (Ser/Thr) on

peptides, where GalNAc is the root carbohydrate. This is mediated by a family of about 20

Golgi-resident polypeptide N-acetylgalactosaminyltransferases (ppGalNAcTs or GALNTs). Core

1 structures result from the attachment of β1-3 linked galactose to the core GalNAc using

C1GALT1 and its chaperone C1GALT1C1. Core 2 structures then form upon addition of β1-6

linked GlcNAc by GCNT1. Modifications of core-3 and core-4 glycans can occur during disease

and thus this classification includes core-3 forming B3GNT6 and core-4 forming GCNT3. Other

O-glycan core-types are rare in nature.

8. Chondroitin Sulfate & Heparan Sulfate Initiation: Chondroitin and heparan sulfate

glycosaminoglycans all have a common core carbohydrate sequence attaching them to their

proteins. These are constructed by the activity of specific Xylotransferases (XYLT1, XYLT2),

galactosyltransferses B4GALT7 and B3GALT6 that sequentially add two galactose residues to

Xylose, and the Glucuronyltransferase B3GAT3 then adds glucuronic acid to the terminal

galactose. Also involved in the formation of this core is FAM20B, a kinase that 2-O-

phosphorylates Xylose. At this point, the addition of GalNAc to GlcA by CSGALNACT1 & 2

results in the initiation of chondroitin sulfates chains. The attachment of GlcNAc by EXTL3 to the

same GlcA results in heparan sulfates.

9. Chondroitin/dermatan sulfate extension: Chondroitin sulfates and dermatan sulfates are

extended via the addition of GalNAc-GlcA repeat units. This is catalyzed by CSGALNACT1

which is better suited for the initial GalNAc attachment followed by CSGALNACT2 which is

preferred for synthesizing disaccharide repeats. CHSY1, CHSY3, CHPF and CHPF2, all exhibit

dual β1,3GlcAT and β1,4GlcAT activity. Additional enzymes mediate sulfation. Epimerization of

glucuronic acid to iduronic acid by DSE and DSEL results in the conversion of chondroitin

sulfates to dermatan sulfates.

10. Heparan sulfate extension: EXT1 and EXT2 both have GlcUA and GlcNAc transferase

activities and are together responsible for HS chain polymerization. EXTL1-3 are additional

enzymes with GlcNAc transferase activity that facilitate heparin sulfate biosynthesis. Additional

enzymes that are critical for heparin sulfate function include the HS2/3/6ST sulfotransferases,

the GlcA epimerase GLCE and additional enzymes mediating N-sulfation (NDSTs).

11. Hyaluronan Synthesis: This pathway consists of the three hyaluronan synthases HAS1-3.

12. GPI Anchor Extension: This pathway includes glycogenes responsible for the synthesis of

glycosphosphatidylinositol (GPI) anchored proteins in the ER. This involves synthesis of a

glycan-lipid precursor that is en bloc transferred to proteins.

13. O-Mannose: This is initiated by the addition of mannose to Ser/Thr using POMT1 or

POMT2. β1-2 or β1-4 GlcNAc linkages can then be made using POMGNT1 or POMGNT2 to

yield M1 or M3 O-linked mannose structures, respectively. MGAT5B can facilitate β1-4 GlcNAc

linkage onto the M1 structure to yield the M2 core. Additional carbohydrates typically found on

complex N-linked glycan antennae can then attached. In particular, such extensions may be

initiated by members of the B4GALT family or B3GALNT2. Specific variants are noted on α-

dystroglycans.

14. O-linked Fucose: This pathway includes POFUT1, the enzyme responsible for the addition

of fucose to Ser/Thr residues. MFNG, LFNG and RFNG which can attach β3GlcNAc to this

fucose.

15. Type 1 & 2 LacNAc: These enzymes help construct either Galβ1,3GlcNAc (Type 1) or

Galβ1,4GlcNAc (Type 2) lactosamine chains on antennae of N-linked glycan, O-linked glycans

and glycolipids. Also included are GCNT1-3 that can facilitate formation of I-branches on N-

glycans.

16. Sialylation: This group encompasses all kinds of sialyltransferases: ST6GAL, ST3GAL,

ST8SIA, and ST6GALNACs. Enrichments to this pathway capture overall increase in sialylation

regardless of context.

17. Fucosylation: These include α1-2 (FUT1, 2) and α1-3 (FUT3, 4, 5, 6, 7, 9)

fucosyltransferases that can act on N-glycans, O-glycans and glycolipids.

18. ABO blood Group Synthesis: These are enzymes involved in the biosynthesis of ABO

antigens

19. LacDiNAc: Glycogenes involved in the synthesis of LacDiNac structures.

20. Sulfated glycan epitopes: This includes the enzymes attaching sulfate to different types of

carbohydrates

Establishing TF–glycogene relationships in Cistrome Cancer: Regulatory potential and

gene correlation data were downloaded from the Cistrome Cancer database in tab-delimited

form (http://cistrome.org/CistromeCancer/CancerTarget/)[14].   TF-gene relationships were

filtered for the 341 glycogenes in this manuscript (**Supplemental Table S1**). In total, the full

dataset contained 41,771 TF-to-glycogene relationships, including relational data for 568 unique

TFs found in the 29 cancer systems across all the glycogenes. Positive regulatory relationships

between TFs and glycogenes were selected based on RP ≥ 0.5 and ρ ≥ 0.4 (**Fig 2**). This filtering

resulted in 20,617 TF-glycogene relationships including 524 unique TFs across 29 cancer types.

Cytoscape was used to visualize TF-glycogene regulatory relationships [40]. To achieve

this, all TF-glycogene relationship data were loaded into cytoscape as a network. These data

were filtered based on RP and _ρ_ thresholds defined previously. A binding potential (BP) score

was computed by taking the product of RP and ρ for each TF-glycogene relationship. TF-

glycogene relationships for each cancer type were separated into sub-networks. The prefuse

force directed layout algorithm in cytoscape was used to arrange nodes in each cancer sub-

network. The closeness of nodes to one another is weighted by 1-BP. Thus, nodes with high

BPs will be placed closer together, whereas smaller BPs will be placed further away. Since

there are two classes of nodes (TFs and glycogenes), we treated TF-glycogene networks as

bipartite, and applied the corresponding procedure for community detection [20]. Firstly, the

bipartite TF-glycogene graphs are projected into two different unipartite graphs, where TFs and

glycogenes are placed into separate graphs. The edge weights connecting TFs is computed as

the number of shared glycogenes they regulate. The TF unipartite graph was then subjected to

a greedy modularity optimization-based approach implemented in the R package igraph [41].

TF-glycogene interactions in each community were subjected to overrepresentation analyses to

identify enriched signaling and glycosylation pathways.

Relating TF-glycogene interactions to glycosylation and signaling pathways: A Fisher's

Exact Test was applied to determine if a particular TF disproportionately regulates one of the 20

glycosylation pathways described in **Supplemental Table S3**. Input data to the test consisted

of all TF-gene interactions that passed the RP and ρ thresholds for the cancer type being

analyzed. TFs were considered to be disproportionately regulating a glycosylation pathway if

the Fisher's exact test resulted in a p-value≤ 0.05 .

TFs enriched to glycosylation pathways were associated with putative regulatory

pathways using Reactome DB over-representation analysis API, which uses a Fisher's exact

test, to associate the TFs with signaling pathways[12]. Signaling pathway enrichments with

adjusted p (FDR)<0.1 were kept. A high p-value cutoff was chosen to allow users to gain a

high-level perspective as to what potential pathways may be regulating enriched TFs. The

connection between cell signaling pathways and TFs, and that between the TFs and

glycosylation pathways were visualized using alluvial plots generated using the R package

ggalluvial. Only signaling pathways with < 30 members are presented for brevity. A

comprehensive listing of enriched signaling pathways is available in Supplemental Table S5.

SUPPORTING INFORMATION
Supplementary Table S1:
File Name: TableS1_Glycogenes.xlsx
File Format: XLXS
Title: List of 341 glycogenes used for cystoscape maps
10
11 Supplementary Table S2:
12 File Name: TableS2_CancerTypes.xlsx
13 File Format: XLSX
14 Title: Cancer type list
15
16 Supplementary Table S3:
17 File Name: TableS1_Glycogene_Pathway_Lists.xlsx
18 File Format: XLSX
19 Title: Glycogene pathway lists
20
21 Supplementary Table S4:
22 File Name: TableS4_Fishers_exact_test_summary.xlsx
23 File Format: XLXS
24 Title: Fisher's exact test to infer TF-glycosylation pathway relation (p<0.05 data are highlighted)
25
26 Supplementary Table S5:
27 File Name: TableS5_Reactome_Enrich_Pathways.xlsx
28 File Format: XLXS
29 Title: Reactome pathway enrichments for all TFs
30
31 Supplementary Table S6:
32 File Name: TableS6_GlycoPathway_Network_Enrichment.xlsc
33 File Format: XLXS
34 Title: Network Cluster Enriched Glycosylation Pathways
35
36 File Name: TableS7_Signaling_Network_Enrichment. xls
37 File Format: XLXS
38 Title: Network Cluster Enriched Signaling Pathways from Reactome
39
40 Supplementary File S1:
41 File Name : FileS1_supplemental_alluvials.docx
42 File Format: docx
43 Title: Alluvial plots for all cancer types
44
45 Supplementary File S2:
46 File Name: FileS2_CancerNetworks_Cistrome.cys
47 File Format : cys (Cytoscape Session File)

1   Title: Cistrome Cancer TF-to-glycogene subnetworks
2
3   R scripts used for analysis are available at:
4   https://github.com/neel-lab/TF-glycogene
5

8   **ACKNOWLEDGEMENT**

9   We gratefully acknowledge helpful discussions with Prof. Rudiyanto Gunawan

11   **FUNDING**

12   This work was supported US National Institutes of Health grants HL103411, GM133195 and

13   GM126537.

1 **REFERENCES:**
(1)   Zhu, Y.; Groth, T.; Kelkar, A.; Zhou, Y.; Neelamegham, S. *Glycobiology* **2020**. https://doi.org/10.1093/glycob/cwaa074.
(2)   Neelamegham, S.; Mahal, L. K. *Curr. Opin. Struct. Biol.* **2016**, 145–152. https://doi.org/10.1016/j.sbi.2016.09.013.
(3)   Lambert, S. A.; Jolma, A.; Campitelli, L. F.; Das, P. K.; Yin, Y.; Albu, M.; Chen, X.; Taipale, J.; Hughes, T. R.; Weirauch, M. T. *Cell* **2018**, 650–665. https://doi.org/10.1016/j.cell.2018.01.029.
(4)   Namba, S.; Sato, K.; Kojima, S.; Ueno, T.; Yamamoto, Y.; Tanaka, Y.; Inoue, S.; Nagae, G.; Iinuma, H.; Hazama, S.; Ishihara, S.; Aburatani, H.; Mano, H.; Kawazu, M. *Cancer Sci.* **2019**, 1096–1104. https://doi.org/10.1111/cas.13937.
(5)   Malta, T. M.; de Souza, C. F.; Sabedot, T. S.; Silva, T. C.; Mosella, M. S.; Kalkanis, S. N.; Snyder, J.; Castro, A. V. B.; Noushmehr, H. *Neuro. Oncol.* **2018**, 608–620. https://doi.org/10.1093/neuonc/nox183.
(6)   Chen, L.; Zhang, W.; Fregien, N.; Pierce, M. *Oncogene* **1998**, 2087–2093. https://doi.org/10.1038/sj.onc.1202124.
(7)   Svensson, E. C.; Conley, P. B.; Paulson, J. C. *J. Biol. Chem.* **1992**, 3466–3472.
(8)   Jiang, Z.; Liu, Z.; Zou, S.; Ni, J.; Shen, L.; Zhou, Y.; Hua, D.; Wu, S. *Oncol. Rep.* **2016**, 1353–1360. https://doi.org/10.3892/or.2016.4959.
(9)   Muramoto, K.; Tange, R.; Ishii, T.; Miyauchi, K.; Sato, T. *Biol. Pharm. Bull.* **2017**, 1282–1288. https://doi.org/10.1248/bpb.b17-00212.
(10)   Weiss, R. J.; Spahn, P. N.; Toledo, A. G.; Chiang, A. W. T.; Kellman, B. P.; Li, J.; Benner, C.; Glass, C. K.; Gordts, P. L. S. M.; Lewis, N. E.; Esko, J. D. *Proc. Natl. Acad. Sci. U. S. A.* **2020**, 9311–9317. https://doi.org/10.1073/pnas.1920880117.
(11)   Mei, S.; Qin, Q.; Wu, Q.; Sun, H.; Zheng, R.; Zang, C.; Zhu, M.; Wu, J.; Shi, X.; Taing, L.; Liu, T.; Brown, M.; Meyer, C. A.; Liu, X. S. *Nucleic Acids Res.* **2016**, D658–D662. https://doi.org/10.1093/nar/gkw983.
(12)   Jassal, B.; Matthews, L.; Viteri, G.; Gong, C.; Lorente, P.; Fabregat, A.; Sidiropoulos, K.; Cook, J.; Gillespie, M.; Haw, R.; Loney, F.; May, B.; Milacic, M.; Rothfels, K.; Sevilla, C.; Shamovsky, V.; Shorser, S.; Varusai, T.; Weiser, J.; Wu, G.; Stein, L.; Hermjakob, H.; D'Eustachio, P. *Nucleic Acids Res.* **2020**, D498–D503. https://doi.org/10.1093/nar/gkz1031.
(13)   Grossman, R. L.; Heath, A. P.; Ferretti, V.; Varmus, H. E.; Lowy, D. R.; Kibbe, W. A.; Staudt, L. M. *N. Engl. J. Med.* **2016**, 1109–1112. https://doi.org/10.1056/NEJMp1607591.
(14)   Mei, S.; Meyer, C. A.; Zheng, R.; Qin, Q.; Wu, Q.; Jiang, P.; Li, B.; Shi, X.; Wang, B.; Fan, J.; Shih, C.; Brown, M.; Zang, C.; Liu, X. S. *Cancer Res.* **2017**, e19–e22. https://doi.org/10.1158/0008-5472.CAN-17-0327.
(15)   Tang, Q.; Chen, Y.; Meyer, C.; Geistlinger, T.; Lupien, M.; Wang, Q.; Liu, T.; Zhang, Y.; Brown, M.; Liu, X. S. *Cancer Res.* **2011**, 6940–6947. https://doi.org/10.1158/0008-5472.CAN-11-2091.
(16)   Wang, S.; Sun, H.; Ma, J.; Zang, C.; Wang, C.; Wang, J.; Tang, Q.; Meyer, C. A.; Zhang, Y.; Liu, X. S. *Nat. Protoc.* **2013**, 2502–2515. https://doi.org/10.1038/nprot.2013.150.
(17)   Bernard, P. S.; Parker, J. S.; Mullins, M.; Cheung, M. C. U.; Leung, S.; Voduc, D.; Vickery, T.; Davies, S.; Fauron, C.; He, X.; Hu, Z.; Quackenhush, J. F.; Stijleman, I. J.; Palazzo, J.; Matron, J. S.; Nobel, A. B.; Mardis, E.; Nielsen, T. O.; Ellis, M. J.; Perou, C. M. *J. Clin. Oncol.* **2009**, 1160–1167. https://doi.org/10.1200/JCO.2008.18.1370.
(18)   Neelamegham, S.; Aoki-Kinoshita, K.; Bolton, E.; Frank, M.; Lisacek, F.; Lütteke, T.; O'Boyle, N.; Packer, N. H.; Stanley, P.; Toukach, P.; Varki, A.; Woods, R. J. *Glycobiology* **2019**, 620–624. https://doi.org/10.1093/glycob/cwz045.
(19)   Cheng, K.; Zhou, Y.; Neelamegham, S. *Glycobiology* **2017**, 200–205. https://doi.org/10.1093/glycob/cww115.

(20) Zhou, T.; Ren, J.; Medo, M.; Zhang, Y.-C. *Phys. Rev. E* **2007**, 46115. https://doi.org/10.1103/PhysRevE.76.046115.

(21) Miklossy, G.; Hilliard, T. S.; Turkson, J. *Nat. Rev. Drug Discov.* **2013**, 611–629. https://doi.org/10.1038/nrd4088.

(22) Bi, X.; Hameed, M.; Mirani, N.; Pimenta, E. M.; Anari, J.; Barnes, B. J. *Breast Cancer Res.* **2011**, R111. https://doi.org/10.1186/bcr3053.

(23) Yanai, H.; Negishi, H.; Taniguchi, T. *Oncoimmunology* **2012**, 1376–1386. https://doi.org/10.4161/onci.22475.

(24) Khan, H. A.; Margulies, C. E. *Front. Genet.* **2019**, 591. https://doi.org/10.3389/fgene.2019.00591.

(25) Sampieri, L.; Di Giusto, P.; Alvarez, C. *Front. Cell Dev. Biol.* **2019**, 123. https://doi.org/10.3389/fcell.2019.00123.

(26) Romagnoli, M.; Belguise, K.; Yu, Z.; Wang, X.; Landesman-Bollag, E.; Seldin, D. C.; Chalbos, D.; Barillé-Nion, S.; Jézéquel, P.; Seldin, M. L.; Sonenshein, G. E. *Cancer Res.* **2012**, 6268–6278. https://doi.org/10.1158/0008-5472.CAN-12-2270.

(27) Dimitroff, C. J. *Proc. Natl. Acad. Sci.* **2019**, 13729–13737. https://doi.org/10.1073/pnas.1900268116.

(28) Yu, W.; Huang, C.; Wang, Q.; Huang, T.; Ding, Y.; Ma, C.; Ma, H.; Chen, W. *Tumour Biol. J. Int. Soc. Oncodevelopmental Biol. Med.* **2014**, 10943–10951. https://doi.org/10.1007/s13277-014-2403-1.

(29) Nan, X.; Ng, H. H.; Johnson, C. A.; Laherty, C. D.; Turner, B. M.; Eisenman, R. N.; Bird, A. *Nature* **1998**, 386–389. https://doi.org/10.1038/30764.

(30) Jones, P. L.; Veenstra, G. J.; Wade, P. A.; Vermaak, D.; Kass, S. U.; Landsberger, N.; Strouboulis, J.; Wolffe, A. P. *Nat. Genet.* **1998**, 187–191. https://doi.org/10.1038/561.

(31) Chen, F.; Liu, X.; Bai, J.; Pei, D.; Zheng, J. *Oncol Rep* **2016**, 1227–1236. https://doi.org/10.3892/or.2015.4515.

(32) Tian, F.; DaCosta Byfield, S.; Parks, W. T.; Yoo, S.; Felici, A.; Tang, B.; Piek, E.; Wakefield, L. M.; Roberts, A. B. *Cancer Res.* **2003**, 8284–8292.

(33) Petersen, M.; Pardali, E.; van der Horst, G.; Cheung, H.; van den Hoogen, C.; van der Pluijm, G.; Ten Dijke, P. *Oncogene* **2010**, 1351–1361. https://doi.org/10.1038/onc.2009.426.

(34) Narimatsu, Y.; Joshi, H. J.; Nason, R.; Van Coillie, J.; Karlsson, R.; Sun, L.; Ye, Z.; Chen, Y.-H.; Schjoldager, K. T.; Steentoft, C.; Furukawa, S.; Bensing, B. A.; Sullam, P. M.; Thompson, A. J.; Paulson, J. C.; Büll, C.; Adema, G. J.; Mandel, U.; Hansen, L.; Bennett, E. P.; Varki, A.; Vakhrushev, S. Y.; Yang, Z.; Clausen, H. *Mol. Cell* **2019**, 394-407.e5. https://doi.org/10.1016/j.molcel.2019.05.017.

(35) Yevshin, I.; Sharipov, R.; Kolmykov, S.; Kondrakhin, Y.; Kolpakov, F. *Nucleic Acids Res.* **2019**, D100–D105. https://doi.org/10.1093/nar/gky1128.

(36) Marbach, D.; Lamparter, D.; Quon, G.; Kellis, M.; Kutalik, Z.; Bergmann, S. *Nat. Methods* **2016**, 366–370. https://doi.org/10.1038/nmeth.3799.

(37) Sonawane, A. R.; Platig, J.; Fagny, M.; Chen, C.-Y.; Paulson, J. N.; Lopes-Ramos, C. M.; DeMeo, D. L.; Quackenbush, J.; Glass, K.; Kuijjer, M. L. *Cell Rep.* **2017**, 1077–1088. https://doi.org/10.1016/j.celrep.2017.10.001.

(38) Taniguchi, N.; Honke, K.; Fukuda, M.; Narimatsu, H.; Yamaguchi, Y.; Angata, T. 2nd ed.; Springer, 2014; Vol. 1–2. https://doi.org/10.1007/978-4-431-54240-7.

(39) Varki, A.; Cummings, R. D.; Esko, J. D.; Stanley, P.; Hart, G. W.; Aebi, M.; Darvill, A. G.; Kinoshita, T.; Packer, N. H.; Prestegard, J. H.; Schnaar, R. L.; Seeberger, P. H. .

(40) Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N. S.; Wang, J. T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T. *Genome Res.* **2003**, 2498–2504. https://doi.org/10.1101/gr.1239303.

(41) Csardi, G.; Nepusz, T. *InterJournal* **2006**, 1695.