Multifractality in postural sway supports quiet eye training in aiming tasks: A study of golf putting

Noah Jacobson[1], Quinn Berleman-Paul[1], Madhur Mangalam[2], Damian G. Kelty-Stephen[1], and Christopher Ralston[1]

[1]Department of Psychology, Grinnell College, Grinnell, IA 50112, USA

[2]Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115, USA

Author for correspondence:

Damian G. Kelty-Stephen

e-mail: keltysda@grinnell.edu

Ethics statement. All participants provided verbal and written informed consent approved by the Institutional Review Board (IRB) at Grinnell College (Grinnell, IA).

Data accessibility. All data analyzed in the present study are available upon request.

Author contributions: Noah Jacobson: Conceptualization, Methodology, Software, Investigation, Resources, Data curation, Writing - original draft, Writing - review & editing. Quinn Berleman-Paul: Methodology, Software, Investigation, Resources, Data curation, Writing - review & editing. Madhur Mangalam: Conceptualization, Methodology, Software, Validation, Formal analysis, Data curation, Writing - review & editing, Visualization, Supervision, Project administration. Damian G. Kelty-Stephen: Conceptualization, Methodology, Validation, Formal analysis, Writing - original draft, Writng - review & editing, Supervision. Christopher Ralston: Conceptualization, Methodology, Validation, Resources, Supervision, Project administration, and Funding Acquisition.

Competing interests. The authors have no competing interests to declare.

Supplementary materials

Supplementary Table S1. Subset of Poisson regression model output including main and interaction effects of time and perturbation of eye height.

Abstract

The 'quiet eye' (QE) approach to visually-guided aiming behavior invests fully in perceptual
information's potential to organize coordinated action. Sports psychologists refer to QE as the stillness
of the eyes during aiming tasks and increasingly into self- and externally-paced tasks. Amidst the
'noisy' fluctuations of the athlete's body, quiet eyes might leave fewer saccadic interruptions to the
coupling between postural sway and optic flow. Postural sway exhibits fluctuations whose multifractal
structure serves as a robust predictor of visual and haptic perceptual responses. Postural sway generates
optic flow centered on an individual's eye height. We predicted that perturbing the eye height by
attaching wooden blocks below the feet would perturb the putting more so in QE-trained participants
than participants trained technically. We also predicted that QE's efficacy and responses to
perturbation would depend on multifractality in postural sway. Specifically, we predicted that less
multifractality would predict more adaptive responses to the perturbation and higher putting accuracy.
Results showed that lower multifractality led to more accurate putts, and the perturbation of eye height
led to less accurate putts, particularly for QE-trained participants. Models of radial error (i.e., the
distance between the ball's final position and the hole) indicated that lower estimates of multifractality
due to nonlinearity coincided with a more adaptive response to the perturbation. These results suggest
that reduced multifractality may act in a context-sensitive manner to restrain motoric degrees of
freedom to achieve the task goal.

Keywords: biotensegrity, degree of freedom, fractal, posture, tensegrity, visual adaptation

1. Introduction

Aiming our behaviors into the visible world requires an ongoing awareness of how our body fits in the world around it. The difficulty of sorting out this bodily fit finds its classic encoding in the 'degrees of freedom' problem: even if a fine-grained choreography of goal-directed movement that stipulates all individual motoric degrees of freedom were possible, this fine-grained choreography would probably be too rigid to adapt to the task or context (Bernstein, 1967). However, another perspective suggests that the abundance of degrees of freedom is not a problem but a 'blessing' (Latash, 2012). According to this view, there is no need to stipulate the infinitude of motoric variables at very fine grains, and on the contrary, task or intention organizes the movement system at the relatively coarse grain of the whole body and context (Latash, 2020). The infinitude thus becomes a guarantee of adaptive flexibility. This positive view of motor abundance appears in recent studies of aiming behaviors like the golf swing, whereby a clearly defined target in the context draws visual attention away from specific postures of the body, and indeed, experts with their eyes on the target show no single optimized mode of execution (Morrison et al., 2016). Hence, instead of requiring explicit choreography, the movement system's organization may readily emerge from perceptual linkages to the surrounding context (Stoffregen, 1985, 1986; Stoffregen & Riccio, 1990).

1.1. Quiet eye vs. technical training

Supporting task performance by emphasizing perceptual linkages vs. explicit choreography is a clear theme of research on 'quiet eye' (QE). Sports psychologists refer to QE as the stillness of the eyes during aiming tasks, specifically the final gaze fixation to a single location or object in the task for a minimum of 100 ms within 3 degrees of visual angle immediately before the movement is initiated (Rienhoff et al., 2016). In the case of a golf swing, 'technical training' that communicates an explicit choreography of the arm and leg placement leads to less accuracy than a QE strategy. Expert golfers show more frequent and longer gaze fixations on specific aspects of the putting situation (e.g., golf ball, putting surface, target hole). Novices tend to fixate more introvertedly (e.g., on the clubhead during the backswing phase) and focus on the ball only after it has been hit with the club (Moore et al., 2012; Vickers, 1992, 1996; Vine et al., 2013, 2017). Visual guidance by external focus of attention, that is, focusing on the target leads to a more effective golf swing than technical training of the postures and limb movements composing a golf swing. Indeed, in contrast with technical training, QE training of golf putting leads to improved putting accuracy as well as to a reduction in heart rate and muscle activity at the moment of impact (Moore et al., 2012). Thus, it is believed that instructing athletes by directing their perceptual linkages outward to the visual field supports higher task performance as well as leads to more expert-like movement and physiology.

QE research opens a fascinating vantage point on the perceptual linkages anchoring organisms to their visual contexts and the synergies transforming visual information into action (e.g., Kelty-Stephen et al., 2020; Mangalam, Lee, et al., 2021; Profeta & Turvey, 2018). Research on QE has mostly examined the visual-cognitive bases for the relatively more extraverted gaze into the task context (Panchuk & Vickers, 2006; Rienhoff et al., 2016; Vickers, 2011), and the relationship of QE to the body, the brain, and the nervous system has received relatively less attention (Davids & Araújo, 2016; Renshaw et al., 2019; Shine et al., 2018). To fill this gap, a 'postural-kinematic hypothesis' has begun to complement the canonical 'visual hypothesis' that QE was a primarily visual-cognitive mechanism (Gallicchio & Ring, 2020). So far, this postural-kinematic hypothesis emphasizes only slower movement times, that is, longer duration offering the foundation for longer fixations. This postural-kinematic hypothesis is a constructive first step. In this study, we aim to embed more of the postural dynamics into this hypothesis, beyond simply movement duration. Additionally, instead of reporting on gaze measurements, we will use QE as a training manipulation and focus on investigating how postural sway supports the effect of the training.

1.2. Optic flow and postural sway can support the quiet eye strategy

A crucial consideration for postural-kinematic contributions to visually guided tasks is optic flow (Stoffregen, 1985, 1986; Stoffregen & Riccio, 1990). Amidst the 'noisy' fluctuations of the athlete's body, quieting the eyes might leave fewer saccadic interruptions to the coupling between optic flow and postural sway. Optic flow consists of the differential expansion, contraction, or rotation of visible objects and surfaces as an individual moves about. It is specific to both the individual's movement and the contextual layout of objects and surfaces. Indeed, virtual visual displays that impose artificial optic flow also engender anticipatory behaviors (e.g., steering, braking) as if the objects and surfaces entailed by the optic flow were out there—not just for humans but for other animal species and even robots (Serres & Ruffier, 2017). Although most prominent in locomotion-related contexts, postural sway is no less capable of generating informative optic flow in non-locomotion contexts (Stoffregen et al., 2005). Because research on QE has mostly focused on the fixation of the eye gaze, QE's relationship to optic flow remains a critical knowledge gap (de Oliveira, 2016).

On the face of it, the preceding arguments might ring hollow at first. Certainly, optic flow requires movement, and it could seem wholly unrelated to the issue of visual fixations supporting wholly internal processing of incoming information into a motor program for the body's subsequent control. This gap between QE and optic flow follows closely from ongoing controversy in the literature between ecological vs. information-processing explanations. Optic flow has ecological roots in Gibson's (1979) disavowal of information-processing. Those roots could potentially raise fears that this work might disregard the extensive, diligent work on the information-processing account of QE. Indeed, a rare mention of 'optic flow' in the information-processing approach to QE weds it firmly to 'the teachings of Gibson,' so concerned with invariance as to 'treat every gaze as the same,' and reluctant to respect the fine-grained dynamics of gaze (Vickers, 2009, p. 286). We make full disclosure: the present use of multifractal modeling of postural access to optic flow is inspired by Gibson's appeal to flows and nestings these flows across many scales (Kelty-Stephen & Wallot, 2017). However, we come to this dialogue with first-hand experience examining the rich heterogeneity of gaze dynamics (e.g., Kelty-Stephen & Mirman, 2013; Stephen & Mirman, 2010) and with such interest in cognitive process as for Gibsonians to openly doubt that some of us could be Gibsonians (Raja & Anderson, 2019). Indeed, our interest in how exploratory behaviors might change perceptual response sets us firmly outside one of the stricter readings of Gibsonian invariance (e.g., Turvey, 1977).

Rather than digging into old trenches, we think that optic flow has a theoretical reach that might pick up where this information-processing account of QE remains incomplete. The theoretical reach of optic flow has grown well beyond Gibson's original intent and usage. For instance, we have already cited cases of optic flow that would have been startling through the strictly Gibsonian lens: information-processing robots use optic flow as well as Gibson's organisms might (Serres & Ruffier, 2017), and optic flow does not require locomotion but can thrive on postural sway (Stoffregen et al., 2005). In the spirit of making the so far dialogue between ecological and information-processing approaches more constructive, the preceding two examples suggest that optic flow might be much more subtle (e.g., as in standing 'still') but also perfectly amenable to information processing (e.g., as in supporting computational models for controlling action).

Meanwhile, without optic flow to support it, QE's information-processing accounts are incomplete and unmoored from their own explicit requirements and proposals. According to a recent summary, the information-processing view of QE is that the eye's stillness provides time and attentional resources useful for the preprogramming of movements used in the aimed behavior (Vickers, 2009). This point is defensible but incomplete: visual-cognitive processes benefit only partially from a motor program's stipulation. Even non-ecological explanations have become newly interested in emergent control parameters that sit outside of any simple stipulation by an internal motor

program (Vecera et al., 2014). Visual attention is most effective when it is most task- and context-
sensitive. The generic usefulness of QE points to some long-unanswered questions. Are the motor
programs implicated in QE flexible and vast enough to accommodate both bodily degrees of freedom
and contextual ones? Or are these motor programs elegantly minimal enough to leave room for non-
programmed factors to support? Neither option is unappealing on its face, but they both reflect an
underdeveloped delivery on the information-processing account's proposals about QE. Yes, longer
looking times can support stronger, more enriched motor programs (e.g., Witt & Proffitt, 2008).
However, it is not always so simple: either looking time is not always a reliable predictor of motor
planning, or the motor program is flexible enough to be rewritten on the fly during the program's
execution (Hajnal et al., 2014, 2016).

Specifically, just as it neglects the finer print about its motor programs, QE's information-
processing account has also neglected its claims about the role of action. The information-processing
account offers QE a bidirectional link between gaze, attention, and action (Vickers, 2009). However,
whereas this view may respect bidirectionality between attention and gaze (i.e., attentive search for
information and looking to what cues attention), the action appears strictly as the outcome passively
inheriting whatever the preprogramming entails and only following motor commands. If motor
programs generate the consequent swings, then the information-processing approach has yet to
elaborate upon two points: first, what bodywide movement coordination supports the postural control
allowing the steady visual fixation and, second, how the preprogramming contributes to the externally
focused movement. For instance, especially as participants repeat the aiming tasks in front of
potentially the same experimental display, the evidence suggests more variety in expert movements
than sameness, mainly as they engage visually in the external focus of attention (Morrison et al., 2016).
This point raises some questions. If QE is merely the same sort of fixation every time, irrespective of
movement, then why should programs carrying similar, static information give rise to varied bodily
execution of the program? If the motor programs that QE supports vary, what is the possible source of
variability? In the former case, the programs might underspecify the movements, and in the latter case,
the programs would seem to embody more variety than the fixity of the visual scene. The whole issue
entailed by the Bernsteinian 'degrees of freedom' concern is that there is no clear programmatic path
from intention to bodily execution. Suppose action did participate in the same bidirectional relationship
that information-processing approaches to QE observe between attention and gaze. In that case, there
might be a way forward, allowing for action to participate in the tuning of programs to ongoing bodily
postures and physiological changes throughout repeated behaviors.

To summarize, the gap between QE and optic flow is critical because the information-
processing approach aligns the fixity of the eyes with the accuracy of aimed behavior. However, there
is a little account of the body between the fixations and the accurate behaviors. This omission of the
body is immensely curious and potentially problematic because movement exhibits rich variability. It
has been difficult for all proposed motor programs to specify accurate goal-directed behavior
(Feldman, 2018) and no less difficult when eye movements are implicated (Feldman & Zhang, 2020).
Moreover, if we can leave off old associations and let optic flow continue to take on shapes beyond the
original Gibsonian imaginings, we might reinstate the body to an account of QE without any threat of
thwarting or disregarding prior art. We aim to rekindle this dialogue between cognitive psychology and
ecological psychology, but we aim to do so without requiring disavowal of any mutually opposed pre-
theoretical commitments.

The present work aims to replicate the beneficial effects of QE on putting performance, and
more importantly, to understand how eye height and optic flow might support QE. We used a single
session in the style of Moore et al.'s (2012) golf-putting training paradigm to test for rapid attunement
to changes in information from optic flow and postural sway. Optic flow centers on an external focus

of control, aligned with their eye height (e.g., Warren, 1984), and when we perturb eye height, we
destabilize optic flow, and the organism needs to learn new relationships between bodily movement
and movement of visible surfaces (Frenz et al., 2003). So, we aimed to test whether perturbing the optic
-flow parameter of eye height perturbs the effect of QE training in golf putting more so than technical
training. In non-golf contexts, for instance, manipulating eye height by adding wooden blocks to
participants' shoes alters participants' affordance judgments significantly, leading to an initial
underestimation of the optimal climbing height for a stair and sitting height for a chair (Warren, 1984).
However, participants adjust to the new eye height and make more accurate decisions shortly afterward
(Mark, 1987; Mark et al., 1990). Hence, we expected that adding wooden blocks—(henceforth, 'clogs',
to distinguish from 'blocks' of experimental trials)—to the feet might upset putting accuracy in the
QE-trained participants initially, but that this accuracy would rebound as participants adjust to new eye
height (Rienhoff et al., 2016).

1.3. Multiscaled organization of postural sway entails a multifractal structure

Besides manipulating eye height, our expectations about the optic-flow foundation for QE also
included a hypothesis about postural sway moderating effects of perturbed eye height. If optic flow
could support QE, we expected that QE-trained response to eye height perturbation would depend on
postural sway. Indeed, in the affordance research on reattunement to eye-height perturbations, postural
sway supported the adaptation of the eye-height relationship to judgments of optimal sitting height
(Mark et al., 1990). However, the exact aspects of sway supporting adaptation have been unclear. It
was not simply that more sway facilitated this adaptation because, for instance, an 'awkward stance'
requiring more compensatory postural adjustments did not support visual adaptation as much as
comfortable standing did. An additional challenge in identifying adaptation-supporting aspects of sway
could be that sway unfolds across multiple timescales. At longer scales, QE training itself might
influence sway by training an explicit use of visual information. At shorter timescales, adding clogs
might destabilize posture initially, and then at the shortest timescales, sway might be implicated in
adapting to this perturbation of eye height. Hence, judicious accounting of sway's contribution to
visual adaptation requires an appreciation of sway's multi-scaled organization.

Then again, the multi-scaled organization of sway may itself reveal a key to predicting—if not
explaining—how the movement system adapts its visually-guided actions. Interactions across multiple
timescales entail a 'multifractal' structure in sway. Multifractality refers to multiple (i.e., 'multi-')
relationships among fluctuations and timescales, with these relationships following power laws with
potentially fractional (i.e., '-fractal') exponents. Interactions across timescales engender a nonlinear
patterning of fluctuations that multifractal analysis can diagnose and quantify (Ihlen & Vereijken,
2010). Specifically, multifractal analysis estimates a spectrum of fractal exponents whose width W_{MF}
quantifies "multifractality." When we compare measured series' W_{MF} to spectrum widths of a set of
best-fitting linear models ('linear surrogates') of those series, multifractal evidence of nonlinearity
manifests as a significant difference between W_{MF} for the original series and the average W_{MF} for linear
surrogates. The t-statistic, t_{MF}, comparing the original series' W_{MF} to the surrogates' W_{MF} serves as an
estimate of what we call 'multifractal nonlinearity' (i.e., how much multifractality in the original series
is due to nonlinear interactions across timescales).

1.4. Multifractal nonlinearity predicts perceptual responses to perturbations of task constraints

Multifractality, W_{MF}, in postural sway is a robust predictor of perceptual responses in both visual and
haptic media (Doyon et al., 2019; Hajnal et al., 2018; Teng et al., 2016). Multifractal nonlinearity, t_{MF},
in postural sway predicts perceptual responses as well, not just in the case of nonvisual judgments of
length and heaviness of manually-wielded objects (Mangalam & Kelty-Stephen, 2020) but also in
visuomotor tasks (Bell et al., 2019; Booth et al., 2018) and in response to perturbations, such as those

due to prismatic goggles in a visual aiming task (Carver et al., 2017). Now, we aimed to test whether t_{MF} could predict the responses to perturbations of eye height in QE-trained golf putting.

We expected that postural sway most adaptive to the perturbation of eye height would show relatively narrow multifractal spectra that would still differ significantly from those of the surrogates. So far, postural sway shows relatively narrower multifractal spectra (i.e., smaller W_{MF}) for younger participants (Munafo et al., 2016), those less likely to experience motion sickness (Koslucher et al., 2016), and those focusing on more proximal surfaces in optic flow (Li et al., 2020). Hence, greater W_{MF} in postural sway might counteract effects of QE. All these findings are qualified by a failure of W_{MF} to correlate with canonical measures of sway like standard deviation. However, postural stability is only ambiguously related to the standard deviation of sway, requiring neither too much nor too little variability in sway (Chen et al., 2008; Rajachandrakumar et al., 2018). Meanwhile, both empirical and theoretical work suggests that t_{MF} is more closely related to postural stabilizing. Postural stabilization following a perturbation to the base of support shows a rapid reduction of t_{MF} (Kelty-Stephen, 2018). Theoretical simulation also predicts that random perturbations to individual scales of a nested system lead to narrower multifractal spectra that still differ from linear surrogates' spectra (i.e., smaller W_{MF} and smaller-yet-significant t_{MF}; Lee & Kelty-Stephen, 2017). Hence, adapting to the posturally perturbed visual information might involve exhibiting smaller-yet-significant t_{MF}.

1.5. Golf-putting task and hypothesis

The participants were instructed to take golf putts towards a circular, hole-sized target painted on a green putting surface in the laboratory—over four blocks of ten trials each. One group received QE training, and the other group received technical training specifying how to move their limbs to make the golf putt. Wooden clogs were attached to the feet of all participants to perturb their eye height. Putting performance was treated continuously in terms of unsigned radial distance from the target (i.e., unsigned radial error).

1.5.1. Hypothesis-1: Multifractal fluctuations in postural sway will lead an eye height perturbation to interrupt QE-trained performance more so than technical-trained performance

We hypothesized that the eye-height perturbation due to clogs would lead to lower putting accuracy characterized by higher unsigned radial error following QE training than technical training. We predicted that the clog's effects on QE would depend on the multifractal nonlinearity, t_{MF}, of sway above and beyond any effects of multifractality (i.e., W_{MF}) alone (Hypothesis-1a). We also predicted that lower t_{MF} of sway would be associated with more adaptive responses to the eye-height perturbation, leading to gradual increase and subsequent decay of radial error across trials within the blocks with the clogs on (Hypothesis-1b).

2. Methods

2.1. Participants

Twenty undergraduate students (11 women, *mean*±1*s.d.* age = 19.5±1.2 years, all right-handed, Oldfield, 1971) with normal or corrected vision participated in this study. All participants provided written informed consent approved by the Institutional Review Board (IRB) at Grinnell College (Grinnell, IA).

2.2. Procedure

In an hour-long session, the participants completed a single putt on each of 40 trials. Prior to the training trials, half of the participants received QE instructions (Table 1, left column) while the other half received technical putting instruction (Table 1, right column), replicating Moore et al.'s (2012)

methodology. The technical instructions mirrored the QE instructions but did not describe the eye-
related behaviors, hence mimicking standard golf instruction. After verbally confirming that they
understood the instructions, the participants watched a video demonstrating a QE-specific putting
stroke (https://www.youtube.com/watch?v=lfWuL1klbsY&feature=youtu.be) or a standard putting
stroke (https://www.youtube.com/watch?v=HOYZn4MYmDI). The researcher directed the QE-
strained participants to the key features of the golfer's gaze control and the technically-trained
participants to the putting mechanics.

The participant performed the golf putts on a 6×15 feet standard indoor All Turf Matts^{TM}
artificial turf mat using a 1.68-inch diameter PING Sigma 2 Anser Stealth^{TM} putter and regular-sized
white golf balls. The participants completed straight putts ten feet away towards a 2.5-inch diameter
circular orange dot, designed to fit the golf ball. This format allowed for more stringent measurement
and gave the participants a more focused target.

The participants completed 20 training trials (treated as two blocks of 10 trials per block in
regression modeling). Following a short 5-minute break, the participants completed two blocks of ten
test trials per block (i.e., 20 test trials total) in the absence of instructions. The participants were
randomly assigned to complete either their first or second block of ten trials while wearing 5-cm-raised
wooden blocks (i.e., clogs) attached to their shoes (they completed the other block of ten test trials
under the same conditions as the 20 training trials). Before the testing began, the participants were
allowed to walk around and adjust to the clogs to reduce fall risk.

2.3. Measures

The task performance was defined in terms of the radial error (the distance from the target at which the
ball stopped, in feet) and the number of putts successfully 'holed.' Because the target was not a real
hole, the putt was counted as holed in either of two events: the ball came to rest within the orange dot
on the floor or within 2 feet directly beyond the hole (Cooke et al., 2010; Vine et al., 2011). Postural
sway was measured using a smartphone with the Physics Toolbox Suite application
(http://vieryasoftware.net) strapped to the participants' lower back. The app recorded the acceleration
of the torso along x-, y-, and z-dimensions at 100 Hz. An experimenter manually started and stopped
recording for each putt. Recording began after the participant set up over the ball and stopped once the
ball stopped moving.

2.4. Multifractal analysis

2.4.1 Multifractal spectrum width of accelerometer displacement series

For each putt, we computed an accelerometer displacement series for multifractal analysis as follows.
The accelerometer data encodes an N-length series of accelerations of the torso along x-, y-, and z-
dimensions centered on an origin on the smartphone. A displacement series of length $N-1$ was
computed by: 1) taking the first differences for the variable (e.g., if we use the letters i and j to indicate
individual values in sequence, $x_{diff}(j) = x(i) - x(i-1)$ for each ith value of x, for $2 < i < N$,
$1 < j = i - 1 < N - 1$, 2) squaring and summing the differences of each dimension's displacements
(i.e., $x^2_{diff} + y^2_{diff} + z^2_{diff}$), and 3) taking the square root of this sum (i.e., $\sqrt{(x^2_{diff} + y^2_{diff} + z^2_{diff})}$).

Multifractal spectra $f(\alpha)$ were estimated for each accelerometer displacement series, using
Chhabra and Jensen's (Chhabra & Jensen, 1989) method. The first step was to partition the series into
subsets (or 'bins') of various sizes, from four points to a fourth of series' length (Fig. 1, left). The
second step was to examine how the proportion of displacements within each bin changed with bin size
or timescale (Fig. 1, right). The third step was to estimate two distinct exponents: an exponent α
describing how the average proportion grows with bin size and an exponent f describing how

Shannon's (Shannon, 1948) entropy of bin proportion changes with bin size. The later steps were all
iterations of this third step, calculating a mass μ by applying an exponent q to bin proportions and
using $\mu(q)$ to weight proportions selectively according to proportion size. q greater than 1 weights
larger-proportion bins, and q lesser than 1 weights smaller-proportion bins (Fig. 2, left). q-based mass
$\mu(q)$-weighting generalizes single α and f estimates from earlier steps into continual $\alpha(q)$ and $f(q)$—
with $\alpha(q)$ describing how mass-weighted average bin proportion changes with bin size and $f(q)$
describing how Shannon's entropy of bin masses $\mu(q)$ with bin size (Fig. 2, top right).

Multifractal analysis quantifies heterogeneity as variety in $\alpha(q)$ and $f(q)$. The ordered pairs
$(\alpha(q), f(q))$ constitutes the multifractal spectrum (Fig. 2, bottom right). We included $\alpha(q)$ and $f(q)$
only when mass-weighted proportions were defined and when both of their corresponding relationships
to bin size correlated at $r > 0.995$ on logarithmic plots, for bin sizes $4, 8, 12, \ldots N/4$ and $-300 < q <$
300, where N is the number of samples in the series.

2.4.2. Surrogate comparison

Multifractal spectrum width is sensitive to nonlinear interactions across timescales as well as to linear
temporal structure. Hence, evidence of nonlinear interactions across timescales requires comparing the
original series' multifractal spectrum width W_{MF} to those computed for a sample of linear surrogate
series (Fig. 3). These linear surrogate series retain the same value as the original series but in a
different order that preserves the original series' linear autocorrelation but destroys original sequence
(i.e., iterative amplitude-adjusted Fourier-transform or IAAFT (Ihlen & Vereijken, 2010; Schreiber &
Schmitz, 1996); Figs. 4a to d). Comparing W_{MF} of the original and surrogate series (Figs. 4e, f)
produces a t-statistic t_{MF}, a measure of nonlinear interactions across timescales that is standardized
across series with differing linear temporal structure. t_{MF} was computed for each original series.

2.5. Mixed-effects modeling

348 The data were submitted to two kinds of mixed-effect modeling, namely, logistic and Poisson, using
349 the glmer() function in *R*-package *lme4* (Bates et al., 2007). The subsequent subsections provide details
350 of the predictors and the models.

2.5.1. Effects of time

The effects of time were modeled using trial number within a block (trial$_{block}$), block number (block),
and interactions of trial number with block number (trial$_{block}$ × block), and trial number across blocks
(trial$_{exp}$). The model used orthogonal polynomials of each of both predictors up to 3^{rd} order (i.e., cubic).
The primary reason for the polynomial terms was to reveal how and when the perturbation of wearing
clogs might manifest and consequently subside. Cubic polynomials allow the simplest way to
accomplish this goal by allowing a peak and subsequent decay to a flat, quiescent portion as part of the
same continuous function. Additionally, unlike the quadratic function, the cubic peak is not required to
be symmetric and centered on the midpoint. Hence, the cubic function controls for any nonlinearities in
the normal course of trials without such perturbations. Whereas block is a linear effect across blocks,
including trial$_{exp}$ in the model allowed controlling for nonlinear change across blocks.

There are two important features to note about the treatment of time in this model. First, the
Poisson model encoded both classifications of trial numbers (i.e., trial$_{exp}$ and trial$_{block}$) in terms of an
orthogonal 3^{rd}-order polynomial, using the function 'poly' in *R*, to include the linear, quadratic and
cubic growth of trial number with all three terms phase-shifted so as to be uncorrelated with one
another. Second, the lack of model convergence limited the appearance of block, such that block was
neither treated nonlinearly nor in interaction with perturbation. Attempting to include the orthogonal
3^{rd}-order polynomials for block number led to a failure of model convergence, so block only appeared

as a linear term. The model also failed to converge when it included interactions of block and
perturbation because these two terms were strongly collinear.

2.5.2. Effects of manipulations

372   Manipulations included quiet-eye instruction, QE (QE = 1 for receiving QE instruction, and QE = 0 for
373   not receiving QE instruction) and perturbation of eye height, perturbation (perturbation = 1 for all ten
374   trials during the block when participants wore the clogs, and perturbation = 0 for all other trials).

2.5.3. Sway: Linear predictors

376   The accelerometer data encodes accelerations of the torso along x-, y-, and z-dimensions. Any single
377   posture corresponds to a specific point in this 3D space. Hence, sway corresponds to the deviation of
378   this 3D acceleration around the mean posture. The present analyses encoded this variability as MSD_{acc}
379   or $RMS_{acc,}$ depending on which predictor supported a convergent model, along with mean and standard
380   deviation of 3D displacement ($Mean_{disp}$ and SD_{disp}, respectively).

2.5.4. Regression modeling

A mixed-effects Poisson regression modeled the radial error (in inches) of all misses, with makes as
defined above treated as having zero radial error. Predictors included QE × perturbation × $trial_{block}$ ×
t_{MF} to our hypothesis by specifically addressing trials within each block, QE × $trial_{block}$ and QE ×
$trial_{exp}$ to control for differences in learning due to QE across a linear progression of block; a nonlinear
function of trials, QE × perturbation × W_{MF} × t_{MF}, to control for the effects of multifractal spectral
width and its relationship with t_{MF}; and MSD_{acc} × $Mean_{disp}$ × SD_{disp} to control for the effects of linear
description of sway. Modeling included lower-order interactions and component main effects (e.g.,
Allison, 1977).

3. Results

3.1.1. Non-multifractal effects on radial error

This section details the non-multifractal effects described in Supplementary Table S1 and effects
preliminary to the test of our hypothesis in Table 2. This model of radial error found a significant main
effect of QE training, specifically, a negative effect ($b = -4.40 \times 100$, $s.e.m. = 4.81 \times 10^{-1}$, $p < 0.0001$)
indicating that QE did promote greater accuracy in terms of proximity of the ball's final position to the
hole. Radial error grew linearly and quadratically across trials but showed negative cubic variation with
trial—both for trials across the experiment and within a block (Fig. 5a). The QE strategy prompted
steady decay of error across the experiment, canceling out the linear and cubic pattern of error over
trials within a block (Fig. 5b). The perturbation of eye height elicited no main effect but increased the
growth of error over trials within a block, with the QE instruction accentuating this nonlinear growth of
error later in the block. The higher-order interaction MSD_{acc} × $Mean_{disp}$ × SD_{disp} and its main
component effects and lower-order interactions showed mostly negative effects (only two of the two-
way interactions showed positive coefficients).

3.1.2. Total multifractality of sway, W_{MF}, increased radial error, but multifractality due to nonlinearity, t_{MF}, attenuated radial error at the beginning and the end of blocks

No matter what portion of multifractality is attributable to linear or nonlinear structure, the total width
W_{MF} of original series' multifractal spectra is associated with greater radial error ($b = 1.80 \times 10^{0}$, $s.e.m.$
$= 1.29 \times 10^{-1}$, $p < 0.0001$). We did not find any main effect of multifractality due to nonlinearity, t_{MF},
nor any interaction of W_{MF} × t_{MF}. However, the negative effect for t_{MF} × $trial_{block}$ (Quadratic; $b = -$
8.93×10^{-2}, $s.e.m. = 1.15 \times 10^{-2}$, $p < 0.0001$) indicated an association between multifractality due to

nonlinearity with a significant reduction of the positive quadratic profile for radial error over trials
within a block (i.e., trial$_{block}$(Quadratic); Section 3.1.1.).

3.1.3. Testing Hypothesis-1a: Perturbation of eye height increased radial error only in
combination with QE training, with greater multifractality due to nonlinearity t_{MF}, or with both

The multifractality of sway promoted the tendency of QE to increase perturbation-related radial error.
Perturbation showed a significant effect through the interaction of perturbation $\times$ W_{MF} ($b = -1.50\times10^{0}$,
$s.e.m. = 3.12\times10^{-1}$, $p < 0.0001$) canceled out the positive effect of W_{MF} on error (Section 3.1.2),
suggesting that the perturbation reduced the error-eliciting effect of W_{MF}. The interaction QE $\times$ W_{MF}
reduced error due to multifractality by roughly half ($b = -1.04\times10^{0}$, $s.e.m. = 2.03\times10^{-1}$, $p < 0.0001$),
suggesting that QE promoted greater accuracy but only in the absence of perturbation. However, the
higher-order interaction with multifractality indicates that QE increased the error attributable to the
perturbation. For instance, the QE $\times$ perturbation $\times$ W_{MF} ($b = 2.12\times10^{0}$, $s.e.m. = 4.91\times10^{-1}$, $p < 0.0001$)
reverses the apparently beneficial error-reducing interaction effect of perturbation $\times$ W_{MF}. Hence, QE
reinstated the error due to total multifractality (Section 3.1.2), and it did so only in the face of the
perturbation.

Above and beyond the effects of W_{MF} in moderating the interaction effect of QE $\times$ perturbation,
t_{MF} promoted the error-increasing effect of perturbation. Specifically, besides the moderating effect of
QE on the effect of perturbation $\times$ W_{MF}, perturbation elicited even greater radial error with increases in
t_{MF} (perturbation $\times$ W_{MF} $\times$ t_{MF}: $b = 1.54\times10^{-1}$, $s.e.m. = 3.15\times10^{-2}$, $p < 0.0001$). QE did reverse this
effect (QE $\times$ perturbation $\times$ W_{MF} $\times$ t_{MF}: $b = -2.21\times10^{-1}$, $s.e.m. = 4.97\times10^{-2}$, $p < 0.0001$). However, QE
also increased error in combination with perturbation and t_{MF} even in the absence of any moderating
effect of W_{MF} (QE $\times$ perturbation $\times$ t_{MF}: $b = 8.62\times10^{-3}$, $s.e.m. = 4.18\times10^{-3}$, $p < 0.05$). This result
supports Hypothesis-1a.

Despite the absence of a simple QE $\times$ perturbation interaction, the eye height perturbation
produced a greater model-predicted increase in radial error for QE-trained participants. To illustrate
this point, we can compare the model predictions for participants in each condition with high and low
W_{MF} and t_{MF} (with 'low' and 'high' values defined as 1st and 3rd quartiles). For participants with 'low'
W_{MF}, the eye height manipulation prompted 2.27% to 11.14% increase in radial error in QE-trained
compared to technical-trained participants for low to high t_{MF}, respectively. For participants with 'high'
W_{MF}, the eye height manipulation prompted 8.80% to 18.23% increase in radial error in QE-trained
compared to technical-trained participants for low to high t_{MF}, respectively.

3.1.4. Testing Hypothesis-1b: the QE group with greater multifractal nonlinearity in sway
showed a greater initial but progressively slower increase in radial error due to the perturbation
of eye height

The interactions of QE $\times$ trial$_{block}$ with multifractal nonlinearity followed a similar but weaker pattern
as the interactions of QE $\times$ trial$_{block}$ with perturbation. Much like the perturbation of eye height, the
interaction of QE with t_{MF} accentuated the radial error late in a block of ten trials (QE $\times$ trial$_{block}$ $\times$ t_{MF}):
it weakened the linear term ($b = -3.73\times10^{-2}$, $s.e.m. = 1.46\times10^{-2}$, $p < 0.05$), contributing to a negative-
quadratic peak in the middle of the block ($b = -1.85\times10^{-1}$, $s.e.m. = 1.66\times10^{-2}$, $p < 0.0001$) and a
positive cubic increase at the end of the block ($b = 1.13\times10^{-1}$, $s.e.m. = 1.73\times10^{-2}$, $p < 0.0001$). Hence,
t_{MF} and perturbation showed independent effects on the progression of error across trials in block.

However, the similarity of these effects independently on error across trials within a block did
not extend to their interaction. Indeed, most relevant to testing Hypothesis-1b is that the interaction of
perturbation and t_{MF} served to speed up the growth of error, and QE slowed it down across the block.
Coefficients for perturbation $\times$ trial$_{block}$ $\times$ t_{MF} indicated that in the face of perturbation, multifractal
nonlinearity removed the observed peak in error due to earlier negative quadratic effects ($b = 3.83\times10^{-}$

^{1}, *s.e.m.* $= 5.28\times10^{-2}$, $p < 0.0001$) and increased error across the linear ($b = 4.07\times10^{-1}$, *s.e.m.* $=$
5.20×10^{-2}, $p < 0.0001$)and cubic terms ($b = 5.39\times10^{-1}$, *s.e.m.* $= 4.45\times10^{-2}$, $p < 0.0001$). On the other
hand, coefficients for perturbation $\times$ trial$_{block}$ $\times$ t_{MF} indicated that, in the face of perturbation for the QE
participants, the linear increase was shallower ($b = -2.71\times10^{-1}$, *s.e.m.* $= 6.11\times10^{-2}$, $p < 0.0001$), the
positive quadratic growth further removed the observed peak in error ($b = 4.24\times10^{-1}$, *s.e.m.* $= 6.14\times10^{-}$
^{2}, $p < 0.0001$), and the cubic term was negative ($b = -7.90\times10^{-1}$, *s.e.m.* $= 5.79\times10^{-2}$, $p < 0.0001$).

The outcome of these differences distinguishes how multifractal nonlinearity predicted changes
in adaptive responses to the perturbation of eye height. Greater t_{MF} exhibited less adaptive response to
the perturbation in the QE case (i.e., initially greater error and a smaller change in error across trials),
and less-but-still-nonzero t_{MF} predicted a more gradual appearance of error and then a subsequent
decay of this error (Fig. 5*b*), suggesting better adaptation. It is evident in figure 5*b* that although the
error in the high-t_{MF} cases with the perturbation was smaller than the delayed increase in the low-t_{MF}
cases with the perturbation, the high-t_{MF} cases showed more sustained error and no adaptation to the
perturbation relative to high-t_{MF} cases without the perturbation. This result supports Hypothesis-1b.

We include example plots of model predictions for individual participants (Fig. 6).

3.2. Logistic regression of 'makes' vs. misses showed less sensitivity to effects

Poisson regression may be too ornate for some readers. However, in contrast to Poisson modeling of a
continuous outcome measure of performance, the investment of logistic regression to model
dichotomous variable limits the number of valid predictors (e.g., van Smeden et al., 2016) and so
affords us less statistical power with which to test for changes in the performance. A mixed-effect
logistic regression model of successful putting (i.e., the 'makes') returned the coefficients reported in
Table 3. Predictors for t_{MF} and for interactions of QE with all terms other than perturbation failed to
improve model fit significantly and so were omitted from the final model. This model failed to return
an effect of QE ($b = 0.20$, *s.e.m.* $= 0.30$, $p = 0.51$), of the perturbation alone ($b = -0.22$, *s.e.m.* $= 0.43$, p
$= 0.60$) or the interaction of QE $\times$ perturbation ($b = -1.16$, *s.e.m.* $= 0.65$, $p = 0.07$).

We found effects of experience and trial-by-trial sway. Block showed a positive effect ($b =$
0.24, *s.e.m.* $= 0.13$, $p = 0.06$), suggesting that odds of a successful putt was 1.27 times higher with each
successive block of trials. RMS_{acc} showed a positive effect ($b = 4.05$, *s.e.m.* $= 1.34$, $p < 0.01$), and total
multifractality W_{MF} showed a negative effect ($b = -7.25$, *s.e.m.* $= 2.94$, $p < 0.05$), suggesting that
successful putts were more likely with more variable and less multifractal accelerations of sway. These
effects resembled some of those found in the Poisson regression model, but it appears that the
continuous measure of radial error afforded more sensitivity within which to model significant fixed
effects.

4. Discussion

We tested the effects of postural stability and its endogenous multifractal fluctuations on a continuous
outcome of putting performance (radial error) with and without QE training. We hypothesized that QE-
trained participants' response to an eye height perturbation would depend on the multifractality
attributable to nonlinear interactions across scales, t_{MF}. That is, more multifractal fluctuations in
postural sway would make eye height perturbation a greater threat to QE-trained performance but not
to technically trained performance. Specifically, we predicted that lower estimates of t_{MF} would
coincide with a more adaptive response, i.e., with slower growth of radial error and subsequent decay
of radial error over trials. Results supported this hypothesis: when we perturbed eye height, accuracy
with QE was associated with reducing multifractal nonlinearity in postural sway. This model included
a negative main effect of QE training, indicating more accurate putts in radial error. However, despite

the absence of a simple interaction of QE with clogs, QE's contingency on multifractal aspects of
postural sway entailed a greater decrement in accuracy for QE-trained participants following eye height
perturbation.

A limitation of these results is that our focus on the postural support of QE-trained performance
came at the expense of focusing on gaze tracking. We used QE training as a manipulation, but we
failed to include a manipulation check. The absence here of gaze-tracking data leaves us unable to
explain with complete confidence why a logistic model of putting performance construed
dichotomously (e.g., as 'makes' vs. 'misses') failed to find a significant effect of QE. On the one hand,
some participants may have followed QE instruction incompletely. On the other hand, a good reason
why the main effect may not appear is the brevity of our design: whereas recent analogues of our
research design already in the literature required participants to conduct 4000 putts over 7 days (Moore
et al., 2012), we only sampled 40 putts in one hour-long session. Radial error may be sufficiently
sensitive to show QE's reliable effect in a more compact design than dichotomous measures. Future
research would no doubt benefit from incorporating gaze-tracking into similarly compact experimental
designs.

These results offer two major insights for research into perception-action in visual aiming tasks.
First, it shows that QE training has roots in postural sway, vindicating and elaborating earlier proposals
that quiet eye is not simply about stabilizing the eyes (Liphart et al., 2015; Williams, 2016). Indeed, it
has long been known that simply stabilizing the retinal position compromises the persistence of a
visual image (Barlow, 1963; Coppola & Purves, 1996; Ditchburn & Ginsborg, 1952). And contrary to a
colloquial understanding of 'quiet eye,' more recent work confirms that those small movements within
gaze fixations called 'microsaccades' might prevent fading (Martinez-Conde et al., 2006) and support
visual attentional processes (Krueger et al., 2019). We might draw a comparison between the 'quiet'
suggested by 'quiet eye' and that indicated by 'quiet standing.' To be precise, we mean to draw the
comparison of 'quiet eye' specifically only to 'quiet standing' in the case of instructing participants to
'stand as still as possible' (i.e., rather than to 'stand as relaxed as possible'; (Houdijk et al., 2015). Both
'quietudes' provide the needed instruction to a participant or athlete to 'please move as little as
possible,' but below the polite clarity for verbal instruction, both 'quietudes' depend upon a rich texture
of fluctuations. Throughout the body, fluctuations support exploration during the 'still as possible'
quiet standing (I.-C. Lee et al., 2019; Murnaghan et al., 2016; Palatinus et al., 2013, 2014).

Here, the theoretical ground may shift underfoot as we try to see a path through these
considerations. Do we mean to say that the achievement of a precision demand could be exploratory?
At first blush, it could sound absurd that something not moving is also exploring. However, here, we
may hit upon a deeper divergence between ecological vs. information-processing perspectives on QE.
To the point, it is true that fixing the gaze and standing as still as possible are themselves
accomplishments of motor control. However, this achievement does not entail that either quiet eye or
quiet stance are the results. Unless we are much mistaken, it is the external focus and the intention of
getting the ball into the net or hole or target that warrants quiet eye in the first place. That is to say, the
quiet eye is a means to an end, and the end is never just quiet eye.

Precision might invite the appearance of achieving fixity, but achieving fixity might be no
guarantee against exploration. For a contrary example, 'static holding' is a haptic case of holding an
object 'as still as possible,' and it is widely recognized as a valid exploratory procedure for haptic
perception (Burton & Turvey, 1990; Carello et al., 1992; Lederman et al., 1996). The only debates are
what kind of information this apparent-fixity mode of exploration has access to. Crucially, apparent
fixity is not necessarily actual fixity, and while there is no need for the body to explore the quiet-eye
achievement of ocular fixity, it would strike us as a strange eventuality to rule quiet-eye strategy out as
a strategy pertaining to exploration. Quiet eye resembles to us an ocular version of static holding that

holds the visual image as close as possible. As noted above, despite their apparent fixity, fixations are a
class of eye movements, and these fixational eye movements are exactly what the quiet-eye strategy
uses to build a motor program. Certainly, if the QE strategy is collecting information that supports the
construction of a motor program, ruling quiet eye out as a form of exploration would strike us as a
problematic move. For instance, consider the pioneering astronomers gazing up at the heavens, aiming
a telescope as quietly as possible at a star or planet. Can we tell them that, no, because they are engaged
in precision work, they are not exploring? We suspect not. Again, to summarize this point: a precision
demand can change motor behavior (Balasubramaniam et al., 2000), but that change is not itself the
end of the story. Despite experimenter intentions to train it, quiet eye is not a goal in itself but only a
strategy for informing the participant's accurate behavior beyond their bodies for precision wherever
they direct their external focus.

Of course, the preceding proposal that QE is exploratory remains conjecture that reaches
beyond the current evidence. Demonstrating an exploratory structure in QE will need empirical
evidence from future research to back up the preceding metaphor and analogy. Specifically, empirical
validation would require at least one of the following: (1) eye movement data indicating exploratory
movements within the fixation time that are similar or relational to those from the haptic or postural
research, (2) performance outcome data that indicates information is obtained that is of value during
the quiet eye fixation period, and/or (3) self-reporting of the subjects (perhaps quantified through
perceptual judgments) indicating information gained following QE training that differs from those who
do not go through the training. We look forward to future research to the opportunity to test these ideas
and revisit the question of whether QE is exploratory behavior.

We may summarize this first insight as follows: The value of QE is not necessarily a net
reduction of fluctuations but rather a way to train athletes to orient their movements towards the optic
flow generated by postural sway. Whereas the postural-kinematic hypothesis had previously only
referred to longer movement duration (Gallicchio & Ring, 2020), we now report that movements are
not merely slower but evince a sort of nonlinearity constricting the degrees of freedom across scales of
the movement-system hierarchy. In one of the subsections below, we discuss how this orienting
towards optic flow might coordinate with mechanisms invoked by the more elaborate visual hypothesis
(e.g., Gallicchio et al., 2016, 2017).

The second insight from the present results is that the multifractality in sway is an essential
aspect of postural adaptations contributing to visually-guided action. It was already known that
damping out postural perturbations could occasion a ratcheting down of multifractality in postural
sway (Kelty-Stephen, 2018). The present results indicate that this reduction of multifractality is
associated with more accurate responses in visually-guided aiming. Note that this reduced
multifractality remains different from corresponding surrogates. Hence, nonlinear interactions across
scales can constrict movement variability, making reduced multifractality more adaptive. This
distinction is a departure from any shorthand presumption that 'more multifractal is better' supported
by early studies of heart rate variability (HRV; Ivanov et al., 2001). Indeed, this departure reflects a
growing clarity about the value of multifractal fluctuations for predicting and explaining outcomes in
perception-action.

4.1. Clarifying the value of multifractal fluctuations for perceiving-acting systems

The shorthand 'more multifractal is better' presumptions from early studies on HRV (Ivanov et al.,
2001) have always underestimated the more elaborate systematicity in the physiology of the human
movement system with upright posture. Early work proposed that a 'loss of [fractal or multifractal]
complexity' would coincide with aging or with pathology (Lipsitz & Goldberger, 1992). Indeed, early
evidence showed that less healthy outcomes co-occurred with more-than-fractal levels of temporal
correlation (Lipsitz, 2002). However, the task has always been known to influence multifractality in

upright posture: multifractality of quiet standing can reduce with pathology (Morales & Kolaczyk, 2002; Shimizu et al., 2002), but multifractality in gait can also increase with pathology (Muñoz-Diosdado, 2005), as well as with walking speeds faster or slower than self-selected comfortable speed (West & Scafetta, 2003). Over the lifespan, the maturation of gait involves loss of multifractality, beginning with the stabilization of toddler gait into young adult gait and continuing into older age (Ashkenazy et al., 2002). And this lifelong progression of multifractality in gait actually is at odds with more recent evidence that multifractality in HRV remains stable with healthy aging (Schmitt & Ivanov, 2007).

Militating strongly against simple 'more multifractal is better' kind of presumptions are a variety of distinctions. Some of these distinctions may sound patently obvious but need recognition if we want to identify for what, after all, multifractality is good. These distinctions include, for instance, that hearts are not upright bipedal bodies, that age is not itself a pathology, and that walking is not standing. The remaining distinctions are more subtle. In the promise of fractal/multifractal analysis, the discourse about 'complexity,' 'dynamical stability,' and the role of fractality/multifractality in 'optimality' (Friston et al., 2012; Oftadeh et al., 2014; Schlesinger & Klafter, 1986) has made no claims on all definitions of optimality and so has not been responsible for acknowledging that some multifractality is not optimal (Reynolds, 2015). Indeed, studies in behavioral sciences have always justified the capacity of fractal variety to offer new insights (Diniz et al., 2011; Dixon et al., 2012). Any notion of a privileged level or direction of fractal variety (Kello, 2018) might be best understood as a reaction to critical, unreceptive perspectives resistant to seeing any fractality or multifractality appear in behavioral-science discourse at all (Wagenmakers et al., 2012). Measurements may wax and wane in their multifractal structure. The only case in which we think more multifractality is always better than less is in its inclusion as a construct in the scientific discourse on the coordination of perceiving-acting systems.

4.2. The value of multifractal fluctuation is specific to task constraints

What is coming into focus is that the value of nonlinearity-driven multifractality depends on task context. More multifractal nonlinearity is adaptive when tasks invite exploration and anticipation. Less multifractal nonlinearity is adaptive when the task involves constraining behavior, drawing degrees of freedom inward around a base of support. As an example of the former, accumulating information as a reader can require keeping an open mind to follow the discourse wherever it may lead, and more multifractality t_{MF} in the pacing of word-by-word reading supports more fluent reading if the story has a twist in its plot (Booth et al., 2018). Similarly, more multifractality W_{MF} in circle-tracing behaviors make for poor precision of tracing but is associated with better performance on neurophysiological tests of flexibility with rule switching (Kelty-Stephen et al., 2016). The freewheeling benefits of more multifractality can also compete alongside the constraint-, repetition-promoting benefit of less multifractality. In a Fitts task, greater multifractality t_{MF} in hand and head movements predicted less stable contact with the targets, but it became a predictor of more stable contact when participants had the diffuse warning that they might be asked to close their eyes and continue the task (Bell et al., 2019). Hence, what would have upset task performance became a resource for exploring the task space for the eventuality of a major loss of sensory information. And now, we see that a less multifractal postural system can be more effective (e.g., smaller t_{MF} in Kelty-Stephen (2018); and smaller W_{MF} in Koslucher et al. (2016) and Li et al. (2020)) and, as present findings indicate, more accurate in how it orients behaviors to a visually-guided aiming task.

Some of the task-specificity remains unclear, e.g., manual wielding of an object is more accurate with greater multifractality in postural sway, but hefting an object to perceive heaviness compared to a reference object is more accurate with less multifractality (Mangalam & Kelty-Stephen, 2020). This difference could have to do with the qualities of length and heaviness, but the simpler

interpretation could be that intending to perceive length directs attention away from the center of
pressure and intending to perceive heaviness directs attention inwards toward the center of pressure,
lending length perceptions and heaviness perception to less and more postural constraint, respectively.
This issue of directing attention is again a reason we will need to consider how to link this postural-
kinematic work on QE back with the visual-hypothesis work on QE.

The fact that reduced multifractality can be beneficial offers a unique distinction between
possibly two related but distinct interpretations of multifractal results: one as a biomarker of pathology
and the other as an adaptation. Indeed, the loss of multifractality can be a biomarker for diagnosing a
pathology (Meyer et al., 2003). The physiological wisdom informing this usage is that the heart has to
be, in effect, poised to absorb and rebound from perturbations, avoiding fragility due to insult at any
single characteristic scale (e.g., Lipsitz & Goldberger, 1992). Implicit in this wisdom is that, within the
healthy body, the heart's task is deeply anticipatory and exploratory, much like those task settings in
which whole organisms benefit from more multifractality. However, concluding that loss of
multifractality is strictly symptomatic of postural systems with deficits (Shimizu et al., 2002) may be
premature. What looks like a deficit may sometimes be a creative constraint, built and used to solve a
task problem (Harrison & Stergiou, 2015; Holt et al., 2010; Vaz et al., 2017). The present results
indicate that reduced multifractality could be an adaptive response preserving stability and promoting
task performance in the face of abrupt change in the organism's mechanical relationship with the base
of support. Hence, far from being the signature of diseased posture, reductions in multifractality that
maintain a significant t_{MF} may be an advantageous loss of multifractality. Though more commonly
known for their capacity to amplify variability, nonlinear interactions across scales acting context-
sensitively may also restrain motoric degrees of freedom so as to achieve the task goal.

Different task constraints prompt different modes of movement variability, and if multifractal
depictions of the body seem to physicalize or mechanize the movement system, they do so only in the
way that nonlinear-dynamical physical mechanisms might better explain context-sensitive behavior.
Surely, good-faith pursuit into nonlinear-dynamical complexity should avoid any catch-all
simplification (e.g., 'more is better, less is worse') and keep a critical eye on the reference frames of
nonlinear-dynamical results. Nonlinear dynamics has long profited from considering its measures in
different reference frames: experimentally breaking apart the coincidence of relative-phase in spatial
frames from relative-phase in muscular frames revealed novel, more generic structure in the Haken-
Kelso-Bunz (1985) law for multi-limb coordination (Park & Turvey, 2008). Acknowledging the task-
based reference frames for our interpretation of multifractal estimates may be equally crucial for
generalizing predictions for a multifractal foundation for perception-action. Low multifractality in the
heart signals a bad prognosis to the clinician when the task is cardiovascular circulation, but low
multifractality in posture supports good task performance in visually-guided aiming, particularly for an
organism beset by malevolent experimenters uprooting their optic flow.

4.3. The value of multifractality is also specific to different tissues of the body.

Just as different tissues of the same body can differently support the same task, we expect that the
multifractal structure spanning these different tissues supports overarching integrity to the organismal-
level behavior differently (Stoffregen et al., 2017). Here, comparing present results with similar work
using a relatively less-mechanical, visual perturbation (Carver et al., 2017) may tee up considerations
for future work that could elaborate an explanation of QE by integrating the postural-kinematic
hypothesis with the visual hypothesis. The present results resemble those of Carver et al. (2017) who
found that, without any training, more multifractality in head sway led to greater error in a low-
difficulty Fitts task (compare W_{MF} in Table 2) but led to smaller error in the case of a visual
perturbation (compare perturbation $\times W_{MF}$ in Table 2). So, multifractality in head sway became a
stabilizing, accuracy-promoting asset to task performance when the target was far out of reach. Future

research could probe the visual hypothesis for QE for multifractal foundations, and based on Carver et al.'s (2017) results, we predict that QE-trained performance would improve with greater multifractality in head sway or eye movements. Monofractal analysis of eye movements has successfully predicted visual attention to text (Wallot et al., 2015), and multifractality of gaze fluctuations may support an extension of the multifractal support for QE. The present results do not include gaze data, but we hope it is informative, especially in support of the postural-kinematic hypothesis, that the QE instructions enlist the nonlinear interactions across scales of the movement system without drawing the athlete's attention to the body.

Ultimately, this task- and anatomically-specific portrait of multifractal fluctuations rests on two even more generic facts. First, multifractal fluctuations spread across the body, and second, estimates of this spread predict the accuracy of the perceptual outcomes (Mangalam, Carver, et al., 2020a, 2020b). Hence, rather than inventorying body parts and task constraints prompting different values of multifractality, the longer view of investigating visually-guided aiming must examine the flow of multifractal fluctuations. Carver et al. (2017) found that multifractality spread from hip to head, as well as from head and hip toward the throwing hand. Indeed, in all these studies, the body appears to direct multifractal fluctuations towards those body parts most clearly engaged in the focal tasks. We might expect similar results in golf putting. QE training may promote this spread by reducing intentional movements of the head and eyes that might interfere with upstream flows of multifractality. The direction of attention towards a distant target in the visual field may depend on stabilizing the postural grasp of the surface underfoot and, perhaps, shunting multifractal fluctuations toward the head and eyes. Indeed, executive functions directing attention to specific features of the visual field benefits from a rich substrate of multifractal fluctuations in the head (Kelty-Stephen & Dixon, 2014), hand (Anastas et al., 2014; Stephen et al., 2012), and brain (Kardan, Adam, et al., 2020; Kardan, Layden, et al., 2020) that vary systematically with access to and mastery of the rules supporting task performance. When we consider that eye movements show multifractality (Kelty-Stephen & Mirman, 2013) and that variability in the monofractal structure of eye movements can predict visual attention (Wallot et al., 2015), we see early glimpses of a multifractal bridge from the postural-kinematic support to the visual support for QE.

The novelty of multifractal modeling to behavioral sciences can make this promise of a multifractal infrastructure supporting visually guided behaviors sound too ethereal and misty to be practical. But we can root this proposal in the known physiology spanning the very same bones, muscles, bones, joints, brain, and eyes whose coordination generates the winning golf putt. Specifically, the connective tissues composing the fascia operate to coordinate context-sensitive behavior across so wide a variety of scales (e.g. (Ingber, 2006)) that multifractality is one of the few current compelling frameworks for modeling how it might support perception-action (Schleip et al., 2014; Turvey & Fonseca, 2014). The flow of multifractal fluctuations through the body could inform future explanations of how QE training and other visual aiming tasks depend on this bodywide network of connective tissues.

References

Allison, P. D. (1977). Testing for interaction in multiple regression. *American Journal of Sociology*, *83*(1), 144–153. https://doi.org/10.1086/226510

Anastas, J. R., Kelty-Stephen, D. G., & Dixon, J. A. (2014). Executive function as an interaction-dominant process. *Ecological Psychology*, *26*(4), 262–282. https://doi.org/10.1080/10407413.2014.957985

Ashkenazy, Y., M. Hausdorff, J., Ch. Ivanov, P., & Eugene Stanley, H. (2002). A stochastic model of human gait dynamics. *Physica A: Statistical Mechanics and Its Applications*, *316*(1), 662–670. https://doi.org/10.1016/S0378-4371(02)01453-X

Balasubramaniam, R., Riley, M. A., & Turvey, M. T. (2000). Specificity of postural sway to the demands of a precision task. *Gait & Posture*, *11*(1), 12–24. https://doi.org/10.1016/S0966-6362(99)00051-X

Barlow, H. B. (1963). Slippage of contact lenses and other artefacts in relation to fading and regeneration of supposedly stable retinal images. *Quarterly Journal of Experimental Psychology*, *15*(1), 36–51. https://doi.org/10.1080/17470216308416550

Bates, D., Sarkar, D., Bates, M., & Matrix, L. (2007). *The lme4 package* (R package version 2).

Bell, C., Carver, N., Zbaracki, J., & Kelty-Stephen, D. (2019). Nonlinear amplification of variability through interaction across scales supports greater accuracy in manual aiming: Evidence from a multifractal analysis with comparisons to linear surrogates in the Fitts task. *Frontiers in Physiology*, *10*, 998. https://doi.org/10.3389/fphys.2019.00998

Bernstein, N. A. (1967). *The Co-ordination and Regulation of Movement*. Pergamon Press.

Booth, C. R., Brown, H. L., Eason, E. G., Wallot, S., & Kelty-Stephen, D. G. (2018). Expectations on hierarchical scales of discourse: Multifractality predicts both short- and long-range effects of violating gender expectations in text reading. *Discourse Processes*, *55*(1), 12–30. https://doi.org/10.1080/0163853X.2016.1197811

Burton, G., & Turvey, M. T. (1990). Perceiving the lengths of rods that are held but not wielded. *Ecological Psychology*, *2*(4), 295–324. https://doi.org/10.1207/s15326969eco0204_1

Carello, C., Fitzpatrick, P., Domaniewicz, I., Chan, T.-C., & Turvey, M. T. (1992). Effortful touch with minimal movement. *Journal of Experimental Psychology: Human Perception and Performance*, *18*(1), 290–302. https://doi.org/10.1037/0096-1523.18.1.290

Carver, N. S., Bojovic, D., & Kelty-Stephen, D. G. (2017). Multifractal foundations of visually-guided aiming and adaptation to prismatic perturbation. *Human Movement Science*, *55*, 61–72. https://doi.org/10.1016/j.humov.2017.07.005

Chen, L.-C., Metcalfe, J. S., Chang, T.-Y., Jeka, J. J., & Clark, J. E. (2008). The development of infant upright posture: Sway less or sway differently? *Experimental Brain Research*, *186*(2), 293–303. https://doi.org/10.1007/s00221-007-1236-1

Chhabra, A., & Jensen, R. V. (1989). Direct determination of the $f(\alpha)$ singularity spectrum. *Physical Review Letters*, *62*(12), 1327–1330. https://doi.org/10.1103/PhysRevLett.62.1327

Cooke, A., Kavussanu, M., McIntyre, D., & Ring, C. (2010). Psychological, muscular and kinematic factors mediate performance under pressure. *Psychophysiology*, *47*(6), 1109–1118. https://doi.org/10.1111/j.1469-8986.2010.01021.x

Coppola, D., & Purves, D. (1996). The extraordinarily rapid disappearance of entopic images. *Proceedings of the National Academy of Sciences*, *93*(15), 8001–8004. https://doi.org/10.1073/pnas.93.15.8001

Davids, K., & Araújo, D. (2016). What could an ecological dynamics rationale offer Quiet Eye research? Comment on Vickers. *Current Issues in Sport Science (CISS)*, *1*, 104. https://doi.org/10.15203/CISS_2016.100

de Oliveira, R. F. (2016). Visual perception in expert action. In M Raab, B. Lobinger, S. Hoffman, A. Pizzera, & S. Laborde (Eds.), *Performance Psychology: Perception, Aaction, Cognition, Emotion* (pp. 254–270). Academic Press.

Diniz, A., Wijnants, M. L., Torre, K., Barreiros, J., Crato, N., Bosman, A. M. T., Hasselman, F., Cox, R. F. A., Van Orden, G. C., & Delignières, D. (2011). Contemporary theories of 1/f noise in motor control. *Human Movement Science*, *30*(5), 889–905. https://doi.org/10.1016/j.humov.2010.07.006

Ditchburn, R. W., & Ginsborg, B. L. (1952). Vision with a stabilized retinal image. *Nature*, *170*(4314), 36–37. https://doi.org/10.1038/170036a0

Dixon, J. A., Holden, J. G., Mirman, D., & Stephen, D. G. (2012). Multifractal dynamics in the emergence of cognitive structure. *Topics in Cognitive Science*, *4*(1), 51–62. https://doi.org/10.1111/j.1756-8765.2011.01162.x

Doyon, J. K., Hajnal, A., Surber, T., Clark, J. D., & Kelty-Stephen, D. G. (2019). Multifractality of posture modulates multisensory perception of stand-on-ability. *PLOS ONE*, *14*(2), e0212220. https://doi.org/10.1371/journal.pone.0212220

Feldman, A. G. (2018). Indirect, referent control of motor actions underlies directional tuning of neurons. *Journal of Neurophysiology*, *121*(3), 823–841. https://doi.org/10.1152/jn.00575.2018

Feldman, A. G., & Zhang, L. (2020). Eye and head movements and vestibulo-ocular reflex in the context of indirect, referent control of motor actions. *Journal of Neurophysiology*, *124*(1), 115–133. https://doi.org/10.1152/jn.00076.2020

Frenz, H., Bremmer, F., & Lappe, M. (2003). Discrimination of travel distances from 'situated' optic flow. *Vision Research*, *43*(20), 2173–2183. https://doi.org/10.1016/S0042-6989(03)00337-7

Friston, K., Breakspear, M., & Deco, G. (2012). Perception and self-organized instability. *Frontiers in Computational Neuroscience*, *6*, 44. https://doi.org/10.3389/fncom.2012.00044

Gallicchio, G., Cooke, A., & Ring, C. (2016). Lower left temporal-frontal connectivity characterizes expert and accurate performance: High-alpha T7-Fz connectivity as a marker of conscious processing during movement. *Sport, Exercise, and Performance Psychology*, *5*(1), 14–24. https://doi.org/10.1037/spy0000055

Gallicchio, G., Cooke, A., & Ring, C. (2017). Practice makes efficient: Cortical alpha oscillations are associated with improved golf putting performance. *Sport, Exercise, and Performance Psychology*, *6*(1), 89–102. https://doi.org/10.1037/spy0000077

Gallicchio, G., & Ring, C. (2020). The quiet eye effect: A test of the visual and postural-kinematic hypotheses. *Sport, Exercise, and Performance Psychology*, *9*(1), 143–159. https://doi.org/10.1037/spy0000162

Gibson, J. J. (1979). *The Ecological Approach to Visual Perception*. Houghton Mifflin.

Hajnal, A., Bunch, D. A., & Kelty-Stephen, D. G. (2014). Going for distance and going for speed: Effort and optical variables shape information for distance perception from observation to response. *Attention, Perception, & Psychophysics*, *76*(4), 1015–1035. https://doi.org/10.3758/s13414-014-0629-z

Hajnal, A., Bunch, D. A., & Kelty-Stephen, D. G. (2016). Pulling out all the stops to make the distance: Effects of effort and optical information in distance perception responses made by rope pulling. *Attention, Perception, & Psychophysics*, *78*(2), 685–699. https://doi.org/10.3758/s13414-015-1035-x

Hajnal, A., Clark, J. D., Doyon, J. K., & Kelty-Stephen, D. G. (2018). Fractality of body movements predicts perception of affordances: Evidence from stand-on-ability judgments about slopes. *Journal of Experimental Psychology: Human Perception and Performance*, *44*(6), 836–841. https://doi.org/10.1037/xhp0000510

Haken, H., Kelso, J. A. S., & Bunz, H. (1985). A theoretical model of phase transitions in human hand movements. *Biological Cybernetics*, *51*(5), 347–356. https://doi.org/10.1007/BF00336922

Harrison, S. J., & Stergiou, N. (2015). Complex adaptive behavior and dexterous action. *Nonlinear Dynamics, Psychology, and Life Sciences*, *19*(4), 345–394. https://pubmed.ncbi.nlm.nih.gov/26375932

Holt, K. G., Wagenaar, R. O., & Saltzman, E. (2010). A dynamic systems: Constraints approach to rehabilitation. *Brazilian Journal of Physical Therapy*, *14*, 446–463. https://doi.org/10.1590/S1413-35552010000600002

Houdijk, H., Brown, S. E., & van Dieën, J. H. (2015). Relation between postural sway magnitude and metabolic energy cost during upright standing on a compliant surface. *Journal of Applied Physiology*, *119*(6), 696–703. https://doi.org/10.1152/japplphysiol.00907.2014

Ihlen, E. A. F., & Vereijken, B. (2010). Interaction-dominant dynamics in human cognition: Beyond 1/*f*
fluctuation. *Journal of Experimental Psychology: General*, *139*(3), 436–463.
https://doi.org/10.1037/a0019098

Ingber, D. E. (2006). Cellular mechanotransduction: Putting all the pieces together again. *The FASEB*
Journal, *20*(7), 811–827. https://doi.org/10.1096/fj.05-5424rev

Ivanov, P. C., Nunes Amaral, L. A., Goldberger, A. L., Havlin, S., Rosenblum, M. G., Stanley, H. E.,
& Struzik, Z. R. (2001). From 1/*f* noise to multifractal cascades in heartbeat dynamics. *Chaos: An*
Interdisciplinary Journal of Nonlinear Science, *11*(3), 641–652.
https://doi.org/10.1063/1.1395631

Kardan, O., Adam, K. C. S., Mance, I., Churchill, N. W., Vogel, E. K., & Berman, M. G. (2020).
Distinguishing cognitive effort and working memory load using scale-invariance and alpha
suppression in EEG. *NeuroImage*, *211*, 116622.
https://doi.org/10.1016/j.neuroimage.2020.116622

Kardan, O., Layden, E., Choe, K. W., Lyu, M., Zhang, X., Beilock, S. L., Rosenberg, M. D., &
Berman, M. G. (2020). Scale-invariance in brain activity predicts practice effects in cognitive
performance. *BioRxiv*, 114959. https://doi.org/10.1101/2020.05.25.114959

Kello, C. T. (2018). Editor's introduction and review: Coordination and context in cognitive science.
Topics in Cognitive Science, *10*(1), 6–17. https://doi.org/10.1111/tops.12302

Kelty-Stephen, D. G. (2018). Multifractal evidence of nonlinear interactions stabilizing posture for
phasmids in windy conditions: A reanalysis of insect postural-sway data. *PLOS ONE*, *13*(8),
e0202367. https://doi.org/10.1371/journal.pone.0202367

Kelty-Stephen, D. G., & Dixon, J. A. (2014). Interwoven fluctuations during intermodal perception:
Fractality in head sway supports the use of visual feedback in haptic perceptual judgments by
manual wielding. *Journal of Experimental Psychology: Human Perception and Performance*,
40(6), 2289–2309. https://doi.org/10.1037/a0038159

Kelty-Stephen, D. G., Lee, I.-C., Carver, N. S., Newell, K. M., & Mangalam, M. (2020). Visual effort
moderates a self-correcting nonlinear postural control policy. *BioRxiv*, 209502.
https://doi.org/10.1101/2020.07.17.209502

Kelty-Stephen, D. G., & Mirman, D. (2013). Gaze fluctuations are not additively decomposable: Reply
to Bogartz and Staub. *Cognition*, *126*(1), 128–134.
https://doi.org/10.1016/j.cognition.2012.09.002

Kelty-Stephen, D. G., Stirling, L. A., & Lipsitz, L. A. (2016). Multifractal temporal correlations in
circle-tracing behaviors are associated with the executive function of rule-switching assessed by
the Trail Making Test. *Psychological Assessment*, *28*(2), 171–180.
https://doi.org/10.1037/pas0000177

Kelty-Stephen, D. G., & Wallot, S. (2017). Multifractality versus (mono-) fractality as evidence of
nonlinear interactions across timescales: Disentangling the belief in nonlinearity from the
diagnosis of nonlinearity in empirical data. *Ecological Psychology*, *29*(4), 259–299.
https://doi.org/10.1080/10407413.2017.1368355

Koslucher, F., Munafo, J., & Stoffregen, T. A. (2016). Postural sway in men and women during
nauseogenic motion of the illuminated environment. *Experimental Brain Research*, *234*(9), 2709–
2720. https://doi.org/10.1007/s00221-016-4675-8

Krueger, E., Schneider, A., Chavaillaz, A., Groner, R., Sawyer, B. D., Sonderegger, A., & Hancock, P.
A. (2019). Microsaccades distinguish looking from seeing. *Journal of Eye Movement Research*,
12(6), 2.

Latash, M. L. (2012). The bliss (not the problem) of motor abundance (not redundancy). *Experimental
Brain Research*, *217*(1), 1–5. https://doi.org/10.1007/s00221-012-3000-4

Latash, M. L. (2020). On primitives in motor control. *Motor Control*, *24*(2), 318–346.
https://doi.org/10.1123/mc.2019-0099

Lederman, S. J., Ganeshan, S. R., & Ellis, R. E. (1996). Effortful touch with minimum movement:
Revisited. *Journal of Experimental Psychology: Human Perception and Performance*, *22*(4), 851–
868. https://doi.org/10.1037/0096-1523.22.4.851

Lee, I.-C., Pacheco, M. M., & Newell, K. M. (2019). The precision demands of viewing distance
modulate postural coordination and control. *Human Movement Science*, *66*, 425–439.
https://doi.org/10.1016/j.humov.2019.05.019

Lee, J. T., & Kelty-Stephen, D. G. (2017). Cascade-driven series with narrower multifractal spectra
than their surrogates: Standard deviation of multipliers changes interactions across scales.
Complexity, *2017*, 7015243. https://doi.org/10.1155/2017/7015243

Li, R., Walter, H. J., & Stoffregen, T. A. (2020). The role of visual feedback about motion of the
ground on postural sway. *Journal of Motor Behavior*, *52*(3), 352–359.
https://doi.org/10.1080/00222895.2019.1627281

Liphart, J., Gallichio, J., Tilson, J. K., Pei, Q., Wu, S. S., & Duncan, P. W. (2015). Concordance and
discordance between measured and perceived balance and the effect on gait speed and falls
following stroke. *Clinical Rehabilitation*, *30*(3), 294–302.
https://doi.org/10.1177/0269215515578294

Lipsitz, L. A. (2002). Dynamics of stability: The physiologic basis of functional health and frailty. *The
Journals of Gerontology: Series A*, *57*(3), B115–B125. https://doi.org/10.1093/gerona/57.3.B115

Lipsitz, L. A., & Goldberger, A. L. (1992). Loss of "complexity" and aging: Potential applications of
fractals and chaos theory to senescence. *JAMA*, *267*(13), 1806–1809.
https://doi.org/10.1001/jama.1992.03480130122036

Mangalam, M., Carver, N. S., & Kelty-Stephen, D. G. (2020a). Global broadcasting of local fractal fluctuations in a bodywide distributed system supports perception via effortful touch. *Chaos, Solitons & Fractals*, *135*, 109740. https://doi.org/10.1016/j.chaos.2020.109740

Mangalam, M., Carver, N. S., & Kelty-Stephen, D. G. (2020b). Multifractal signatures of perceptual processing on anatomical sleeves of the human body. *Journal of The Royal Society Interface*, *17*(168), 20200328. https://doi.org/10.1098/rsif.2020.0328

Mangalam, M., & Kelty-Stephen, D. G. (2020). Multiplicative-cascade dynamics supports whole-body coordination for perception via effortful touch. *Human Movement Science*, *70*, 102595. https://doi.org/10.1016/j.humov.2020.102595

Mangalam, M., Lee, I.-C., Newell, K. M., & Kelty-Stephen, D. G. (2020). Visual effort moderates postural cascade dynamics. *BioRxiv*, 209486. https://doi.org/10.1101/2020.07.17.209486

Mark, L. S. (1987). Eyeheight-scaled information about affordances: A study of sitting and stair climbing. *Journal of Experimental Psychology: Human Perception and Performance*, *13*(3), 361–370. https://doi.org/10.1037/0096-1523.13.3.361

Mark, L. S., Balliett, J. A., Craver, K. D., Douglas, S. D., & Fox, T. (1990). What an actor must do in order to perceive the affordance for sitting. *Ecological Psychology*, *2*(4), 325–366. https://doi.org/10.1207/s15326969eco0204_2

Martinez-Conde, S., Macknik, S. L., Troncoso, X. G., & Dyar, T. A. (2006). Microsaccades counteract visual fading during fixation. *Neuron*, *49*(2), 297–305. https://doi.org/10.1016/j.neuron.2005.11.033

Meyer, M., Stiedl, O., & Kerman, B. (2003). Discrimination by multifractal spectrum estimation of human heartbeat interval dynamics. *Fractals*, *11*(02), 195–204. https://doi.org/10.1142/S0218348X03002063

Moore, L. J., Vine, S. J., Cooke, A., Ring, C., & Wilson, M. R. (2012). Quiet eye training expedites motor learning and aids performance under heightened anxiety: The roles of response programming and external attention. *Psychophysiology*, *49*(7), 1005–1015. https://doi.org/10.1111/j.1469-8986.2012.01379.x

Moore, L. J., Vine, S. J., Freeman, P., & Wilson, M. R. (2013). Quiet eye training promotes challenge appraisals and aids performance under elevated anxiety. *International Journal of Sport and Exercise Psychology*, *11*(2), 169–183. https://doi.org/10.1080/1612197X.2013.773688

Morales, C. J., & Kolaczyk, E. D. (2002). Wavelet-based multifractal analysis of human balance. *Annals of Biomedical Engineering*, *30*(4), 588–597. https://doi.org/10.1114/1.1478082

Morrison, A., McGrath, D., & Wallace, E. S. (2016). Motor abundance and control structure in the golf swing. *Human Movement Science*, *46*, 129–147. https://doi.org/10.1016/j.humov.2016.01.009

Munafo, J., Curry, C., Wade, M. G., & Stoffregen, T. A. (2016). The distance of visual targets affects
the spatial magnitude and multifractal scaling of standing body sway in younger and older adults.
Experimental Brain Research, *234*(9), 2721–2730. https://doi.org/10.1007/s00221-016-4676-7

Muñoz-Diosdado, A. (2005). A non linear analysis of human gait time series based on multifractal
analysis and cross correlations. *Journal of Physics: Conference Series*, *23*, 87–95.
https://doi.org/10.1088/1742-6596/23/1/010

Murnaghan, C. D., Carpenter, M. G., Chua, R., & Inglis, J. T. (2016). Keeping still doesn't "make
sense": Examining a role for movement variability by stabilizing the arm during a postural control
task. *Journal of Neurophysiology*, *117*(2), 846–852. https://doi.org/10.1152/jn.01150.2015

Oftadeh, R., Haghpanah, B., Vella, D., Boudaoud, A., & Vaziri, A. (2014). Optimal fractal-like
hierarchical honeycombs. *Physical Review Letters*, *113*(10), 104301.
https://doi.org/10.1103/PhysRevLett.113.104301

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Palatinus, Z., Dixon, J. A., & Kelty-Stephen, D. G. (2013). Fractal fluctuations in quiet standing predict
the use of mechanical information for haptic perception. *Annals of Biomedical Engineering*, *41*(8),
1625–1634. https://doi.org/10.1007/s10439-012-0706-1

Palatinus, Z., Kelty-Stephen, D. G., Kinsella-Shaw, J., Carello, C., & Turvey, M. T. (2014). Haptic
perceptual intent in quiet standing affects multifractal scaling of postural fluctuations. *Journal of
Experimental Psychology: Human Perception and Performance*, *40*(5), 1808–1818.
https://doi.org/10.1037/a0037247

Panchuk, D., & Vickers, J. N. (2006). Gaze behaviors of goaltenders under spatial–temporal
constraints. *Human Movement Science*, *25*(6), 733–752.
https://doi.org/10.1016/j.humov.2006.07.001

Park, H., & Turvey, M. T. (2008). Imperfect symmetry and the elementary coordination law. In A.
Fuchs & V. K. Jirsa (Eds.), *Coordination: Neural, Behavioral and Social Dynamics* (pp. 3–25).
Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-540-74479-5_1

Profeta, V. L. S., & Turvey, M. T. (2018). Bernstein's levels of movement construction: A
contemporary perspective. *Human Movement Science*, *57*, 111–133.
https://doi.org/10.1016/j.humov.2017.11.013

Raja, V., & Anderson, M. L. (2019). Radical embodied cognitive neuroscience. *Ecological Psychology*,
31(3), 166–181. https://doi.org/10.1080/10407413.2019.1615213

Rajachandrakumar, R., Mann, J., Schinkel-Ivy, A., & Mansfield, A. (2018). Exploring the relationship
between stability and variability of the centre of mass and centre of pressure. *Gait & Posture*, *63*,
254–259. https://doi.org/10.1016/j.gaitpost.2018.05.008

Renshaw, I., Davids, K., Araújo, D., Lucas, A., Roberts, W. M., Newcombe, D. J., & Franks, B. (2019). Evaluating weaknesses of "perceptual-cognitive training" and "brain training" methods in sport: An ecological dynamics critique. *Frontiers in Psychology*, *9*, 2468. https://www.frontiersin.org/article/10.3389/fpsyg.2018.02468

Reynolds, A. M. (2015). Liberating Lévy walk research from the shackles of optimal foraging. *Physics of Life Reviews*, *14*, 59–83. https://doi.org/10.1016/j.plrev.2015.03.002

Rienhoff, R., Tirp, J., Strauß, B., Baker, J., & Schorer, J. (2016). The 'quiet eye' and motor performance: A systematic review based on Newell's constraints-led model. *Sports Medicine*, *46*(4), 589–603. https://doi.org/10.1007/s40279-015-0442-4

Schleip, R., Mechsner, F., Zorn, A., & Klingler, W. (2014). The bodywide fascial network as a sensory organ for haptic perception. *Journal of Motor Behavior*, *46*(3), 191–193. https://doi.org/10.1080/00222895.2014.880306

Schlesinger, M. F., & Klafter, J. (1986). Lévy walks versus Lévy flights. In H. E. Stanley & N. Ostrowski (Eds.), *Growth and Form* (Stanley, H, pp. 279–283). Martinus Nijhoff.

Schmitt, D. T., & Ivanov, P. C. (2007). Fractal scale-invariant and nonlinear properties of cardiac dynamics remain stable with advanced age: A new mechanistic picture of cardiac control in healthy elderly. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *293*(5), R1923–R1937. https://doi.org/10.1152/ajpregu.00372.2007

Schreiber, T., & Schmitz, A. (1996). Improved surrogate data for nonlinearity tests. *Physical Review Letters*, *77*(4), 635–638. https://doi.org/10.1103/PhysRevLett.77.635

Serres, J. R., & Ruffier, F. (2017). Optic flow-based collision-free strategies: From insects to robots. *Arthropod Structure & Development*, *46*(5), 703–717. https://doi.org/10.1016/j.asd.2017.06.003

Shannon, C. E. (1948). A mathematical theory of communication. *The Bell System Technical Journal*, *27*(3), 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x

Shimizu, Y. U., Thurner, S., & Ehrenberger, K. (2002). Multifractal spectra as a measure of complexity in human posture. *Fractals*, *10*(01), 103–116. https://doi.org/10.1142/S0218348X02001130

Shine, J. M., Aburn, M. J., Breakspear, M., & Poldrack, R. A. (2018). The modulation of neural gain facilitates a transition between functional segregation and integration in the brain. *ELife*, *7*, e31130. https://doi.org/10.7554/eLife.31130

Stephen, D., Anastas, J., & Dixon, J. (2012). Scaling in cognitive performance reflects multiplicative multifractal cascade dynamics. *Frontiers in Physiology*, *3*, 102. https://doi.org/10.3389/fphys.2012.00102

Stephen, D. G., & Mirman, D. (2010). Interactions dominate the dynamics of visual cognition. *Cognition*, *115*(1), 154–165. https://doi.org/10.1016/j.cognition.2009.12.010

Stoffregen, T. A. (1985). Flow structure versus retinal location in the optical control of stance. *Journal of Experimental Psychology: Human Perception and Performance*, *11*(5), 554–565. https://doi.org/10.1037/0096-1523.11.5.554

Stoffregen, T. A. (1986). The role of optical velocity in the control of stance. *Perception & Psychophysics*, *39*(5), 355–360. https://doi.org/10.3758/BF03203004

Stoffregen, T. A., Mantel, B., & Bardy, B. G. (2017). The senses considered as one perceptual system. *Ecological Psychology*, *29*(3), 165–197. https://doi.org/10.1080/10407413.2017.1331116

Stoffregen, T. A., & Riccio, G. E. (1990). Responses to optical looming in the retinal center and periphery. *Ecological Psychology*, *2*(3), 251–274. https://doi.org/10.1207/s15326969eco0203_3

Stoffregen, T. A., Yang, C.-M., & Bardy, B. G. (2005). Affordance judgments and nonlocomotor body movement. *Ecological Psychology*, *17*(2), 75–104. https://doi.org/10.1207/s15326969eco1702_2

Teng, D. W., Eddy, C. L., & Kelty-Stephen, D. G. (2016). Non-visually-guided distance perception depends on matching torso fluctuations between training and test. *Attention, Perception, & Psychophysics*, *78*(8), 2320–2328. https://doi.org/10.3758/s13414-016-1213-5

Turvey, M T. (1977). Contrasting orientations to the theory of visual information processing. *Psychological Review*, *84*(1), 67–88. https://doi.org/10.1037/0033-295X.84.1.67

Turvey, Michael T, & Fonseca, S. T. (2014). The medium of haptic perception: A tensegrity hypothesis. *Journal of Motor Behavior*, *46*(3), 143–187. https://doi.org/10.1080/00222895.2013.798252

van Smeden, M., de Groot, J. A. H., Moons, K. G. M., Collins, G. S., Altman, D. G., Eijkemans, M. J. C., & Reitsma, J. B. (2016). No rationale for 1 variable per 10 events criterion for binary logistic regression analysis. *BMC Medical Research Methodology*, *16*(1), 163. https://doi.org/10.1186/s12874-016-0267-3

Vaz, D. V., Silva, P. L., Mancini, M. C., Carello, C., & Kinsella-Shaw, J. (2017). Towards an ecologically grounded functional practice in rehabilitation. *Human Movement Science*, *52*, 117–132. https://doi.org/10.1016/j.humov.2017.01.010

Vecera, S. P., Cosman, J. D., Vatterott, D. B., & Roper, Z. J. J. (2014). The control of visual attention: Toward a unified account. In B. H. Ross (Ed.), The Psychology of Learning and Motivation (pp. 303–347). Academic Press.

Vickers, J. N. (1992). Gaze control in putting. *Perception*, *21*(1), 117–132. https://doi.org/10.1068/p210117

Vickers, J. N. (1996). Visual control when aiming at a far target. *Journal of Experimental Psychology: Human Perception and Performance*, *22*(2), 342–354. https://doi.org/10.1037/0096-1523.22.2.342

Vickers, J. N. (2009). Advances in coupling perception and action: the quiet eye as a bidirectional link between gaze, attention, and action. In Markus Raab, J. G. Johnson, & H. R. B. T.-P. in B. R.

Heekeren (Eds.), *Mind and Motion: The Bidirectional Link between Thought and Action* (Vol. 174, pp. 279–288). Elsevier. https://doi.org/10.1016/S0079-6123(09)01322-3

Vickers, J. N. (2011). Mind over muscle: The role of gaze control, spatial cognition, and the quiet eye in motor expertise. *Cognitive Processing*, *12*(3), 219–222. https://doi.org/10.1007/s10339-011-0411-2

Vine, S. J., Lee, D. H., Walters-Symons, R., & Wilson, M. R. (2017). An occlusion paradigm to assess the importance of the timing of the quiet eye fixation. *European Journal of Sport Science*, *17*(1), 85–92. https://doi.org/10.1080/17461391.2015.1073363

Vine, S. J., Lee, D. O. N., Moore, L. E. E. J., & Wilson, M. R. (2013). Quiet eye and choking: Online control breaks down at the point of performance failure. *Medicine & Science in Sports & Exercise*, *45*(10), 1988–1994. https://doi.org/10.1249/MSS.0b013e31829406c7

Vine, S., Moore, L., & Wilson, M. (2011). Quiet eye training facilitates competitive putting performance in elite golfers. *Frontiers in Psychology*, *2*, 8. https://doi.org/0.3389/fpsyg.2011.00008

Wagenmakers, E.-J., van der Maas, H. L. J., & Farrell, S. (2012). Abstract concepts require concrete models: Why cognitive scientists have not yet embraced nonlinearly coupled, dynamical, self-organized critical, synergistic, scale-free, exquisitely context-sensitive, interaction-dominant, multifractal, interdependent b. *Topics in Cognitive Science*, *4*(1), 87–93. https://doi.org/10.1111/j.1756-8765.2011.01164.x

Wallot, S., O'Brien, B., Coey, C. A., & Kelty-Stephen, D. (2015). Power-law fluctuations in eye movements predict text comprehension during connected text reading. In D. C. Noelle, R. Dale, A. S. Warlaumont, J. Yoshimi, T. Matlock, & C. D. Jennings (Eds.), *Proceedings of the 37th Annual Meeting of the Cognitive Science Society* (pp. 2583–2588).

Warren, W. H. (1984). Perceiving affordances: Visual guidance of stair climbing. *Journal of Experimental Psychology: Human Perception and Performance*, *10*(5), 683–703. https://doi.org/10.1037/0096-1523.10.5.683

West, B. J., & Scafetta, N. (2003). Nonlinear dynamical model of human gait. *Physical Review E*, *67*(5), 51917. https://doi.org/10.1103/PhysRevE.67.051917

Williams, A. M. (2016). Quiet eye vs. noisy brain: The eye like the brain is always active–comment on Vickers. *Current Issues in Sport Science (CISS)*, *1*, 116. https://doi.org/10.15203/CISS_2016.116

Witt, J. K., & Proffitt, D. R. (2008). Action-specific influences on distance perception: A role for motor simulation. *Journal of Experimental Psychology: Human Perception and Performance*, *34*(6), 1479–1492. https://doi.org/10.1037/a0010781

Table 1. Training instructions given to the quiet-eye and technical groups during the training phase.

quiet-eye (QE) training instructions	technical training instructions

assume your stance, and ensure your gaze is located on the back of the ball	take your stance with your legs shoulder-width apart
after setting up over the ball, fix your gaze on the hole	set your position so that your head is directly above the ball looking down
make no more than three fixations towards the hole	keep your clubhead square to the ball
your final fixation should be a quiet eye on the back of the ball. The onset of the quiet eye should occur before the stroke begins and last for 2 to 3s	allow your arms and shoulders to remain loose
ensure you direct gaze towards the clubhead during the putting stroke	the putting action should be pendulum-like, making sure that you accelerate through the ball
the quiet eye should remain on the green for 200 to 300ms after the club contacts the ball	after contact, follow through but keep your head still and facing down

1075   Source: Adapted from (Moore et al., 2013).

1076 **Table 2.** Subset of Poisson regression model including main and interaction effects non-multifractal
1077 and multifractal measures of sway, perturbation of eye height, and time

predictor	b	s.e.m.	p*
effects of non-multifractal measures of sway			
(see Section 3.1.1.)			
MSD_{acc}	-8.80×10^0	1.67×10^{-1}	**< 0.0001**
$Mean_{disp}$	-2.91×10^2	1.33×10^1	**< 0.0001**
SD_{disp}	4.43×10^0	1.64×10^0	**< 0.01**
$MSD_{acc} \times Mean_{disp}$	1.97×10^4	1.00×10^1	**< 0.0001**
$MSD_{acc} \times SD_{disp}$	-3.11×10^3	1.03×10^1	**< 0.0001**
$Mean_{disp} \times SD_{disp}$	2.56×10^4	1.05×10^1	**< 0.0001**
$MSD_{acc} \times Mean_{disp} \times SD_{disp}$	-1.04×10^5	1.08×10^1	**< 0.0001**
effects of multifractal measures of sway and interactions with time			
(see Section 3.1.2.)			
W_{MF}	1.80×10^0	1.29×10^{-1}	**< 0.0001**
t_{MF}	2.21×10^{-4}	1.05×10^{-3}	0.83
$W_{MF} \times t_{MF}$	-5.93×10^{-3}	8.27×10^{-3}	0.48
$t_{MF} \times trial_{block}(Linear)$	1.36×10^{-2}	9.82×10^{-3}	0.17
$t_{MF} \times trial_{block}(Quadratic)$	-8.93×10^{-2}	1.15×10^{-2}	**< 0.0001**
$t_{MF} \times trial_{block}(Cubic)$	-1.08×10^{-2}	1.14×10^{-2}	0.34
interactions of multifractal measures of sway with perturbation			
(see Section 3.1.3.)			
$QE \times W_{MF}$	-1.04×10^{-0}	2.03×10^{-1}	**< 0.0001**
$QE \times t_{MF}$	-1.85×10^{-3}	1.30×10^{-3}	0.16
$QE \times W_{MF} \times t_{MF}$	-4.59×10^{-3}	1.09×10^{-2}	0.67
$perturbation \times W_{MF}$	-1.50×10^0	3.12×10^{-1}	**< 0.0001**
$perturbation \times t_{MF}$	4.57×10^{-3}	2.85×10^{-3}	0.11
$perturbation \times W_{MF} \times t_{MF}$	1.54×10^{-1}	3.15×10^{-2}	**< 0.0001**
$QE \times perturbation \times W_{MF}$	2.12×10^0	4.91×10^{-1}	**< 0.0001**
$QE \times perturbation \times t_{MF}$	8.62×10^{-3}	4.18×10^{-3}	**< 0.05**
$QE \times perturbation \times W_{MF} \times t_{MF}$	-2.21×10^{-1}	4.97×10^{-2}	**< 0.0001**
interactions of multifractal measures with perturbation and time			
(see Section 3.1.4.)			
$QE \times trial_{block}(Linear) \times t_{MF}$	-3.73×10^{-2}	1.46×10^{-2}	**< 0.05**
$QE \times trial_{block}(Quadratic) \times t_{MF}$	-1.85×10^{-1}	1.66×10^{-2}	**< 0.0001**
$QE \times trial_{block}(Cubic) \times t_{MF}$	1.13×10^{-1}	1.73×10^{-2}	**< 0.0001**
$perturbation \times trial_{block}(Linear) \times t_{MF}$	3.83×10^{-1}	5.28×10^{-2}	**< 0.0001**
$perturbation \times trial_{block}(Quadratic) \times t_{MF}$	4.07×10^{-1}	5.20×10^{-2}	**< 0.0001**
$perturbation \times trial_{block}(Cubic) \times t_{MF}$	5.39×10^{-1}	4.45×10^{-2}	**< 0.0001**
$QE \times perturbation \times trial_{block}(Linear) \times t_{MF}$	-2.71×10^{-1}	6.11×10^{-2}	**< 0.0001**
$QE \times perturbation \times trial_{block}(Quadratic) \times t_{MF}$	5.24×10^{-1}	6.14×10^{-2}	**< 0.0001**
$QE \times perturbation \times trial_{block}(Cubic) \times t_{MF}$	-7.90×10^{-1}	5.79×10^{-2}	**< 0.0001**

1078 *boldfaced values indicate statistical significance at the alpha level of 0.05.

1079   **Table 3.** Logistic regression modeling of successful shots.

predictor	b	s.e.m.	p*
intercept	–2.85	0.48	**< 0.0001**
QE	0.20	0.30	0.51
perturbation	–0.22	0.43	0.60
QE × perturbation	–1.16	0.65	0.07
block	0.24	0.13	**0.06**
RMS_{Acc}	4.05	1.34	**< 0.01**
W_{MF}	–7.25	2.94	**< 0.05**

1080   *boldfaced values indicate statistical significance at the alpha level of 0.05.

1081   **Figure 1.** Schematic of Chhabra and Jensen's (Chhabra & Jensen, 1989) method used to estimate
1082   multifractal spectrum $f(\alpha)$ (Part-1). See Section 2.4.1. for further details.

1083   **Figure 2.** Schematic of Chhabra and Jensen's (Chhabra & Jensen, 1989) method used to estimate
1084   multifractal spectrum $f(\alpha)$ (Part-2). See Section 2.4.1. for further details.

option A: multifractal spectrum width reflects strictly linear properties of the original series

snip   snip   snip   snip

option B: multifractal spectrum width reflects effects unfloding across multiple, nested scales

samples [1/100 Hz]

shuffle data to destroy nonlinearity but preserve linearity (overall form and short-lag relationships)

flip   flip   flip   flip   flip

samples [1/100 Hz]

null hypothesis: multifractal spectrum width of original series does not differ from that of surrogate series

use one-sample *t*-test to compare original series' multifractal spectrum width to those of 30 surrogate series

if one-sample *t*-test **is not** significantly large, fail to reject null: **series is linear**

if one-sample *t*-test **is** significantly large, fail to reject null: **series exhibits cascades**

1085   **Figure 3.** Surrogate analysis used to identify whether a series exhibits multifractality due to
1086   nonlinearity (cascades). See Section 2.4.2. for further details.

Figure 4. Multifractal analysis. (*a*, *b*) Representative original accelerometer displacement series. (*c*, *d*) Representative IAAFT surrogate series of the original accelerometer displacement series in *a* and *b*. (*e*, *f*) Multifractal spectrum ($f(\alpha)$, α) of original series in *a* and *b*, and those of their five representative IAAFT surrogates.

Figure 5. Trial-by-trial model predictions of radial error for the two participant groups. (*a*) Technical-trained group. (*b*) QE-trained group.

Figure 6. The observed trial-by-trial radial error and trial-by-trial model predictions on the same axis for four participants each from the technical (top four panels) and QE-trained (bottom four panels) groups, respectively. The representation includes two participants each with the highest (left panels) and lowest (right panels) Pearson's correlation coefficient *r* indicating the strength of relationship between the observed and predicted radial errors.