Multifractality in postural sway supports quiet eye training in aiming tasks: A study of golf putting

Noah Jacobson[1], Quinn Berleman-Paul[1], Madhur Mangalam[2] and Damian G. Kelty-Stephen[1]

[1]Department of Psychology, Grinnell College, Grinnell, IA 50112, USA

[2]Department of Physical Therapy, Movement and Rehabilitation Sciences, Northeastern University, Boston, MA 02115, USA

Author for correspondence:

Damian G. Kelty-Stephen

e-mail: keltysda@grinnell.edu

Ethics statement. All participants provided verbal and written informed consent approved by the Institutional Review Board (IRB) at Grinnell College (Grinnell, IA).

Data accessibility. All data analyzed in the present study are available upon request.

Author contributions: N.J., Q.B-P., and D.G.K-S. conceived and designed research; N.J., Q.B-P., and D.G.K-S. performed experiments; N.J., Q.B-P., and D.G.K-S. analyzed data; N.J., Q.B-P., M.M., and D.G.K-S. interpreted results of experiments; M.M. and D.G.K-S. prepared figures; N.J. and D.G.K-S. drafted manuscript; N.J., Q.B-P., M.M., and D.G.K-S. edited and revised manuscript; N.J., Q.B-P., M.M., and D.G.K-S. approved final version of manuscript.

Competing interests. The authors have no competing interests to declare.

Supplementary materials

Supplementary Table S1. Subset of Poisson regression model output including main and interaction effects of time and perturbation of eye height.

Abstract

The 'quiet eye' (QE) approach to visually-guided aiming behavior invests fully in perceptual
information's potential to organize coordinated action. Sports psychologists refer to QE as the stillness
of the eyes during aiming tasks and increasingly into self- and externally-paced tasks. Amidst the
'noisy' fluctuations of the athlete's body, quiet eyes might leave fewer saccadic interruptions to the
coupling between postural sway and optic flow. Postural sway exhibits fluctuations whose multifractal
structure serves as a robust predictor of visual and haptic perceptual responses. Postural sway generates
optic flow centered on an individual's eye height. We predicted that perturbing the eye height by
attaching wooden blocks below the feet would perturb the putting more so in QE-trained participants
than those trained technically. We also predicted that QE's efficacy and responses to this perturbation
would depend on multifractality in postural sway. Specifically, we predicted that less multifractality
would predict more adaptive responses to the perturbation and higher putting accuracy. Results showed
that lower multifractality led to more frequent successful putts, and the perturbation of eye height led to
less frequent successful putts, particularly for QE-trained participants. Models of radial error (i.e., the
distance between the ball's final position and the hole) indicated that lower estimates of multifractality
due to nonlinearity coincided with a more adaptive response to the perturbation. These results suggest
that reduced multifractality may act in a context-sensitive manner to restrain motoric degrees of
freedom to achieve the task goal.

Keywords: biotensegrity, degree of freedom, fractal, posture, tensegrity, visual adaptation

1. Introduction

Aiming our behaviors into the visible world requires an ongoing awareness of how our body fits in the world around it. The difficulty of sorting out this bodily fit finds its classic encoding in the 'degrees of freedom' problem: even if a fine-grained choreography of goal-directed movement that stipulates all individual motoric degrees of freedom at fine grains were possible, this fine-grained choreography would probably be too rigid to adapt to the task or context (Bernstein, 1967). However, another perspective suggests that the abundance of degrees of freedom is not a problem but a 'blessing' (Latash, 2012). According to this view, there is no need to stipulate the infinitude of motoric variables at very fine grains, and on the contrary, task or intention organizes the movement system at the relatively coarse grain of the whole body and context (Latash, 2020). The infinitude thus becomes a guarantee of adaptive flexibility. This positive view of motor abundance appears in recent studies of aiming behaviors like the golf swing, whereby a clearly defined target in the context draws visual attention away from specific postures of the body, and indeed, experts with their eyes on the target show no single optimized mode of execution (Morrison et al., 2016). Hence, instead of requiring explicit choreography, the movement system's organization may readily emerge from perceptual linkages to the surrounding context (Stoffregen, 1985, 1986; Stoffregen & Riccio, 1990).

1.1. Quiet eye vs. technical training

Supporting task performance by emphasizing perceptual linkages vs. explicit choreography is a clear research theme on 'quiet eye' (QE). Sports psychologists refer to QE as the stillness of the eyes during aiming tasks, specifically the final gaze fixation to a single location or object in the task for a minimum of 100 ms within 3 degrees of visual angle immediately before the movement is initiated (Rienhoff et al., 2016). In the case of a golf swing, 'technical training' that communicates an explicit choreography of the arm and leg placement leads to less accuracy than a 'quiet eye' (QE) strategy. Experts show more frequent and longer gaze fixations on specific aspects of the putting situation (e.g., golf ball, putting surface, target hole). Novices tend to fixate more introvertedly (e.g., on the clubhead during the backswing phase) and focus on the ball only after it has been hit with the club (Moore et al., 2012; Vickers, 1992, 1996; Vine et al., 2013, 2017). Visual guidance by external focus of attention, that is, focusing on the target leads to a more effective golf swing than technical training of the postures and limb movements composing a golf swing. Indeed, in contrast with technical training, QE training of golf putting leads to improved putting accuracy as well as to a reduction in heart rate and muscle activity at the moment of impact (Moore et al., 2012). Thus, it is believed that instructing athletes by directing their perceptual linkages outward to the visual field supports higher task performance as well as leads to more expert-like movement and physiology.

QE research opens a fascinating vantage point on the perceptual linkages anchoring organisms to their visual contexts and the synergies transforming visual information into action (e.g., Kelty-Stephen et al., 2020; Mangalam, Lee, et al., 2020; Profeta & Turvey, 2018). Research on QE has mostly examined the visual-cognitive bases for the relatively more extraverted gaze into the task context (Panchuk & Vickers, 2006; Rienhoff et al., 2016; Vickers, 2011), and the relationship of QE to the body, the brain, and the nervous system has received relatively less attention (Davids & Araújo, 2016; Renshaw et al., 2019; Shine et al., 2018). To fill this gap, a 'postural-kinematic hypothesis' has begun to complement the canonical 'visual hypothesis' that QE was a primarily visual-cognitive mechanism (Gallicchio & Ring, 2020). So far, this postural-kinematic hypothesis emphasizes only slower movement times, that is, longer duration offering the foundation for longer fixations. This postural-kinematic hypothesis is a constructive first step. In this study, we aim to embed more of the postural dynamics into this hypothesis, beyond simply movement duration. Additionally, instead of reporting on gaze measurements, we will use QE as a training manipulation and focus on investigating how postural sway supports the effect of the training.

1.2. Optic flow and postural sway can support the quiet eye strategy

A crucial consideration for postural-kinematic contributions to visually guided tasks is optic flow (Stoffregen, 1985, 1986; Stoffregen & Riccio, 1990). Amidst the 'noisy' fluctuations of the athlete's body, quieting the eyes might leave fewer saccadic interruptions to the coupling between optic flow and postural sway. Optic flow consists of the differential expansion, contraction, or rotation of visible objects and surfaces as an individual moves about. It is specific to both the individual's movement and the contextual layout of objects and surfaces. Indeed, virtual visual displays that impose artificial optic flow also engender anticipatory behaviors (e.g., steering, braking) as if the objects and surfaces entailed by the optic flow are out there—not just for humans but for other animal species and even robots (Serres & Ruffier, 2017). Although most prominent in locomotion-related contexts, postural sway is no less capable of generating informative optic flow in non-locomotion contexts (Stoffregen et al., 2005). Because research on QE has mostly focused on the fixation of the eye gaze, QE's relationship to optic flow remains a critical gap (de Oliveira, 2016).

On the face of it, the preceding arguments might ring hollow at first. Certainly, optic flow requires movement, and it could seem wholly unrelated to the issue of visual fixations supporting wholly internal processing of incoming information into a motor program for the body's subsequent control. This gap between QE and optic flow follows closely from ongoing controversy in the literature about the relative relevance of ecological vs. information-processing explanations. Optic flow has ecological roots in Gibson's (1979) work that has been antithetical to information-processing, potentially raising the fears that this work might have unseating or simple designs, disregarding the extensive, diligent work on the information-processing account of QE. Indeed, a rare mention of 'optic flow' in the information-processing approach to QE wedded 'optic flow' to 'the teachings of Gibson,' so concerned with invariance as to 'treat every gaze as the same,' and reluctant to respect the fine-grained dynamics of gaze (Vickers, 2009, p. 286). We make full disclosure: the present use of multifractal modeling of postural access to optic flow is inspired by Gibson's appeal to flows and nestings these flows across many scales (Kelty-Stephen & Wallot, 2017), but we come to this dialogue with the first-hand experience examining the rich heterogeneity of gaze dynamics (e.g., Kelty-Stephen & Mirman, 2013; Stephen & Mirman, 2010) and with such interest in cognitive process as for Gibsonians to openly doubt that some of us could be Gibsonians (Raja & Anderson, 2019). Indeed, our interest in how exploratory behaviors might change perceptual response sets us firmly outside one of the stricter readings of Gibsonian invariance (Turvey, 1977).

Rather than digging into old trenches, we think that optic flow has a theoretical reach that might pick up where this information-processing account remains incomplete. Optic flow has grown to reach well beyond Gibson's original intent and usage. For instance, we have already cited cases of optic flow that would have been startling through the strictly Gibsonian lens: information-processing robots use optic flow as well as Gibson's organisms might (Serres & Ruffier, 2017), and optic flow does not require locomotion but can thrive on postural sway (Stoffregen et al., 2005). In the spirit of making the so far dialogue between ecological and information-processing approaches more constructive, the preceding two examples suggest that optic flow might be much more subtle (e.g., as in standing 'still') but also perfectly amenable to information processing (e.g., as in supporting computational models for controlling action).

Meanwhile, without optic flow to support it, we see QE's information-processing accounts relatively unmoored from their explicit requirements and proposals. According to a recent summary, the information-processing view of QE is that the eye's stillness provides time and attentional resources useful for the preprogramming of movements used in the aimed behavior (Vickers, 2009). This point is defensible but incomplete: visual-cognitive processes benefit only partially from a motor program's stipulation. Even non-ecological explanations have become newly interested in emergent

control parameters that sit outside of any simple stipulation by an internal motor program. Visual
attention is most effective when it is most task- and context-sensitive, and the whole issue entailed by
the Bernsteinian 'degrees of freedom' concern is the absence of a clear programmatic path from
intention to bodily execution. The usefulness of QE in various domains is a reason to recommend this
strategy, but it points to some long-unanswered questions. Are the motor programs implicated in QE
flexible and vast enough to accommodate bodily degrees of freedom and contextual ones? Or are these
motor programs elegantly minimal enough to leave room for non-programmed factors to support?
Neither option is unappealing on its face, but they both reflect an underdeveloped delivery on the
information-processing account's proposals about QE. Yes, longer looking times can support stronger,
more enriched motor programs (Witt & Proffitt, 2008). However, it is not always so simple: either
looking time is not always a reliable predictor of motor planning, or the motor program is flexible
enough to be rewritten on the fly during the program's execution (Hajnal et al., 2014, 2016).

Specifically, just as it has neglected to specify the finer print about its motor programs, QE's
information-processing account has neglected its claims about the role of action. The information-
processing account offers QE a bidirectional link between gaze, attention, and action (Vickers, 2009).
However, whereas this view may respect bidirectionality between attention and gaze (i.e., attentive
search for information and looking to what cues attention), the action appears strictly as the outcome
passively inheriting whatever the preprogramming entails but only following motor commands. If
motor programs generate the consequent swings, then the information-processing approach has yet to
elaborate upon two points: first, what bodywide movement coordination supports the postural control
allowing the steady visual fixation and, second, how the preprogramming contributes to the externally
focused movement. For instance, especially as participants repeat the aiming tasks in front of
potentially the same experimental display, the evidence suggests more variety in expert movements
than sameness, mainly as they engage visually in the external focus of attention (Morrison et al., 2016).
This point raises some questions. If QE is merely the same sort of fixation every time, irrespective of
movement, then why should programs carrying similar, static information give rise to varied bodily
execution of the program? If the motor programs that QE supports vary, what is the possible source in
the programs? In the former case, the programs might underspecify the movements, and in the latter
case, the programs would seem to embody more variety than the fixity of the visual scene. Suppose
action did participate in the same bidirectional relationship that information-processing approaches to
QE observe between attention and gaze. In that case, there might be a way forward, allowing for action
to participate in the tuning of programs to ongoing bodily postures and physiological changes
throughout repeated behaviors.

To summarize, the gap between QE and optic flow is critical because the information-
processing approach aligns the fixity of the eyes on one side with the accuracy of aimed behavior on
the other. However, there is a little account of the body between these fixations and accurate behaviors.
This omission of the body is immensely curious and potentially problematic because of the
movement's rich variability. To the point, it has been difficult for all motor programs to specify
accurate goal-directed behavior (Feldman, 2018) and no less difficult when eye movements are
implicated (Feldman & Zhang, 2020). Moreover, if we can leave off old associations and let optic flow
continue to take on shapes beyond the original Gibsonian imaginings, there is not so much a threat of
thwarting or disregarding prior art filling in the gaps left by information-processing. We aim to
rekindle this dialogue between cognitive psychology and ecological psychology, but we aim to do so
without requiring disavowal of any mutually opposed pre-theoretical commitments.

The present work aims to replicate the beneficial effects of QE on putting performance, and
more importantly, to understand how eye height and optic flow might support QE. Specifically,
assuming that QE might depend on the optic flow, we aim to test whether perturbing the optic flow

parameter of eye height perturbs the effect of QE training in golf putting more so than technical
training. Indeed, experimentally varying eye height is known to complicate optic flow in ways that
briefly perturb visual perception (e.g., Frenz et al., 2003). We used a single session in the style of
Moore et al.'s (2012) golf-putting training paradigm to test for rapid attunement to changes in
information from optic flow and postural sway. In non-golf contexts, for instance, manipulating eye
height by adding wooden blocks to participants' shoes alters participants' affordance judgments
significantly, leading to an initial underestimation of the optimal climbing height for a stair and sitting
height for a chair (Warren, 1984). However, participants adjust to the new eye height and make more
accurate decisions later on (Mark, 1987; Mark et al., 1990). Hence, we expect that adding wooden
blocks—(henceforth, 'clogs', to distinguish from 'blocks' of experimental trials)—to the feet might
upset putting accuracy in the QE-trained participants initially, but that this accuracy will rebound as
participants adjust to new eye height (Rienhoff et al., 2016). Optic flow is the structured change as
people move around in a visual world, often very subtle, occurring through minuscule postural sway
during 'quiet standing.' What is crucial is its centering on an external focus of control, outwards around
the organism, and aligned with their eye height. Hence, eye height is a crucial parameter of optic flow
(e.g., Warren, 1984), and when we perturb eye height, we destabilize optic flow, and the organism
needs to learn new relationships between bodily movement and movement of visible surfaces.

1.3. Multiscaled organization of postural sway entails a multifractal structure

Besides our manipulation of eye height, our expectations about the optic-flow foundation for QE also
include a hypothesis about which specific aspects of postural sway should moderate the effects of
perturbed eye height. If optic flow supports QE, we expect that QE-trained response to eye height
perturbation will depend on postural sway. Indeed, in the affordance research on reattunement to
perturbation of eye height, postural sway supported the adaptation of the relationship of eye height to
judgments of optimal chair height for sitting (Mark et al., 1990). However, it was not simply that more
sway facilitated this adaptation because, for instance, an 'awkward stance' requiring more
compensatory postural adjustments did not support visual adaptation as much as comfortable standing
did. An additional challenge in identifying what portion of sway could support adaptation to altered eye
height is that sway depends on events at multiple timescales. At longer scales, QE training itself might
influence sway by training an explicit use of visual information. At shorter timescales, adding clogs
might destabilize posture initially, and then at the shortest timescales, sway might be implicated in
adapting to this perturbation of eye height. Hence, judiciously accounting the sway's contribution to
visual adaptation due to perturbation of eye height requires an appreciation of sway's multi-scaled
organization.

The multi-scaled organization of sway may itself reveal a key to predicting—if not explaining
—how the movement system adapts its visually-guided actions. Interactions across multiple timescales
noted above are currently understood to entail a 'multifractal' structure in sway. Multifractality refers
to multiple (i.e., 'multi-') relationships among fluctuations and timescales, with these relationships
following power laws with potentially fractional (i.e., '-fractal') exponents. Interactions across
timescales engender a nonlinear patterning of fluctuations that multifractal analysis is well-poised to
diagnose and quantify (Ihlen & Vereijken, 2010). Specifically, multifractal analysis estimates a
spectrum of fractal exponents whose width W_{MF} quantifies multifractality. When we compare measured
series' W_{MF} to spectrum widths of a set of best-fitting linear models ('linear surrogates') of those series,
multifractal evidence of nonlinearity manifests as a significant difference between W_{MF} for the original
series and the average W_{MF} for linear surrogates. The t-statistic, t_{MF}, comparing the original series' W_{MF}
to the surrogates' W_{MF} serves as an estimate of what we call 'multifractal nonlinearity' (i.e., how much
multifractality in the original series is due to nonlinear interactions across timescales).

1.4. Multifractal nonlinearity predicts perceptual responses to perturbations of task constraints

Multifractality, W_{MF}, in postural sway is a robust predictor of perceptual responses in both visual and
haptic media (Doyon et al., 2019; Hajnal et al., 2018; Teng et al., 2016). Multifractal nonlinearity, t_{MF},
in postural sway predicts perceptual responses as well, not just in the case of nonvisual judgments of
length and heaviness of manually-wielded objects (Mangalam & Kelty-Stephen, 2020) but also in
visuomotor tasks (Bell et al., 2019; Booth et al., 2018) and in response to perturbations, such as those
due to prismatic goggles in a visual aiming task (Carver et al., 2017). Now, we aim to test whether t_{MF}
can predict the responses to perturbations of eye height in QE-trained golf putting.

We expected that postural sway most adaptive to the perturbation of eye height would show
relatively narrow multifractal spectra that would still differ significantly from those of the surrogates.
So far, postural sway shows relatively narrower multifractal spectra (i.e., smaller W_{MF}) for younger
participants (Munafo et al., 2016), those less likely to experience motion sickness (Koslucher et al.,
2016), and those focusing on more proximal surfaces in optic flow (Li et al., 2020). Hence, greater W_{MF}
in postural sway might counteract any effects of QE. All these findings are qualified by a failure of
W_{MF} to correlate with canonical measures of sway like standard deviation. However, postural stability
is only ambiguously related to the standard deviation of sway, requiring neither too much nor too little
variability in sway (Chen et al., 2008; Rajachandrakumar et al., 2018). Meanwhile, both empirical and
theoretical work suggests that t_{MF} is more closely related to postural stabilizing. Evidence shows that
postural stabilization following a perturbation to the base of support shows a rapid reduction of t_{MF}
(Kelty-Stephen, 2018). The theoretical simulation also predicts that random perturbations to individual
scales of a nested system lead to narrower multifractal spectra that still differ from linear surrogates'
spectra (i.e., smaller W_{MF} and smaller-yet-significant t_{MF}; Lee & Kelty-Stephen, 2017). Hence, adapting
to the posturally perturbed visual information must involve exhibiting smaller-yet-significant t_{MF}.

1.5. Golf-putting task and hypothesis

The participants were instructed to take golf putts towards a circular, hole-sized target painted on a
green putting surface in the laboratory—over four blocks of ten trials each. One group received QE
training, and the other group received technical training specifying how to move their limbs to make
the golf putt. Wooden clogs were attached to the feet of all participants to perturb their eye height.
Putting performance was treated continuously (i.e., in terms of radial error).

1.5.1. Hypothesis-1: Multifractal fluctuations in postural sway will lead an eye height
perturbation to interrupt QE-trained performance more so than technical-trained performance

Putting performance was treated continuously in terms of unsigned radial distance from the target (i.e.,
unsigned radial error). We hypothesized that the eye height perturbation due to clogs would lead to
lower putting accuracy characterized by higher unsigned radial error following QE training than
technical training. We predicted that the clog's effects on QE would depend on the multifractal
nonlinearity, t_{MF}, of sway beyond any effects of multifractality (i.e., W_{MF}) alone (Hypothesis-1a). We
also predicted that lower t_{MF} of sway would be associated with more adaptive responses to the eye
height perturbation, leading to a gradual increase but a subsequent decay of radial error across trials
within the blocks with the clogs on (Hypothesis-1b).

2. Methods

2.1. Participants

Twenty undergraduate students (11 women, *mean*±1*s.d.* age = 19.5±1.2 years, all right-handed,
Oldfield, 1971) with normal or corrected vision participated in this study. All participants provided
written informed consent approved by the Institutional Review Board (IRB) at Grinnell College
(Grinnell, IA).

2.2. Procedure

In an hour-long session, the participants completed a single putt on each of 40 trials. Prior to the training trials, half of the participants received QE instructions (Table 1, left column) while the other half received technical putting instruction (Table 1, right column), replicating Moore et al.'s (2012) methodology. The technical instructions mirrored the QE instructions but did not describe the eye-related behaviors, hence mimicking standard golf instruction. After verbally confirming that they understood the instructions, the participants watched a video demonstrating a QE-specific putting stroke (https://www.youtube.com/watch?v=lfWuL1klbsY&feature=youtu.be) or a standard putting stroke (https://www.youtube.com/watch?v=HOYZn4MYmDI). The researcher directed the quiet-eye trained participants to the key features of the golfer's gaze control and the technically-trained participants to the putting mechanics.

The participant performed the golf putts on a 6×15 feet standard indoor All Turf Matts™ artificial turf mat using a 1.68-inch diameter PING Sigma 2 Anser Stealth™ putter and regular-sized white golf balls. The participants completed straight putts ten feet away towards a 2.5-inch diameter circular orange dot, designed to fit the golf ball. This format allowed for more stringent measurement and gave the participants a more focused target.

The participants completed 20 training trials (treated as two blocks of 10 trials per block in regression modeling). Following a short 5-minute break, the participants completed two blocks of ten test trials per block (i.e., 20 test trials total) in the absence of instructions. The participants were randomly assigned to complete either their first or second block of ten trials while wearing 5-cm-raised wooden blocks (i.e., clogs) attached to their shoes (they completed the other block of ten test trials under the same conditions as the 20 training trials). Before the testing began, the participants were allowed to walk around and adjust to the clogs to reduce fall risk.

2.3. Measures

The task performance was defined in terms of the radial error (the distance from the target at which the ball stopped, in feet) and the number of putts successfully 'holed.' Because the target was not a real hole, the putt was counted as holed in either of two events: the ball came to rest within the orange dot on the floor or within 2 feet directly beyond the hole (Cooke et al., 2010; Vine et al., 2011). Postural sway was measured using a smartphone with the Physics Toolbox Suite application (http://vieryasoftware.net) strapped to the participants' lower back. The app recorded the acceleration of the torso along x-, y-, and z-dimensions at 100 Hz. An experimenter manually started and stopped recording for each putt. Recording began after the participant set up over the ball and stopped once the ball stopped moving.

2.4. Multifractal analysis

2.4.1 Multifractal spectrum width of accelerometer displacement series

For each putt, we computed an accelerometer displacement series for multifractal analysis as follows. The accelerometer data encodes an N-length series of accelerations of the torso along x-, y-, and z-dimensions centered on an origin on the smartphone. A displacement series of length $N-1$ was computed by: 1) taking the first differences for the variable (e.g., if we use the letters i and j to indicate individual values in sequence, $x_{diff}(j) = x(i) - x(i-1)$ for each ith value of x, for $2 < i < N$, $1 < j = i - 1 < N - 1$, 2) squaring and summing the differences of each dimension's displacements (i.e., $x_{diff}^2 + y_{diff}^2 + z_{diff}^2$), and 3) taking the square root of this sum (i.e., $\sqrt{(x_{diff}^2 + y_{diff}^2 + z_{diff}^2)}$).

Multifractal spectra $f(\alpha)$ were estimated for each accelerometer displacement series, using Chhabra and Jensen's (Chhabra & Jensen, 1989) method. The first step was to partition the series into

subsets (or 'bins') of various sizes, from four points to a fourth of series' length (Fig. 1, left). The second step was to examine how the proportion of displacements within each bin changed with bin size or timescale (Fig. 1, right). The third step was to estimate two distinct exponents: an exponent α describing how the average proportion grows with bin size and an exponent f describing how Shannon's (Shannon, 1948) entropy of bin proportion changes with bin size. The later steps were all iterations of this third step, calculating a mass μ by applying an exponent q to bin proportions and using $\mu(q)$ to weight proportions selectively according to proportion size. q greater than 1 weights larger-proportion bins, and q lesser than 1 weights smaller-proportion bins (Fig. 2, left). q-based mass $\mu(q)$-weighting generalizes single α and f estimates from earlier steps into continual $\alpha(q)$ and $f(q)$— with $\alpha(q)$ describing how mass-weighted average bin proportion changes with bin size and $f(q)$ describing how Shannon's entropy of bin masses $\mu(q)$ with bin size (Fig. 2, top right).

Multifractal analysis quantifies heterogeneity as variety in $\alpha(q)$ and $f(q)$. The ordered pairs ($\alpha(q)$, $f(q)$) constitutes the multifractal spectrum (Fig. 2, bottom right). We included $\alpha(q)$ and $f(q)$ only when mass-weighted proportions were defined and when both of their corresponding relationships to bin size correlated at $r > 0.995$ on logarithmic plots, for bin sizes $4, 8, 12, \ldots N/4$ and $-300 < q < 300$, where N is the number of samples in the series.

2.4.2. Surrogate comparison

Multifractal spectrum width is sensitive to nonlinear interactions across timescales as well as to linear temporal structure. Hence, evidence of nonlinear interactions across timescales requires comparing the original series' multifractal spectrum width W_{MF} to those computed for a sample of linear surrogate series (Fig. 3). These linear surrogate series retain the same value as the original series but in a different order that preserves the original series' linear autocorrelation but destroys original sequence (i.e., iterative amplitude-adjusted Fourier-transform or IAAFT (Ihlen & Vereijken, 2010; Schreiber & Schmitz, 1996); Figs. 4a to d). Comparing W_{MF} of the original and surrogate series (Figs. 4e, f) produces a t-statistic t_{MF}, a measure of nonlinear interactions across timescales that is standardized across series with differing linear temporal structure. t_{MF} was computed for each original series.

2.5. Mixed-effects modeling

The data were submitted to two kinds of mixed-effect modeling, namely, logistic and Poisson, using the glmer() function in R-package *lme4* (Bates et al., 2007). The subsequent subsections provide details of the predictors and the models.

2.5.1. Effects of time

The effects of time were modeled using trial number within a block (trial$_{block}$), block number (block), and interactions of trial number with block number (trial$_{block}$ × block), and trial number across blocks (trial$_{exp}$). The model used orthogonal polynomials of each of both predictors up to 3rd order (i.e., cubic). The primary reason for the polynomial terms was to reveal how and when the perturbation of wearing clogs might manifest and consequently subside. Cubic polynomials allow the simplest way to accomplish this goal by allowing a peak and a flat, quiescent portion as part of the same continuous function. Additionally, unlike the quadratic function, the cubic peak is not required to be symmetric and occurring at the midpoint. Hence, the cubic function controls for any nonlinearities in the normal course of trials without such perturbations. Whereas block is a linear effect across blocks, including trial$_{exp}$ in the model allowed controlling for nonlinear change across blocks.

There are two important features to note about the treatment of time in this model. First, the Poisson model encoded both classifications of trial numbers (i.e., trial$_{exp}$ and trial$_{block}$) in terms of an orthogonal 3rd-order polynomial, using the function 'poly' in R, to include the linear, quadratic and cubic growth of trial number with all three terms phase-shifted so as to be uncorrelated with one

368 another. Second, the lack of model convergence limited the appearance of block, such that block was
369 neither treated nonlinearly nor in interaction with perturbation. Attempting to include the orthogonal
370 3^{rd}-order polynomials for block number led to a failure of model convergence, so block only appeared
371 as a linear term. The model also failed to converge when it included interactions of block and
372 perturbation because these two terms were strongly collinear.

2.5.2. Effects of manipulations

374 Manipulations included quiet-eye instruction, QE (QE = 1 for receiving QE instruction, and QE = 0 for
375 not receiving QE instruction) and perturbation of eye height, perturbation (perturbation = 1 for all ten
376 trials during the block when participants wore the clogs, and perturbation = 0 for all other trials).

2.5.3. Sway: Linear predictors

378 The accelerometer data encodes accelerations of the torso along x-, y-, and z-dimensions. Any single
379 posture corresponds to a specific point in this 3D space. Hence, sway corresponds to the deviation of
380 this 3D acceleration around the mean posture. The present analyses encoded this variability as MSD_{acc}
381 or $RMS_{acc,}$ depending on which predictor supported a convergent model, along with mean and standard
382 deviation of 3D displacement ($Mean_{disp}$ and SD_{disp}, respectively).

2.5.4. Regression modeling

A mixed-effects Poisson regression modeled the radial error (in inches) of all misses, with makes as
defined above treated as having zero radial error. Predictors included QE × perturbation × $trial_{block}$ × t_{MF}
to our hypothesis by specifically addressing trials within each block, QE × $trial_{block}$ and QE × $trial_{exp}$ to
control for differences in learning due to QE across a linear progression of block; a nonlinear function
of trials, QE × perturbation × W_{MF} × t_{MF}, to control for the effects of multifractal spectral width and its
relationship with t_{MF}; and MSD_{acc} × $Mean_{disp}$ × SD_{disp} to control for the effects of linear description of
sway. Modeling included lower-order interactions and component main effects (e.g., Allison, 1977).

3. Results

3.1.1. Non-multifractal effects on radial error

This section details the non-multifractal effects described in Supplementary Table S1 and effects
preliminary to the test of our hypothesis in Table 2. This model of radial error found a significant main
effect of QE training, specifically, a negative effect (b = –4.40×100, $s.e.m.$ = 4.81×10^{-1}, p < 0.0001)
indicating that QE did promote greater accuracy in terms of proximity of the ball's final position to the
hole. Radial error grew linearly and quadratically across trials but showed negative cubic variation with
trial—both for trials across the experiment and within a block (Fig. 5a). The QE strategy prompted
steady decay of error across the experiment, canceling out the linear and cubic pattern of error over
trials within a block (Fig. 5b). The perturbation of eye height elicited no main effect but increased the
growth of error over trials within a block, with the QE instruction accentuating this nonlinear growth of
error later in the block. The higher-order interaction MSD_{acc} × $Mean_{disp}$ × SD_{disp} and its main component
effects and lower-order interactions showed mostly negative effects (only two of the two-way
interactions showed positive coefficients).

**3.1.2. Total multifractality of sway, W_{MF}, increased radial error, but multifractality due to
nonlinearity, t_{MF}, attenuated radial error at the beginning and the end of blocks**

No matter what portion of multifractality is attributable to linear or nonlinear structure, the total width
W_{MF} of original series' multifractal spectra is associated with greater radial error (b = 1.80×10^0, $s.e.m.$ =
1.29×10^{-1}, p < 0.0001). We did not find any main effect of multifractality due to nonlinearity, t_{MF}, nor
any interaction of W_{MF} × t_{MF}. However, the negative effect for t_{MF} × $trial_{block}$ (Quadratic; b = –8.93×10^{-2},

s.e.m. $= 1.15 \times 10^{-2}$, $p < 0.0001$) indicated an association between multifractality due to nonlinearity
with a significant reduction of the positive quadratic profile for radial error over trials within a block
(i.e., trial$_{block}$(Quadratic); Section 3.1.1.).

3.1.3. Testing Hypothesis-1a: Perturbation of eye height increased radial error only in combination with QE training, with greater multifractality due to nonlinearity t_{MF}, or with both

Much like the effect of trials within a block (Section 3.2.3.), the multifractality of sway promoted the
tendency of QE to increase perturbation-related radial error. Perturbation showed a significant effect
through its interactions with other factors, the earliest being its interaction with trial$_{block}$ (Section 3.2.3.);
the interaction of perturbation $\times$ W_{MF} ($b = -1.50 \times 10^{0}$, *s.e.m.* $= 3.12 \times 10^{-1}$, $p < 0.0001$) canceled out the
positive effect of W_{MF} on error (Section 3.1.2), suggesting that the perturbation reduced the error-
eliciting effect of W_{MF}. The interaction QE $\times$ W_{MF} reduced error by roughly half ($b = -1.04 \times 10^{0}$, *s.e.m.*
$= 2.03 \times 10^{-1}$, $p < 0.0001$), suggesting that QE promoted greater accuracy but only in the absence of
perturbation. However, the interaction with multifractality indicates that QE increased the error
attributable to the perturbation. For instance, the QE $\times$ perturbation $\times$ W_{MF} ($b = 2.12 \times 10^{0}$, *s.e.m.* $=$
4.91×10^{-1}, $p < 0.0001$) reverses the apparently beneficial error-reducing interaction effect of
perturbation $\times$ W_{MF}. Hence, QE reinstated the error due to total multifractality (Section 3.1.2), and it did
so only in the face of the perturbation.

Above and beyond the effects of W_{MF} in moderating the interaction effect of QE $\times$ perturbation,
t_{MF} promoted the error-increasing effect of perturbation. Specifically, besides the moderating effect of
QE on the effect of perturbation $\times$ W_{MF}, perturbation elicited even greater radial error with increases in
t_{MF} (perturbation $\times$ W_{MF} $\times$ t_{MF}: $b = 1.54 \times 10^{-1}$, *s.e.m.* $= 3.15 \times 10^{-2}$, $p < 0.0001$). QE did reverse this effect
(QE $\times$ perturbation $\times$ W_{MF} $\times$ t_{MF}: $b = -2.21 \times 10^{-1}$, *s.e.m.* $= 4.97 \times 10^{-2}$, $p < 0.0001$). However, QE also
increased error in combination with perturbation and t_{MF} even in the absence of any moderating effect
of W_{MF} (QE $\times$ perturbation $\times$ t_{MF}: $b = 8.62 \times 10^{-3}$, *s.e.m.* $= 4.18 \times 10^{-3}$, $p < 0.05$). This result supports
Hypothesis-1a.

Despite the absence of a simple QE $\times$ perturbation interaction, the eye height perturbation
produced a greater increase in radial error for QE-trained participants. To illustrate this point, we can
compare the model predictions for participants in each condition with high and low W_{MF} and t_{MF} (with
'low' and 'high' values defined as 1st and 3rd quartiles). For participants with 'low' W_{MF}, the eye height
manipulation prompted 2.27% to 11.14% increase in radial error in QE-trained compared to technical-
trained participants for low to high t_{MF}, respectively. For participants with 'high' W_{MF}, the eye height
manipulation prompted 8.80% to 18.23% increase in radial error in QE-trained compared to technical-
trained participants for low to high t_{MF}, respectively.

3.1.4. Testing Hypothesis-1b: the QE group with greater multifractal nonlinearity in sway showed a greater initial but progressively slower increase in radial error due to the perturbation of eye height

The interactions of QE $\times$ trial$_{block}$ with multifractal nonlinearity followed a similar but weaker pattern as
the interactions of QE $\times$ trial$_{block}$ with perturbation. Much like the perturbation of eye height, the
interaction of QE with t_{MF} accentuated the radial error late in a block of ten trials (QE $\times$ trial$_{block}$ $\times$ t_{MF}):
it weakened the linear term ($b = -3.73 \times 10^{-2}$, *s.e.m.* $= 1.46 \times 10^{-2}$, $p < 0.05$), contributing to a negative-
quadratic peak in the middle of the block ($b = -1.85 \times 10^{-1}$, *s.e.m.* $= 1.66 \times 10^{-2}$, $p < 0.0001$) and a
positive cubic increase at the end of the block ($b = 1.13 \times 10^{-1}$, *s.e.m.* $= 1.73 \times 10^{-2}$, $p < 0.0001$). Hence,
t_{MF} and perturbation showed independent effects on the progression of error across trials in block.

However, the similarity of these effects independently on error across trials within a block did
not extend to their interaction. Indeed, most relevant to testing Hypothesis-1b is that the interaction of
perturbation and t_{MF} served to speed up the growth of error, and QE slowed it down across the block.

Coefficients for perturbation $\times$ trial$_{block}$ $\times$ t_{MF} indicated that in the face of perturbation, multifractal
nonlinearity removed the observed peak in error due to earlier negative quadratic effects ($b = 3.83 \times 10^{-1}$,
$s.e.m. = 5.28 \times 10^{-2}$, $p < 0.0001$) and increased error across the linear ($b = 4.07 \times 10^{-2}$, $s.e.m. = 5.20 \times 10^{-2}$,
$p < 0.0001$)and cubic terms ($b = 5.39 \times 10^{-1}$, $s.e.m. = 4.45 \times 10^{-2}$, $p < 0.0001$). On the other hand,
coefficients for perturbation $\times$ trial$_{block}$ $\times$ t_{MF} indicated that, in the face of perturbation for the QE
participants, the linear increase was shallower ($b = -2.71 \times 10^{-1}$, $s.e.m. = 6.11 \times 10^{-2}$, $p < 0.0001$), the
positive quadratic growth further removed the observed peak in error ($b = 4.24 \times 10^{-1}$, $s.e.m. = 6.14 \times 10^{-2}$,
$p < 0.0001$), and the cubic term was negative ($b = -7.90 \times 10^{-1}$, $s.e.m. = 5.79 \times 10^{-2}$, $p < 0.0001$).

The outcome of these differences distinguishes how multifractal nonlinearity predicted changes
in adaptive responses to the perturbation of eye height. Greater t_{MF} exhibited less adaptive response to
the perturbation in the QE case, and less-but-still-nonzero t_{MF} predicted a more gradual appearance of
error and then a subsequent decay of this error. Greater t_{MF} led the QE group to exhibit initially greater
error and a smaller change in error across trials, and lower levels of t_{MF} led the QE group to show a
delayed growth in error and subsequent decay in error (Fig. 5b), suggesting adaptation. It is evident in
figure 5b that although the error in the high-t_{MF} cases with the perturbation was smaller than the
delayed increase in the low-t_{MF} cases with the perturbation, the high-t_{MF} cases showed more sustained
error and no adaptation to the perturbation relative to high-t_{MF} cases without the perturbation. This
result supports Hypothesis-1b.

3.2. Logistic regression of 'makes' vs. misses showed less sensitivity to effects

Poisson regression may be too ornate and elaborate to model the behavioral responses to the
manipulations or effects of endogenous multifractal fluctuations. However, in contrast to Poisson
modeling of a continuous outcome measure of performance, the investment of logistic regression to
model dichotomous variable limits the number of valid predictors (e.g., van Smeden et al., 2016) and
so affords us less statistical power with which to test for changes in the performance. A mixed-effect
logistic regression model of successful putting (i.e., the 'makes') returned the coefficients reported in
Table 3. Predictors for t_{MF} and for interactions of QE with all terms other than perturbation failed to
improve model fit significantly and so were omitted from the final model. This model failed to return
an effect of QE ($b = 0.20$, $s.e.m. = 0.30$, $p = 0.51$), of the perturbation alone ($b = -0.22$, $s.e.m. = 0.43$, p
$= 0.60$) or the interaction of QE $\times$ perturbation ($b = -1.16$, $s.e.m. = 0.65$, $p = 0.07$).

We found effects of experience and trial-by-trial sway. Block showed a positive effect ($b =$
0.24, $s.e.m. = 0.13$, $p = 0.06$), suggesting that odds of a successful putt was 1.27 times higher with each
successive block of trials. RMS_{acc} showed a positive effect ($b = 4.05$, $s.e.m. = 1.34$, $p < 0.01$), and total
multifractality W_{MF} showed a negative effect ($b = -7.25$, $s.e.m. = 2.94$, $p < 0.05$), suggesting that
successful putts were more likely with more variable and less multifractal accelerations of sway. These
effects resembled some of those found in the Poisson regression model, but it appears that the
continuous measure of radial error afforded more sensitivity within which to model significant fixed
effects.

4. Discussion

We tested the effects of postural stability and its endogenous multifractal fluctuations on a continuous
outcome of putting performance (radial error) with and without QE training. We hypothesized that QE-
trained participants' response to an eye height perturbation would depend on the multifractality
attributable to nonlinear interactions across scales, t_{MF}. That is, the multifractal fluctuations in postural
sway would make eye height perturbation a threat to QE-trained performance but not to technically
trained performance. Specifically, we predicted that lower estimates of t_{MF} would coincide with a more
adaptive response, with slower growth of radial error and subsequent decay of radial error over trials.

Results supported the first hypothesis: when we perturbed eye height, accuracy with QE was associated
with reducing multifractal nonlinearity in postural sway. This model included a negative main effect of
QE training, indicating more accurate putts in radial error. However, despite the absence of a simple
interaction of QE with clogs, QE's contingency on multifractal aspects of postural sway entailed a
greater decrement in accuracy following eye height perturbation.

A limitation of these results is that our focus on the postural support of QE-trained performance
came at the expense of focusing on gaze tracking. We used QE training as a manipulation, but we
failed to include a manipulation check. The absence here of gaze-tracking data leaves us unable to
explain with complete confidence why a logistic model of putting performance construed
dichotomously (e.g., as 'makes' vs. 'misses') failed to find a significant effect of QE. On the one hand,
some participant may have followed QE instruction incompletely. On the other hand, a good reason
why the main effect may not appear is the brevity of our design: whereas recent analogues of our
research design already in the literature required participants to conduct 4000 putts over 7 days (Moore
et al., 2012), we only sampled 40 putts in one hour-long session. Radial error may be sufficiently
sensitive to show QE's reliable effect in a more compact design than dichotomous measures. Future
research would no doubt benefit from incorporating gaze-tracking into similarly compact experimental
designs.

These results offer two major insights for research into perception-action in visual aiming tasks.
First, it shows that QE training has roots in postural sway, vindicating and elaborating earlier proposals
that quiet eye is not simply about stabilizing the eyes (Liphart et al., 2015; Williams, 2016). Indeed, it
has long been known that simply stabilizing the retinal position compromises the persistence of a visual
image (Barlow, 1963; Coppola & Purves, 1996; Ditchburn & Ginsborg, 1952). And contrary to a
colloquial understanding of 'quiet eye,' more recent work confirms that those small movements within
gaze fixations called 'microsaccades' might prevent fading (Martinez-Conde et al., 2006) and support
visual attentional processes (Krueger et al., 2019). We might draw a comparison between the 'quiet'
suggested by 'quiet eye' and that indicated by 'quiet standing.' To be precise, we mean to draw the
comparison of 'quiet eye' specifically only to 'quiet standing' in the case of instructing participants to
'stand as still as possible' (i.e., rather than to 'stand as relaxed as possible'; (Houdijk et al., 2015). Both
'quietudes' provide the needed instruction to a participant or athlete to 'please move as little as
possible,' but below the polite clarity for verbal instruction, both 'quietudes' depend upon a rich texture
of fluctuations. Fluctuations throughout the body support exploration during the 'still as possible' quiet
standing (I.-C. Lee et al., 2019; Murnaghan et al., 2016; Palatinus et al., 2013, 2014).

Here, the theoretical ground may shift underfoot as we try to see a path through these
considerations. Do we mean to say that the achievement of a precision demand could be exploratory?
At first blush, it could sound absurd that something not moving is also exploring. However, here, we
may hit upon a deeper divergence between ecological vs. information-processing perspectives on QE.
To the point, it is true that fixing the gaze and standing as still as possible are themselves
accomplishments of motor control. However, this achievement does not entail that either quiet eye or
quiet stance are the results. Unless we are much mistaken, it is the external focus and the intention of
getting the ball into the net or hole or target that warrants quiet eye in the first place. That is to say, the
quiet eye is a means to an end, and the end is never just quiet eye.

Precision might invite the appearance of achieving fixity, but achieving fixity might be no
guarantee against exploration. For a contrary example, 'static holding' is a haptic case of holding an
object 'as still as possible,' and it is widely recognized as a valid exploratory procedure for haptic
perception (Burton & Turvey, 1990; Carello et al., 1992; Lederman et al., 1996). The only debates are
what kind of information this apparent-fixity mode of exploration has access to. Crucially, apparent
fixity is not necessarily actual fixity, and while there is no need for the body to explore the quiet-eye

achievement of ocular fixity, it would strike us as a strange eventuality to rule quiet-eye strategy out as
a strategy pertaining to exploration. Quiet eye resembles to us an ocular version of static holding that
holds the visual image as close as possible. As noted above, despite their apparent fixity, fixations are a
class of eye movements, and these fixational eye movements are exactly what the quiet-eye strategy
uses to build a motor program. Certainly, if the QE strategy is collecting information that supports the
construction of a motor program, ruling quiet eye out as a form of exploration would strike as a
problematic move. For instance, consider the pioneering astronomers gazing up at the heavens, aiming
a telescope as quietly as possible at a star or planet. Can we tell them that, no, because they are engaged
in precision work, they are not exploring? We suspect not. Again, to summarize this point: a precision
demand can change motor behavior (Balasubramaniam et al., 2000), but that change is not itself the
end of the story. Despite experimenter intentions to train it, quiet eye is not a goal in itself but only a
strategy for informing the participant's accurate behavior beyond their bodies for precision wherever
they direct their external focus.

Of course, the preceding proposal that QE is exploratory remains conjecture that reaches
beyond the current evidence. Demonstrating an exploratory structure in QE will need empirical
evidence from future research to back up the preceding metaphor and analogy. Specifically, empirical
validation would require at least one of the following: (1) eye movement data indicating exploratory
movements within the fixation time that are similar or relational to those from the haptic or postural
research, (2) performance outcome data that indicates information is obtained that is of value during
the quiet eye fixation period, and/or (3) self-reporting of the subjects (perhaps quantified through
perceptual judgments) indicating information gained following QE training that differs from those who
do not go through the training. We look forward to future research to the opportunity to test these ideas
and revisit the question of whether QE is exploratory behavior.

The value of QE is not necessarily a net reduction of fluctuations but rather a way to train
athletes to orient their movements towards the optic flow generated by postural sway. Whereas the
postural-kinematic hypothesis had previously only referred to longer movement duration (Gallicchio &
Ring, 2020), we now report that movements are not merely slower but evince a sort of nonlinearity
constricting the degrees of freedom across scales of the movement-system hierarchy. In one of the
subsections below, we discuss how this orienting towards optic flow might coordinate with
mechanisms invoked by the more elaborate visual hypothesis (e.g., Gallicchio et al., 2016, 2017).

The second insight is that the multifractality in sway is an essential aspect of postural
adaptations contributing to visually-guided action. It was already known that damping out postural
perturbations could occasion a ratcheting down of multifractality in postural sway (Kelty-Stephen,
2018). The present results indicate that this reduction of multifractality is associated with more accurate
responses in visually-guided aiming. Note that this reduced multifractality remains different from
corresponding surrogates. Hence, nonlinear interactions across scales can constrict movement
variability, making reduced multifractality more adaptive. This distinction is a departure from any
shorthand presumption that 'more multifractal is better' supported by early studies of heart rate
variability (HRV; Ivanov et al., 2001). Indeed, this departure reflects a growing clarity about the value
of multifractal fluctuations for predicting and explaining outcomes in perception-action.

4.1. Clarifying the value of multifractal fluctuations for perceiving-acting systems

The shorthand 'more multifractal is better' presumptions from early studies on HRV (Ivanov et al.,
2001) have always underestimated the more elaborate systematicity in the physiology of the human
movement system with upright posture. Early work proposed that a 'loss of [fractal or multifractal]
complexity' would occasion or coincide with aging or with pathology (Lipsitz & Goldberger, 1992).
Indeed, early evidence showed that less healthy outcomes co-occurred with more-than-fractal levels of
temporal correlation (Lipsitz, 2002). However, the task has always been known to influence

multifractality in upright posture: multifractality of quiet standing can reduce with pathology (Morales
& Kolaczyk, 2002; Shimizu et al., 2002), but multifractality in gait can increase with pathology
(Muñoz-Diosdado, 2005), as well as with walking speeds faster or slower than self-selected
comfortable speed (West & Scafetta, 2003). Over the lifespan, the maturation of gait involves loss of
multifractality, beginning with the stabilization of toddler gait into young adult gait and continuing into
older age (Ashkenazy et al., 2002). And this lifelong progression of multifractality in gait actually is at
odds with more recent evidence that multifractality in HRV remains stable with healthy aging (Schmitt
& Ivanov, 2007).

Militating strongly against simple 'more multifractal is better' kind of presumptions are a
variety of distinctions. Some of these distinctions may sound patently obvious but need recognition if
we want to identify for what, after all, multifractality is good. These distinct points include, for
instance, that hearts are not upright bipedal bodies, that age is not itself a pathology, and that walking is
not standing. The remaining distinctions are more subtle. In the promise of fractal/multifractal analysis,
the discourse about 'complexity,' 'dynamical stability,' and the role of fractality/multifractality in
'optimality' (Friston et al., 2012; Oftadeh et al., 2014; Schlesinger & Klafter, 1986) has made no
claims on all definitions of optimality and so has not been responsible for acknowledging that some
multifractality is not optimal (Reynolds, 2015). Indeed, studies in behavioral sciences have always
justified the capacity of fractal variety to offer new insights (Diniz et al., 2011; Dixon et al., 2012).
Any notion of a privileged level or direction of fractal variety (Kello, 2018) might be best understood
as a reaction to critical, unreceptive perspectives resistant to seeing any fractality or multifractality
appear in behavioral-science discourse at all (Wagenmakers et al., 2012). Measurements may wax and
wane in their multifractal structure. The only case in which we think more multifractality is always
better than less is in the scientific discourse on the coordination of movement systems engaged in
perception-action.

4.2. The value of multifractal fluctuation is specific to task constraints

What is coming into focus is that the value of nonlinearity-driven multifractality depends on task
context. More of it is adaptive when tasks invite exploration and anticipation. Less of it is adaptive
when the task involves constraining behavior, drawing degrees of freedom inward around a base of
support. As an example of the former, accumulating information as a reader can require keeping an
open mind to follow the discourse wherever it may lead, and more multifractality t_{MF} in the pacing of
word-by-word reading supports more fluent reading if the story has a twist in its plot (Booth et al.,
2018). Similarly, more multifractality W_{MF} in circle-tracing behaviors make for poor tracing but is
associated with better performance on neurophysiological tests of flexibility with rule switching
(Kelty-Stephen et al., 2016). The freewheeling benefits of more multifractality can also compete
alongside the constraint-, repetition-promoting benefit of less multifractality. In a Fitts task, greater
multifractality t_{MF} in hand and head movements predicted less stable contact with the targets, but it
became a predictor of more stable contact when participants had the diffuse warning that they might be
asked to close their eyes and continue the task (Bell et al., 2019). Hence, what would have upset task
performance became a resource for exploring the task space for the eventuality of a major loss of
sensory information. And now, we see that a less multifractal postural system can be more stable (e.g.,
smaller t_{MF} in Kelty-Stephen (2018); and smaller W_{MF} in Koslucher et al. (2016) and Li et al. (2020)
and, as present findings indicate, more accurate in how it orients behaviors to a visually-guided aiming
task.

Some of the task-specificity remains unclear, e.g., manual wielding of an object is more
accurate with greater multifractality in postural sway, but hefting an object to perceive heaviness
compared to a reference object is more accurate with less multifractality (Mangalam & Kelty-Stephen,
2020). This difference could have to do with the qualities of length and heaviness, but the simpler

interpretation could be that intending to perceive directs attention away from the center of pressure and intending to perceive heaviness directs attention inwards toward the center of pressure, lending length perceptions and heaviness perception to more and less postural instability. This issue of directing attention is again a reason we will need to consider how to link this postural-kinematic work on QE back with the visual-hypothesis work on QE.

The fact that reduced multifractality can be beneficial offers a unique distinction between possibly two related but distinct interpretations of multifractal results: one as a biomarker of pathology and the other as adaptation. Indeed, the loss of multifractality can be a biomarker for diagnosing a pathology (Meyer et al., 2003). The physiological wisdom informing this usage is that the heart has to be, in effect, poised to absorb and rebound from perturbations, avoiding fragility due to insult at any single characteristic scale (e.g., Lipsitz & Goldberger, 1992). Implicit in this wisdom is that, within the healthy body, the heart's task is deeply anticipatory and exploratory, much like those task settings in which whole organisms benefit from more multifractality. However, concluding that loss of multifractality is strictly symptomatic of postural systems with deficits (Shimizu et al., 2002). What looks like a deficit may sometimes be a creative constraint, built and used to solve a task problem (Harrison & Stergiou, 2015; Holt et al., 2010; Vaz et al., 2017). The present results indicate that reduced multifractality could be an adaptive response preserving stability and promoting task performance in the face of abrupt change in the organism's mechanical relationship with the base of support. Hence, far from being the signature of diseased posture, reductions in multifractality that maintain a significant tMF may be an advantageous loss of multifractality. Though more commonly known for their capacity to amplify variability, nonlinear interactions across scales acting context-sensitively may also restrain motoric degrees of freedom so as to achieve the task goal.

Different task constraints prompt different modes of movement variability, and if multifractal depictions of the body seem to physicalize or mechanize the movement system, they do so only in the way that nonlinear-dynamical physical mechanisms might better explain context-sensitive behavior. Surely, good-faith pursuit into nonlinear-dynamical complexity should avoid any catch-all simplification (e.g., 'more is better, less is worse') and keep a keen eye on what sort of multifractality is good for what end. Nonlinear dynamics has also profited from consideration of its measures in different reference frames before: experimentally breaking apart the coincidence of relative-phase in spatial frames from relative-phase in muscular frames revealed novel, more generic structure in the Haken-Kelso-Bunz (1985) law for multi-limb coordination (Park & Turvey, 2008). Finally, considering the task frames for our interpretation of multifractal estimates may be equally crucial for generalizing predictions for a multifractal foundation for perception-action. Low multifractality in the heart signals a bad prognosis to the clinician, but low multifractality in posture supports good task performance in visually-guided aiming, particularly for an organism beset by malevolent experimenters uprooting their optic flow.

4.3. The value of multifractality is also specific to different tissues of the body.

Just as different tissues of the same body can differently support the same task, we expect that the multifractal structure that spans across these different tissues supports overarching integrity to the organismal-level behavior we measure in the laboratory (Stoffregen et al., 2017). Here, a comparison of present results with similar work using a relatively less-mechanical, visual perturbation (Carver et al., 2017), may tee up considerations for future work that could elaborate an explanation of QE by integrating the postural-kinematic hypothesis with the visual hypothesis. The present results resemble those of Carver et al. (2017) who found that, without any training (i.e., neither QE nor technical), more multifractality in torso sway led to greater error in a low-difficulty Fitts task (compare W_{MF} in Table 2) led to smaller error in the case of a visual perturbation (compare perturbation $\times W_{MF}$ in Table 2), that is, multifractality in head sway became a stabilizing, accuracy-promoting asset to task performance when

the target was far out of reach. Future research could also probe the visual hypothesis for QE for multifractal foundations, and based on Carver et al.'s (2017) results, we predict that QE-trained performance would improve with greater multifractality in head sway or eye movements. Monofractal analysis of eye movements has successfully predicted visual attention to text (Wallot et al., 2015), and such fluctuations in quiet eye likely support an extension of the multifractal support for QE. The present results do not include gaze data, but we hope it is informative, especially in support of the postural-kinematic hypothesis, that the QE instructions enlist the nonlinear interactions across scales of the movement system without drawing the athlete's attention to the body.

Ultimately, this task- and anatomically-specific portrait of multifractal fluctuations rests on the even more generic facts that multifractal fluctuations spread across the body and that estimates of this spread predict the accuracy of the perceptual outcomes (Mangalam, Carver, et al., 2020a, 2020b). Hence, rather than inventorying body parts and task constraints prompting different values of multifractality, the longer view of investigating visually-guided aiming must examine the flow of multifractal fluctuations. Carver et al. (2017) found that multifractality spread from hip to head, as well as from head and hip toward the throwing hand. Indeed, in all these studies, the body appears to direct multifractal fluctuations towards those body parts most clearly engaged in the focal tasks. We might expect similar results in golf putting. QE training may promote this spread by reducing intentional movements of the head and eyes that might interfere with upstream flows of multifractality. The direction of attention towards a distant target in the visual field may depend on stabilizing the postural grasp of the surface underfoot and shunting multifractal fluctuations toward the head and eyes. Indeed, executive functions directing attention to specific features of the visual field benefits from a rich substrate of multifractal fluctuations in the head (Kelty-Stephen & Dixon, 2014), hand (Anastas et al., 2014; Stephen et al., 2012), and brain (Kardan, Adam, et al., 2020; Kardan, Layden, et al., 2020) that vary systematically with access to and mastery of the rules supporting task performance. When we consider that eye movements show multifractality (Kelty-Stephen & Mirman, 2013) and that variability in the monofractal structure of eye movements can predict visual attention (Wallot et al., 2015), we see early glimpses of a multifractal bridge from the postural-kinematic support to the visual support for QE.

The novelty of multifractal modeling to behavioral sciences can make this promise of a multifractal infrastructure supporting visually guided behaviors sound too ethereal and misty to be practical. But we can root this proposal in the known physiology spanning the very same bones, muscles, bones, joints, brain, and eyes whose coordination generates the winning golf putt. Specifically, the connective tissues composing the fascia operate to coordinate context-sensitive behavior across so wide a variety of scales (e.g. (Ingber, 2006)) that multifractality is one of the few current compelling frameworks for modeling how it might support perception-action (Schleip et al., 2014; Turvey & Fonseca, 2014). Examining how task constraints moderate a flow of multifractal fluctuations through the body could inform future explanations for QE training and other visual aiming tasks.

References

Allison, P. D. (1977). Testing for interaction in multiple regression. *American Journal of Sociology*, *83*(1), 144–153. https://doi.org/10.1086/226510

Anastas, J. R., Kelty-Stephen, D. G., & Dixon, J. A. (2014). Executive function as an interaction-dominant process. *Ecological Psychology*, *26*(4), 262–282. https://doi.org/10.1080/10407413.2014.957985

Ashkenazy, Y., M. Hausdorff, J., Ch. Ivanov, P., & Eugene Stanley, H. (2002). A stochastic model of human gait dynamics. *Physica A: Statistical Mechanics and Its Applications*, *316*(1), 662–670. https://doi.org/10.1016/S0378-4371(02)01453-X

Balasubramaniam, R., Riley, M. A., & Turvey, M. T. (2000). Specificity of postural sway to the demands of a precision task. *Gait & Posture*, *11*(1), 12–24. https://doi.org/10.1016/S0966-6362(99)00051-X

Barlow, H. B. (1963). Slippage of contact lenses and other artefacts in relation to fading and regeneration of supposedly stable retinal images. *Quarterly Journal of Experimental Psychology*, *15*(1), 36–51. https://doi.org/10.1080/17470216308416550

Bates, D., Sarkar, D., Bates, M., & Matrix, L. (2007). *The lme4 package* (R package version 2).

Bell, C., Carver, N., Zbaracki, J., & Kelty-Stephen, D. (2019). Nonlinear amplification of variability through interaction across scales supports greater accuracy in manual aiming: Evidence from a multifractal analysis with comparisons to linear surrogates in the Fitts task. *Frontiers in Physiology*, *10*, 998. https://doi.org/10.3389/fphys.2019.00998

Bernstein, N. A. (1967). *The Co-ordination and Regulation of Movement*. Pergamon Press.

Booth, C. R., Brown, H. L., Eason, E. G., Wallot, S., & Kelty-Stephen, D. G. (2018). Expectations on hierarchical scales of discourse: Multifractality predicts both short- and long-range effects of violating gender expectations in text reading. *Discourse Processes*, *55*(1), 12–30. https://doi.org/10.1080/0163853X.2016.1197811

Burton, G., & Turvey, M. T. (1990). Perceiving the lengths of rods that are held but not wielded. *Ecological Psychology*, *2*(4), 295–324. https://doi.org/10.1207/s15326969eco0204_1

Carello, C., Fitzpatrick, P., Domaniewicz, I., Chan, T.-C., & Turvey, M. T. (1992). Effortful touch with minimal movement. *Journal of Experimental Psychology: Human Perception and Performance*, *18*(1), 290–302. https://doi.org/10.1037/0096-1523.18.1.290

Carver, N. S., Bojovic, D., & Kelty-Stephen, D. G. (2017). Multifractal foundations of visually-guided aiming and adaptation to prismatic perturbation. *Human Movement Science*, *55*, 61–72. https://doi.org/10.1016/j.humov.2017.07.005

Chen, L.-C., Metcalfe, J. S., Chang, T.-Y., Jeka, J. J., & Clark, J. E. (2008). The development of infant upright posture: Sway less or sway differently? *Experimental Brain Research*, *186*(2), 293–303. https://doi.org/10.1007/s00221-007-1236-1

Chhabra, A., & Jensen, R. V. (1989). Direct determination of the $f(\alpha)$ singularity spectrum. *Physical Review Letters*, *62*(12), 1327–1330. https://doi.org/10.1103/PhysRevLett.62.1327

Cooke, A., Kavussanu, M., McIntyre, D., & Ring, C. (2010). Psychological, muscular and kinematic factors mediate performance under pressure. *Psychophysiology*, *47*(6), 1109–1118. https://doi.org/10.1111/j.1469-8986.2010.01021.x

Coppola, D., & Purves, D. (1996). The extraordinarily rapid disappearance of entopic images. *Proceedings of the National Academy of Sciences*, *93*(15), 8001–8004. https://doi.org/10.1073/pnas.93.15.8001

Davids, K., & Araújo, D. (2016). What could an ecological dynamics rationale offer Quiet Eye research? Comment on Vickers. *Current Issues in Sport Science (CISS)*, *1*, 104. https://doi.org/10.15203/CISS_2016.100

de Oliveira, R. F. (2016). Visual perception in expert action. In M Raab, B. Lobinger, S. Hoffman, A. Pizzera, & S. Laborde (Eds.), *Performance Psychology: Perception, Aaction, Cognition, Emotion* (pp. 254–270). Academic Press.

Diniz, A., Wijnants, M. L., Torre, K., Barreiros, J., Crato, N., Bosman, A. M. T., Hasselman, F., Cox, R. F. A., Van Orden, G. C., & Delignières, D. (2011). Contemporary theories of 1/f noise in motor control. *Human Movement Science*, *30*(5), 889–905. https://doi.org/10.1016/j.humov.2010.07.006

Ditchburn, R. W., & Ginsborg, B. L. (1952). Vision with a stabilized retinal image. *Nature*, *170*(4314), 36–37. https://doi.org/10.1038/170036a0

Dixon, J. A., Holden, J. G., Mirman, D., & Stephen, D. G. (2012). Multifractal dynamics in the emergence of cognitive structure. *Topics in Cognitive Science*, *4*(1), 51–62. https://doi.org/10.1111/j.1756-8765.2011.01162.x

Doyon, J. K., Hajnal, A., Surber, T., Clark, J. D., & Kelty-Stephen, D. G. (2019). Multifractality of posture modulates multisensory perception of stand-on-ability. *PLOS ONE*, *14*(2), e0212220. https://doi.org/10.1371/journal.pone.0212220

Feldman, A. G. (2018). Indirect, referent control of motor actions underlies directional tuning of neurons. *Journal of Neurophysiology*, *121*(3), 823–841. https://doi.org/10.1152/jn.00575.2018

Feldman, A. G., & Zhang, L. (2020). Eye and head movements and vestibulo-ocular reflex in the context of indirect, referent control of motor actions. *Journal of Neurophysiology*, *124*(1), 115–133. https://doi.org/10.1152/jn.00076.2020

Frenz, H., Bremmer, F., & Lappe, M. (2003). Discrimination of travel distances from 'situated' optic flow. *Vision Research*, *43*(20), 2173–2183. https://doi.org/10.1016/S0042-6989(03)00337-7

Friston, K., Breakspear, M., & Deco, G. (2012). Perception and self-organized instability. *Frontiers in Computational Neuroscience*, *6*, 44. https://doi.org/10.3389/fncom.2012.00044

Gallicchio, G., Cooke, A., & Ring, C. (2016). Lower left temporal-frontal connectivity characterizes expert and accurate performance: High-alpha T7-Fz connectivity as a marker of conscious processing during movement. *Sport, Exercise, and Performance Psychology*, *5*(1), 14–24. https://doi.org/10.1037/spy0000055

Gallicchio, G., Cooke, A., & Ring, C. (2017). Practice makes efficient: Cortical alpha oscillations are associated with improved golf putting performance. *Sport, Exercise, and Performance Psychology*, *6*(1), 89–102. https://doi.org/10.1037/spy0000077

Gallicchio, G., & Ring, C. (2020). The quiet eye effect: A test of the visual and postural-kinematic hypotheses. *Sport, Exercise, and Performance Psychology*, *9*(1), 143–159. https://doi.org/10.1037/spy0000162

Gibson, J. J. (1979). *The Ecological Approach to Visual Perception*. Houghton Mifflin.

Hajnal, A., Bunch, D. A., & Kelty-Stephen, D. G. (2014). Going for distance and going for speed: Effort and optical variables shape information for distance perception from observation to response. *Attention, Perception, & Psychophysics*, *76*(4), 1015–1035. https://doi.org/10.3758/s13414-014-0629-z

Hajnal, A., Bunch, D. A., & Kelty-Stephen, D. G. (2016). Pulling out all the stops to make the distance: Effects of effort and optical information in distance perception responses made by rope pulling. *Attention, Perception, & Psychophysics*, *78*(2), 685–699. https://doi.org/10.3758/s13414-015-1035-x

Hajnal, A., Clark, J. D., Doyon, J. K., & Kelty-Stephen, D. G. (2018). Fractality of body movements predicts perception of affordances: Evidence from stand-on-ability judgments about slopes. *Journal of Experimental Psychology: Human Perception and Performance*, *44*(6), 836–841. https://doi.org/10.1037/xhp0000510

Haken, H., Kelso, J. A. S., & Bunz, H. (1985). A theoretical model of phase transitions in human hand movements. *Biological Cybernetics*, *51*(5), 347–356. https://doi.org/10.1007/BF00336922

Harrison, S. J., & Stergiou, N. (2015). Complex adaptive behavior and dexterous action. *Nonlinear Dynamics, Psychology, and Life Sciences*, *19*(4), 345–394. https://pubmed.ncbi.nlm.nih.gov/26375932

Holt, K. G., Wagenaar, R. O., & Saltzman, E. (2010). A dynamic systems: Constraints approach to rehabilitation. *Brazilian Journal of Physical Therapy*, *14*, 446–463. https://doi.org/10.1590/S1413-35552010000600002

Houdijk, H., Brown, S. E., & van Dieën, J. H. (2015). Relation between postural sway magnitude and metabolic energy cost during upright standing on a compliant surface. *Journal of Applied Physiology*, *119*(6), 696–703. https://doi.org/10.1152/japplphysiol.00907.2014

834 Ihlen, E. A. F., & Vereijken, B. (2010). Interaction-dominant dynamics in human cognition: Beyond 1/
835  *f* fluctuation. *Journal of Experimental Psychology: General*, *139*(3), 436–463.
836  https://doi.org/10.1037/a0019098

837 Ingber, D. E. (2006). Cellular mechanotransduction: Putting all the pieces together again. *The FASEB*
838  *Journal*, *20*(7), 811–827. https://doi.org/10.1096/fj.05-5424rev

839 Ivanov, P. C., Nunes Amaral, L. A., Goldberger, A. L., Havlin, S., Rosenblum, M. G., Stanley, H. E.,
840  & Struzik, Z. R. (2001). From 1/*f* noise to multifractal cascades in heartbeat dynamics. *Chaos: An*
841  *Interdisciplinary Journal of Nonlinear Science*, *11*(3), 641–652.
842  https://doi.org/10.1063/1.1395631

843 Kardan, O., Adam, K. C. S., Mance, I., Churchill, N. W., Vogel, E. K., & Berman, M. G. (2020).
844  Distinguishing cognitive effort and working memory load using scale-invariance and alpha
845  suppression in EEG. *NeuroImage*, *211*, 116622.
846  https://doi.org/10.1016/j.neuroimage.2020.116622

847 Kardan, O., Layden, E., Choe, K. W., Lyu, M., Zhang, X., Beilock, S. L., Rosenberg, M. D., &
848  Berman, M. G. (2020). Scale-invariance in brain activity predicts practice effects in cognitive
849  performance. *BioRxiv*, 114959. https://doi.org/10.1101/2020.05.25.114959

850 Kello, C. T. (2018). Editor's introduction and review: Coordination and context in cognitive science.
851  *Topics in Cognitive Science*, *10*(1), 6–17. https://doi.org/10.1111/tops.12302

852 Kelty-Stephen, D. G. (2018). Multifractal evidence of nonlinear interactions stabilizing posture for
853  phasmids in windy conditions: A reanalysis of insect postural-sway data. *PLOS ONE*, *13*(8),
854  e0202367. https://doi.org/10.1371/journal.pone.0202367

855 Kelty-Stephen, D. G., & Dixon, J. A. (2014). Interwoven fluctuations during intermodal perception:
856  Fractality in head sway supports the use of visual feedback in haptic perceptual judgments by
857  manual wielding. *Journal of Experimental Psychology: Human Perception and Performance*,
858  *40*(6), 2289–2309. https://doi.org/10.1037/a0038159

859 Kelty-Stephen, D. G., Lee, I.-C., Carver, N. S., Newell, K. M., & Mangalam, M. (2020). Visual effort
860  moderates a self-correcting nonlinear postural control policy. *BioRxiv*, 209502.
861  https://doi.org/10.1101/2020.07.17.209502

862 Kelty-Stephen, D. G., & Mirman, D. (2013). Gaze fluctuations are not additively decomposable: Reply
863  to Bogartz and Staub. *Cognition*, *126*(1), 128–134.
864  https://doi.org/10.1016/j.cognition.2012.09.002

865 Kelty-Stephen, D. G., Stirling, L. A., & Lipsitz, L. A. (2016). Multifractal temporal correlations in
866  circle-tracing behaviors are associated with the executive function of rule-switching assessed by
867  the Trail Making Test. *Psychological Assessment*, *28*(2), 171–180.
868  https://doi.org/10.1037/pas0000177

Kelty-Stephen, D. G., & Wallot, S. (2017). Multifractality versus (mono-) fractality as evidence of
nonlinear interactions across timescales: Disentangling the belief in nonlinearity from the
diagnosis of nonlinearity in empirical data. *Ecological Psychology*, *29*(4), 259–299.
https://doi.org/10.1080/10407413.2017.1368355

Koslucher, F., Munafo, J., & Stoffregen, T. A. (2016). Postural sway in men and women during
nauseogenic motion of the illuminated environment. *Experimental Brain Research*, *234*(9), 2709–
2720. https://doi.org/10.1007/s00221-016-4675-8

Krueger, E., Schneider, A., Chavaillaz, A., Groner, R., Sawyer, B. D., Sonderegger, A., & Hancock, P.
A. (2019). Microsaccades distinguish looking from seeing. *Journal of Eye Movement Research*,
12(6), 2.

Latash, M. L. (2012). The bliss (not the problem) of motor abundance (not redundancy). *Experimental
Brain Research*, *217*(1), 1–5. https://doi.org/10.1007/s00221-012-3000-4

Latash, M. L. (2020). On primitives in motor control. *Motor Control*, *24*(2), 318–346.
https://doi.org/10.1123/mc.2019-0099

Lederman, S. J., Ganeshan, S. R., & Ellis, R. E. (1996). Effortful touch with minimum movement:
Revisited. *Journal of Experimental Psychology: Human Perception and Performance*, *22*(4), 851–
868. https://doi.org/10.1037/0096-1523.22.4.851

Lee, I.-C., Pacheco, M. M., & Newell, K. M. (2019). The precision demands of viewing distance
modulate postural coordination and control. *Human Movement Science*, *66*, 425–439.
https://doi.org/10.1016/j.humov.2019.05.019

Lee, J. T., & Kelty-Stephen, D. G. (2017). Cascade-driven series with narrower multifractal spectra
than their surrogates: Standard deviation of multipliers changes interactions across scales.
Complexity, *2017*, 7015243. https://doi.org/10.1155/2017/7015243

Li, R., Walter, H. J., & Stoffregen, T. A. (2020). The role of visual feedback about motion of the
ground on postural sway. *Journal of Motor Behavior*, *52*(3), 352–359.
https://doi.org/10.1080/00222895.2019.1627281

Liphart, J., Gallichio, J., Tilson, J. K., Pei, Q., Wu, S. S., & Duncan, P. W. (2015). Concordance and
discordance between measured and perceived balance and the effect on gait speed and falls
following stroke. *Clinical Rehabilitation*, *30*(3), 294–302.
https://doi.org/10.1177/0269215515578294

Lipsitz, L. A. (2002). Dynamics of stability: The physiologic basis of functional health and frailty. *The
Journals of Gerontology: Series A*, *57*(3), B115–B125. https://doi.org/10.1093/gerona/57.3.B115

Lipsitz, L. A., & Goldberger, A. L. (1992). Loss of "complexity" and aging: Potential applications of
fractals and chaos theory to senescence. *JAMA*, *267*(13), 1806–1809.
https://doi.org/10.1001/jama.1992.03480130122036

Mangalam, M., Carver, N. S., & Kelty-Stephen, D. G. (2020a). Global broadcasting of local fractal fluctuations in a bodywide distributed system supports perception via effortful touch. *Chaos, Solitons & Fractals*, *135*, 109740. https://doi.org/10.1016/j.chaos.2020.109740

Mangalam, M., Carver, N. S., & Kelty-Stephen, D. G. (2020b). Multifractal signatures of perceptual processing on anatomical sleeves of the human body. *Journal of The Royal Society Interface*, *17*(168), 20200328. https://doi.org/10.1098/rsif.2020.0328

Mangalam, M., & Kelty-Stephen, D. G. (2020). Multiplicative-cascade dynamics supports whole-body coordination for perception via effortful touch. *Human Movement Science*, *70*, 102595. https://doi.org/10.1016/j.humov.2020.102595

Mangalam, M., Lee, I.-C., Newell, K. M., & Kelty-Stephen, D. G. (2020). Visual effort moderates postural cascade dynamics. *BioRxiv*, 209486. https://doi.org/10.1101/2020.07.17.209486

Mark, L. S. (1987). Eyeheight-scaled information about affordances: A study of sitting and stair climbing. *Journal of Experimental Psychology: Human Perception and Performance*, *13*(3), 361–370. https://doi.org/10.1037/0096-1523.13.3.361

Mark, L. S., Balliett, J. A., Craver, K. D., Douglas, S. D., & Fox, T. (1990). What an actor must do in order to perceive the affordance for sitting. *Ecological Psychology*, *2*(4), 325–366. https://doi.org/10.1207/s15326969eco0204_2

Martinez-Conde, S., Macknik, S. L., Troncoso, X. G., & Dyar, T. A. (2006). Microsaccades counteract visual fading during fixation. *Neuron*, *49*(2), 297–305. https://doi.org/10.1016/j.neuron.2005.11.033

Meyer, M., Stiedl, O., & Kerman, B. (2003). Discrimination by multifractal spectrum estimation of human heartbeat interval dynamics. *Fractals*, *11*(02), 195–204. https://doi.org/10.1142/S0218348X03002063

Moore, L. J., Vine, S. J., Cooke, A., Ring, C., & Wilson, M. R. (2012). Quiet eye training expedites motor learning and aids performance under heightened anxiety: The roles of response programming and external attention. *Psychophysiology*, *49*(7), 1005–1015. https://doi.org/10.1111/j.1469-8986.2012.01379.x

Moore, L. J., Vine, S. J., Freeman, P., & Wilson, M. R. (2013). Quiet eye training promotes challenge appraisals and aids performance under elevated anxiety. *International Journal of Sport and Exercise Psychology*, *11*(2), 169–183. https://doi.org/10.1080/1612197X.2013.773688

Morales, C. J., & Kolaczyk, E. D. (2002). Wavelet-based multifractal analysis of human balance. *Annals of Biomedical Engineering*, *30*(4), 588–597. https://doi.org/10.1114/1.1478082

Morrison, A., McGrath, D., & Wallace, E. S. (2016). Motor abundance and control structure in the golf swing. *Human Movement Science*, *46*, 129–147. https://doi.org/10.1016/j.humov.2016.01.009

Munafo, J., Curry, C., Wade, M. G., & Stoffregen, T. A. (2016). The distance of visual targets affects
the spatial magnitude and multifractal scaling of standing body sway in younger and older adults.
Experimental Brain Research, *234*(9), 2721–2730. https://doi.org/10.1007/s00221-016-4676-7

Muñoz-Diosdado, A. (2005). A non linear analysis of human gait time series based on multifractal
analysis and cross correlations. *Journal of Physics: Conference Series*, *23*, 87–95.
https://doi.org/10.1088/1742-6596/23/1/010

Murnaghan, C. D., Carpenter, M. G., Chua, R., & Inglis, J. T. (2016). Keeping still doesn't "make
sense": Examining a role for movement variability by stabilizing the arm during a postural control
task. *Journal of Neurophysiology*, *117*(2), 846–852. https://doi.org/10.1152/jn.01150.2015

Oftadeh, R., Haghpanah, B., Vella, D., Boudaoud, A., & Vaziri, A. (2014). Optimal fractal-like
hierarchical honeycombs. *Physical Review Letters*, *113*(10), 104301.
https://doi.org/10.1103/PhysRevLett.113.104301

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Palatinus, Z., Dixon, J. A., & Kelty-Stephen, D. G. (2013). Fractal fluctuations in quiet standing predict
the use of mechanical information for haptic perception. *Annals of Biomedical Engineering*, *41*(8),
1625–1634. https://doi.org/10.1007/s10439-012-0706-1

Palatinus, Z., Kelty-Stephen, D. G., Kinsella-Shaw, J., Carello, C., & Turvey, M. T. (2014). Haptic
perceptual intent in quiet standing affects multifractal scaling of postural fluctuations. *Journal of
Experimental Psychology: Human Perception and Performance*, *40*(5), 1808–1818.
https://doi.org/10.1037/a0037247

Panchuk, D., & Vickers, J. N. (2006). Gaze behaviors of goaltenders under spatial–temporal
constraints. *Human Movement Science*, *25*(6), 733–752.
https://doi.org/10.1016/j.humov.2006.07.001

Park, H., & Turvey, M. T. (2008). Imperfect symmetry and the elementary coordination law. In A.
Fuchs & V. K. Jirsa (Eds.), *Coordination: Neural, Behavioral and Social Dynamics* (pp. 3–25).
Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-540-74479-5_1

Profeta, V. L. S., & Turvey, M. T. (2018). Bernstein's levels of movement construction: A
contemporary perspective. *Human Movement Science*, *57*, 111–133.
https://doi.org/10.1016/j.humov.2017.11.013

Raja, V., & Anderson, M. L. (2019). Radical embodied cognitive neuroscience. *Ecological Psychology*,
31(3), 166–181. https://doi.org/10.1080/10407413.2019.1615213

Rajachandrakumar, R., Mann, J., Schinkel-Ivy, A., & Mansfield, A. (2018). Exploring the relationship
between stability and variability of the centre of mass and centre of pressure. *Gait & Posture*, *63*,
254–259. https://doi.org/10.1016/j.gaitpost.2018.05.008

Renshaw, I., Davids, K., Araújo, D., Lucas, A., Roberts, W. M., Newcombe, D. J., & Franks, B.
(2019). Evaluating weaknesses of "perceptual-cognitive training" and "brain training" methods in
sport: An ecological dynamics critique. *Frontiers in Psychology*, *9*, 2468.
https://www.frontiersin.org/article/10.3389/fpsyg.2018.02468

Reynolds, A. M. (2015). Liberating Lévy walk research from the shackles of optimal foraging. *Physics*
of Life Reviews, *14*, 59–83. https://doi.org/10.1016/j.plrev.2015.03.002

Rienhoff, R., Tirp, J., Strauß, B., Baker, J., & Schorer, J. (2016). The 'quiet eye' and motor
performance: A systematic review based on Newell's constraints-led model. *Sports Medicine*,
46(4), 589–603. https://doi.org/10.1007/s40279-015-0442-4

Schleip, R., Mechsner, F., Zorn, A., & Klingler, W. (2014). The bodywide fascial network as a sensory
organ for haptic perception. *Journal of Motor Behavior*, *46*(3), 191–193.
https://doi.org/10.1080/00222895.2014.880306

Schlesinger, M. F., & Klafter, J. (1986). Lévy walks versus Lévy flights. In H. E. Stanley & N.
Ostrowski (Eds.), *Growth and Form* (Stanley, H, pp. 279–283). Martinus Nijhoff.

Schmitt, D. T., & Ivanov, P. C. (2007). Fractal scale-invariant and nonlinear properties of cardiac
dynamics remain stable with advanced age: A new mechanistic picture of cardiac control in
healthy elderly. *American Journal of Physiology-Regulatory, Integrative and Comparative*
Physiology, *293*(5), R1923–R1937. https://doi.org/10.1152/ajpregu.00372.2007

Schreiber, T., & Schmitz, A. (1996). Improved surrogate data for nonlinearity tests. *Physical Review*
Letters, *77*(4), 635–638. https://doi.org/10.1103/PhysRevLett.77.635

Serres, J. R., & Ruffier, F. (2017). Optic flow-based collision-free strategies: From insects to robots.
Arthropod Structure & Development, *46*(5), 703–717. https://doi.org/10.1016/j.asd.2017.06.003

Shannon, C. E. (1948). A mathematical theory of communication. *The Bell System Technical Journal*,
27(3), 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x

Shimizu, Y. U., Thurner, S., & Ehrenberger, K. (2002). Multifractal spectra as a measure of complexity
in human posture. *Fractals*, *10*(01), 103–116. https://doi.org/10.1142/S0218348X02001130

Shine, J. M., Aburn, M. J., Breakspear, M., & Poldrack, R. A. (2018). The modulation of neural gain
facilitates a transition between functional segregation and integration in the brain. *ELife*, *7*,
e31130. https://doi.org/10.7554/eLife.31130

Stephen, D., Anastas, J., & Dixon, J. (2012). Scaling in cognitive performance reflects multiplicative
multifractal cascade dynamics. *Frontiers in Physiology*, *3*, 102.
https://doi.org/10.3389/fphys.2012.00102

Stephen, D. G., & Mirman, D. (2010). Interactions dominate the dynamics of visual cognition.
Cognition, *115*(1), 154–165. https://doi.org/10.1016/j.cognition.2009.12.010

Stoffregen, T. A. (1985). Flow structure versus retinal location in the optical control of stance. *Journal
of Experimental Psychology: Human Perception and Performance*, *11*(5), 554–565.
https://doi.org/10.1037/0096-1523.11.5.554

Stoffregen, T. A. (1986). The role of optical velocity in the control of stance. *Perception &
Psychophysics*, *39*(5), 355–360. https://doi.org/10.3758/BF03203004

Stoffregen, T. A., Mantel, B., & Bardy, B. G. (2017). The senses considered as one perceptual system.
Ecological Psychology, *29*(3), 165–197. https://doi.org/10.1080/10407413.2017.1331116

Stoffregen, T. A., & Riccio, G. E. (1990). Responses to optical looming in the retinal center and
periphery. *Ecological Psychology*, *2*(3), 251–274. https://doi.org/10.1207/s15326969eco0203_3

Stoffregen, T. A., Yang, C.-M., & Bardy, B. G. (2005). Affordance judgments and nonlocomotor body
movement. *Ecological Psychology*, *17*(2), 75–104. https://doi.org/10.1207/s15326969eco1702_2

Teng, D. W., Eddy, C. L., & Kelty-Stephen, D. G. (2016). Non-visually-guided distance perception
depends on matching torso fluctuations between training and test. *Attention, Perception, &
Psychophysics*, *78*(8), 2320–2328. https://doi.org/10.3758/s13414-016-1213-5

Turvey, M T. (1977). Contrasting orientations to the theory of visual information processing.
Psychological Review, *84*(1), 67–88. https://doi.org/10.1037/0033-295X.84.1.67

Turvey, Michael T, & Fonseca, S. T. (2014). The medium of haptic perception: A tensegrity
hypothesis. *Journal of Motor Behavior*, *46*(3), 143–187.
https://doi.org/10.1080/00222895.2013.798252

van Smeden, M., de Groot, J. A. H., Moons, K. G. M., Collins, G. S., Altman, D. G., Eijkemans, M. J.
C., & Reitsma, J. B. (2016). No rationale for 1 variable per 10 events criterion for binary logistic
regression analysis. *BMC Medical Research Methodology*, *16*(1), 163.
https://doi.org/10.1186/s12874-016-0267-3

Vaz, D. V., Silva, P. L., Mancini, M. C., Carello, C., & Kinsella-Shaw, J. (2017). Towards an
ecologically grounded functional practice in rehabilitation. *Human Movement Science*, *52*, 117–
132. https://doi.org/10.1016/j.humov.2017.01.010

Vickers, J. N. (1992). Gaze control in putting. *Perception*, *21*(1), 117–132.
https://doi.org/10.1068/p210117

Vickers, J. N. (1996). Visual control when aiming at a far target. *Journal of Experimental Psychology:
Human Perception and Performance*, *22*(2), 342–354. https://doi.org/10.1037/0096-1523.22.2.342

Vickers, J. N. (2009). Advances in coupling perception and action: the quiet eye as a bidirectional link
between gaze, attention, and action. In Markus Raab, J. G. Johnson, & H. R. B. T.-P. in B. R.
Heekeren (Eds.), *Mind and Motion: The Bidirectional Link between Thought and Action* (Vol.
174, pp. 279–288). Elsevier. https://doi.org/10.1016/S0079-6123(09)01322-3

Vickers, J. N. (2011). Mind over muscle: The role of gaze control, spatial cognition, and the quiet eye
in motor expertise. *Cognitive Processing*, *12*(3), 219–222. https://doi.org/10.1007/s10339-011-
0411-2

Vine, S. J., Lee, D. H., Walters-Symons, R., & Wilson, M. R. (2017). An occlusion paradigm to assess
the importance of the timing of the quiet eye fixation. *European Journal of Sport Science*, *17*(1),
85–92. https://doi.org/10.1080/17461391.2015.1073363

Vine, S. J., Lee, D. O. N., Moore, L. E. E. J., & Wilson, M. R. (2013). Quiet eye and choking: Online
control breaks down at the point of performance failure. *Medicine & Science in Sports & Exercise*,
45(10), 1988–1994. https://doi.org/10.1249/MSS.0b013e31829406c7

Vine, S., Moore, L., & Wilson, M. (2011). Quiet eye training facilitates competitive putting
performance   in   elite   golfers.   *Frontiers   in   Psychology*,   *2*,   8.
https://doi.org/0.3389/fpsyg.2011.00008

Wagenmakers, E.-J., van der Maas, H. L. J., & Farrell, S. (2012). Abstract concepts require concrete
models: Why cognitive scientists have not yet embraced nonlinearly coupled, dynamical, self-
organized critical, synergistic, scale-free, exquisitely context-sensitive, interaction-dominant,
multifractal,   interdependent   b.   *Topics   in   Cognitive   Science*,   *4*(1),   87–93.
https://doi.org/10.1111/j.1756-8765.2011.01164.x

Wallot, S., O'Brien, B., Coey, C. A., & Kelty-Stephen, D. (2015). Power-law fluctuations in eye
movements predict text comprehension during connected text reading. In D. C. Noelle, R. Dale,
A. S. Warlaumont, J. Yoshimi, T. Matlock, & C. D. Jennings (Eds.), *Proceedings of the 37th*
Annual Meeting of the Cognitive Science Society (pp. 2583–2588).

Warren, W. H. (1984). Perceiving affordances: Visual guidance of stair climbing. *Journal of*
Experimental   Psychology: Human   Perception   and   Performance,   *10*(5),   683–703.
https://doi.org/10.1037/0096-1523.10.5.683

West, B. J., & Scafetta, N. (2003). Nonlinear dynamical model of human gait. *Physical Review E*,
67(5), 51917. https://doi.org/10.1103/PhysRevE.67.051917

Williams, A. M. (2016). Quiet eye vs. noisy brain: The eye like the brain is always active–comment on
Vickers. *Current Issues in Sport Science (CISS)*, *1*, 116. https://doi.org/10.15203/CISS_2016.116

Witt, J. K., & Proffitt, D. R. (2008). Action-specific influences on distance perception: A role for motor
simulation. *Journal of Experimental Psychology: Human Perception and Performance*, *34*(6),
1479–1492. https://doi.org/10.1037/a0010781

Table 1. Training instructions given to the quiet-eye and technical groups during the training phase.

quiet-eye (QE) training instructions	technical training instructions
assume your stance, and ensure your gaze is located on the back of the ball	take your stance with your legs shoulder-width apart
after setting up over the ball, fix your gaze on the hole	set your position so that your head is

	directly above the ball looking down
make no more than three fixations towards the hole	keep your clubhead square to the ball
your final fixation should be a quiet eye on the back of the ball. The onset of the quiet eye should occur before the stroke begins and last for 2 to 3s	allow your arms and shoulders to remain loose
ensure you direct gaze towards the clubhead during the putting stroke	the putting action should be pendulum-like, making sure that you accelerate through the ball
the quiet eye should remain on the green for 200 to 300ms after the club contacts the ball	after contact, follow through but keep your head still and facing down

1073 Source: Adapted from (Moore et al., 2013).

1074 **Table 2.** Subset of Poisson regression model including main and interaction effects non-multifractal
1075 and multifractal measures of sway, perturbation of eye height, and time

predictor	b	s.e.m.	p^*
effects of non-multifractal measures of sway			
(see Section 3.2.1.)			
MSD_{acc}	-8.80×10^0	1.67×10^{-1}	**< 0.0001**
$Mean_{disp}$	-2.91×10^2	1.33×10^1	**< 0.0001**
SD_{disp}	4.43×10^0	1.64×10^0	**< 0.01**
$MSD_{acc} \times Mean_{disp}$	1.97×10^4	1.00×10^1	**< 0.0001**
$MSD_{acc} \times SD_{disp}$	-3.11×10^3	1.03×10^1	**< 0.0001**
$Mean_{disp} \times SD_{disp}$	2.56×10^4	1.05×10^1	**< 0.0001**
$MSD_{acc} \times Mean_{disp} \times SD_{disp}$	-1.04×10^5	1.08×10^1	**< 0.0001**
effects of multifractal measures of sway and interactions with time			
(see Section 3.2.2.)			
W_{MF}	1.80×10^0	1.29×10^{-1}	**< 0.0001**
t_{MF}	2.21×10^{-4}	1.05×10^{-3}	0.83
$W_{MF} \times t_{MF}$	-5.93×10^{-3}	8.27×10^{-3}	0.48
$t_{MF} \times trial_{block}(Linear)$	1.36×10^{-2}	9.82×10^{-3}	0.17
$t_{MF} \times trial_{block}(Quadratic)$	-8.93×10^{-2}	1.15×10^{-2}	**< 0.0001**
$t_{MF} \times trial_{block}(Cubic)$	-1.08×10^{-2}	1.14×10^{-2}	0.34
interactions of multifractal measures of sway with perturbation			
(see Section 3.2.3.)			
$QE \times W_{MF}$	-1.04×10^{-0}	2.03×10^{-1}	**< 0.0001**
$QE \times t_{MF}$	-1.85×10^{-3}	1.30×10^{-3}	0.16
$QE \times W_{MF} \times t_{MF}$	-4.59×10^{-3}	1.09×10^{-2}	0.67
$perturbation \times W_{MF}$	-1.50×10^0	3.12×10^{-1}	**< 0.0001**
$perturbation \times t_{MF}$	4.57×10^{-3}	2.85×10^{-3}	0.11
$perturbation \times W_{MF} \times t_{MF}$	1.54×10^{-1}	3.15×10^{-2}	**< 0.0001**
$QE \times perturbation \times W_{MF}$	2.12×10^0	4.91×10^{-1}	**< 0.0001**
$QE \times perturbation \times t_{MF}$	8.62×10^{-3}	4.18×10^{-3}	**< 0.05**
$QE \times perturbation \times W_{MF} \times t_{MF}$	-2.21×10^{-1}	4.97×10^{-2}	**< 0.0001**
interactions of multifractal measures with perturbation and time			
(see Section 3.2.4.)			
$QE \times trial_{block}(Linear) \times t_{MF}$	-3.73×10^{-2}	1.46×10^{-2}	**< 0.05**
$QE \times trial_{block}(Quadratic) \times t_{MF}$	-1.85×10^{-1}	1.66×10^{-2}	**< 0.0001**
$QE \times trial_{block}(Cubic) \times t_{MF}$	1.13×10^{-1}	1.73×10^{-2}	**< 0.0001**
$perturbation \times trial_{block}(Linear) \times t_{MF}$	3.83×10^{-1}	5.28×10^{-2}	**< 0.0001**
$perturbation \times trial_{block}(Quadratic) \times t_{MF}$	4.07×10^{-1}	5.20×10^{-2}	**< 0.0001**
$perturbation \times trial_{block}(Cubic) \times t_{MF}$	5.39×10^{-1}	4.45×10^{-2}	**< 0.0001**
$QE \times perturbation \times trial_{block}(Linear) \times t_{MF}$	-2.71×10^{-1}	6.11×10^{-2}	**< 0.0001**
$QE \times perturbation \times trial_{block}(Quadratic) \times t_{MF}$	5.24×10^{-1}	6.14×10^{-2}	**< 0.0001**
$QE \times perturbation \times trial_{block}(Cubic) \times t_{MF}$	-7.90×10^{-1}	5.79×10^{-2}	**< 0.0001**

1076 *boldfaced values indicate statistical significance at the alpha level of 0.05.

1077 **Table 3.** Logistic regression modeling of successful shots.

predictor	b	s.e.m.	p^*
intercept	−2.85	0.48	**< 0.0001**
QE	0.20	0.30	0.51
perturbation	−0.22	0.43	0.60
QE × perturbation	−1.16	0.65	0.07
block	0.24	0.13	**0.06**
RMS_{Acc}	4.05	1.34	**< 0.01**
W_{MF}	−7.25	2.94	**< 0.05**

1078 *boldfaced values indicate statistical significance at the alpha level of 0.05.

1079   **Figure 1.** Schematic of Chhabra and Jensen's (Chhabra & Jensen, 1989) method used to estimate
1080   multifractal spectrum $f(\alpha)$ (Part-1). See Section 2.4.1. for further details.

Figure 2. Schematic of Chhabra and Jensen's (Chhabra & Jensen, 1989) method used to estimate multifractal spectrum $f(\alpha)$ (Part-2). See Section 2.4.1. for further details.

option A: multifractal spectrum width reflects strictly linear properties of the original series

option B: multifractal spectrum width reflects effects unfloding across multiple, nested scales

snip   *snip*   *snip*   *snip*

samples [1/100 Hz]

shuffle data to destroy nonlinearity but preserve linearity (overall form and short-lag relationships)

flip   *flip*   *flip*   *flip*   *flip*

samples [1/100 Hz]

null hypothesis: multifractal spectrum width of original series does not differ from that of surrogate series

use one-sample *t*-test to compare original series' multifractal spectrum width to those of 30 surrogate series

if one-sample *t*-test **is not** significantly large, fail to reject null: **series is linear**

if one-sample *t*-test **is** significantly large, fail to reject null: **series exhibits cascades**

1083 **Figure 3.** Surrogate analysis used to identify whether a series exhibits multifractality due to
1084 nonlinearity (cascades). See Section 2.4.2. for further details.

Figure 4. Multifractal analysis. (*a*, *b*) Representative original accelerometer displacement series. (*c*, *d*) Representative IAAFT surrogate series of the original accelerometer displacement series in *a* and *b*. (*e*, *f*) Multifractal spectrum (*f*(*α*), *α*) of original series in *a* and *b*, and those of their five representative IAAFT surrogates.

Figure 5. The observed trial-by-trial radial error and trial-by-trial model predictions on the same axis for four participants each from the technical (top four panels) and QE-trained (bottom four panels) groups, respectively. The representation includes two participants each with the highest (left panels) and lowest (right panels) Pearson's correlation coefficient r indicating the strength of relationship between the observed and predicted radial errors.

1094   **Figure 6.** Trial-by-trial model predictions of radial error for the two participant groups. (*a*) Technical-
1095   trained group. (*b*) QE-trained group.