Running title: Dynamics of transcriptional programs and chromatin

accessibility in mouse spermatogonial cells from early postnatal to adult life

Irina Lazar-Contes[1,2]*, Deepak K. Tanwar[1,2]*, Rodrigo G. Arzate-Mejia[1,2]*, Leonard

C. Steg[1,2], Olivier Ulrich Feudjio[3], Marion Crespo[3], Pierre-Luc Germain[1,2], Isabelle M.

Mansuy[1,2#]

1. Laboratory of Neuroepigenetics, Brain Research Institute, Medical Faculty of the

University of Zurich and Institute for Neuroscience, Department of Health Science

and Technology of the ETH Zurich, Zurich, Switzerland

2. Center for Neuroscience Zurich, ETH and University Zurich

3. ADLIN Science, Pépinière «Genopole Entreprises», Evry, France

*Equal contribution

[#]Correspondence: mansuy@hifo.uzh.ch

Abstract

In mammals, spermatogonial cells (SCs) are undifferentiated male germ cells in

testis quiescent until birth that self-renew and differentiate to produce spermatogenic

cells and functional sperm across life. The transcriptome of SCs is highly dynamic

and timely regulated during postnatal development. We examined if such dynamics

involves changes in chromatin organization by profiling the transcriptome and

chromatin accessibility in SCs from early postnatal stages to adulthood in mice using

RNA-seq and ATAC-seq. By integrating transcriptomic and epigenomic features, we

show that SCs undergo massive chromatin remodeling during postnatal

development that correlates with distinct gene expression profiles and transcription

factors (TF) motif enrichment. We identify genomic regions with significantly

different chromatin accessibility in adult SCs that are marked by histone

modifications associated with enhancers and promoters. Some of the regions with

increased accessibility correspond to transposable element subtypes enriched in

multiple TFs motifs and close to differentially expressed genes. Our results

underscore the dynamics of chromatin organization in developing germ cells and the

involvement of the regulatory genome.

Introduction

Spermatogonial cells (SCs) are the initiators and supporting cellular foundation of spermatogenesis in testis in many species including mammals. In mice, SCs become active one to two days after birth, when they exit mitotic arrest and start dividing to populate the basement membrane of seminiferous tubules. During the first week of postnatal life, a population of SCs continues to proliferate to give rise to undifferentiated A_{single} (A_s), A_{paired} (A_{pr}) and $A_{aligned}$ (A_{al}) cells. The remaining SCs differentiate to form chains of daughter cells that become primary and secondary spermatocytes around postnatal day (PND) 10 to 12. Spermatocytes then undergo meiosis and give rise to haploid spermatids that develop into spermatozoa. Spermatozoa are then released in the lumen of seminiferous tubules and continue to mature in the epididymis until becoming capable of fertilization by PND 42-48 (De Rooij, 2017; Kubota and Brinster, 2018; Oatley and Griswold, 2017).

Recent work showed that SCs in early postnatal life have distinct transcriptional signatures (Green et al., 2018; Hammoud et al., 2014; Hermann et al., 2018; Law et al., 2019). During the first week of postnatal development, SCs display unique features necessary for the rapid establishment and expansion of the cell population along the basement membrane. This includes high expression of genes involved in cell cycle regulation, stem cell proliferation, transcription and RNA processing (Grive et al., 2019). In comparison, genes expressed in adult SCs are involved in the maintenance of a balance between proliferation and differentiation, and help constitute a steady cell population that ensures sperm formation across life. Previous transcriptome analyses have revealed that adult SCs prioritize pathways related to paracrine signaling and niche communication, as well as mitochondrial functions and oxidative phosphorylation (Grive et al., 2019; Hermann et al., 2018). Epigenetic changes such as histone tail modifications and DNA methylation have also been reported in SCs during postnatal development (Hammoud et al., 2014; Hammoud et al., 2015). Today however, the relationship between the transcriptome dynamics and chromatin landscape in SCs during the transition from early postnatal to adult stage has not been fully characterized.

We characterized the transcriptome and the profile of chromatin accessibility in SCs
during the transition from early postnatal to adult stages. The results reveal
extensive changes in transcription and in chromatin accessibility in particular, an
increase in chromatin accessibility at enhancer regions in adult SCs compared to
postnatal SCs. Regions with changes in chromatin accessibility are enriched in
binding motifs of several TFs that are differentially expressed between postnatal and
adult stages, suggesting a possible role in changes in chromatin accessibility.
Analyses of chromatin accessibility at transposable elements (TEs) identified
previously uncharacterized changes at long terminal repeats (LTR) and LINE L1
subtypes between developing and adult SCs. Together, these findings suggest a
functional link between transcriptional dynamics and chromatin accessibility in SCs
during development, and underscore the plasticity of genome organization in germ
cells.

Results

FACS enriches SCs collected from postnatal and adult mouse testis

We collected testes from 8- and 15-day old pups (PND8 and 15) and adult males
and prepared cell suspensions from each animal by enzymatic digestion. The
preparations were enriched for SCs by fluorescence-activated cell sorting (FACS)
using specific surface markers (Kubota et al., 2004) (Fig. 1A, B). The purity of sorted
cells was evaluated by immunocytochemistry using PLZF (ZBTB16), a well-
established marker for undifferentiated SCs (Costoya et al., 2004). FACS enriched
cell populations from 3-6% PLZF+ (before FACS) to 85-95% PLZF+ (Fig. 1B),
suggesting a high SCs enrichment.

To validate the molecular identity of enriched SCs populations, we profiled their
transcriptome at PND8, PND15 and adulthood by total RNA sequencing (RNA-seq)
(n=6 for each group) and examined the expression of known spermatogonial stem
cell and somatic markers (Leydig and Sertoli cells) (Fig. 1C and Fig. S1A, B).
Classical SCs markers such as *c-kit* (Schrans-Stassen et al., 1999)*, Id4* (Helsel et
al., 2017; Sun et al., 2015), *Lin28a* (Chakraborty et al., 2014; Wang et al., 2020),
Zbtb16 (Costoya et al., 2004; Song et al., 2020) and *Gfra1* (He et al., 2007) were
robustly expressed in all SCs samples at each developmental stage (Fig. 1C and
Fig. S1A, B), indicating high purity of sorted cells. In contrast, low to negligible

expression of somatic cells markers such as *Vim* (Bernardino et al., 2018) and
Tspan17 (Gewiss et al., 2021) for Sertoli cells, and *Fabp3* and *Hsd3b1* for Leydig
cells (Sararols et al., 2021) (Fig. 1C) was detected. To further validate the samples
purity, we manually curated a list of spermatogonial and somatic cell markers
derived from recent single-cell RNA-seq datasets (Cao et al., 2021; Sararols et al.,
2021) and determined their expression level in our datasets. Again, we detected
robust and consistent expression of all reported spermatogonial markers and low
expression of somatic cells markers in all samples at each developmental stage (Fig.
S1A). These results confirm the efficiency of our FACS-based enrichment method.
104

Dynamic transcriptomic states characterize postnatal and adult SCs
We examined the transcriptome dynamics of SCs from postnatal to adult stages by
identifying differentially expressed genes (DEGs) in the RNA-seq datasets. We used
a stringent cut-off (adjusted-P≤0.05, abs Log_2 fold change (FC)≥1) on 17,000 genes
expressed during at least one developmental stage (See Methods). A total of 663
DEGs were identified between PND8 and PND15 (146 down-regulated and 517 up-
regulated) and 2483 DEGs between PND15 and adult stage (914 down-regulated
and 1579 up-regulated) (Fig. 1D-E and Table S1). Consistent with previous reports
(Hammoud et al., 2015), we observed a dynamic regulation of germ cell factors,
transcription factors (TF) involved in core pluripotency pathways and signaling
molecules important for self-renewal (Fig. S1B). For instance, *Pou5f1 (Oct4)*, a TF
necessary for pluripotency (Nichols et al., 1998), is significantly down-regulated in
adult SCs while the TFs *Klf4* and *Sox2*, also needed for pluripotency (An et al.,
2019), are expressed similarly in postnatal and adult stages although at different
level (i.e *Klf4* is expressed more than *Sox2*) (Fig. S1B).
120

We conducted Gene Ontology (GO) enrichment analyses to identify the biological
processes which DEGs are involved in. GO analyses showed that DEGs between
PND8 and PN15 are involved in cell differentiation, cilium movement, sperm motility
and transcription and include for instance, TFs such as *Junb*, *Hoxb6*, *Cebpa* and
Pparg (Fig. 1F and Table S2). Further, genes coding for histone proteins such as
histone H2B *Hist2h3b* and epigenetic modifiers like the methyltransferase *Pygo1* and
histone acetyl-transferase *Ncoa1* are also differentially expressed between PND8
and 15. Interestingly, genes down- or up-regulated between postnatal stages are

involved in different processes. While down-regulated genes are implicated in cell

differentiation, cell migration and regulation of proliferation for instance, *Cdkn1a* that

is down-regulated at PND15 regulates genetic diversity during spermatogenesis

(Kanatsu-Shinohara et al., 2022), up-regulated genes are rather involved in germ cell

development, cell signalling, insulin secretion and transcription (Fig. S1C and Table

S2). Further, DEGs between PND15 and adulthood have roles in reproduction,

spermatogenesis, cell differentiation, DNA replication, glycolysis and extracellular

matrix (ECM)-receptor interaction pathways (Fig. 1G, Fig. S1D and Table S2). For

example, the DNA methyltransferase *Dnmt1*, necessary for genome regulation

(Edwards et al., 2017) and *Col4a2,* a subunit of type IV collagen and component of

the basal membrane (Reissig et al., 2019), are specifically down-regulated in adult

SCs (Fig. S1C). Interestingly, expression of collagen has been associated to a high

proliferative potential and the ability to form germ cell colonies in SCs (He et al.,

2005), suggesting that regulation of collagen genes in adult SCs may decrease

germ-stem cell potential. Genes up-regulated in adult SCs are involved in cilium

movement, germ cell development and glycolysis (Table S2).

We then examined the differences and similarities of transcriptional profiles across

the three developmental stages. To identify unique or common changes in gene

expression during the transition from early postnatal to adult stages, we compared

the list of DEGs at PND8 versus PND15 and at PND15 versus adulthood.

Remarkably, the vast majority of DEGs show significant stage-specific changes in

transcription (Fig. S2A). For instance, 75% (495/663) of DEGs between PND8 and

PND15 are not statistically changed in the transition to adult SCs (Fig. S2B and

Table S1 and S2). Similarly, 93% (2325/2493) of DEGs in adult SCs are not

statistically changed when compared to PND8 versus PND15 SCs (Fig. 1H and

Table S1 and S2). GO enrichment analyses showed that adult-specific down-

regulated genes are involved in cell differentiation, cell migration, regulation of signal

transduction and Wnt signalling, regulation of DNA replication and transcription as

well as collagen biosynthesis and laminin interactions while adult-specific up-

regulated genes are involved in sexual reproduction, male gamete generation, germ

cell development, cilium movement and glycolysis (Fig. 1H and Table S1 and S2).

Interestingly, a small fraction of all DEGs (4.9%) are detected as significantly
changed in the same direction and magnitude (adjusted-P≤0.05, absLog$_2$FC≥1)
across the three developmental stages (Fig. S2C and Table S1 and S2). For
instance, 121 genes are consistently up-regulated from PND8 to adult stage, while
just 26 are consistently down-regulated from PND8 to adult stage (Fig. S2C). Such
DEGs are involved in different biological pathways like chromatin organization (Table
S2). In particular, the histone gene clusters displayed significant down-regulation
across postnatal development and in adulthood (Table S2).

170   Landscape of chromatin accessibility in SCs during postnatal development

Cellular differentiation is generally accompanied by changes in chromatin
accessibility at regulatory elements (Atlasi et al., 2017). We examined how chromatin
accessibility in SCs is modified during development using omni-ATAC-seq (Corces
et al., 2017). We focused on SCs at PND15 and adulthood, stages that showed the
highest changes in gene expression (Fig. 1). Accessible regions were identified by
peak-calling on merged nucleosome-free fragments (NFF) as proxy for genomic
regions with potential regulatory activity. We identified 158,977 peaks with clear
ATAC-seq signal compared with surrounding genomic regions (Fig. S3A and Table
S3). Most accessible regions were located at distal intergenic regions (38%), introns
(28%) and putative promoter regions (23%) (Fig. S3B, C) and encompassed
sequences enriched for histone PTMs associated with active chromatin such as
H3K4 mono-, di- and tri-methylation (Fig. S2D). Notably, signal enrichment was
higher for H3K4 methylation than for other histone PTMs (Fig. S3D). These results
are consistent with previous observations that ATAC-seq peaks are identified at
regulatory elements such as enhancers and promoters and are preferentially located
at genomic regions with nucleosomes carrying H3K4me (Henikoff et al., 2020).

We then examined differences in chromatin accessibility between PND15 and adult
SCs by differential accessibility analysis. 3212 differentially accessible regions
(DARs) were identified (adjusted-P≤0.05, absLog$_2$FC≥1) with a total of 760 regions
with decreased chromatin accessibility (DARs-down) and 2452 regions with
increased accessibility (DARs-up) in adult SCs (Figure 2A, B and Table S3). DARs
were predominantly localized in distal intergenic regions (54% DARs-down, 37%
DARs-up) and introns (33% DARs-down, 36% DARs-up) with a minority located in

putative promoter regions (8% DARs-down, 12% DARs-up), consistent with the
genomic distribution of all detected ATAC-seq peaks (Fig. 2C). GO analyses of the
closest genes assigned to each DAR showed that these genes are involved in male
gonad development, cell adhesion, sex differentiation and regulation of cell
communication among others (Fig. 2D, E).

Epigenomic annotation of DARs using high-quality publicly available ChIP-seq
datasets for postnatal SCs (Cheng et al., 2020) revealed that DARs are
predominantly located in regions enriched for histone PTMs associated with
enhancers and promoters. 51% of regions with increased accessibility are located at
genomic loci that carry histone PTMs since early postnatal stages, and about half
(48%) overlap with regions significantly enriched in H3K4me1, a histone PTM
associated to enhancers. In contrast, 80% of regions with decreased accessibility
are located in regions with H3K4me1 and 1/3 (33%) also overlap with the repressive
histone PTM, H3K27me3 (Fig. 2F). Interestingly, while 16% of regions with
increased accessibility are located at potential active regulatory elements that carry
H3K27ac, none of the regions with decreased accessibility overlap with H3K27ac
(Fig. 2F). Therefore, our data indicate that the transition from postnatal to adult SCs
is accompanied by discrete, yet robust, changes in chromatin accessibility at
potential regulatory elements, suggesting their involvement in the control of gene
transcription.

DARs are associated with binding sites for distinct families of TFs

We examined the relationship between differences in chromatin accessibility and TF
binding by TF motif analysis using DARs as input. We observed that regions with
decreased accessibility in adult SCs are enriched for binding motifs of members of
specific TF families such as KLF (KLF2, KLF5, KLF11, KLF12, KLF15, KLF16), SP
(SP1-5, SP9), FOXO (FOXO1, FOXO3), ETS (ETV1-6) among others (Fig. 3A and
Table S4). Expression analyses showed that many of these TFs are differentially
expressed in postnatal or adult SCs (adjusted-P<0.05, 10 down-regulated and 11 up-
regulated) (Fig. 3B, C). For example, KLF11,12 and 15, TFs that regulate stem cell
maintenance and development (Bialkowska et al., 2017), are upregulated in adult
SCs compared to PND15 (Fig. 3C). These TFs and their motif match top enriched
motifs detected in DARs in adult SCs (Fig. 3A). The SP family of TFs also show

differential expression and enrichment of binding motifs in regions with decreased
accessibility. SP5, which has its lowest expression in adult SCs (Fig. 3B, C),
promotes self-renewal of mESCs by directly regulating *Nanog* (Tang et al., 2017) .

Members of the FOXO family of TFs also have an enrichment of motifs in regions
with decreased accessibility. While FOXO3 has increased expression from PND15
onwards, FOXO1 is decreased in adult compared to postnatal SCs (Fig. 3B, C).
Interestingly, FOXO1 and FOXO3 regulate spermatogonial stem cell function and
maintenance in mouse (Goertz et al., 2011). Finally, ETS-type of TFs also show
differential expression during SCs development and their motif is enriched in regions
with decreased accessibility. ELK4 is up-regulated in adult SCs and can act as a
transcriptional repressor via recruitment of the HDAC Sirt7 and deacetylation of
H3K18ac (Fig. 3B-D) (Barber et al., 2012). EKL4 can also act as transcriptional
activator of immediate early genes such as *c-Fos* (Dalton and Treisman, 1992).
Interestingly, *c-Fos* itself codes for a TF important for the regulation of proliferation
(He et al., 2008) and shows a trend towards up-regulation in adult SCs (Table S1). In
contrast, GABPA is downregulated in adult SCs compared to PND15, and has been
involved in the regulation of proliferation in mESCs (Ueda et al., 2017).

Regions with decreased chromatin accessibility also show an enrichment for the
binding motif of the TF DMRT1. *Dmrt1* is progressively repressed during SCs
postnatal development and is the lowest in adult SCs (Fig 3B, C and E). DMRT1 can
act either as a repressor or activator and controls testis development and male germ
cell proliferation (Zhang et al., 2016). DMRT1 can inhibit meiosis and promote
mitosis in SCs by repressing Stra8 (Matson et al., 2010). Consistently, we observed
transcriptional repression of *Stra8* in adult SCs (Table S1).

Next, we identified motifs overrepresented in regions with increased chromatin
accessibility in adult SCs. The identified motifs correspond to binding motifs of
members of specific TF families such as POU (POU2F2, POU1F1, POU5F1), RFX
(RXF1-7), DMRT (DMRT1, DMRT3, DMRTA1-2, DMRTC2), SOX (SOX2, SOX3,
SOX5, SOX13), NFY (NFYA, NFYB, NFYC) and AP-1 (JUN, JUNB, JUND, FOS,
ATF3, BATF) (Fig. 3F and Table S4). A subset of them is also differentially
expressed (adjusted-P<0.05, 13 down-regulated and 5 up-regulated). The most
overrepresented motif is similar to the binding motif of the POU family of TFs, which

are critical regulators of stem cells (Nichols et al., 1998). *Pou1f1* and *Pou5f1* are transcriptionally repressed in adult SCs while *Pou2f2* is maximally expressed in PND15 SCs and down-regulated in adult cells (Fig. 3G, H). We also identified motifs for members of the RFX family of TFs, which are master regulators of ciliogenesis (Choksi et al., 2014) implicated in regulation of neural stem cells (Kawase et al., 2014). RFX2 is robustly expressed in adult SCs (Fig. 3G, H and I) and has been reported to induce the expression of ciliary genes in association with the TF FOXJ, which has a trend towards up-regulation in adult SCs (Fig. S4) (Quigley and Kintner, 2017). Interestingly, ciliary genes are among the top genes specifically up-regulated in adult SCs (Fig. 1H), suggesting a regulatory relationship between RFX2 and ciliary genes expression in adult SCs.

Other binding motifs enriched at regions with increased accessibility correspond to members of the NF-Y complex, NF-YA, NF-YB and NF-YC. In mESCs, NF-Y TFs facilitate a permissive chromatin conformation and play an important role in the expression of core ESC pluripotency genes (Oldfield et al., 2014). NF-YA/B motif enrichment has also been found in regions of open chromatin in human SCs (Guo et al., 2017). Interestingly, the expression of NF-YA and NF-YC is progressively down-regulated starting at PND15 (Fig. 3G, H). We also detected an enrichment for the binding motifs of members of the AP-1 family of TFs, which are involved in many processes, from regulation of cell proliferation to differentiation and acute responses to environmental clues (Bejjani et al., 2019; Vierbuchen et al., 2017). Interestingly, except for *Fosl2*, other TFs do not show any significant change in transcription in postnatal or adult SCs and are either constitutively expressed or repressed. For instance, c-FOS and JUN are robustly expressed in postnatal and adult SCs (Fig. 3J) consistent with their role in promoting the proliferative potential of spermatogonial stem cells (He et al., 2008; Wang et al., 2018). Together, these data reveal that a shift in TFs repertoire accompanies chromatin reorganization occurring in SCs during postnatal to adult development.

A subset of DARs is associated with differential gene expression

Thus far, our results show that in SCs, the transition from postnatal to the adult stages is accompanied by changes in gene expression and chromatin accessibility.

In some instances, differences in gene expression can be correlated with differences

in chromatin accessibility of regulatory elements. Since DARs in postnatal and adult

SCs overlap with putative enhancer and promoter elements, we examined if these

changes in chromatin accessibility correlate with changes in gene expression. First,

we tested if promoter regions of DEGs (+/-2.5kb from TSS) have differential

chromatin accessibility by identifying all possible ATAC-seq peaks located in

promoters and calculating an average accessibility signal. No significant difference

was detected for down-regulated genes in adult SCs, despite a trend for decreased

accessibility around the TSS (Fig. 4A). However, down-regulated genes had a

significant increase in chromatin accessibility in their body (Fig. 4B). Up-regulated

genes had a significant increase in chromatin accessibility both at the promoter

region and gene body (Fig. 4A, B). Then, we conducted motif enrichment analyses

and identified families of TFs with overrepresented binding motifs at the promoter of

DEGs. Binding motifs for members of KLF and SP family of TFs were

overrepresented in the promoter of both down- and up-regulated DEGs, while motifs

for members of TF families SOX and NFY were detected only in the promoter of

down-regulated genes. Promoters of up-regulated genes were specifically enriched

in motifs for RFX and AP-1 (Fig. 4C). These observations suggest that changes in

chromatin accessibility, in particular at the promoter of up-regulated DEGs may be

associated with differential binding of RFX and AP-1 TFs.

Next, we examined the overlap between DARs and the promoter of DEGs.

Unexpectedly, only 3.2% of all promoters of DEGs overlap with at least one DAR

(Fig. 4D). The majority of overlapping DARs (71%) is associated with promoters of

up-regulated genes in adult SCs (Fig. 4D). For example, *Gpx4*, a peroxidase

involved in the control of stemness (Hu et al., 2021), is transcriptionally active in

adult SCs and has an open chromatin at its promoter region (Fig. 4E). Since the

majority of DARs are located at distal intergenic regions and have histone PTMs

associated with active regulatory elements, we extended our search of their target

genes by assigning a large region (+/-100kb) around each DEG. We observed that a

third of DEGs were associated with at least one DAR (Fig. 4F), with the majority of

DARs (69%) overlapping regulatory elements of up-regulated genes. For instance,

Braf is transcriptionally up-regulated in adult SCs, which correlates with increased

chromatin accessibility at distal regulatory elements (Fig. 4G). Our results overall

show that only a fraction of DEGs is associated with DARs, suggesting that changes

in gene expression are not always mirrored at the level of chromatin accessibility,

consistent with recent observations (Kiani et al., 2022).

327 Chromatin accessibility at transposable elements (TEs) undergoes remodeling

328 during the transition from early postnatal to adulthood in SCs

TEs are tightly regulated in the germline by coordinated epigenetic mechanisms

involving DNA methylation, chromatin silencing and PIWI proteins-piRNA pathway

(Deniz et al., 2019). SCs were recently shown to have a unique landscape of

chromatin accessibility at long-terminal repeats (LTRs) retrotransposons, compared

with other stages of germ cells in testis (Sakashita et al., 2020). We examined

chromatin accessibility at TEs in PND15 and adult SCs by quantifying ATAC-seq

reads overlapping TEs defined by UCSC RepeatMasker then analyzing differential

accessibility at subtypes level (see Methods section). These analyses showed that

SCs transitioning from PND15 to adult stages have significantly different chromatin

accessibility at 135 TEs subtypes (adjusted-P≤0.05, absLog$_2$FC≥1) (Fig. 5A, B and

Table S5). Most TEs subtypes have decreased chromatin accessibility (93/135,

68,9%) (Fig. 5A) and 42 have increased accessibility (Fig. 5B) in adult SCs.

Differentially accessible TEs loci are predominantly located in intergenic (68%) and

intronic (25%) regions and 6% re close to a gene (+/-1kb from TSS) (Fig. 5C).

LTR retrotransposons were the most abundant TEs with altered chromatin

accessibility, specifically ERVK and ERV1 subtypes (Fig. 5A, B). Exemplary ERVK

subtypes with less accessible chromatin included RLTR17, RLTR9A3, RLTR12B and

RMER17B (Table S5). Enrichment of RLTR17 and RLTR9 repeats was previously

reported in mESCs, specifically at TFs important for pluripotency maintenance such

as *Oct4* and *Nanog* (Fort et al., 2014). Interestingly, we identified the promoter

region of the long non-coding RNA (lncRNA) *Lncenc1*, an important regulator of

pluripotency in mESCs (Fort et al., 2014; Sun et al., 2018), to harbor several LTR

loci with decreased accessibility in adult SCs. One of these loci, RLTR17, overlap

with the TSS of *Lncenc1*, and its decreased accessibility correlates with markedly

lower *Lncenc1* expression in adult SCs (Fig. 5D). *Lncenc1* (also known as *Platr18*) is

part of the pluripotency-associated transcript (*Platr*) family of lncRNAs recently

identified as potential regulators of pluripotency-associated genes *Oct4*, *Nanog* and

Zfp42 in mESCs (Bergmann et al., 2015). We also identified several other *Platr*

genes such as *Platr27* and *Platr14*, whose TSS overlap with LTRs with reduced

accessibility, RLTR17 and RLTR16B_MM, respectively (Fig. 5D and Table S5).

These two pluripotency-associated transcripts tended to be downregulated in adult

SCs but were unchanged at PND8 and PND15 (Fig. 5D and Table S1). The other

LTR subtypes with decreased accessibility in adult SCs belong to ERV1, ERVL and

MaLR families (Fig. 5A). A few non-LTR TEs also had decreased chromatin

accessibility particularly 7 DNA element subtypes, 2 satellite subtypes and 1 LINE

subtype, respectively (Table S5).

TE subtypes with increased accessibility mostly included members of ERVK, ERVL

and ERV1 families (24/42, 57.1%) (Table S5). Interestingly, many LINE L1 subtypes

had increased chromatin accessibility in adult SCs (Table S5). When parsing the

data for more accessible loci within L1 subtypes, we found several L1 loci situated

less than +/-5kb from the TSS of olfactory (*Olfr*) genes, particularly *Olfr* gene clusters

on chromosome 2, 7 and 11 (Table S5).

TEs are known to provide tissue-specific substrates for TF binding (Sundaram and

Wysocka, 2020). We examined the regulatory potential of differentially accessible

LTR subtypes by determining the enrichment of TF motifs in these regions. We

grouped LTR subtypes by family (EVK, ERV1, ERVL and ERVL-MaLR families) and

observed that among families with less accessible chromatin, ERVKs had the

highest number of enriched TF motifs in adult SCs (Table S5). Top hits included TFs

with known regulatory functions in cell proliferation and differentiation such as

FOXL1 and FOXQ1, stem cell maintenance factors ELF1, EBF1 and THAP11 and

TFs important for spermatogenesis PBX3, ZNF143 and NFYA/B (Table S5). ERVLs

displayed motif enrichment for very few TFs, among which ETV2, recently reported

to be the spermatogonial stem cell factor ZBTB7A and the testis-specific CTCF

paralog CTCFL (Table S5) (Green et al., 2018).

More accessible LINE L1 subtypes were highly enriched in TF motifs, particularly in

multiple members of ETS, E2F and FOX families (Table S5). The most significant

motifs belonged to spermatogonial stem cell maintenance and stem cell potential

regulators FOXO1 and ZEB1, as well as TFs recently associated with active

enhancers of the stem cell-enriched population of SCs such as ZBTB17 and KLF5

(Table S5) (Cheng et al., 2020). More accessible ERV1s were also enriched in

several TF binding sites, including spermatogenesis-related TFs (PBX3, PRDM1,
NFYA/B), hypoxia inducible HIF1A and cytokine regulators STAT5A/B, suggesting
different metabolic demand between postnatal and adult stage (Table S5).

Discussion

This study examines the transcriptomic landscape and chromatin accessibility of
mouse SCs from early postnatal development to adulthood. Using FACS-enriched
SCs populations at PND8, PND15 and adulthood, it shows that the transcriptome of
SCs is dynamically regulated during development and that many factors including
pluripotency-related TFs and signaling molecules are differentially expressed. This
correlates with changes in chromatin accessibility although not all DEGs have
altered accessibility. TEs such as LTR retrotransposons have modified chromatin
accessibility, which in some cases affects pluripotency-associated lncRNAs. TF motif
enrichment in differentially accessible TEs suggested various regulatory roles in
SCs.

Different transcriptional states characterize the postnatal development of SCs

SCs are known to undergo genome-wide transcriptional changes during
development (Grive et al., 2019; Hammoud et al., 2015; Hermann et al., 2018). The
present data complement previous findings by showing that postnatal maturation of
SCs is accompanied by an overall decrease in the expression of pluripotency
markers and differential expression of signaling molecules and regulators of cell
cycle and spermatogenesis. The high quality and depth of our RNA-seq datasets
allowed us to uncover specific transitional states of SCs postnatal maturation that
were not known before. For instance, we observed that while the transition from
early to mid-postnatal stages (PND8 to PND15) is accompanied by changes in gene
expression, the transition from mid-postnatal to adult stages is more extensive, both
in number of regulated genes and magnitude of changes. We also observed a clear
trend towards up-regulation of gene expression as postnatal maturation proceeds,
that may reflect gradual transcriptional increase for some genes and possibly sharp
changes at certain developmental stages for others.

Consistently, many transcriptional regulators and chromatin modifiers are
differentially regulated during SCs development. Different families of TFs including
factors with a specific role in SCs like RFX and DMRT, and factors with more
widespread functions across cell types such as members of the AP-1 family are
modulated, suggesting the contribution of multiple classes of TFs to SCs
development. Notably, several transcriptional repressors and epigenetic silencers
are differentially down-regulated which may explain the marked trend towards
transcriptional up-regulation in our datasets (Fig. 1D). Interestingly, we also
observed a global down-regulation of histone genes starting at PND15 (Table S1),
which may modify the composition and structure of chromatin and be associated with
chromatin openness (Prado et al., 2017). Overall, our transcriptional data highlight
the yet unappreciated dynamic regulation of TFs and epigenetic/chromatin modifiers
in SCs during postnatal development.

Chromatin accessibility in SCs is modified at enhancers during postnatal
development

Our results suggest that the transition of SCs from early postnatal to adult life is
accompanied by changes in chromatin accessibility. In particular, accessibility is
increased in regions enriched in H3K4me1, a histone PTM typically associated with
primed enhancers, but also to a lesser extent, in regions enriched in H3K27ac, a
mark associated with active regulatory elements (Gasperini et al., 2020; Shlyueva et
al., 2014). This suggests that the transition of SCs from postnatal to adult stages
involves changes in the regulatory genome, in particular at primed enhancers.
Enhancer priming has recently been proposed as an important mechanism for SCs
differentiation (McCarrey, 2023). Differences in chromatin accessibility may be
partially explained by differential binding of TFs to regulatory elements (Klemm et al.,
2019). This may involve at least three non-mutually exclusive events: 1) a change in
the abundance of TFs, 2) a change in the epigenetic status of TF binding sites due to
different amount and/or activity of epigenetic modifiers, and 3) a change in
nucleosome positioning around TF binding sites due to different amount and/or
activity of chromatin remodelers. For instance, CpG methylation and nucleosome
positioning can influence TF binding at regulatory elements and affect chromatin
accessibility (Barisic et al., 2019; Kaluscha et al., 2022). The fact that *Dnmt1* is
downregulated in adult SCs concomitantly with differential expression of TFs and

chromatin modifiers may explain changes in chromatin accessibility at potential
enhancers.

Modest correspondence between transcription and chromatin accessibility in SCs

Despite robust changes in gene expression in adult SCs, only a subset of DARs
could be assigned to DEGs. This suggests that transcriptional regulation and
chromatin accessibility at regulatory elements are not always linked and can be
dissociated, as previously reported (Chereji et al., 2019; Kiani et al., 2022). This
implies that chromatin accessibility may be modified without any effects on
transcription and conversely, transcription can be activated or repressed without
requiring any change in chromatin accessibility. In turn, it is possible that a repressor
can be exchanged with an activator at a regulatory element but have no detectable
consequences. It is also possible that changes in chromatin accessibility have no
immediate effects on transcription but reflect an intermediate state that primes a
gene or genomic locus for later activation or repression such as observed with
developmentally- or experience-dependent primed enhancers (Marco et al., 2020;
Rada-Iglesias et al., 2011). Consistently, the majority of DARs with increased
accessibility overlap with regions enriched in H3K4me1, a histone PTM characteristic
of primed enhancers thought to control future transcriptional responses in different
cell types (Rada-Iglesias et al., 2011). These regions may be important for
responses to external cues in adult SCs. The apparent lack of correspondence
between transcription and accessibility may also be technical and due to the
stringency of filters we used to assign genomic loci to DARs. We cannot exclude the
possibility that modest changes in accessibility (<log2FC1) correspond to changes in
TF occupancy. Chromatin accessibility has been suggested to not be the primary
determinant of chromatin-mediated gene regulation (Chereji et al., 2019). Such
possibility could be addressed by examining genome-wide maps of TFs, cofactors
and RNA-Pol II occupancy that would provide more accurate information about the
regulatory genome and its relationship with observed gene expression programs.

Chromatin accessibility at TEs during SCs maturation
Our results reveal differences in chromatin accessibility and TFs motif landscape at
different TEs subtypes between PND15 and adult SCs. ERVK and ERV1 subtypes
were the most abundant categories of TEs that become less accessible in adult SCs,
whilst LINE L1 subtypes gained accessibility. Although the majority of these TEs

reside in intergenic and intronic regions, we could detect specific loci belonging to
differentially accessible ERVK and LINE L1 subtypes localized nearby TSS of
distinct gene families. Notably, ERVKs are known to play an important role in the
regulation of mRNA and lncRNAs transcription during spermatogenesis (Davis et al.,
2017; Sakashita et al., 2020). The landscape of chromatin accessibility at LTRs loci
in SCs is known to be unique compared to other mitotic germ cells and meiotic
gametes in testis (Sakashita et al., 2020).

RLTR17, one of the LTR subtypes with decreased chromatin accessibility in adult
SCs overlaps with the TSS of several downregulated lncRNAs from the *Platr* family.
Platr genes, including *Lncenc1* and *Platr14* identified in our study*,* are LTR-
associated lncRNAs important for gene expression programs in ESCs (Bergmann et
al., 2015). Interestingly, RLTR17 has also previously been linked to pluripotency
maintenance. In mESCs, it is highly expressed and enriched in open chromatin
regions and has been shown to provide binding sites for the pluripotency factors
Oct4 and Nanog (Fort et al., 2014). Therefore, we suggest that RLTR17 chromatin
organization may play a role in regulating pluripotency programs between early
postnatal and adult SCs.

In contrast to decreased accessibility at LTRs loci, LINE L1 subtype had increased
accessibility in adult SCs. Some of these L1 loci were located close to olfactory
receptor genes with upregulated mRNA expression. Recent findings in mouse and
human ESCs suggest a non-random genomic localization of L1 elements,
specifically at genes encoding proteins with specialized functions (Lu et al., 2020).
Among them, the *Olfr* gene family was the most enriched in L1 elements (Lu et al.,
2020). Although their role in SCs is currently not established, Olfr proteins have been
implicated in swimming behaviour of sperm cells (Fukuda and Touhara, 2005). Given
their dynamic regulation in SCs, we speculate that *Olfr* genes play additional roles in
spermatogenesis, other than in sperm physiology. Together with the high number of
enriched TF motifs identified at differentially accessible ERVKs and LINE L1
elements, these data highlight a regulatory role for chromatin organization of TEs in
SCs during the transition from development to adult stage (Sundaram et al., 2014;
Sundaram and Wysocka, 2020).

Limitations

One limitation of our study is the partial purity of SCs populations obtained by FACS, due to the fact that it does not fully remove other testicular cells from the samples. This therefore does not exclude the possible influence of contaminating cells on the data and their interpretation. Such influence may explain differences between our datasets and datasets in the literature. Nevertheless, by comparing chromatin landscape in developing and adult SCs, the study reveals an age-dependent dynamic reorganization of chromatin accessibility in these cells. When integrated with our transcriptomic datasets and published histone PTMs profiles, the data provide novel insight into transcriptome-chromatin dynamics of mouse SCs from early postnatal life to adulthood. This represents an important resource for future studies on mouse germ cells.

Methods

Mouse husbandry

Male C57Bl/6J mice were purchased from Janvier Laboratories (France) and bred in-house to C57Bl/6J primiparous females to generate males for the experiments. All animals were kept on a reversed 12-h light/12-h dark cycle in a temperature- and humidity-controlled facility with food (M/R Haltung Extrudat, Provimi Kliba SA, Switzerland) and water provided *ad libitum*. Cages were changed once weekly. Animals from 2 independent cohorts generated separately were used for the experiments.

Germ cells isolation

Germ cells were isolated at postnatal day (PND) 8 or 15 for RNA-seq and ATAC-seq, and at postnatal week 20 (adult) for ATAC-seq. Testicular single-cell suspensions were prepared as previously described with slight modifications (Kubota et al., 2004a, 2004b). For PND8 and PND15 cells, testes from 2 animals were pooled for each sample while for adults, testes from individual males were used. Pup testes were collected in sterile HBSS on ice. Tunica albuginea was gently removed from each testis, making sure to keep seminiferous tubules as intact as possible. Tubules were enzymatically digested in 0.25% trypsin-EDTA (ThermoFisher Scientific) and 7mg/ml DNase I (Sigma-Aldrich) solution for 5 min at 37oC. The

suspension was vigorously pipetted up and down 10 times and incubated again for 3
min at 37oC. The digestion was stopped by adding 10% fetal bovine serum
(ThermoFisher Scientific) and cells were passed through a 20μm-pore-size cell
strainer (Miltenyi Biotec) and pelleted by centrifugation at 600g for 7 min at 4oC.
Cells were resuspended in PBS-S (PBS with 1% PBS, 10 mM HEPES, 1 mM
pyruvate, 1mg/ml glucose, 50 units/ml penicillin and 50 μg/ml streptomycin) and
used for sorting. Adult testes were collected and the tunica was removed.
Seminiferous tubules were digested in 1 mg/mL collagenase type IV (Sigma Aldrich)
and then in 0.25 % Trypsin-EDTA (Gibco), both times for 8-12 minutes and in the
presence of DNase I solution (Sigma Aldrich). FBS (Cytiva HyClone) was added to a
concentration of 10 % and the cell suspension was filtered through a 40μm-pore-size
cell strainer (Corning) and washed with DPBS-S (1 % FBS, 10 mM HEPES (Gibco),
1mM sodium pyruvate (Gibco), 1mg/mL Glucose (Gibco) and 1X Penicillin-
Streptomycin (Gibco) in DPBS (Gibco). 5 mL of the single cell suspension was then
overlayed on 2 mL 30 % Percoll (Sigma) and centrifuged with disabled break at 600g
for 8 minutes at 4 °C. The supernatant was removed and cell suspension was
washed twice in DPBS-S and used for sorting.

SCs enrichment by FACS

For pup testis, dissociated cells were stained with BV421-conjugated anti-β2M,
biotin-conjugated anti-THY1 (53-2.1) and PE-conjugated anti-αv-integrin (RMV-7)
antibodies. THY1 was detected by staining with Alexa Fluor 488-Sav. For adult
testes, cells were stained with anti-α6-integrin (CD49f; GoH3), BV421-conjugated
anti-β2 microglobulin (β2M; S19.8) and R-phycoerythrin (PE)-conjugated anti-THY1
(CD90.2; 30H-12) antibodies. α6-Integrin was detected by Alexa Fluor 488-SAv after
staining with biotin-conjugated rat anti-mouse IgG1/2a (G28-5) antibody. Prior to
FACS, 1 μg/ml propidium iodide (Sigma) was added to cell suspensions to
discriminate dead cells. All antibody incubations were performed in PBS-S for at
least 30 min at 4oC followed by washing in PBS-S. Antibodies were obtained from
BD Biosciences (San Jose, United States) unless otherwise stated. Cell sorting was
performed at 4oC on a FACS Aria III 5L using an 85μm nozzle at the Cytometry
Facility of University of Zurich. For RNA-seq at PND8 and PND15, cells were
collected in 1.5 ml Eppendorf tubes in 500 μL PBS-S, immediately pelleted by
centrifugation and snap frozen in liquid N2. Cell pellets were stored at -80oC until

RNA extraction. For RNA-seq of adults, 1000 spermatogonial cells (MHC I negative,
alpha-6-integrin and Thy1 positive) per male were sorted into PBS. For Omni-ATAC
at PND15, 25'000 cells were collected in a separate tube, pelleted by centrifugation
and immediately processed using a library preparation protocol (Corces et al., 2017).
For Omni-ATAC in adults, 5000 cells from each animal were collected in a separate
tube and processed using the same protocol.

Immunocytochemistry

The protocol used for assessing SCs enrichment after sorting was kindly provided by
Jon Oatley at Washington State University, Pullman, USA (Yang et al., 2013).
Briefly, 30,000-50,000 cells were adhered to poly-L-Lysine coated coverslips
(Corning Life Sciences) in 24-well plates for 1 h. Cells were fixed in freshly prepared
4% PFA for 10 min at room temperature then washed in PBS with 0.1% Triton X-100
(PBS-T). Non-specific antibody binding was blocked by incubation with 10% normal
goat serum for 1 h at room temperature. Cells were incubated overnight at 4oC with
mouse anti-PLZF (0.2 µg/ml, Active Motif) primary antibody. Alexa488 goat anti-
mouse IgG (1 µg/mL, ThermoFisher Scientific) was used for secondary labelling at
4oC for 1 h. Coverslips were washed 3 times, mounted onto glass slides with
VectaShield mounting medium containing DAPI (Vector Laboratories) and examined
by fluorescence microscopy. SCs enrichment was determined by counting PLZF+
cells in 10 random fields of view from each coverslip and dividing by the total number
of cells present in the same fields of view (DAPI-stained nuclei). The number of
PLZF+ and PLZF-cells from the 10 different fields of view was averaged.

RNA extraction and RNA-seq library preparation

For RNA-seq at PND8 and PND15, total RNA was extracted from sorted cells using
AllPrep RNA/DNA Micro kit (Qiagen). RNA quality was assessed using a Bioanalyzer
2100 (Agilent Technologies). Samples were quantified using Qubit RNA HS Assay
(ThermoFisher Scientific). RNA sequencing libraries were prepared using SMARTer
Stranded Total RNA-Seq Kit v3 (Takara Bio USA, Inc) following the recommended
protocol with minor adjustments. For PND8 and PND15 SCs libraries, RNA was
fragmented for 4 minutes at 94 °C. After reverse transcription, samples were
barcoded using SMARTer Unique Dual Index Kit (Takara Bio USA, Inc). Five PCR
cycles were run and PCR products were purified using AMPure XP reagent

(Beckman Coulter Life Sciences; bead:sample ratio 0.8X). Ribosomal cDNA was
depleted following the protocol and final library amplification was performed with 12
PCR cycles. Libraries were purified twice using AMPure XP reagent (bead:sample
ratio 1X) and eluted in Tris buffer. Adult SCs libraries were prepared using the option
to start directly from cells (1000 cells as input). Cells were incubated for 6 minutes at
85 °C and processed as indicated for PND8 and PND15 SCs. Samples were pooled
at equal molarity, as determined in a 100-800 bp window on the Bioanalyzer.

Omni-ATAC library preparation and sequencing

Chromatin accessibility was profiled from PND15 and adult SCs. Libraries were
prepared according to Omni-ATAC protocol, starting from 25,000 PND15 and 5000
adult sorted SCs (Corces et al., 2017). Briefly, sorted cells were lysed in cold lysis
buffer (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% NP40, 0.1%
Tween-20, and 0.01% digitonin) and nuclei were pelleted and transposed using
Nextera Tn5 (Illumina) for 30 min at 37oC in a thermomixer with shaking at 1000
rpm. Transposed fragments were purified using MinElute Reaction Cleanup Kit
(Qiagen). Following purification, libraries were generated by PCR amplification using
NEBNext High-Fidelity 2X PCR Master Mix (New England Biolabs) and purified
using Agencourt AMPure XP magnetic beads (Beckman Coulter) to remove primer
dimers (78bp) and fragments of 1000-10,000 bp. Library quality was assessed on an
Agilent High Sensitivity DNA chip using the Bioanalyzer 2100 (Agilent Technologies).
6 samples were sequenced from PND15 and 5 from adult SCs.

RNA-seq analysis

Quality control and alignment: 100bp single end sequencing was performed at the
Functional Genomics Center Zurich (FGCZ) using a Novaseq SP flowcell on the
Novaseq 6000 platform. Quality assessment of FASTQ files was done using FastQC
(Andrews et al., 2012) (version 0.11.8). TrimGalore (Krueger, 2015) (version 0.6.2)
was used to trim adapters and low-quality ends from reads with Phred score less
than 30 (-q 30) and to discard trimmed reads shorter than 30bp (--length 30).
Trimmed reads were pseudo-aligned using Salmon (Patro et al., 2017) (version
0.9.1) with automatic detection of the library type (-l A), correcting for sequence-
specific bias (--seqBias) and correcting for fragment GC bias correction (--gcBias) on
a transcript index prepared for the Mouse genome (GRCm38) from GENCODE

(version M18) (Harrow et al., 2012), with additional piRNA precursors and TEs

(concatenated by family) from Repeat Masker as in (Gapp et al., 2018).

Downstream analysis: Data analyses and plotting were conducted with R (R Core

Team, 2020) (version 3.6.2) using packages from The Comprehensive R Archive

Network (CRAN) (https://cran.r-project.org) and Bioconductor (Huber et al., 2015).

Pre-filtering of genes was done using the filterByExpr function from edgeR

(Robinson et al., 2009) (version 3.28.1) with a design matrix requiring at least 15

counts (min.counts=15). Normalization factors were obtained using TMM

normalization (Robinson and Oshlack, 2010) from edgeR package and differential

gene expression analyses were done using limma-voom (Law et al., 2014) pipeline

from limma (Ritchie et al., 2015) (version 3.42.2). GO enrichment analyses were

performed using g:Profiler (https://biit.cs.ut.ee/gprofiler/gost) querying against Mus

musculus database.

659 ATAC-seq analysis

Quality control, alignment, and peak calling: Paired-end (PE) sequencing was

performed on PND15 and adult SCs on Illumina HiSeq2500 platform at FGCZ.

FASTQ files were assessed for quality using FastQC (Andrews et al., 2012) (version

0.11.8). QC was performed using TrimGalore (Krueger, 2015) (version 0.6.2) in PE

mode (--paired), trimming adapters, low-quality ends (-q 30) and discarding reads

shorter than 30bp after trimming (--length 30). Alignment on GRCm38 genome was

performed using Bowtie2 (Langmead and Salzberg, 2012) (version 2.3.5) with the

following parameters: allowing fragments up to 2kb to align (-X 2000), entire read

alignment (--end-to-end), suppressing unpaired alignments for paired reads (--no-

mixed), suppressing discordant alignments for paired reads (--no-discordant) and

minimum acceptable alignment score with respect to read length (--score-min L,-

0.4,-0.4). Using alignmentSieve (version 3.3.1) from deepTools (Ramirez et al.,

2016) (version 3.4.3), aligned data (BAM files) were adjusted for read start sites to

represent the center of the transposon cutting event (--ATACshift), and filtered for

reads with a high mapping quality (--minMappingQuality 30). Reads mapping to the

mitochondrial chromosome and ENCODE blacklisted regions were filtered out. To

call nucleosome-free regions, all aligned files were merged within groups (PND15

and adult), sorted and indexed using SAMtools (Li et al., 2009) (version 0.1.19) and

nucleosome-free fragments (NFFs) were obtained by selecting alignments with a

template length between 40 and 140bp in length. Peak calling on NFFs was

performed using MACS2 (Zhang et al., 2008) (version 2.2.7.1) with mouse genome

size (-g 2744254612) and PE BAM file format (-f BAMPE).

Differential accessibility analysis: These analyses were conducted in R (version

3.6.2) using packages from CRAN (https://cran.r-project.org) and Bioconductor

(Huber et al., 2015). Peaks were annotated based on overlap with GENCODE

(version M18) (Harrow et al., 2012) transcript and/or distance to the nearest TSS

(https://github.com/mansuylab/SC_postnatal_adult/bin/annoPeaks.R). The number

of extended reads overlapping in peak regions was calculated using csaw package

(Lun and Smyth, 2015) (version 1.20.0). Peak regions that did not have at least 15

reads in at least 40% of the samples were filtered out. Normalization factors were

obtained on filtered peak regions using TMM normalization method (Robinson and

Oshlack, 2010) and differential analysis on peaks (PND15 versus adult) was

performed using Genewise Negative Binomial Generalized Linear Models with

Quasi-likelihood (glmQLFit) Tests from edgeR package (Robinson et al., 2009)

(version 3.28.1). Peak regions with an absLog2FC≥1 and adjusted-P≤0.05 were

categorized as DARs.

Downstream analysis: Heatmaps of normalized ATAC-seq signal were created using

deepTools with default parameters. Genomic annotation of ATAC-seq peak and

DARs and identification of closest gene DARs were performed using ChIPpeakAnno

(Zhu et al., 2010). GO enrichment analysis of genes associated to DARs was

performed using g:Profiler (Raudvere et al., 2019) querying against the *Mus*

musculus database. For the epigenomic annotation of DARs, publicly available

signal files and peak files for all histone marks were used (Cheng et al., 2020).

Heatmaps of ChIP-seq signal for histone PTMs around ATAC-seq peaks and DARs

were generated with deepTools. Overlap between ChIP-seq peaks and DARs was

generated using Intervene (Khan et al., 2017). TF motif analysis was performed

using MEME-ChIP (Ma et al., 2014) and motif identification was done querying

identified motifs against JASPAR database (Castro-Mondragon et al., 2021).

Quantification and identification of differences in chromatin accessibility between

promoter regions of DEGs was performed with deepStats (Richard 2019). To

quantify differences in chromatin accessibility between DEGs, bamscale cov (Pongor

et al., 2020) was employed using ATAC-seq BAM files as input and genomic

coordinates of up-regulated and down-regulated genes. Kruskal-Wallis test was used
to test for significant differences in chromatin accessibility between each category of
DEGs at postnatal and adult stage.

Differential accessibility analysis at TEs: TE gene transfer format (GTF) file was
obtained from http://labshare.cshl.edu/shares/mhammelllab/www-
data/TEtranscripts/TE_GTF/mm10_rmsk_TE.gtf.gz on 03.02.2020. This file provides
hierarchical information about TEs: Class (level 1, eg. LTR), family (level 2, eg. LTR -
>L1), subtype (level 3, eg. LTR-> L1->L1_Rod), and locus (level 4, eg. LTR->L1 ->
L1_Rod -> L1_Rod_dup1). TE loci were annotated based on overlap with
GENCODE (version M18) as described above for ATAC-seq peaks. Filtered BAM
files (without reads mapping to blacklisted or mitochondrial regions) were used to
analyze TEs. Mapped reads were assigned to TEs using featureCounts from R
package Rsubread (Liao et al., 2019) (version 2.0.1) and were summarized to
Subtypes (level 3), allowing for multi-overlap with fractional counts, while ignoring
duplicates. The number of extended reads overlapping at TE loci were obtained
using csaw package (Lun and Smyth, 2015) (version 1.20.0). Subtypes without at
least 15 reads, and loci without at least 5 reads in at least 40% of samples were
filtered out. Normalization and differential accessibility analysis were performed as
described above. Subtypes which had an absolute Log2FC≥0.5 and adjusted-P≤0.05
were categorized as differentially accessible subtypes and loci with an absLog2FC≥1
and adjusted-P≤0.05 were categorized as differentially accessible loci. For further
downstream data analysis, only differentially accessible loci or differentially
accessible subtypes were considered. TF motif enrichment analysis was performed
using the marge package (Amezquita, 2018) (version 0.0.4.9999), which is a
wrapper around the Homer tool (Heinz et al., 2010) (version 4.11.1).

Authors contribution: ILC and IMM designed and conceived the study. ILC
performed all RNA-seq, ICC and ATAC-seq experiments with support of LCS. DKT
and RGA-M analyzed the RNA-seq, ATAC-seq, ChIP-seq with significant support
from PLG. OUF and MC performed chromatin accessibility analysis of DEGs. ILC,
DKT and RGA-M prepared figures. ILC and RGA-M interpreted the data with
significant input from DKT, PLG and IMM. ILC, DKT, RGA-M and IMM wrote the

manuscript, with significant input from PLG. All authors read and accepted the final
version of the manuscript.

Acknowledgments: We thank Martin Roszkowski and Francesca Manuella for
taking care of animals breeding, Yvonne Zipfel for animal care, Silvia Schelbert and
Alberto Corcoba for taking care of animal licenses and lab organization. We thank
Catherine Aquino and Emilio Yángüez from Functional Genomics Center Zurich for
support and advice with libraries preparation and sequencing. We are very grateful
to Jon Oatley, Melissa Oatley, Tessa Lord and Nathan Law for conceptual advice,
hands-on training, and for providing detailed protocols for testis dissection and
preparation, and immunocytochemistry of SCs. We thank Zuguang Gu for support
with heatmaps. We thank Service and Support for Science IT (www.s3it.uzh.ch) for
computational infrastructure.

Competing interest: The authors declare that they have no competing interests.

Funding: The Mansuy lab is funded by the University Zürich, the ETH Zürich, the
Swiss National Science Foundation grant number 31003A_175742/1, the National
Centre of Competence in Research (NCCR) RNA&Disease funded by the Swiss
National Science Foundation (grant number 182880/Phase 2 and 205601/Phase 3),
ETH grants (ETH-10 15-2 and ETH-17 13-2), European Union Horizon 2020
Research Innovation Program Grant number 848158, European Union projects
FAMILY and HappyMums funded by the Swiss State Secretariat for Education,
Research and Innovation (SERI), FreeNovation grant from Novartis
Forschungsstiftung, and the Escher Family Fund. RGA-M received an ETH
Postdoctoral Fellow grant number 20-1 FEL-28. DKT received a Swiss Government
Excellence Scholarship.
768
Data and material availability: Sequencing data is available via the EBI BioStudies
database within the ArrayExpress collection (RNA-seq: E-MTAB-12721; ATAC-seq:
E-MTAB-12722). The code employed for data analysis is available from
https://github.com/mansuylab/SC_postnatal_adult/.   Additional   information   is
available   upon   request.

References

Amezquita RA. (2018) **marge: An API for Analysis of Motifs Using HOMER in R**.
bioRxiv doi:10.1101/249268.

An Z, Liu P, Zheng J, Si C, Li T, Chen Y, Ma T, Zhang MQ, Zhou Q, Ding S. (2019)
Sox2 and Klf4 as the Functional Core in Pluripotency Induction without
Exogenous Oct4. *Cell Rep* **29**:1986-2000.e8.

Andrews S, Krueger F, Segonds-Pichon A, Biggins L, Krueger C, Wingett S. (2012).
FastQC. A quality control tool for high throughput sequence data.

Argelaguet R, Clark SJ, Mohammed H, Stapel LC, Krueger C, Kapourani CA, Imaz-
Rosshandler I, Lohoff T, Xiang Y, Hanna CW, Smallwood S, Ibarra-Soria X,
Buettner F, Sanguinetti G, Xie W, Krueger F, Göttgens B, Rugg-Gunn PJ,
Kelsey G, Dean W, Nichols J, Stegle O, Marioni JC, Reik W. (2019) **Multi-**
omics profiling of mouse gastrulation at single-cell resolution. *Nature*
576:487–491.

Arzate-Mejia RG, Mansuy IM. (2023). **Remembering through the genome: the**
role of chromatin states in brain functions and diseases. *Translational*
Psychiatry **13**:1–11.

Atlasi YS, Stunnenberg H. (2017) **The interplay of epigenetic marks during stem**
cell differentiation and development. *Nat Rev Genet* **18**:643-658

Barber MF, Michishita-Kioi E, Xi Y, Tasselli L, Kioi M, Moqtaderi Z, Tennen RI,
Paredes S, Young NL, Chen K, Struhl K, Garcia BA, Gozani1 O, Li W, Chua KF.
(2012) **SIRT7 links H3K18 deacetylation to maintenance of oncogenic**
transformation. *Nature* **487**:114–118.

Barisic D, Stadler MB, Iurlaro M, Schübeler D. (2019**) Mammalian ISWI and**
SWI/SNF selectively mediate binding of distinct transcription factors.
Nature **569**:136–140.

Bejjani F, Evanno E, Zibara K, Piechaczyk M, Jariel-Encontre I. (2019) **The AP-1**
transcriptional complex: Local switch or remote command? *Biochimica et*
Biophysica Acta (BBA) - Reviews on Cancer **1872**:11–23.

Bergmann JH, Li J, Eckersley-Maslin MA, Rigo F, Freier SM, Spector DL. (2015)
Regulation of the ESC transcriptome by nuclear long noncoding RNAs.
Genome Res **25**:1336–1346.

Bernardino RL, Alves MG, Oliveira PF. (2018) **Evaluation of the purity of sertoli**
cell primary cultures. *Methods in Molecular Biology* **1748**:9–15.

Bialkowska AB, Yang VW, Mallipattu SK. (2017). **Krüppel-like factors in**
mammalian stem cells and development. *Development* **144**:737.

Bleckwehl T, Crispatzu G, Schaaf K, Respuela P, Bartusel M, Benson L, Clark SJ,
Dorighi KM, Barral A, Laugsch M, van IJcken WFJ, Manzanares M, Wysocka J,
Reik W, Rada-Iglesias Á. (2021) **Enhancer-associated H3K4 methylation**
safeguards in vitro germline competence. *Nature Communications 2021 12:1*
12:1–19.

Cao C, Ma Q, Mo S, Shu G, Liu Q, Ye J, Gui Y. (2021) **Single-Cell RNA**
Sequencing Defines the Regulation of Spermatogenesis by Sertoli-Cell
Androgen Signaling. *Front Cell Dev Biol* **9**:3095.

Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, Lemma RB, Turchi L,
Blanc-Mathieu R, Lucas J, Boddie P, Khan A, Manosalva Pérez N, Fornes O,
Leung TY, Aguirre A, Hammal F, Schmelter D, Baranasic D, Ballester B,
Sandelin A, Lenhard B, Vandepoele K, Wasserman WW, Parcy F, Mathelier A.
(2022) **JASPAR 2022: the 9th release of the open-access database of**
transcription factor binding profiles. *Nucleic Acids Res.* **50**(D1):D165-D173.

Chakraborty P, Buaas FW, Sharma M, Snyder E, de Rooij DG, Braun RE. (2014)
LIN28A marks the spermatogonial progenitor population and regulates its
cyclic expansion. *Stem Cells* **32**:860.

Cheng K, Chen IC, Cheng CHE, Mutoji K, Hale BJ, Hermann BP, Geyer CB, Oatley
JM, McCarrey JR. (2020). **Unique Epigenetic Programming Distinguishes**

Regenerative Spermatogonial Stem Cells in the Developing Mouse Testis. *iScience* **23**.

Chereji R V., Eriksson PR, Ocampo J, Prajapati HK, Clark DJ. (2019) **Accessibility of promoter DNA is not the primary determinant of chromatin-mediated gene regulation**. *Genome Res* **29**:1985–1995.

Choksi SP, Lauter G, Swoboda P, Roy S. (2014) **Switching on cilia: transcriptional networks regulating ciliogenesis**. *Development* **141**:1427–1441.

Chu JS, Baillie DL, Chen N. (2010) **Convergent evolution of RFX transcription factors and ciliary genes predated the origin of metazoans**. *BMC Evol Biol* **10**.

Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, Satpathy AT, Rubin AJ, Montine KS, Wu B, Kathiria A, Cho SW, Mumbach MR, Carter AC, Kasowski M, Orloff LA, Risca VI, Kundaje A, Khavari PA, Montine TJ, Greenleaf WJ, Chang HY. (2017) **An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues**. *Nat Methods* **14**:959–962

Costoya JA, Hobbs RM, Barna M, Cattoretti G, Manova K, Sukhwani M, Orwig KE, Wolgemuth DJ, Pandolfi PP. (2004) **Essential role of Plzf in maintenance of spermatogonial stem cells**. *Nat Genet* **36**:653–659.

Dalton S, Treisman R. (1992) **Characterization of SAP-1, a protein recruited by serum response factor to the c-fos serum response element**. *Cell* **68**:597–612.

Dann CT, Alvarado AL, Molyneux LA, Denard BS, Garbers DL, Porteus MH. (2008). **Spermatogonial Stem Cell Self-Renewal Requires OCT4, a Factor Downregulated During Retinoic Acid-Induced Differentiation**. *Stem Cells* **26**:2928–2937.

Davis MP, Carrieri C, Saini HK, Dongen S, Leonardi T, Bussotti G, Monahan JM, Auchynnikava T, Bitetti A, Rappsilber J, Allshire RC, Shkumatava A, O'Carroll

D, Enright AJ. (2017). **Transposon-driven transcription is a conserved**
feature of vertebrate spermatogenesis and transcript evolution. *EMBO Rep*
18:1231–1247.

De Rooij DG. (2017) **The nature and dynamics of spermatogonial stem cells.**
Development (Cambridge). doi:10.1242/dev.146571

Deniz Ö, Frost JM, Branco MR. (2019) **Regulation of transposable elements by**
DNA modifications. *Nat Rev Genet*. doi:10.1038/s41576-019-0106-6

Edwards JR, Yarychkivska O, Boulard M, Bestor TH. (2017) **DNA methylation and**
DNA methyltransferases. *Epigenetics Chromatin* **10**:1–10.

Fort A, Hashimoto K, Yamada D, Salimullah M, Keya CA, Saxena A, Bonetti A,
Voineagu I, Bertin N, Kratz A, Noro Y, Wong CH, De Hoon M, Andersson R,
Sandelin A, Suzuki H, Wei CL, Koseki H, Hasegawa Y, Forrest ARR, Carninci P.
(2014) **Deep transcriptome profiling of mammalian stem cells supports a**
regulatory role for retrotransposons in pluripotency maintenance. *Nat*
Genet **46**:558–566.

Fukuda N, Touhara K. (2005) **Developmental expression patterns of testicular**
olfactory receptor genes during mouse spermatogenesis. *Genes to Cells*
11:71–81.

Gapp K, van Steenwyk G, Germain PL, Matsushima W, Rudolph KLM, Manuella F,
Roszkowski M, Vernaz G, Ghosh T, Pelczar P, Mansuy IM, Miska EA. **(2018)**
Alterations in sperm long RNA contribute to the epigenetic inheritance of
the effects of postnatal trauma. *Molecular Psychiatry* **25**:2162–2174.

Gasperini M, Tome JM, Shendure J. (2020) **Towards a comprehensive catalogue**
of validated and target-linked human enhancers. *Nature Reviews Genetics*
2020 21:5 **21**:292–310.

Gewiss RL, Law NC, Helsel AR, Shelden EA, Griswold MD. (2021) **Two distinct**
Sertoli cell states are regulated via germ cell crosstalk. *Biol Reprod*
105:1591.

Goertz MJ, Wu Z, Gallardo TD, Hamra FK, Castrillon DH. (2011) **Foxo1 is required**
in mouse spermatogonial stem cells for their maintenance and the
initiation of spermatogenesis. *J Clin Invest* **121**:3456–3466.

Green CD, Ma Q, Manske GL, Shami AN, Zheng X, Marini S, Moritz L, Sultan C,
Gurczynski SJ, Moore BB, Tallquist MD, Li JZ, Hammoud SS. (2018) **A**
Comprehensive Roadmap of Murine Spermatogenesis Defined by Single-
Cell RNA-Seq. *Dev Cell* **46**:651-667.

Grive KJ, Hu Y, Shu E, Grimson A, Elemento O, Grenier JK, Cohen PE. (2019)
Dynamic transcriptome profiles within spermatogonial and spermatocyte
populations during postnatal testis maturation revealed by single-cell
sequencing. *PLoS Genet* **15**:e1007810.

Guo J, Grow EJ, Yi C, Goriely A, Hotaling JM, Cairns Correspondence BR. (2017)
Chromatin and Single-Cell RNA-Seq Profiling Reveal Dynamic Signaling
and Metabolic Transitions during Human Spermatogonial Stem Cell
Development. *Cell Stem Cell* **21**:533–546.

Haggerty C, Kretzmer H, Riemenschneider C, Kumar AS, Mattei AL, Bailly N,
Gottfreund J, Giesselmann P, Weigert R, Brändl B, Giehr P, Buschow R,
Galonska C, von Meyenn F, Pappalardi MB, McCabe MT, Wittler L, Giesecke-
Thiel C, Mielke T, Meierhofer D, Timmermann B, Müller FJ, Walter J, Meissner
A. (2021) **Dnmt1 has de novo activity targeted to transposable elements**.
Nature Structural & Molecular Biology 2021 28:7 **28**:594–603.

Hammoud SS, Low DHP, Yi C, Carrell DT, Guccione E, Cairns BR. (2014)
Chromatin and Transcription Transitions of Mammalian Adult Germline
Stem Cells and Spermatogenesis. *Cell Stem Cell* **15**:239–253.

Hammoud SS, Low DHP, Yi C, Lee CL, Oatley JM, Payne CJ, Carrell DT, Guccione
E, Cairns BR. (2015). **Transcription and imprinting dynamics in developing**
postnatal male germline stem cells. *Genes Dev* **29**:2312–24.

Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, Aken
BL, Barrell D, Zadissa A, Searle S, Barnes I, Bignell A, Boychenko V, Hunt T,

Kay M, Mukherjee G, Rajan J, Despacio-Reyes G, Saunders G, Steward C,
Harte R, Lin M, Howald C, Tanzer A, Derrien T, Chrast J, Walters N,
Balasubramanian S, Pei B, Tress M, Rodriguez JM, Ezkurdia I, Van Baren J,
Brent M, Haussler D, Kellis M, Valencia A, Reymond A, Gerstein M, Guigó R,
Hubbard TJ. (2012) **GENCODE: The reference human genome annotation**
for the ENCODE project. *Genome Research* **22**:1760–1774.

He Z, Feng L, Zhang X, Geng Y, Parodi DA, Suarez-Quian C, Dym M. (2005)
Expression of Col1a1, Col1a2 and procollagen I in germ cells of immature
and adult mouse testis. *Reproduction* **130**:333–341.

He Z, Jiang J, Hofmann M-C, Dym M. (2007). **Gfra1 Silencing in Mouse**
Spermatogonial Stem Cells Results in Their Differentiation Via the
Inactivation of RET Tyrosine Kinase 1. *Biol Reprod* **77**:723–733.

He Z, Jiang J, Kokkinaki M, Golestaneh N, Hofmann M-C, Dym M. (2008) **Gdnf**
upregulates c-Fos transcription via the Ras/Erk1/2 pathway to promote
mouse spermatogonial stem cell proliferation. *Stem Cells* **26**:266–278.

Heintzman ND, Hon GC, Hawkins RD, Kheradpour P, Stark A, Harp LF, Ye Z, Lee
LK, Stuart RK, Ching CW, Ching KA, Antosiewicz-Bourget JE, Liu H, Zhang X,
Green RD, Lobanenkov V V., Stewart R, Thomson JA, Crawford GE, Kellis M,
Ren B. (2009) **Histone modifications at human enhancers reflect global**
cell-type-specific gene expression. *Nature* **459**. doi:10.1038/nature07829

Helsel AR, Yang QE, Oatley MJ, Lord T, Sablitzky F, Oatley JM. (2017) **ID4 levels**
dictate the stem cell state in mouse spermatogonia. *Development* **144**:624–
634.

Henikoff S, Henikoff JG, Kaya-Okur HS, Ahmad K. (2020) **Efficient chromatin**
accessibility mapping in situ by nucleosome-tethered tagmentation. *Elife*
9:1–19.

Hermann BP, Cheng K, Singh A, Roa-De La Cruz L, Mutoji KN, Chen I-C,
Gildersleeve H, Lehle JD, Mayo M, Westernströer B, Law NC, Oatley MJ, Velte
EK, Niedenberger BA, Fritze D, Silber S, Geyer CB, Oatley JM, McCarrey JR.

(2018) **The Mammalian Spermatogenesis Single-Cell Transcriptome, from**
Spermatogonial Stem Cells to Spermatids. *Cell Rep* **25**:1650-1667.e8.

Hu Q, Zhang Y, Lou H, Ou Z, Liu J, Duan W, Wang H, Ge Y, Min J, Wang F, Ju Z.
(2021) **GPX4 and vitamin E cooperatively protect hematopoietic stem and**
progenitor cells from lipid peroxidation and ferroptosis. *Cell Death &*
Disease 2021 12:7 **12**:1–9.

Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, Bravo HC,
Davis S, Gatto L, Girke T, Gottardo R, Hahne F, Hansen KD, Irizarry RA,
Lawrence M, Love MI, MacDonald J, Obenchain V, Oles AK, Pages H, Reyes A,
Shannon P, Smyth GK, Tenenbaum D, Waldron L, Morgan M. (2015)
Orchestrating high-throughput genomic analysis with Bioconductor.
Nature Methods **12**:115–121.

Kaluscha S, Domcke S, Wirbelauer C, Stadler MB, Durdu S, Burger L, Schübeler D.
(2022) **Evidence that direct inhibition of transcription factor binding is the**
prevailing mode of gene and repeat repression by DNA methylation. *Nature*
Genetics 2022 54:12 **54**:1895–1906.

Kanatsu-Shinohara M, Naoki H, Tanaka T, Tatehana M, Kikkawa T, Osumi N,
Shinohara T. (2022) **Regulation of male germline transmission patterns by**
the Trp53-Cdkn1a pathway. *Stem Cell Reports* **17**:1924–1941.

Kawase S, Kuwako K, Imai T, Renault-Mihara F, Yaguchi K, Itohara S, Okano H.
(2014) **Regulatory Factor X Transcription Factors Control Musashi1**
Transcription in Mouse Neural Stem/Progenitor Cells. *Stem Cells Dev*
23:2250.

Khan A, Mathelier A. (2017) **Intervene: a tool for intersection and visualization of**
multiple gene or genomic region sets. *BMC Bioinformatics* **18**, 287.

Kiani K, Sanford EM, Goyal Y, Raj A. (2022) **Changes in chromatin accessibility**
are not concordant with transcriptional changes for single-factor
perturbations. *Mol Syst Biol* **18**:e10979.

Klemm SL, Shipony Z, Greenleaf WJ. (2019) **Chromatin accessibility and the**
regulatory epigenome. *Nature Reviews Genetics 2018 20:4* **20**:207–220.

Krueger F. (2015). **Trim Galore. A wrapper tool around Cutadapt and FastQC to**
consistently apply quality and adapter trimming to FastQ files,
www.bioinformatics.babraham.ac.uk/projects/trim_galore/.

Kubota H, Avarbock MR, Brinster RL. (2004) **Culture Conditions and Single**
Growth Factors Affect Fate Determination of Mouse Spermatogonial Stem
Cells. *Biol Reprod* **71**:722–731.

Kubota H, Brinster RL. (2018) **Spermatogonial stem cells**. *Biol Reprod.*
doi:10.1093/biolre/ioy077

Langmead B, Salzberg SL. (2012) **Fast gapped-read alignment with Bowtie 2**.
Nature Methods **9**:357–359.

Law CW, Chen Y, Shi W, Smyth GK. (2014) **voom: precision weights unlock**
linear model analysis tools for RNA-seq read counts. *Genome Biology*
15:R29.

Law NC, Oatley MJ, Oatley JM. (2019). **Developmental kinetics and**
transcriptome dynamics of stem cell specification in the spermatogenic
lineage. *Nat Commun* **10**:2787.

Liao Y, Smyth GK, Shi W. (2019) **The R package Rsubread is easier, faster,**
cheaper and better for alignment and quantification of RNA sequencing
reads. *Nucleic Acids Research* **47**:e47–e47.

Lu JY, Shao W, Chang L, Yin Y, Li T, Zhang H, Hong Y, Percharde M, Guo L, Wu Z,
Liu L, Liu W, Yan P, Ramalho-Santos M, Sun Y, Shen X. (2020) **Genomic**
Repeats Categorize Genes with Distinct Functions for Orchestrated
Regulation. *Cell Rep* **30**:3296-3311.e5.

Lun ATL, Smyth GK. (2015) **csaw: a Bioconductor package for differential**
binding analysis of ChIP-seq data using sliding windows. *Nucleic Acids*
Research **44**:e45.

Ma W, Noble W, Bailey T. (2014) **Motif-based analysis of large nucleotide data**
sets using MEME-ChIP. *Nat Protoc* **9**, 1428–1450.

Marco A, Meharena HS, Dileep V, Raju RM, Davila-Velderrain J, Zhang AL,
Adaikkan C, Young JZ, Gao F, Kellis M, Tsai LH. (2020). **Mapping the**
epigenomic and transcriptomic interplay during memory formation and
recall in the hippocampal engram ensemble. *Nature Neuroscience 2020*
23:12 **23**:1606–1617.

Matson CK, Murphy MW, Griswold MD, Yoshida S, Bardwell VJ, Zarkower D. (2010).
The mammalian Doublesex homolog DMRT1 is a transcriptional
gatekeeper that controls the mitosis versus meiosis decision in male germ
cells. *Dev Cell* **19**:612.

McCarrey, JR. (2023) **Epigenetic priming as a mechanism of predetermination**
of spermatogonial stem cell fate. *Andrology*. 11: 918– 926.

Nichols J, Zevnik B, Anastassiadis K, Niwa H, Klewe-Nebenius D, Chambers I,
Schöler H, Smith A. (1998) **Formation of pluripotent stem cells in the**
mammalian embryo depends on the POU transcription factor Oct4. *Cell*
95:379–391.

Oatley JM, Griswold MD. (2017) **The biology of mammalian spermatogonia, The**
Biology of Mammalian Spermatogonia. doi:10.1007/978-1-4939-7505-1

Oldfield AJ, Yang P, Conway AE, Cinghu S, Freudenberg JM, Yellaboina S, Jothi R.
(2014) **Histone-Fold Domain Protein NF-Y Promotes Chromatin**
Accessibility for Cell Type-Specific Master Transcription Factors. *Mol Cell*
55:708–722.

Pongor LS, Gross JM, Vera Alvarez R, Murai J, Jang SM, Zhang H, Redon C, Fu H,
Huang SY, Thakur B, Baris A, Marino-Ramirez L, Landsman D, Aladjem MI,
Pommier Y. (2020) **BAMscale: quantification of next-generation sequencing**
peaks and generation of scaled coverage tracks. *Epigenetics &*
Chromatin **13**, 21.

Prado F, Jimeno-González S, Reyes JC (2017) **Histone availability as a strategy**
to control gene expression. RNA Biology, 14:3, 281-286.

Quigley IK, Kintner C. (2017) **Rfx2 Stabilizes Foxj1 Binding at Chromatin Loops**
to Enable Multiciliated Cell Gene Expression. *PLoS Genet* **13**.

Rada-Iglesias A, Bajpai R, Swigut T, Brugmann SA, Flynn RA, Wysocka J. (2011) **A**
unique chromatin signature uncovers early developmental enhancers in
humans. *Nature* **470**.

Ramirez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S,
Dündar F, Manke T. (2016) **deepTools2: a next generation web server for**
deep-sequencing data analysis. *Nucleic Acids Research* **44**:W160--W165.

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J. (2019)
g:Profiler: a web server for functional enrichment analysis and
conversions of gene lists. *Nucleic Acids Research* **47**:W191–W198

Reissig LF, Herdina AN, Rose J, Gesek BM, Lane JL, Prin F, Wilson R, Hardman E,
Galli A, Tudor C, Tuck E, Mazzeo CI, White JK, Ryder E, Gleeson D, Adams DJ,
Geyer SH, Mohun TJ, Weninger WJ. (2019) **The Col4a2em1(IMPC)Wtsi**
mouse line: Lessons from the Deciphering the Mechanisms of
Developmental Disorders program. *Biol Open* **8**.

Richard G. (2019). **gtrichard/deepStats: deepStats 0.3.1 (Version 0.3.1)**.
Zenodo. https://doi.org/10.5281/zenodo.3336593

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. (2015) **limma**
powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Research* **43**:e47--e47.

Robinson MD, McCarthy DJ, Smyth GK. (2009). **edgeR: a Bioconductor package**
for differential expression analysis of digital gene expression data.
Bioinformatics **26**:139–140.

Robinson MD, Oshlack A. (2010). **A scaling normalization method for differential**
expression analysis of RNA-seq data. *Genome Biology* **11**:R25.

Sakashita A, Maezawa S, Alavattam K, Yukawa M, Barski A, Pavlicev M, Namekawa
S. (2020) **Endogenous retroviruses drive species-specific germline**
transcriptomes in mammals. *Nat Struct Mol Biol* **27**, 967–977.

Sararols P, Stévant I, Neirijnck Y, Rebourcet D, Darbey A, Curley MK, Kühne F,
Dermitzakis E, Smith LB, Nef S. (2021) **Specific Transcriptomic Signatures**
and Dual Regulation of Steroidogenesis Between Fetal and Adult Mouse
Leydig Cells. *Front Cell Dev Biol* **9**:1726.

Schrans-Stassen BHGJ, van de Kant HJG, de Rooij DG, van Pelt AMM. (1999)
Differential expression of c-kit in mouse undifferentiated and
differentiating type A spermatogonia. *Endocrinology* **140**:5894–5900.

Shaulian E, Karin M. (2002). **AP-1 as a regulator of cell life and death**. *Nat Cell*
Biol **4**:E131–E136.

Shlyueva D, Stampfel G, Stark A. (2014) **Transcriptional enhancers: from**
properties to genome-wide predictions. *Nature Reviews Genetics 2014 15:4*
15:272–286.

Song W, Shi X, Xia Q, Yuan M, Liu J, Hao K, Qian Y, Zhao X, Zou K. (2020) **PLZF**
suppresses differentiation of mouse spermatogonial progenitor cells via
binding of differentiation associated genes. *J Cell Physiol* **235**:3033–3042.

Sun F, Xu Q, Zhao D, Chen CD. (2015) **Id4 Marks Spermatogonial Stem Cells in**
the Mouse Testis. *Scientific Reports* **5**:17594.

Sun Z, Zhu M, Lv P, Cheng L, Wang Q, Tian P, Yan Z, Wen B. (2018). **The Long**
Noncoding RNA Lncenc1 Maintains Naive States of Mouse ESCs by
Promoting the Glycolysis Pathway. *Stem Cell Reports* **11**:741–755.

Sundaram V, Cheng Y, Ma Z, Li D, Xing X, Edge P, Snyder MP, Wang T. (2014)
Widespread contribution of transposable elements to the innovation of
gene regulatory networks. *Genome Res* **24**:1963–1976.

Sundaram V, Wysocka J. (2020) **Transposable elements as a potent source of**
diverse cis-regulatory sequences in mammalian genomes. *Philosophical*

Transactions of the Royal Society B: Biological Sciences.
doi:10.1098/rstb.2019.0347

Tang L, Wang M, Liu D, Gong M, Ying QL, Ye S. (2017). **Sp5 induces the**
expression of Nanog to maintain mouse embryonic stem cell self-renewal.
PLoS One **12**.

Thompson PJ, Macfarlan TS, Lorincz MC. (2016) **Long Terminal Repeats: From**
Parasitic Elements to Building Blocks of the Transcriptional Regulatory
Repertoire. *Mol Cell.* doi:10.1016/j.molcel.2016.03.029

Ueda A, Akagi T, Yokota T. (2017) **GA-Binding Protein Alpha Is Involved in the**
Survival of Mouse Embryonic Stem Cells. *Stem Cells* **35**:2229–2238.

Vierbuchen T, Ling E, Cowley CJ, Couch CH, Wang X, Harmin DA, Roberts CWM,
Greenberg ME. (2017) **AP-1 Transcription Factors and the BAF Complex**
Mediate Signal-Dependent Enhancer Selection. *Mol Cell* **68**:1067-1082.e12.

Wang M, Liu X, Chang G, Chen Yidong, An G, Yan L, Gao S, Xu Y, Cui Y, Dong J,
Chen Yuhan, Fan X, Hu Y, Song K, Zhu X, Gao Y, Yao Z, Bian S, Hou Y, Lu J,
Wang R, Fan Y, Lian Y, Tang W, Wang Y, Liu J, Zhao L, Wang L, Liu Z, Yuan
R, Shi Y, Hu B, Ren X, Tang F, Zhao XY, Qiao J. (2018) **Single-Cell RNA**
Sequencing Analysis Reveals Sequential Cell Fate Transition during
Human Spermatogenesis. *Cell Stem Cell* **23**:599-614.e4.

Wang Mei, Yu L, Wang S, Yang F, Wang Min, Li L, Wu X. (2020) **LIN28A binds to**
meiotic gene transcripts and modulates their translation in male germ
cells. *J Cell Sci* **133**.

Wu F, Liu Y, Wu Q, Li D, Zhang L, Wu X, Wang R, Zhang D, Gao S, Li W. (2018).
Long non-coding RNAs potentially function synergistically in the cellular
reprogramming of SCNT embryos. *BMC Genomics* **19**:631.

Wu X, Oatley JM, Oatley MJ, Kaucher A V., Avarbock MR, Brinster RL. (2010). **The**
POU domain transcription factor POU3F1 is an important intrinsic
regulator of GDNF-induced survival and self-renewal of mouse
spermatogonial stem cells. *Biol Reprod* **82**:1103–1111.

Xiong L, Tolen EA, Choi J, Velychko S, Caizzi L, Velychko T, Adachi K, Maccarthy
CM, Lidschreiber M, Cramer P, Schöler HR. (2022) **Oct4 differentially**
regulates chromatin opening and enhancer transcription in pluripotent
stem cells. *Elife* **11**. doi:10.7554/ELIFE.71533

Yamane M, Ohtsuka S, Matsuura K, Nakamura A, Niwa H. (2018). **Overlapping**
functions of krüppel-like factor family members: Targeting multiple
transcription factors to maintain the naïve pluripotency of mouse
embryonic stem cells. *Development (Cambridge)* **145**.

Zhang T, Oatley J, Bardwell VJ, Zarkower D. (2016). **DMRT1 Is Required for**
Mouse Spermatogonial Stem Cell Maintenance and Replenishment. *PLoS*
Genet **12**:e1006293.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C,
Myers RM, Brown M, Li W, Liu XS. (2008) **Model-based Analysis of ChIP-Seq**
(MACS). *Genome Biology* **9**:R137.

Zhu L, Gazin C, Lawson N, Pagès H, Lin S, Lapointe D, Green M. (2010)
ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-
chip data. *BMC Bioinformatics*, **11**, 237.

Figure legends

Figure 1. Transcriptome dynamics between early postnatal and adult SCs.
(A) Schematic representation of the experimental strategy to isolate and analyse
SCs from postnatal or adult male mice.
(B) Upper panel: Representative dot plots of the sorting strategy for enrichment of
postnatal and adult SCs. Gating based on side scatter/forward scatter (SSC-
A/FSC-A) and forward scatter (height/forward scatter) (FSC-H/FSC-A) area
was conducted to exclude cell debris and clumps. Lower panel:
Representative immunocytochemistry on unsorted and sorted PND15 cells
with anti-PLZF antibody and DAPI (nuclei) showing enrichment of PLZF+ cells
after FACS.
(C) Heatmap of expression profile of selected markers of SCs and different
testicular somatic cells extracted from total RNA-seq data on PND8, PND15

and adult samples (n=6 for each group). Key genes for stem cell potential,
stem and progenitor spermatogonia and Leydig and Sertoli cells were chosen
to assess the enrichment of SCs in the sorted cell populations. Each row in
the heatmap represents a biological replicate from two experimental batches
(B1 and B2). At the bottom are shown Integrative Genomics Viewer (IGV)
tracks for aggregated RNA-seq signal for PND8, PND15 and adult over key
genes used as markers. Gene expression is represented as Log2CPM
(counts per million).

(D) Volcano plot of differentially expressed genes (DEGs) (adjusted-P≤0.05 and
absolute Log2FC≥1) between PND8 and PND15 (left) and PND15 and adult
(right). Figures in grey boxes indicate the number of down- and up-regulated
DEGs in each comparison.

(E) Genomic IGV snapshots of exemplary DEGs showing aggregated RNA-seq
signal for PND8, PND15 and adult SCs.

(F) Left: Bar-plot of GO categories enriched in DEGs between PND8 and PND15
(adjusted-P≤0.05). Dotted line indicates threshold value for significance of
0.05. Right: Heatmap of DEGs between PND8 and PND15 belonging to the
GO category "Transcription". Each row represents a biological replicate from
two experimental batches. Shown are Log2FC with respect to average at
PND8.

(G) Left: Bar-plot of GO categories enriched in DEGs between PND15 and adult
(adjusted-P≤0.05). Dotted line indicates threshold value for significance of
0.05. Right: Heatmap of DEGs between PND8 and PND15 belonging to the
GO category "Spermatogenesis". Each row represents a biological replicate
from two experimental batches. Shown are Log2FC with respect to average at
PND8.

(H) Left: Heatmap of all DEGs specific to adult SCs. Each row represents a
biological replicate from two experimental batches (B1 and B2). Shown are
Log2FC with respect to average at PND8. Right, Bar-plots of GO categories
enriched in up-regulated (adjusted-P≤0.05) (top) or down-regulated (adjusted-
P≤0.05) (bottom) genes in adult SCs. Dotted line indicates threshold value for
significance of adjusted-P=0.05.

Figure 2. Dynamics of chromatin accessibility of SCs during the transition from
postnatal to adult stage

(A) Volcano plot of DARs obtained by ATAC-seq (adjusted-P≤0.05 and absolute
Log2FC≥1) between PND15 and adult SCs. Figures in grey boxes indicate the
number of DARs with decreased (Down) or increased (Up) chromatin
accessibility in PND15 compared to adult SCs.

(B) Heatmaps showing normalized ATAC-seq signal for all identified DARs
comparing PND15 and adult stages. Each row represents a 2kb genomic region
extended 1kb down- and upstream from the centre of each identified DAR (as
shown in A). Rows are ordered in a decreasing level of mean accessibility.

(C) Bar plot illustrating the genomic distribution of DARs between PND15 and adult
SCs.

(D) Bar plot of GO categories for associated genes assigned by proximity to all
DARs. Dotted line indicates threshold value for significance of adjusted-P=0.05.

(E) IGV tracks for ATAC-seq signal for PND15 and adult SCs showing DARs (red
box) located at promoters and intergenic regions.

(F) Heatmaps showing normalized ChIP-seq signal for different histone marks in
adult SCs (public data derived from Cheng et al., 2020) at all identified ATAC-seq
peaks from PND15 and adult stages. Each row in the heatmap represents a 2kb
genomic region extended 1kb down- and up-stream from the centre of each
identified peak. Shown data correspond to non-regenerative SCs as stated in
Cheng et al., 2020.

(G) Pie charts showing the overlap of all identified DARs with genomic regions
significantly enriched for different histone marks in adult SCs (public data derived
from Cheng et al., 2020).

Figure 3. TF binding motifs at DARs and their transcriptional dynamics during the
transition from postnatal to adult SCs

(A) Table of top five enriched motifs in genomic regions with decrease in chromatin
accessibility in adult SCs and TF matching motifs. A full list of TF motifs is
provided in Table S4.

(B) Heatmap of expression profile of TFs with motifs in genomic regions with
decreased chromatin accessibility shown in (A). Each row represents a biological
replicate from two experimental batches. Unsupervised hierarchical clustering

was applied to each row and a dendrogram indicating similarity of expression

profiles among genes is shown. Shown are Log2FC with respect to average of

PND8 SCs.

(C) Line-plots of average expression of each gene displayed in the heatmap in (B)

showing the dynamics of gene expression across PND8, PND15 and adult SCs.

(D, E) Genomic snapshot from IGV showing aggregated RNA-seq signal from PND8,

PND15 and adult SCs for the TFs D) *Dmrt1* and E) *Elk4.*

(F) Table of top five enriched motifs in genomic regions with increase in chromatin

accessibility in adult SCs and TF matching motifs. A full list of TF motifs is

provided in Table S4.

(G)Heatmap of expression profile of TFs with motifs in genomic regions with increase

in chromatin accessibility shown in (F). Shown are Log2FC with respect to the

average of PND8 SCs.

(H)Line-plots of average expression of each gene displayed in the heatmap in (G)

showing the dynamics of gene expression across SCs development.

(I, J) Genomic snapshot from IGV showing the aggregated RNA-seq signal from

PND8, PND15 and adult SCs for the TFs I) *Jun* and J) *Rfx2.*

Figure 4. Chromatin accessibility dynamics at DEGs during the transition from

postnatal to adult SCs

(A)   *Top*, profile plots of ATAC-seq signal from PND15 and adult SCs over all

ATAC-seq peaks located within a genomic region of 5kb surrounding the TSS of

DEGs. Mean (line) and bootstrap confidence interval (area, shadow lines)

computed over all non-overlapping 50bp genomic regions. n.s, non-significant

differences at TSS region. *, significant difference in signal between PND15 and

adult at p-value<0.05. PC, peak center from ATAC-seq data. Profile plots for all

ATAC-seq peaks surrounding the TSS of down-regulated (top left) or up-regulated

(top right) DEGs.. *Middle,* profile plots of ATAC-seq signal from PND15 and adult

SCs over all ATAC-seq peaks located within a genomic region of 500bp

surrounding the TSS of DEGs. *Bottom*, heatmaps showing normalized ATAC-seq

signal from PND15 and adult for all ATAC-seq peaks surrounding the TSS of all

down-regulated (left) or up-regulated (right) DEGs. Each row in the heatmap

represents a 2kb genomic region extended 1kb down- and up-stream from the

center of each identified peak. Row Are ordered in a decreasing level of mean
accessibility taking PND15 as reference.

(B)   Violin plots of average ATAC-seq signal for PND15 and adult SCs over gene
body of down-regulated (left) or up-regulated (right) DEGs. ****p<0.0001.

(C)   TF motif enrichment analysis for all ATAC-seq peaks surrounding the TSS of
all down-regulated (left) or up-regulated (right) DEGs.

(D)   *Left*, pie chart of percentage of DARs overlapping the extended promoter of
DEGs (+/- 5kb from TSS). *Right*, pie chart of percentage of DEGs associated with a
DAR.

(E)   IGV tracks for an example of DAR associated to the extended promoter (+/-
5kb from TSS) of a DEG. Shown are tracks for RNA-seq and ATAC-seq signal for
PND15 and adult SCs generated in this study and signal track for H3K4me3 from
SCs derived from public ChIP-seq data (Cheng et al., 2020).

(F)   *Left*, pie chart of percentage of DARs overlapping potential regulatory
sequences of DEGs (+/-100kb from start and end of gene). *Right*, pie chart of the
percentage of DEGs associated with a DAR.

(G)   IGV tracks for an example of DAR associated with potential regulatory
sequences (+/-100kb from start and end of gene) of a DEG.

1258

Figure 5. Differential chromatin accessibility at TEs in PND15 and adult SCs.

(A) Heatmap of LTR and LINE subtypes with decreased accessibility between
PND15 and adult SCs extracted from Omni-ATAC data (adjustedP≤0.05 and
Log2FC≥0.5)

(B) Heatmap of LTR and LINE subtypes with increased accessibility between
PND15 and adult SCs extracted from Omni-ATAC data (adjustedP≤0.05 and
Log2FC≥0.5). Expression changes of these subtypes between adult and PND15
SCs RNA-seq from literature are represented as Log2FC.

(C) Bar plot illustrating the genomic distribution of DARs between PND15 and adult
SCs.

(D) Genomic snapshot from IVG showing aggregated ATAC-seq and RNA-seq
signal from PND8, PND15 and adult SCs for *Lncenc1* and *Platr14.* Below signal
tracks is shown the RepeatMasker track for the mouse genome, and TEs with
changes in chromatin accessibility are highlighted.

Supplementary Figure 1. Expression of SCs markers and their dynamics between
early postnatal and adult stage.

(A) Heatmap of the expression profile of an extended list of selected markers of
spermatogonial and different testicular somatic cells extracted from total RNA-seq
data on PND8, PND15 and adult samples (n=6 for each group). Each row in the
heatmap represents a biological replicate from two experimental batches. Gene
names in bold correspond to cellular markers presented in Figure 1C. Gene
expression is represented as Log2CPM (counts per million).

(B) *Left*, Heatmap of the expression profile of genes involved in germ-cell
maintenance, pluripotency and signalling as reported by Hammoud, 2015. Each
row in the heatmap represents a biological replicate from two experimental
batches. Unsupervised hierarchical clustering was applied to each of the rows of
the heatmaps and a dendrogram indicating similarity of expression profiles among
genes in each biological category is shown with each heatmap. *Right*, line-plots of
the average expression of each gene displayed in the heatmaps showing the
dynamics and breadth of gene expression across SCs postnatal development.

(C) Genomic snapshots from IVG of exemplary DEGs showing aggregated RNA-seq
signal from PND8, PND15 and adult stages.

(D) *Left*, bar-plot of GO KEGG pathways categories enriched in DEGs between
PND15 and adult SCs (adjusted-P≤0.05). Dotted line indicates a threshold value
for significance of 0.05. *Right*, Heatmap of DEGs between PND8 and PND15
belonging to the GO category "ECM-receptor interaction". Each row in the
heatmap represents a biological replicate from two experimental batches. Shown
are Log2FC with respect to the average of PND8.

Supplementary Figure 2. Specific transcriptional programs between postnatal and
adult SCs.

(A) Four-way Venn diagram of DEGs detected between PND8-PND15 and PND15-
adult comparisons.

(B) Heatmap of all DEGs specific to PND8 stage. Each row in the heatmap represents
a biological replicate from two experimental batches. Shown are the Log2 FC with
respect to the average of PND8.

(C) Heatmap of all DEGs with significant changes across the 3 developmental time
points. Each row in the heatmap represents a biological replicate from two

experimental batches. Shown are the Log2 FC with respect to the average of
PND8.

Supplementary Figure 3. Chromatin accessibility landscape of SCs from PND15 to
adult stage.
(A) Heatmaps showing normalized ATAC-seq signal for all identified ATAC-seq
peaks from PND15 and adult stages. Each row in the heatmap represents a 2kb
genomic region extended 1kb down- and upstream from the center of each
identified peak. Each row is ordered in a decreasing level of average
accessibility. On top of each heatmap is shown a plot of the signal profile over all
ATAC-seq peaks and their extended genomic region.
(B) Bar plot of the genomic distribution of all identified ATAC-seq peaks from PND15
and adult stages.
(C) IGV tracks for ATAC-seq signal for PND15 and adult SCs showing ATAC-seq
peaks located at promoters and intragenic regions of genes with important
functions in SCs. Highlighted genomic segments correspond to ATAC-seq
peaks.
(D) Heatmaps showing normalized ChIP-seq signal for different histone marks in
adult Scs (public data derived from Cheng et al., 2020) at DARs from PND15
and adult stages. Each row in the heatmap represents a 2kb genomic region
extended 1kb down- and upstream from the centre of each identified peak.
Shown data corresponds to non-regenerative SCs as stated in Cheng et al.,
2020.

1328

Figure 1. Lazar-Contes et al.

Figure 2. Lazar-Contes et al.

BP: Biological Process; MF: Molecular Function: CC: Cellular Component

Figure 3. Lazar-Contes et al.

Figure 4. Lazar-Contes et al.

Figure 5. Lazar-Contes et al.

A

PND15 Adult

- LTR-ERV1
- LTR-ERVK
- LTR-ERVL
- LTR-MaLR
- LTR-Gypys
- LINE-BovB

log2FC
-2 0 -1 0.5

B

PND15 Adult

- LTR-ERV1
- LTR-ERVK
- LTR-ERVL
- LTR-MaLR
- LINE-L1

log2FC
-2 0 2

C

Percentage of TE loci with changes in chromatin accessibility (%)

- Promoter
- Exon
- Intron
- Distal Intergenic

Increased accessibility (n=6197)

Decreased accessibility (n=5881)

D

ATAC-seq [0 - 3.52] PND15
[0 - 3.52] Adult
RNA-seq [0 - 0.72] PND15
[0 - 0.72] Adult
Repeats **RLTR17**
Lncenc1

ATAC-seq [0 - 8.91] PND15
[0 - 8.91] Adult
RNA-seq [0 - 0.43] PND15
[0 - 0.43] Adult
Repeats **RLTR16B**
Gm13884 *Platr14*