Targeted knockdown of ribulose-1, 5-bisphosphate carboxylase-oxygenase in rice mesophyll cells impact on photosynthesis and growth

Chirag Maheshwari[1], Robert A Coe[1], Shanta Karki[1], Sarah Covshoff[2], Ronald Tapia[1], Aruna Tyagi[3], Julian M. Hibberd[2], Robert T. Furbank[4], W Paul Quick[1,5], Hsiang-Chun Lin[1]*

[1]C_4 Rice Centre, International Rice Research Institute (IRRI), Los Baños, Philippines.

[2]Department of Plant Sciences, University of Cambridge, Cambridge, CB2 3EA, United Kingdom.

[3]Division of Biochemistry, ICAR-Indian Agricultural Research Institute, New Delhi, India.

[4]ARC Centre of Excellence for Translational Photosynthesis, Research School of Biology, The Australian National University, Acton, 2601, Australia

[5]Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, UK.

*To whom correspondence should be addressed: h.lin@irri.org

Abstract

We generated antisense constructs targeting two of the five Rubisco small subunit genes (*OsRBCS2 and 4*) which account for between 30-40% of the RBCS transcript abundance in leaf blades. The constructs were driven by a maize phosphoenolpyruvate carboxylase (PEPC) promoter known to have enriched expression in mesophyll cells (MCs). In the resulting lines leaf Rubisco protein content was reduced by between 30- 50% and CO_2 assimilation rate was limited under photorespiratory and non-photorespiratory conditions. A relationship between Rubisco protein content and CO_2 assimilation rate was found. This was associated with a significant reduction in dry biomass accumulation and grain yield of between 37 to 70%. In addition to serving as a resource for reducing Rubisco accumulation in a cell-preferential manner, these lines allow us to characterize gene function and isoform specific suppression on photosynthesis and

growth. Our results suggest that the knockdown of multiple genes is required to completely

reduce Rubisco accumulation in MCs.

Abbreviation: A, net rate of CO_2 assimilation; ACi, photosynthesis CO_2 response curves; BSC,

bundle sheath cell; C_a, ambient CO_2 concentration; CE, carboxylation efficiency; Chl,

chlorophyll; C_i, intercellular CO_2 concentration; rbcL, large subunit of Rubisco; MC, mesophyll

cell; PPFD, photosynthetic photon flux density; R_d, dark respiration rate; Rubisco, ribulose-1,5-

bisphosphate carboxylase/oxygenase; RuBP, ribulose 1,5-bisphosphate; RBCS, small subunit of

Rubisco.

Keywords: *Oryza sativa*, antisense, Rubisco, RBCS, C_4 rice.

Introduction

In C_3 photosynthesis the first step of CO_2 fixation is carried out by ribulose-1, 5-bisphosphate

carboxylase-oxygenase (Rubisco, EC 4.1.1.39) in the chloroplast of mesophyll cells (MCs).

Photosynthetic rate is limited by the activity of Rubisco because of its extremely low catalytic

turnover rate and competing oxygenation reaction, which leads to the formation of toxic

metabolites which must be broken-down by a series of reactions in a process known as

photorespiration (von Caemmerer and Quick, 2000). This consumes energy leading to the loss of

previously fixed CO_2. As the temperature increases the rate of oxygenation increases due to a

decline in the ratio of CO_2 to O_2 in the leaf cell, as such rates of photorespiration are highest in

hot climates such as the major rice growing regions of the world. To compensate for these

catalytic inefficiencies, Rubisco is produced in large amounts accounting for up to 50% of total

leaf protein and forming a major sink for leaf nitrogen (Parry et al., 2013).

Some species suppress the oxygenation reaction allowing Rubisco to operate close to its

maximal carboxylase rate (Whitney et al., 2011). In C_4 plants Rubisco is localized to the bundle

sheath cells (BSCs) allowing CO_2 to be concentrated at high levels at the exclusion of oxygen.

As a result, radiation use efficiency (RUE) and yield of C_4 species such as maize and sorghum

can be up to 50% higher than those of C_3 species such as rice and wheat (Sheehy et al., 2007).

Water use efficiency (WUE) and nitrogen use can be up to 50% lower (Ghannoum et al., 2011).

The potential efficiencies associated with C_4 photosynthesis has stimulated interest in converting

rice (*Oryza sativa* L.) from a C_3 to a C_4 photosynthetic pathway (Hibberd et al., 2008, Kajala et

al., 2011; Ermakova et al. 2019). The C_4 Rice Consortium (https://c4rice.com/; von Caemmerer

et al., 2012) was established with this goal in mind. To investigate the feasibility of such an

endeavor a toolkit of transgenic resources is being assembled (Kajala et al., 2011; Ermakova et

al., 2019). Among these, there are lines in which parts of the photorespiratory cycle have been

selectively downregulated to test the hypothesis of whether this primes a plant for C_4

photosynthesis (Lin et al., 2016). Here we report on the downregulation and translocation of

Rubisco from MCs to BSCs and the effect on rice photosynthesis and growth.

In rice the Rubsico holoenzyme is made up of eight large Rubisco subunits (rbcLs) encoded by a

single gene on the chloroplast DNA and eight small Rubisco subunits (RBCSs) encoded by a

five multigene family on the nucleus DNA (Dean et al., 1989; Rodermel 1999; Sasanuma 2001).

OsRBCS1 is located on chromosome 2 but is not expressed in the leaf blade (Morita et al., 2014;

Suzuki et al., 2007, 2009). The remaining four genes (*OsRBCS2-OsRBCS5*) are in a tandem

array on chromosome 12 and are highly expressed in photosynthetic active tissues such as leaf

blades (Suzuki et al., 2009). The deduced amino acid sequence without the transit peptide for

targeting the chloroplast are completely identical, leading to the assumption that there are no

functional differences within the *RBCS* gene family within a species. It has previously been

shown that suppression of a *RBCS* gene leads to a reduction in the holoenzyme (Hudson et al.,

1992; Makino et al., 1997; Rodermel et al., 1988) with the expression of *rbcL* regulated at its

transcript level in response to the availably of RBCS protein (Makino et al., 1997; Suzuki and

Makino 2012). Although RBCS protein has no catalytical function for CO_2 fixation, it is

important for maximal activity and structural stability for the Rubisco holoenzyme (Andersson

and Backlund, 2008).

To reduce Rubisco protein accumulation, we generated antisense constructs targeting two of the

five *RBCS* genes (*OsRBCS2* and *4*) expressed them in photosynthetic tissue. Together these

account for between 30-40% of the *RBCS* transcript abundance in leaf blades (Suzuki et al.,

2009), allowing a substantial reduction in Rubisco protein accumulation to be achieved without a

lethal phenotype developing. Lines must produce viable seed in order to be crossed with lines

expressing C_4 enzyme genes facilitating gradual replacement of C_3 with C_4 photosynthesis

(Kajala et al., 2011). The antisense genes against *OsRBCS2* and *4* are driven by a maize

phosphoenolpyruvate carboxylase (PEPC) promoter known to have enriched expression in MCs
(Matsuoka et al., 1994). In addition to serving as a resource for reducing Rubisco accumulation
in a cell-specific manner, these lines allow us to characterize gene function and isoform specific
suppression on photosynthesis and growth.

Materials and Methods

Generation of *OsRBCS2* and *OsRBCS4* knockdown transgenic rice lines

To individually reduce *OsRBCS2* (Os12g17600) and *OsRBCS4* (Os12g19470) expression in rice
MCs, the full coding sequences (CDS) of LOC_Os12g17600.1 and LOC_Os12g19470.2 were
cloned separately in the antisense direction via RT-PCR using the following primers: 5'–
CACCTTAGTTGCCACCAGACTCCT and 5'–ATGGCCCCCTCCGTGATGGC for *OsRBCS2*,
5'–CACCTTAGTTGCCGCCTGACTCCT and 5'–ATGGCTCCCTCGGTGATGGC for
OsRBCS4. The CACC at the 5' end of the forward primer allowed directional cloning into
pENTR/D-TOPO (ThermoFisher Scientific, USA) and subsequent Gateway cloning into the
pSC110 expression vector. The pSC110 vector was created by Gibson Assembly and contained
the B73 *ZmPEPC* promoter to drive gene expression in rice MCs (Gibson et al., 2009; Osborn et
al., 2017; Matsuoka et al., 1994). Both antisense *OsRBCS2* and *OsRBCS4* vectors were verified
by sequencing and transformed into rice (*Oryza sativa* indica *cv*. IR64) using *Agrobacterium*-
mediated transformation following the method described by Yin et al. (2019).

Plant growth conditions

Plants were grown at the International Rice Research Institute (IRRI), Los Baños, Philippines,
14°9'53.58"N, 121°15'32.19"E. Seeds were sown and germinated in 100 ml rootrainers
(http://rootrainers.co.uk/), after one-week plants were transplanted into 7-liter pots filled with
soil from the IRRI upland farm. Plants were grown in a containment transgenic screen house
with a day/night temperature of $35 \pm 3°C/28 \pm 3°C$ and relative humidity of 70-90%. Maximum
light intensity was 2000 µmol photons m^{-2} s^{-1} on a clear sunny day.

PCR screening

Transgenic plants were subjected to genomic PCR screening to confirm the presence of *OsRBCS*
antisense DNA sequence. PCR was carried out by using the KAPA 3G plant PCR kit (Kapa
Biosystem, USA; https://www.sigmaaldrich.com/life-science/roche-biochemical-reagents/kapa-
genomics-reagents.html) with the gene-specific primers (5´-ACGACTCCCCATCCCTATTT
and 5´-TGCATTCGTCCGTATCATC for *OsRBCS2*; 5´-CATTGATCACCAATCGCATC and
5´-ACGTCAGCAATGGCGGAA for *OsRBCS4*). Plasmid DNA containing antisense *OsRBCS2*
and *OsRBC4* were used as positive controls and non-transgenic rice and water as negative
controls. PCR conditions were as follows: pre-denaturation for 5 min at 95°C, 30 cycles of the
polymerization reaction consisting of a denaturation step for 20 s at 95°C, an annealing step for
15 s at 60°C and an extension step for 1 min at 72°C, and a final extension step for 5 min at
72°C.

DNA blot analysis

127 Large-scale genomic DNA was extracted from leaves at the mid-tillering stage. DNA blot
128 analysis was carried out as described by Lin et al., (2016) except genomic DNA was digested
129 with *BgL*II restriction endonuclease (New England Biolabs, USA;
130 https://international.neb.com/).

Quantitative RT-PCR

Total RNA was extracted and analyzed as described in Lin et al., (2016) except for the primer
pairs used. The mRNA level of each *OsRBCS* gene was quantified on a fold change basis in
comparison with wild-type plants. 3´- untranslated regions (UTRs) of the *OsRBCS* gene family
have different nucleotide sequences and so were utilized for amplification. The *Elongation factor*
1-alpha gene (*OsEF1α*; Os03g0177500) was used as an internal standard. Primers pairs were:
5'-CTTCGGCAACGTCAGCAATG and 5'-AGCACGGCCGGTAAAATCA for *OsRBCS2*; 5'-
TTCCAAGGGCTCAAGTCCAC and 5'-ATAGGCAGCAATCCACCAGC for *OsRBCS3*; 5'-
GTGGCCGATTGAGGGCATTA and 5'-TCACCGTCACCGAACGATAG for *OsRBCS4*; 5'-
GGTGTGGCCGATTGAAGG and 5'-ACTGGGAAAGGGGAAACCAAT for *OsRBCS5*; 5'-
CCACTGGTGGTTTTGAGGC and 5'GGCCCTTGTACCAGTCAAG for *OsEF1α*.

Soluble leaf protein

Leaf samples for soluble protein were harvested between 09:00 and 11:00 h from the fourth fully

expanded leaf. Proteins were extracted and fractionated as described previously Lin et al. (2016).

Samples were loaded based on an equal leaf area (0.396 mm^2 for rbcL and RBCS). After

electrophoresis, proteins were electroblotted onto a polyvinylidene difluoride membrane and

probed with antisera against Rubisco protein (provided by Richard Leegood, Sheffield

University, UK) at a dilution 1:200. A peroxidase-conjugated secondary antibody was used at a

dilution of 1:200 and immunoreactive bands were visualized with ECL Western Blotting

Detection Reagents (GE Healthcare, USA; https://www.gelifesciences.com). Rubisco protein

content was quantified from SDS-page gels using ALPHA-Ease FC software (Alpha Innotech,

USA), values are expressed as the percentage protein reduction compared to the wild-type.

Leaf chlorophyll content, plant growth analysis and destructive harvesting

Leaf chlorophyll content was measured with a SPAD 502 Chlorophyll Meter (SPAD, Konica

Minolta; https://www.konicaminolta.com) at the mid-tillering stage on the upper fully expanded

leaves. Values given are the average ± SE of three measurements from six plants per line. Plant

height was measured from the soil surface to the tip of the youngest fully expanded leaf. Total

tiller number was counted prior to harvesting for destructive measurements. All above-ground

biomass (leaves, stems, and sheaths) were harvested, weighed, and placed in paper bags, and

oven dried at 70°C until a constant dry biomass weight was achieved. Values of plant height,

total tiller number, and dry biomass are presented as the average ± SE of ten plants per line.

Gas exchange measurement

Leaf gas exchange measurements were made at IRRI using a LI-6400XT portable photosynthesis

system (LI-COR Biosciences, USA; https://www.licor.com) as described in Lin et al. (2016)

during the tillering stage for each line and wild type, 60-65 days post-germination.

Immunolocalization

Leaf samples were harvested between 09:00 h and 11:00 h from 9-week-old plants. The middle

portion of the seventh fully expanded leaf was dissected and fixed for four hours in a solution

containing 4% paraformaldehyde, 0.2% glutaraldehyde, and 25 mM sodium phosphate buffer pH

7.2. After fixation, sections were rinsed four times in 25 mM phosphate buffer over a period of

60 min. Thin leaf sections were cut using a razor blade and blocked in TBST buffer (0.1%

Tween 20, 20mM Tris, 154 mM NaCl) containing 3% milk for two hours at room temperature.

Sections were probed with antisera against Rubisco protein diluted 1:100 in blocking solution

and incubated overnight at 4°C. The remaining steps were performed at room temperature.

Sections were washed six times with blocking solution and then incubated for two hours with

Alexa Fluor 488 goat anti-rabbit IgG (Invitrogen, USA;

https://www.thermofisher.com/ph/en/home/brands/invitrogen.html) at 37°C in the dark. Sections

were washed six times with blocking solution, post-stained with 0.05% calcofluor white for five

min and washed with distilled water. Sections were mounted on microscope slides in 50%

glycerol and examined on a BX61 Disk Scanning microscope (Olympus, USA;

https://www.olympus-global.com) with florescence function under DAPI, RFP and GFP filters.

Leaf Anatomy Analysis

Leaf width was measured at the middle portion of the fully expanded penultimate leaf. Between

2 to 4 mm^2 of the middle portion of the penultimate leaf was cut and fixed in formaldehyde-

acetic acid-alcohol (FAA) solution under vacuum (20 psi) at room temperature for at least 12 h.

Leaf sections were rinsed twice in water for 60 min. Leaf sections were dehydrated using an

ethanol series, incubated twice in 70% ethanol for 30 min; once in 80% ethanol for 30 min; once

in 90% ethanol for 30 min, and finally three times in 100% ethanol for 30 min. Leaf sections

were infiltrated in 10% ethanol and varying concentrations of Spurr's resin solution (10% to

100%). Sections were incubated for at least 60 min at each concentration solution. Leaf sections

were then finally infiltrated twice for 4 h with 100% Spurr's resin solution. Sections were placed

in fresh 100% Spurr's resin solution in molds. Samples were polymerized overnight at 70°C.

Embedded leaf sections were cut into 10-15 μm thick sections on a microtome (MT2-B, DuPont-

Instruments-Sorvall, USA) and then dried. Sections were stained with 0.05% toluidine blue O

stain in 0.1% sodium carbonate, pH 11.1 for 5 min then rinsed four times with distilled water,

each for 5 min. Sections were dried and mounted on slides using Permount^{TM} (Fischer Scientific,

USA; https://www.fishersci.com/us/en/home.html). Transverse images were acquired using a

BX51 Olympus microscope (Olympus, USA; https://www.olympus-global.com/) at 4X and 40X

magnification. Mesophyll cell number was counted by calculating the numbers of mesophyll

cells in between minor veins. Leaf thickness, interveinal distance (IVD), vein number (major and

minor), mesophyll cell length was measured as described in Chatterjee et al. (2016).

Statistics

Statistical analyses were performed using Statistical Tool for Agricultural Research (STAR)
software (International Rice Research Institute, Philippines) using a one-way analysis of variance
(ANOVA) or a Student's t-test with a P-value of < 0.05.

Results

OsRBCS2 and Os*RBCS4* specific knockdown lines

A total of 190 T_0 plants were PCR positive for the antisense-Os*RBCS2* construct and 30 T_0 plants
for the antisense-*OsRBCS4* construct. Plants with reduced Rubisco protein accumulation relative
to the wild-type rice were selected for DNA blot analysis in order to confirm T-DNA integration.
Two *OsRBCS2* knockdown events (*rbcs2*-165 and *rbcs2*-266) and two *OsRBCS4* knockdown
events (*rbcs4*-022 and *rbcs4*-053) with the lowest Rubisco accumulation were selected from T_1
progeny. These were advanced generating a total of 40 PCR positive T_3 generation *OsRBCS2*
knockdown plants (from two events *rbcs2*-165 and *rbcs2*-266) and 40 T_2 generation *OsRBCS4*
knockdown plants (from two events *rbcs4*-022 and *rbcs4*-053) (Fig. S1). DNA blot analysis
showed that these transgenic lines carried between two and four copies of the antisense
constructs (Fig. S2). This led to a reduction in the accumulation of *OsRBCS2* transcripts of
between 15 and 24-fold (Fig. 1A) and *OsRBCS4* transcripts of between 6 and 12-fold in the
respective knockdown lines (Fig. 1C). Reduction of either isoform was also associated with a
slight increase in the accumulation of *OsRBCS3* and *OsRBCS5* transcripts (Fig 1B & 1D). These
results indicate that the *OsRBCS2* and *OsRBCS4* genes were selectively suppressed by the
antisense approach. Correspondingly immunoblot analysis showed a reduction in the
accumulation of RBCS in both *OsRBCS2* and *OsRBCS4* knockdown lines and no detectable
RBCS protein in event *rbcs2*-266 and *rbcs4*-53 (Fig. 2B). There are coordinated reductions of
RBCS and rbcL in both *rbcs2* and *rbcs4* lines and rbcL protein were reduced about 28% in
rbcs2-165, 41% in *rbcs2*-266, 50% in *rbcs4*-022 and 39% *rbcs4*-53 compared to wild-type
plants (Fig. 2A; Table 1). Immunolocalization confirmed reduction of Rubisco protein
preferentially in the MCs (Fig. S3). Reduction in Rubisco content was correlated with the
number of antisense inserts. *rbcs2*-266 and *rbcs4*-022 carried four and three antisense inserts,

respectively, and their Rubisco protein contents were lower comparable to the events carrying
fewer antisense inserts.

Photosynthetic perturbations associated with reduced Rubisco

The response of the net rate of CO_2 assimilation (A) to intercellular CO_2 concentration (Ci) was
measured under photorespiratory (21% O_2; Fig. 3A) and non-photorespiratory conditions (2%
O_2; Fig. 3B) at high irradiance (2,000 µmol photon m^{-2} s^{-1}) where RuBP-saturated rates of
Rubisco are limiting to photosynthesis. Under photorespiratory conditions, A was significantly
lower in both $rbcs2$ and $rbcs4$ lines at all intercellular CO_2 concentrations compared to wild-type
rice, although photosynthesis was not saturated in any of the lines. Rates of CO_2 assimilation
were much lower in $rbcs4$-022 and $rbcs2$-226 lines compared to the other lines for the same
gene; this was related to percentage reduction in Rubisco protein as $rbcs4$-22 and $rbcs2$-226 had
lowest Rubisco content among four $rbcs$ knockdown lines (Fig. 1). The responses under non-
photorespiratory conditions were similar to those under photorespiratory conditions, except for a
slight decrease in curvature and convergence in $rbcs2$-165 and wild-type response at the highest
intercellular CO_2 concentrations (Fig. 3B). These results suggest that photosynthesis was limited
by Rubisco content under almost all conditions. Consistent with this, carboxylation efficiency
(CE) was significantly reduced under both conditions, but most markedly under photorespiratory
conditions (Table 2), indicative of Rubisco limited capacity in the leaf. CO_2 compensation points
($\mathit{\Gamma}$) were significantly higher in event $rbcs2$-266 and $rbcs4$-022 compared with wild-type plants
under both conditions (Table 2). In response to changes in irradiance, CO_2 assimilation in wild-
type plants was not saturated at 2,000 µmol photon m^{-2} s^{-1} (Fig. 3C). In the knockdown lines,
photosynthesis was saturated at a of PPFD~1000 µmol photon m^{-2} s^{-1}, with CO_2 assimilation rates
less than half those of wild-type plants. There was no discernable relationship between CO_2
assimilation rate and Rubisco protein reduction, although there were line specific differences,
with notably higher rates in $rbcs4$-053 compared to other antisense lines. Quantum efficiency (Φ)
and respiration rates (R_d) were statistically significantly lower in all $rbcs$ lines compared with
wild-type plants (Table 2).

Phenotypic perturbations associated with reduced Rubisco

Reduction of Rubisco protein in *rbcs2* and *rbcs4* knockdown lines led to a statistically significant reduction in grain yield of between 37 to 70% (Table 3), with considerable variation observed between individual lines. In all lines except *rbcs2*-165, there was a corresponding statistically significant reduction in dry biomass, with *rbcs4* lines exhibiting the largest decrease relative to the wild-type plants. There were no statistically significant differences in tiller number but *rbcs4* lines were slightly shorter than wild-type plants (Table 3). There were no consistent differences in leaf chlorophyll contents, with small but statistically significant reductions in *rbcs4* lines and a higher chlorophyll content in *rbcs2*-266. The difference in phenotype perturbations between lines follows protein abundance with *rbcs2*-165 exhibiting the lowest Rubisco protein reduction and smallest phenotypic perturbations and *rbcs4*-022 the largest Rubisco protein reduction and most significant phenotypic perturbations.

Effect of reduced Rubisco on leaf anatomy

There were statistically significant differences in interveinal distance (IVD) between individual *rbcs* knockdown lines (Table 4), with *rbsc2*-165 having the widest IVD and *rbcs4*-022 line the lowest. Mesophyll cell number was significantly reduced in all the *rbcs* knockdown lines. The mesophyll cell length was increased in all *rbcs* knockdown lines but was only statistically significant in *rbcs2*-165 (Table 4). No difference in the number and thickness of major or minor veins were found. Leaves of both *rbcs4* knockdown lines were significantly narrower than either the wild-type or *rbcs2* lines, which correspond with reduced IVD and MC number.

Discussion

To mimic the downregulation of part of the Calvin–Benson cycle in MCs that is required for C_4 photosynthesis, we have successfully targeted two *OsRBCS* genes, *OsRBCS2* and *OsRBCS4*. Genetic redundancy within the *OsRBCS* gene family does not completely compensate for a reduction in the expression of a single multigene family member (Kanno et al., 2017, Ogawa et al., 2012). Constructs designed to independently reduce expression of *OsRBCS2* and *OsRBCS4* in the MCs led to a significant reduction in gene transcript for the target gene with modest

increases in the other multigene family members (Fig. 1). These results are consistent with
previous studies showing that the expressions of each *OsRBCS* genes were regulated
independently from other *OsRBCS* genes (Kanno et al., 2017, Ogawa et al., 2012). In leaf blades
of wild-type plants *OsRBCS2* transcripts accumulate to similar levels as those of *OsRBCS4*
(Suzuki et al., 2009). However, the effect of suppression of *OsRBCS2* antisense was stronger
with a larger reduction in *OsRBCS2* transcript accumulation in *rbcs2* knockdown lines. In *rbcs2*
knockdown lines Rubsico protein accumulation was 30-40% lower, which is consistent with
reports from previous studies where Rubisco content was reduced by up to 45% in *OsRBCS2*
cDNA antisense rice lines (Makino et al., 1997; Makino et al., 2000). Although we did not
generate lines targeting all the *OsRBCS* multigene family, these results suggest that targeting
multiple genes is required to completely reduce Rubisco accumulation in MCs.

Reducing Rubsico protein content by 30% or more in the antisense rice plants limits CO_2
assimilation rates under photorespiratory conditions (21% O_2). A relationship between the
reduction in Rubisco protein content and CO_2 assimilation rate was observed, which is consistent
with previous studies (Makino et al., 1997; Suzuki et al., 2012). On the other hand, other studies
also reported that a relatively small decrease in Rubisco content (65-90% of wild-type Rubisco)
can lead to an increase in CO_2 assimilation rate of between 5–15% at elevated CO_2 due to the
excessive amount of Rubisco for photosynthesis in rice (Kanno et al., 2017; Makino et al., 1997;
Makino et al., 2000). However here both *rbcs2* and *rbcs4* knockdown lines did not have higher
CO_2 assimilation rate than wild-type plants in elevated CO_2 conditions. The differences between
results in this study and previous studies may due to reduction of *OsRBCS2* and *OsRBCS4* being
preferentially targeted to the MCs leading to a more severe reduction in Rubisco content in the
MCs. On a protein quantity basis, the reduction of *rbcs2* lines had a more significant impact on
CO_2 assimilation rate than *rbcs4* lines given the percentage reduction in Rubisco protein.

A reduction in Rubisco protein content and CO_2 assimilation was associated with a significant
reduction in dry biomass accumulation and grain yield. The magnitude of the reduction was
proportional to the reduction in Rubisco protein content. The *rbcs* knockdown lines had no
difference in number of major and minor veins or leaf thickness compared to wild-type plants
(Table 4). Evan et al. (1994) reported a small reduction (92% of wild-type) in leaf thickness in
the *rbcs* antisense tobacco lines when Rubisco was removed by more than 50%. We also

observed that the numbers of MCs in between minor veins was reduced and the mesophyll cell length was slightly increased in *rbcs* knockdown lines (Table 4). This suggests that there may be anatomical changes to the leaf associated with altering Rubisco protein content.

Our results show that preferentially MC targeted antisense-*rbcs* plants can be used to reduce Rubisco accumulation in rice to levels that induce a phenotype without lethality. The resulting knockdown plants have reduced CO_2 assimilation rates with moderate effects on plant growth. This provides a useful foundation for installing C_4 photosynthesis in rice.

References

Andersson, I., & Backlund, A. (2008). Structure and function of Rubisco. Plant Physiology and Biochemistry 46: 275-291.

Dean, C., Favreau, M., Bedbrook, J., & Dunsmuir, P. (1989). Sequences 5' to translation start regulate expression of petunia *rbcS* genes. The Plant Cell, 1(2), 209-215.

Ermakova, M., Danila, F.R., Furbank, R.T., & von Caemmerer, S. (2020). On the road to C_4 rice: advances and perspectives. The Plant Journal, 101, 940–950. doi: 10.1111/tpj.14562.

Evans, J.R., von Caemmerer, S., Setchell, B.A., & Hudson, G.S. (1994). The Relationship Between CO_2 Transfer Conductance and Leaf Anatomy in Transgenic Tobacco With a Reduced Content of Rubisco. Australian Journal of Plant Physiology, 21(4), 475 – 495.

Furbank, R. T., Chitty, J. A., von Caemmerer, S., & Jenkins, C. L. (1996). Antisense RNA inhibition of *RbcS* gene expression reduces Rubisco level and photosynthesis in the C_4 plant *Flaveria bidentis*. Plant Physiology, 111(3), 725-734.

Ghannoum, O., Evans, J. R., & von Caemmerer, S. (2011). Nitrogen and water use efficiency in C_4 plants. In A. S. Raghavendra and R. F. Sage (Eds), C_4 Photosynthesis and Related CO_2 Concentrating Mechanisms, Advances in Photosynthesis. Springer Verlag, Berlin, 129–46.

345   Gibson, D. G., Young, L., Chuang, R. Y., Venter, J. C., Hutchison III, C. A., & Smith, H. O.
346   (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases.
347   Nature methods, 6(5), 343.

348   Hibberd, J. M., Sheehy, J. E., & Langdale, J. A. (2008). Using C_4 photosynthesis to increase the
349   yield of rice-rationale and feasibility. Current opinion in plant biology, 11(2),
350   228-231.

351   Hudson, G. S., Evans, J. R., von Caemmerer, S., Arvidsson, Y. B., & Andrews, T. J. (1992).
352   Reduction of ribulose-1, 5-bisphosphate carboxylase/oxygenase content by
353   antisense RNA reduces photosynthesis in transgenic tobacco plants. Plant
354   Physiology, 98(1), 294-302.

355   Kajala, K., Covshoff, S., Karki, S., Woodfield, H., Tolley, B. J., Dionora, M. J. A., ... & Quick,
356   W. P. (2011). Strategies for engineering a two-celled C_4 photosynthetic pathway
357   into rice. Journal of experimental botany, 62(9), 3001-3010.

358   Kanno, K., Suzuki, Y., & Makino, A. (2017). A small decrease in Rubisco content by individual
359   suppression of *RBCS* genes leads to improvement of photosynthesis and greater
360   biomass production in rice under conditions of elevated CO_2. Plant and Cell
361   Physiology, 58(3), 635-642.

362   Lin, H., Karki, S., Coe, R. A., Bagha, S., Khoshravesh, R., Balahadia, C. P., ... & Quick, W. P.
363   (2016). Targeted knockdown of GDCH in rice leads to a photorespiratory-
364   deficient phenotype useful as a building block for C_4 rice. Plant and Cell
365   Physiology, 57(5), 919-932.

366   Makino, A., Shimada, T., Takumi, S., Kaneko, K., Matsuoka, M., Shimamoto, K., ... &
367   Yamamoto, N. (1997). Does decrease in ribulose-1, 5-bisphosphate carboxylase
368   by antisense *RbcS* lead to a higher N-use efficiency of photosynthesis under
369   conditions of saturating CO_2 and light in rice plants? Plant physiology, 114(2),
370   483-491.

Makino, A., Nakano, H., Mae, T., Shimada, T. & Yamamoto, N. (2000). Photosynthesis, plant
growth and N allocation in transgenic rice plants with decreased Rubisco under
CO_2 enrichment, Journal of Experimental Botany, 51, 383–389.

Matsuoka, M., Kyozuka, J., Shimamoto, K., & Kano-Murakami, Y. (1994). The promoters of
two carboxylases in a C_4 plant (maize) direct cell-specific, light-regulated
expression in a C_3 plant (rice). The Plant Journal, 6(3), 311-319.

Morita, K., Hatanaka, T., Misoo, S., & Fukayama, H. (2014). Unusual small subunit that is not
expressed in photosynthetic cells alters the catalytic properties of Rubisco in rice.
Plant Physiology, 164(1), 69-79.

Osborn, H. L., Alonso-Cantabrana, H., Sharwood, R. E., Covshoff, S., Evans, J. R., Furbank, R.
T., & von Caemmerer, S. (2017). Effects of reduced carbonic anhydrase activity
on CO_2 assimilation rates in *Setaria viridis*: a transgenic analysis. Journal of
Experimental Botany, 68(2), 299-310.

Parry, M. A., Andralojc, P. J., Scales, J. C., Salvucci, M. E., Carmo-Silva, A. E., Alonso, H., &
Whitney, S. M. (2013). Rubisco activity and regulation as targets for crop
improvement. Journal of experimental botany, 64(3), 717-730.

Quick, W. P., Schurr, U., Fichtner, K., Schulze, E. D., Rodermel, S. R., Bogorad, L., & Stitt, M.
(1991). The impact of decreased Rubisco on photosynthesis, growth, allocation
and storage in tobacco plants which have been transformed with antisense *rbcS*.
The Plant Journal, 1(1), 51-58.

Rodermel, S. (1999). Subunit control of Rubisco biosynthesis–a relic of an endosymbiotic past?
Photosynthesis Research, 59(2-3), 105-123.

Rodermel, S. R., Abbott, M. S., & Bogorad, L. (1988). Nuclear-organelle interactions: nuclear
antisense gene inhibits ribulose bisphosphate carboxylase enzyme levels in
transformed tobacco plants. Cell, 55(4), 673-681.

Sasanuma, T. (2001). Characterization of the *rbcS* multigene family in wheat: subfamily
classification, determination of chromosomal location and evolutionary analysis.
Molecular genetics and genomics, 265(1), 161-171.

Sheehy, J.E., Ferrer, A.B., Mitchell, P.L., Elmido-Mabilangen, A., Pablico, P., & Dionora,
M.J.A. (2007). How the rice crop works and why it needs a new engine. In:
Sheehy JE, Mitchell PL, Hardy B, eds. Charting new pathways to C$_4$ rice. Los
Banos, Philippines: International Rice Research Institute, 3–26.

Suzuki, Y., & Makino, A. (2012). Availability of Rubisco small subunit up-regulates the
transcript levels of large subunit for stoichiometric assembly of its holoenzyme in
rice. Plant physiology, 160(1), 533-540.

Suzuki, Y., Nakabayashi, K., Yoshizawa, R., Mae, T., & Makino, A. (2009). Differences in
expression of the *RBCS* multigene family and Rubisco protein content in various
rice plant tissues at different growth stages. Plant and Cell Physiology, 50(10),
1851-1855.

Suzuki, Y., Ohkubo, M., Hatakeyama, H., Ohashi, K., Yoshizawa, R., Kojima, S., ... & Makino,
A. (2007). Increased Rubisco content in transgenic rice transformed with the
'sense' *rbcS* gene.

Suzuki, Y., Tomonori K. D., Kawazu, T., Miyake, C., & Makino, A. (2010). Differences in
Rubisco content and its synthesis in leaves at different positions in *Eucalyptus*
globulus seedlings. Plant, cell & environment, 33(8), 1314-1323.

von Caemmerer, S., & Quick, W. P. (2000). Rubisco: physiology in vivo. In Photosynthesis (pp.
85-113). Springer, Dordrecht.

von Caemmerer, S., Quick, W. P., & Furbank, R. T. (2012). The development of C$_4$ rice: current
progress and future challenges. Science, 336(6089), 1671-1672.

Whitney, S., Houtz, R. L., & Alonso, H. (2011). Advancing our Understanding and capacity to
engineer nature's CO$_2$-sequestering Enzyme, Rubisco. Plant Physiology, 155, 27-
35.

Yin, X., Anand, A., Quick, W. P., & Bandyopadhyay A. (2019). Editing a Stomatal
Developmental Gene in Rice with CRISPR/Cpf1. Yiping Qi (ed.), Plant Genome
Editing with CRISPR Systems: Methods and Protocols, Methods in Molecular
Biology, vol. 1917, 257-268.

Fig. 1. Transcript accumulation of four *OsRBCS* isoforms (*OsRBCS*2-5) as measured by quantitative real-time PCR in the leaf blade of *rbcs* knockdown lines. Values are expressed as fold changes in transcript accumulation relative to wild-type (mean ± SE). $2^{-\Delta\Delta CT}$ was used to quantify relative abundance against *OsEF-1α* transcript (Os03g0177500). Values are the mean ± SE of three leaves from three plants per genotype. Different letters within groups denote statistically significant differences , ANOVA *p*-value ≤0.05.

441 **A**

446 **B**

447 rbcL

448 RBCS

50 kDa

12 kDa

451 **Fig. 2.** Soluble leaf protein. (A) SDS-PAGE and (B) immunoblots with antibodies against rbcL
452 and RBCS. Wild-type (WT) and plants of *rbcs2* and *rbcs4* knockdown lines (numbers). Total
453 extracted proteins were separated by 12% SDS-gel. Samples were loaded based on equal leaf
454 area of 3.96 mm^2.

Table 1. Rubisco large subunit protein content.

	Percentage abundance to WT			
rbcs2-165	72	±	0.20	*
rbcs2-266	59	±	0.19	*
rbcs4-022	50	±	0.25	*
rbcs4-053	61	±	0.30	*

Values are the average ± SE of one leaf from four plants of *rbcs2* and *rbcs4* knockdown lines and wild-type (WT). Asterisk denotes statistically significant differences with WT, Student t-test, *p*-value ≤0.05.

Fig. 3. Net CO_2 assimilation rate (A) in wild-type and *rbcs2* and *rbcs4* knockdown lines. In response to intercellular CO_2 concentration (Ci) measured at (A) 21% O_2 and (B) 2% O_2. Measurements were made at a light intensity of 2,000 μmol photons m^{-2} s^{-1} and a leaf temperature of 30 °C. In response to photosynthetic photon flux density (PPFD; C). Measurements were made at CO_2 concentration of 400 μmol CO_2 mol air^{-1} and a leaf temperature of 30 °C. Values in legend are percentage reduction in Rubisco protein compared to the wild-type. Values are the mean ± SE of three leaves from three plants per genotype.

509 **Table 2. Comparison of photosynthesis parameters.**

	Γ	CE	R_d	Φ
	$\mu mol\ CO_2\ m^{-2}\ s^{-1}$	$\mu mol\ CO_2\ m^{-2}\ s^{-1}$ $\mu mol\ CO_2\ mol^{-1}$	$\mu mol\ CO_2\ m^{-2}\ s^{-1}$	
21 % O_2				
Wild-type	51.69 ± 1.59^c	0.126 ± 0.02^a	0.53 ± 0.05^a	0.046 ± 0.03^a
rbcs2-165	51.92 ± 0.15^c	0.067 ± 0.0^b	0.41 ± 0.03^b	0.039 ± 0.02^b
rbcs2-266	66.72 ± 1.45^a	0.046 ± 0.0^b	0.39 ± 0.07^b	0.031 ± 0.00^b
rbcs4-022	57.77 ± 1.58^b	0.05 ± 0.01^b	0.35 ± 0.02^b	0.030 ± 0.00^b
rbcs4-053	49.85 ± 2.71^c	0.06 ± 0.01^b	0.48 ± 0.05^b	0.025 ± 0.00^b
2 % O_2				
Wild-type	13.32 ± 0.95^c	0.166 ± 0.04^a	-	-
rbcs2-165	14.67 ± 1.11^c	0.086 ± 0.0^b	-	-
rbcs2-266	24.88 ± 1.54^a	0.057 ± 0.0^b	-	-
rbcs4-022	19.54 ± 0.71^b	0.06 ± 0.01^b	-	-
rbcs4-053	14.24 ± 3.12^c	0.08 ± 0.01^b	-	-

CO_2 compensation point (Γ), carboxylation efficiency (CE), respiration rate (R_d) and quantum
efficiency (Φ) of wild type and *rbcs2* and *rbcs4* antisense lines. Measurements for CE and Γ were
made at a photosynthetic photon flux density (PPFD) of 2,000 μmol photons $m^{-2}\ s^{-1}$, leaf
temperature 30 °C and either 21% or 2% O_2. Measurements of R_d were made after 30 min in the
dark. Measurements of Φ were made at a CO_2 concentration of 400 μmol $CO_2\ mol^{-1}$ and a leaf
temperature of 30 °C. Values represent the mean ± SE of three leaves from three plants per
genotype. Different letters within groups indicate statistical significance from a one-way
ANOVA (p≤0.05) and a least significant difference test (LSD) for post-hoc pairwise comparison.

Table 3. Leaf chlorophyll content, plant height, tiller number, dry biomass and grain yield of wild-type and *rbcs* knockdown lines.

	Chl (SPAD value)	Plant height cm	Tiller number	Dry biomass g	Grain yield g
Wild-type	43.14 ± 0.75[b]	105 ± 1.4[a]	26.1 ± 4.5[abc]	58.16 ± 10.93[a]	28.84 ± 7.68[a]
rbcs2-165	42.30 ± 0.71[bc]	105.6 ± 1.9[a]	29.2 ± 3.9[a]	57.36 ± 6.81[a]	18.07 ± 5.59[b]
rbcs2-266	44.04 ± 0.93[a]	103.6 ± 1.2[a]	27.5 ± 4.4[ab]	44.32 ± 5.12[b]	14.33 ± 4.52[b]
rbcs4-022	41.84 ± 1.20[c]	101.4 ± 1.8[b]	22.6 ± 4.9[bc]	32.60 ± 8.76[c]	8.90 ± 2.44[b]
rbcs4-053	42.00 ± 0.97[c]	99.4 ± 3.0[b]	23.4 ± 4.8[c]	33.19 ± 8.23[c]	13.80 ± 5.39[b]

Chlorophyll contents (SPAD value) are the average ± SE of three leaves from ten plants per genotype. Plant height, tiller number, dry biomass and grain yield are the average ± SE of ten individual plants per line; 77 days post-germination. Biomass is the total dry weight of leaf, stem and sheath tissue. Different letters within groups indicate statistical significance based on a one-way ANOVA ($p \leq 0.05$) with a Least Significant Difference (LSD) test for post-hoc pairwise comparison.

Table 4. Leaf width, leaf thickness, interveinal distance (IVD), vein number and mesophyll cell characteristics of wild-type and *rbcs* knockdown lines.

	Leaf width (cm)	Leaf thickness (µm)		IVD (µm)	No. of major vein	No. of minor vein	No. of Mesophyll cell	Mesophyll cell length (µm)
		Major vein	Minor vein					
Wild-type	1.4 ± 0.08^a	177.45 ± 7.86^a	88.82 ± 5.44^a	240.94 ± 11.75^{ab}	10.4 ± 0.9^a	40.2 ± 1.4^a	8.22 ± 0.4^a	23.82 ± 0.29^b
rbcs2-165	1.36 ± 0.06^a	175.10 ± 11.42^a	87.93 ± 4.63^a	245.80 ± 9.27^a	11.6 ± 0.5^a	40.8 ± 2.5^a	7.63 ± 0.5^b	26.68 ± 1.50^a
rbcs2-266	1.34 ± 0.05^a	167.05 ± 8.86^a	87.26 ± 3.12^a	230.11 ± 4.74^{bc}	11.6 ± 0.5^a	42.4 ± 3.7^a	7.31 ± 0.1^{bc}	25.35 ± 0.67^{ab}
rbcs4-022	1.16 ± 0.07^b	174.35 ± 7.98^a	85.82 ± 1.76^a	216.38 ± 7.11^c	10.6 ± 0.7^a	43.2 ± 1.0^a	7.12 ± 0.2^c	25.18 ± 0.84^{ab}
rbcs4-053	1.2 ± 0.09^b	159.95 ± 9.56^a	82.70 ± 3.71^a	226.89 ± 7.52^{bc}	10.6 ± 0.6^a	40.8 ± 2.6^a	7.53 ± 0.2^{bc}	24.50 ± 1.22^b

Leaf width is the average $\pm$ SE of two fully expanded penultimate leaves from ten individual plants per genotype. Leaf thickness of major veins is the average $\pm$ SE of four major veins of one fully expanded penultimate leaf from five individual plants per genotype. Leaf thickness of minor veins is the average $\pm$ SE of 10-12 minor veins of one fully expanded penultimate leaf from five individual plants per genotype. Interveinal distance (IVD) is the average $\pm$ SE of 5-6 transverse sections at both the left- and right-hand side of one fully expanded penultimate leaf from five individual plants per genotype. Major and minor vein numbers are the average $\pm$ SE of five individual plants per genotype. Mesophyll cell number and Mesophyll cell length (µm) are the average $\pm$ SE of 10-12 different sections of one fully expanded penultimate leaf from five individual plants per genotype. Different letters indicate statistical significance based on a one-way ANOVA ($p \leq 0.05$) with a Least Significant Difference (LSD) test for post-hoc pairwise comparison.

600 **Fig. S1.** PCR analysis of *rbcs2* and *rbcs4* knockdown lines (numbers). Wild-type (WT) was used as a negative control and the vector
601 plasmid (P) as positive control. B indicates blank (H$_2$O), M indicates a ladder (100bp).

Fig. S2. DNA blot analysis of *rbcs2* and *rbcs4* knockdown lines (numbers). The genomic DNA of each plant was extracted from leaves with the CTAB method and digested with *BgL*II. Wild-type plants (WT) were used as a negative control and the vector plasmid (P1 for antisense-*OsRBCS2* construct and P2 for antisense-*OsRBCS2* construct) as a positive control. A DIG-labeled DNA molecular marker indicates the molecular weight (MW).

Fig. S3. Representative images of immunolocalization of Rubisco protein in leaves from wild-type (WT), *rbcs2* and *rbcs4* knockdown lines. Anti-Rubisco rabbit polyclonal primary antibody diluted 1:200 plus Alexa Fluor 488 goat anti-rabbit IgG as secondary antibody diluted 1:200 was used to probe for rbcL (shown in green color). Chlorophyll is seen as a red autofluorescence. The cell wall was visualized by co-staining with calcofluor white and is shown in blue. Magnification: 200x. Scale bar: 20 μm. BSC: Bundle sheath cell. MC: mesophyll cell.