Exploiting oxalotrophic bacteria to control *Aspergillus niger* lung infections

Running title: Bacteria-induced control of aspergillosis

Fabio Palmieri[1], Ilona Palmieri[1], Nourine Noormamode[1], Aislinn Estoppey[1], M. Omar Ishak[2],

Julia M. Kelliher[2], Armelle Vallat[3], Rashi Iyer[4], Saskia Bindschedler[1], Karen Davenport[2],

Patrick S. G. Chain[2], Jennifer Foster Harris[2], Pilar Junier[1]*.

[1] Laboratory of Microbiology, Institute of Biology, University of Neuchâtel, Neuchâtel,

Switzerland.

[2] Bioscience Division, Los Alamos National Laboratory, Los Alamos, NM, USA.

[3] Neuchâtel Platform of Analytical Chemistry, University of Neuchâtel, Neuchâtel,

Switzerland

[4] Analytics, Intelligence and Technology Division, Los Alamos National Laboratory, Los

Alamos, NM, USA.

*Corresponding author

E-mail: pilar.junier@unine.ch (PJ)

Abstract

The opportunistic pathogen *Aspergillus* affects a record number of people worldwide. Many aspects of the ecology of *Aspergillus* spp. are linked to human pathogenesis, but not their ability to manipulate their environmental pH and enhance infectious potential via the secretion of oxalic acid. Here we showed that co-culturing with bacteria able to degrade oxalic acid (oxalotrophic bacteria) effectively suppress the pH shift induced by *A. niger*, and thereby arrest fungal growth. In 3D-cell culture systems infected with *A. niger*, co-inoculation with as little as 10 cells of an oxalotrophic bacterium was sufficient to return pH, Ca^{2+} and oxalic acid concentrations to the physiological values of uninfected cells. This is the first demonstration of the use of bacteria to control the infection potential of *A. niger*. This approach shifts the focus to the manipulation of the host environment as part of potential interventions to fight opportunistic fungal pathogens.

Introduction

Fungal diseases are estimated to kill more than 1.5 million people every year, with the number of affected patients expected to increase even further (Bongomin et al., 2017; Fisher et al., 2020). Indeed, over the last three decades, the intensification in the burden of fungal disease on human health has correlated with an increase in the number of individuals from at-risk populations (Ramirez-Ortiz and Means, 2012). As the size of these populations (for instance, immunosuppressed patients) is expected to keep increasing in the future (Bongomin et al., 2017), tackling fungal pathogenesis is urgent. However, only a very limited number of antifungal drugs is used nowadays to control fungal pathogens. Because of this restricted chemical arsenal, the same classes of molecules are used in human health, animal husbandry, and agriculture, leading to the rise, and rapid spread, of resistance in fungal pathogens that can

affect both plant and animal hosts (Fisher et al., 2018). In spite of this, tackling fungal disease

has been largely neglected (Brown et al., 2012).

The most prevalent fungal pathogens affecting humans are airborne opportunists such as

Aspergillus spp., *Cryptococcus* spp., *Pneumocystis* spp., and human-associated commensals

like *Candida albicans* (Brunke et al., 2016; Pfaller and Diekema, 2004). These fungal species

are responsible for approximately 90% of the deaths due to fungal infection (Fisher et al., 2020).

The ecology of all these organisms plays a very significant role in their ability to transition from

harmless to pathogenic. A good example of this is the genus *Aspergillus*, whose members are

widespread in the environment and are commonly isolated from both the outdoor and indoor

environments (Kousha et al., 2011; Paulussen et al., 2017). The remarkable plasticity in the

ecology and stress-response of *Aspergillus* spp. is believed to form the basis of its success as

an opportunistic pathogen (Paulussen et al., 2017). Indeed, species from this genus, and in

particular *Aspergillus fumigatus*, are thought to cause about 450'000 deaths worldwide every

year (Brown et al., 2012).

Although many aspects of the ecology of *Aspergillus* spp. have been connected to pathogenicity

(e.g. nutritional plasticity, reproductive success, and tolerance to environmental stressors), the

ability to manipulate pH via the secretion of low molecular weight organic acids, and in

particular, oxalic acid, has been largely ignored in the context of human pathogenesis. Several

clinical reports have shown the presence of calcium oxalate crystals in pulmonary aspergillosis

in animals and humans (Kurrein et al., 1975; Maeno et al., 2015; Muntz, 1999; Oda et al., 2013;

Payne et al., 2017; Yi et al., 2019). Oxalic acid and oxalate crystals are thought to directly cause

damage to the host tissues (including pulmonary blood vessels), and to generate free radicals

which can harm cells indirectly (Payne et al., 2017). A recent case report of invasive pulmonary

aspergillosis in a 69-year old man with lymphoma and pneumonia indicated the presence of

calcium oxalate crystals around blood vessels and within the blood vessel walls. This suggests

a potential mechanical role of oxalate crystals in the angioinvasion of *Aspergillus* (Yi et al.,

2019). However, a link between oxalic acid production and pathogenicity has not yet been made

in fungi from this genus or in any other fungal human pathogen; despite having been widely

acknowledged in the case of fungal plant pathogens, where pH manipulation by the secretion

of oxalic acid plays a direct role in pathogenesis (Dutton and Evans, 1996; Lehner et al., 2008;

Palmieri et al., 2019). Acidification of the local environment is exploited by plant pathogens to

weaken the cell wall structure, facilitate infection, inhibit plant defenses, and induce

programmed cell death (de Oliveira Ceita et al., 2007; Dutton and Evans, 1996).

Oxalic acid is a ubiquitous compound in the environment. Its production and consumption by

microorganisms have been directly associated with pH regulation in soil (Graustein et al.,

1977). In such ecosystems, oxalic acid is often found complexed with divalent cations,

especially calcium (Martin et al., 2012). Despite its chemical stability (K_{sp} calcium oxalate

monohydrate $= 2.32 \times 10^{-9}$), calcium oxalate rarely accumulates (Certini et al., 2000; Hofmann

and Bernasconi, 1998). This is because calcium oxalate is used by soil oxalotrophic bacteria as

both a carbon and energy source. The transformation of a strong organic acid into a weaker one

(oxalic acid versus carbonic acid; $pK_a = 1.25$ and 4.14, respectively), creates a local increase in

soil pH (Braissant et al., 2004).

We hypothesize that oxalic acid production may be needed for *Aspergillus* spp. to manipulate

pH during lung infection and that interfering with this process may limit pathogenicity. Such

intervention can be accomplished in two different ways. One method is to interfere with the

ability of the fungus to produce oxalic acid, for instance by knocking out one or more essential

genes in the synthesis pathway. Although valid under laboratory conditions, this option cannot

be considered for a therapeutic intervention. The second approach could aim at reducing the

concentration of oxalic acid *in situ* by directly removing this compound. This can be achieved

by using oxalotrophic bacteria. This approach offers an alternative that can be envisioned for a

future therapeutic intervention, based on the manipulation of the lung microbiota (Fig. 1).

Therefore, we chose to investigate this second alternative in order to offer a proof-of-concept

of the potential of using oxalotrophic bacteria to control the course of fungal pathogenesis.

To test our hypothesis, we selected *Aspergillus niger* as a model. This organism has been

extensively studied for its ability to produce oxalic acid (Cameselle et al., 1998; Ruijter et al.,

1999) and is a known agent of aspergillosis in humans (Paulussen et al., 2017). We first

confirmed that *A. niger* secretes oxalic acid and measured the effect of co-cultivation with

oxalotrophic bacteria *in vitro*. Then, we investigated the impact of co-cultivation on human

bronchial epithelial cells (HBEC) cultures where we determined a dose-response effect for both

the fungus and the bacteria. Finally, we used Transwells® inserts and bronchioles-on-a-chip

(BoC) systems to measure the effect of the fungus and the co-cultivation with an oxalotrophic

bacterium on pH, calcium concentration, oxalic acid concentration, and tissue integrity.

Results

Detection of oxalic acid produced by *Aspergillus niger* in different culture conditions

Since the metabolism of fungi can change significantly depending on the nutritional conditions

of the growth medium, we first tested whether the strain of *A. niger* used for the experiments

produces oxalic acid on different standard laboratory media. Ultra-High Performance Liquid

Chromatography (UHPLC) analysis confirmed that the fungus produced oxalic acid as the sole

low molecular weight organic acid when grown in malt and in Reasoner's 2 liquid media (Fig.

S1A). Acidification and the presence of calcium oxalate crystals could also be detected upon

growth in water yeast agar (WYA) (Fig. S1B). We thus concluded that *A. niger* consistently

produced oxalic acid and acidified its medium under laboratory growth conditions.

Confrontation assays between *A. niger* and selected bacterial strains

We proceeded to test how the presence of oxalotrophic bacteria impacts the growth of *A. niger* and the evolution of the pH in different growth media. Two oxalotrophic bacteria (*Cupriavidus necator* and C. *oxalaticus*) and one non-oxalotrophic bacterium (*Pseudomonas putida*) were used in confrontation assays. In malt agar diluted 10 times (MA 1/10), *A. niger* colonized the entire Petri dish, including the area in which the bacterial inocula were applied. All the three bacteria inoculated did not survive in the area of interaction with the fungus (Fig. S2A). In Reasoner's 2 agar (R2A), *A. niger* did not colonize the area beyond the barrier formed by the bacterial inoculum, but some microcolonies formed in the co-culture with *P. putida* beyond the bacterial inoculation zone (Fig. S2B). In water yeast agar (WYA), fungal growth was also restricted to the area delimited by the inocula in the co-culture with the two oxalotrophic bacteria, while *P. putida* did not survive the interaction with the fungus, which instead, colonized the entire plate. Moreover, in WYA medium, which contained a pH indicator, the acidification of the medium by the fungus grown alone or in co-culture with the non-oxalotrophic bacterium was clearly visible. In contrast, in the co-cultures with oxalotrophic bacteria, the pH of the medium did not change (Fig. S2C). The control of fungal growth was particularly remarkable in the case of *C. oxalaticus* (Fig. 2). The results in R2A and WYA validate our hypothesis *in vitro* by demonstrating that fungal growth and pH can be controlled by oxalotrophic bacteria.

Growth tests in lung cells culture medium and confrontation assay between *A. niger* and *C. oxalaticus*

In order to obtain evidence that oxalotrophic bacteria can control the growth of *A. niger* in a
lung infection model, we set up experiments on bronchial epithelial cell cultures. Given the
impact of the medium on the control of fungal growth, we repeated the growth and
confrontation assays in Air-Liquid Interface (ALI) medium, which is the medium used for
differentiation of lung cells (Fulcher et al., 2005). The non-oxalotrophic bacterial model (*P.*
putida) acidified the medium after growth, while the oxalotrophic bacteria (*C. necator* and *C.*
oxalaticus) did not acidify the medium (Fig S3A). *C. oxalaticus* was found to also inhibit
conidia germination and mycelial development (Fig. S3B). Moreover, the presence of *C.*
oxalaticus in co-culture with the fungus stabilized the pH of the culture medium at a neutral pH
(Fig. S3C), consistent with decreased oxalic acid concentration in the medium (Fig. S3D). We
therefore conclude that oxalotrophic bacteria have a significant inhibitory effect on *A. niger*
growth in ALI medium.

Establishment of a dose-response curve on submerged cultures

We established a dose-response curve for increasing conidial and/or bacterial loads on the
morphology of HBECs in submerged cultures. This aimed at identifying the optimal conidial
and/or bacterial load to perform additional experiments in Transwells® and BoC systems. After
24 hours, the overall size, shape and integrity of the lung cells changed at an absolute load of
500 conidia and above. The HBECs shrank in size due to actin agglomeration. Moreover, from
a conidial load of >=1000, fungal growth also had an adverse effect on tissue integrity (Fig.
S4). The same experiment was performed with *C. oxalaticus*. A strong morphological change
was induced by a total load of 500 bacterial cells and above. The HBECs became rounder, and
actin agglomeration increased compared with the cells-only control (Fig. S5). This indicates
that a dose of either 500 conidia or 500 bacterial cells is detrimental. To analyze the effect of

co-culturing *A. niger* with the oxalotrophic bacterium on HBECs integrity, we performed a test

with 10 and 500 conidia confronted with 10 bacterial cells. After 72h, *A. niger* induced

morphological changes (size reduction and actin agglomeration), with a stronger effect for 500

conidia, confirming the results obtained in the 24 h experiment. With the co-inoculation of as

few as 10 *C. oxalaticus* cells, the morphology of the HBECs was similar to the morphology of

HBECs of bacteria-only control, suggesting the inhibition of fungal development (Fig. S6). We

concluded that a conidial and bacterial load of 10 conidia/cells was ideal to monitor the

interaction of *A. niger* and *C. oxalaticus* in differentiated HBECs in Transwells® and BoC

systems.

Biocontrol assay of *A. niger* bronchial cells infection by *C. oxalaticus*

After establishing a dose-response curve on HBECs in submerged cultures (Fig. S4-S6), the

effect of inoculation of 10 conidia of *A. niger* alone or in co-culture with 10 *C. oxalaticus* cells

was assessed in differentiated bronchial tissue in Transwells® inserts and BoC systems. In the

presence of the fungus alone, changes in three key environmental factors were observed: pH,

Ca^{2+} concentration, and concentration of soluble oxalic acid. The pH dropped significantly from

7.5 down to 4.5 in Transwells® and from 7.3 to 6.8 in BoC systems (Fig. 3A). Ca^{2+}

concentrations changed from 1 mM to around 0.2 mM in both culture systems (Fig. 3B). The

level of soluble oxalic acid dropped from 500 μM (cells alone) to 75 μM (cells inoculated with

the fungus) (Fig. 3C). pH, Ca^{2+} and free oxalic acid levels were statistically indistinguishable

when oxalotrophic bacteria were co-cultured with the fungus or in the controls with cells and

bacteria or cells alone. In addition, calcium oxalate crystals were observed in the cultures in

which the fungus developed (Fig. 3D), but not when the fungus was co-cultured with

oxalotrophic bacteria (Fig. 3E). This suggests that the lower levels of soluble oxalic acid

measured in the treatment with the fungus were likely the result of complexation of oxalic acid

and Ca^{2+} and corroborates the pH and Ca^{2+} concentration data. Moreover, the absence of

calcium oxalate crystals when the fungus was in co-culture with *C. oxalaticus* agrees with the

pH, Ca^{2+} concentration and soluble oxalic concentrations measured. These results validated our

hypothesis that oxalotrophic bacteria can be used to manipulate the microenvironment created

by *A. niger* and to hamper its development.

In addition to the changes in the environmental parameters measured above, we also observed

a cytopathic effect when conidia of *A. niger* developed into mycelia (Fig. 4A and B). The

absence of lactate dehydrogenase (LDH) activity measured in the medium in the treatment with

A. niger (Fig. 4C), supports the cytopathic effect and aggressiveness observed upon infection

by *A. niger*. LDH measurements were elevated in response to the presence of oxalotrophic

bacteria. Indeed, *C. oxalaticus* is a soil bacterium that is completely foreign to the lung

environment, thus potentially inducing an inflammatory response, something that needs to be

addressed for any future therapeutic application.

Discussion

By demonstrating the biocontrol activity that oxalotrophic bacteria exerted on *A. niger*, our

study represents a novel paradigm in which the manipulation of the lung environment can

become the focus of an anti-fungal medical intervention. This contrasts with the prevailing use

(and abuse) of antimicrobial compounds, which are instead directed to the pathogen and

inevitably result in the emergence of resistance (Toprak et al., 2011). To our knowledge, this is

the first demonstration of such an alternative approach for combatting human fungal infections

and is a significant step towards an integrated disease management strategy that is urgently

required to slow the emergence of resistance in fungal pathogens. Moreover, this approach can

be also relevant to *A. fumigatus*, the major causative agent of pulmonary aspergillosis, as the

presence of calcium oxalate crystals has been reported in the literature, suggesting also the

production of oxalic acid production by this pathogen (Kousha et al., 2011; Maeno et al., 2015;

Pabuçcuoğlu, 2005; Payne et al., 2017). Currently there is an increasing interest in applying

ecological theory in lung microbiome research (Foesel et al., 2019). Although considered sterile

for a long time, the lung is now known to harbor a diverse microbiota (Enaud et al., 2020;

Mitchell et al., 2016). In the healthy lung, this microbiota is dominated by the genera *Prevotella*,

Streptococcus and *Veillonella* (Charlson et al., 2011; Foesel et al., 2019). Its composition differs

in a diseased state (Cui et al., 2015; Dickson et al., 2014; van Woerden et al., 2013) and after

transplantation (Charlson et al., 2012). This dysbiosis of the lung microbiota, which can be

associated with comorbidities in at-risk patients, could make the lung environment more

permissive to fungal pathogens such as *Aspergillus* spp. This could in part explain the current

epidemic of nosocomial fungal infections (Suleyman and Alangaden, 2016).

Oxalate-degrading capabilities have been previously reported in strains of the genera

Lactobacillus (Turroni et al., 2007), *Streptococcus* (Miller and Dearing, 2013), *Prevotella*

(Suryavanshi et al., 2016; Ticinesi et al., 2020) and *Veillonella* (Suryavanshi et al., 2016), all

of which are represented in the lung microbiota. In this context, assessing the oxalotrophic

potential of the lung microbiome and its subsequent stimulation could offer an endogenous

alternative to fight aspergillosis. Indeed, as aspergillosis mainly affects immunocompromised

patients, the administration of an exogenous bacterial species as therapeutic intervention could

be unviable. Nevertheless, complementing the microbiota with a specific bacterial strain, or

even a consortium, has been proven successful in protecting the host from *Clostridium difficile*

infection in a mouse model (Buffie et al., 2015; Lawley et al., 2012). Another potential venue

is the administration of a completely abiotic delivery system of oxalate-degrading enzymes

directly in the lungs by inhalation.

Our results also provide an indication of potential additional roles of oxalic acid in fungal

pathogenesis in humans. Secretion of oxalic acid by the fungus has a clear effect on the

concentration of free Ca^{2+} in the medium. Calcium is known to be a secondary messenger in

many cell types, playing a central role in the activation of the immune response (Vig and Kinet,

2009). During immune stimulation, the concentration of intracellular Ca^{2+} in immune cells

increases thanks to Ca^{2+} entry from the extracellular media or cellular compartments (Grinstein

and Klip, 1989). Accordingly, the use of a chelating agent of divalent cations has been shown

to inhibit the immune response *in vitro* (Diamantstein and Odenwald, 1974; Diamantstein and

Ulmer, 1975). This suggests that chelation of Ca^{2+} in calcium oxalate crystals upon fungal

infection could inhibit the immune response. Moreover, the detection of calcium oxalate

crystals is also meaningful in the case of a chronic inflammatory response, as calcium oxalate

crystals have been involved in inflammation in renal cells (Mulay et al., 2013). A recent study

also shows that calcium oxalate crystals induce the production of inflammatory cytokines in

primary and THP-1 monocytes (Dominguez-Gutierrez et al., 2018). Finally, a recent study from

McClintock *et al.* (McClintock et al., 2020) showed the role of calcium in epithelial tissue

cohesion and barrier function, which could be disrupted upon fungal infection and subsequent

Ca^{2+} sequestration in the form of calcium oxalate.

The results presented here are the stepping-stones for the development of a pre-clinical *in vivo*

animal model and the subsequent validation of the proposed approach in order to translate these

results into a potential clinical application. Indeed, a more holistic approach with the integration

of the immune system components will be key to understand the interplay between the host, the

lung microbiota, the environment, and the pathogen.

Materials and Methods

Bacterial and Fungal cultures

All bacterial and fungal strains used in this study come from the collection of the Laboratory of Microbiology of the University of Neuchâtel (LAMUN) and are summarized in Table 1. *P. putida* KT2440 was kindly provided by Dr. Arnaud Dechesne (Technical University of Denmark). *C. necator* JMP289 was kindly provided by Prof. Jan van der Meer (University of Lausanne). *C. oxalaticus* Ox1 was tagged in-house using insertion with a MiniTn7 system. Bacterial strains were routinely cultured on Nutrient Agar (NA, Carl Roth, Karlsruhe, Germany) medium, whereas *Aspergillus niger* was routinely cultured on Malt Agar (MA) medium. Potato Dextrose Agar (PDA, Carl Roth, Karlsruhe, Germany) was used for *A. niger* conidia production. MA medium was composed of 12 g/L malt extract (Sios Homebrewing GmbH, Wald, Switzerland) and 15 g/L agar (Biolife Italiana, Milano, Italy). Brain Heart Infusion Agar (BHIA, Brain Heart Broth, Sigma-Aldrich, Darmstadt, Germany, 15 g/L agar) was used to have mycelium-only colony edge without any conidia.

Table 1. Bacterial and fungal strains used

Code	Strain #	Species	Fluorescent tag	References
Pp	NEU 1264	*Pseudomonas putida* KT2440	GFP	
Cn	NEU 1286	*Cupriavidus necator* JMP289	GFP	(Sentchilo et al., 2009)
Co	NEU 1287	*Cupriavidus oxalaticus* Ox1	mCherry	(Şahin et al., 2000)

An	NEU M8	*Aspergillus niger*		-	(Ferro, 2012)

All the bacterial and fungal strains used in this study come from the collection of the Laboratory

of Microbiology of the University of Neuchâtel.

Low Molecular Weight Organic Acids (LMWOA) detection by UHPLC and by

colorimetric pH indicator-based Petri dish assay

In order to highlight the production of LMWOA by *A. niger*, we used Ultra-High-Performance

Liquid Chromatography (UHPLC), as well as a culture-based assay by using a pH indicator-

containing agar culture medium.

For the UHPLC analysis, 500 µl of 30 mM H_2SO_4 were added to 1 mL of a two-week liquid

culture in malt 1/10 and Reasoner's 2 liquid media in triplicate, to obtain 20 mM H_2SO_4 final

concentration. The samples were incubated at 60°C for two hours in order to dissolve

precipitated metal oxalate crystals, and then centrifuged at 3000 g for 10 min. All the samples

were filtered at 0.22 µm (13mm syringe filters, PTFE, hydrophilic) and 200 µl were added into

HPLC vials with 250 µl conical inserts. UHPLC (Ultimate 3000 RS-Dionex, Thermo Fisher

Scientific, MA, USA) was coupled with DAD detector set at 210 ± 2 nm. A 5 µL of sample

was injected onto a Prevail™ organic acid column (5 µm particle size, 150 x 4.6 mm, Grace

Davison Discovery Sciences, Deerfield, IL, USA) with the temperature kept at 40°C. The

mobile phase consisted of 50 mM phosphate buffer adjusted to pH 2.5 with phosphoric acid

with a flow rate of 1 mL/min. Pure oxalic acid (Merck KGaA, Darmstadt, Germany) was

identified by the retention time and was quantified by an external standard curve, linear

regression from five calibration points (0.2 to 5 mg/mL).

For the culture-based assay, water yeast agar supplemented with bromocresol purple

(WYA+BP) (Mela et al., 2011) was used as a pH indicator-containing medium. WYP+BP was

composed of 1 g K_2HPO_4, 5 g NaCl, 0.1 g yeast extract, 10 mg bromocresol purple and 20 g

agar, per liter of Milli-Q® water. After one week of incubation at room temperature, the

presence of typical bi-pyramidal shaped calcium oxalate crystals was assessed by observing a

thin slice of agar medium sampled at the edge of the colony and stained with lactophenol cotton

blue under a Leica DM4 B optical microscope connected to a Leica DFC7000 T camera.

Confrontation assays on solid media

Confrontations assays were performed between *A. niger* and three bacterial strains:

Pseudomonas putida, *Cupriavidus necator* and *Cupriavidus oxalaticus*, on three culture media

differing in their nutrient content. Malt agar diluted ten times (MA 1/10) was used as nutrient-

rich medium and was composed of 1.2 g/L malt extract (Sios Homebrewing GmbH, Wald,

Switzerland) and 15 g/L (Biolife Italiana, Milano, Italy). Reasoner's 2 agar (R2A) (Reasoner

and Geldreich, 1985) was used as nutrient-poor medium and the composition per liter of Milli-

Q® water was the following: 0.5 g yeast extract, 0.5 g Bacto Peptone, 0.5 g casamino acids,

0.5 g glucose, 0.5 g soluble starch, 0.3 g Na-pyruvate, 0.3 g K_2HPO_4, 0.05 g $MgSO_4 \cdot 7H_2O$

and 15 g agar. WYA+BP (Mela et al., 2011) was also used. *A. niger* was inoculated in the center

of the plates containing the different culture media. For that, a plug coming from the apical part

of an active colony was sampled using the wider end of a Pasteur pipette. The three different

bacterial strains were inoculated as opposite lines on either side of the fungal inoculum. Plates

were incubated at room temperature for 20 days, and pictures of the plates were taken. Pictures

of the bacterial inocula were taken using a Nikon SMZ18 epifluorescence stereoscope,

connected to a Nikon DS-Ri2 camera, in order to assess the viability of the bacterial strains

thanks to constitutively expressed fluorescent proteins.

Growth tests and confrontation assay in Air-Liquid Interface (ALI) medium

Air-Liquid Interface (ALI) medium was composed of DMEM/F12 (Cat.# 11320033, Thermo

Fisher Scientific) and LHC Basal Medium (Cat.# 12677019, Thermo Fisher Scientific) at a 1:1

ratio according to Fulcher *et al.* (Fulcher et al., 2005). Bacterial growth in ALI medium was

tested for 3 days at room temperature. Increasing concentrations of Ca^{2+}, from 1 mM to 10 mM,

was also tested on bacterial and fungal growth in a 96-well plate. Given that ALI medium

contains 1 mM Ca^{2+}, a 10 mM Ca^{2+} stock solution of ALI medium was prepared by adding 1.32

g/L $CaCl_2$ to have an added Ca^{2+} concentration of 9 mM. This stock solution was used to prepare

8, 7.5, 6, 4.5, 3, and 2.5 mM dilutions of $CaCl_2$ ALI-containing media. To produce fungal

conidia, *A. niger* was cultured on PDA for 10 days at room temperature. Conidia were then

harvested by using Dulbecco's Phosphate Buffer Saline (DPBS) supplemented with 0.01%

(v/v) Tween 80. Harvested conidia were washed tree times with DPBS following centrifugation

at 2000xg for 5 min at room temperature. Finally, conidia were resuspended in 2 mL DPBS and

quantified with an Improved Neubauer counting chamber. Two μL of 16'000 conidia/bacterial

cell per μL suspensions were inoculated in 200 μL ALI medium containing the Ca^{2+}

concentrations indicated above. To test mycelial growth in the different media, small agar plugs

(approximately 3 x 3 mm) coming from the edge of a colony of *A. niger* on BHIA were used.

The 96-well plate was incubated at room temperature for 7 days and growth was visually

assessed.

Confrontation of *A. niger* with *C. oxalaticus* was performed in 100-mm Corning® tissue culture

plate containing 12 mL ALI medium to allow the fungus to attach during growth. The fungal

inoculum was taken from the apical part of an active colony grown on BHIA by using the wider

end of a Pasteur pipette. For the confrontation assay, the fungal inoculum was placed in the

medium after addition and mixing of 100 μL inoculum of an overnight culture of *C. oxalaticus*

in ALI medium. A fungus-only plate was used as control. The plates were incubated at room

temperature for 7 days. Oxalic acid concentration was quantified by using the Oxalic Acid

Colorimetric Assay Kit (Sigma-Aldrich, Darmstadt, Germany, Cat.# MAK179), following the

manufacturer instructions.

Preparation and sterilization of the bronchiole-on-a-chip (BoC)

The chip was assembled as described in Hsieh, Nath and Huang (Hsieh et al., 2019). Each unit

of the culture platform was fabricated by using a layer-by-layer stacking technique (Lin et al.,

2019). The devices were designed using Solid Edge 2D software (ST9, Siemens PLM

Software), and each layer of the prelaminated polymeric sheet was obtained using a CO_2 laser

cutter (Universal Laser System). The prelaminated polymeric sheets were combined with

biocompatible adhesive tapes (9122, 3M Company) with PMMA (1.5 and 3 mm thick) or PET

(0.1 and 0.25mm thick). After cutting, each layer was aligned and assembled using a seam roller

to complete the devices. The culture chip includes a Y-shaped apical and a basal part separated

by a porous PET membrane (pore size = 0.4μm) prepared as described in Arefin *et al.* (Arefin

et al., 2018). The PET membrane was sandwiched between two PET sheets using adhesive

transfer tape to create the cell culture surface. This allows nutrients to pass from the media to

the cells through the porous membrane. The open design of the tissue chip makes cell seeding

procedure easy and accessible.

For sterilization, each chip was placed in a 100 mm Petri dish and sterilized with 5% hydrogen

peroxide solution for one hour. The chips were then rinsed 2-3 times with sterile deionized

water for 15 min between each rinse. Once all liquid was removed, the chips were let dry overnight under a laminar flow hood. The next day, the inlet and outlet of the chip were connected with a sterile tubing and rinsed 3 more times with sterile deionized water as explained before. After the last rinse, 200 μl sterile DPBS was added in the channel (apical part) and 5 mL in the basolateral part (bottom part) of the chip, and the chip was placed in a humidified incubator at 37°C with 5% CO_2 overnight. The next morning, peroxide contamination was checked in each chip using a CG8+ i-STAT cartridge (Abbott, USA, Cat.# 03P88-25). Rinses with sterile deionized water and overnight incubation with sterile DPBS were repeated until peroxide was no longer detected in the devices.

Primary normal human bronchial epithelial cell culture

Primary normal human bronchial epithelial cells (Lifeline Cell Technology, Cat.# FC-0035) were expanded in a T-75 cell culture flask with vent cap (Corning, Cat.# 3290) containing 13 mL BronchiaLife™ B/T complete medium (Lifeline Cell Technology, Cat.# LL-0023) supplemented with 3 mL/L 0.5% Phenol Red solution (Sigma-Aldrich, Cat.# P0290, 15 mg/L final concentration) to 70-80% confluence in a humidified incubator at 37°C with 5% CO_2. Culture medium was changed every other day. Cells were used until passage 2 maximum for all experiments. Cells were then harvested by trypsinization with 0.05% Trypsin / 0.02% EDTA (Lifeline Cell Technology, Cat.# CM-0017), followed by the addition of Trypsin Neutralizing buffer (Lifeline Cell Technology, Cat.# CM-0018), and counted using a hemocytometer after centrifugation at 100xg for 5 min and resuspension of the cell pellet in BronchiaLife™ medium.

For submerged undifferentiated tissue culture in 96-well plates, 30'000 cells were seeded in 200 μl BronchiaLife™ medium per well. The 96-well plate was placed in a humidified

incubator at 37°C with 5% CO_2 for 2-3 days until confluence and formation of a monolayer of

bronchial cells.

For air-lifted differentiated tissue culture, Transwell® inserts in 24-well plates (Corning, Cat.#

3470) and BoC devices have first to be coated with collagen in order to allow proper cell

attachment onto the porous membrane. For that, 200 μL of 30 μg/mL rat collagen (Corning,

Cat.# 354236) diluted in DPBS were added to the apical side of Transwell® inserts and BoC

devices and the latter have been placed at 37 °C for 4 h in a 5% CO_2 humidified incubator.

Excess collagen was then removed, and all inserts and devices were rinsed once with sterile

DPBS and stored in a refrigerator until seeding of the cells.

Transwell® inserts and BoC devices were seeded on the apical side with 50'000 cells and

86'000 cells in 200 μL BronchiaLife™ complete medium, respectively. 600 μL of

BronchiaLife™ complete medium was added to the basolateral side of Transwell® inserts in

the 24-well plates and 3 mL in the basolateral side of the BoC device. All the inserts and device

were then placed in a humidified incubator at 37°C with 5% CO_2 for 2-3 days until confluence.

When bronchial cells formed a monolayer, they were shifted to air-liquid interface.

BronchiaLife™ complete medium was removed carefully from the apical part and replaced by

Air-Liquid Interface (ALI) Epithelial Differentiation Medium (Lifeline Cell Technology, Cat.#

LM-0050) supplemented with 3 mL/L 0.5% Phenol Red solution (Sigma-Aldrich, Cat.# P0290,

15 mg/L final concentration). The same was done for the medium in the basolateral side.

Finally, the medium on the apical side was removed and the inserts and devices were placed in

a humidified incubator at 37°C with 5% CO_2 for 21 days. Medium was changed every other

day as described previously. The cultures were observed daily using an EVOS™ XL Core

bright field inverted microscope (Thermo Fisher Scientific, CA, USA).

Determination of conidial and bacterial load and confrontation assay on submerged undifferentiated bronchial epithelial cell cultures

In order to determine the optimal conidial and bacterial load to be used for confrontation on bronchial tissue cultures, increasing conidial and bacterial loads were tested to assess their effect on the morphology of bronchial epithelial cells in submerged cultures. *A. niger* was cultured on PDA for 7 days at 37°C in order to produce conidia. *A. niger* conidia were then harvested as already described. Finally, conidia were resuspended in 2 mL DPBS and quantified with an Improved Neubauer counting chamber. Bacteria were cultured in BronchiaLife™ complete medium at 37°C overnight and quantified with an Improved Neubauer counting chamber. A stock suspension of *A. niger* conidia was made at 1'000'000 conidia/mL that was diluted further to obtain suspensions at 500'000, 100'000, 50'000, 10'000, 5000 and 1000 conidia/mL. The same was done for *C. oxalaticus* from a stock suspension at 100'000 bacterial cells/mL diluted until 1000 bacterial cells/mL. 10 μL of each suspension was added to submerged undifferentiated bronchial tissue in a 96-well plate in order to have 10'000, 5000, 1000, 500, 100, 50 and 10 conidia/well (200 μL) for *A. niger*, and 1000, 500, 100, 50, 10 bacterial cells/well (200 μL) for *C. oxalaticus*. The plates were placed in a humidified incubator at 37°C with 5% CO_2 for 24h. For the confrontations assay, cells were infected with 10 or 500 *A. niger* conidia and were put in confrontation with 10 bacterial cells. Controls with only cells, fungus or bacteria were also prepared. The plate was placed in a humidified incubator at 37°C with 5% CO_2 for 72h.

After 24h or 72h incubation, the cells were fixed by adding 100 μL 4% paraformaldehyde in DPBS per well and the plates were let sit for 15 min at room temperature. The cells were then rinsed 3 times, with 2 min waiting time between each rinse. Two drops of ActinGreen™ 488 ReadyProbes™ Reagent (actin stain, Thermo Fisher Scientific, Cat.# R37110) and one drop of NucBlue™ Live ReadyProbes™ Reagent (nuclei counterstain, Thermo Fisher Scientific, Cat.

R37605) was added in 1 mL of DPBS to prepare the staining solution. 200 μL of staining

solution were added per well and the plates were let sit for 30 min at room temperature. Finally,

the cells were washed 3 times with DPBS before imaging with a Zeiss Axio Observer Z1

fluorescence inverted microscope (Carl Zeiss AG, Oberkochen, Germany).

Confrontations on differentiated bronchial tissues in Transwell® inserts and BoC devices

Differentiated bronchial tissues in Transwell® inserts and in BoC devices were infected with

10 μL of 1000 conidia or bacterial cell to get 10 conidia or bacterial cells. Fungus and bacteria

were co-inoculated for the confrontation and controls with only medium, cells, fungus or

bacteria were included. Each condition was done in triplicate for the Transwell® inserts and

one unique replicate for each condition was done for the BoC devices. All Transwell® inserts

and BoC devices were incubated in a humidified incubator at 37°C with 5% CO_2 for 72h.

Immunofluorescence staining

Differentiated bronchial tissues in Transwell® inserts and BoC devices were fixed by adding

100 μL 4% paraformaldehyde in DPBS per insert or device and the 24-well plate and devices

were let sit for 15 min at room temperature. The cells were then rinsed 3 times with 200 μL

DPBS, with 2 min waiting time between each rinse. Cells were permeabilized with 100 μL

0.5% Triton X-100 in DPBS for 15 min at room temperature and rinsed 3 times with 200 μL

DPBS, with 2 min waiting time between each rinse. After that, cells were blocked with 100 μL

3% BSA in DPBS for 1h at room temperature. Anti-Mucin 5AC mouse monoclonal antibody

(Abcam, Cat.# ab218466) was prepared in DPBS (1/100). The actin stain (ActinGreen™ 488

ReadyProbes™ Reagent, Thermo Fisher Scientific, Cat.# R37110) and the nuclei counterstain

(NucBlue™ Live ReadyProbes™ Reagent, Thermo Fisher Scientific, Cat. # R37605) were

added to the same buffer (2 drops/mL and 1 drop/mL, respectively). Anti-Aspergillus rabbit

polyclonal antibody (Abcam, Cat.# ab20419, RRID:AB_445571) was also added (1/200) in the

staining buffer for the conditions where *A. niger* conidia were inoculated. Fixed cells were

incubated with 100 µL buffer containing the stains and Anti-Mucin 5AC and Anti-Aspergillus

antibodies overnight at 4°C. The next day, fixed cells were washed 3 times with DPBS and

secondary antibodies were applied. Goat anti-Mouse IgG antibody (1/250) conjugated with

Alexa Fluor 546 (Thermo Fisher Scientific, Cat.# A-11003, RRID:AB_2534071) directed

against Anti-Mucin 5AC antibody and Goat anti-Rabbit IgG antibody (1/500) conjugated with

Alexa Fluor 594 (Thermo Fisher Scientific, Cat.# A-11012, RRID:AB_2534079) directed

against Anti-Aspergillus antibody were prepared in DPBS. Fixed cells were incubated with 100

µL buffer containing the secondary antibodies overnight at 4°C. The following day, fixed cells

were once again washed 3 times with DPBS and the membranes from the inserts were carefully

cut out with a sharp knife. The membranes were mounted on a glass slide using Fluoromount-

G™ Mounting Medium (Thermo Fisher Scientific, Cat.# 00-4958-02) and imaged with a Zeiss

Axio Observer Z1 fluorescence inverted microscope (Carl Zeiss AG, Oberkochen, Germany).

pH measurements and quantification of calcium, oxalic acid and lactate dehydrogenase

(LDH)

pH measurements of the culture medium after 72h incubation were done directly after taking

the samples out of the incubator using pH-indicator strips (Merck, Germany, Cat.# 109535) for

the Transwell® inserts samples, and CG8+ i-STAT cartridges (Abbott, USA, Cat.# 03P88-25)

for the BoC devices, as the pH change of the culture medium indicator was less visible. Free

calcium (Calcium Colorimetric Assay, Sigma-Aldrich, Darmstadt, Germany, Cat.# MAK022),

free oxalic acid (Oxalic Acid Colorimetric Assay Kit, Sigma-Aldrich, Darmstadt, Germany,

Cat.# MAK179), and LDH (ScienCell Research Laboratories, CA, USA, Cat. # 8078) were

quantified in the culture medium using colorimetric assay kits following the manufacturer

instructions.

Statistical analyses

Statistical significance of the data from the confrontations on differentiated bronchial tissues in

Transwell® inserts (3 replicates, n = 3) was tested with unpaired two-tailed Student t-tests in

Microsoft® Excel (Version 16.37). The statistical significance threshold was set to 5%.

Acknowledgments

We would like to thank Diego Gonzales and Ted Turlings for critical review of the paper, and Pulak Nath from the Materials Physics and Applications Division of the Los Alamos National Laboratory for providing the equipment and laboratory infrastructure for the BoC devices fabrication. BoC devices were developed under Defense Threat Reduction Agency (DTRA) interagency agreement CBMXCEL-XL1-2-0001. **Author contributions:** FP, PSC, JFH and PJ conceived and designed the study. FP, IP and NN performed the *in-vitro* experiments. AV performed the UHPLC analysis. FP performed the experiments related to cell work. JMK performed the microscopic observation of the CaOx crystals in the differentiated bronchial epithelial tissue. MOI fabricated and assembled the BoC devices. FP, JFH and PJ analyzed the data. FP and PJ wrote the paper, and all the authors reviewed and edited the paper. **Data and materials availability:** All data is available in the main text or the supplementary materials. **Competing interests:** Authors declare no competing interests.

References

Arefin A, McCulloch Q, Martinez R, Martin SA, Singh R, Ishak OM, Higgins EM, Haffey KE, Huang J-H, Iyer S, Nath P, Iyer R, Harris JF. 2018. Micromachining of Polyurethane Membranes for Tissue Engineering Applications. *ACS Biomaterials Science & Engineering* **4**:3522–3533. doi:10.1021/acsbiomaterials.8b00578

Bongomin F, Gago S, Oladele RO, Denning DW. 2017. Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *J Fungi (Basel)* **3**. doi:10.3390/jof3040057

Braissant O, Cailleau G, Aragno M, Verrecchia EP. 2004. Biologically induced mineralization in the tree Milicia excelsa (Moraceae): its causes and consequences to the environment. *Geobiology* **2**:59–66. doi:10.1111/j.1472-4677.2004.00019.x

Brown GD, Denning DW, Gow NAR, Levitz SM, Netea MG, White TC. 2012. Hidden Killers: Human Fungal Infections. *Science Translational Medicine* **4**:165rv13. doi:10.1126/scitranslmed.3004404

Brunke S, Mogavero S, Kasper L, Hube B. 2016. Virulence factors in fungal pathogens of man. *Curr Opin Microbiol* **32**:89–95. doi:10.1016/j.mib.2016.05.010

Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu H, Kinnebrew M, Viale A, Littmann E, van den Brink MRM, Jenq RR, Taur Y, Sander C, Cross JR, Toussaint NC, Xavier JB, Pamer EG. 2015. Precision microbiome reconstitution restores bile acid mediated resistance to Clostridium difficile. *Nature* **517**:205–208. doi:10.1038/nature13828

Cameselle C, Bohlmann JT, Núñez MJ, Lema JM. 1998. Oxalic acid production by Aspergillus niger. *Bioprocess Engineering* **19**:247–252. doi:10.1007/pl00009017

Certini G, Corti G, Ugolini FC. 2000. Vertical trends of oxalate concentration in two soils

under Abies alba from Tuscany (Italy). *Journal of Plant Nutrition and Soil Science* **163**:173–

177. doi:10.1002/(SICI)1522-2624(200004)163:2<173::AID-JPLN173>3.0.CO;2-H

Charlson ES, Bittinger K, Haas AR, Fitzgerald AS, Frank I, Yadav A, Bushman FD, Collman

RG. 2011. Topographical Continuity of Bacterial Populations in the Healthy Human

Respiratory Tract. *American Journal of Respiratory and Critical Care Medicine* **184**:957–

963. doi:10.1164/rccm.201104-0655OC

Charlson ES, Diamond JM, Bittinger K, Fitzgerald AS, Yadav A, Haas AR, Bushman FD,

Collman RG. 2012. Lung-enriched Organisms and Aberrant Bacterial and Fungal Respiratory

Microbiota after Lung Transplant. *American Journal of Respiratory and Critical Care*

Medicine **186**:536–545. doi:10.1164/rccm.201204-0693OC

Cui L, Lucht L, Tipton L, Rogers MB, Fitch A, Kessinger C, Camp D, Kingsley L, Leo N,

Greenblatt RM, Fong S, Stone S, Dermand JC, Kleerup EC, Huang L, Morris A, Ghedin E.

2015. Topographic Diversity of the Respiratory Tract Mycobiome and Alteration in HIV and

Lung Disease. *American Journal of Respiratory and Critical Care Medicine* **191**:932–942.

doi:10.1164/rccm.201409-1583OC

de Oliveira Ceita G, Macêdo JNA, Santos TB, Alemanno L, da Silva Gesteira A, Micheli F,

Mariano AC, Gramacho KP, da Costa Silva D, Meinhardt L, Mazzafera P, Pereira GAG, de

Mattos Cascardo JC. 2007. Involvement of calcium oxalate degradation during programmed

cell death in Theobroma cacao tissues triggered by the hemibiotrophic fungus Moniliophthora

perniciosa. *Plant Science* **173**:106–117. doi:10.1016/j.plantsci.2007.04.006

Diamantstein T, Odenwald MV. 1974. Control of the immune response in vitro by calcium

ions. I. The antagonistic action of calcium ions on cell proliferation and on cell

differentiation. *Immunology* **27**:531–541.

Diamantstein T, Ulmer A. 1975. The control of immune response in vitro by Ca2+. II. The

Ca2+-dependent period during mitogenic stimulation. *Immunology* **28**:121–125.

Dickson RP, Martinez FJ, Huffnagle GB. 2014. The role of the microbiome in exacerbations

of chronic lung diseases. *The Lancet* **384**:691–702. doi:10.1016/S0140-6736(14)61136-3

Dominguez-Gutierrez PR, Kusmartsev S, Canales BK, Khan SR. 2018. Calcium Oxalate

Differentiates Human Monocytes Into Inflammatory M1 Macrophages. *Front Immunol*

9:1863. doi:10.3389/fimmu.2018.01863

Dutton MV, Evans CS. 1996. Oxalate production by fungi: its role in pathogenicity and

ecology in the soil environment. *Canadian Journal of Microbiology* **42**:881–895.

doi:10.1139/m96-114

Enaud R, Prevel R, Ciarlo E, Beaufils F, Wieërs G, Guery B, Delhaes L. 2020. The Gut-Lung

Axis in Health and Respiratory Diseases: A Place for Inter-Organ and Inter-Kingdom

Crosstalks. *Frontiers in Cellular and Infection Microbiology* **10**.

doi:10.3389/fcimb.2020.00009

Ferro KI. 2012. The impact of oxalogenic plants on soil carbon dynamics - formation of a

millennium carbon storage as calcium carbonate. *Faculty of Sciences*. Neuchâtel: University

of Neuchâtel.

Fisher MC, Gurr SJ, Cuomo CA, Blehert DS, Jin H, Stukenbrock EH, Stajich JE, Kahmann

R, Boone C, Denning DW, Gow NAR, Klein BS, Kronstad JW, Sheppard DC, Taylor JW,

Wright GD, Heitman J, Casadevall A, Cowen LE. 2020. Threats Posed by the Fungal

Kingdom to Humans, Wildlife, and Agriculture. *mBio* **11**. doi:10.1128/mBio.00449-20

Fisher MC, Hawkins NJ, Sanglard D, Gurr SJ. 2018. Worldwide emergence of resistance to

antifungal drugs challenges human health and food security. *Science* **360**:739–742.

doi:10.1126/science.aap7999

580 Foesel BU, Pfeiffer S, Raj ACD, Etschmann SK, Schloter M. 2019. Applying ecological

581 theories in lung microbiome research: lessons learned from microbial ecology and evolution?

582 In: Cox MJ, Ege MJ, von Mutius E, editors. The Lung Microbiome. pp. 50–66.

583 doi:10.1183/2312508x.10015718

584 Fulcher ML, Gabriel S, Burns KA, Yankaskas JR, Randell SH. 2005. Well-Differentiated

585 Human Airway Epithelial Cell Cultures In: Picot J, editor. Methods in Molecular Medicine.

586 Graustein WC, Cromack K Jr, Sollins P. 1977. Calcium oxalate: occurrence in soils and effect

587 on nutrient and geochemical cycles. *Science (Washington, DC)* 1252–1254.

588 Grinstein S, Klip A. 1989. Calcium homeostasis and the activation of calcium channels in

589 cells of the immune system. *Bull N Y Acad Med* **65**:69–79.

590 Hofmann BA, Bernasconi SM. 1998. Review of occurrences and carbon isotope geochemistry

591 of oxalate minerals: implications for the origin and fate of oxalate in diagenetic and

592 hydrothermal fluids. *Chemical Geology* **149**:127–146. doi:10.1016/S0009-2541(98)00043-6

593 Hsieh H-L, Nath P, Huang J-H. 2019. Multistep Fluidic Control Network toward the

594 Automated Generation of Organ-on-a-Chip. *ACS Biomaterials Science & Engineering*

595 **5**:4852–4860. doi:10.1021/acsbiomaterials.9b00912

596 Kousha M, Tadi R, Soubani AO. 2011. Pulmonary aspergillosis: a clinical review. *European*

597 *Respiratory Review* **20**:156–174. doi:10.1183/09059180.00001011

598 Kurrein F, Green GH, Rowles SL. 1975. Localized Deposition of Calcium Oxalate around a

599 Pulmonary Aspergillus niger Fungus Ball. *American Journal of Clinical Pathology* **64**:556–

600 563. doi:10.1093/ajcp/64.4.556

601 Lawley TD, Clare S, Walker AW, Stares MD, Connor TR, Raisen C, Goulding D, Rad R,

602 Schreiber F, Brandt C, Deakin LJ, Pickard DJ, Duncan SH, Flint HJ, Clark TG, Parkhill J,

Dougan G. 2012. Targeted Restoration of the Intestinal Microbiota with a Simple, Defined

Bacteriotherapy Resolves Relapsing Clostridium difficile Disease in Mice. *PLoS Pathog*

8:e1002995. doi:10.1371/journal.ppat.1002995

Lehner A, Meimoun P, Errakhi R, Madiona K, Barakate M, Bouteau F. 2008. Toxic and

signalling effects of oxalic acid. *Canadian Journal of Microbiology* **3**:746–748.

doi:10.1093/jxb/ern166.746

Lin CK, Hsiao YY, Nath P, Huang JH. 2019. Aerosol delivery into small anatomical airway

model through spontaneous engineered breathing. *Biomicrofluidics* **13**:044109.

doi:10.1063/1.5121188

Maeno T, Sasaki M, Shibue Y, Mimura K, Oka H. 2015. Calcium oxalate in the sputum may

aid in the diagnosis of pulmonary aspergillosis: A report of two cases. *Med Mycol Case Rep*

8:32–6. doi:10.1016/j.mmcr.2015.01.003

Martin G, Guggiari M, Bravo D, Zopfi J, Cailleau G, Aragno M, Job D, Verrecchia E, Junier

P. 2012. Fungi, bacteria and soil pH: The oxalate-carbonate pathway as a model for metabolic

interaction. *Environmental Microbiology* **14**:2960–2970. doi:10.1111/j.1462-

2920.2012.02862.x

McClintock SD, Attili D, Dame MK, Richter A, Silvestri SS, Berner MM, Bohm MS,

Karpoff K, McCarthy CL, Spence JR, Varani J, Aslam MN. 2020. Differentiation of human

colon tissue in culture: Effects of calcium on trans-epithelial electrical resistance and tissue

cohesive properties. *PLoS ONE* **15**:e0222058. doi:10.1371/journal.pone.0222058

Mela F, Fritsche K, de Boer W, van Veen JA, de Graaff LH, van den Berg M, Leveau JH.

2011. Dual transcriptional profiling of a bacterial/fungal confrontation: Collimonas

fungivorans versus Aspergillus niger. *ISME J* **5**:1494–504. doi:10.1038/ismej.2011.29

Miller A, Dearing D. 2013. The Metabolic and Ecological Interactions of Oxalate-Degrading

Bacteria in the Mammalian Gut. *Pathogens* **2**:636–652. doi:10.3390/pathogens2040636

Mitchell AB, Oliver BGG, Glanville AR. 2016. Translational Aspects of the Human

Respiratory Virome. *American Journal of Respiratory and Critical Care Medicine* **194**:1458–

1464. doi:10.1164/rccm.201606-1278CI

Mulay SR, Kulkarni OP, Rupanagudi KV, Migliorini A, Darisipudi MN, Vilaysane A,

Muruve D, Shi Y, Munro F, Liapis H, Anders HJ. 2013. Calcium oxalate crystals induce renal

inflammation by NLRP3-mediated IL-1beta secretion. *J Clin Invest* **123**:236–46.

doi:10.1172/JCI63679

Muntz FH. 1999. Oxalate-producing pulmonary aspergillosis in an alpaca. *Vet Pathol*

36:631–2. doi:10.1354/vp.36-6-631

Oda M, Saraya T, Wakayama M, Shibuya K, Ogawa Y, Inui T, Yokoyama E, Inoue M,

Shimoyamada H, Fujiwara M, Ota T, Takizawa H, Goto H. 2013. Calcium oxalate crystal

deposition in a patient with Aspergilloma due to Aspergillus niger. *J Thorac Dis* **5**:E174-8.

doi:10.3978/j.issn.2072-1439.2013.08.45

Pabuççuoğlu U. 2005. Aspects of oxalosis associated with aspergillosis in pathology

specimens. *Pathology Research and Practice* **201**:363–368. doi:10.1016/j.prp.2005.03.005

Palmieri F, Estoppey A, House GL, Lohberger A, Bindschedler S, Chain PSG, Junier P. 2019.

Oxalic acid, a molecule at the crossroads of bacterial-fungal interactions. *Adv Appl Microbiol*

106:49–77. doi:10.1016/bs.aambs.2018.10.001

Paulussen C, Hallsworth JE, Alvarez-Perez S, Nierman WC, Hamill PG, Blain D, Rediers H,

Lievens B. 2017. Ecology of aspergillosis: insights into the pathogenic potency of Aspergillus

fumigatus and some other Aspergillus species. *Microb Biotechnol* **10**:296–322.

doi:10.1111/1751-7915.12367

Payne CL, Dark MJ, Conway JA, Farina LL. 2017. A retrospective study of the prevalence of

calcium oxalate crystals in veterinary Aspergillus cases. *J Vet Diagn Invest* **29**:51–58.

doi:10.1177/1040638716672254

Pfaller MA, Diekema DJ. 2004. Rare and Emerging Opportunistic Fungal Pathogens:

Concern for Resistance beyond Candida albicans and Aspergillus fumigatus. *Journal of*

Clinical Microbiology **42**:4419–4431. doi:10.1128/jcm.42.10.4419-4431.2004

Ramirez-Ortiz ZG, Means TK. 2012. The role of dendritic cells in the innate recognition of

pathogenic fungi (A. fumigatus, C. neoformans and C. albicans). *Virulence* **3**:635–46.

doi:10.4161/viru.22295

Reasoner DJ, Geldreich EE. 1985. A new medium for the enumeration and subculture of

bacteria from potable water. *Applied and Environmental Microbiology* **49**:1.

Ruijter GJG, van de Vondervoort PJI, Visser J. 1999. Oxalic acid production by Aspergillus

niger: an oxalate-non-producing mutant produces citric acid at pH 5 and in the presence of

manganese. *Microbiology* **145**:2569–2576. doi:10.1099/00221287-145-9-2569

Şahin N, Işik K, Tamer AÜ, Goodfellow M. 2000. Taxonomic Position of "Pseudomonas

oxaliticus" Strain Ox1T (DSM 1105T) (Khambata and Bhat, 1953) and its Description in the

Genus Ralstonia as Ralstonia oxalatica comb. nov. *Systematic and Applied Microbiology*

23:206–209. doi:10.1016/s0723-2020(00)80006-5

Sentchilo V, Czechowska K, Pradervand N, Minoia M, Miyazaki R, van der Meer JR. 2009.

Intracellular excision and reintegration dynamics of the ICEclc genomic island of

Pseudomonas knackmussii sp. strain B13. *Mol Microbiol* **72**:1293–306. doi:10.1111/j.1365-

2958.2009.06726.x

Suleyman G, Alangaden GJ. 2016. Nosocomial Fungal Infections: Epidemiology, Infection

Control, and Prevention. *Infect Dis Clin North Am* **30**:1023–1052.

doi:10.1016/j.idc.2016.07.008

Suryavanshi MV, Bhute SS, Jadhav SD, Bhatia MS, Gune RP, Shouche YS. 2016.

Hyperoxaluria leads to dysbiosis and drives selective enrichment of oxalate metabolizing

bacterial species in recurrent kidney stone endures. *Scientific Reports* **6**.

doi:10.1038/srep34712

Ticinesi A, Nouvenne A, Chiussi G, Castaldo G, Guerra A, Meschi T. 2020. Calcium Oxalate

Nephrolithiasis and Gut Microbiota: Not just a Gut-Kidney Axis. A Nutritional Perspective.

Nutrients **12**. doi:10.3390/nu12020548

Toprak E, Veres A, Michel JB, Chait R, Hartl DL, Kishony R. 2011. Evolutionary paths to

antibiotic resistance under dynamically sustained drug selection. *Nat Genet* **44**:101–5.

doi:10.1038/ng.1034

Turroni S, Vitali B, Bendazzoli C, Candela M, Gotti R, Federici F, Pirovano F, Brigidi P.

2007. Oxalate consumption by lactobacilli: Evaluation of oxalyl-CoA decarboxylase and

formyl-CoA transferase activity in Lactobacillus acidophilus. *Journal of Applied*

Microbiology **103**:1600–1609. doi:10.1111/j.1365-2672.2007.03388.x

van Woerden HC, Gregory C, Brown R, Marchesi JR, Hoogendoorn B, Matthews IP. 2013.

Differences in fungi present in induced sputum samples from asthma patients and non-atopic

controls: a community based case control study. *BMC Infectious Diseases* **13**:69.

doi:10.1186/1471-2334-13-69

Vig M, Kinet J-P. 2009. Calcium signaling in immune cells. *Nature Immunology* **10**:21–27.

doi:10.1038/ni.f.220

Yi Y, Cho SY, Lee DG, Jung JI, Park YJ, Lee KY. 2019. Invasive Pulmonary Aspergillosis

Due to Aspergillus awamori: Role of Calcium Oxalate Crystal Precipitation Mimicking

Mucormycosis. *Mycopathologia*. doi:10.1007/s11046-019-00405-z

Fig. 1. Schematic summary of the proposed strategy to control *Aspergillus niger* infection by introducing oxalotrophic bacteria to modify the *A. niger* environmental niche. (A) *A. niger* conidia (depicted in dark grey) arrive in the respiratory system through breathing. (B) During a normal infection process in a susceptible host, *A. niger* modifies the environment by secreting oxalic acid (or oxalate Ox^{2-}) which decreases pH and chelates free calcium in the form of calcium oxalate (CaOx; depicted in light grey) crystals. This results in the infection of the host's tissue. (C) The biocontrol strategy proposed here takes advantage of the ability of oxalotrophic bacteria (cells depicted in red) to consume CaOx and thus reestablish physiological pH and free calcium concentrations.

Fig. 2. Comparison of the growth of *Aspergillus niger* (An) alone and in confrontation with

***Cupriavidus oxalaticus* (Co) in different culture media.** The red line next to each picture

represents the extent of *A. niger* growth. On MA 1/10 (A), there is no significant growth

inhibition of *A. niger*, as it kills *C. oxalaticus*. *A. niger* growth is highly restricted to the center

of the plate when co-cultured with *C. oxalaticus* on R2A (B). The growth inhibition of *A. niger*

when co-cultured with *C. oxalaticus* is less pronounced on WYA + BP (C).

Fig. 3. Influence of the interaction between *Aspergillus niger* and the oxalotrophic bacterium *Cupriavidus oxalaticus* on environmental parameters of differentiated bronchial tissue in Transwell® inserts and bronchiole-on-a-chip (BoC) devices. In the presence of the fungus, the pH (A) decreases, as compared to all other treatments. This pH decrease is correlated with a drastic decrease in the concentration of free Ca^{2+} (B) (*p*-value between C and C+F = $8,951 \times 10^{-7}$). Free oxalic acid concentrations were lower in the presence of the fungus, compared with the basal level secreted by the bronchial cells (C) (*p*-value between C and C+F = $4,587 \times 10^{-8}$). These results are supported by the detection of calcium oxalate crystals in the presence of the fungus (D). In the co-culture with the oxalotrophic bacterium, pH, free Ca^{2+} and free oxalic acid concentrations return to physiological levels, and this was concomitant with the absence of crystals (E) (*p*-values between C+F and C+F+B for free Ca^{2+} and free oxalic acid = $0,002$ and $1,415 \times 10^{-7}$, respectively). C: lung cells; C+F: lung cells + fungus; C+B: lung cells + bacteria; C+F+B: lung cells + fungus + bacteria. For A, B, C, the results represent the mean + sd of three independent measurements for the Transwell® inserts for each condition (three biological replicates, n = 3). For A, pH results for the BoC devices represent a unique measurement per condition (one replicate, n = 1). For B and C, Ca^{2+}

734   and oxalic acid results for the BoC devices represents the mean + sd of two measurements per

735   condition (one replicate).

Fig. 4. Cytopathic effect of *Aspergillus niger* on differentiated bronchial tissue and lactate

dehydrogenase (LDH) measurements of the co-culture between *A. niger* and *Cupriavidus*

***oxalaticus* in Transwell® inserts and bronchiole-on-a-chip (BoC) devices.** (A) Control

showing healthy differentiated bronchial epithelial cells. (B) Bronchial epithelial cells infected

with *A. niger*. (C) LDH leakage was measured as a proxy for cell damage. In the presence of

the fungus, no LDH has been detected, probably because of the destruction of the tissue by *A.*

niger (B). *C. oxalaticus* cause significantly more LDH leakage that the basal LDH level of

control cells (p-value = 0,004). C: lung cells; C+F: lung cells + fungus; C+B: lung cells +

bacteria; C+F+B: lung cells + fungus + bacteria. For C, the results represent the mean + sd of

three independent measurements for the Transwell® inserts for each condition (three biological

replicates, n = 3). LDH results for the BoC devices represents the mean + sd of three

measurements per condition (one replicate, n = 1).