Title: Virucidal and antiviral activity of astodrimer sodium against SARS-CoV-2 *in vitro.*

Authors: Jeremy R.A. Paull[a],*; Graham P. Heery [a], Michael D. Bobardt[b], Alex Castellarnau[a];

Carolyn A. Luscombe[a]; Jacinth K. Fairley[a]; Philippe A. Gallay[b]

Author affiliations:

[a] Starpharma Pty Ltd, 4-6 Southampton Crescent, Abbotsford, Victoria 3067, Australia

[b] Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA

92307, USA

* Corresponding author: jeremy.paull@starpharma.com (JP)

Declaration of interest:

The authors declare the following financial interests that may be considered as potential

competing interests: J.R.A.P., J.K.F. and G.P.H. are paid employees of Starpharma Pty Ltd. A.C.

and C.A.L. are paid consultants to Starpharma Pty Ltd.

Word count:

Abstract: 284

Text: 3463

Inserts: 4 tables, 5 figures

Abstract

An effective response to the ongoing coronavirus disease (COVID-19) pandemic caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) will involve a range of complementary preventive modalities. The current studies were conducted to evaluate the *in vitro* SARS-CoV-2 antiviral and virucidal activity of astodrimer sodium, a dendrimer with broad spectrum antimicrobial activity, including against enveloped viruses in *in vitro* and *in vivo* models, that is marketed for antiviral and antibacterial applications. We report that astodrimer sodium inhibits replication of SARS-CoV-2 in Vero E6 and Calu-3 cells, with 50% effective concentrations (EC_{50}) for i) reducing virus-induced cytopathic effect of 0.002 to 0.012 mg/mL in Vero E6 cells, and ii) infectious virus release by plaque assay of 0.019 to 0.031 mg/mL in Vero E6 cells and 0.031 to 0.045 mg/mL in Calu-3 cells. The selectivity index (SI) in these assays was as high as 2197. Astodrimer sodium was also virucidal, reducing SARS-CoV-2 infectivity by >99.9% (>3 $\log_{10}$) within 1 minute of exposure, and up to >99.999% (>5 $\log_{10}$) shown at astodrimer sodium concentrations of 10 to 30 mg/mL in Vero E6 and Calu-3 cell lines. Astodrimer sodium also inhibited infection in a primary human airway epithelial cell line. The data were similar for all investigations and were consistent with the potent antiviral and virucidal activity of astodrimer sodium being due to inhibition of virus-host cell interactions, as previously demonstrated for other viruses. Further studies will confirm if astodrimer sodium binds to SARS-CoV-2 spike protein and physically blocks initial attachment of the virus to the host cell. Given the *in vitro* effectiveness and significantly high SI, astodrimer sodium warrants further investigation for potential as a nasally administered or inhaled antiviral agent for SARS-CoV-2 prevention and treatment applications.

Keywords: Astodrimer; COVID-19; dendrimer; antiviral; SARS-CoV-2; SPL7013

1. Introduction

The ongoing pandemic coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection, has resulted in unprecedented efforts to rapidly develop strategies to contain infection rates for the protection of vulnerable populations. An effective public health response to the current pandemic will involve currently available vaccines being complemented by supplementary preventive modalities.

SARS-CoV-2 receptors and coreceptors have been shown to be highly expressed in nasal epithelial cells (Sungnak et al., 2020). This finding is consistent with the virus infectivity or replication pattern along the respiratory tract, which peaks proximally (nasal cavity) and is relatively minimal in the distal alveolar regions (Hou et al., 2020). These findings suggest that nasal carriage of the virus is a key feature of transmission, and that nasally administered therapeutic modalities could be potentially effective in helping to prevent spread of infection.

Astodrimer sodium (SPL7013) is a generation-four lysine dendrimer with a polyanionic surface charge (McCarthy et al., 2005) that is active against several enveloped and non-enveloped viruses including human immunodeficiency virus-1 (HIV-1) (Lackman-Smith et al., 2008, Tyssen et al., 2010), herpes simplex virus (HSV)-1 and -2 (Gong et al., 2005), H1N1 and H3N2 influenza virus, human respiratory syncytial virus (HRSV), human papillomavirus (HPV), adenovirus and Zika virus (unpublished data). Astodrimer sodium also has antibacterial properties. Both size and surface charge contribute to the function of the compound (Tyssen et al., 2010), and when administered topically, astodrimer sodium is not absorbed systemically (Chen et al., 2009; O'Loughlin et al., 2010; McGowan et al., 2011).

Vaginally administered astodrimer sodium protected macaques from infection with chimeric

simian-HIV-1 (SHIV)$_{89.6P}$ (Jiang et al., 2005), and mice and guinea pigs from HSV-2 infection

(Bernstein et al., 2003) in vaginal infection challenge models. Astodrimer 1% Gel (10 mg/mL

astodrimer sodium) administered vaginally has been shown to be safe and effective in phase 2

and large phase 3 trials for treatment and prevention of bacterial vaginosis (BV) (Chavoustie et

al., 2020; Waldbaum et al., 2020; Schwebke et al., 2021) and is marketed in Europe, Australia,

New Zealand and several countries in Asia.

The current studies were conducted to assess the antiviral and virucidal activity of astodrimer

sodium against SARS-CoV-2 *in vitro*, to determine its potential as a reformulated, nasally

administered or inhaled antiviral agent to help prevent spread of SARS-CoV-2 infection.

2. Materials and methods

2.1 Virus, cell culture, astodrimer sodium and controls

SARS-CoV-2 hCoV-19/Australia/VIC01/2020 was a gift from Melbourne's Peter Doherty

Institute for Infection and Immunity (Melbourne, Australia). Virus stock was generated at

360Biolabs (Melbourne, Australia) by two passages in Vero cells in virus growth media, which

comprised Minimal Essential Medium (MEM) without L-glutamine supplemented with 1% (w/v)

L-glutamine, 1.0 µg/mL of L-(tosylamido-2-phenyl) ethyl chloromethyl ketone (TPCK)-treated

trypsin (Worthington Biochemical, NJ, USA), 0.2% bovine serum albumin (BSA) and 1%

insulin-transferrin-selenium (ITS).

SARS-CoV-2 2019-nCoV/USA-WA1/2020 strain was isolated from an oropharyngeal swab

from a patient with a respiratory illness who developed clinical disease (COVID-19) in January

2020 in Washington, US, and sourced from BEI Resources (NR-52281). Virus was derived from

African green monkey kidney Vero E6 cells or lung homogenates from human angiotensin

converting enzyme 2 (hACE2) transgenic mice.

Vero E6 and human Calu-3 cell lines were cultured in MEM without L-glutamine supplemented

with 10% (v/v) heat-inactivated fetal bovine serum (FBS) and 1% (w/v) L-glutamine. Vero E6

and Calu-3 cells were passaged for a maximum of 10 passages for antiviral and virucidal studies.

Hank's balanced salt solution (HBSS) with 2% FBS was used for infection. The 2019-

nCoV/USA-WA1/2020 strain antiviral assays were performed with a multiplicity of infection

(MOI) of 0.1.

The virus inoculums for virucidal assays were 10^4, 10^5, and 10^6 pfu/mL. After defined incubation

periods, the solution was pelleted through a 20% sucrose cushion (Beckman SW40 Ti rotor) and

resuspended in 1.5 mL MEM, which was then added to 2.5×10^4 cells/well.

Primary human bronchial epithelial cells (HBEpC) (Sigma-Aldrich, MO, USA) were grown and

maintained in HBEpC/HTEpC growth medium (Cell Applications, CA, USA). These primary

cells express the ACE2 receptor and are permissive to SARS-CoV-2 infection. These cells were

used to determine the antiviral effect of astodrimer sodium against SARS-CoV-2 in a primary

human airway epithelial cell line. Cells were infected with SARS-CoV-2 2019-nCoV/USA-

WA1/2020 at 10^3 pfu/mL with 1 mL added to 2.5×10^4 cells/well. The positive control was

addition of 10 µg/mL of SARS-CoV-2 spike protein antibody (pAb, T01KHuRb)

(ThermoFisher, MA, USA) at the time of infection.

Astodrimer sodium was prepared as 100 mg/mL in water and stored at 4°C. Astodrimer sodium

has a molecular weight of 16581.57 g/mol and the structure is described and illustrated in Tyssen

et al., 2010. The purity of the compound used in these studies was assessed by ultra-high-

performance liquid chromatography (UPLC) to be 98.79%.

Remdesivir (MedChemExpress, NJ, USA) was used as a positive control in the virus-induced

cytopathic effect (CPE) inhibition and plaque assays.

Iota-carrageenan (Sigma-Aldrich, MO, USA) was used in the primary epithelial cell

nucleocapsid and plaque assays to compare the antiviral activity of this substance with

astodrimer sodium. Concentrations used are those reported to show activity against SARS-CoV-

2 (Bansal et al., 2020).

2.2 Virus-induced cytopathic effect inhibition assay

Vero E6 (ATCC-CRL1586) cell stocks were generated in cell growth medium, which

comprised MEM without L-glutamine supplemented with 10% (v/v) heat-inactivated FBS and

1% (w/v) L-glutamine. Vero E6 cell monolayers were seeded in 96-well plates at

$2x10^4$ cells/well in 100 µL growth medium (MEM supplemented with 1% (w/v) L-glutamine,

2% FBS) and incubated overnight at 37°C in 5% CO_2. SARS-CoV-2 infection was established

by using an MOI of 0.05 to infect cell monolayers.

Astodrimer sodium or remdesivir were serially diluted 1:3, 9 times and each compound

concentration was assessed for both antiviral efficacy and cytotoxicity in triplicate.

Astodrimer sodium was added to Vero E6 cells 1 hour prior to infection or 1 hour post-

infection with SARS-CoV-2. Cell cultures were incubated at 37°C in 5% CO_2 for 4 days prior

to assessment of CPE. The virus growth media was MEM supplemented with 1% (w/v) L-

glutamine, 2% FBS, and 4 µg/mL TPCK-treated trypsin. On Day 4, viral-induced CPE and

cytotoxicity of the compound were determined by measuring the viable cells using the

methylthiazolyldiphenyl-tetrazolium bromide (MTT) assay (MP Biomedicals, NSW,

Australia). Absorbance was measured at 540-650 nm on a plate reader.

2.3 Antiviral plaque assay evaluation and nucleocapsid ELISA

For the antiviral evaluation, astodrimer sodium was added to cells 1 hour prior to, at the time of,

and 1 hour after exposing the cells to virus. For both the antiviral and virucidal assays, at 6 hours

after infection, cells were washed to remove astodrimer sodium and/or any virus remaining in the

supernatant, in such way that a

Following initial infection, cell cultures were incubated and supernatants recovered after 16

hours or 4 days. The amount of virus in the supernatants was determined by plaque assay (plaque

forming unit [pfu]) and by nucleocapsid enzyme-linked immunosorbent assay (ELISA). The

plaque assay used was as described in van den Worm et al., 2012, utilizing 2% sodium

carboxymethyl cellulose overlay, fixation of cells by 4% paraformaldehyde and staining with

0.1% crystal violet. The nucleocapsid ELISA assay was as described by Bioss Antibodies, USA

(BSKV0001).

The assessment of astodrimer sodium cytotoxicity occurred on Day 4 by measuring lactate

dehydrogenase (LDH) activity in the cytoplasm using an LDH detection kit (Cayman Chemical),

with 0.5% saponin used as the positive cytotoxic control.

2.4 Virucidal assay

For the virucidal evaluation, concentrations of astodrimer sodium (0.0046 to 30 mg/mL) were

incubated with SARS-CoV-2 2019-nCoV/USA-WA1/2020 for times ranging from 5 seconds to

2 hours. To neutralize the effect of astodrimer sodium, unbound compound was separated from

the astodrimer:virus mixture by pelleting the preincubated mixture through a 20% sucrose

cushion (Beckman SW40 Ti rotor). The astodrimer sodium-containing supernatant was removed

(i.e., neutralising the effect of SPL7013) and then the pelleted virus was gently resuspended and

added to Vero E6 or Calu-3 cell cultures. Virus infection, cell culture and cytotoxicity

assessment was as described for the plaque assay described in Section 2.3.

All antiviral and virucidal assays were performed in triplicate, except where indicated in the

results.

2.5 Determination of 50% effective concentration (EC_{50}) and cytotoxicity (CC_{50})

The concentration of compound that gives a 50% reduction in viral-induced CPE, infectious

virus (pfu/mL), or secreted viral nucleocapsid (EC_{50}) was calculated using the formula of

Pauwels et al., 1998.

The concentration of compound that resulted in a 50% reduction in cell viability (CC_{50}) after

4 days of culture was also calculated by the formula of Pauwels et al., 1998.

2.6 Primary Epithelial Cell Assay

Astodrimer sodium (0, 1.1, 3.3 and 10 mg/mL) or iota-carrageenan (0, 6, 60 and 600 µg/mL)

were added to HBEpC cells 1 hour prior to infection with SARS-CoV-2. Cells were cultured for

4 days post-infection and the cell supernatant was analysed for the amount of secreted SARS-

CoV-2 nucleocapsid by ELISA, and infectious virus was quantitated by plaque assay, as

described in Section 2.3.

3. Results

3.1 Virus-induced cytopathic effect inhibition

In two independent virus-induced CPE inhibition assays, astodrimer sodium inhibited SARS-

CoV-2 (hCoV-19/Australia/VIC01/2020) replication in Vero E6 cells in a dose dependent

manner (Table 1). Astodrimer sodium inhibited viral replication when added either 1 hour prior

to infection, or 1 hour post-infection with SARS-CoV-2.

Astodrimer sodium was initially tested in the range of 0.0013 to 8.63 mg/mL. In the repeat set of

assays, astodrimer sodium was tested in the range of 0.0001 to 0.86 mg/mL to help further

characterize the lower end of the dose response curve. The effective and cytotoxic concentrations,

and selectivity indices from the assays are shown individually and as means in Table 1 for CPE

determined readouts.

The selectivity index (SI) for astodrimer sodium against SARS-CoV-2 in the CPE studies ranged

from 793 to 2197 for the initial assays where compound was added 1 hour prior to infection and

1 hour after infection, respectively, and was >70 to >80 in the repeat assays, in which

cytotoxicity was not observed up to the highest concentration tested (0.86 mg/mL).

The positive control, remdesivir, was also active in the CPE inhibition assay, with a SI of >33.

3.2 Antiviral efficacy

To determine the ability of astodrimer sodium to inhibit globally diverse SARS-CoV-2 strains,

the compound was evaluated against the 2019-nCoV/USA-WA1/2020 virus in Vero E6 cells and

human Calu-3 cells. Antiviral readouts were based on virological endpoints of infectious virus or

viral nucleocapsid released into the supernatant post-infection. As shown in Table 2 and Figures

1 and 2, astodrimer inhibited the 2019-nCoV/USA-WA1/2020 strain with an EC_{50} 0.019 to 0.031

mg/mL and 0.031 to 0.045 mg/mL for infectious virus release as determined by plaque assay in

Vero E6 cell for Calu-3 cells, respectively. These data are consistent with the inhibition by

astodrimer of the replication of the Australian SARS-CoV-2 isolate *in vitro*. The dose response

data for the nucleocapsid released into the supernatant by ELISA were similar to the infectious

virus release data in each cell line (data not shown). The positive control, remdesivir, was also

active in the plaque assay.

3.3 Virucidal efficacy

Virucidal assays investigated if astodrimer sodium could reduce viral infectivity by irreversibly

inactivating SARS-CoV-2 prior to infection of Vero E6 cells and human airway Calu-3 cells.

Following incubation of virus with astodrimer for up to 2 hours and neutralization of astodrimer,

the astodrimer-exposed virus was added to cell cultures. After either 16 hours or 96 hours (Day

4), the cell culture supernatant was collected for assessment of progeny viral infectivity as

determined by the amount of secreted infectious virus and nucleocapsid. The SARS-CoV-2

replication lifecycle is completed in approximately 8 hours (Ogando et al., 2020) and in these

studies, we sampled at 16 hours (2 lifecycles) or Day 4 (12 lifecycles) post-infection.

Enabling a possible 12 rounds of infection, the Day 4 (96 hour) sampling time point identified

that exposure of 10^6 pfu/mL SARS-CoV-2 to astodrimer sodium for 1 to 2 hours resulted in a

dose-dependent reduction in viral infectivity, with 10 to 30 mg/mL astodrimer sodium achieving

up to >99.999% (>5 $\log_{10}$) reduced infectivity in Vero E6 cells and >99.9% (>3 $\log_{10}$) reduced

infectivity in Calu-3 cells (data not shown) compared to untreated virus. SARS-CoV-2

infectivity was also reduced by up to >99.999% in Vero E6 cells when the incubation time of

astodrimer (10 to 30 mg/mL) with 10^6 pfu/mL virus was reduced to 15 to 30 minutes (data not

shown).

Incubation of astodrimer sodium with viral inoculums of 10^4, 10^5 and 10^6 pfu/mL for as little as

5 seconds resulted in evidence of reduced infectivity, with 10 to 15 minutes exposure being

sufficient to achieve >99.9% reduction in virus infectivity, and greater reduction achieved with

lower viral inoculum (>99.999%, 10^4 pfu/mL viral inoculum, 10 to 30 mg/mL astodrimer

sodium, and 10 to 15 min incubation time) (Table 3, Figure 3).

When assessed 16 hours post-infection of cells with astodrimer-exposed virus, it was found that

≥10 mg/mL astodrimer sodium inactivated >99.9% SARS-CoV-2 (10^4 pfu/mL) within as little as

1 minute of exposure (Table 4, Figure 4).

3.4 Antiviral Efficacy in Primary Human Airway Epithelial Cells

To determine the ability of astodrimer sodium to prevent SARS-CoV-2 infection of primary

human epithelial cells, the compound was evaluated against the 2019-nCoV/USA-WA1/2020

strain in HBEpC cell culture. Astodrimer sodium was found to reduce infection of HBEpC

primary cells by SARS-CoV-2 by up to 98% vs virus control by nucleocapsid ELISA (Figure

5A), and by up to 95% in the plaque assay (data not shown). In contrast, treatment with iota-

carrageenan had minimal antiviral effect against SARS-CoV-2 in this cell line, with the highest

concentration tested reducing infection by just 17% by nucleocapsid ELISA (Figure 5B), and just

21% in the plaque assay (data not shown). The maximum level of inhibition with astodrimer

sodium was comparable to inhibition achieved with the SARS-CoV-2 spike protein antibody

(pAb, T01KHuRb) positive control (see Figure 5A and B).

4. Discussion

Astodrimer sodium demonstrated potent antiviral activity against globally diverse SARS-CoV-2

strains *in vitro*. Antiviral activity was demonstrated by reduction in CPE, release of infectious

virus and release of viral nucleocapsid protein. Antiviral activity was demonstrated when

astodrimer sodium was added to cells prior to infection of cells and when the compound was

added to cells already exposed to SARS-CoV-2. Irreversible virucidal activity was demonstrated

when astodrimer sodium was mixed with virus for as little as 1 minute.

Of note is a significantly high SI for astodrimer sodium in the antiviral assays relative to other

antiviral compounds under investigation for SARS-CoV-2 activity (Pizzorno et al., 2020).

Remdesivir was used as the antiviral positive control for the CPE inhibition and antiviral assays

and the experimental EC_{50} was consistent with published data generated with a different clinical

isolate of SARS-CoV-2 (Wang et al., 2020).

Astodrimer sodium inhibited infection of a human airway primary epithelial cell by SARS-CoV-

2, whereas iota-carrageenan, which is a polyanionic compound in marketed nasal spray

formulations, failed to provide significant inhibition at concentrations that have previously been

shown to reduce SARS-CoV-2 infection in Vero E6 cells (Bansal et al., 2020). The unique

structure of astodrimer sodium, a sulphonated, roughly spherical molecule with a core and

densely packed branches radiating out from the core, appears to provide potential benefits over

other polyanionic compounds such as iota-carrageenan and heparin, which are linear sulphated

molecules with a distribution of molecular weight. The authors are not aware of data showing

that iota-carrageenan is virucidal, while heparin has demonstrated a lack of irreversible, virucidal

interaction with HSV virion components (Ghosh et al., 2009).

The antiviral data are consistent with astodrimer sodium being a potent inhibitor of early events

in the virus lifecycle. The virucidal assay data suggest that astodrimer sodium antiviral activity

was consistent with the proposed mechanism of action of binding to virus, thereby irreversibly

inactivating virus and blocking infection.

The virucidal activity of astodrimer sodium demonstrated that it irreversibly inhibits the early

phase of virus infection and replication. These findings suggest potent inhibition of viral

attachment, fusion and entry of the virus, which prevents virus replication and release of

infectious virus progeny.

Astodrimer sodium has been previously found to be an effective antiviral that exerts its inhibition

in the early virus-host receptor recognition interactions (Tyssen et al., 2010; Telwatte et al.,

2011), and its potential mechanism of action against SARS-CoV-2 is likely similar to that

identified for other pathogens. Astodrimer sodium was found to bind to HIV-1 by strong

electrostatic forces to positively charged clusters of highly conserved amino acids on HIV-1

gp120 protein and/or positively charged amino acid regions located between the stems of V1/V2

and V3 loops, which are exposed by conformational changes to gp120 after viral binding to the

receptor/co-receptor complex (Tyssen et al., 2010; Connell and Lortat-Jacob, 2013).

Many viruses utilize negatively charged heparan sulfate proteoglycans (HS) on the cell plasma

membrane as an initial means to scan the surface of the cell, and to attach in order to chaperone

the virus onto the receptor complex prior to viral entry (Sarrazin et al., 2011; Connell and Lortat-

Jacob, 2013). The receptor interactions occur in a sequential manner with virus-HS interactions

preceding receptor/co-receptor binding, which combined leads to fusion of the viral envelope

and the cell membrane.

Data indicate that astodrimer sodium-gp120 interaction may physically block initial HIV-1

association with HS and thereby block the subsequent virus-receptor complex functions.

Virucidal studies of astodrimer sodium determined that it did not disrupt the HIV-1 particle or

cause the loss of gp120 spike protein from the viral surface (Telwatte et al., 2011).

A report by Liu et al., 2020, described that densely glycosylated trimeric SARS-CoV-2 spike (S)

protein subunit S1, which is important for receptor binding, binds to HS. To engage with the

ACE2 receptor, the S protein undergoes a hinge-like conformational change that transiently

hides or exposes the determinants of receptor binding (Wrapp et al., 2020). Recent studies have

identified the binding of heparin to the receptor binding domain (RBD) of S1 resulting in a

conformational change to the S protein (Mycroft-West et al., 2020a, b and c). Mutations in the S

protein that are distal from the RBD also impact on viral transmission (Walls et al., 2020; Korber

et al., 2020; Yuan et al., 2020). Non-RBD polybasic cleavage sites, including S1/S2 loop

(Hoffmann et al., 2020a), have been described on SARS-CoV-2 S protein (Qiao and Olvera de la

Cruz, 2020) and may also be a site of potential interaction with astodrimer sodium.

SARS-CoV-2 utilizes the ACE2 receptor for viral infection of host cells (Hoffmann et al.,

2020b). Human CoV-NL63 also utilizes HS and ACE2 as its cellular receptor complex

(Milewska et al., 2014). The importance of HS for viral infectivity was also demonstrated for

close genetically related pseudo-typed SARS-CoV (Lang et al., 2011). The potential dependence

of SARS-CoV-2 on HS for attachment and entry combined with antiviral data from other viruses

suggest that negatively charged astodrimer sodium may have antiviral activity against SARS-

CoV-2 *in vitro* by blocking the early virus-receptor recognition events.

Astodrimer sodium is a polyanionic dendrimer reformulated for use as a topical, nasally

administered antiviral agent to inactivate SARS-CoV-2 before infection can occur. The potential

advantages of astodrimer sodium over other technologies include its lack of systemic absorption

following topical application (Chen et al., 2009; O'Loughlin et al., 2010; McGowan et al., 2011).

In addition, the SI of astodrimer sodium for SARS-CoV-2 is high and in a vaginal gel

formulation (10 mg/mL), the compound has been shown to be safe and effective in phase 2 and

large phase 3 trials for treatment and prevention of BV (Chavoustie et al., 2020; Waldbaum et al.,

2020; Schwebke et al., 2021) and is now marketed in Europe, Australia, New Zealand and

several countries in Asia. Astodrimer sodium is also the active antiviral substance in VivaGel®

condom products that have marketing authorization in Europe, Japan, Australia/New Zealand

and Canada. However, these current formulations are not appropriate for use to protect the

respiratory tract from SARS-CoV-2 infection.

5. Conclusions

Data from the current studies, taken together with studies of astodrimer sodium antiviral activity

against HIV-1, and HSV-1 and -2, indicate that the compound exerts its antiviral activity against

geographically diverse SARS-CoV-2 isolates by interfering with the early virus-cell recognition

events. Astodrimer sodium is a potent virucidal agent that reduces the infectivity of SARS-CoV-

2 by >99.9% after 1 minute of exposure to the virus. These studies support astodrimer sodium

being able to prevent early virus entry steps such as attachment, thereby reducing or preventing

viral infection or cell-cell spread.

An antiviral agent such as astodrimer sodium that blocks binding of the virus to target cells could

potentially be used as a preventive and/or a therapeutic agent against SARS-CoV-2. These

antiviral studies suggest that reformulation of astodrimer sodium for delivery to the respiratory

tract may be an effective preventive strategy to block SARS-CoV-2 transmission and augment

other protective and therapeutic strategies.

The potent antiviral and virucidal activity of astodrimer sodium against SARS-CoV-2 warrants

further investigation.

Tables

Table 1: Antiviral efficacy, measured by a reduction in CPE in virus-infected cells at Day 4 post-infection, and selectivity of astodrimer sodium against SARS-CoV-2 (hCoV-19/Australia/VIC01/2020) infection of Vero E6 cells

Compound / Assay Type	Repeat	EC_{50} (mg/mL)	CC_{50} (mg/mL)	SI
Astodrimer sodium added 1-hour pre-infection	Assay 1	0.004	2.93	793
	Assay 2	0.012	>0.863	>70
	Mean ± SD	0.008 ± 0.006	NC	>431[a]
Astodrimer sodium added 1-hour post-infection	Assay 1	0.002	3.29	2197
	Assay 2	0.011	>0.863	>80
	Mean ± SD	0.006 ± 0.007	NC	>1138[a]
Remdesivir added 1-hour post-infection	Assay 1	0.59 µM	>20 µM	>33
	Assay 2	0.61 µM	>20 µM	>33

EC_{50}=50% effective concentration; CC_{50}=50% cytotoxic concentration; SI=selectivity index (CC_{50}/EC_{50});
SD=standard deviation; NC=not calculated
[a] Average of Assay 1 and Assay 2 SI

Table 2: Antiviral efficacy, measured by a reduction in mean infectious virus (Log_{10} pfu/mL), and selectivity of astodrimer sodium against SARS-CoV-2 (2019-nCoV/USA-WA1/2020) on Day 4 post-infection

Compound / Assay Type	Cell Line	EC_{50} (mg/mL)	CC_{50} (mg/mL)	SI
Astodrimer sodium added 1-hour pre-infection	Vero E6	0.032	15.09	472
	Calu-3	0.037	21.76	588
Astodrimer sodium added at time of infection	Vero E6	0.020	15.09	755
	Calu-3	0.035	21.76	622
Astodrimer sodium added 1-hour post-infection	Vero E6	0.019	15.09	794
	Calu-3	0.030	21.76	725
Remdesivir added 1-hour post-infection	Vero E6	0.791 µM	N/A	N/A
	Calu-3	0.589 µM	N/A	N/A

EC_{50}=50% effective concentration; CC_{50}=50% cytotoxic concentration; SI=selectivity index (CC_{50}/EC_{50}); N/A=not applicable

349 **Table 3: Virucidal efficacy of 10 mg/mL astodrimer sodium against SARS-CoV-2 (2019-nCoV/USA-WA1/2020), measured by**

350 **a reduction in mean infectious virus (Log$_{10}$ pfu/mL), at 96 hours post-infection**

Viral Load (pfu/mL)	Virus:Astodrimer Incubation Time	Reduction vs. Virus Control (Log$_{10}$ ± SD)	Reduction vs. Virus Control (%)
10^6	5 sec	0.10 ± 0.20	20.567
	10 sec	0.03 ± 0.06	7.388
	30 sec	0.10 ± 0.10	20.567
	1 min	0.33 ± 0.12	53.584
	10 min	2.20 ± 0.10	99.369
	15 min	3.67 ± 0.23	99.979
10^5	5 sec	0.33 ± 0.21	53.584
	10 sec	0.23 ± 0.06	41.566
	30 sec	0.30 ± 0.17	49.881
	1 min	0.47 ± 0.21	65.855
	10 min	3.70 ± 0.26	99.980
	15 min	4.60 ± 0.10	99.998
10^4	5 sec	-0.13 ± 0.21	-35.936
	10 sec	0.07 ± 0.29	14.230
	30 sec	0.10 ± 0.10	20.567
	1 min	0.10 ± 0.00	20.567
	10 min	5.07 ± 0.25	>99.999
	15 min	5.83 ± 0.12	>99.999

351 Shading indicates data points where virucidal efficacy is >99.9% (3 log$_{10}$ reduction) vs. virus control; virus control=untreated virus, 0 mg/mL astodrimer sodium;
352 SD=standard deviation

353 Table 4: Virucidal efficacy of 10 mg/mL astodrimer sodium against SARS-CoV-2 (2019-nCoV/USA-WA1/2020), measured by

354 a reduction in mean infectious virus (Log$_{10}$ pfu/mL), at 16 hours post-infection

Viral Load (pfu/mL)	Virus:Astodrimer Incubation Time	Reduction vs. Virus Control (Log$_{10}$ ± SD)	Reduction vs. Virus Control (%)
10^5	30 sec	0.00 ± 0.36	0.000
	1 min	2.63 ± 0.15	99.767
	5 min	4.63 ± 0.31	99.998
	15 min	4.60 ± 0.10	99.998
10^4	30 sec	0.20 ± 0.20	36.904
	1 min	3.17 ± 0.12	99.932
	5 min	3.67 ± 0.21	99.979
	15 min	4.00 ± 0.10	99.990

355 Shading indicates data points where virucidal efficacy is >99.9% (3 log$_{10}$ reduction) vs. virus control; virus control=untreated virus, 0 mg/mL astodrimer sodium
356 SD=standard deviation
357
358

Figure Captions

Figure 1. Dose-response and cytotoxicity analysis of SARS-CoV-2 (2019-nCoV/USA-WA1/2020) antiviral activity of astodrimer sodium in Vero E6 cells as measured by infectious virus release (Log$_{10}$ pfu/mL) on Day 4 post-infection

Astodrimer sodium (0.0005 to 10 mg/mL) was added to cell cultures 1 hour prior to (t = -1h), at the time of (t = 0), and 1 hour post-infection (t = +1h). Cytotoxicity was assessed by LDH detection (OD @ 490 nm), with 0.5% saponin used as the positive cytotoxic control. Points and error bars represent mean ± SD of triplicate readings.

Figure 2. Dose-response and cytotoxicity analysis of SARS-CoV-2 (2019-nCoV/USA-WA1/2020) antiviral activity of astodrimer sodium in Calu-3 cells as measured by infectious virus release (Log$_{10}$ pfu/mL) on Day 4 post-infection

Astodrimer sodium (0.0005 to 10 mg/mL) was added to cell cultures 1 hour prior to (t = -1h), at the time of (t = 0), and 1 hour post-infection (t = +1h). Cytotoxicity was assessed by LDH detection (OD @ 490 nm), with 0.5% saponin used as the positive cytotoxic control. Points and error bars represent mean ± SD of triplicate readings.

Figure 3. Virucidal efficacy of astodrimer sodium against SARS-CoV-2 (2019-nCoV/USA-WA1/2020) measured by a reduction in mean infectious virus (Log$_{10}$ pfu/mL), at 96 hours post-infection in Vero E6 cells

Astodrimer sodium (1 to 30 mg/mL) was incubated with SARS-CoV-2 (2019-nCoV/USA-WA1/2020) for 5 sec up to 15 min. Treated virus was added to Vero E6 cells and the amount of infectious virus in the supernatant was determined by plaque assay 96 hours post-infection.

Graph shows dose-response of astodrimer sodium virucidal activity using 10^4 pfu/mL virus

inoculum. Points and error bars represent mean ± SD of triplicate readings. Dotted line indicates

level of mean infectious virus when untreated virus was added to Vero E6 cells (virus control).

Figure 4. Virucidal efficacy of astodrimer sodium against SARS-CoV-2 (2019-nCoV/USA-

WA1/2020) measured by a reduction in mean infectious virus (Log$_{10}$ pfu/mL), at 16 hours

post-infection in Vero E6 cells

Astodrimer sodium (0.0046 to 30 mg/mL) was incubated with SARS-CoV-2 (2019-nCoV/USA-

WA1/2020) for 30 sec, 1 min, 5 min and 15 min. Treated virus was added to Vero E6 cells and

the amount of infectious virus in the supernatant was determined by plaque assay 16 hours post-

infection. Graph shows dose-response of astodrimer sodium virucidal activity using 10^4 pfu/mL

virus inoculum. Points and error bars represent mean ± SD of triplicate readings. Dotted line

indicates level of mean infectious virus when untreated virus was added to Vero E6 cells (virus

control).

Figure 5. Antiviral efficacy of astodrimer sodium and iota-carrageenan against SARS-

CoV-2 (2019-nCoV/USA-WA1/2020) measured by a reduction in nucleocapsid (ng/mL), at

Day 4 post-infection in human bronchial epithelial primary cells (HBEpC)

Astodrimer sodium (0, 1.1, 3.3 and 10 mg/mL) or iota-carrageenan (0, 6, 60 and 600 µg/mL)

were added to cell cultures 1 hour prior to infection.

A. Dose-response of astodrimer sodium antiviral activity. Points and error bars represent mean ±

SD of triplicate readings.

B. Dose-response of carrageenan antiviral activity. Points represent one replicate.

Dotted lines indicates level of inhibition achieved with positive control, SARS-CoV-2 pAb.

Fig. 1

Fig. 2

409
410

411   **Fig 3.**

413   **Fig 4.**

416   **Fig 5.**

417   **A**

419   **B**

ACE2	angiotensin converting enzyme 2	LLOD	lower limit of detection
BSA	bovine serum albumin	MEM	minimal essential medium
BV	bacterial vaginosis	MOI	multiplicity of infection
CC_{50}	50% cytotoxic concentration	MTT	methylthiazolyldiphenyl-tetrazolium bromide
COVID-19	coronavirus disease 2019	NC	not calculated
CPE	cytopathic effect	OD	optical density
EC_{50}	50% effective concentration	pfu	plaque forming unit
ELISA	enzyme-linked immunosorbent assay	RBD	receptor binding domain
FBS	fetal bovine serum	S protein	spike protein
HBEpC	human bronchial epithelial primary cells	SARS-CoV	severe acute respiratory syndrome coronavirus
HIV-1	human immunodeficiency virus type 1	SARS-CoV-2	severe acute respiratory syndrome coronavirus 2
HPV	human papillomavirus	SD	standard deviation
HRSV	human respiratory syncytial virus	SHIV	simian-human immunodeficiency virus
HS	heparan sulfate proteoglycan	SI	selectivity index
HSV	herpes simplex virus	$TCID_{50}$	median tissue culture infective dose
HTEpC	human tracheal epithelial primary cells	TOA	time of addition
ITS	insulin-transferrin-selenium	TPCK	L-(tosylamido-2-phenyl) ethyl

| | | | chloromethyl ketone |
| LDH | lactate dehydrogenase | UPLC | ultra-high-performance liquid chromatography |

Acknowledgments

The authors were fully responsible for the content, editorial decisions, and opinions expressed in

the current article. The authors would like to acknowledge 360Biolabs Pty Ltd (Melbourne,

Australia) for the conduct of the CPE assays and Scripps Research Institute for the conduct of the

antiviral and virucidal assays. The authors would also like to thank the Peter Doherty Institute for

Infection and Immunity (Melbourne, Australia) for the gift to 360Biolabs of SARS-CoV-2

hCoV-19/Australia/VIC01/2020 used in these studies.

Funding

433 The research was funded by Starpharma Pty Ltd, which was responsible for study design,

434 interpretation of data, writing the manuscript and decision to submit the article for publication.

References

Bernstein, D.I., Stanberry, L.R., Sacks, S., Ayisi, N.K., Gong, Y.H., Ireland, J., Mumper, R.J.,

Holan, G., Matthews, B, McCarthy, T., Bourne, N., 2003. Evaluations of unformulated and

formulated dendrimer-based microbicide candidates in mouse and guinea pig models of

genital herpes. Antimicrob. Agents Chemother. 47(12), 3784-3788.

https://doi.org/10.1128/aac.47.12.3784-3788.2003

Bansal, S., Jonsson, C.B., Taylor, S.L., Figueroa, J.M., Vanesa, A.D., Palacios, C., Vega, J.C.,

2020. Iota-carrageenan and xylitol inhibit SARS-CoV-2 in cell culture. bioRxiv

2020.08.19.225854. https://doi.org/10.1101/2020.08.19.225854

Chavoustie, S.E., Carter, B.A., Waldbaum, A.S., Donders, G.G.G., Peters, K.H., Schwebke, J.R.,

Paull, J.R.A, Price, C.F., Castellarnau, A., McCloud, P., Kinghorn, G.R., 2020. Two phase

3, double-blinded, placebo-controlled studies of the efficacy and safety of Astodrimer 1%

Gel for the treatment of bacterial vaginosis. Eur. J. Obstet. Gynecol. Reprod. Biol. 245, 13-

18. https://doi.org/10.1016/j.ejogrb.2019.11.032

Chen, M.Y., Millwood, I.Y., Wand, H., Poynten, M, Law, M., Kaldor, J.M., Wesselingh, S.,

Price, C.F., Clark, L.J., Paull, J.R., Fairley, C.K., 2009. A randomized controlled trial of

the safety of candidate microbicide SPL7013 gel when applied to the penis. J. Acquir.

Immune Defic. Syndr. 50(4), 375-380. https://doi.org/10.1097/QAI.0b013e318198a7e6

Connell, B.J., Lortat-Jacob, H., 2013. Human immunodeficiency virus and heparan sulfate: from

attachment to entry inhibition. Frontiers Immunol. 4, 385.

https://doi.org/10.3389/fimmu.2013.00385

Ghosh, T., Chattopadhyay, K., Marschall, M., Karmakar, P., Mandal, P., Ray, B., 2009. Focus on

antivirally active sulfated polysaccharides: from structure-activity analysis to clinical

evaluation. Glycobiology 19(1), 2-15. https://doi.org/10.1093/glycob/cwn092

Gong, E., Matthews, B., McCarthy, T., Chu, J., Holan, G., Raff, J., Sacks, S., 2005. Evaluation

of dendrimer SPL7013, a lead microbicide candidate against herpes simplex virus.

Antiviral Res. 68(3), 139-146. https://doi.org/10.1016/j.antiviral.2005.08.004

Hoffmann, M., Kleine-Weber, H., Pöhlmann, S., 2020a. A multibasic cleavage site in the spike

protein of SARS-CoV-2 is essential for infection of human lung cells. Molecular Cell

78(4), 779-784.e5. https://doi.org/10.1016/j.molcel.2020.04.022

Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens,

T.S., Herrler, G., Wu, N.H., Nitsche, A., Müller, M.A., Drosten, C., Pöhlmann, S., 2020b.

SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically

proven protease inhibitor. Cell 181(2), 271-280.e8.

https://doi.org/10.1016/j.cell.2020.02.052

Hou, Y.J., Okuda, K., Edwards, C.E., et al. 2020. SARS-CoV-2 reverse genetics reveals a

variable infection gradient in the respiratory tract. Cell 182(2), 429-446.e14.

https://doi.org/10.1016/j.cell.2020.05.042

Jiang, Y.H., Emau, P., Cairns, S., Flanary, L., Morton, W.R., McCarthy, T., Tsai, C.C., 2005.

SPL7013 gel as a topical microbicide for prevention of vaginal transmission of $SHIV_{89.6P}$

in macaques. AIDS Res. Hum. Retroviruses 21(3), 207-213.

https://doi.org/10.1089/aid.2005.21.207

Korber, B., Fischer, W.M., Gnanakaran, S., Yoon, H., Theiler, J., Abfalterer, W., Foley, B.,

Giorgi, E.E., Bhattacharya, T., Parker, M.D., Partridge, D.G., Evans, C.M., de Silva, T.I.,

LaBranche, C.C., Montefiori, D.C., 2020. Spike mutation pipeline reveals the emergence

of a more transmissible form of SARS-CoV-2. bioRxiv.

https://doi.org/10.1101/2020.04.29.069054

Lackman-Smith, C., Osterling, C., Luckenbaugh, K., Mankowski, M., Snyder, B., Lewis, G.,

Paull, J., Profy, A., Ptak, R.G., Buckheit Jr., R.W., Watson, K.M., Cummins Jr., J.E.,

Sanders-Beer, B.E., 2008. Development of a comprehensive human immunodeficiency

virus type-1 screening algorithm for discovery and preclinical testing of topical

microbicides. Antimicrob. Agents Chemother. 52(5), 1768-1781.

https://doi.org/10.1128/AAC.01328-07

Lang, J., Yang, N., Deng, J., Liu, K., Yang, P., Zhang, G., Jiang, C., 2011. Inhibition of SARS

pseudovirus cell entry by lactoferrin binding to heparan sulfate proteoglycans. PLoS ONE

6(8), e23710. https://doi.org/10.1371/journal.pone.0023710

Liu, L., Chopra, P., Li, X., Wolfert, M.A., Tompkins, S.M., Boons, G.J., 2020. SARS-CoV-2

spike protein binds heparan sulfate in a length- and sequence-dependent manner. bioRxiv.

https://doi.org/10.1101/2020.05.10.087288

McCarthy, T.D., Karellas, P., Henderson, S.A., Giannis, M., O'Keefe, D.F., Heery, G., Paull,

J.R.A., Matthews, B.R., Holan, G., 2005. Dendrimers as drugs: discovery and preclinical

and clinical development of dendrimer microbicides for HIV and STI prevention. Mol.

Pharm. 2(4), 312-318. https://doi.org/10.1021/mp050023q

McGowan, I., Gomez, K., Bruder, K., Febo, I, Chen, B.A., Richardson, B.A., Husnik, M.,

Livant, E., Price, C., Jacobson, C., 2011. Phase 1 randomized trial of the vaginal safety and

acceptability of SPL7013 gel (VivaGel[®]) in sexually active young women (MTN-004).

AIDS 25(8), 1057-1064. https://doi.org/10.1097/QAD.0b013e328346bd3e

Milewska, A., Zarebski, M., Nowak, P., Stozek, K., Potempa, J., Pyrc, K., 2014. Human

coronavirus NL63 utilizes heparan sulfate proteoglycans for attachment to target cells. J.

Virol. 88(22), 13221-13230. https://doi.org/10.1128/JVI.02078-14

Mycroft-West, C., Su, D., Elli, S., Li, Y., Guimond, S., Miller, G., Turnbull, J., Yates, E.,

Guerrini, M., Fernig, D., Lima, M., Skidmore, M., 2020a. The 2019 coronavirus (SARS-

CoV-2) surface protein (spike) S1 receptor binding domain undergoes conformational

change upon heparin binding. bioRxiv. https://doi.org/10.1101/2020.02.29.971093

Mycroft-West, C.J., Su, D., Li, Y., Guimond, S., Rudd, T.R., Elli, S., Miller, G., Nunes, Q.M.,

Procter, P., Bisio, A., Forsyth, N.R., Turnbull, J.E., Guerrini, M., Fernig, D., Yates, E.A.,

Lima, M.A., Skidmore, M.A., 2020b. SARS CoV-2 spike S1 receptor binding domain

undergoes conformational change upon interaction with low molecular weight heparins.

bioRxiv. https://doi.org/10.1101/2020.04.29.068486

Mycroft-West, C.J., Su, D., Pagani, I., Rudd, T.R., Elli, S., Guimond, S., Miller, G., Meneghetti,

M.C.Z., Nader, H.B., Li, Y., Nunes, Q.M., Procter, P., Mancini, N., Clementi, M., Forsyth,

N.R., Turnbull, J.E., Guerrini, M., Fernig, D., Vicenzi, E., Yates, E.A., Lima, M.A.,

Skidmore, M.A., 2020c. Heparin inhibits cellular invasion of SARS-CoV-2: structural

dependence on the interaction of the surface protein (spike) S1 receptor binding domain

with heparin. bioRxiv 2020.2004.2028.066761. https://doi.org/10.1101/2020.04.28.066761

O'Loughlin, J., Millwood, I.Y., McDonald, H.M., Price, C.F., Kaldor, J.M., Paull, J.R.A., 2010.

Safety, tolerability, and pharmacokinetics of SPL7013 gel (VivaGel®): a dose ranging

phase 1 study. Sex. Transm. Dis. 37(2), 100-104.

https://doi.org/10.1097/OLQ.0b013e3181bc0aac

Ogando, N.S., Dalebout, T.J., Zevenhoven-Dobbe, J.C., Limpens, R.W.A.L., van der Meer, Y.,

Caly, L., Druce, J., de Vries, J.J.C., Kikkert, M., Bárcena, M., Sidorov, I., Snijder, J., 2020.

SARS-coronavirus-2 replication in Vero E6 cells: replication kinetics, rapid adaption and

cytopathology. J. General Virol. 10.1099/jgv.0.001453. Advance online publication.

https://doi.org/10.1099/jgv.0.001453

Pauwels, R., Balzarini, J., Baba, M., Snoeck, R., Schols, D., Herdewijn, P., Desmyter, J., De

Clercq, E., 1988. Rapid and automated tetrazolium-based colorimetric assay for the

detection of anti-HIV compounds. J. Virol. Methods 20(4), 309-321.

https://doi.org/10.1016/0166-0934(88)90134-6

Pizzorno, A., Padey, B., Dubois, J., Julien, T., Traversier, A., Dulière, V., Brun, P., Lina, B.,

Rosa-Calatrava, M., Terrier, O., 2020. In vitro evaluation of antiviral activity of single and

combined repurposable drugs against SARS-CoV-2. Antiviral Res. 104878. Advance

online publication. https://doi.org/10.1016/j.antiviral.2020.104878

Qiao, B., Olvera de la Cruz, M., 2020. The distal polybasic cleavage sites of SARS-CoV-2 spike

protein enhance spike protein-ACE2 binding. bioRxiv June 10.

https://doi.org/10.1101/2020.06.09.142877

Sarrazin, S., Lamanna, W.C., Esko, J.D., 2011. Heparan sulfate proteoglycans. Cold Spring Harb.

Perspect. Biol. 3(7), a004952. https://doi.org/10.1101/cshperspect.a004952

Schwebke, J., Carter, B., Waldbaum, A., Price, C., Castellarnau, A., Paull, J., McCloud, P.,

Kinghorn, G., 2021. A phase 3, randomized, controlled trial of Astodrimer 1% Gel for

preventing recurrent bacterial vaginosis. Eur. J. Obstet. Gynecol. Reprod. Biol. X 10,

100121. https://doi.org/10.1016/j.eurox.2021.100121

Sungnak, W., Huang, N., Bécavin, C., Berg, M., Queen, R., Litvinukova, M., Talavera-López, C.,

Maatz, H., Reichart, D., Sampaziotis, F., Worlock, K.B., Yoshida, M., Barnes, J.L., 2020.

SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with

innate immune genes. Nat. Med. 26(5), 681-687. https://doi.org/10.1038/s41591-020-0868-

6

Telwatte, S., Moore, K., Johnson, A., Tyssen, D., Sterjovski, J., Aldunate, M., Gorry, P.R.,

Ramsland, P.A., Lewis, G.R., Paull, J.R.A., Sonza, S., Tachedjian, G., 2011. Virucidal

activity of the dendrimer microbicide SPL7013 against HIV-1. Antiviral Res. 90(3), 195-

199. https://doi.org/10.1016/j.antiviral.2011.03.186

Tyssen, D., Henderson, S.A., Johnson, A., Sterjovski, J, Moore, K., La, J., Zanin, M., Sonza, S.,

Karellas, P., Giannis, M.P., Krippner, G., Wesselingh, S., McCarthy, T., Gorry, P.R.,

Ramsland, P.A., Cone, R., Paull, J.R.A., Lewis, G.R., Tachedjian, G., 2010. Structure

activity relationship of dendrimer microbicides with dual action antiviral activity. PLoS

ONE 5(8), e12309. https://doi.org/10.1371/journal.pone.0012309

van den Worm, S.H., Eriksson, K.K., Zevenhoven, J.C., Weber, F., Zust, R., Kuri, T., Dijkman,

R., Chang, G., Siddell, S.G., Snijder, E., Thiel, V., Davidson, A.D., 2012. Reverse genetics

of SARS-related coronavirus using vaccinia virus-based recombination. PLoS ONE

7(3):e32857. https://doi.org/10.1371/journal.pone.0032857

Waldbaum, A.S., Schwebke, J.R., Paull, J.R.A., Price, C.F., Edmondson, S.R., Castellarnau, A.,

McCloud, P., Kinghorn, G.R., 2020. A phase 2, double-blind, multicenter, randomized,

placebo-controlled, dose-ranging study of the efficacy and safety of Astodrimer Gel for the

treatment of bacterial vaginosis. PLoS ONE 15(5), e0232394.

https://doi.org/10.1371/journal.pone.0232394

Walls, A.C., Park, Y.J., Tortorici, M.A., Wall, A., McGuire, A.T., Veesler, D., 2020. Structure,

function and antigenicity of the SARS-CoV-2 spike glycoprotein. Cell 181(2), 281-292.e6

https://doi.org/10.1016/j.cell.2020.02.058

Wang, M., Cao, R., Zhang, L., Yang, X., Liu, J., Xu, M/., Shi, Z., Hu, Z., Zhong, W., Xiao, G.,

2020. Remdesivir and chloroquine effectively inhibit the recently emerged novel

coronavirus (2019-nCoV) in vitro. Cell Res. 30(3), 269-271.

https://doi.org/10.1038/s41422-020-0282-0

Wrapp, D., Wang, N., Corbett, K.S., Goldsmith, J.A., Hsieh, C.L., Abiona, O., Graham, B.S.,

McLellan, J.S., 2020. Cryo-EM structure of the 2019-nCoV spike in the prefusion

conformation. Science 367(6483), 1260-1263. https://doi.org/10.1126/science.abb2507

Yuan, M., Wu, N.C., Zhu, X., Lee, C.C.D., So, R.T.Y., Lv, H., Mok, C.K.P., Wilson, I.A., 2020.

A highly conserved cryptic epitope in the receptor binding domains of SARS-CoV-2 and

SARS-CoV. Science 368(6491), 630-633. https://doi.org/10.1126/science.abb7269

582