Consistent variations in personality traits and their potential for genetic improvement in the

biocontrol agent *Trichogramma evanescens*

Silène Lartigue[a,b,c], Myriam Yalaoui[b], Jean Belliard[b], Claire Caravel[b], Louise Jeandroz[b],

Géraldine Groussier[b], Vincent Calcagno[b], Philippe Louâpre[c], François-Xavier Dechaume-

Moncharmont[d], Thibaut Malausa[b] and Jérôme Moreau[c, e]

[a] ENGREF AgroParisTech, Paris, France

[b] UMR Institut Sophia Agrobiotech, INRAE, UCA, CNRS, 06903 Sophia Antipolis, France

[c] UMR CNRS 6282 Biogéosciences, Université Bourgogne Franche-Comté, 6 Boulevard

Gabriel, 21000 Dijon, France

[d] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, 69622

Villeurbanne, France

[e] Centre d'Études Biologiques de Chizé, UMR 7372, CNRS & La Rochelle Université, 79360

Villiers-en-bois, France

**\* Corresponding author:** S. Lartigue, UMR Institut Sophia Agrobiotech, INRAE, UCA,

CNRS, Sophia Antipolis, France. E-mail address: silene.lartigue@gmail.com

**Abstract**

Improvements in the biological control of agricultural pests require improvements in the phenotyping methods used by practitioners to select efficient biological control agent (BCA) populations in industrial rearing or field conditions. Consistent inter-individual variations in behaviour (i.e. animal personality) probably affect BCA efficiency, but have never been taken into account in the development of phenotyping methods, despite having characteristics useful for phenotyping: repeatable (by definition), often heritable, etc. We developed a video-tracking method targeting animal personality traits and evaluated the feasibility of its use for genetic improvement in the BCA *Trichogramma evanescens*, by phenotyping 1,049 individuals from 24 isogenic lines. We found consistent individual variations in boldness, activity and exploration. Personality differences between the 24 isogenic lines suggested a genetic origin of the variations in activity and exploration (broad-sense heritability estimates of 0.06 to 0.11) and revealed a trade-off between exploration and fecundity.

**Key words:** biocontrol, animal personality, *Trichogramma*, genetic improvement, intraspecific variation, behavioural ecology, behavioural syndrome, genetic correlations, pace-of-life

**Introduction**

The demand for more sustainable agriculture is increasing worldwide (Godfray et al., 2010; Willer & Lernoud, 2019). Various elements can be used in the development of sustainable strategies, and biological control (BC) is one such element that is currently attracting considerable attention (van Lenteren 2012). Most BC methods are based on the choice, rearing and introduction of biological control agent (BCA) populations able to control the target pests (Eilenberg, Hajek, & Lomer, 2001). Choosing the right BCA is key to the success of pest regulation programmes and is based on (i) the ability of the BCA to control pest populations in the field, (ii) its potential to adapt to the release environment, (iii) its expected impact on local biodiversity, and (iv) the feasibility of mass-rearing and storing the BCA in industrial conditions (Briese, 2000; Kruitwagen, Beukeboom, & Wertheim, 2018; Sforza, 2010). The identification of BCA species or populations with as many of the desired features as possible is time-consuming and complex, particularly given that the choice of non-indigenous species before use as BCAs is constrained by increasingly strict regulations for the protection of biodiversity (Lommen, Jong, & Pannebakker, 2017).

Phenotyping is key for (i) the efficient characterisation of traits related to the desirable features of BCAs listed above, (ii) smart choices of BC taxa when screening the available natural enemy diversity and (iii) the management of phenotypic evolution in industrial contexts involving rearing procedures and quality control (Kruitwagen et al., 2018; Lommen et al., 2017). However, the phenotyping methods currently used in the choice of BCAs or for quality control are mostly low-throughput and based on single proxies of fitness, such as predation or parasitism rate, size, sex ratio, longevity, or developmental rate (Hopper, Roush, & Powell, 1993; Prezotti, Parra, Vencovsky, Coelho, & Cruz, 2004; Roitberg, Boivin, & Vet, 2001; Smith, 1996). These proxies are intuitively correlated with fitness under laboratory conditions, but their actual relevance for biocontrol, in industrial mass-rearing or field conditions, remains a

matter of debate (Lommen et al., 2017; Roitberg et al., 2001). This situation calls for drastic

improvements in the phenotyping capacities of the community involved in BC research and

innovation.

Behavioural traits are among the most promising of the traits to which more attention

could be paid in BCA phenotyping procedures. Most behavioural traits are likely to affect the

performance of BCA both during industrial mass rearing and in the field (Roitberg, 2007;

Wajnberg, 2009; Wajnberg, Roitberg, & Boivin, 2016). Indeed, studies of BCA behavioural

traits have suggested that these traits could (i) facilitate the selection of BCAs that are specific

to the targeted pest, (ii) improve release strategies (through studies of the BCA response to pre-

release handling or BCA mating behaviour, for example), and (iii) predict the efficiency of

target pest suppression by the BCA (Mills & Kean, 2010). However, there have been few

studies of BCA behavioural traits for the development of phenotyping methods, and behaviour

has been largely neglected by those using BC (Mills & Kean, 2010; Wajnberg, Bernstein, &

Alphen, 2008).

As a consequence, the current state-of-the-art for insect behavioural studies displays

several key limitations. The first limitation is the lack of diversity of possible target traits for

phenotyping. Indeed, although many studies have focused on traits relating to foraging

behaviour (Lirakis & Magalhães, 2019; Mills & Wajnberg, 2008), tools for measuring other

aspects of behaviour remain scarce. A second limitation is the insufficient focus on the

intraspecific variation of traits. Such variation has been comprehensively investigated for only

a limited number of BCA species and a limited number of traits (Kruitwagen et al. 2018; Lirakis

and Magalhães 2019). This situation is detrimental because the investigation of only a fraction

of the available intraspecific variability makes it difficult to identify the populations displaying

the highest performance for biocontrol, and prevents the development of efficient genetic

improvement programmes based on selective breeding and controlled evolution (Wajnberg

2004; Bolnick et al. 2011; Lommen et al. 2017; Kruitwagen et al. 2018, Lirakis and Magalhães

2019). A third limitation is the reliance of most choices in BC exclusively on comparisons

between average trait values for species or populations (Lommen et al., 2017). Published studies

have suggested that individual variation can affect the characteristics of the population thought

to be important for BC (Biro & Stamps, 2008; Michalko, Pekár, & Entling, 2019; Réale, Reader,

Sol, McDougall, & Dingemanse, 2007; Wolf & Weissing, 2012).

One way to overcome each of these three limitations would be to apply approaches used

in the field of animal personality to BC. Indeed, these approaches provide a framework offering

(i) sets of behavioural traits rarely studied in BC and displaying features (repeatability,

heritability) that make them good candidates for use in genetic improvement for BC, and (ii)

phenotyping methods suitable for analyses of intraspecific variation, including inter-individual

variation. Animal personality research focuses on inter-individual differences in behaviour that

are consistent over time and context (Dingemanse, Kazem, Reale, & Wright, 2009; Réale et al.,

2007). Interest in animal personality has increased over the last few decades, and studies have

been performed on diverse taxa, including insects (Amat, Desouhant, Gomes, Moreau, &

Monceau, 2018; Bell, Hankison, & Laskowski, 2009; Dingemanse et al., 2009; Gosling, 2001;

Kralj-fiser & Schuett, 2014; Mazué, Dechaume-Moncharmont, & Godin, 2015; Monceau et al.,

2017; Réale et al., 2007; Sih, Bell, & Johnson, 2004; van Ooers & Sinn, 2011) and, more

specifically, insects used as BC agents (Gomes, Desouhant, & Amat, 2019; Michalko et al.,

2019). Réale et al. (2007) described five main categories of personality traits: boldness,

exploration, activity, aggressiveness and sociability. Personality traits have been shown to be

correlated with traits relevant for pest control, such as foraging capacity, fecundity, growth,

survival (Biro & Stamps, 2008), dispersal ability (Cote, Fogarty, Weinersmith, Brodin, & Sih,

2010) and insecticide resistance (Royauté, Buddle, & Vincent, 2014). These traits are probably,

therefore, of interest in the context of BC. Moreover, personality traits are repeatable, by

definition, and can be heritable (Dochtermann, Schwab, & Sih, 2014; Réale et al., 2007;

Stirling, Reale, & Roff, 2002), making them suitable tools for genetic improvement. From a

methodological point of view, animal personality provides valuable information for the design

of phenotyping and genetic improvement strategies in BC. Indeed, animal personality studies

are based on standardised methods designed to measure inter-individual variation and to

investigate correlations between traits (e.g. by looking for behavioural syndromes) (Réale et

al., 2007; Sih et al., 2004). This is particularly relevant to the objective of selecting several

combined BC traits rather than a single trait, as recently recommended by Lommen et al. (2017)

and Kruitwagen et al. (2018). The investigation of correlations between traits is also important,

to detect trade-offs that may constrain genetic improvement programmes or affect BC traits if

mass-rearing causes uncontrolled trait selection (Mackauer, 1976).

In this study, we assessed the potential for BCA phenotyping based on concepts and

methods used in the field of animal personality. We used the egg parasitoid *Trichogramma*

*evanescens* Westwood, 1833 (Hymenoptera: Trichogrammatidae) as a test species.

*Trichogramma* micro-wasps are used worldwide in augmentative BC against lepidopteran pests

(Hassan, 1993; van Lenteren, 2012). Their economic importance (Robin & Marchand, 2020;

Thibierge, 2015) justifies investments in research and development aiming to improve their

genetic potential. Our aims were (i) to determine whether behavioural traits meeting the criteria

of personality traits could be measured in these micro-wasps of approximately 0.5 mm in

length; (ii) to investigate the relationships between personality traits and traits classically

measured on BCAs in the laboratory, and (iii) to determine whether personality traits could be

used in genetic improvement strategies for BCAs. We developed a method based on the video-

tracking and measuring, at individual level, of multidimensional behavioural traits relating to

boldness, activity and exploration. We investigated the relationships between these traits, and

between these traits and individual fitness traits relevant for mass rearing (offspring number,

longevity, tibia length). We then compared the traits between 24 near-isogenic strains, to obtain

a first insight into the broad-sense heritability of these traits. We looked for genetic correlations

potentially constraining the use of these traits for genetic improvement.

**Methods**

*Laboratory rearing of T. evanescens*

We used 24 near-isogenic lines (hereafter referred to as "lines") of *Trichogramma*

*evanescens*. Lines were created from inbred crosses in populations established from individuals

sampled in different parts of France (geographic origins detailed in Table 7 in the appendix),

from 2010 to 2016, and reared in the laboratory at $18 \pm 1$ °C, $70 \pm 10\%$ RH and 16:8 h L:D

(details of the protocol followed to create the lines are provided in the appendix). Genetic

diversity within lines was below 1.1 alleles per locus at 19 microsatellite loci (unpublished

data), and individuals within lines were considered genetically identical. We created two

sublines for each line (Lynch & Walsh, 1998), to disentangle the confounding effects of rearing

tubes and lines (which may be caused by maternal effects). We considered variation between

lines to be of genetic origin, and variation within lines to be of environmental origin. We reared

*Trichogramma evanescens* individuals on sterilised *Ephestia kuehniella* Zeller 1879

(Lepidoptera: Pyralidae) eggs, renewed every 10 days, at $25.5 \pm 1$ °C, $70 \pm 10\%$ RH and 16:8

h L:D (Schöller & Hassan, 2001). We kept populations in glass tubes (height: 73 mm, diameter:

11 mm), and fed adults with honey *ad libitum*.

*Measurement of variables*

*General experimental design*

160 The following experimental design was used to measure phenotypic traits in

161 *Trichogramma* females (Figure 1). We used *T. evanescens* females that had mated within the

162 last 24 hours, randomly chosen from each line. We checked the physical integrity of these

females, which were isolated in glass tubes before the beginning of the experiment (height: 73 mm, diameter: 11 mm) and fed with honey, *ad libitum*. On the first two days, we assessed the behavioural traits of the females. We estimated the number of offspring on days 3 to 5, and longevity from day 6. The experiment lasted from May to July 2019, and was split into 17 experimental sessions, in each of which, we used three females per line. The physiological, developmental and behavioural traits of *Trichogramma* wasps, and of *T. evanescens* in particular, are dependent on temperature (Ayvaz, Karasu, Karabörklü, Tunçbilek, 2008; Schöller & Hassan, 2001). Moreover, as Suverkropp et al. (2001) showed that *T. brassicae* individuals have similar levels of activity throughout the day at temperatures of about 25 °C or higher, we assumed that our *T. evanescens* individuals had similar responses to temperature throughout the day. Therefore, we performed the behavioural experiments at 25.5 ± 1 °C, 70 ± 10% RH. We then measured female longevity and offspring number at 18 ± 1 °C, 70 ± 10% RH, to ensure that the females would live long enough for the final stages of the experiment (Cônsoli & Parra, 1995; Schöller & Hassan, 2001).

**Figure 1.** Overview of the experimental design, for one session.

*Behavioural trait measurement*

We observed individuals in an arena composed of two sheets of glass (24 cm x 18 cm), one for the floor and one for the ceiling. The 2 mm space between them was maintained by walls made of a black rubber sheet. We placed this arena on an LCD screen (Samsung© 28" LED, 3840*2160 pixels, 60 Hz) displaying a white circle with a diameter of 5.5 cm on a dark background (Figure 2.a). The LCD screen was turned on one hour before the beginning of the

experiment, to ensure that a stable temperature of 25.5 ± 1 °C was achieved in the area. The

conditions in the growth chamber in which the experimental design was set up were as follows:

22.5 ± 1 °C and 70 ± 10% RH. We used a fine paintbrush to introduce a randomly chosen

female into the centre of the arena while the screen was showing a white background. The glass

ceiling was replaced, and we then switched to a background with a white circle on a dark

background, with the female positioned in the middle of the white circle. We observed the

behaviour of the female for 90 seconds, with video recording at 25 frames per second (with a

resolution of 1080 p), with a Nikon© D750 camera (Figure 2.a).

We then analysed the videos files, determining the 2D spatial position ($x$-$y$ coordinates)

and body orientation (in radians) of the female on each frame, with C-trax software (Branson,

Robie, Bender, Perona, & Dickinson, 2009). We independently determined the exact position

of the border between the white circle and the black background with ImageJ software

(Abràmoff, Magalhães, & Ram, 2004). We thus defined regions of interest of 0.5 cm on either

side of the border, for investigation of the behaviour of the insect near the border (Figure 2.b).

We imported the C-trax and ImageJ output files into R software v.3.6.1 (R Core Team 2019)

and cleaned our data to remove tracking artefacts. We used the "trajr" package (Mclean &

Volponi, 2018) to calculate speed and sinuosity, in each region of interest. We calculated seven

variables linked to three personality traits — boldness, exploration and activity — as defined

by Réale et al. (2007).

Boldness is the reaction of the individual to a risky situation (Réale et al., 2007). We

estimated boldness by measuring three variables. The first was the time until the female first

entered the dark area (area 3 in Figure 2.b). Higher values indicated that the female took longer

to cross the border, which we interpreted as meaning that the female was less bold. The second

and third variables were the absolute difference in speed between areas 2 and 1 (Figure 2.b) and

the absolute difference in sinuosity between areas 2 and 1 (Figure 2.b). We considered high

values for these two variables to indicate a larger change in behaviour at the border, which we

interpreted as meaning that the female was more affected by the border and was, therefore, less

bold. Exploration represents the individual's reaction to a new environment (Réale et al., 2007).

Exploration was estimated in area 1 as (1) the total area explored per unit time, and (2)

the mean sinuosity of the pathway (Figure 2.b). For this variable, we hypothesised that the

females with the most winding pathways remained closer to their release point, indicating a

lower level of exploration.

Finally, we measured activity in area 4, so exploration and activity were measured in

different areas of the experimental arena. Activity was estimated as (i) the proportion of time

the female was considered to be active (with a speed of more than 0.01 centimetres per second),

referred to hereafter as "activity rate", and (ii) mean speed (Figure 2.b), considering faster

movement to be indicative of a higher level of activity.

We estimated the repeatability of measurements, by conducting two observations per

female, with 24 hours between the two measurements, a time interval corresponding to 20% of

the mean lifespan of this species. Females were tested in a random order on day 1, and then in

the same order on day 2, to ensure that all individuals had exactly the same time interval

between two measurements. Between behavioural experiments, each female was placed back

in its glass tube and fed with honey, *ad libitum*, in a growth chamber at $25.5 \pm 1$ °C, $70 \pm 10\%$

RH and 16:8 h L:D. Behavioural trait measurements were obtained for 776 to 996 females from

the 24 lines.

**Figure 2.** Experimental set-up of the behavioural experiment. Fig. 2.a. shows a photograph of the experimental

setup: the LCD screen displaying the white circle on a dark background, the arena and the Nikon[©] D750 camera

above. Fig. 2.b. represents the defined areas of the arena. The grey shading corresponds to the dark background,

the white part indicates the white circle, and the dark cross is the site at which the female was placed at the start

of the experiment. The dotted lines represent the virtual borders defined between areas 1 and 2, and between areas

3 and 4.

*Offspring number, longevity and tibia length measurement*

After the second day of behavioural observation, females were kept in their glass tubes

at $18 \pm 1$ °C, $70 \pm 10\%$ RH and 16:8 h L:D and fed with honey, *ad libidum*. We provided each

female with a piece of paper 4.50 cm x 0.85 cm in size, covered with *E. kuehniella* eggs, *ad*

*libidum*. *E. kuehniella* eggs were removed 72 hours later and placed in conditions of $25.5 \pm 1$

°C, $70 \pm 10\%$ RH and 16:8 h L:D. Three days later, we counted the number of parasitised eggs

(considered as black eggs), to estimate the size of the progeny of each female over a period of

72 hours, providing a proxy for female fitness. From day 6, we measured female longevity (the

females were still kept in the same individual tubes with honey, but with no *E. kuehniella* eggs,

at $18 \pm 1$ °C, $70 \pm 10\%$ RH and 16:8 h L:D). Tubes were checked every day at 5 p.m., until the

death of the female. Dead females were conserved in ethanol, for subsequent measurement of

tibia length on a micrograph (obtained with an Axioplan II microscope), with ImageJ software

(Abràmoff et al., 2004). Images were acquired at the Microscopy Platform of Sophia

Agrobiotech Institute, INRA, UNS, CNRS, UMR 1355-7254, INRA PACA, Sophia Antipolis.

Not all individuals lived long enough for all the phenotypic measurements to be made. We

therefore collected progeny measurements for 929 females, longevity measurements for 655

females and tibia size measurements for 959 females, from all 24 lines.

*Data analysis*

We used the R software v.3.6.1 for all statistical analyses. For each variable, we first

fitted a linear mixed model with the lme4 package (Bates, Maechler, Bolker, & Walker, 2015),

with individual, line, subline and session as random effects. For each variable, data

transformations were chosen after graphical inspection of the distribution of model residuals,

estimated with the "simulateResiduals" function of the DHARMa package (Hartig, 2019). We

performed logarithmic transformations for all behavioural variables except for the area explored

within area 1. We addressed several questions regarding the data, and the data analysis for each

of these questions is presented below.

*Are the measured behavioural traits repeatable?*

We first estimated the repeatability of the behavioural traits measured with generalised

linear mixed models, using the rptR package (Stoffel, Nakagawa, & Schielzeth, 2017). The

"rptGaussian" function of the rptR package was used to provide repeatability estimates. As

repeatability can be defined as the proportion of variation explained by between-individual

variation (Nakagawa & Schielzeth, 2010), we included only two random effects in these

models: individual (assuming that the effects of line and subline on variation were included in

the individual effect) and session, with individual as a grouping factor. In subsequent analyses,

we considered only traits that were significantly repeatable.

*Do the measured traits identify individual behavioural strategies?*

Based on methods generally used in animal personality studies, we first investigated

correlations between behavioural traits and then summarized the data by principal component

analysis (PCA). We first obtained a single value for each trait for each individual, by extracting,

from the linear mixed model described above, linear predictors for each individual, with the

"ranef" function of the lme4 package. We used these values to measure the phenotypic

correlation between traits, by calculating Spearman's rank correlation coefficients, to determine

whether individuals adopted different strategies, or whether it was possible to describe

behavioural syndromes. We estimated bootstrapped 95% confidence intervals from 1000

bootstraps, to assess the significance of the Spearman's rank correlation coefficients obtained

(Nakagawa & Cuthill, 2007), using the "spearman.ci" function of the RVAideMemoire package

(Hervé, 2020). *P*-values were adjusted by the false discovery rate method (Benjamini &

Hochberg, 1995). We then performed PCA with the "PCA" function of the FactoMineR

package (Le, Josse, & Husson, 2008), using both values obtained for each individual (days 1

and 2, when available). We estimated two synthetic personality scores based on the first two

axes of the PCA. We used the "fviz_pca_biplot" function of the factoextra package

(Kassambara & Mundt, 2019) to obtain a graphical representation of the correlation between

repeatable behavioural traits and the distribution of individual values along the two first axes

of the PCA.

*Are the measured traits correlated with fitness-related traits?*

We studied the correlation between behavioural and fitness-related traits, using the same

linear mixed model as described in the introduction to this section. We extracted linear

predictors (using the "ranef" function of the lme4 package (Bates et al., 2015)) for each

individual and each personality score from this model. We assessed the correlation between the

linear predictors of these personality traits and scores, and offspring number, body size and

longevity, by calculating Spearman's rank correlation coefficients. We estimated bootstrapped

95% confidence intervals to assess significance of the Spearman's rank correlation coefficients

obtained, with the same R function and method as described above. *P*-values were adjusted by

the false discovery rate method.

*Are the measured traits heritable?*

We sought to establish a first estimate of broad-sense heritability for each trait. To this

end, we followed the simple design proposed by Lynch and Walsh (1998) for clonal

populations, and approximated the proportion of the variance explained by genetic factors with

an estimate of the proportion of variance explained by the line effect in our generalised linear

mixed models. This estimate was obtained with the "rptGaussian" function of the rptR package

(Stoffel et al., 2017), with models including line, subline, individual and session as random

effects, and line as a grouping factor.

*Do personality traits differentiate the isogenic lines?*

We compared the personality scores of the 24 lines, taking into account variation due to

individual, subline and session effects. With the values of each personality score extracted from

the PCA (see above), we first fitted a linear mixed-effects model with the "lmer" function of

the lme4 package (Bates et al., 2015), with line as a fixed effect and individual, subline and

session effects as random effects. We performed a Tukey all-pairs comparison on lines with the

"glht" function of the multcomp package (Hothorn, Bretz, & Westfall, 2008). We graphically

represented the distribution of each line along the two personality scores, for the same PCA as

described above, estimated from individual values. We then used the "plot.PCA" function of

the FactoMineR package to represent only mean point values for each line on the graph.

*Are personality traits genetically correlated with fitness-related traits?*

We investigated the genetic correlation between genetic traits, using the same linear

mixed model as described in the introduction to this section. We first extracted linear predictors

for each line and trait, with the "ranef" function of the lme4 package. We then used these values

to calculate Spearman's rank correlation coefficients. We estimated bootstrapped 95%

confidence intervals, to assess significance of the Spearman's rank correlation coefficients, and

adjusted the *p*-values as described above.

**Results**

*Are the measured behavioural traits repeatable?*

Repeatability estimates for the seven behavioural traits ranged from 0.04 to 0.35 (Table

1). The repeatability estimates had confidence intervals excluding zero for all traits except for

"time to first crossing of the border between the white and black areas" (Table 1). Only

repeatable traits were considered in the subsequent analysis.

**Table 1.** Estimated repeatability (R) and 95% confidence intervals (between square brackets) for behavioural traits.

Repeatable traits (R-value in bold type) were used to estimate personality scores.

| Personality trait category | Variable assessed | R [95% CI] |
|---|---|---|
| Activity | Mean speed in area 4 | **0.35** [0.29; 0.40] |
| | Activity rate in area 4 | **0.08** [0.01; 0.14] |
| Boldness | Change of speed in the border area (area 2) | **0.10** [0.04; 0.17] |
| | Change of sinuosity in the border area (area 2) | **0.12** [0.04; 0.19] |
| | Time to first crossing of the white/black border | 0.04 [0.00; 0.11] |
| Exploration | Sinuosity in area 1 | **0.24** [0.17; 0.30] |
| | Area explored in area 1 | **0.18** [0.12; 0.24] |

*Do the measured traits identify individual behavioural syndromes?*

All repeatable variables were correlated with at least one other variable (Table 2),

indicating the existence of a behavioural syndrome. We combined these six variables into two

personality scores based on the first two axes of a PCA, which accounted for 56.8% of the

variance (Table 3). The first axis (personality score 1, PC1) was positively correlated with the

"area explored in area 1" and inversely correlated with "sinuosity in area 1" and with the

"change of sinuosity in the border area 2" (Table 3). Highly positive values of PC1

corresponded to a high exploration score (Figure 3). The second axis (personality score 2, PC2)

and correlated mostly with "mean speed in area 4", "activity rate in area 4" and the "change of

speed in border area 2" (Table 3). High positive values of PC2 correspond to high activity scores

(Figure 3).

**Table 2.** Phenotypic correlation between behavioural variables, with Spearman's rank correlation coefficient Rho

and 95 percent confidence intervals (between square brackets), based on a number of individual values from N =

977 to N = 1009. Correlation coefficients with confidence intervals excluding zero are shown in bold, and

correlation coefficients remaining significantly different from zero after Benjamini and Hochberg correction are

indicated with an asterisk. The personality trait category to which each variable belongs is indicated in brackets:

activity (A), boldness (B) and exploration (E).

|  | (B) Change of speed in border area 2 | (A) Mean speed in area 4 | (A) Activity rate | (B) Change of sinuosity in border area 2 | (E) Sinuosity in area 1 |
|---|---|---|---|---|---|
| (A) Mean speed in area 4 | **0.31** **[0.25; 0.37] *** | | | | |
| (A) Activity rate in area 4 | **0.10** **[0.04; 0.16] *** | **0.38 *** **[0.32; 0.43] *** | | | |
| (B) Change of sinuosity in border area 2 | **0.11** **[0.05; 0.17] *** | **0.07** **[0.01; 0.14] *** | **-0.12** **[-0.18; -0.06] *** | | |
| (E) Sinuosity in area 1 | **-0.07** **[-0.14; -0.01] *** | **0.13** **[0.07; 0.19] *** | **-0.16** **[-0.22; -0.10] *** | **0.38** **[0.32; 0.44] *** | |
| (E) Area explored in area 1 | **0.11** **[0.04; 0.16] *** | 0.01 [-0.04; 0.08] | **0.29** **[0.23; 0.34] *** | **-0.28** **[-0.34; -0.22] *** | **-0.56** **[-0.61; -0.52] *** |

356　**Table 3.** Parameters from the first two principal components (PC1 and PC2) of the PCA for the behavioural

357　variables measured. Component loadings represent the relationship between the principal components and the

358　variables from which they are constructed. The personality trait category to which each variable belongs is

359　indicated in brackets: activity (A), boldness (B) and exploration (E).

| Parameter | PC1 | PC2 |
|---|---|---|
| Eigenvalue | 1.87 | 1.54 |
| Percentage of variance explained | 31.23 | 25.58 |
| Component loading | | |
| *(A) Mean speed in area 4* | 0.16 | 0.84 |
| *(A) Activity rate in area 4* | 0.43 | 0.60 |
| *(B) Change of speed in area 2* | 0.19 | 0.51 |
| *(B) Change of sinuosity in area 2* | -0.56 | 0.34 |
| *(E) Area explored in area 1* | 0.81 | -0.09 |
| *(E) Sinuosity in area 1* | -0.81 | 0.27 |

**Figure 3.** Graphical representation of the first two axes of the PCA on individual values (grey points) for

repeatable behavioural traits (in black type).

*Are the measured traits correlated with fitness-related traits?*

Active females (i.e. those with higher PC2 values) had significantly larger numbers of

offspring and significantly longer tibias (Table 4). Higher rates of exploration (i.e. higher PC1

values) were not significantly correlated with any of the fitness-related traits measured. None

of the behavioural variables or personality scores was significantly correlated with longevity

(Table 4).

**Table 4.** Phenotypic correlation between behavioural traits (behavioural variables and personality scores) and other life history traits (with Spearman's rank correlation coefficient Rho and 95% confidence intervals (between square brackets) calculated from 959 individual values). Correlation coefficients with confidence intervals excluding zero are shown in bold, and correlation coefficients that remained significantly different from zero after Benjamini and Hochberg correction are indicated with an asterisk. The personality trait category to which each variable belongs is indicated in brackets: activity (A), boldness (B) and exploration (E).

|  | Offspring number | Longevity | Tibia length |
|---|---|---|---|
| Behavioural variables | | | |
| *(A) Mean speed in area 4* | **0.20 [0.14; 0.26] *** | -0.05 [-0.12; 0.03] | **0.19 [0.12; 0.25] *** |
| *(A) Activity rate in area 4* | -0.01 [-0.08; 0.06] | -0.06 [-0.13; 0.02] | -0.07 [-0.13; 0.00] |
| *(B) Change of speed in border area 2* | **0.13 [0.06; 0.19] *** | -0.08 [-0.15; 0.00] | **0.16 [0.10; 0.21] *** |
| *(B) Change of sinuosity in border area 2* | **0.11 [0.04; 0.17] *** | 0.002 [-0.07; 0.08] | 0.05 [-0.01; 0.11] |
| *(E) Area explored in area 1* | -0.05 [-0.11; 0.01] | -0.05 [-0.13; 0.02] | -0.02 [-0.09; 0.04] |
| *(E) Sinuosity in area 1* | 0.01 [-0.05; 0.07] | 0.05 [-0.02; 0.13] | 0.05 [-0.01; 0.11] |
| Personality scores | | | |
| *Exploration score 1 (PC1)* | -0.01 [-0.07; 0.06] | -0.05 [-0.13; 0.03] | -0.03 [-0.09; 0.03] |
| *Activity score 2 (PC2)* | **0.17 [0.10; 0.23] *** | -0.01 [-0.10; 0.07] | **0.15 [0.09; 0.21] *** |

*Are the measured traits heritable?*

Broad-sense heritability estimates for behavioural traits and personality scores ranged from 0.01 to 0.11. Confidence intervals excluded zero for all traits linked to activity and exploration, whereas they included zero for the two traits linked to boldness (Table 5). Fitness-related traits (offspring number, tibia length and longevity) displayed broad-sense heritability ranging from 0.04 to 0.28, with all confidence intervals excluding zero (Table 5).

**Table 5.** Broad-sense heritability (H²) of traits measured with 95% confidence intervals (between square brackets). Heritability estimates are shown in bold if their 95% confidence interval did not include zero. The personality trait category to which each behavioural variable belongs is indicated in brackets: activity (A), boldness (B) and exploration (E).

| | H² [95% CI] |
|---|---|
| Behavioural variables | |
| *(A) Mean speed in area 4* | **0.11** [0.05; 0.18] |
| *(A) Activity rate in area 4* | 0.02 [0.00; 0.04] |
| *(B) Change of speed in border area 2* | 0.01 [0.00; 0.03] |
| *(B) Change of sinuosity in border area 2* | 0.01 [0.00; 0.03] |
| *(E) Area explored in area 1* | **0.06** [0.02; 0.10] |
| *(E) Sinuosity in area 1* | **0.06** [0.02; 0.11] |
| Personality scores | |
| *Exploration score 1 (PC1)* | **0.08** [0.03; 0.13] |
| *Activity score 2 (PC2)* | **0.05** [0.02; 0.10] |
| Fitness-related traits | |
| *Offspring number* | **0.12** [0.05; 0.19] |
| *Tibia length* | **0.05** [0.01; 0.09] |
| *Longevity* | **0.28** [0.14; 0.39] |

*Do personality traits differentiate between lines?*

We found significant differences in personality scores between lines (Figure 4.a and 4.b), and the 24 lines were distributed along the first two axes of the PCA (Figure 5). We were therefore able to distinguish between lines that were very active and exploratory (e.g., lines 3 and 12), and lines that were less active and exploratory (e.g., lines 14 and 21); we were also able to distinguish between lines that were very exploratory but not very active (e.g., lines 9 and 10) and lines that were active but not very exploratory (for example line 4).

**Figure 4.** Boxplot of the adjusted values of personality score 1 (Fig. 4.a) and personality score 2 (Fig. 4.b) after the elimination of variation due to individual, subline and session effects, and compact letter display after Tukey all-pair comparisons. Two lines with no letters in common are considered to be significantly different (with a p-value <0.05).

**Figure 5.** Distribution of the mean points for the 24 lines (centroids) along the first two axes of the PCA.

*Are personality traits genetically correlated with fitness-related traits?*

       The only genetic correlation between personality scores and fitness-related traits that remained significant after FDR correction was the positive correlation between exploration score (PC1) and offspring number (Table 6).

**Table 6.** Genetic correlation between personality and other life history traits (Spearman's rank correlation coefficient, based on the trait estimates of 24 near-isogenic lines, with associated p-values in brackets). Correlation coefficients with confidence intervals excluding zero are shown in bold, and correlation coefficients that remained significantly different from zero after Benjamini and Hochberg correction are indicated with an asterisk. The personality trait category to which each behavioural variable belongs is indicated in brackets: activity (A), boldness (B) and exploration (E).

|  | Offspring number | Longevity | Tibia length |
| --- | --- | --- | --- |
| Behavioural variables |  |  |  |
| *(A) Mean speed in area 4* | -0.16 [-0.54; 0.33] | -0.32 [-0.62; 0.06] | **0.51 [0.21; 0.70]** |

| | | | |
|---|---|---|---|
| *(A) Activity rate in area 4* | **-0.53 [-0.78; -0.14]** | 0.15 [-0.28; 0.54] | -0.21 [-0.62; 0.28] |
| *(B) Change of speed in border area 2* | -0.03 [-0.44; 0.37] | **-0.45 [-0.72; -0.10]** | 0.35 [-0.05; 0.63] |
| *(B) Change of sinuosity in border area 2* | 0.29 [-0.11; 0.63] | 0.15 [-0.31; 0.56] | 0.19 [-0.20; 0.54] |
| *(E) Area explored in area 1* | **-0.60 [-0.79; -0.26] *** | 0.01 [-0.40; 0.43] | -0.28 [-0.65; 0.16] |
| *(E) Sinuosity in area 1* | **0.63 [0.33; 0.82] *** | 0.25 [-0.20; 0.62] | -0.03 [-0.43; 0.39] |
| Personality scores | | | |
| *Exploration score 1 (PC1)* | **-0.64 [-0.83; -0.29] *** | -0.09 [-0.52; 0.32] | -0.19 [-0.58; 0.24] |
| *Activity score 2 (PC2)* | -0.10 [-0.52; 0.39] | -0.22 [-0.66; 0.19] | **0.41 [0.01; 0.67]** |

**Discussion**

We investigated whether animal personality could be used to develop or improve phenotyping methods for the BCA *Trichogramma evanescens*. We first developed an automated phenotyping method based on automated pathway analysis, providing a set of behavioural trait measures that proved repeatable over time and heritable (i.e. personality traits). We then identified differences in life history strategies between individuals as behavioural traits were correlated together and combined them into personality scores, which were correlated with other life history traits. Finally, we observed differences in personality scores between the 24 genotypes of *T. evanescens* and found a negative genetic correlation between exploration and fecundity.

*Evidence of personality traits in* **Trichogramma evanescens**

Personality has never before been assessed in a species as small as *Trichogramma evanescens*. Based on other video-tracking studies in other species (Branson et al., 2009; Charalabidis, Dechaume-Moncharmont, Petit, & Bohan, 2017), we designed and developed a video-tracking approach measuring a large number of variables relating to the movements of *T. evanescens* individuals during their presence in the different areas (white, black and border areas) within an experimental arena. Here, we chose to work on seven variables that (i) could be calculated with methods commonly used in trajectory and movement studies (speeds, trajectory length and sinuosity estimates) (Mclean & Volponi, 2018) and (ii) we considered to

be associated with some of the commonly defined personality traits defined by Réale et al.

(2007): boldness, exploration and activity.

For each of the seven behavioural variables, we assessed repeatability, broad-sense

heritability and phenotypic and genetic correlations between personality traits and between

these traits and other life history traits, according to methods generally used in animal

personality studies (Réale et al., 2007). For six of the seven behavioural variables, we observed

significant repeatability (R) (values ranging from 0.10 to 0.35, Table 1). The R values obtained

were within the range of R values commonly observed for behavioural traits, although most

were lower than the mean R value obtained for animal behaviour (0.35) (Bell et al., 2009).

However, personality has rarely been studied in parasitoid insects, and a recent study on the

parasitoid wasp *Venturia canescens* reported a similar R value for activity and a lower R value

for exploration (about 0.10, whereas we obtained R values for exploration-related variables of

0.18 and 0.24 (Gomes et al., 2019)).

The broad-sense heritability of the variables (ranging from 0.06 to 0.11, Table 5) was

lower than the mean value for animal behaviour (0.24) in the meta-analysis by Dochtermann et

al. (2019). Stirling et al. (2002) found no significant differences in heritability between

behavioural and life-history traits in their meta-analysis, whereas we found that heritability

values for personality traits were lower than heritability values of two classical fitness-related

traits (offspring number and longevity) in *T. evanescens* (Table 5).

Behavioural traits could be grouped together into two continuums or behavioural

syndromes (Réale et al., 2007; Sih et al., 2004; Sih, Cote, Evans, Fogarty, & Pruitt, 2012): a

continuum extending from individuals with low levels of exploratory behaviour to highly

exploratory individuals, and a continuum extending from individuals with low levels of activity

to highly active individuals (Figure 3). Bold (or shy) behaviour and active behaviour have been

shown to be correlated with fecundity traits in several species (Biro & Stamps, 2008), but rarely

in insects (Monceau et al., 2017). In this study, we found a weak but significant phenotypic correlation between behavioural traits, fecundity and body length, as shy or active females produced more offspring, and had longer tibias (Table 4). Finally, an analysis of genetic correlations showed that the lines with the most exploratory individuals had the smallest numbers of offspring (Table 6). These correlations seem to be compatible with the pace-of-life syndrome (POLS) hypothesis, a currently debated hypothesis (Royauté, Berdal, Garrison, & Dochtermann, 2018), according to which, behavioural traits are related to morphological, physiological and other life-history traits (Réale et al. 2010).

*Potential of personality traits for use in genetic improvement of the biocontrol agent* **Trichogramma evanescens**

In this study, our aim was to evaluate the possibility of using personality traits as traits of interest in biological control, and of integrating these traits into genetic improvement programmes for the BCA *T. evanescens*. The six repeatable behavioural traits we measured were correlated with each other, and could be combined into two continuums. For each individual and continuum, we estimated a personality score corresponding to the position of the individual along the continuum, a common method in animal personality studies (Mazué et al., 2015; Monceau et al., 2017). We found that it was possible to capture a large proportion of the behavioural trait variance with two scores (36.2% of the total variance explained by personality score 1, and 26.4% explained by personality score 2). This finding highlights the utility of calculating a few synthetic indices (or scores), rather than measuring large numbers of variables, to obtain relevant information for BC. We therefore systematically present our results considering all the traits individually and summarized as two personality scores.

The relevance of the behavioural traits or synthetic scores to the context of BC was demonstrated by the phenotypic correlations between these traits and scores and the traits classically measured in BC (fecundity, longevity and body length) (Hopper et al., 1993; Prezotti

et al., 2004; Roitberg et al., 2001; Smith, 1996). In this study, active females (i.e. with high

values for "mean speed in area 4" and "personality score 2") produced more offspring and had

longer tibias (Table 4). By contrast, we found that bold females (i.e. with low values for "change

of speed in border area 2" and "change of sinuosity in border area 2") produced a small number

of offspring (Table 4). In several species, activity and boldness behaviours have been shown to

be correlated with traits of ecological importance, such as dispersal (Sih et al., 2004), which is

also a trait linked to field efficiency in BC (Fournier & Boivin, 2000). Our results indicate that

active females produce more offspring, which is predictive of a high degree of efficiency in

rearing conditions and in the field. Note, however, that we did not assess survival or body

condition in the offspring. The same females also displayed shyer behaviour, which would

presumably have a negative impact on field efficiency. Further studies are, therefore, required

to assess the full ecological relevance of these lines in BC. The relevance of the variables

measured will be confirmed only if they are shown to be correlated with BC performance in

industrial and field conditions.

Most of our data analyses aimed to evaluate the added value of the measured behavioural

traits for genetic improvement strategies, breeding programs. We found that personality scores

differ among isogenic lines (Figure 4.a. and Figure 4.b.) and that these differences highlight

contrasted behaviours, as evidenced by their distribution along the two personality scores in

Figure 5. This may make it possible to differentiate between these behaviours and to select for

them, should they prove relevant in terms of BC efficiency. We also observed a negative genetic

correlation between the personality score relating to exploration and offspring production. It

will probably be important to take this trade-off into account in BC, as it may oppose

performance in rearing and performance in the field. Indeed, as for activity and boldness,

exploration behaviours are also correlated with traits linked to field efficiency in BC, such as

dispersal (Fournier & Boivin, 2000; Sih et al., 2004).

Given these results, and the ease with which all the traits can be assessed and personality scores obtained through short (90 seconds) automated video-tracking measurements, the new method described here may provide useful criteria for the selection of candidate BCA taxa (populations, strains, sibling species, etc.) or for quality control purposes. However, the high level of intra-isogenic line variability observed (Figure 4.a and Figure 4.b), accounting for the relatively low broad-sense heritability of the traits and scores (between 0.01 and 0.11; Table 5), constrains the use of this method, as it may be necessary to phenotype large numbers of individuals for reliable comparisons between taxa or reared populations. The low heritability also constitutes an obstacle to the implementation of ambitious experimental evolution programmes. Oriented experimental evolution may be fastidious for traits displaying such a high degree of environmentally induced variability. As a comparison, breeding programmes for livestock animals generally make use of traits with higher heritability. Heritability values for morphological, physiological, behavioural or other traits linked to fitness and considered in these breeding programmes generally range from 0.17 to 0.70 in sheep, pigs, cows and fish (Juengel et al., 2019; Kavlak & Uimari, 2019; Moretti, de Rezende, Biffani, & Bozzi, 2018; Vargas Jurado, Leymaster, Kuehn, & Lewis, 2016).

**Conclusion**

In conclusion, the use of methods and concepts of animal personality to develop phenotyping methods and associated data analyses for BC led to the rapid phenotyping of traits rarely used in BC that were repeatable, heritable and correlated with fitness-related traits. However, it will be possible to consider the actual potential of these traits and of the phenotyping method satisfactory only after investigating the relationships between the laboratory-measured traits and BC performance indices in real BC situations, in industrial production settings or in field releases. This first study has driven the launch of large-scale field

experiments, which are currently underway and aim to generate field-release performance indices.

**Acknowledgements**

We would like to thank all the members of staff at Institut Sophia Agrobiotech who helped us when we were overwhelmed by the sheer numbers of *Trichogramma* individuals in our experimental tubes, including Léa Tregoat-Bertrand in particular. Special thanks to Maxime Dahirel for helpful advice for statistical analysis, and to Morgane Bequet-Rennes and Simon Vaïsse, who contributed to this study by running previous experiments that led to our decision to design this new phenotyping assay. We thank the biocontrol company BIOLINE, in particular Paloma Martinez and Benoît Franck, and the INRAE biological resource centre "Egg-Parasitoids Collection" (CRB EP-Coll, Sophia Antipolis) for providing field-sampled populations for the establishment of some of the isogenic lines used in this study. We thank the Microscopy Platform of the Sophia Agrobiotech Institute, INRA, UNS, CNRS, UMR 1355-7254, INRA PACA, Sophia Antipolis for providing access to instruments and technical advice. This work was funded by the the French "Agence Nationale de la Recherche" under the "Investissements d'Avenir UCAJEDI" project with reference n ° ANR-15-IDEX-01. Silène Lartigue received financial support from the French Ministry in charge of Agriculture.

**Conflict of interest disclosure**

The authors of this article declare that they have no financial conflict of interest with the content of this article.

**Data and R code**

Data tables and code needed to re-do the analyses and figures are available on Zenodo. (http://doi.org/10.5281/zenodo.4058218).

## References

Abràmoff, M. D., Magalhães, P. J., & Ram, S. J. (2004). Image processing with imageJ. *Biophotonics International*, *11*(7), 36–41.

Amat, I., Desouhant, E., Gomes, E., Moreau, J., & Monceau, K. (2018). Insect personality: what can we learn from metamorphosis? *Current Opinion in Insect Science*, *27*, 46–51.

Ayvaz, A., Karasu, E., Karabörklü, S., Tunçbilek, A. Ş. (2008). Effects of cold storage, rearing temperature, parasitoid age and irradiation on the performance of *Trichogramma evanescens* Westwood (Hymenoptera: Trichogrammatidae). *Journal of Stored Products Research*, *44*, 232–240.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

Bell, A. M., Hankison, S. J., & Laskowski, K. L. (2009). The repeatability of behaviour: a meta-analysis. *Animal Behaviour*, *77*(4), 771–783.

Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300.

Biro, P. A., & Stamps, J. A. (2008). Are animal personality traits linked to life-history productivity? *Trends in Ecology and Evolution*, *23*(7), 361–368.

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., … Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology and Evolution*, *26*(4), 183–192.

Branson, K., Robie, A. A., Bender, J., Perona, P., & Dickinson, M. H. (2009). High-throughput ethomics in large groups of *Drosophila*. *Nature Methods*, *6*(6), 451–457.

Briese, D. T. (2000). Classical biological control. *Australian Weed Management Systems*, 161–192.

Charalabidis, A., Dechaume-Moncharmont, F. X., Petit, S., & Bohan, D. A. (2017). Risk of predation makes foragers less choosy about their food. *PLoS ONE*, *12*(11), 1–18.

Cônsoli, F. L., & Parra, J. R. P. (1995). Effects of constant and alternating temperatures on *Trichogramma galloi* Zucchi (Hym., Trichogrammatidae) biology II.-parasitism capacity and longevity. *Journal of Applied Entomology*, *119*, 667–670.

Cote, J., Fogarty, S., Weinersmith, K., Brodin, T., & Sih, A. (2010). Personality traits and dispersal tendency in the invasive mosquitofish (*Gambusia affinis*). *Proceedings of the Royal Society B: Biological Sciences*, *277*(1687), 1571–1579.

Dingemanse, N. J., Kazem, A. J. N., Reale, D., & Wright, J. (2009). Behavioural reaction norms: animal personality meets individual plasticity. *Trends in Ecology and Evolution*, *25*(2), 81–89.

Dochtermann, N. A., Schwab, T., Anderson Berdal, M., Dalos, J., & Royauté, R. (2019). The heritability of behavior: a meta-analysis. *Journal of Heredity*, *110*(4), 403–410.

Dochtermann, N. A., Schwab, T., & Sih, A. (2014). The contribution of additive genetic variation to personality variation: heritability of personality. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1798).

Eilenberg, J., Hajek, A., & Lomer, C. (2001). Suggestions for unifying the terminology in biological control. *BioControl*, *46*(4), 387–400.

Fournier, F., & Boivin, G. (2000). Comparative dispersal of *Trichogramma evanescens* and *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae) in relation to environmental conditions. *Population Ecology*, *29*(1), 55–63.

Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., … Toulmin, C. (2010). Food security: the challenge of feeding 9 billion people. *Science*, *327*, 812–818.

Gomes, E., Desouhant, E., & Amat, I. (2019). Evidence for risk-taking behavioural types and potential effects on resource acquisition in a parasitoid wasp. *Animal Behaviour*, *154*, 17–28.

Gosling, S. D. (2001). From mice to men: what can we learn about personality from animal research? *Psychological Bulletin*, *127*(1), 45–86.

Hartig, F. (2019). DHARMa: Residual diagnostics for hierarchical (multi-level / mixed) regression models. Retrieved from https://cran.r-project.org/package=DHARMa

Hassan, S. A. (1993). The mass rearing and utilization of *Trichogramma* to control lepidopterous pests, achievements and outlook. *Pesticide Science*, *37*, 387–391.

Hervé, M. (2020). RVAideMemoire: testing and plotting procedures for biostatistics. Retrieved from https://cran.r-project.org/package=RVAideMemoire

Hopper, K. R., Roush, R. T., & Powell, W. (1993). Management of genetics of biological-control introductions. *Annual Review of Entomology*, *38*, 27–51.

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric models. *Biometrical Journal*, *50*(3), 346–363.

Juengel, J. L., Hickey, S. M., Clarke, S. M., Cullen, N. G., McEwan, J. C., & Dodds, K. G. (2019). Heritability of ram mating success in multi-sire breeding situations. *Animal*, *13*(5), 917–923.

Kassambara, A., & Mundt, F. (2019). factoextra: extract and visualize the results of multivariate data analyses. Retrieved from https://cran.r-project.org/package=factoextra

Kavlak, A. T., & Uimari, P. (2019). Estimation of heritability of feeding behaviour traits and their correlation with production traits in Finnish Yorkshire pigs. *Journal of Animal Breeding and Genetics*, *136*(6), 484–494.

Kralj-fiser, S., & Schuett, W. (2014). Studying personality variation in invertebrates: why bother? *Animal Behaviour*, *91*, 41–52.

Kruitwagen, A., Beukeboom, L. W., & Wertheim, B. (2018). Optimization of native biocontrol agents, with parasitoids of the invasive pest *Drosophila suzukii* as an example. *Evolutionary Applications*, *11*(9), 1473–1497.

Le, S., Josse, J., & Husson, F. (2008). FactoMineR: an R package for multivariate analysis. *Journal of Statistical Software*, *25*(1), 1–18.

Lirakis, M., & Magalhães, S. (2019). Does experimental evolution produce better biological control agents? A critical review of the evidence. *Entomologia Experimentalis et Applicata*, *167*(7), 584–597.

Lommen, S. T. E., Jong, P. W. De, & Pannebakker, B. A. (2017). It is time to bridge the gap
between exploring and exploiting: prospects for utilizing intraspecific genetic variation to
optimize arthropods for augmentative pest control – a review. *Entomologia Experimentalis*
*et Applicata*, *168*, 108–123.

Lynch, M., & Walsh, B. (1998). *Genetics and Analysis of Quantitative Traits*. Sunderland, MA:
Sinauer.

Mackauer, M. (1976). Genetic problems in the production of biological control agents. *Annual*
*Review of Entomology*, *21*(1), 369–385.

Mazué, G. P. F., Dechaume-Moncharmont, F. X., & Godin, J. G. J. (2015). Boldness-
exploration behavioral syndrome: Interfamily variability and repeatability of personality
traits in the young of the convict cichlid (*Amatitlania siquia*). *Behavioral Ecology*, *26*(3),
900–908.

Mclean, D. J., & Volponi, M. A. S. (2018). trajr: An R package for characterisation of animal
trajectories, (March), 440–448.

Michalko, R., Pekár, S., & Entling, M. H. (2019). An updated perspective on spiders as
generalist predators in biological control. *Oecologia*, *189*(1), 21–36.

Mills, N. J., & Kean, J. M. (2010). Behavioral studies, molecular approaches, and modeling:
methodological contributions to biological control success. *Biological Control*, *52*(3),
255–262.

Mills, N. J, & Wajnberg, E. (2008). Optimal foraging behaviour and efficient biological control
methods. *Behavioural Ecology of Insect Parasitoids–From Theoretical Approaches to*
*Field Applications*, 3–30.

Monceau, K., Dechaume-Moncharmont, F. X., Moreau, J., Lucas, C., Capoduro, R., Motreuil,
S., & Moret, Y. (2017). Personality , immune response and reproductive success: an
appraisal of the pace-of-life syndrome hypothesis. *Journal of Animal Ecology*, *86*, 932–
942.

Moretti, R., de Rezende, M. P. G., Biffani, S., & Bozzi, R. (2018). Heritability and genetic
correlations between rumination time and production traits in Holstein dairy cows during
different lactation phases. *Journal of Animal Breeding and Genetics*, *135*(4), 293–299.

Nakagawa, S., & Cuthill, I. C. (2007). Effect size, confidence interval and statistical
significance: A practical guide for biologists. *Biological Reviews*, *82*(4), 591–605.

Nakagawa, S., & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists, 935–956.

Prezotti, L., Parra, J. R. P., Vencovsky, R., Coelho, A. S. G., & Cruz, I. (2004). Effect of the
size of the founder population on the quality of sexual populations of *Trichogramma*
*pretiosum*, in laboratory. *Biological Control*, *30*(2), 174–180.

R Core Team. (2019). R: A language and environment for statistical computing. Vienna: R
Foundation for Statistical Computing. Retrieved from https://www.r-project.org/

Réale, D., Garant, D., Humphries, M. M., Bergeron, P., Careau, V., & Montiglio, P. O. (2010).
Personality and the emergence of the pace-of-life syndrome concept at the population
level. *Philosophical Transactions of the Royal Society of London. Series B, Biological*
*Sciences*, *365*(1560), 4051–4063.

Réale, D., Reader, S. M., Sol, D., McDougall, P. T., & Dingemanse, N. J. (2007). Integrating animal temperament within ecology and evolution. *Biological Reviews*, *82*(2), 291–318.

Robin, D. C., & Marchand, P. A. (2020). Macroorganismes de biocontrôle en France, état des lieux. *Innovations Agronomiques*, *79*, 425–439.

Roitberg, B. D. (2007). Why pest management needs behavioral ecology and vice versa. *Entomological Research*, *37*, 14–18.

Roitberg, B. D., Boivin, G., & Vet, L. E. M. (2001). Fitness, parasitoids, and biological control: an opinion. *The Canadian Entomologist*, *133*, 429–438.

Royauté, R., Berdal, M. A., Garrison, C. R., & Dochtermann, N. A. (2018). Paceless life? A meta-analysis of the pace-of-life syndrome hypothesis. *Behavioral Ecology and Sociobiology*, 72–64.

Royauté, R., Buddle, C. M., & Vincent, C. (2014). Interpopulation variations in behavioral syndromes of a jumping spider from insecticide-treated and insecticide-free orchards. *Ethology*, *120*(2), 127–139.

Schöller, M., & Hassan, S. A. (2001). Comparative biology and life tables of *Trichogramma evanescens* and *T. cacoeciae* with *Ephestia elutella* as host at four constant temperatures. *Entomologia Experimentalis et Applicata*, *98*(1), 35–40.

Sforza, R. (2010). Recherche et évaluation des candidats pour la lutte biologique classique. In *La lutte biologique. Vers de nouveaux équilibres écologiques* (Quae-Educa, pp. 280–285).

Sih, A., Bell, A., & Johnson, J. C. (2004). Behavioral syndromes: an ecological and evolutionary overview. *Trends in Ecology and Evolution*, *19*(7), 372–378.

Sih, A., Cote, J., Evans, M., Fogarty, S., & Pruitt, J. (2012). Ecological implications of behavioural syndromes. *Ecology Letters*, *15*, 278–289.

Smith, S. M. (1996). Biological control with *Trichogramma*: advances, successes, and potential of their use. *Annual Review of Entomology*, *41*, 375–406.

Stirling, D. G., Reale, D., & Roff, D. A. (2002). Selection, structure and the heritability of behaviour. *Journal of Evolutionary Biology*, *15*, 277–289.

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods in Ecology and Evolution*, *8*(11), 1639–1644.

Suverkropp, B. P., Bigler, F., & Van Lenteren, J. C. (2001). Temperature influences walking speed and walking activity of *Trichogramma brassicae* (Hym., Trichogrammatidae). *Journal of Applied Entomology*, *125*(6), 303–307.

Tabone, E., Bardon, C., Desneux, N., & Wajnberg, E. (2010). Parasitism of different *Trichogramma* species and strains on *Plutella xylostella* L. on greenhouse cauliflower. *Journal of Pest Science*, *83*(3), 251–256.

Thibierge, J. (2015). Innovations Agronomiques 46 (2015), 27-37, *46*, 27–37.

van Lenteren, J. C. (2012). The state of commercial augmentative biological control: Plenty of natural enemies, but a frustrating lack of uptake. *BioControl*, *57*(1), 1–20.

van Ooers, K., & Sinn, D. L. (2011). Toward a basis for the penotypic gambit: Advances in the

evolutionnary genetics of animal personalities. In *From genes to animal behavior* (pp. 165–183). Springer, Tokyo.

Vargas Jurado, N., Leymaster, K. A., Kuehn, L. A., & Lewis, R. M. (2016). Estimating heritability of wool shedding in a cross-bred ewe population. *Journal of Animal Breeding and Genetics*, *133*(5), 396–403.

Wajnberg, E. (2004). Measuring genetic variation in natural enemies used for biological control: why and how ? In *Genetics, Evolution and Biological Control*.

Wajnberg, E. (2009). Genetics of the behavioral ecology of egg parasitoids. In *Egg Parasitoids in Agroecosystems with Emphasis on Trichogramma* (pp. 149–165). Springer, Dordrecht.

Wajnberg, E., Bernstein, C., & Alphen, J. J. M. Van. (2008). *Behavioral ecology of insect parasitoids: from theoretical approaches to field applications*. Blackwell Publishing.

Wajnberg, E., Curty, C., & Jervis, M. (2012). Intra-population genetic variation in the temporal pattern of egg maturation in a parasitoid wasp. *PLoS ONE*, *7*(9).

Wajnberg, E., Roitberg, B. D., & Boivin, G. (2016). Using optimality models to improve the ef fi cacy of parasitoids in biological control programmes. *Entomologia Experimentalis et Applicata*, *158*, 2–16.

Willer, H., & Lernoud, J. (2019). *The world of organic agriculture: Statistics and emerging trends 2019*. Bonn.

Wolf, M., & Weissing, F. J. (2012). Animal personalities: consequences for ecology and evolution. *Trends in Ecology & Evolution*, *27*(8), 452–461.