Theta and alpha power across fast and slow timescales in cognitive control

Pieter Huycke[1], Pieter Verbeke[1], C. Nico Boehler[1], and Tom Verguts[1]

[1]Department of Experimental Psychology, Ghent University

Author Note

We have no known conflict of interest to disclose.

Correspondence concerning this article should be addressed to Pieter Huycke,

Department of Experimental Psychology, Henri Dunantlaan 2, 9000 Ghent, Belgium. E-mail:

pieter.huycke@ugent.be

Abstract

Theta and alpha frequency neural oscillations are important for learning and cognitive control, but their exact role has remained obscure. In particular, it is unknown whether they operate at similar timescales, and whether they support different cognitive processes. We recorded EEG in 30 healthy human participants while they performed a procedural learning task containing both novel (block-unique) and repeating stimuli. We investigated behavior and electrophysiology at both fast (i.e., within blocks) and slow (i.e., between blocks) time scales. Behaviorally, both response time and accuracy improved over both fast and slow timescales. At the same time, on the spectral level, theta power significantly decreased along the slow timescale, whereas alpha power instead significantly increased along the fast timescale. We thus demonstrate that theta and alpha both play a role during learning, but operate at different timescales. This result poses important empirical constraints for theories on learning, cognitive control, and neural oscillations.

Keywords: cognitive control, procedural learning, frontomedial theta, posterior alpha

Theta and alpha power across fast and slow timescales in cognitive control

In daily life, agents frequently have to remember, apply, and automatize novel stimulus-action mappings, a process which falls under the general umbrella of *cognitive control* (Botvinick et al., 2001; Ridderinkhof, 2004). An example is learning to type using a computer keyboard. The first few attempts at typing a sentence, one is slow and might have to look at the keyboard, but over time complete sentences can be typed without delay or numerous errors.

Previous research has started to identify the neural signatures of cognitive control. In the spectral domain, theta band (4 – 8 Hz) oscillations were originally observed in the hippocampus (Buzsáki, 2002; O'Keefe, 1993; Vertes & Kocsis, 1997) and considered to be crucial for memory formation. More recently, theta was also observed in structures of the cerebral cortex, and became intensively studied in the cognitive-control literature (Cavanagh & Frank, 2014). These theta oscillations are thought to be generated in the frontal-medial (FM) cortex of the human brain (Anguera et al., 2013; Enriquez-Geppert et al., 2014; Sauseng et al., 2019). In their seminal review, Cavanagh and Frank (2014) related theta to cognitive control by showing that theta power increased for novel and difficult stimuli, as well as for negative feedback (stimulus-locked) and errors (response-locked). An emerging consensus suggests that cortical theta coordinates long-range interactions for implementation of cognitive control. For example, using EEG in a Stroop task, Hanslmayr et al. (2008) showed that phase coupling between the anterior cingulate cortex and the left prefrontal cortex lasted longer for incongruent stimuli than for neutral or congruent ones. Several other studies have described theta frequency phase locking between frontal and other areas in humans using EEG (Cohen, 2009; Nigbur et al., 2012; van de Vijver et al., 2011), as well as in non-human animals using intracranial recordings (Narayanan et al., 2013; Phillips et al.,

2014). In short, recent literature suggests that theta plays a fundamental role in the exertion of cognitive control.

Alpha band (8-12 Hz) activity has also been implicated in cognitive control. Two main effects are found when considering the interplay between alpha and cognitive control. First, alpha power increases in task-irrelevant brain areas (Klimesch, 1999; Mazaheri et al., 2009; Pfurtscheller, 2003). To explain this increase, Jensen and Mazaheri (2010) proposed that alpha activity blocks irrelevant processing pathways, which in turn allows the relevant brain areas to process the presented information more efficiently. This hypothesis was named 'gating by inhibition'. Evidence comes from a study by Jokisch and Jensen (2007), in which participants completed a delayed-match-to-sample task, discriminating either the identity or the orientation of a presented picture of a face. Consistent with gating by inhibition, the authors found that alpha activity increased in the brain area that was irrelevant for the current condition. Thus, an increase in alpha power might be linked to inhibition of a targeted (task-irrelevant) area. Second, alpha power decreases in task-relevant brain areas (Händel et al., 2011; Jensen & Mazaheri, 2010; Thut, 2006). Using a go-no-go task, Mazaheri et al. (2009) observed alpha suppression in task-relevant brain areas. Thus, whereas cognitive control is typically accompanied by high FM theta power, it usually co-occurs with low alpha power.

Generally, both theta and alpha are modulated by cognitive control, albeit in a different direction (in- and decrease for theta and alpha, respectively). Also on a trial-to-trial basis, theta and alpha power are (negatively) correlated. Mazaheri et al. (2009) observed in their go-no-go task that post-trial anterior theta power was negatively correlated with post-trial posterior alpha power, and especially so after error trials.

Given these tight couplings between theta and alpha, what is their respective role in cognitive control? One possibility is that theta and alpha simply reflect two elements of the

same mechanism, which manifest in frontal and posterior areas respectively. In this view, theta and alpha would always be correlated across time and conditions, yet originate from different neural areas. An alternative view is that theta and alpha both implement control, but at different timescales. This would not be visible with standard paradigms where the involvement of theta and alpha is not tracked across time for the same stimuli. Indeed, the transition from controlled to automatic processing may require many repetitions and consist of several stages (Ashby et al., 2007), with modulations of different oscillations potentially involved at different stages. Current protocols cannot disentangle these possibilities.

To address this problem, we devised a paradigm that is able to investigate how cognitive control unfolds over both a fast and a slow timescale. We started from a stimulus-action learning paradigm by Ruge and Wolfensteller (2010). In their fMRI experiment, subjects learned four unique stimulus-action mappings at the start of each experimental block through instruction. Each stimulus-action mapping was shown eight times within one experimental block. The authors investigated changes in neural activity as a function of the number of times a stimulus was seen within a block. On the behavioral level, subjects improved with practice, as indicated by both faster reaction times (RTs) and fewer errors. On the neural level, the authors found that areas that were hypothesized to be involved in automatization (striatum and the pre- and postcentral gyri) became more active with practice. In contrast, brain areas that were presumably related to cognitive control (such as the lateral prefrontal cortex and the intraparietal sulcus) became less active with increasing practice.

In the current work, we investigate the timescales of cognitive control, and its transition into automatic processing, and in particular the role of theta and alpha oscillations in this process. For that purpose we presented one set of four stimulus-action mappings repeatedly within a block (fast time scale, as in Ruge and Wolfensteller, (2010)). The same set of mappings also repeated across blocks (slow time scale), interleaved with blocks where

novel mappings were presented. Furthermore, we measured EEG, which allowed us to track theta and alpha power changes over the two timescales, and investigate if theta and alpha show the same pattern across the two timescales, or not.

Methods

The experimental procedure was approved by the Ghent University Ethical committee (Faculty of Psychology and Educational Sciences). All 30 participants signed the informed consent. In return for their participation, a monetary compensation of €25 was provided. All participants reported to be free from neurological conditions and had normal or corrected-to-normal vision. All participants were Dutch native speakers, and three of them were left-handed. The experiment started either at 9:00 A.M. or 1:00 P.M. and always lasted two hours. Data from 24 subjects (18 – 47 years old, 23.1 ± 5.6 (mean ± SD) years (17 females, 22 right-handed) were analyzed and reported. Criteria for exclusion were 1) data loss (6.66%), 2) inferior data quality (10%), and 3) technical issues (3.33%).

Design and Experimental Protocol

The experiment took place in a Faraday cage and consisted of a procedural learning task. Subjects learned stimulus-action mappings through trial-and-error learning. Subjects responded by pressing the "f" and "j" keys on the keyboard of the experiment computer using their left and right index fingers respectively. Each experimental block started with four stimuli that were presented in a vertical order (familiarization phase; no action mapping information provided; see Figure 1A). Participants were instructed at the beginning of the experiment that these would be the stimuli that they would encounter in the following block.

Figure 1. Visualization of the stimuli seen during the experiment, and a trial overview. (A) An example of the stimulus overview that was provided at the start of each experimental block. This overview displayed the four stimuli that would be shown in the following block for general stimulus familiarization (without providing action mapping information). (B) An overview of an experimental trial. Each trial lasted exactly 6500 ms in total.

After stimulus inspection, participants could start the experimental block by pressing the space bar on their keyboard. Following this, a five-second counter started, and the first trial of that block would start. A trial started with a fixation cross colored in either blue or black, balanced across subjects. Participants were informed prior to the start of the experiment what these colored fixation crosses meant. For half of the subjects, a blue fixation cross meant that a stimulus was about to appear, while a black fixation cross signified the end of the trial. For the other half of subjects, the opposite pattern applied. The duration of the first fixation cross was temporally jittered, in that the presentation time was drawn from a uniform distribution (1000-2000 milliseconds (ms)). Then, participants saw the stimulus framed by a black box in the center of the screen. From that moment, the subject had 1500 ms

to press one of the two response buttons. Only when a button was pressed in time, the box would turn grey, indicating that an answer was registered. After 1500 ms, the stimulus disappeared from the screen, and was replaced by three dots which we will refer to as the 'anticipation phase'. This stimulus timing was again temporally jittered, with a presentation time between 500 and 750 ms. Following the anticipation phase, the feedback (right, wrong, or too late) was provided in the form of a simple Dutch text message ("Correct answer", "Wrong answer", "Too slow") in the middle of the screen. The feedback presentation time depended on the presentation time of the anticipation phase. Together, these two phases lasted for 2000 ms. Finally, as mentioned earlier, a colored fixation cross marked the end of the trial. The presentation time of this fixation cross depended on the presentation time of the first fixation cross: Together they were shown for 3000 ms. Figure 1B provides a visual representation of a trial in our experiment. Subjects learned the correct stimulus-action mappings via trial-to-trial feedback.

The stimuli in each block were presented in semi-random order. Specifically, stimulus presentation order was generated so that stimuli that were not presented recently had a higher likelihood to be shown in the next trial. Thus, trial number in an experimental block and the number of times a stimulus is seen in a block are positively correlated. The first two blocks (64 trials in total) were performed as practice before the EEG set-up. Each block comprised of four stimuli that were each shown eight times, yielding a total of 32 trials per block. The experiment (without practice) lasted for 16 blocks, summing to 512 trials in total per subject. Note that at the end of this experiment a subject had seen 72 unique stimuli. Of these 72 stimuli, eight stimuli served as repeating stimuli (with constant response mapping), while the other 64 were unique ('novel') stimuli. Each repeating stimulus was seen 64 times, while the novel stimuli were seen eight times each. The stimuli used in an experimental run were randomly drawn from a collection of 124 stimuli. Half of the blocks (256 trials) contained

repeating stimuli; the other blocks contained novel stimuli. The order in which experimental blocks were presented was again semi-random: participants would never see more than two blocks in a row of the same condition (novel/repeating). Participants could take a break at the end of each block, which was denoted by a message on the screen. No time limits were imposed on these breaks. When they were ready to continue, subjects could press the space button, after which they would see the stimuli from the next block. Participants used a cushioned chinrest during EEG recording. The experiment was created using PsychoPy 3 (Peirce, 2007).

Data Recording and Initial Processing

All EEG data were collected using a 64-channel BioSemi Active Two System (BioSemi, Amsterdam, Netherlands) at 1024 Hz. The electrodes were placed following the standard international 10-20 electrode mapping defined by Jasper (1958). The cap contained a posterior CMS-DRL electrode combination. Six additional external electrodes were placed on the left and right mastoid, lateral canthi of both eyes, and above and below the left eye. The data collected using these electrodes were used as a reference during analysis (the average of the mastoid electrodes), or for detecting and correcting eye movements and eye blinks (the other external electrodes). Data preprocessing was done using the MNE Python software (Gramfort et al., 2014), and custom code. Our pipeline was inspired by the EEGLAB preprocessing pipeline (Delorme & Makeig, 2004), but some adaptations were made. First, the data were re-referenced to the average of the mastoid electrodes. Data was then visually inspected to mark bad channels, which were then interpolated. For 66.66% of analyzed subjects at least one channel was interpolated, with a maximum of six interpolated channels across subjects. The majority of interpolated electrodes were located in the posterior area of the scalp. The next step was to high-pass filter this data at 0.1 Hz, and to split this filtered data in epochs locked to stimulus onset. We then performed a visual inspection of

these epochs and excluded identifiably noisy epochs from further analysis. This exclusion process was performed blind towards the experimental conditions. Our next step would be to run ICA on the data epochs. However, Winkler et al. (2015) demonstrated that the efficiency of ICA in reducing artifacts greatly depends on the chosen preprocessing procedure. Specifically, the authors suggest that high-pass filtering the data above 1 or 2 Hz prior to ICA consistently leads to good results. Thus, in order to improve our artifact correction procedure, our next step was to go back to the continuous EEG data obtained after bad channel interpolation and bandpass-filtering it between 1 and 40 Hz. Afterwards, we again split this filtered data in epochs time-locked to stimulus onset. We then removed the epochs that were marked as bad when visually cleaning the 0.1 Hz high-pass filtered epochs. Following this we used the extended Infomax independent component analysis (ICA) algorithm implemented in MNE (Lee et al., 1999) on the 1-40 Hz bandpass-filtered, epoched data. The resulting ICA components were then used on the 0.1 Hz high-pass filtered data to remove eye movement artifacts. The final step in our preprocessing pipeline was to remove epochs based on the behavioral data: we removed 1) epochs where the RT deviated more than three standard deviations from the overall mean RT, and 2) epochs where the participants were incorrect, even though they had already seen that stimuli 3 times or more (assuming those represented attentional slips). The same pipeline was used to preprocess epochs time-locked to the response. Overall, the complete data cleaning procedure led to an exclusion ranging from 3.32% to 29.30% ($M = 17.34$, $SD = 5.21$) of the recorded epochs across participants.

Statistical Analysis

Behavioral Analysis

All statistical analyses of the behavioral data were performed using the 'lme4' package (Bates et al., 2007), which is embedded in the R software (Version 3.6.3). All

associated plots were created using the visualization libraries 'seaborn' and 'Matplotlib' (Hunter, 2007) in the Python programming language (Version 3.5.6).

For the behavioral results, two dependent variables are considered: RT and error rates. RT was first log scaled prior to statistical tests to increase normality. For each statistical test involving RT as the dependent variable, we built a linear mixed effects model. For errors, a generalized linear mixed effects model was utilized. All models contained a random intercept for subject. The independent variables were always included as fixed effects. Note that in plots the original, untransformed, data is displayed.

In order to elucidate the effects over the fast timescale, we analyzed how the recorded behavioral and neural metrics evolved within blocks. To this end, we analyzed dynamics as a function of stimulus number, i.e. the number of times a stimulus was seen by a subject. For example, when investigating how RT evolved over the fast timescale, we would build a linear mixed effects model, where the independent variables were stimulus number and condition, and the dependent variable RT.

The slow timescale represents changes between blocks. To capture this, we analyzed changes in our recorded data as a function of block number and condition.

EEG Data Analysis

Our analyses focus on the total power, but we also investigate evoked power. Evoked power refers to the phase-consistent part of the EEG signal (Hajihosseini & Holroyd, 2013), which can be accessed by computing the event-related potential (ERP) before spectral analysis.

Before exploring how alpha and theta power fluctuate across timescales, additional processing was done. The nature of this data preparation depended on several factors. First, the timescale (fast or slow) considered in the specific analysis influenced our data analysis.

Second, whether we looked at either total power or evoked power impacted our analysis choices. The third and final factor of our pipeline was the event (stimulus or response) to which the data was time-locked. We will now delineate how each of these decisions was implemented in our analyses.

To investigate how oscillatory alpha and theta power evolve for each timescale, we again used stimulus numbers and block numbers as independent variables for the fast and slow timescales, respectively. Specifically, we started our analysis of oscillatory power by comparing the power values for the end points on each timescale. For the fast timescale, we compared the recorded spectral power for stimulus 1 versus 8. For the slow timescale, the power in experimental block 1 was contrasted with the power in block 8. Thus, in order to equate the relevant power measures, we first split the recorded EEG data for each subject in two extreme parts for each timescale: data representing earlier timepoints (stimulus 1, or block 1) and data from late timepoints (stimulus 8, or block 8). Then, we computed the time-frequency representation (TFR) for each subset of the data on the same timescale using Morlet wavelets. The power estimates were computed for 15 frequency bands, log spaced between 4 Hz and 30 Hz. The number of cycles used for each frequency was equal to the frequency divided by two, to preserve the balance between temporal and frequency precision (Cohen, 2014). This procedure ultimately resulted in TFRs for each timepoint for each subject. To compare the resulting TFRs, a cluster-level statistical permutation test was used to account for multiple comparisons (Maris & Oostenveld, 2007). This permutation test was run with 1000 permutations, and a significance threshold of 5% was used.

For evoked power, the aforementioned permutation test was run on the TFRs of the ERPs of specific subsets. For the fast timescale this meant that we only included the data representing stimulus 1 and 8 in the permutation test. For each of these subsets the ERP was computed, which was then followed by the permutation test. A similar procedure was

followed for the data tracking the slow timescale, but there the subject-specific data was split in a part containing data of block 1, and a part containing the data of block 8.

Stimulus-locked epochs were 2500 ms long (-1000 to 1500 ms relative to stimulus onset). We analyzed the time period ranging from 0 to 1000 ms after stimulus onset for the fast timescale, and the period from 0 to 750 ms after stimulus onset for the slow timescale. The period from 1000 to 250 ms before stimulus onset served as the baseline. For response-locked data, we also used a pre-stimulus baseline period, but this one ranged from 1000 to 500 ms before stimulus onset. We shortened our baseline period because our analyzed time period was shorter: we only consider the data from 235 ms before response onset, to 305 ms after response. The response-locked time window was defined like this because otherwise some overlap might occur with the anticipation phase of the next trial (see Figure 1B), especially for the trials where subjects exhibited later responses.

Based on the results of the cluster-level statistical permutation test, time-frequency plots were created. The time-frequency plots represent the cluster statistics averaged over all channels. This resulted in a two-dimensional plot displaying the computed statistics as a function of time and frequency bands. To assess where power fluctuations originate from on the scalp, we also created topographical plots by subtracting block 1 (stimulus 1) from block 8 (stimulus 8).

In the time-frequency clustering algorithm, we only contrasted stimulus 1 or block 1 with stimulus 8 or block 8 respectively across both timescales. However, in order to further explore the data in between these extremes, we conducted follow-up analyses on the intermediate stimuli 2-7 and blocks 2-7 based on the clusters identified in the above analyses (thus avoiding "double dipping"; Kriegeskorte et al., 2009). To this end, we used the 'lme4' package (Bates et al., 2007), incorporated in the R software (Version 3.6.3). For example,

consider the hypothetical situation where a significant difference in theta power is found in some time window after stimulus onset in the stimulus-locked cluster test comparing stimuli 1 and 8. The first step in the follow-up analysis would be to compute the average theta power in that same time window across trials containing stimuli 2 to 7. Following this, we would build a linear mixed effects model with theta power as dependent variable, subject number as a random intercept, and stimulus number and condition as fixed effects. A similar procedure was used for analyzing effects over blocks 2 to 7.

Figure 2. RT and error rate following the fast and the slow timescale. RT (A) and error rate (B) as a function of stimulus number, respectively. These plots capture behavioral effects following the fast timescale. Note that the plot is based on data of the novel condition and the first block of the repeating condition. RT (C) and error rate (D) as a function of block number, respectively. Thus, these plots show behavioral changes on the slow timescale. A distinction is made based on condition, where 'novel' indicates that new stimuli were shown in that block, and 'repeating' means that stimuli were shown that returned throughout the experiment. The vertical lines represent bootstrapped 95% confidence intervals in all plots.

Code and data sharing

The Python and R scripts used for data processing, plotting and analyses are provided at

https://github.com/phuycke/alpha_theta_timescales. Our (un)processed data are available at

https://osf.io/2q5eh/.

Results

Behavioral Results

Focusing on the fast timescale, stimulus number had a main effect on both RT, $F(7,$

$5561) = 61.48$, $p < .001$, and error rate, $X^2(7, N = 24) = 122.38$, $p < .001$. Both RT and error

rate decreased as stimulus repetitions increased. How RT and error rate evolved as a function

of stimulus repetitions is visualized in Figure 2A and 2B, respectively. Note that stimulus

repetitions is defined by the number of times a stimulus was seen by a subject. Hence,

stimulus repetition 1 means that this is the first time that a stimulus was seen. This plot

(Figure 2A-B) was created using the data from the novel condition, and the first block of the

repeating condition. This selection was made because this subset contained information on

the first eight encounters with the experimental stimuli. Note that accuracy starts off slightly

below 50% because of a dependency between stimuli: If the participant has already seen 3 of

the 4 stimuli (e.g., two with left and one with right response), the response to the fourth

stimulus can be known with certainty (in the example, right response).

Next, we considered the same dependent variables on the slow timescale. Both block

number, $F(7, 10118) = 25.79$, $p < .001$, and condition (novel or repeating), $F(1, 10118) =$

542.87, $p < .001$, had a significant impact on RT. Additionally, the interaction effect between

block number and condition also proved significant, $F(7, 10118) = 14.50$, $p < .001$. A follow-

up pairwise t-test (Benjamini-Hochberg adjusted) showed that in the repeating condition,

block 1 was significantly different from all others (all $p < .001$). In the novel condition, none

of the blocks differed significantly (all $p > .05$). On error rate, block number had a significant impact, $X^2(7, N = 24) = 34.29, p < .001$. The main effect of condition on error rate proved to be non-significant, $X^2(1, N = 24) = 0.03, p = .871$. However, the interaction effect of block number and condition was significant, $X^2(7, N = 24) = 27.30, p < .001$. Specifically, using the same follow-up test, we concluded that in the repeating condition the error rate in block 1 was significantly higher than the error rate in the other blocks (all $p < .001$), but remained constant in the novel condition (all $p > .05$). How RT and error rate evolved following the slow timescale is shown in Figure 2C and 2D, respectively.

EEG Results

Fast timescale

In total power changes on the fast timescale (stimulus 8 minus stimulus 1), we found a significant cluster in the stimulus-locked data. Specifically, we identified a significant alpha cluster ($p = .001$) after cluster correction approximately between 700 and 850 ms after stimulus onset (Figure 3A). This cluster is localized in the posterior part of the scalp (Figure 3B). Follow-up analyses focusing on alpha power (stimulus 2 to 7) in this significant cluster indicated that alpha power (from here on abbreviated as alpha) significantly increased with stimulus number, $F(5, 7684) = 20.34, p < .001$. The main effect of condition also proved to be significant, $F(1, 7684) = 61.70, p < .001$. Finally, the interaction effect between stimulus number and condition was also significant, $F(5, 10118) = 3.40, p = .004$. How alpha evolves as a function of stimulus number and condition is visualized in Figure 3C.

Next, we analyzed stimulus-locked evoked power contrasting stimulus 1 vs. 8. We found a significant alpha cluster between 700 and 800 ms after stimulus onset ($p < .001$). Combined with the previous result, this indicates that the stimulus-locked alpha burst can be seen in both the total and the evoked power, occurring more than 700 ms after the stimulus

onset.

For the response-locked data, we again started by comparing the total power measures for stimulus 1 vs. stimulus 8, performing the same cluster-correction procedure. A significant alpha cluster was observed ($p = .002$) around 200 ms after response onset. No significant results were obtained when analyzing the response-locked, evoked power.

Slow timescale

For the slow timescale, we started by analyzing the total power fluctuations by contrasting block 1 with block 8 (block 1 minus block 8). The permutation test highlighted two significant theta clusters (4 – 8 Hz), one at stimulus onset and one around 400 ms after stimulus onset ($p = .04$) (Figure 4A). The cluster in the time window from 250 to 450 ms originate was observed over a frontal scalp region (Figure 4B). A follow-up test based on this cluster (blocks 2-7) using a linear mixed effects model showed that block number is a significant predictor of theta power, $F(5, 7651) = 2.59$, $p = .02$. In contrast, the main effect of condition was not significant, $F(1, 7561) = 1.12$, $p = .29$. The interaction effect between condition and block number, $F(5, 7651) = 2.32$, $p = .04$, proved significant (Figure 4D).

We also analyzed the contrast between block 1 and block 8 with a focus on stimulus-locked evoked power, response-locked total power and response-locked evoked power. None of these contrasts yielded significant results.

Alpha-theta Dissociation

In order to investigate whether the fluctuations in alpha and theta power are specific to the fast and slow timescales respectively, we additionally tested the interaction between stimulus (block) number and frequency band on each time scale separately. On the fast timescale, the interaction between stimulus number and frequency band (theta, alpha) indeed proved significant, $F(7, 20259) = 22.33$, $p < .001$. Similarly, for the slow timescale the

interaction effect between block and frequency band (theta, alpha) also proved to be significant, $F(7, 20259) = 2.88$, $p = .005$.

We next further investigated the interactions between time point (stimulus or block) and condition (novel or repeating). Indeed, the interactions observed above may originate from the fact that each time point (stimulus repetition or block) was coded as a separate factor in the analysis. Thus, to further clarify these results, we computed the subject-specific slope of the least-squares regression lines created when regressing alpha (or theta) power as a function of stimulus (or block) number. Then, we compared the subject-specific slopes for the novel condition with the slopes for the repeating condition using a paired samples t-test. This test was performed for each combination between frequency band and timescale. We found only one significant difference in subject-specific slopes between conditions: when considering alpha as a function of stimulus number we found that the subject-specific slopes for the novel condition ($M = 0.03$, $SD = 0.02$) compared to the repeating condition ($M = 0.01$, $SD = 0.02$) differed significantly, $t(24) = 3.24$, $p = .004$. All other post-hoc tests yielded non-significant results (all $p > .1$). The results of this post-hoc analysis confirm that only the interaction effect between stimulus number and condition on alpha survives after further statistical testing.

As another follow-up test we investigated whether alpha ($8 – 12$ Hz) significantly changed over the slow timescale. To investigate this, we considered the time period from 700 to 850 ms after stimulus onset, since significant alpha effects were found in this window on the fast timescale. This analysis suggested that alpha did not change as a function of block number, $F(7, 10118) = 1.57$, $p = .14$. The main effect of condition proved significant, $F(1, 10118) = 69.06$, $p < .001$. Finally, the interaction between block number and condition was not significant, $F(7, 10118) = 1.73$, $p = .10$. How alpha behaved on the slow timescale is depicted in Figure 3D. Next, we investigated whether theta frequency effects ($4 – 8$ Hz) could

be observed on the fast timescale. To this end, we analyzed theta power in the time window

ranging from 250 to 400 ms after stimulus onset. This time window was chosen based on our

analyses for the slow timescale, which are described in the previous section. We found a

significant main effect of stimulus number on theta power, $F(7, 10118) = 2.04$, $p = .05$

(Figure 4C). Both the main effect of condition, $F(1, 10118) = 1.12$, $p = .29$, and the

interaction between stimulus number and condition, $F(7, 10118) = 0.89$, $p = .51$, were not

significant. Thus, we argue that on the fast time scale, alpha changed; whereas on the slow

time scale, theta changed across time.

Correlation between neural and behavioral metrics

In a final analysis step, we investigated whether alpha and theta can act as predictors

for RT. To explore this, we created a linear mixed effects model with RT as dependent

variable; subject as random intercept; theta and alpha as random slopes; theta, alpha, stimulus

number and block number as fixed effects; and the interaction effects between stimulus

number and condition and between block number and condition also as fixed effects. Hence,

we included all the experimental variables that could account for RT, and we investigated

whether theta and alpha metrics had a significant contribution in explaining RT. The fitting of

the complete model was followed by model selection using the AIC as criterion. The final

model only contained theta, suggesting that theta, but not alpha, is a significant predictor of

RT, $F(1, 22) = 9.08$, $p = .007$, over and above the experimental variables.

Figure 3. Total alpha power across timescales. (A) This time-frequency plot depicts the cluster statistics which were averaged over all channels, resulting in a two-dimensional plot displaying the computed statistics as a function of time and frequency band. We delineated a cluster of high-value statistics around 800 ms after stimulus onset (timepoint 0). This delineated area was used to conduct our follow-up analyses (see Figure 3C and Figure 4C). (B) The topography of the delineated area in panel A. This figure depicts the topography of the difference of stimulus 8 minus stimulus 1 in the alpha frequency band, averaged across time between 700 and 850 ms after stimulus onset. (C) Alpha (8 – 12 Hz) between 700 and 850 ms after stimulus onset averaged across time and trials as a function of stimulus number. The analyzed time-interval was defined based on panel A. (D) Alpha averaged across time and electrodes between 250 and 400 ms after stimulus onset as a function of block number and condition. We analyzed this time period based on the results depicted in Figure 4A. Note that in both panels C and D averages are represented by dots, while bootstrapped 95%

confidence intervals are represented by vertical lines. The straight lines represent linear

regression lines.

Figure 4. Total theta power across timescales. (A) Channel-averaged time-frequency

representation of power. Statistically significant clusters are circled in black. The cluster

between 250 and 400 ms was used to conduct our follow-up analyses (see Figure 3D and

Figure 4D). (B) The topography of the cluster in panel A. This figure depicts the topography

of the contrast block 8 minus block 1 in the theta frequency band, averaged across time in the

period between 250 and 400 ms after stimulus onset. This contrast yielded an anterior

positive difference and a posterior negative difference. (C) Theta (4 – 8 Hz) between 700 and

850 ms after stimulus onset averaged across time and subjects as a function of stimulus

number. We analyzed this time period based on the results depicted in Figure 3A. (D) Theta

averaged across time and electrodes between 250 and 400 ms after stimulus onset as a

function of block number and condition. The analyzed time-interval was defined based on

panel A. Note that in both panels C and D averages are represented by dots, while

bootstrapped 95% confidence intervals are represented by vertical lines. The straight lines

represent linear regression lines.

Discussion

Previous research indicates that alpha and theta frequencies play instrumental roles in cognitive control, but it remains unclear how they unfold during the implementation of cognitive control. In this study, participants performed a procedural learning task in which stimulus-action associations had to be learned. These associations were block-specific and could be either novel or repeating. Learning took place on two separate timescales: a fast (within blocks) timescale, and a slow (between blocks) timescale. Our results indicate that cognitive control is implemented across both a fast and a slow timescale. On the fast timescale we observed that alpha power significantly increases over time, and more strongly so in the novel than in the repeated condition (see Figure 3C). In contrast, on the slow timescale theta power decreases over time (see Figure 4D), but not differently across conditions. Alpha did not significantly change in the slow timescale, whereas theta significantly decreases on the fast timescale. Follow-up tests show significant interactions between stimulus number and frequency band on the fast time scale, indicating that alpha increased faster than theta on the fast time scale; and between block and frequency band on the slow timescale; showing that theta decreased faster than alpha on the slow time scale. Taken together, we argue that alpha increases are specific to the fast timescale, whereas a decrease in theta power is specific to the slow timescale.

We interpret the results obtained at the fast (within a block) timescale in the 'gating by inhibition' framework proposed by Jensen and Mazaheri (2010), according to which alpha oscillations suppress task-irrelevant brain areas, and thereby indirectly 'gate' task-relevant brain areas. Evidence for this hypothesis comes from studies in memory (Grabner et al., 2003; Sauseng et al., 2005), face processing (Jokisch & Jensen, 2007), visual detection (Hanslmayr et al., 2005), and cognitive control dynamics (Carp & Compton, 2009; Compton et al., 2011; Janssens et al., 2018). For example, Carp and Compton (2009) and Compton et

al. (2011) observed a significantly lower alpha power in the inter-trial interval of a Stroop

task in error relative to correct trials. Similar results were obtained when contrasting high-

conflict with low-conflict trials. From the 'gating by inhibition'-framework, alpha may reflect

inhibition of incorrect stimulus-action mappings. In our task, higher alpha would reflect

stronger inhibition of inappropriate stimulus-action mappings, which in turn leads to better

task performance. This interpretation is in line with the significant interaction between

stimulus number and condition, which shows that the effect of stimulus number on alpha is

different for the novel and the repeating condition. This difference might be due to the

stronger inhibition needed in the novel condition to suppress irrelevant mappings, whereas

this inhibition is less needed in the repeating conditions since learning has already taken

place.

We explain our slow (i.e., across blocks) time scale results in the theta band as

reflecting the role of FM theta in the exertion of cognitive control. In their review paper,

Cavanagh and Frank (2014) observed that cognitive control tasks typically showed a neural

signature in the theta band approximately 250 to 500 ms after stimulus onset. We

hypothesized that this theta band activation originates from the mid-frontal part of the scalp.

In these results, FM theta is interpreted as a signal that marks the need for cognitive control.

Other studies focused on the role that FM theta might play in the coordination of cognitive

control. For instance, it is thought that (FM) theta synchronizes posterior processing areas

depending on the task at hand (Fries, 2015). Several authors (Fries, 2015; Verbeke &

Verguts, 2019; Voloh et al., 2015; Womelsdorf et al., 2006) suggested that communication

between neural areas is achieved using gamma-band synchronization, and the role of theta is

to reset gamma such that the appropriate task-relevant areas may subsequently phase-lock in

the gamma band. Theta-band phase synchrony can be observed between central areas and

more distant brain areas. For example, after errors, enhanced synchrony is observed between

the MFC and the occipital cortex (Cohen, 2009); and after incongruent stimuli during a

Stroop task (Hanslmayr et al., 2008), synchrony is increased between FM and frontolateral

regions. Consistently, we observed decreased theta power when comparing block 8 minus

block 1, in both the novel and the repeating condition. Since theta decreases equally in both

novel and repeating conditions over a slow timescale, our results suggest that theta is

implicated in a general cognitive process, such as mapping stimuli on available actions. We

hypothesize that the decrease in theta reflects that this mapping process enfolds more

efficiently when a subject is more experienced with the task, irrespective of the stimuli or

actions that are currently being mapped. In short, theta reflects a general (i.e., not stimulus- or

action-specific) learning process that takes place on a slow timescale, and becomes optimized

with more training. Finally, the correlation of theta with RT suggests that trials where this

mapping process is implemented more strongly, allow more efficient processing and thus

faster RT.

One might suggest that the significant theta cluster found approximately 250 to 400

ms after stimulus onset simply reflects a P300 response, an event-related potential

hypothesized to be indicative of mental workload (Miller et al., 2011; Wickens et al., 1977).

Our analyses however point out that a significant theta cluster was only found when

comparing total power between blocks 1 and 8, and not when analyzing evoked power.

Therefore, we argue that it is unlikely that our significant theta cluster is a manifestation of a

P300 response to our experimental stimuli.

In principle, the increase in alpha might reflect an increasingly fast response process

as the block develops. However, this interpretation is unlikely. Specifically, RT decreases as

a function of stimulus number from approximately 700 ms (stimulus 1) to approximately 500

ms (stimulus 8; see Figure 2A). However, the alpha effects are localized in a later time

window situated around 800 ms (see Figure 3A), suggesting that these alpha effects occur

well after response onset. Similarly, it could be hypothesized that the theta effects are response-related. However, this is also unlikely because the theta effects occur well before the mean RT.

Alternatively, one might argue that our results reflect cognitive fatigue resulting from prolonged testing on a simple task. A study by Wang et al. (2016) suggests that this interpretation is improbable. In this EEG study, participants were asked to perform a Stroop task for a duration of 160 minutes with the goal to elicit cognitive exhaustion. Wang et al. (2016) report an anterior ERP that seems to be related to compensatory mechanisms used to counteract the effects of cognitive fatigue. Specifically, the authors note an inverted U-shaped relationship between the time on task and the amplitude of this frontal ERP. The signal peaks when the subject attempts to compensate cognitive fatigue, and decreases both during baseline, and when the subject is too tired to engage in compensation. Crucially, the latter was observed in the time window ranging from 120 minutes to 160 minutes after the start of the experiment. Thus, it seems unlikely that our results can be explained by cognitive fatigue, since the neural markers of fatigue occur after a time-on-task greatly exceeding our time-on-task of 60 minutes.

Our results can also be linked to the literature on automatization and extensive training. For instance, in the fMRI study by Ruge and Wolfensteller (2010) on instruction-based learning, the authors noted that brain activation in the central gyri and the striatum increased on a fast timescale, suggesting that these areas are relevant for automatization. On the other hand, activation in the lateral PFC and the intraparietal sulcus (IPS) decreases with more experience with a stimulus. Given the posterior location of IPS, the fact that increased alpha correlates with decreased BOLD (Ritter et al., 2009), and our hypothesized inhibitory role of alpha, as well as our finding that alpha increases with stimulus experience, we hypothesize that the measured increase in posterior alpha may signify the active inhibition of

the IPS. Since the spatial resolution of EEG is too low to verify this theory in an anatomically-specific fashion, a future study might investigate this further using a method that has both high spatial and high temporal resolution (such as intracranial EEG). We furthermore show that theta power changes on a slower timescale, suggesting that automatization might occur on distinct timescales signified by distinct neural markers.

Work on instruction implementation (Hartstra et al., 2011; Ruge & Wolfensteller, 2010) shows that frontoparietal activation is relatively quickly replaced by striatal activation after training on a stimulus-action mapping. However, the automatization literature (e.g., the 'Subcortical Pathways Enable Expertise Development' (SPEED; Ashby et al., 2007) model) suggests that there is yet another automatization stage after striatal involvement. In particular, the SPEED model postulates the existence of at least two distinct neural pathways, both running from the sensory association cortex to the premotor cortex. A subcortical pathway that learns fast, runs via the basal ganglia, making it eligible for dopamine-driven learning. The other, purely cortical and slower-learning pathway learns via Hebbian learning, receiving training signals from the subcortical pathway, so that with more experience the slow learning pathway takes over. We hypothesize that our data document the transfer from a frontoparietal based representation to one based on basal ganglia. According to the SPEED model, there should thus be yet a third stage, where basal ganglia are no longer involved. To test the spectral signatures of the latter stage, future research will have to extend our paradigm to one where participants are trained over consecutive days. This would provide extra information on alpha and theta dynamics, such as at which point in time the power measures stabilize, and whether learning and automatization enfold on other timescales than the ones reported in the current study.

To sum up, we found that theta and alpha frequency both play a role during stimulus-action learning, but their impact differs across timescales. Alpha band dynamics are specific

to the fast timescale, whereas theta changes are observed in the slow timescale, and similar for novel and repeated stimuli. Our data suggest that automatization is not a unitary process but consists of at least two dissociable steps. Which neural structures are involved, how these neural processes evolve with more training, and the nature of their computational role, remain to be investigated in future research.

Acknowledgements

The authors declare no conflict of interest. This work was supported by the EOS grant number G0F3818N by Research Foundation Flanders (to TV and CNB). PV was supported by grant 1102519N from Research Foundation Flanders.

References

Anguera, J. A., Boccanfuso, J., Rintoul, J. L., Al-Hashimi, O., Faraji, F., Janowich, J., Kong, E., Larraburo, Y., Rolle, C., Johnston, E., & Gazzaley, A. (2013). Video game training enhances cognitive control in older adults. *Nature*, *501*(7465), 97–101. https://doi.org/10.1038/nature12486

Ashby, F. G., Ennis, J. M., & Spiering, B. J. (2007). A neurobiological theory of automaticity in perceptual categorization. *Psychological Review*, *114*(3), 632–656. https://doi.org/10.1037/0033-295X.114.3.632

Bates, D., Sarkar, D., Bates, M. D., & Matrix, L. (2007). The lme4 package. *R Package Version*, *2*(1), 74.

Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. D. (2001). Conflict monitoring and cognitive control. *Psychological Review*, *108*(3), 624–652. https://doi.org/10.1037/0033-295X.108.3.624

Buzsáki, G. (2002). Theta Oscillations in the Hippocampus. *Neuron*, *33*(3), 325–340. https://doi.org/10.1016/S0896-6273(02)00586-X

Carp, J., & Compton, R. J. (2009). Alpha power is influenced by performance errors.

Psychophysiology, *46*(2), 336–343. https://doi.org/10.1111/j.1469-8986.2008.00773.x

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control.

Trends in Cognitive Sciences, *18*(8), 414–421.

https://doi.org/10.1016/j.tics.2014.04.012

Cohen, M. X. (2009). Unconscious errors enhance prefrontal-occipital oscillatory synchrony.

Frontiers in Human Neuroscience, *3*. https://doi.org/10.3389/neuro.09.054.2009

Cohen, M. X. (2014). *Analyzing neural time series data: Theory and practice*. MIT press.

Compton, R. J., Arnstein, D., Freedman, G., Dainer-Best, J., & Liss, A. (2011). Cognitive

control in the intertrial interval: Evidence from EEG alpha power: Cognitive control

in the intertrial interval. *Psychophysiology*, *48*(5), 583–590.

https://doi.org/10.1111/j.1469-8986.2010.01124.x

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *Journal of

Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Enriquez-Geppert, S., Huster, R. J., Scharfenort, R., Mokom, Z. N., Zimmermann, J., &

Herrmann, C. S. (2014). Modulation of frontal-midline theta by neurofeedback.

Biological Psychology, *95*, 59–69. https://doi.org/10.1016/j.biopsycho.2013.02.019

Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. *Neuron*, *88*(1),

220–235. https://doi.org/10.1016/j.neuron.2015.09.034

Grabner, R. H., Stern, E., & Neubauer, A. C. (2003). When intelligence loses its impact:

Neural efficiency during reasoning in a familiar area. *International Journal of

Psychophysiology*, *49*(2), 89–98. https://doi.org/10.1016/S0167-8760(03)00095-3

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C.,

Parkkonen, L., & Hämäläinen, M. S. (2014). MNE software for processing MEG and

EEG data. *NeuroImage*, *86*, 446–460.

https://doi.org/10.1016/j.neuroimage.2013.10.027

Hajihosseini, A., & Holroyd, C. B. (2013). Frontal midline theta and N200 amplitude reflect complementary information about expectancy and outcome evaluation: Frontal theta and N200 provide distinct information. *Psychophysiology*, *50*(6), 550–562. https://doi.org/10.1111/psyp.12040

Händel, B. F., Haarmeier, T., & Jensen, O. (2011). Alpha Oscillations Correlate with the Successful Inhibition of Unattended Stimuli. *Journal of Cognitive Neuroscience*, *23*(9), 2494–2502. https://doi.org/10.1162/jocn.2010.21557

Hanslmayr, S., Klimesch, W., Sauseng, P., Gruber, W., Doppelmayr, M., Freunberger, R., & Pecherstorfer, T. (2005). Visual discrimination performance is related to decreased alpha amplitude but increased phase locking. *Neuroscience Letters*, *375*(1), 64–68. https://doi.org/10.1016/j.neulet.2004.10.092

Hanslmayr, S., Pastötter, B., Bäuml, K.-H., Gruber, S., Wimber, M., & Klimesch, W. (2008). The Electrophysiological Dynamics of Interference during the Stroop Task. *Journal of Cognitive Neuroscience*, *20*(2), 215–225. https://doi.org/10.1162/jocn.2008.20020

Hartstra, E., Kühn, S., Verguts, T., & Brass, M. (2011). The implementation of verbal instructions: An fMRI study. *Human Brain Mapping*, *32*(11), 1811–1824. https://doi.org/10.1002/hbm.21152

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science & Engineering*, *9*(3), 90–95. https://doi.org/10.1109/MCSE.2007.55

Janssens, C., De Loof, E., Boehler, C. N., Pourtois, G., & Verguts, T. (2018). Occipital alpha power reveals fast attentional inhibition of incongruent distractors. *Psychophysiology*, *55*(3), e13011. https://doi.org/10.1111/psyp.13011

Jasper, H. H. (1958). The ten-twenty electrode system of the International Federation. *Electroencephalogr. Clin. Neurophysiol.*, *10*, 370–375.

Jensen, O., & Mazaheri, A. (2010). Shaping Functional Architecture by Oscillatory Alpha Activity: Gating by Inhibition. *Frontiers in Human Neuroscience*, *4*. https://doi.org/10.3389/fnhum.2010.00186

Jokisch, D., & Jensen, O. (2007). Modulation of Gamma and Alpha Activity during a Working Memory Task Engaging the Dorsal or Ventral Stream. *Journal of Neuroscience*, *27*(12), 3244–3251. https://doi.org/10.1523/JNEUROSCI.5399-06.2007

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: A review and analysis. *Brain Research Reviews*, *29*(2–3), 169–195. https://doi.org/10.1016/S0165-0173(98)00056-3

Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S. F., & Baker, C. I. (2009). Circular analysis in systems neuroscience: The dangers of double dipping. *Nature Neuroscience*, *12*(5), 535–540. https://doi.org/10.1038/nn.2303

Lee, T.-W., Girolami, M., & Sejnowski, T. J. (1999). Independent Component Analysis Using an Extended Infomax Algorithm for Mixed Subgaussian and Supergaussian Sources. *Neural Computation*, *11*(2), 417–441. https://doi.org/10.1162/089976699300016719

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Mazaheri, A., Nieuwenhuis, I. L. C., van Dijk, H., & Jensen, O. (2009). Prestimulus alpha and mu activity predicts failure to inhibit motor responses. *Human Brain Mapping*, *30*(6), 1791–1800. https://doi.org/10.1002/hbm.20763

Miller, M. W., Rietschel, J. C., McDonald, C. G., & Hatfield, B. D. (2011). A novel approach

to the physiological measurement of mental workload. *International Journal of

Psychophysiology*, *80*(1), 75–78. https://doi.org/10.1016/j.ijpsycho.2011.02.003

Narayanan, N. S., Cavanagh, J. F., Frank, M. J., & Laubach, M. (2013). Common medial

frontal mechanisms of adaptive control in humans and rodents. *Nature Neuroscience*,

16(12), 1888–1895. https://doi.org/10.1038/nn.3549

Nigbur, R., Cohen, M. X., Ridderinkhof, K. R., & Stürmer, B. (2012). Theta Dynamics

Reveal Domain-specific Control over Stimulus and Response Conflict. *Journal of

Cognitive Neuroscience*, *24*(5), 1264–1274. https://doi.org/10.1162/jocn_a_00128

O'Keefe, J. (1993). Hippocampus, theta, and spatial memory. *Current Opinion in

Neurobiology*, *3*(6), 917–924. https://doi.org/10.1016/0959-4388(93)90163-S

Peirce, J. W. (2007). PsychoPy—Psychophysics software in Python. *Journal of Neuroscience

Methods*, *162*(1–2), 8–13. https://doi.org/10.1016/j.jneumeth.2006.11.017

Pfurtscheller, G. (2003). Induced Oscillations in the Alpha Band: Functional Meaning.

Epilepsia, *44*(s12), 2–8. https://doi.org/10.1111/j.0013-9580.2003.12001.x

Phillips, J. M., Vinck, M., Everling, S., & Womelsdorf, T. (2014). A Long-Range Fronto-

Parietal 5- to 10-Hz Network Predicts "Top-Down" Controlled Guidance in a Task-

Switch Paradigm. *Cerebral Cortex*, *24*(8), 1996–2008.

https://doi.org/10.1093/cercor/bht050

Ridderinkhof, K. R. (2004). The Role of the Medial Frontal Cortex in Cognitive Control.

Science, *306*(5695), 443–447. https://doi.org/10.1126/science.1100301

Ritter, P., Moosmann, M., & Villringer, A. (2009). Rolandic alpha and beta EEG rhythms'

strengths are inversely related to fMRI-BOLD signal in primary somatosensory and

motor cortex. *Human Brain Mapping*, *30*(4), 1168–1187.

https://doi.org/10.1002/hbm.20585

Ruge, H., & Wolfensteller, U. (2010). Rapid Formation of Pragmatic Rule Representations in

the Human Brain during Instruction-Based Learning. *Cerebral Cortex*, *20*(7), 1656–

1667. https://doi.org/10.1093/cercor/bhp228

Sauseng, P., Klimesch, W., Doppelmayr, M., Pecherstorfer, T., Freunberger, R., &

Hanslmayr, S. (2005). EEG alpha synchronization and functional coupling during top-

down processing in a working memory task. *Human Brain Mapping*, *26*(2), 148–155.

https://doi.org/10.1002/hbm.20150

Sauseng, P., Tschentscher, N., & Biel, A. L. (2019). Be Prepared: Tune to FM-Theta for

Cognitive Control. *Trends in Neurosciences*, *42*(5), 307–309.

https://doi.org/10.1016/j.tins.2019.02.006

Thut, G. (2006). Band Electroencephalographic Activity over Occipital Cortex Indexes

Visuospatial Attention Bias and Predicts Visual Target Detection. *Journal of

Neuroscience*, *26*(37), 9494–9502. https://doi.org/10.1523/JNEUROSCI.0875-

06.2006

van de Vijver, I., Ridderinkhof, K. R., & Cohen, M. X. (2011). Frontal Oscillatory Dynamics

Predict Feedback Learning and Action Adjustment. *Journal of Cognitive

Neuroscience*, *23*(12), 4106–4121. https://doi.org/10.1162/jocn_a_00110

Verbeke, P., & Verguts, T. (2019). Learning to synchronize: How biological agents can

couple neural task modules for dealing with the stability-plasticity dilemma. *PLOS

Computational Biology*, *15*(8), e1006604.

https://doi.org/10.1371/journal.pcbi.1006604

Vertes, R. P., & Kocsis, B. (1997). Brainstem-diencephalo-septohippocampal systems

controlling the theta rhythm of the hippocampus. *Neuroscience*, *4*(81), 893–926.

Voloh, B., Valiante, T. A., Everling, S., & Womelsdorf, T. (2015). Theta–gamma

coordination between anterior cingulate and prefrontal cortex indexes correct

attention shifts. *Proceedings of the National Academy of Sciences*, *112*(27), 8457–

8462. https://doi.org/10.1073/pnas.1500438112

Wang, C., Trongnetrpunya, A., Samuel, I. B. H., Ding, M., & Kluger, B. M. (2016).

Compensatory Neural Activity in Response to Cognitive Fatigue. *The Journal of*

Neuroscience, *36*(14), 3919–3924. https://doi.org/10.1523/JNEUROSCI.3652-

15.2016

Wickens, C. D., Isreal, J., & Donchin, E. (1977). The Event Related Cortical Potential as an

Index of Task Workload. *Proceedings of the Human Factors Society Annual Meeting*,

21(4), 282–286. https://doi.org/10.1177/107118137702100404

Winkler, I., Debener, S., Muller, K.-R., & Tangermann, M. (2015). On the influence of high-

pass filtering on ICA-based artifact reduction in EEG-ERP. *2015 37th Annual*

International Conference of the IEEE Engineering in Medicine and Biology Society

(EMBC), 4101–4105. https://doi.org/10.1109/EMBC.2015.7319296

Womelsdorf, T., Fries, P., Mitra, P. P., & Desimone, R. (2006). Gamma-band

synchronization in visual cortex predicts speed of change detection. *Nature*,

439(7077), 733–736. https://doi.org/10.1038/nature04258

A

B

	3000 ms
Fixation cross 1000 ... 2000 ms Jittered	
Stimulus presentation 1500 ms Response window 1500 ms	
...	Anticipation 500 ... 750 ms Jittered
Correct answord	Feedback 2000 - (750 ... 500) ms Jittered
	Relaxation 3000 - (2000 ... 1000) ms Jittered

2000 ms

Fast timescale

A RT (ms)

B Error (in %)

Stimulus repetitions

Slow timescale

C RT (ms)

D Error (in %)

Block number

Novel
Repeating