MECOM permits pancreatic acinar cell dedifferentiation avoiding cell death

under stress conditions

Running title: MECOM permits acinar cell dedifferentiation

Backx Elyne [1], Wauters Elke [2], Baldan Jonathan[2], Van Bulck Mathias[1], Michiels Ellis[1],

Heremans Yves[1,3], De Paep Dieter Luc [4], Bouwens Luc [2], Kurokawa Mineo [5], Goyama

Susumu [5], Jacquemin Patrick [6], Houbracken Isabelle[1,2], Rooman Ilse[1]

[1] Laboratory of Medical and Molecular Oncology, Oncology Research Centre, Vrije Universiteit Brussel,

Brussels, Belgium

[2] Cell Differentiation Laboratory, Vrije Universiteit Brussel, Brussels, Belgium

[3] Laboratory of Beta Cell Neogenesis, Vrije Universiteit Brussel, Brussels, Belgium

[4] Beta Cell Bank, Diabetes Research Center, UZ Brussel, Brussels, Belgium

[5] Department of Hematology and Oncology, Graduate School of Medicine, The University of

Tokyo, Tokyo, Japan

[6] Laboratory of Liver and Pancreatic Differentiation, De Duve Institute, Université Catholique de Louvain,

Brussels, Belgium

Corresponding author: Ilse Rooman, irooman@vub.be

ABSTRACT

Maintenance of the pancreatic acinar cell phenotype suppresses tumor formation. Hence, repetitive acute or chronic pancreatitis, stress conditions in which the acinar cells dedifferentiate, predispose for cancer formation in the pancreas. Dedifferentiated acinar cells acquire a large panel of duct cell specific markers. However, it remains unclear to what extent dedifferentiated acini differ from native duct cells and which genes are uniquely regulating acinar cell dedifferentiation. Moreover, most studies have been performed in mouse since the availability of human cells is scarce.

Here, we applied a non-genetic lineage tracing method in our culture model of human pancreatic exocrine cells that allowed cell-type specific gene expression profiling by RNA sequencing. Subsequent to this discovery analysis, one transcription factor that was unique for dedifferentiated acinar cells was functionally characterized using *in vitro* and *in vivo* genetic loss-of-function experimental models.

RNA sequencing analysis showed that human dedifferentiated acinar cells expressed genes in 'Pathways of cancer' with prominence of the transcription factor *MECOM (EVI-1)* that was absent from duct cells. During mouse embryonic development, pre-acinar cells transiently expressed MECOM and MECOM was re-expressed in experimental *in vivo* models of acute and chronic pancreatitis *in vivo,* conditions in which acinar cells dedifferentiate. MECOM expression correlated with and was directly regulated by SOX9. MECOM loss-of-function in mouse acinar cells *in vitro* and *in vivo* impaired cell adhesion resulting in more prominent acinar cell death and suppressed acinar cell dedifferentiation by limiting ERK signaling.

In conclusion, we transcriptionally profiled the two major human pancreatic exocrine cell types, acinar and duct cells, during experimental stress conditions. We provide insights that in dedifferentiated acinar cells, cancer pathways are upregulated in which

46 MECOM is a critical regulator that suppresses acinar cell death by permitting cellular

47 dedifferentiation.

49 INTRODUCTION

Few cancers have such poor survival rates as pancreatic ductal adenocarcinoma

(PDAC) (1). Understanding the molecular mechanisms that are at play in early stage

cancer formation is important to improve disease outcomes, but clinical samples of

early disease are rare. Hence, genetically modified mouse models are pivotal to

provide us insight into the contribution of the different pancreatic cell types to tumor

formation (2-4). Accumulating evidence points to acinar cells as cells of origin for

pancreatic tumor development (5-10). In order to become susceptible to

tumorigenesis, acinar cells have to lose their differentiated phenotype to some extent.

Consequently, tumorigenesis. is restrained by mechanisms that control acinar cell

differentiation (5-10). Loss of differentiation can result from repetitive acute or chronic

pancreatitis whereby dedifferentiated acinar cells acquire a panel of markers that are

typical for (embryonic) duct cells (11-13). Notably, duct cells can also give rise to

tumors but likely through on other mechanisms (14-16).

To date, knowledge is lacking to discriminate the mechanisms that govern the duct-

like dedifferentiated acinar cell phenotype versus the 'native' duct cell phenotype, the

latter being more resistant to stress- or inflammation-associated tumorigenesis.

Previously, we had established *in vitro* experimental models that recapitulate a human

(12) and rodent (13) acinar dedifferentiation. Specifically, upon suspension culture

acinar cells lose their acinar-specific markers while gaining (embryonic) duct-like

features, thereby mimicking chronic pancreatitis(12, 13).Here, we investigated the

differential gene expression in adult human dedifferentiated acinar cells compared to

native duct cells from the same cell preparations. The transcription factor *MECOM/EVI-*

1 was found to be uniquely re-expressed in dedifferentiated acinar cells since

embryonic development.

Although *MECOM* transcription factor is a known oncogene with a role in apoptosis

(17-19) and pancreatic tumor formation (20, 21), its function in acinar cell

(de)differentiation is still unknown. We analyzed its role in the context of acinar cell

dedifferentiation through caerulein-induced pancreatitis modeling in transgenic *Evi-1*

knock-out mice. We provide mechanistic insights that mouse acinar cells are

dependent on MECOM expression to prevent cell death by allowing them acquisition

of the dedifferentiated cell state.

MATERIALS AND METHODS

Human donor material and cell culture

Human pancreata from deceased donors were processed by the Beta Cell Bank of

the Centre for Beta Cell Therapy in Diabetes (Brussels, Belgium), affiliated to the

Eurotransplant Foundation (Leiden, The Netherlands). Consent for the use of

residual donor material for research was obtained according to the legislation in

country of organ procurement. This project was approved by the Medical Ethical

Committee of UZ Brussel - Vrije Universiteit Brussel (B.U.N. 143201732260).

Human exocrine cells were cultured as previously described (11, 12).

Mouse strains and experiments

ElaCreERT mice (22) ElaCreERT;Sox9$^{f/f}$ (23) (kindly provided by P. Jacquemin (UCL,

Belgium) originally from D. Stoffers (UPenn, USA)) were crossed with EVI-1exon4 loxP

mice (*Mecom*$^{f/f}$) (24) (kindly provided by M. Kurokawa, UPenn), resulting in

ElaCreERT;*Mecom*^(f/f) mice (Mecom KO). ElaCreERT;*Sox9*^(f/f) (23) were kindly provided

by P. Jacquemin (UCL, Belgium). Six to eight-weeks-old ElaCreERT;*Mecom*^(f/f) and

ElaCreERT;Sox9^(f/f) mice received tamoxifen and (Z)-4-Hydroxytamoxifen (all from

Sigma-Aldrich, St-Louis, MO, USA) and were treated with 1,25mg/kg body weight

caerulein (Invitrogen Waltham, Massachusetts, USA) according to (23). Acinar cell

isolation was adapted from (13). Embryonic mouse tissue was isolated from

ElaCreERT mice at E16.5 timepoint of embryonic development. All animal experiments

were approved by the Ethical Committee for Animal Testing at the Vrije Universiteit

Brussel (#17-637-1).

Fluorescent activated cell sorting (FACS)

FITC-conjugated UEA-1 (*Ulex Europaeus* Agglutinin-1, Sigma-Aldrich) lectin labeling

of acinar cells within the human exocrine cell fraction was performed as in (12). At day

4 of suspension culture, cell clusters were dissociated following the protocol of (11).

Analysis and cell sorting were performed on a BD FACSAria (BD Biosciences, Franklin

Lakes, NJ, USA). Viable, single cells were gated based on forward and side scatter.

More details on experimental design and statistics can be found in supplementary

materials and methods.

Results

MECOM is a cancer-related transcription factor uniquely expressed in human

dedifferentiated acinar cells

Given the difference in propensity of dedifferentiated acinar cells and duct cells for

pancreatitis-associated tumor development (5), we were specifically interested in the

differential gene expression of these two populations. Using the acinar-specific UEA-

1 lectin in combination with the duct cell marker CA19.9, we FACS-purified both cell

populations (11, 12) (Figure 1a,b). We confirmed the mutually exclusive expression

pattern of CD142, a progenitor marker (11) in the UEA1+ dedifferentiated acinar cells,

and of CA19.9 in the UEA1- duct cells (Figure 1b). We validated the different

expression levels of the acinar cell markers MIST-1 and chymotrypsin, the progenitor

marker CD142 and the duct-cell marker CK19 in the original mixed exocrine cell

fraction and in the purified dedifferentiated acinar cells (UEA1+/CA19.9-) and native

duct cells (UEA1-/CA19.9+) (Supplementary Figure 1). Comparing dedifferentiated

acinar cells with native duct cells by RNAseq analysis highlighted 7953 differentially

expressed genes (Adj. p<0,01). Of these, 1219 genes were expressed higher in

dedifferentiated acinar cells (log fold change (FC) of ≤-2; Adj. p<0,01) (Supplementary

Table 1) and were enriched for the KEGG pathways (Adj. p<0.01): 'Protein digestion

and absorption' and 'Pancreatic secretion', including genes such as carboxypeptidase

A and amylase, pointing to gene expression reminiscent of the abundant acinar cell

transcriptome (25). In addition, 'Pathways in cancer' belonged to the top enriched

pathways such as those of the WNT signaling pathway, which is well-known for its

function in pancreatic tumorigenesis. Other enriched pathways included 'PI3K-AKT

signaling', which is involved in acinar cell dedifferentiation and pancreatic

tumorigenesis (26-28) and 'Complement and coagulation cascades' encompassing

genes belonging to the tissue damage pathway (Figure 1c) and that code for F8,

fibrinogen and F3 (CD142), the marker used above (11) (Figure 1b).

Dedifferentiated acinar and duct cells did have equal expression of the embryonic

precursor gene *PDX1*, confirming our previous work (13). *PTF1A*, *NR5A2*, *RBPJL*,

FOXA3 and *GATA4* were expressed higher in the dedifferentiated acini reminiscent of

their fully differentiated state. Of note, *GATA6* expression, which is required for embryonic acinar cell differentiation (29, 30) and a suppressor of pancreatic carcinogenesis (31, 32), was not different from the duct cells.

Conversely, duct cell-enriched genes included *CFTR*, *SOX9*, *NKX6.1*, *HES1*, *HNF1B* and *ONECUT2* (*HNF6B*), established markers of duct cell differentiation. The most significantly duct cell-enriched gene was *DCDC2* (Doublecortin Domain Containing 2), demonstrated to bind tubulin and enhance microtubule polymerization (33) (Figure 1d). In the duct cell fraction, we found KEGG pathway enrichment for 'ECM receptor pathway interaction' and 'O-linked glycosylation'. Also, here, 'Pathways in cancer' featured here but the genes in this pathway differed from the ones in the dedifferentiated acinar cell fraction (Supplementary Table 1).

Among the differentially expressed genes were 71 transcription factors. The most significantly enriched transcription factor in the dedifferentiated acinar cell fraction was *MECOM* (log2FC = -6,09; Adj. p = 1,04E-123) (Figure 1d, Supplementary Table 1). *MECOM*, a known oncogene (34-36). It has been described as an activator of KRAS and glypican-1 in pancreatic cancer (20, 21). During our study, it was reported in relation to acinar cell-derived tumorigenesis (37). Hence, we confirmed *MECOM* expression in human PDAC samples (N=3) (Supplementary Figure 2a). We also used RNAseq analysis of stage 1 and 2 PDAC (N=103) where it correlated positively with overall survival (Supplementary figure 2b). In the same cohort, we previously reported a similar pattern for *SOX9* and a correlation in expression between *SOX9* and *MECOM* (Pearson R=0,49) (23). Pathway analysis of genes correlating with MECOM (Supplementary figure 2c) revealed similar pathways as those reported in our study on SOX9 where regulation of ErbB signaling was highlighted (23).

In conclusion, we FACS-purified human dedifferentiated acinar and duct cells and

analyzed their transcriptome. The dedifferentiated acinar cell fraction showed a

signature reminiscent of the acinar cell identity and specifically expressed *MECOM*, a

transcription factor that featured in the pathway 'Pathways of cancer'. MECOM was

expressed in PDAC where it correlated with SOX9. We hypothesized that MECOM

plays a role during acinar cell dedifferentiation, an event that initiates pancreatic tumor

development.

MECOM is expressed in developing acinar cells and is re-expressed during

pancreatitis-associated acinar cell dedifferentiation

We comprehensively assessed expression of MECOM during acinar cell differentiation

or loss thereof.

First, we confirmed the differential expression of *MECOM* in the purified human

dedifferentiated acinar and duct cells from the experiments above (Figure 2a). Using

RISH, *MECOM* mRNA appeared expressed at low levels in the acinar compartment of

normal human pancreas (0,24±0,09 mRNA dots/cell; N=3), but increased strongly in

samples of chronic pancreatitis (1,46±0,10 mRNA dots/cell; N=3) (Figure 2b,c).

Within the human purified dedifferentiated acinar cell fraction, *MECOM* showed strong

correlation with *CD142* (Pearson R = 0,926, adj. P < 0,01), as discussed before (Figure

1). Immunofluorescent co-staining of MECOM and CD142 after 4 days of culture

showed strong colocalization of the two proteins (56,94±10,63% MECOM+/CD142+

cells; N=5), while MECOM did not co-localize with the duct cell marker CK19

(5,69±2,32% MECOM+/CK19+ cells; N=5) (Figure 2d,e).

Next, we assessed the expression pattern (Figure 2f-i) and levels (Figure 2j) of *Mecom*

mRNA in murine pancreas. In contrast to normal adult mouse pancreas where *Mecom*

was expressed at very low levels (Figure 2f,j) a significantly increased expressed was

noted in E16.5 pancreas where it localized to pro-acinar tip cells (Figure 2g,j) (38, 39).

Similar to human, *Mecom* became re-expressed during acute and chronic pancreatitis

in mouse (Figure 2h,i,j).

Altogether, these data show that *Mecom* is transiently expressed in developing acinar

cells and becomes re-expressed during pancreatitis in dedifferentiated acinar cells.

SOX9 regulates *Mecom* expression in dedifferentiated acinar cells

SOX9 is an important marker of acinar cell dedifferentiation and is indispensable for

tumor formation in the pancreas (23). SOX9 correlates with MECOM in PDAC patients,

as shown above, but also in other tissues (40, 41). Hence, we investigated whether

these two transcription factors might also correlate in the context of acinar cell

dedifferentiation. Indeed, we observed a strong correlation between *MECOM* and

SOX9 (Pearson R = 0,83; *p=0,015; N=6) (Figure 3a) in the human exocrine cell

cultures in which acinar cells gradually dedifferentiate, a finding which was confirmed

at the protein level (Figure 3b).

Based on the observed induction of *Mecom* after 48 hours of caerulein-induced acute

pancreatitis (Figure 2h), we investigated *Mecom* expression in *Sox9* loss-of-function

mice (Figure 3c) (13, 42). ElaCreERT;*Sox9*$^{f/f}$ mice showed significant reduction in

Mecom compared to control mice (ElaCreERT) (19,48 ± 4,54% reduction in mRNA

dots) (Figure 3c). suggestive for SOX9 to act as an upstream regulator of *Mecom*.

To further study whether SOX9 directly regulated *Mecom* in the acinar cell context, we

performed a promoter-reporter assay on the partially differentiated mouse acinar cell

line 266-6 (that expresses *Mecom*, as shown further). We observed low increase in

luciferase activity when cells were transfected with the *Mecom* enhancer expression

plasmid alone. After *Sox9* overexpression, luciferase activity increased significantly

(Figure 3d), indicating that SOX9 regulates the Mecom enhancer in partially

differentiated acinar cells.

In conclusion, we showed that *Mecom* is (directly) regulated by SOX9 in pancreatic

acinar cell context.

***Mecom* maintains cell adhesion and cell viability in a partially differentiated**

acinar cell line and in human and mouse acinar cell cultures

A lentiviral all-in-one CRISPR/Cas9-mediated approach comparing 3 different gRNAs

designed to knock out (KO) *Mecom* in a partially differentiated acinar cell line (266-6)

(Supplementary Figure 3), enabled us to study the role of Mecom in acinar cell

dedifferentiation. RISH for *Mecom* in the loss-of-function condition (Figure 4a) showed

significant reduction of mRNA expression (27,30±4,25% dots/cell decrease). In the

Mecom KO condition, the cells were not able to organize into the typical 'acinar dome-

like' structures that were observed in the cells transduced with non-targeting gRNA

(Figure 4b), suggestive of loss of proper cell-cell contacts. We also observed a

decrease in absolute cell number (Figure 4c), an overall diminished cell viability (Figure

4d) and a significant increase in cleaved caspase 3+ cells (Figure 4e), all indicative of

cell death.

We further studied downstream transcriptional effects of Mecom loss in this cell line by

RNAseq analysis. Differential gene expression analysis revealed downregulation of

genes enriched in 'Focal adhesion', 'ECM receptor interaction' and 'Pathways in

cancer' (Figure 4f). This supported our observations of diminished dome formation due

to potentially impaired cell adhesion. We also compared the differentially expressed

genes from the Mecom KO condition with a publicly available MECOM ChIP-Seq

dataset (43) and found several overlapping genes (Figure 4g) that we further

considered, in particular integrins that link the extracellular matrix (ECM) components

such as collagen to the cytoskeleton. The data suggested that loss of *Mecom* might

compromise cell-cell interaction by diminished Collagen IV-Integrin a2 interaction and

downstream ERK-mediated signaling (Figure 4h), the latter considered crucial in acinar

cell dedifferentiation (44).

Diminished expression of collagen type IV and reduced ERK signaling upon Mecom

loss-of-function was confirmed by immunofluorescent staining (Figure 4i,j).

In conclusion, *Mecom* is indispensable to maintain cell adhesion and cell viability in a

partially differentiated acinar cell line.

Next, we crossed ElaCreERT deletor mice with mice carrying a floxed exon 4 of *Evi*,

an exon which is conserved in all transcript variants (24) (Figure 5a). We studied

effects in dedifferentiated acinar cells isolated from these mice using an established

cell culture model, similar to the human cell culture model described above (13).

Immunohistochemical staining for MECOM showed 42,71±13,42% less MECOM

positive cells in freshly isolated (day 0) MECOM KO exocrine cells, consistent with

previous reports on the recombination efficiency of the ElaCreERT strain (45-47).

Interestingly, by day 4, we detected only 23,89±13,42% less MECOM positive cells in

the KO condition suggesting selective loss of MECOM deficient cells (Supplementary

Figure 4).

We observed that after cell isolation the control cells remained organized in small

acinar spheroid-like structures while in the MECOM KO condition they often presented

as single cells. After 4 days of culture, clusters did form in the ElaCreERT;*Mecom*[f/f]

mouse cells but cluster diameter was significantly smaller compared to those of

ElaCreERT control mice (57,21±15,73% reduction in average diameter) (Figure 5b,e),

suggestive of diminished cell adhesion and cell death of the single cells. In our

experience acinar cells cannot survive as single cells (e.g. after FACS purification).

Similar to our results in the 266-6 cells (Figure 4), an increased fraction of acinar cells

in the MECOM KO cultures stained positive for cleaved caspase 3 (81,21±26,47%

increase in cleaved caspase 3+ cells) (Figure 5c,f). Notably, the MECOM KO cultures

also showed fewer expression of CK19+ cells in comparison to the control cells

(25,79±6,33% reduction in CK19+ cells), suggesting a lower capacity for

dedifferentiation (Figure 5d,g).

These observations were confirmed by shRNA-mediated knockdown in cultured

human exocrine cells in culture. At 24h, 4 and 7 days after transduction, average

cluster diameter was significantly decreased (Supplementary Figure 5), underscoring

the compromised cell-cell adhesion.

In conclusion, MECOM is indispensable for maintenance of cell-cell adhesion and cell

viability in primary cultures of mouse and human acinar cells.

Mice with *Mecom*-deficient acinar cells show impaired recovery from acute

pancreatitis

Since the primary cell culture models above recapitulate acinar cell dedifferentiation

as it occurs during pancreatitis (11-13), we subjected control and MECOM KO mice to

caerulein-induced acute pancreatitis. In wild-type mice, acute pancreatitis peaks

around day 4 and resolves by day 11 (Figure 6a). We observed no macroscopic

differences in the pancreas nor in weight of the pancreas or body weight when

comparing Mecom-deficient with control mice (Supplementary Figure 6).

Haematoxylin-eosin staining however showed more inter-acinar spaces at day 11 in

the Mecom KO mice (Figure 6b,f). Similar to the phenotype observed in the 266-6 cells

(Figure 4) and the *ex vivo* cultures (Figure 5), this was preceded by an increased

incidence of apoptosis (Figure 6c,g). Since we observed more inter-acinar space in the

Mecom KO mice, we analyzed immune cell infiltration. Similar to a previous report (37),

at day 11, CD3 positive area in MECOM KO mice remained at a level comparable to

the acute phase of pancreatitis (day 4) whereas in the control mice it was significantly

decreased (Figure 6e,i). F4/80+ macrophages showed a similar trend (not shown).

Remarkably, no difference in collagen deposition could be detected between control

and MECOM KO mice (Figure 6d,h) indicating that the inter-acinar space was not filled

with collagen. The histological pattern showing accumulation of excessive liquid in the

interstitial space was consistent with that of edema. Also, we did not observe any

changes in acinar-to-ductal transdifferentiation markers at day 11 (not shown).

Altogether, these results indicate that MECOM-deficiency in pancreatic acinar cells

under the experimental stress of acute pancreatitis leads to increased acinar cell death

with ensuing immune cell infiltration and edema.

Discussion

When subjected to stress, such as occurring during chronic pancreatitis, acinar cells

dedifferentiate and acquire a ductal cell-like phenotype, rendering them prone to *Kras*

induced transformation, in contrast to the native duct cells which seem more resistant

to neoplastic development. Where studies in the past focused on the acinar cell

plasticity and on how the dedifferentiated acinar cells become duct-like (13), the

inherent differences between the two cell populations that may underly a different

propensity to tumor formation remained unclear. Therefore, we profiled, for the first

time to our knowledge, the transcriptome of human dedifferentiated acinar cells versus

their native duct cell counterpart. This comparative transcriptomic analysis provided

several new insights, including the fact that both cell populations were enriched in

'Pathways of cancer' although the related genes differed in both fractions. We identified

MECOM, a transcription factor known to be involved in (pancreatic) cancer, as the

most uniquely expressed gene in the dedifferentiated acinar cells.

MECOM is known to regulate many aspects of tumorigenesis including differentiation,

proliferation and apoptosis (17-19, 40). It is a regulator of stomach-specific genes,

some of which become upregulated in dedifferentiated acinar cells (23). Previously, a

role for MECOM in pancreas cancer was proposed (20, 21, 37). Here, we focused on

its role in acinar cell dedifferentiation. We uncovered that MECOM is uniquely

expressed in dedifferentiated mouse and human acinar cells and absent from native

duct cells, both in cell cultures and in tissue samples. Notably, MECOM expression

levels appeared even higher in embryonic mouse pancreas, specifically in the pro-

acinar tip cells again suggesting a relation to an immature acinar cell phenotype.

Interestingly, SOX9 is a known regulator of acinar cell dedifferentiation and a driver of

pancreatic tumor formation (21, 48). Here, SOX9 expression correlated with that of

MECOM. An interplay of MECOM and SOX9 has been described in other tissues (12,

13, 44) and we confirmed that SOX9 transactivates the *Mecom* enhancer in an acinar

cell context. Still, it requires further investigation why SOX9 does not transactivate

MECOM expression in duct cells where SOX9 expression is the highest.

We observed a clear phenotype when culturing MECOM deficient acinar cells (266-6

cells and primary mouse and human acinar cells), i.e. impaired formation of acinar

domes or cell clusters, suggesting impaired cell adhesion and higher rates of

apoptosis. We showed that MECOM depletion reduced the cell-cell interactions

through *Itga2* and its ligands such as *Col4a2*. In our MECOM KO cells, the expression

of several integrins (*Itga2, Itga4, Itga5 and Itga9*) was decreased. Importantly,

available ChIP-Seq data of MECOM in human ovarian carcinoma cells also showed direct binding to the promotor regions of *Col4a2* and *Itga2* (43). We confirmed this in MECOM KO cells, where reduced activity of the ERK signaling pathway could be directly linked to reduced acinar cell dedifferentiation, a process for which this signaling pathway is critical (12, 44). In a caerulein-induced acute pancreatitis model in MECOM-deficient mice, we again make similar observations and demonstrated a prolonged immune cell infiltration, similar as reported recently (37).

We propose that normal acinar cells have two options when they are subjected to stress, either to subside in an "incognito" reversible dedifferentiated state, or to die. MECOM seems indispensable for the former option. Ye et al., who, like us did not provide formal proof by lineage tracing, claimed that MECOM-deficient acinar cells transdifferentiate more extensively during pancreatitis (37). We suggest that MECOM-deficient acinar cells undergo apoptosis and therefore cannot contribute any longer to the CK19+ transdifferentiated cell pool. This is supported by our observations increased inter-acinar spaces filled with fluid (edema) in the acute pancreatitis model. Such phenotype has also been reported in EGF knockout mice (49). We also did not detect any increase in collagen deposition around the recovering acini, which clearly differs from acinar to ductal transdifferentiation that is accompanied by typical collagen deposition (50), further supporting our conclusion.

Altogether, these results show that dedifferentiated acinar cells have a clearly distinct gene expression profile compared to native duct cells. Under stress, MECOM expression is turned on, downstream of SOX9, where it prevents acinar cell death and allowing cellular dedifferentiation.

Acknowledgements

Biospecimens and clinical data were provided by the Australian Pancreatic Cancer Genome Initiative (APGI, www.pancreaticcancer.net.au), which is supported by an Avner Pancreatic Cancer Foundation Grant (www.avnersfoundation.org.au). This work was supported by the Research Foundation Flanders (FWO) (Odysseus project and FWO research project to IR (FWOODYS12 and FWOAL931)). We thank Gunter Leuckx, Emmy De Blay, Veerle Laurysens, Geert Stangé, the laboratory of Liver Cell Biology Research Group at Vrije Universiteit Brussel and Flemish Institute of Biotechnology for expert technical assistance. We thank Patrick Jacquemin for providing the ElaCreERT mice, and Susumu Goyama and Mineo Kurokawa for providing the Evi-1 floxed mice.

Conflict of interest

The authors declare no competing financial interests

References

1.   Rahib L, Smith BD, Aizenberg R, Rosenzweig AB, Fleshman JM, Matrisian LM. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. Cancer Res. 2014;74(11):2913-21.

2.   Krah NM, De La OJ, Swift GH, Hoang CQ, Willet SG, Chen Pan F, et al. The acinar differentiation determinant PTF1A inhibits initiation of pancreatic ductal adenocarcinoma. Elife. 2015;4.

3.   Direnzo D, Hess DA, Damsz B, Hallett JE, Marshall B, Goswami C, et al. Induced Mist1 expression promotes remodeling of mouse pancreatic acinar cells. Gastroenterology. 2012;143(2):469-80.

4.   von Figura G, Morris JPt, Wright CV, Hebrok M. Nr5a2 maintains acinar cell

differentiation and constrains oncogenic Kras-mediated pancreatic neoplastic

initiation. Gut. 2014;63(4):656-64.

5.   Kopp JL, von Figura G, Mayes E, Liu FF, Dubois CL, Morris JPt, et al.

Identification of Sox9-dependent acinar-to-ductal reprogramming as the principal

mechanism for initiation of pancreatic ductal adenocarcinoma. Cancer Cell.

2012;22(6):737-50.

6.   De La OJ, Emerson LL, Goodman JL, Froebe SC, Illum BE, Curtis AB, et al.

Notch and Kras reprogram pancreatic acinar cells to ductal intraepithelial neoplasia.

Proc Natl Acad Sci U S A. 2008;105(48):18907-12.

7.   Habbe N, Shi G, Meguid RA, Fendrich V, Esni F, Chen H, et al. Spontaneous

induction of murine pancreatic intraepithelial neoplasia (mPanIN) by acinar cell

targeting of oncogenic Kras in adult mice. Proc Natl Acad Sci U S A.

2008;105(48):18913-8.

8.   Tuveson DA, Zhu L, Gopinathan A, Willis NA, Kachatrian L, Grochow R, et al.

Mist1-KrasG12D knock-in mice develop mixed differentiation metastatic exocrine

pancreatic carcinoma and hepatocellular carcinoma. Cancer Res. 2006;66(1):242-7.

9.   Grippo PJ, Nowlin PS, Demeure MJ, Longnecker DS, Sandgren EP.

Preinvasive pancreatic neoplasia of ductal phenotype induced by acinar cell targeting

of mutant Kras in transgenic mice. Cancer Res. 2003;63(9):2016-9.

10.   Guerra C, Schuhmacher AJ, Canamero M, Grippo PJ, Verdaguer L, Perez-

Gallego L, et al. Chronic pancreatitis is essential for induction of pancreatic ductal

adenocarcinoma by K-Ras oncogenes in adult mice. Cancer Cell. 2007;11(3):291-302.

11.   Baldan J, Houbracken I, Rooman I, Bouwens L. Adult human pancreatic acinar

cells dedifferentiate into an embryonic progenitor-like state in 3D suspension culture.

Sci Rep. 2019;9(1):4040.

12.   Houbracken I, de Waele E, Lardon J, Ling Z, Heimberg H, Rooman I, et al.

Lineage tracing evidence for transdifferentiation of acinar to duct cells and plasticity of

human pancreas. Gastroenterology. 2011;141(2):731-41, 41 e1-4.

13.   Pinho AV, Rooman I, Reichert M, De Medts N, Bouwens L, Rustgi AK, et al.

Adult pancreatic acinar cells dedifferentiate to an embryonic progenitor phenotype with

concomitant activation of a senescence programme that is present in chronic

pancreatitis. Gut. 2011;60(7):958-66.

14.   Roy N, Malik S, Villanueva KE, Urano A, Lu X, Von Figura G, et al. Brg1

promotes both tumor-suppressive and oncogenic activities at distinct stages of

pancreatic cancer formation. Genes Dev. 2015;29(6):658-71.

15.   Lee AYL, Dubois CL, Sarai K, Zarei S, Schaeffer DF, Sander M, et al. Cell of

origin affects tumor development and phenotype in pancreatic ductal adenocarcinoma.

Gut. 2018.

16.   Pinho AV, Chantrill L, Rooman I. Chronic pancreatitis: a path to pancreatic

cancer. Cancer Lett. 2014;345(2):203-9.

17.   Konantz M, Andre MC, Ebinger M, Grauer M, Wang H, Grzywna S, et al. EVI-1

modulates leukemogenic potential and apoptosis sensitivity in human acute

lymphoblastic leukemia. Leukemia. 2013;27(1):56-65.

18.   Rommer A, Steinmetz B, Herbst F, Hackl H, Heffeter P, Heilos D, et al. EVI1

inhibits apoptosis induced by antileukemic drugs via upregulation of

CDKN1A/p21/WAF in human myeloid cells. PLoS One. 2013;8(2):e56308.

19. Glass C, Wuertzer C, Cui X, Bi Y, Davuluri R, Xiao YY, et al. Global Identification

of EVI1 Target Genes in Acute Myeloid Leukemia. PLoS One. 2013;8(6):e67134.

20. Tanaka M, Ishikawa S, Ushiku T, Morikawa T, Isagawa T, Yamagishi M, et al.

EVI1 modulates oncogenic role of GPC1 in pancreatic carcinogenesis. Oncotarget.

2017;8(59):99552-66.

21. Tanaka M, Suzuki HI, Shibahara J, Kunita A, Isagawa T, Yoshimi A, et al. EVI1

oncogene promotes KRAS pathway through suppression of microRNA-96 in

pancreatic carcinogenesis. Oncogene. 2014;33(19):2454-63.

22. Desai BM, Oliver-Krasinski J, De Leon DD, Farzad C, Hong N, Leach SD, et al.

Preexisting pancreatic acinar cells contribute to acinar cell, but not islet beta cell,

regeneration. J Clin Invest. 2007;117(4):971-7.

23. Grimont A, Pinho AV, Cowley MJ, Augereau C, Mawson A, Giry-Laterriere M,

et al. SOX9 regulates ERBB signalling in pancreatic cancer development. Gut.

2015;64(11):1790-9.

24. Goyama S, Yamamoto G, Shimabe M, Sato T, Ichikawa M, Ogawa S, et al. Evi-

1 is a critical regulator for hematopoietic stem cells and transformed leukemic cells.

Cell Stem Cell. 2008;3(2):207-20.

25. MacDonald RJ, Swift GH, Real FX. Transcriptional control of acinar

development and homeostasis. Prog Mol Biol Transl Sci. 2010;97:1-40.

26. Baer R, Cintas C, Dufresne M, Cassant-Sourdy S, Schonhuber N, Planque L,

et al. Pancreatic cell plasticity and cancer initiation induced by oncogenic Kras is

completely dependent on wild-type PI 3-kinase p110alpha. Genes Dev.

2014;28(23):2621-35.

27.   Elghazi L, Weiss AJ, Barker DJ, Callaghan J, Staloch L, Sandgren EP, et al.

Regulation of pancreas plasticity and malignant transformation by Akt signaling.

Gastroenterology. 2009;136(3):1091-103.

28.   Wu CY, Carpenter ES, Takeuchi KK, Halbrook CJ, Peverley LV, Bien H, et al.

PI3K regulation of RAC1 is required for KRAS-induced pancreatic tumorigenesis in

mice. Gastroenterology. 2014;147(6):1405-16 e7.

29.   Shi ZD, Lee K, Yang D, Amin S, Verma N, Li QV, et al. Genome Editing in

hPSCs Reveals GATA6 Haploinsufficiency and a Genetic Interaction with GATA4 in

Human Pancreatic Development. Cell Stem Cell. 2017;20(5):675-88 e6.

30.   Tiyaboonchai A, Cardenas-Diaz FL, Ying L, Maguire JA, Sim X, Jobaliya C, et

al. GATA6 Plays an Important Role in the Induction of Human Definitive Endoderm,

Development of the Pancreas, and Functionality of Pancreatic beta Cells. Stem Cell

Reports. 2017;8(3):589-604.

31.   Martinelli P, Carrillo-de Santa Pau E, Cox T, Sainz B, Jr., Dusetti N, Greenhalf

W, et al. GATA6 regulates EMT and tumor dissemination, and is a marker of response

to adjuvant chemotherapy in pancreatic cancer. Gut. 2017;66(9):1665-76.

32.   Martinelli P, Madriles F, Canamero M, Pau EC, Pozo ND, Guerra C, et al. The

acinar regulator Gata6 suppresses KrasG12V-driven pancreatic tumorigenesis in

mice. Gut. 2016;65(3):476-86.

33.   Longoni N, Kunderfranco P, Pellini S, Albino D, Mello-Grand M, Pinton S, et al.

Aberrant expression of the neuronal-specific protein DCDC2 promotes malignant

phenotypes and is associated with prostate cancer progression. Oncogene.

2013;32(18):2315-24, 24 e1-4.

34.   Izutsu K, Kurokawa M, Imai Y, Maki K, Mitani K, Hirai H. The corepressor CtBP

interacts with Evi-1 to repress transforming growth factor beta signaling. Blood.

2001;97(9):2815-22.

35.   Yuasa H, Oike Y, Iwama A, Nishikata I, Sugiyama D, Perkins A, et al. Oncogenic

transcription factor Evi1 regulates hematopoietic stem cell proliferation through GATA-

2 expression. EMBO J. 2005;24(11):1976-87.

36.   Buonamici S, Chakraborty S, Senyuk V, Nucifora G. The role of EVI1 in normal

and leukemic cells. Blood Cells Mol Dis. 2003;31(2):206-12.

37.   Ye J, Huang A, Wang H, Zhang AMY, Huang X, Lan Q, et al. PRDM3 attenuates

pancreatitis and pancreatic tumorigenesis by regulating inflammatory response. Cell

Death & Disease. 2020;11(3):187.

38.   Schaffer AE, Freude KK, Nelson SB, Sander M. Nkx6 transcription factors and

Ptf1a function as antagonistic lineage determinants in multipotent pancreatic

progenitors. Dev Cell. 2010;18(6):1022-9.

39.   Zhou Q, Law AC, Rajagopal J, Anderson WJ, Gray PA, Melton DA. A

multipotent progenitor domain guides pancreatic organogenesis. Dev Cell.

2007;13(1):103-14.

40.   Garside VC, Cullum R, Alder O, Lu DY, Vander Werff R, Bilenky M, et al. SOX9

modulates the expression of key transcription factors required for heart valve

development. Development. 2015;142(24):4340-50.

41.   Mateo F, Arenas EJ, Aguilar H, Serra-Musach J, de Garibay GR, Boni J, et al.

Stem cell-like transcriptional reprogramming mediates metastatic resistance to mTOR

inhibition. Oncogene. 2017;36(19):2737-49.

42.   Jensen JN, Cameron E, Garay MV, Starkey TW, Gianani R, Jensen J.

Recapitulation of elements of embryonic development in adult mouse pancreatic

regeneration. Gastroenterology. 2005;128(3):728-41.

43.   Bard-Chapeau EA, Jeyakani J, Kok CH, Muller J, Chua BQ, Gunaratne J, et al.

Ectopic viral integration site 1 (EVI1) regulates multiple cellular processes important

for cancer and is a synergistic partner for FOS protein in invasive tumors. Proc Natl

Acad Sci U S A. 2012;109(6):2168-73.

44.   Collins MA, Yan W, Sebolt-Leopold JS, Pasca di Magliano M. MAPK signaling

is required for dedifferentiation of acinar cells and development of pancreatic

intraepithelial neoplasia in mice. Gastroenterology. 2014;146(3):822-34 e7.

45.   Means AL, Meszoely IM, Suzuki K, Miyamoto Y, Rustgi AK, Coffey RJ, Jr., et

al. Pancreatic epithelial plasticity mediated by acinar cell transdifferentiation and

generation of nestin-positive intermediates. Development. 2005;132(16):3767-76.

46.   Stanger BZ, Stiles B, Lauwers GY, Bardeesy N, Mendoza M, Wang Y, et al.

Pten constrains centroacinar cell expansion and malignant transformation in the

pancreas. Cancer Cell. 2005;8(3):185-95.

47.   Strobel O, Dor Y, Alsina J, Stirman A, Lauwers G, Trainor A, et al. In vivo lineage

tracing defines the role of acinar-to-ductal transdifferentiation in inflammatory ductal

metaplasia. Gastroenterology. 2007;133(6):1999-2009.

48.   Tanaka M, Shibahara J, Ishikawa S, Ushiku T, Morikawa T, Shinozaki-Ushiku

A, et al. EVI1 expression is associated with aggressive behavior in intrahepatic

cholangiocarcinoma. Virchows Arch. 2019;474(1):39-46.

49.   Ardito CM, Grüner BM, Takeuchi KK, Lubeseder-Martellato C, Teichmann N,

Mazur PK, et al. EGF receptor is required for KRAS-induced pancreatic tumorigenesis.

Cancer Cell. 2012;22(3):304-17.

50.   Hamada S, Masamune A. Elucidating the link between collagen and pancreatic

cancer: what's next? Expert Rev Gastroenterol Hepatol. 2018;12(4):315-7.

Figure Legends

Figure 1. RNAseq analysis identifies *MECOM* as uniquely expressed in

dedifferentiated acinar cells. (a) Schematic overview of FACS-purification of human

dedifferentiated acinar and duct cells, using acinar-specific UEA-1 lectin. (b)

Immunofluorescent staining of FACS-purified dedifferentiated acinar (UEA-1+/CA19.9-

) and duct (UEA-1-/CA19.9+) cells at day 4 with CD142 and CA19.9. DNA is labelled

in blue. (c) Table of genes uniquely expressed in human dedifferentiated acinar cells

(green) and duct cells. Selection limited to genes with BaseMean ≥ 1000, Log2FC ≥ 2

or ≤ -2 and P.adj ≤ 0,01. (d) KEGG pathways enriched (Adj. p-value < 0,01) in human

dedifferentiated acinar cells (N=5).

Figure 2. MECOM is uniquely upregulated in dedifferentiated acinar cells and not

in duct cells. (a) qPCR for MECOM on purified human dedifferentiated acinar (UEA-

1+/CA19.9-) versus duct cells (UEA-1-/CA19.9+) after FACS isolation at day 4 of

culture, compared to the mixed exocrine fraction at day of isolation (D0) (N=5; mean ±

SD; ***p<0,005; One-way ANOVA with post-hoc Bonferroni correction). (b,c) RNA-ISH

for MECOM on tissue from normal human pancreas (b) and human chronic pancreatitis

(c) Scale = 50µM. Arrows indicate individual mRNA dots. (d,e) Immunofluorescent co-

staining of human mixed exocrine fraction at day 4 of culture for MECOM and CD142

(d) and for MECOM and CK19 (e). Scale = 50µM. (f-i) RNA-ISH for *Mecom* and (j)

Mecom expression level in normal mouse pancreas (f; dotted line indicates a normal

duct, striped line indicates the outer edge of an islet), embryonic pancreas (g); E16.5

embryonic pancreas; dashed line indicates groups of tip cells) and in a mouse model

for acute (h) and chronic (i) pancreatitis. Scale = 50µM. (j) 5 individual microscopic

fields per replicate were quantified. (N=5; mean ± SD; *p<0,05; **p<0,01; One-way

ANOVA with post-hoc Bonferroni correction;).

Figure 3. SOX9 regulates MECOM during acinar cell dedifferentiation. (a) qPCR

for MECOM and SOX9 expression on human exocrine mixed fraction in culture over

time (day 1 to day 8). Each donor is indicated as a different colored line. (b) Immunoblot

for MECOM, SOX9 and actin as loading control on human exocrine mixed fraction in

culture over time. Representative blot from N=3 human donor samples. The different

isoforms of MECOM locus are indicated by arrows. (c) Generation of acinar-specific

(Elastase) Cre deletor mice with Sox9^{ex5} flanked by loxP sites. Acinar-specific deletion

of SOX9 is performed in ElaCreERT;*Sox9*$^{f/f}$ versus control (ElaCreERT) littermates by

administering tamoxifen 3 times over a period of 5 days, followed by a two-week

washout. Acute pancreatitis is induced by administering caerulein by 7 hourly injections

for 3 days over a period of 5 days. Mice are sacrificed at day 4 after caerulein

administration. Immunohistochemical staining for SOX9 and RNA in situ hybridization

of *Mecom* mRNA expression in mouse pancreatic tissue after experimentally induced

acute pancreatitis in control (ElaCreERT) and acinar-specific Sox9 knockout

(ElaCreERT;*Sox9*$^{f/f}$) mice. Scale = 50µM. (N=3; **p<0.01; unpaired two-tailed t-test)

(d) Luciferase reporter assay in 266-6 cells, 48h after transfection. (mean ± SD; N=3;

***p<0,001; one-way ANOVA with post-hoc Bonferroni correction)

Figure 4. *Mecom* retains cell adhesion and survival in 266-6 cells. (a) RNA in situ

hybridization (RISH) on control (NT-gRNA) and Mecom KO (gRNA3) 266-6 cells. Scale

= 200µM. (N=3; **p<0,01; unpaired two-tailed t-test). (b) Phase contrast images of

control (NT-gRNA) and Mecom KO (gRNA3) 266-6 cells. Scale = 200µM. (c) Absolute

number of cells at 96h after seeding 50.000 cells (mean ± SD; N=3; **p<0,01; unpaired

two-tailed t-test) (d) Cell viability measured relative to background levels of the culture

medium by CellTiter Glo assay (mean ± SD; N=3; ***p<0,001; unpaired two-tailed t-

test) (e) Immunofluorescent staining and quantification of cleaved caspase 3+ cells.

(N=3; **p<0,01; unpaired two-tailed t-test) (f) KEGG pathway analysis on differential

gene expression profiles after RNAseq analysis of MECOM wildtype versus MECOM

KO cells (N=3) (g) Venn diagrams showing genes overlapping between differentially

expressed genes in MECOM KO 266-6 cells and publicly available ChIP-seq dataset

from Bard-Chapeau et al. (43) (h) Schematic representation of our hypothesis. (i)

Collagen IV immunofluorescent staining on 266-6 NTgRNA and gRNA3-transduced

cultured cells. (j) Immunoblot for phospho-ERK and total ERK and quantifications of

pERK relative to total protein stain. (mean ± SD; N=3; *p<0,05; Unpaired two-tailed t-

test)

Figure 5. *Mecom* retains cell adhesion and survival in cultured acinar cells. (a)

Acinar cell-specific deletion of MECOM is performed by administering tamoxifen 3

times over a period of 5 days both subcutaneously and by gavage, followed by a two-

week washout period. Acute pancreatitis is induced by administering caerulein by 7

hourly injections for 3 days over a period of 5 days. Mice are sacrificed at day 4 after

caerulein treatment. (b,e) Phase contrast images and quantification of mean cluster

surface of ElaCreERT and ElaCreERT;*Mecom*^f/f isolated acinar cells immediately after

isolation (D0) and at day 4 of culture (D4). Scale = 100µM. (c,f) Cleaved caspase 3

staining and quantification on ElaCreERT and ElaCreERT;*Mecom*^f/f acinar cells at day

4 of culture. Scale = 25µM. *(d,g)* CK19 staining and quantification of ElaCreERT and

ElaCreERT;*Mecom*^f/f acinar cells at day 4 of culture. Scale = 25µM. (mean ± SD; N=6;

*p<0,05; **p<0,01; two-way ANOVA with post-hoc Bonferroni correction.)

Figure 6. *Mecom* depletion in acinar cells *in vivo* causes pronounced cell death

during recovery after caerulein-induced acute pancreatitis. (a) Acinar-specific

deletion of MECOM is performed by administering tamoxifen 3 times over a period of

5 days, followed by a two-week washout period. Acute pancreatitis is induced by

administering caerulein by 7 hourly injections for 3 days over a period of 5 days. Mice

are sacrificed at day 4 and day 11 after caerulein administration. (b,f) H&E staining

and quantification of acinar/non-acinar area ratio relative to day 4. (c,g) Cleaved

caspase 3 immunohistochemical staining and quantification at day 4. (d,h) Collagen

staining and quantification relative to day 4. Scale = 50µM. (e,i) CD3 staining and

quantification relative to day 4 for ElaCreERT and ElaCreERT;*Mecom*^f/f mice. Scale =

50µM. (mean ± SD, N=6, **p<0,01, ns = non-significant. two-way ANOVA with post-

hoc Bonferroni correction)

Figure 1

a

Isolation exocrine cells
UEA-1 labelling

CA19.9 labelling

4 days culture

FACS sorting

UEA-1 lectin
Exocrine cell clusters

Dedifferentiated acinar cells
UEA-1+/CA19.9-

Duct cells
UEA-1-/CA19.9+

b

Dedifferentiated acinar cells
UEA-1+/CA19.9-

Duct cells
UEA-1-/CA19.9+

UEA-1 CD142
CA19.9 HOECHST

c

Dedifferentiated acinar cells (FC≤-2, adj.P≤0,01)	
Pathway	Adj.P
Protein digestion and absorption	1,185208E-11
Pancreatic secretion	1,267413E-10
Metabolic pathways	8,920072E-04
Pathways in cancer	2,494694E-03
Complement and coagulation cascades	2,704916E-03
Metabolism of xenobiotics by cytochrome P450	3,886956E-03
Fat digestion and absorption	3,953574E-03
Mineral absorption	3,953574E-03
PI3K-Akt signaling pathway	4,032070E-03
Drug metabolism - cytochrome P450	5,712517E-03
Steroid hormone biosynthesis	8,027015E-03
Glycine, serine and threonine metabolism	9,867525E-03

d

symbol	log2FoldChange	padj
DCD2	4,178944	7,066393E-126
MXRA5	-5,436366	1,103561E-129
GSTA2	-5,679022	2,094860E-159
IGFBP2	-5,777801	8,550959E-126
MGST1	-5,895840	1,446389E-147
MECOM	-6,091834	1,040978E-123
LGALS2	-4,859710	7,753555E-130

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6