Title: Effects of scent enrichment on behavioural and physiological indicators of stress in zoo

primates – a preliminary study on red-ruffed lemurs and Lar gibbons

Short title: Effect of scent enrichment on the welfare of red-ruffed lemur and Lar gibbon

Authors: Stefano Vaglio[1,2], Stefano S. K. Kaburu[3*], Christopher Young[1], Richard Pearce[1],

Sarah Smith[1], Beth Tompkins[1], Emily Elwell[1], Sara Fontani[1], Giovanna Marliani[4] & Pier

Attilio Accorsi[4]

Affiliations of Authors:

[1]Department of Biology, Chemistry and Forensic Science, University of Wolverhampton,

Wolverhampton, UK

[2]Department of Anthropology & Behaviour, Ecology and Evolution Research (BEER)

Centre, Durham University, Durham, UK

[3]Department of Biomedical Science & Physiology, University of Wolverhampton,

Wolverhampton, UK

[4]Dipartimento di Scienze Mediche Veterinarie, Università di Bologna, Bologna, Italy

***Corresponding Author:**

Dr Stefano S. K. Kaburu

Department of Biomedical Science & Physiology, Faculty of Science and Engineering,

University of Wolverhampton, City Campus South, Wulfruna Street, Wolverhampton, WV1

1LY, UK

E-mail s.kaburu@wlv.ac.uk

Abstract

Almost half of the primate species recognized today worldwide are classified as endangered or critically endangered in the wild. Captive breeding is vital for primate conservation, with modern zoos serving a crucial role as breeders of populations acting as buffers against extinction, ambassadors of endangered species, and educators of the general public. However, captive populations may experience welfare issues that may also undermine their breeding success. In order to design and test a new scent enrichment programme to enhance the breeding success and well-being of critically endangered zoo primates, we carried out a preliminary study to assess the effects of lavender on captive red-ruffed lemurs (*Varecia rubra*) and Lar gibbons (*Hylobates lar*) as these species presently show a low success rate in captive breeding and are critically endangered in the wild. We combined behavioural observations and faecal endocrinology analyses to assess the effects of lavender on two captive troops (N = 8) housed at Dudley Zoo and Castle (UK). We recorded observations of natural species-specific and abnormal stress-related behaviours for 144 hr using instantaneous scan sampling. We collected 78 faecal samples and measured the faecal cortisol concentrations using radioimmunoassay. We found a significant effect by the scent enrichment on social interactions and stress-related behaviours (i.e., pacing and self-scratching), with both species reducing their rates of stress-related behaviours after they were exposed to lavender. We also found that both species displayed a significant increase in faecal glucocorticoids following the exposure to lavender. Our findings suggest that lavender may work as scent enrichment to decrease the stress levels of zoo primates across the major lineages, but its effects might change depending on the species-specific social lives and olfactory repertoires of primates.

Keywords: lavender, stereotypic behaviours, faecal cortisol, *Varecia rubra*, *Hylobates lar*

Introduction

Almost half of the total primate species recognized today worldwide are classified as endangered or critically endangered in the wild – primarily due to human activities [1]. Therefore, raising global scientific and public awareness of the plight of the world's primates is now vital [1]. The most important actions needed for ensuring the survival of these irreplaceable species are conservation, research, public education and outreach. Across all these contexts, zoos play a major role [2], as zoo animal populations are usually managed to educate the public regarding wildlife and their habitats, and to preserve endangered species through captive breeding and reintroduction programmes [3]. In this context, the maintenance of the genetic variation of such captive populations is imperative [4]. However, captive populations, potentially serving as buffers against extinction, experience problems that impair them from being viable for reintroduction into the wild. More specifically, zoo animal populations face reproductive challenges which have so far inhibited them from serving as viable 'reserve populations' [5]. Additionally, managing zoo populations is challenging because of the mismatch between natural and captive environments [6]. Primates have evolved distinct behavioural patterns, and difficulty in engaging in these behaviours can cause frustration or boredom, which, in turn, can lead to stress and development of abnormal behaviours [7] that may undermine their individual welfare and their breeding success.

To maintain captive healthy populations, with good genetic variability and thus high survival rates in case of reintroduction, modern zoos take part in conservation breeding programmes. As reproductive success is linked to how closely captive environmental conditions mirror those that primates would be experiencing in the wild [5], zoos also use environmental enrichments to manage captive populations. Environmental enrichments and conservation breeding programmes are strictly related, as enrichment is a dynamic process

that changes an animal's environment, increasing its behavioural choices and prompting a

wider range of natural and species-specific behaviours and abilities [8]. Furthermore,

enrichment can contribute to promoting resiliency to stress, which helps animals recovering

from adverse stimuli [9], improving both the exhibit from the visitor perspective and the

reproductive performance of the hosted animals [10]. Enrichment can also foster the essential

skills that animals need for their survival if reintroduced into their habitat [11].

Scent-based enrichments can be effective at increasing active behaviours in zoo

animals and potentially improve their welfare [9,12,13]. However, some authors reported

findings that are less clear or indicate that scent enrichment has little effect [14,15]. The

delivery mechanism of the scent and the type of scents used are crucial for the

implementation of novel olfactory enrichment programmes [16]. The majority of studies have

used spices or essential oils rather than focusing on natural or biological scents, but this may

not necessarily be appropriate for all species [14]. The main goal of olfactory enrichment is

to improve welfare of animals in captive environments, but there is also the possibility that

the use of scents can have further positive impacts in addition to this. For instance, scents

may elicit behavioural and physiological responses and thus it is important to consider the

use of olfactory enrichment to promote potential beneficial effects on reproductive success

[17].

Primates are traditionally considered "microsmatic" (i.e., with a poor sense of smell)

[18] and only a small proportion of studies on olfactory enrichment have been undertaken on

primate species [19] with the majority of scents currently used in these studies remaining

largely focused on essential oils, spices or herbs [14]. However, various lines of evidence

suggest that chemical communication may be important in primates [20]. In particular, it has

become increasingly clear that the sense of smell plays a crucial role in primate socio-sexual

communication, with semiochemicals being important for kin recognition, mate choice and

the regulation of socio-sexual behaviours [21]. However, little is known about the overall

effects of olfactory enrichment on primate species.

The overarching aim of our work is to design and test a new scent enrichment

programme to enhance the breeding success and well-being of critically endangered zoo

primates. In this context, we carried out a preliminary study which aimed to assess the effects

of lavender (which was chosen based on its effectiveness in humans and domestic animals;

reviewed in [22]) on resting and social behaviour as well as on both behavioural and

physiological indicators of stress in two captive primate species: red-ruffed lemurs (*Varecia*

rubra) and Lar gibbons (*Hylobates lar*). These species are currently classified as either

critically endangered (red-ruffed lemurs [23]) or endangered (Lar gibbons [24]), largely due

to the deforestation, logging and hunting activities that threaten the habitat and survival of

these species [23,24]. Therefore, designing and implementing strategies that improve welfare

and breeding success of these species in captivity is particularly crucial. In this study, we

predicted that lavender would reduce the stress levels of zoo red-ruffed lemurs and Lar

gibbons, which would be reflected by significant changes in behavioural (i.e., decrease of the

frequency of stress-related behaviours, and increase of the frequency of social behaviours)

and physiological indicators of well-being when comparing before (i.e., baseline period) and

after (i.e., post enrichment period) the scent enrichment.

Material and methods

Study subjects and housing

We studied two captive troops of red-ruffed lemurs (n = 3) and Lar gibbons (n = 5)

housed at Dudley Zoological Gardens (UK). The troop of red-ruffed lemurs consisted of two

related (brothers) adult males (15 y/o) and one unrelated adult female (14 y/o). The troop of

Lar gibbons consisted of one adult male (16 y/o), one adult females (17 y/o) and their

offspring – one adult female (6 y/o), two young males (2 y/o and 8 y/o, respectively).

We carried out behavioural observations and faecal sampling from September to

December 2019 (red-ruffed lemurs) and from September to December 2018 (Lar gibbons).

Both troops lived in an indoor enclosure (heated to 28 °C) with access to an outdoor

enclosure (i.e., "visitor walkthrough" enclosure in the case of red-ruffed lemurs).

Study protocol

We divided the overall study period in three periods: baseline, scent enrichment, post

enrichment. We collected behavioural data and faecal samples (see below "Assessment of

welfare") for four days per study period (12 days in total) over three months.

Scent enrichment

We cut white cotton sheets into 75 cm long and 5 cm wide strips, which we then

soaked with 20 drops naissance 100% pure lavender essential oil diluted with 12 ml of cold

boiled water. We prepared the scent cotton strips at early morning of each sampling day over

the scent enrichment period. We positioned these strips around both indoor and outdoor

enclosures; particularly, and we tied them approximately 1 m from the ground around the

climbing frames as these were the most used areas of the enclosures.

Assessment of welfare

We have used both behavioural (e.g., naturalistic species-specific behaviours,

stereotypic behaviours) and physiological (e.g., corticosteroid levels, regulated by

hypothalamic-pituitary-adrenal-HPA axis) methods to assess the effects of scent enrichment.

Behavioural data collection

We collected behavioural data by instantaneous scan sampling [25] of some

behaviours (Table 1), with behaviours recorded at 5-min intervals over the duration of six

hours from 9am to 3pm, four days per study period (baseline, scent enrichment, post

enrichment) over three months. We recorded a total of 144 hr of observations over the study

period (excluding the pilot study), with 50 scan samples each sampling day on each group

(see supporting information).

Table 1 - Ethogram

Behaviour	Description
Resting	Lying or sitting while awake, with eyes open and arms down by side of body.
Sleeping	Lying on back, front or side, eyes closed and whole body is relaxed
Grooming	Using fingers or mouth to pick through the coat, removing any foreign bodies from a conspecific
Play	Animal is engaging in activities such as chasing others, leaping around the enclosure etc. in a playful context
Self-scratching	An animal rubs their own body at a fast pace
Pacing	Animal walks back and forth in a distinct, unchanging pattern within the enclosure.

Hormone sampling and measurements

We collected faecal samples of all subjects every morning before behavioural

observations, whenever defecation was observed and the identity of the animal was known.

In total, we collected 78 samples (red-ruffed lemurs = 25; Lar gibbons = 53). The samples

were stored in a freezer at 20°C right after collection. At the end of the study period, the

collated samples were fully prepared by adding biological hazard labels onto each pot before

being delivered in a refrigerated (dry ice) to the Department of Veterinary Medical Sciences

and Animal Production Science of Bologna University for radioimmunoassay (RIA).

Cortisol concentrations were determined by RIA. All concentrations were expressed

in pg/mg of faecal matter. The extraction methodology followed the methods of Fontani and

colleagues [26]. In brief, five millilitres of a methanol:water (4:1 v/v) solution were added to

60 mg (wet weight) of faeces in capped-glass tube vials. The vials were then vortexed for 30

min using a multitube pulsing vortexer. After centrifugation at 1,500 g for 15 min, 5 ml ethyl

ether (BDH Italia, MI, Italy) and 0.2 ml NaHCO3 (5%; Sigma Chemical Co., St. Louis, MO)

were added to 1 ml of supernatant. This preparation was vortexed for 1 min and centrifuged

for 5 min at 1,500 g. The ether portion was aspirated with a pipette, and evaporated under an

airstream suction hood at 37°C. The dry residue was redissolved into 0.5 ml of 0.05 M

phosphate-buffered saline (PBS; pH 7.5).

Cortisol was assayed in the faecal samples according to the method of [27]. The

validation parameters of the analyses were as follows:

Cortisol. Sensitivity 3.10 pg/100 l; intra-assay variability 6.8%; interassay variability

9.3%; specificity (%), cortisol 100, corticosterone 9.5, 11$_\zeta$-hydroxyprogesterone 8.3,

cortisone 5.3, 11$_\zeta$-desoxycortisol 5.0, progesterone 0.6, desoxycorticosterone 0.5, 20$_\zeta$-

dihydrocortisone 0.4, testosterone 0.3, aldosterone 0.1, dehydroepiandrosterone <

0.0001, 5$_\zeta$-pregnenolone <0.0001, 17$^\vee$-estradiol < 0.0001, and cholesterol < 0.0001.

Statistical analyses

In order to assess the effect of scent enrichment on red-ruffed lemurs' and Lar

gibbons' behaviour and FGCs, we first generated three behavioural categories from the

individual behavioural measures we collected. More specifically, we generated: 1) a resting

category by adding up our data on resting and sleeping behaviours; 2) a social category by

combining our data on grooming and play; and, finally, 3) a stress category by combining our

data on pacing and self-scratching behaviours. We included scratching in this category since

this is commonly considered an indicator of anxiety [28]. We then used the *aov* function in R

(v 4.0.2) [28] to run a 2x2 Analysis of Variance (ANOVA), in which the rates of resting,

social and stress-related behaviours as well as FGC concentrations were set as dependent

variables in separate models, while species (Lar gibbon vs red-ruffed lemur) and enrichment

condition (before vs. after exposure to scent enrichment), along with their interaction, were

entered as independent factors in all models. We excluded from the analysis an outlier from

the faecal glucocorticoid data set. All models met the assumptions of homogeneity of

variance and normality of residuals.

Results

The 2x2 ANOVA model that included resting rates as the outcome variable revealed

only a significant main effect of Species, with red-ruffed lemurs resting more frequently than

Lar gibbons, but there was no significant main effect of Enrichment condition or the

interaction between Enrichment and Species on resting rates (Table 2).

Table 2. Results of the 2x2 ANOVA testing the effect of Enrichment condition, Species and their interaction on resting rates. Significant result is shown in bold.

Dependent variable	Independent variables	df	Sum square	F-value	P-value	Partial η^2
Resting	Enrichment	1	0.003	0.48	0.499	0.17
	Species	**1**	**0.266**	**51.23**	**< 0.001**	**0.8**
	Enrichment * Species	1	0.00	0.00	0.999	< 0.01
	Residuals	13	0.07			

By contrast, we found a significant effect of the interaction between enrichment condition and species on rates of social interaction (Table 3). Follow-up post-hoc t-tests revealed that, while Lar gibbons increased their rates of social interactions following their exposure to lavender (t(3.2) = 7, p = 0.005, d = 5.7, Figure 1), red-ruffed lemurs showed a reduction in rates of social interactions after being exposed to lavender (t(8.3) = 2.8, p = 0.02, d = 1.6, Fig. 1).

Table 3. Results of the 2x2 ANOVA testing the effect of Enrichment condition, Species and their interaction on social rates. Significant result is shown in bold.

Dependent variable	Independent variables	df	Sum square	F-value	P-value	Partial $\eta2$
Social	Enrichment	1	0.002	3.04	0.10	0.22
	Species	1	0.001	1.65	0.22	0.11
	Enrichment * Species	**1**	**0.005**	**6.46**	**0.02**	**0.33**
	Residuals	13	0.01			

Fig. 1. Effect of enrichment condition on rates of social interactions in red-ruffed lemur and Lar gibbon. Height of the boxes represents mean while whiskers represent standard error.

Rates of stress-related behaviours were significantly predicted by enrichment

condition but not by species or the interaction between species and enrichment condition

(Table 4). More specifically, both species reduced their rates of stress-related behaviours

after they were exposed to lavender (Fig. 2).

Table 4. Results of the 2x2 ANOVA testing the effect of Enrichment condition, Species and

their interaction on rates of stress-related behaviours. Significant result is shown in bold.

Dependent variable	Independent variables	df	Sum square	F-value	P-value	Partial η2
Stress-related behaviours	**Enrichment**	**1**	**0.0013**	**5.42**	**0.037**	**0.30**
	Species	1	0.0001	0.23	0.641	0.02
	Enrichment * Species	1	0.0001	0.41	0.533	0.03
	Residuals	13	0.0032			

Fig. 2. Effect of enrichment condition on rates of stress-related behaviours in both red-ruffed

lemur and Lar gibbon. Height of the boxes represents mean while whiskers represent

standard error.

Finally, we found that enrichment condition significantly predicted FGCs (Table 5):

following the exposure to lavender, both species displayed a significant increase in FGC

concentration, compared to before being exposed to the scent enrichment (Fig. 3).

Table 5. Results of the 2x2 ANOVA testing the effect of Enrichment condition, Species and

their interaction on FGC concentration. Significant result is shown in bold.

Dependent variable	Independent variables	df	Sum square	F-value	P-value	Partial η2
FGC	**Enrichment**	**1**	**9.89**	**11.23**	**0.002**	**0.26**
	Species	1	2.06	2.34	0.135	0.06
	Enrichment * Species	1	1.13	1.28	0.265	0.03
	Residuals	36	31.72			

Fig. 3. Effect of enrichment condition on FGC concentration in both red-ruffed lemur and

Lar gibbon. Height of the boxes represents mean while whiskers represent standard error.

Discussion

The effects of olfactory enrichment have been tested on several domestic, farm,

laboratory and zoo-housed animals [29,30]. However, olfactory stimulation is still one of the

least studied forms of enrichment (reviewed in [31]). In addition, there are mixed and

conflicting conclusions regarding the benefits of olfactory enrichment on animal welfare, and

this is particularly the case of primate species among which the overall effects of olfactory

enrichment is currently unclear and understudied (reviewed in [22]).

In this study, we found a significant effect of the scent enrichment on social

interactions, with red-ruffed lemurs showing a reduction and Lar gibbons exhibiting an

increase in rates of social interactions after being exposed to lavender. Generally, several

studies have found that scent enrichments may be effective at increasing active behaviours in

individuals in zoo-housed animals, such as cheetahs (*Acinonyx jubatus*) [9], Californian sea

lions (*Zalophus californianus*) [13] and Rothschild giraffes (*Giraffa camelopardalis*

rothschildi) [12] but not in meerkats (*Suricata suricatta*) [15]. Regarding primates, Gronqvist

and colleagues [32] showed that, among captive Javan gibbons (*Hylobates moloch*), olfactory

enrichment significantly increased the frequency of natural species-specific behaviours,

although individuals' interest in the olfactory enrichment decreased rapidly after the first day.

By contrast, Baker et al. [16] tested the effects of olfactory enrichment on ring-tailed lemurs

(*Lemur catta*) and found no significant effects on individuals' behaviours in the presence of

the odours, a result which was also shown in a study with essential oils on gorillas (*Gorilla*

gorilla gorilla) [14].

The opposite effect that scent enrichment exerted on rates of social interactions in the

two study species, with increased rates of social behaviour in Lar gibbons and decreased rates

of social behaviour in red-ruffed lemurs following exposure to the scent enrichment, might be

related to the fact that these two species spent most of their time in different types of social

behaviour. More specifically, Lar gibbons spent 72% of their social time playing, while red-

ruffed lemurs spent 78% of their social time grooming. A wide range of studies has shown

that frequency of play behaviour is strongly affected by the levels of exposure to stress the

animal is experiencing. In rats (*Rattus rattus*), for example, prenatal exposure to a stressor

significantly increases the latency to play [33]. Similarly, studies on both squirrel monkeys

(*Saimiri sp.*), rhesus macaque (*Macaca mulatta*) and common marmoset (*Callithrix jacchus*)

showed that play rates were higher in individuals who experienced low stress levels [34–36].

In this context, our findings that Lar gibbon spent more time playing after the introduction of

the scent enrichment compared to before the exposure to lavender might reflect the fact that

this species might have experienced lower stress levels after the exposure to the scent, which

is consistent with our prediction. On the other hand, social grooming, a common affiliative

behaviour in primates, has long been shown to have a stress-reducing effect. Work conducted

on macaques has shown, for example, that either receiving or giving grooming lowers stress

levels in the individuals involved in the interaction [36–42]. We, therefore, suggest that red-

ruffed lemurs reduced their rates of social grooming following the exposure to the scent

enrichment because the presence of lavender might have decreased their overall stress levels.

As we predicted, we found a significant effect on the scent enrichment on stress-

related behaviours, with both red-ruffed lemurs and Lar gibbons reducing their rates of stress-

related behaviours after they were exposed to lavender. This is consistent with our prediction

that scent enrichment reduces behavioural indicators of stress (i.e., pacing and self-

scratching). Similar results were reported in other species. Uccheddu and colleagues [43], for

example, exposed domestic dogs to a variety of essentials oils, and found that some scents

increased frequencies of behavioural indicators of relaxation while others decreased

behavioural indicators of stress such as pacing and over-grooming. Similarly, a study on

cheetahs and Sumatran tigers (*Panthera tigris sumatrae*) found that stereotypic pacing

behaviour was significantly decreased in the presence of a hay ball with cinnamon [44].

309   Finally, our finding that both red-ruffed lemurs and Lar gibbons displayed a

significant increase in FGCs following the exposure to lavender is at first puzzling and in

direct opposite to our findings related to the behavioural indicator of stress (see above). This

result, therefore, suggests that changes in behavioural indicators of stress such as pacing and

self-scratching do not necessarily reflect physiological indicators of stress, such as FGCs.

This is consistent with the work conducted by Higham and colleagues [45] showing that,

among olive baboons (*Papio anubis*), both short-term (i.e., day-to-day) and long-term

variation in rates of self-directed variation does not reflect changes in FGCs. This mismatch

might be due to the fact that self-directed behaviours and FGCs may reflect two different

types of stress. More specifically, self-directed behaviours likely reflect low-level acute

stress, or anxiety [46], while FGCs likely reflect high-level chronic stress. Accordingly, self-

directed behaviours have been found to increase in anxiety-inducing context, such as when

animals are given anxiogenic drugs [47], or after aggression [48]. On the other hand, FGC

levels have been shown to increase when animals are exposed to high levels of stress, such as

in the presence of tourists [49] or when exposed to the odour of a predator [50]. Therefore,

this suggests that researchers should, ideally, use *both* behavioural *and* physiological

indicators of stress and anxiety in order to have a complete picture on the levels of stress an

animal experiences and that behavioural indicators of anxiety should not necessarily be

interpreted as indicator of glucocorticoid production.

The question remains, however, on why both red-ruffed lemurs and Lar gibbons

exhibited higher FGC levels after exposure to scent enrichment compared to before scent

enrichment was introduced. The main function of glucocorticoids is to mobilize the energy

into the bloodstream from storage sites inside the body. This energy can then be used by

tissues and organs, such as muscle, and brain. Consequently, glucocorticoids are generally

produced every time the body needs energy, such as during stress response, when the animal

might need energy, for example, to escape from a predator. However, there are other non-

stress-related contexts in which animal body might need energy, such as early in the morning

when an animal needs to start their physical activity. Accordingly, glucocorticoid levels tend

to peak early in the morning. Here we argue that FGCs increased after the exposure to

lavender because the introduction of the scent enrichment might have increased the animals'

general activity pattern (e.g., locomotion, foraging), which might have required an increase in

glucocorticoid production by the animals' body.

In general, it is crucial to consider the ecological/biological relevance of the scent

enrichment to the species as this is likely to affect the results. Many scents, including

lavender, are chosen based on their effectiveness in humans or domestic animals, but this

may not necessarily be appropriate for all animal species [22]. In particular, as previous

authors have suggested, important factors to consider for the implementation of novel

olfactory enrichment programmes are the delivery of the scent and the type of scents used

[16]. The majority of studies have used spices or essential oils rather than focusing on natural

or biological scents; however, as with all types of enrichment, the biology of the species

should be taken into account and its effectiveness should be continually monitored to inform

best practices. For example, several studies have suggested that the use of either natural

prey/predator odours or odours from conspecifics should be researched further [32]. Other

studies have suggested the use of diffusers as a delivery method may be more effective [16].

Additionally, further studies have indicated that scents could be used in a number of

combinations and introduced randomly to continue to add novelty to the enrichment

programme and avoid the problem of habituation [9].

Interestingly, we found that scent enrichment exerted both behavioural and

physiological effects on both red-ruffed lemurs and Lar gibbon. While the use of chemical

communication and olfactory signals in lemurs has long been established [51–53], apes are

commonly considered microsmatic. However, accumulating evidence seems to suggest that

apes may also rely on olfactory signals and chemical communication. Henkel and Setchell

[54], for example, recently showed that chimpanzees (*Pan troglodytes*) spend more time

sniffing the urine of outgroup individuals than that of group members. Similarly, accounts of

border patrolling among wild chimpanzees report how individuals are often seen sniffing the

ground when near the territory of a neighbouring group [55]. Our findings showing that scent

enrichment exerts both behavioural and physiological changes in Lar gibbon confirms that

olfactory communication plays a key role in social interaction and physiological regulation

not only in chimpanzees but also in other apes.

In conclusion, our preliminary study provides strong evidence of a beneficial effect of

scent enrichment on animal welfare. Following exposure to the scent enrichment, animals

spend more time play, and less time grooming and exhibit lower rates of stress-related

behaviours. Our finding that both red-ruffed lemurs and Lar gibbon increased their levels of

FGCs after exposure to lavender might be associated to an increase in activity after animals

were exposed to the scent enrichment. Future work will need to expand the investigation of

the effect of scent enrichment on animal welfare by focusing on a wider range of primate

species and on a bigger sample size.

Acknowledgements

We are grateful to Dudley Zoo and Castle (especially David Beeston, Chris Leeson, Pat

Stevens, and primate keepers) and Twycross Zoo (especially Mat Liptovszky, Manuela

Townsend, Freisha Patel, and primate keepers) for their support to the project and assistance

with sample collection. We thank Jemma Billingsley, Stephanie Courten, Ailie McAuley,

Alexandria Lott, Amy Quirke, Demi Jayne Sheppard, Henry Swain, and Luke Bryant for

collecting data (which were not included in this paper) on howler monkeys, Diana monkeys

and orangutans at Dudley and Twycross zoos. This research work was supported by the

Faculty of Science and Engineering, University of Wolverhampton (equipment & laboratory

consumables), and the Department of Veterinary Medical Sciences, University of Bologna

(laboratory analyses).

Author contribution

Conceived and designed the experiment: SV. Performed the experiment: SS, BT, RP, GM,

PAA. Analysed the data: SSKK. Wrote the paper: SV, SSKK, CY, EE, SF.

Supporting information caption

S1 Text. Dataset

The table includes both the raw data and the rates of resting, social and stress-related

behaviours as well as the faecal glucocorticoid concentration (pg/mg) exhibited by red-ruffed

lemur and Lar gibbon before and after exposure to the scent enrichment (lavender).

References

1.   Estrada A, Garber PA, Rylands AB, Roos C, Fernandez-Duque E, Di Fiore A,

et al. Impending extinction crisis of the world's primates: Why primates matter. Science

advances. 2017;3: e1600946.

2.   Mellor DJ, Hunt S, Gusset M. Caring for wildlife: the world zoo and aquarium animal

welfare strategy. WAZA Executive Office. 2015.

3.   Schulte-Hostedde AI, Mastromonaco GF. Integrating evolution in the management of

captive zoo populations. Evolutionary applications. 2015;8: 413–422.

4.   Lacy RC. Stopping evolution. Conservation genetics in the age of genomics. New

York: Columbia University Press New York; 2009. pp. 58–81.

5.   Meier K. Blue Eyes on Red Lists: Conservation and the Future of the Blue-eyed

Black Lemur. Award Winning Anthropology Papers, Macalester College. 2016.

6.   Carroll SP, Jørgensen PS, Kinnison MT, Bergstrom CT, Denison RF, Gluckman P, et

al. Applying evolutionary biology to address global challenges. Science. 2014;346: 1245993.

7.   Hosey GR. How does the zoo environment affect the behaviour of captive primates?

Applied Animal Behaviour Science. 2005;90: 107–129.

8.   Ben-Ari ET. What's New at the Zoo? Zoo biologists are taking a scientific approach

to improving the quality of life for captive animals. BioScience. 2001;51: 172–177.

9.   Quirke T, O'Riordan RM. The effect of a randomised enrichment treatment schedule

on the behaviour of cheetahs (*Acinonyx jubatus*). Applied Animal Behaviour Science.

2011;135: 103–109.

10.   Carlstead K, Shepherdson D. Effects of environmental enrichment on reproduction.

Zoo biology. 1994;13: 447–458.

11.   Danial Rioldi E. The influence of conservation breeding programs on animal

communication and behaviour–a literary review. Mälardalen University. 2013.

12. Fay C, Miller L. Utilizing scents as environmental enrichment: preference assessment

and application with Rothschild giraffe. Animal Behavior and Cognition. 2015;2: 285–291.

13. Samuelson MM, Lauderdale LK, Pulis K, Solangi M, Hoffland T, Lyn H. Olfactory

enrichment in California sea lions (*Zalophus californianus*): an effective tool for captive

welfare? Journal of Applied Animal Welfare Science. 2017;20: 75–85.

14. Wells DL, Hepper PG, Coleman D, Challis MG. A note on the effect of olfactory

stimulation on the behaviour and welfare of zoo-housed gorillas. Applied Animal Behaviour

Science. 2007;106: 155–160.

15. Myles S, Montrose VT. The effects of olfactory stimulation on the behaviour of

captive meerkats (*Suricata suricatta*). Journal of Zoo and Aquarium Research. 2015;3: 37–

42.

16. Baker B, Taylor S, Montrose VT. The effects of olfactory stimulation on the behavior

of captive ring-tailed lemurs (*Lemur catta*). Zoo biology. 2018;37: 16–22.

17. Rafacz ML, Santymire RM. Using odor cues to elicit a behavioral and hormonal

response in zoo-housed African wild dogs. Zoo Biology. 2014;33: 144–149.

18. Heymann EW. Scent marking strategies of New World primates. American Journal of

Primatology. 2006;68: 650–661.

19. Clark F, King AJ. A critical review of zoo-based olfactory enrichment. Chemical

signals in vertebrates 11. Springer; 2008. pp. 391–398.

20. Setchell JM, Vaglio S, Moggi-Cecchi J, Boscaro F, Calamai L, Knapp LA. Chemical

composition of scent-gland secretions in an Old World monkey (*Mandrillus sphinx*):

influence of sex, male status, and individual identity. Chemical Senses. 2010;35: 205–220.

21. Vaglio S, Minicozzi P, Romoli R, Boscaro F, Pieraccini G, Moneti G, et al. Sternal

gland scent-marking signals sex, age, rank, and group identity in captive mandrills. Chemical

senses. 2016;41: 177–186.

22.   Wells DL. Sensory stimulation as environmental enrichment for captive animals: a

review. Applied Animal Behaviour Science. 2009;118: 1–11.

23.   Borgerson C, Eppley TM, Patel E, Johnson S, Louis E, Razafindramanana J. Red

ruffed lemur (*Varecia rubra*). The IUCN Red List of Threatened Species 2020. 2020. Report

No.: 2307–8235.

24.   Brockelman W, Geissmann T. *Hylobates lar*. The IUCN Red List of Threatened

Species 2020

25.   Altmann S. The structure of primate social communication. Social communication

among primates. Chicago: University of Chicago Press; 1967. pp. 325–362.

26.   Fontani S, Vaglio S, Beghelli V, Mattioli M, Bacci S, Accorsi PA. Fecal

concentrations of cortisol, testosterone, and progesterone in cotton-top tamarins housed in

different zoological parks: relationships among physiological data, environmental conditions,

and behavioral patterns. Journal of Applied Animal Welfare Science. 2014;17: 228–252.

27.   Tamanini C, Giordano N, Chiesa F, Seren E. Plasma cortisol variations induced in the

stallion by mating. European Journal of Endocrinology. 1983;102: 447–450.

28.   R Core Team. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. 2020.

29.   Heitman K, Rabquer B, Heitman E, Streu C, Anderson P. The use of lavender

aromatherapy to relieve stress in trailered horses. Journal of Equine Veterinary Science.

2018;63: 8–12.

30.   Blackie N, de Sousa M. The Use of Garlic Oil for Olfactory Enrichment Increases the

Use of Ropes in Weaned Pigs. Animals. 2019;9: 148.

31.   Campbell-Palmer R, Rosell F. The importance of chemical communication studies to

mammalian conservation biology: a review. Biological Conservation. 2011;144: 1919–1930.

32.   Gronqvist G, Kingston-Jones M, May A, Lehmann J. The effects of three types of

environmental enrichment on the behaviour of captive Javan gibbons (*Hylobates moloch*).

Applied Animal Behaviour Science. 2013;147: 214–223.

33.   Takahashi LK, Haglin C, Kalin NH. Prenatal stress potentiates stress-induced

behavior and reduces the propensity to play in juvenile rats. Physiology & behavior. 1992;51:

319–323.

34.   Biben M, Champoux M. Play and stress: cortisol as a negative correlate of play in

Saimiri. Play Culture Stud. 1999;2: 191–208.

35.   Norscia I, Palagi E. When play is a family business: adult play, hierarchy, and

possible stress reduction in common marmosets. Primates. 2011;52: 101–104.

36.   Wooddell LJ, Hamel AF, Murphy AM, Byers KL, Kaburu SSK, Meyer JS, et al.

Relationships between affiliative social behavior and hair cortisol concentrations in semi-free

ranging rhesus monkeys. Psychoneuroendocrinology. 2017;84: 109–115.

37.   Aureli F, Yates K. Distress prevention by grooming others in crested black macaques.

Biology letters. 2010;6: 27–29.

38.   Shutt K, MacLarnon A, Heistermann M, Semple S. Grooming in Barbary macaques:

better to give than to receive? Biology Letters. 2007;3: 231–233.

39.   Boccia ML, Reite M, Laudenslager M. On the physiology of grooming in a pigtail

macaque. Physiology & Behavior. 1989;45: 667–670.

40.   Aureli F, Preston SD, de Waal F. Heart rate responses to social interactions in free-

moving rhesus macaques (*Macaca mulatta*): a pilot study. Journal of Comparative

Psychology. 1999;113: 59–65.

41.   Gust DA, Gordon TP, Hambright K, Wilson ME. Relationship between social factors

and pituitary-adrenocortical activity in female rhesus monkeys (*Macaca mulatta*). Hormones

and Behavior. 1993; 27: 318–331.

42.   Kaburu SSK, MacLarnon A, Majolo B, Qarro M, Semple S. Dominance rank and

self-scratching among wild female Barbary macaques (*Macaca sylvanus*). African Zoology.

2012;47: 74–79.

43.   Uccheddu S, Mariti M, Sannen A, Vervaecke H, Arnout H, Gutierrez RJ, et al.

Behavioural and cortisol responses of shelter dogs to a cognitive bias test after olfactory

enrichment with essential oils. Dog Behavior. 2018;2: 1–14.

44.   Damasceno J, Genaro G, Quirke T, McCarthy S, McKeown S, O'Riordan R. The

effects of intrinsic enrichment on captive felids. Zoo Biology. 2017;36: 186–192.

45.   Higham JP, MacLarnon AM, Heistermann M, Ross C, Semple S. Rates of self-

directed behaviour and faecal glucocorticoid levels are not correlated in female wild olive

baboons (*Papio hamadryas anubis*). Stress. 2009;12: 526–532.

46.   Maestripieri D, Schino G, Aureli F, Troisi A. A modest proposal: displacement

activities as an indicator of emotions in primates. Animal Behaviour. 1992;44: 967–979.

47.   Schino G, Perretta G, Taglioni AM, Monaco V, Troisi A. Primate displacement

activities as an ethopharmacological model of anxiety. Anxiety. 1996;2: 186–191.

48.   Schino G. Reconciliation in domestic goats. Behaviour. 1998;135: 343–356.

49.   Barja I, Silván G, Rosellini S, Piñeiro A, González-Gil A, Camacho L, et al. Stress

physiological responses to tourist pressure in a wild population of European pine marten. The

Journal of Steroid Biochemistry and Molecular Biology. 2007;104: 136–142.

50.   Monclús R, Rödel HG, Palme R, Von Holst D, De Miguel J. Non-invasive

measurement of the physiological stress response of wild rabbits to the odour of a predator.

Chemoecology. 2006;16: 25–29.

51.   Scordato ES, Dubay G, Drea CM. Chemical composition of scent marks in the

ringtailed lemur (*Lemur catta*): glandular differences, seasonal variation, and individual

signatures. Chemical Senses. 2007;32: 493–504.

52. Janda ED, Perry KL, Hankinson E, Walker D, Vaglio S. Sex differences in
scent-marking in captive red-ruffed lemurs. American Journal of Primatology. 2019;81:
e22951.

53. Hayes RA, Morelli T, Wright P. Anogenital gland secretions of Lemur catta and
Propithecus verreauxi coquereli: a preliminary chemical examination. American Journal of
Primatology: Official Journal of the American Society of Primatologists. 2004;63: 49–62.

54. Henkel S, Setchell JM. Group and kin recognition via olfactory cues in chimpanzees
(*Pan troglodytes*). Proceedings of the Royal Society B. 2018;285: 20181527.

55. Samuni L, Preis A, Mundry R, Deschner T, Crockford C, Wittig RM. Oxytocin
reactivity during intergroup conflict in wild chimpanzees. Proceedings of the National
Academy of Sciences. 2017;114: 268–273.

Figure 1

Figure 2

Figure 3