A CAR RNA FISH assay reveals functional and spatial heterogeneity of

chimeric antigen receptor T cells in tissue

Karsten Eichholz[1], Alvason Zhenhua Li[1], Kurt Diem[2], Semih U. Tareen[3], Michael C. Jensen[4,5],

Jia Zhu[1,2], Lawrence Corey[1,2,6*]

[1]Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Research Center, Seattle, WA,
USA.

[2]Department of Laboratory Medicine, University of Washington, Seattle, WA, USA.

[3]Work performed while employed at Juno Therapeutics, Seattle, WA; currently employed at Sana
Biotechnology, Seattle, WA, USA.

[4]Clinical Research Division, Fred Hutchinson Cancer Research Center (FHCRC), Seattle, WA,
USA.

[5]Ben Towne Center for Childhood Cancer Research, Seattle Children's Research Institute, Seattle,
WA, USA.

[6]Department of Medicine, University of Washington, Seattle, WA, USA.

*correspondence: L. Corey, MD, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave

N, MS E3-300, Seattle, WA 98190 (Lcorey@fredhutch.org).

One Sentence Summary: We developed a CAR RNA FISH assay to study chimeric antigen

receptor T cell trafficking and function in human and mouse tissue.

Short title: A CAR RNA FISH assay to study CAR T cells in tissues

Abstract

Chimeric antigen receptor (CAR) T cells are engineered cells used in cancer therapy and are

studied to treat infectious diseases. Trafficking and persistence of CAR T cells is an important

requirement for efficacy to target cancer and HIV sanctuary sites. Here, we describe a CAR RNA

FISH histocytometry platform combined with a dnnRRS image analysis algorithm to quantitate

spatial distribution and in vivo functional ability of a CAR T cell population at a single cell

resolution. In situ, CAR T cell exhibited a heterogenous effector gene expression and this was

related to the distance from tumor cells, allowing a quantitative assessment of the potential in vivo

effectiveness. The platform offers the potential to study immune functions engineered cells in situ

with their target cells in tissues with high statistical power and thus, as an important tool for

preclinical and potentially clinical assessment of CAR T cell effectiveness.

Impact statement

We developed an imaging platform and analysis pipeline to study large populations of engineered

cells on a single cell level in situ.

Introduction

Chimeric antigen receptors (CAR) T cells are engineered T cells that are under development to treat various cancers and infectious diseases. The CAR is a recombinant fusion protein that combines an extracellular antigen-specific single chain variable fragment (scFv) with a membrane domain and an intracellular signaling domain to promote targeted killing. The scFv is directed against extracellular antigens, provides high specificity and induces cytolysis when the CAR T cell interacts with its target. Recently, two CAR T cell therapies based on the anti-CD19 scFv clone FMC63 have been approved for treatment of B-cell acute lymphoblastic leukemia (B-ALL) and non-Hodgkin lymphoma and numerous constructs are in pre-clinical and clinical development for a variety of hematological and solid tumors.

CAR T cell-based therapies are also in pre-clinical development to target the viral reservoir in individuals infected with the human immune deficiency virus-1 (HIV-1) to provide a functional cure. Anti-HIV CAR T cells either use a modified version of CD4, the cell surface protein responsible for HIV attachment, or a scFv based on various HIV envelope protein gp120-specific broadly-neutralizing antibodies, e.g. VRC01, VRC07-523, or PGT128[1–3].

While CAR T cell therapies have shown some success in hematological cancers, the application in solid tumors settings is still hampered by the immunosuppressive tumor microenvironment, poor tumor infiltration, the choice of an efficient scFv-tumor antigen pair without off-target effects and adverse effects like cytokine release syndrome[4–6].

CAR T cell performance, proliferation and persistence in preclinical in vivo studies and clinical application are mainly studied by flow cytometry, quantitative PCR, and for preclinical cancer mouse models only, whole body bioluminescence imaging of the tumor burden. These techniques are favorable to measure CAR T cell phenotype, functionality and expansion with high statistical

power and specificity and can assess global anti-tumor efficacy, however, the location of the CAR

T cells in the tissue with respect to the tumor cells cannot be determined.

To track CAR T cells by flow cytometry *in vitro* and *in situ,* various polypeptide tags like truncated

epidermal growth factor receptor (EGFRt) are co-expressed with the CAR*(7)*. These polypeptide

tags, like EGFRt, are not expressed on non-engineered T cells and show low immunogenicity.

EGFRt has been successfully used to study CAR T cells in vivo by flow cytometry in mice, humans

and nonhuman primates*(7–9)*, and the anti-EGFR mouse monoclonal antibody clone 31G7 has

been used to stain for CAR T cells in tissue *(7)*. However, to study CAR T cell trafficking and

function *in situ*, the use of anti-EGFRt antibodies is complicated by the ubiquitous expression of

EGFR in various healthy and tumor tissues as well as the limited possibility to multiplex different

target proteins available for fluorescence microscopy compared to multi-color flow cytometry. An

alternative technique is to use RNA *in situ* hybridization (RNA ISH) to track the CAR T cells.

Recently, RNA ISH approaches have been successfully applied to track CAR T cells in patients

with glioblastoma*(10)*, in a case study in a patient with B-ALL, who died 5 days post-anti-CD19

CAR T cell infusion*(11)* and in the clinical samples from the ZUMA-1 trial that tested the anti-

CD19 CAR Axicabtagene ciloleucel *(12)*.

Automated image analysis requires reliable image segmentation, however, commonly used

commercial and open source software mainly rely on nuclei detection and watershed algorithms.

The weak or discontinuous cell membrane staining, overlapping cells tissue- or fixation-derived

autofluorescence and other fluorescent artifacts have limited the use of current software packages

for automated image analysis. To overcome the limitations, stochastic algorithms that use hidden-

Markov-model (HMM) statistics proved to be useful to tackle the more advanced requirements of

automated image analysis in tissue *(13–17)*. More recently, we developed a robust and efficient

Random-Reaction-Seed method (RRS) that employs the maximum likelihood method to estimate

probabilistic functions of Markov chains which showed efficient tracking of neurite net-like

structures in fluorescence microscopy images *(18)*.

This article describes a dnnRRS-assisted RNA *in situ* hybridization assay to track and phenotype

tumor cells and anti-CD19 and anti-HIV envelope CAR T cells in tissues. We use this technique

to detect CAR T cells localization in tissue in a quantitative manner in both humans and preclinical

animal models. Our data indicate that proximity to tumor tissue influences in vivo killing potential

as defined by in situ IFNγ and GZMB mRNA expression; thus, providing direct in situ evidence

of in vivo activity and illustrating the importance of spatial interactions between the tumor

microenvironment and in vivo functional activity of adoptively transferred T cells The CAR RNA

FISH imaging platform can be easily modified for novel CAR-antigen pairs and the study of the

interactions between CAR T cell and the infiltrating tumor microenvironment and provide both

preclinical and clinical insights in determinants of potential CAR T cell effectiveness of this

emerging cellular therapy platform using commonly available technologies.

Material and methods

Cell culture

TM-LCL, Jurkat and VRC07-523 CAR Jurkat cells were cultured in RPMI1640 (Gibco), 10%

fetal bovine serum and antibiotics. The VRC07-523 CAR Jurkat cell line was a kind gift from

Hans-Peter Kiem.

CAR T cells expansion

Anti-CD19 CD8 CAR T cells were cultured with TM-LCL feeder cells as previously described

with minor modifications[19]. Briefly, 10^6 anti-CD19 CD8 CAR T cells were expanded in

presence of 7 x 10^6 irradiated (8000 cGy) TM-LCL feeder cells in RPMI1640 containing 50 IU/mL

human recombinant IL-2 (Peprotech), 10 ng/ml recombinant human lL-15 (Peprotech), 10% fetal

bovine serum, 50 µM β-mercaptoethanol (SigmaAldrich) and 25 mM Hepes, pH 7.4 (Gibco) and

antibiotics.

Tissue samples

The mouse tissues used for the CAR RNA FISH in situ validation contain a BE2 cell

neuroblastoma xenograft and were described in a previous study[20].

Flow cytometry

Day 12 CD8 CAR T cells TM-LCL co-cultures were stained with LIVE/DEAD Fixable Aqua

Dead Cell Stain (ThermoFisher) followed by antibody staining against CD8 (FITC, clone RPA-

T8, BD, 555366), CD19 (BV786, clone SJ25C1, biolegend) and EGFRt (PE, clone erbitux,

provided by Juno Therapeutics) and analyzed on a LSR II flow cytometer. Gating on CD8+

EGFRt+ cells was done with fluorescence minus one-controls for CD8 and EGFRt.

CAR RNA FISH – probes

zz probes for CAR mRNA were custom-made by Acdbio and are directed against the scFv, the

signaling domain-coding region and in some cases the 3'UTR for CAR detection in tissues that

124 are devoid of any other lentiviral constructs that may harbor the wPRE (Fig. 1A, Fig. S1A). The

125 individual probes were compared *in silico* with the human, mouse and macaque (*M. mulatta* and

126 *M. nemestrina*) genome to avoid cross-detection of endogenous counterparts of the CAR (e.g.

127 CD3zeta).

128 **CAR RNA FISH sample preparation – cytospins**

129 To prepare cytospin slides to validate the probe specificity, 10^7 anti-CD19 CD8 CAR T cells or

130 Jurkat cells were washed twice with 40 mL PBS and centrifuged at 250 x g for 10 min. The cell

131 pellets were resuspended in 10 mL 4% paraformaldehyde, PBS, fixed at 37°C for 1 h, and washed

132 two times with 10 mL PBS. The cell pellet was then resuspended at 10^6 cell/mL in 70% ethanol

133 and stored for up to 7 days for further cytospin processing.

134 To obtain different densities of CAR T cells, CAR T cells were mixed with Jurkat cells at different

135 ratio (0 – 100%). The cell suspensions were then centrifuged onto RNAse-free Superfrost plus

136 glass slides (Electron microscopy sciences) with a cytospin centrifuge (cytospin 3,

137 ThermoShandon) set to 800 rpm for 10 min to prepare the cytospins.

138 The glass slides were air dried for 20 min at ambient temperature and dehydrated stepwise in 50%,

139 70% and absolute ethanol for 5 min and stored in absolute ethanol at -20ºC until further processing.

140 **CAR RNA FISH sample preparation – tissues**

141 Fresh frozen OCT-embedded tissues blocks were sectioned to 10 μm thickness onto RNAse-free

142 Superfrost plus glass slides (Electron microscopy sciences) on a Leica CM1950 cryostat at -20°C.

143 The slides were kept at -20°C for 1 h and transferred to -80°C or directly processed for FISH.

144 OCT sections were incubated in 4% PFA, PBS pre-chilled to 4°C for 15 min, washed 2 times

145 briefly with PBS and dehydrated in a graded series of ethanol for 5 min each (50%, 70%, 2 x

146 100%).

FFPE tissue blocks were sectioned to 4 μm thickness on a Leica RM2255 microtome. Sections

were heated at 60ºC for 1 h, dewaxed and processed for RNAscope pretreatment on a Leica Bond

RX according to manufacturer's recommendations with Leica Bond Epitope Retrieval Solution 2

(ACDbio) for 15 min at 95ºC. Subsequently, FFPE sections were processed manually as described

in the following section.

Fluorescent CAR RNA FISH ISH procedure

Cytospin slides or tissue section slides were removed from 100% ethanol after dehydration and

air-dried for 30 min at 40°C in a hybridization oven or 5 min at ambient temperature, respectively.

A hydrophobic barrier was created around the sample on the slides, air dried for 1 min, treated

with H_2O_2 for 10 min (for RNAscope® Multiplex Fluorescent Assay version 2) followed with

protease (30 min protease III at 40°C for cytospin samples, 15 min protease IV at ambient

temperature for mouse tissue sections in OCT-embedded tissues, or 30 min protease plus at 40°C

for FFPE tissues) (ACDbio).

After protease digestion, samples were washed twice in PBS and anti-RNA zz probes against CAR,

CD4, CD8, IFNγ, GZMB, PPIB and DapB were hybridized at 40°C for 2 h (Table 1 suppl.). As a

negative control, the slides were treated with 4 μg/mL RNAase A and 100 U/mL RNAse T1

(RNase A/T1 mix, ThermoFisher) for 30 min between the protease digestion and the probe

hybridization step. For the signal amplification after the probe hybridization, we used initially the

RNAscope Multiplex Fluorescent Assay version 1 with the fluorophores Alexa 488 and Atto 550

and in later experiments the RNAscope Multiplex Fluorescent Assay version 2 with the

fluorophores Opal 570 and Opal 650 (PerkinElmer) according the manufacturer's protocol for

multiplex fluorescent in situ hybridization kit with minor modifications. We added an additional

counter staining for the plasma membrane before the nuclear counter staining or omitted the

nuclear counterstaining completely. In brief, the sample slides were washed 2 times for 2 min with

PBS, incubated with 10 µg/mL wheat germ agglutinin conjugated with Texas Red-X or Oregon

Green 488 (Invitrogen) for 10 min at ambient temperature, washed again 2 times for 2 min with

PBS before proceeding to the nuclear counter staining step.

Digital droplet PCR detection of CAR T cells in tissues

gDNA and RNA were extracted with QiAmp DNA kit (Qiagen) and RNeasy Mini extraction kit

(Qiagen), respectively. 10 µL of DNA or RNA were used for ddPCR.

Primer sequences were as follows for wPRE (5′- CGCAACCCCCACTGGTT -3′ [wPRE forward],

5′-AAAGCGAAAGTCCCGGAAAG-3′ [wPRE reverse] and 5′- FAM-CCACCACCTGTCAGC-

MGB-3′ [wPRE probe]) and the CAR ORF (5′- AGCTGCCGATTTCCAGAAGA -3′ [CAR

forward], 5′- CCAGGTTCAGCTCGTTGTACAG -3′ [CAR reverse] and 5′- FAM-

TCAGCAGAAGCGCCGACGCC-QSY -3′ [CAR probe])

The DNA ddPCR was used with ddPCR Supermix for probe (Biorad) and the PCR conditions

were 95°C for 10 min, then 40 cycles at 94°C for 30 s, 60°C for 1 min followed by 98°C for

10min. The RNA-ddPCR was used with One-Step RT-ddPCR Advanced kit for Probes (Biorad).

The thermal conditions were 50°C for 60 min for reverse transcription, 95°C for 10 min, then 40

cycles at 94°C for 30 s, 60°C for 1 min for the ddPCR followed by 98°C for 10 min.

Immunohistochemistry

To combine IHC with the CAR RNA FISH procedure we followed a previously described protocol

with minor modification*(21)*. After the last RNA target was developed (step Amp4 in the

RNAscope Multiplex Fluorescent Assay version 1, TSA amplification step in the RNAscope

Multiplex Fluorescent Assay version 2), the cells and tissue sections were washed with PBS and

blocked with 5% donkey or goat serum (Jackson ImmunoResearch), 2% BSA (Gemini Bio-

Products) +0.2% Triton X-100 (SigmaAldrich) in 5X Casein (Vector Laboratories) for 30 min,

the mouse anti-huCD19 antibody (dilution 1:100,Clone ID: UMAB103, origene) overnight at 4°C

for the mouse xenograft tissues.

Slides were washed, incubated with secondary donkey anti-mouse IgG-Alexa 647 or Alexa 488

(1:100 Molecular Probes/ThermoFisher Scientific) or a goat anti-mouse BV480 (Becton

Dickinson) for 1 hour at room temperature, and washed 2 times for 5 minutes in PBS + tween

(0.05% v/v). Subsequently, samples were counterstained with DAPI and mounted with ProLong

Gold antifade (Thermofisher).

Image acquisition

Images were acquired with a PerkinElmer Ultraview Vox spinning disk confocal microscope

equipped with 4 laser lines 405, 488, 561 and 640 nm and a Hamamatsu ImageEM EMCCD

camera (512 x 512 pixels, 16 micron pixel size, 90% Quantum efficiency) with a 40 x objective, a

Leica SP8 confocal microscope equipped with laser lines at 405 nm, 440 nm, and adjustable white

light laser, or a Nikon eclipse Ti epifluorescence microscope equipped with a sola light machine,

a Hamamatsu ORCA-Flash4.0 LT+ C11440 (2048 ×2048 pixels, 82 % Quantum efficiency)

and Plan Apo 10x 0.45, Plan Apo 20x 0.75 and Plan Fluo 40x 1.3 objectives. Images that

were analyzed with dlRSS algorithm were not modified post-acquisition unless otherwise stated

in the description of the algorithm. Images for publication figures that were acquired with the

spinning disk confocal microscope were size-adjusted to 2048 x 2048 pixels and z-stacks were

projected with the Max intensity function in the Fiji software package*(22)*.

A Nikon Eclipse E800 with a Megapixel Nikon DigitalSight DS 5Mc color camera was used for

CAR in situ hybridization microscopy in human specimen.

CAR T cell detection algorithm

Imaging data was analyzed on a GPU workstation with a 32 core CPU, 64 GB RAM and a

QUADPRO P5000 GPU. To perform image segmentation based on WGA-plasma membrane

staining, we extended a previously described algorithm for identification of neurite net-like

structures and to generate the ground truth automatically for each image *(18)*. It is a robust and

efficient Random-Reaction-Seed method (RRS) that employs the maximum likelihood method to

estimate probabilistic functions of Markov chain. A U-net-based auto encoder decoder

convolutional deep neuronal network*(23)* performs image segmentation and phenotyping based on

the presence of nuclei, RNA and/or protein.

To detect the cell membrane two subsequent steps are performed by the algorithm. First, it extracts

a net-like draft of the cellular border based on the dnnRRS results and second, it performs cell-

loop segmentation of the net-like draft and thus, also removes false positive membranes from the

draft. Each tile of a tile scan is analyzed separately and to avoid any redundancy of areas that

overlap on the border of two adjacent images a 48-pixel wide area was omitted from the analysis

on each side of the tile. This equals to (4 x 48 x 2048 pixels)/(2048 x 2048 pixels) = 0.09 or 9% of

each slide that were not used for analysis. To overcome the optical vignetting effect in the corners

of the images were normalized using a Gaussian function.

For RNA FISH and CD19 IHC detection, a conventional local-maxima method is used and are

then combined with the membrane map. For the comparison of GZMB and IFNγ expression in

CAR T cells in the CD19high and CD19low areas, CD19high and CD19low areas were defined

by generating a circle for each CD19+ cells that corresponds to the actual area covered by the cell

at its x y position. The circles were than expanded approximately 16 times the radius of the circle

and connective groups were created by overlapping the expanded circles. The largest connective

group was denoted as CD19high area whereas the remainder of the tissue section was defined as

CD19low area. The spatial density of CAR+ GZMB-, CAR+ GZMB+, CAR+ IFNγ -, CAR+ IFNγ

+ was than calculated for the CD19high and CD19low area of each individual tissue section.

The RNA and protein expression levels as well as the location of the cells was exported as a

comma-separated values file and loaded into the flow cytometry data analysis software FlowJo 10

(Becton Dickinson). In FlowJo, the imaging data was analyzed with the gating and plotting tools.

To compare cell counts of the dnnRRA and manual operators, manual operators used the cell

counter function in imageJ to track cell counts.

Statistical analysis

The data were analyzed with graphpad prism software with the statistical tests stated in the figure

legends. One-way ANOVA combined with a Tukey test was used to compare multiple groups. *,

, *, **** correspond to $p < 0.05$, $p < 0.01$, $p < 0.001$ and $p < 0.0001$, respectively. Paired t

test was used to compare the spatial density within the CD19high and CD19low areas. *, **, ***,

**** correspond to $p < 0.05$, $p < 0.01$, $p < 0.001$ and $p < 0.0001$, respectively.

Box and whiskers plots show the median, the 5th to 95th percentile for the whiskers and the 25th

to 75th percentile for the box.

Results

Rationale for CAR RNA FISH probe design

We designed the CAR RNA probes to be directed against the wPRE in the 3'UTR and the scFv of

the anti-CD19 CAR FMC63 and the anti-HIV env CAR VRC07-523 (Fig. 1A, Fig. 1A suppl.).

The scFv of the CAR is based on the variable region of the two mRNA species that translate to the

Ig heavy and light chain of an antibody. To rule out that the probes against the scFv cross-hybridize

with a large number of B cells in potential target tissues we stained human tonsil sections with a

pan-IGHG RNA FISH or a CAR FMC63 15ZZ probes. IGHG RNA FISH showed an ubiquitous

and bright staining throughout the tissue section (Fig. 2A suppl). By contrast, the CAR FMZ 63

15ZZ probes stained only a minority of cells in the tonsils tissue (Fig. 2B suppl. and inlet) and this

signal could be abrogated by RNAse pretreatment prior to CAR FMZ 63 15ZZ RNA FISH (Fig.

2C suppl.). We concluded that the number of B cells that stained false-positive for CAR FMZ 63

15ZZ RNA FISH was sufficiently small to optimize the CAR RNA FISH approach in combination

with various T cell markers for the anti-CD19 CAR based on the mouse monoclonal antibody

clone FMC63*(24)* and the anti-HIV envelope CAR based on the broadly-neutralizing antibodies

VRC07-523*(25)* and combined it with a recently published image segmentation and analysis

algorithm*(18)*.

CAR RNA FISH detects CAR T cells in *in vitro* cultures

For *in vitro* experiments, T cells were co-cultured with γ-irradiated CD19-expressing TM-LCL

lymphoblastoid cells for 12 days and purity was assessed by flow cytometry (~90%) (Fig. 3 suppl.).

To validate the CAR FMC63 26ZZ probes, anti-CD19 CD8 CAR T cells or Jurkat cells were

processed for cytospins and stained for CAR mRNA (Fig. 1B). The anti-CD19 CD8 CAR T cells

stained positive with the CAR FMC63 26ZZ probe set. By contrast, Jurkat cells stained with the

CAR FMC63 26ZZ probe set were negative for CAR RNA spots. RNA FISH against housekeeping

gene PPIB (Peptidyl-prolyl cis-trans isomerase B) and the bacterial gene DapB as positive and

negative controls, respectively, for RNA integrity and probe specificity, yielded characteristic

RNA FISH staining only in cells with relevant genes (Fig. 1B).

RNA FISH-based imaging techniques detect the target mRNA mostly in the cytosol that surrounds

the nucleus as demonstrated with the CAR mRNA (Fig. 4A suppl.). The density of tissue impedes

automated image segmentation to define the quantity of T cells expressing the majority of the

extranuclear CAR mRNA and cannot be defined based on the nuclear DAPI staining. Traditional

algorithms rely on nuclei detection and assume a slightly extended perimeter for the cytoplasm to

detect the RNA. To define cellular boundaries and to facilitate computational image segmentation,

we used fluorescently labeled wheat germ agglutinin as a second counterstain to label N-acetyl-D-

glucosamine and sialic acids on cell surface glycoproteins (Fig.1C). This allowed to analyze the

images with a modified version of a random reaction seed algorithm recently published by our

group *(18)* (Fig. 1D). In brief, the acquired images are partitioned and, several 1000 seeds are

distributed on the WGA plasma membrane stain images (Fig 1D). The algorithm searches for the

plasma membrane high fluorescence intensity signal, follows the signal using hidden Markov

model statistics and tries to close a loop, which would represent a cell and thus, creates a membrane

map*(18)*. Once the cell circumference has been determined, the algorithm uses classical Laplacian

of Gaussian local maxima detection for nuclear and RNA staining within this perimeter (Fig. 1D).

The output data contains the location and RNA expression levels of each individual cell of the

input image (Fig. 1D).

To validate the CAR RNA FISH approach in combination with the dnnRRS algorithm, we stained

anti-CD19 CD8 CAR T and Jurkat cells mixed at different ratio (1, 0.75, 0.5, 0.25 and 0) with the

CAR FMC63 26zz probe set and acquired images. The number of cells stained positive by CAR

RNA FISH and the number of cells detected with dnnRRS increased with incrementing number

of input anti-CD19 CD8 CAR T cells (Fig. 1E). The input CAR T cells were 90% EGFRt+ by

flow cytometry (Fig 1A suppl.), hence, we expected to detect ~ 90%, 67.5%, 45%, 22.5% and 0%

CAR+ cells for the corresponding ratio of 1, 0.75, 0.5, 0.25 and 0. The dnnRRS algorithm detected

~5 – 60% CAR+ cells in the CD19 CAR T cell data set. There was a strong correlation between

detection of the algorithm and the expected values (R^2=0.95) (Fig. 1F) and a significantly higher

number of CAR RNA dots/cells compared to Jurkat cells alone (Fig. 1G). We chose the same

approach to validate CAR RNA FISH for a VRC07-523 CAR in a VRC07-523 CAR Jurkat cell

line and confirmed that it specifically stained these VRC07-523 CAR Jurkat cells but not control

Jurkat cells (Fig. 5A suppl.). In addition, in serial dilution experiments an increasing number of

cells stained positive for the VRC07-523 CAR RNA and this was detectable by dnnRRS (Fig. 5B

suppl.). The ratio of detected cells by dnnRRS (7- 68%) correlated with the expected values by

flow cytometry (0-97%) (R^2=0.98) (Fig. 5C suppl.) and we observed a significantly higher number

of CAR RNA dots/cell between the VRC07-523 CAR Jurkat cells at each ratio and non-transduced

Jurkat cells (Fig. 5D suppl.). Combined, these data indicate that our CAR RNA FISH probe design

and dnnRRS image analysis is highly specific and quantitative for detection of CAR T cells in cell

culture samples.

CAR RNA FISH detect CAR T cells *in situ* in mouse xenografts

To validate CAR RNA FISH for *in situ* applications in preclinical animal models, we used a BE2

neuroblastoma xenograft model of anti-CD19 CAR T cell-infused mice or control mice without

CAR T cell infusion*(20)*. The BE2 cell line was transduced with multiple lentiviruses to express

CD19 and luciferase and may contain the common wPRE sequence. We performed digital droplet

PCR with primers directed against the signaling domain and the wPRE in genomic DNA. We

detected the wPRE portion in both CAR+ and CAR- BE2 tumor tissue whereas the CAR signaling

domain was only present in CAR+ BE2 tumor tissues (Fig. 6 suppl.). Therefore, to exclude false-

positive cell detection due to cross-hybridization artifacts during the *in situ* setup of the CAR RNA

FISH on mouse xenograft tissues, we only used a reduced probe set of total 15 zz probes, coined

FMC63-15ZZ, and omitted the wPRE probes (Fig. 1A suppl.).

We observed a characteristic RNA FISH pattern with the CAR RNA FISH in CD19+ BE2

xenograft tissues from mice infused with anti-CD19 CAR T cells but not from control mice (Fig.

2A). To ensure that mRNA integrity was of sufficient quality in the mouse xenograft and the RNA

FISH staining procedure gave only low unspecific staining, we used probes against the house

keeping gene PPIB (Fig. 7A suppl.) and the bacterial gene DapB (Fig. 7B suppl.), which gave a

ubiquitous staining pattern or did not result in any positive signal in the corresponding area on

adjacent slides, respectively. To further confirm the specificity of the CAR RNA FISH signal, no

FISH signal could be observed when we omitted the FMC63 15ZZ probe set (Fig. 7C suppl.), or

after pretreatment of CAR T cell-positive tissues with RNAse A/H (Fig. 7D suppl.). These data

indicate that the CAR RNA FISH approach can be applied for CAR T cell detection in formalin

fixation and paraffin embedded tissues. We performed similar experiments in CAR T cell-positive

and -negative mouse xenograft tissue section that went through the process of fresh-frozen OCT

embedding and obtained similar results (data not shown).

To quantify the RNA expression of the CAR and any other mRNA of interest on a per cell basis

on the complete tissue section we used a novel and unique deep neuronal network random reaction

seed (dnnRRS) image analysis algorithm to perform computational image analysis *(18)*. WGA, a

lectin that binds N-acetyl-D-glucosamine and sialic acids on ubiquitous glycoproteins on the cell

surface allowed us to reliably define the cell boundaries of each cell in the tumor section (Fig. 2B).

The dnnRRS algorithm uses local fluorescence maxima detection for nuclei and RNA and random

reaction seed WGA-based image segmentation to quantify the spatial distribution and frequency

any mRNA of interest on a per cells and the data can be exported and analyzed in flow cytometry

analysis software like Flowjo by the individual researcher (Fig. 2C). To exemplify the performance

of the dnnRRS algorithm, we chose 10 images from our data set (Fig. 8A suppl.) that showed

typical fluorescence intensities for nuclei and membrane staining for different imaging situations.

Three operators counted the total cell number manually in imageJ based on membrane staining

and nuclei staining and the dnnRRS approach (Fig. 8B suppl.). The observed values obtained by

manual counting or dnnRRS correlated across the example images (Pearson's r<0.9 for each of

the operators) (Fig. 8C suppl.). The average time required per image was 15:31 min ± 8:02 min

for manual counting and 1:28 min ± 0:18 min for the dnnRRS algorithm (Fig. 8D suppl.). In

addition to the total number of cells, the dnnRRS algorithm also segments and quantifies RNA

FISH and IHC fluorescence data for T cell- or tumor-related genes, x-y position of the individual

cell and cellular borders and integrates and stitches adjacent micrograph images. Notably, the

image data sets recorded for the individual tissue sections frequently exceeded 300 multichannel

images per sections (data not shown) and we were thus able to interrogate multiplexed mRNA

expression levels in 150000 to 250000 cells per tissue section on a per cell basis.

We analyzed image data sets of a CAR RNA FISH or PPIB RNA FISH combined with

immunohistochemistry for CD19 from CAR T cell-positive and negative BE2 xenograft tissues

with the dnnRRS algorithm and used FlowJo for data analysis. We detected 32.6% and 0.13%

CAR+ in the CAR+ (Fig. 2D) and CAR- BE2 (Fig. 2E) tumor, respectively. CD19 expression was

below 1% and above 30% in CAR+ BE2 tumor and CAR- BE2 xenograft (Fig. 2D, E),

respectively. In the CAR- BE2 tumor, PPIB and CD19 expression showed homogenous

distribution (Fig. 2F). By contrast, the areas that stained positive for CD19 were almost devoid of

PPIB expression in the CAR+ BE2 tumor and showed signs of recent tissues disruption or lysis

(Fig. 3G).

The distribution of CAR T cells *in vivo* differed spatially, and a substantial number of cells were

negative by CAR RNA FISH and CD19IHC in CAR+ BE2 tumor sections. To better characterize

this heterogeneity of the tumor composition, we performed additional combinational RNA FISH

for the CAR, CD19 and CD171, another highly expressed protein in neuroblastomas and in the

BE2 neuroblastoma cell line *(20, 26)*. We found up to 38% CD171RNA+/CD19- and less than

0.01% CD171RNA+/CD19+ cells in the CAR+ BE2 tumor and in stark contrast ~75%

CD171RNA+/CD19+ in the CAR- BE2 tumor (data not shown).

These data indicated that the CAR RNA FISH detects specifically CAR T cells in tissues from

preclinical *in vivo* models and that the dnnRRS algorithm can be used to analyze CAR T cell

abundance, spatial distribution and efficacy.

CD4 and CD8 CAR T cells are widely distributed in the tumor and display differential

functional and spatial gene expression

To further characterize the functionality of the CAR T cells, we then set out to do combinations of

multiplex RNA FISH for the CAR, the T cell lineage marker CD4 and CD8α, as well as the effector

genes IFNγ and GZMB and analyzed the image data sets with the dnnRRS algorithm to phenotype

the tissue.

Independent CAR RNA FISH in combination with CD4 or CD8α showed that the CD4 and CD8

CAR T cells are evenly distributed within the CAR+ foci (Fig. 3A, B). We then subjected the

image data to computational analysis with the dnnRRS algorithm. For the CAR and CD8α RNA

FISH, 20.6% of the total cells were CAR+/CD8α+ and 19.9% CAR+/CD8- (presumably

CAR+/CD4+ T cells). For the CAR and CD4 RNA FISH, 17.2% of the total cells were

CAR+/CD4+ and 24.4% CAR+/CD4- (presumably CAR+/CD8+ T cells) (Fig 3C, D). Thus, in

this model the tissue-based distribution was relatively similar between the CD4+ and Cd8+ CAR

T cells.

To further characterize the functionality and heterogeneity of the CAR T cell pool within the

CAR+ BE2 tumor, we performed IHC for CD19, CAR RNA FISH combined with either GZMB

or IFNγ RNA FISH. CAR+ cells that express high levels of GZMB RNA (Fig. 4A) or IFNγ RNA

(Fig. 4B) can be observed by confocal microscopy close to the residual tumor. Concomitantly,

CAR+ cells that are located in areas with low or absence of CD19 expression only express low

levels of either GZMB or IFNγ RNA (Fig. 9 suppl. A and B). Whole-tissue tile scans of the CD19,

CAR RNA FISH co-staining with either GZMB RNA FISH (n=5) (Fig. 4C) or IFNγ RNA FISH

(n=5) (Fig. 4D) also indicated that the highest expression of GZMB and IFNγ is in close proximity

to the tumor. Examples images from this scan are also shown in the data supplement (Fig. 9 suppl.

C and D). The data sets were analyzed by dnnRRS and the x y position of CD19+ and CAR+

GZMB-/CAR+ GZMB+ (Fig. 4E and Fig. 10 suppl.) or CAR+ IFNγ-/CAR+ IFNγ+ (Fig. 5F and

Fig. 11 suppl.) cell were plotted separately or combined in Flowjo.

Around 2500 or 2800 CD19+ cells were detected in the GZMB and IFNγ datasets, respectively

(Fig. 5A). We detected approximately >31000 CAR+ GZMB- and 700 CAR+ GZMB+ cells/tissue

section as well as >36000 CAR+IFNγ- and 40 CAR+ IFNγ+ cells/tissue section (Fig. 5B).

Therefore, the frequency and spatial density of CAR+ GZMB+ was approximately 2% or 37

cells/mm2 whereas for CAR+IFNγ+ cells it was below 0.2% or 1.8 cells/mm2 (Fig. 5C, D). The

frequency and spatial density of CAR+ cells that were negative for either GZMB or IFNγ was

above 98% or around 1500 CAR+/mm2 (Fig. 5C, D).

We divided the spatial datasets in a CD19high and CD19low area to further analyze the effect of

tumor antigen expression on the spatial expression of the CAR T cell effector genes GZMB and

IFNγ (Fig. 5E). The spatial density of CD19+ cells/mm2 in the CD19high and CD19low area in

the GZMB and IFNγ dataset were significantly different (Fig. 5F, G) indicating that CAR T cells

had a higher probability for antigen exposure in the CD19high area. For the GZMB dataset, the

number of total CAR+ (Fig. 5H), CAR+ GZMB- (Fig. 5I) and CAR+GZMB+ cells/mm2 (Fig. 5J)

was not significantly different between the CD19high and CD19low area, however, there was a

trend towards higher spatial density of total CAR+ and CAR+ GZMB- cells/mm2 in the CD19low

area and an opposite trend of higher spatial density for CAR+GZMB+ cells/mm2 in the CD19high

area. The ratio of CAR+ GZMB+ to CAR+ GZMB- cells/mm2 was significantly higher in the

CD19high area (Fig. 6K). For the IFNγ dataset, the number of total CAR+ (Fig. 6L), CAR+ IFNγ-

(Fig. 6M) and CAR+ IFNγ+ cells/mm2 (Fig. 6N) was significantly higher in the CD19low area.

The ratio of CAR+ IFNγ+ to CAR+ IFNγ- cells/mm2 was not significant in a comparison of the

CD19high and CD19low area (Fig. 6O). In summary, these data indicate that the dnnRRS CAR

RNA FISH approach in combination with RNA FISH for T cell lineage and effector genes can

provide important insights into the spatial distribution and functionality of CAR T cells on a single

cell level and indicate that there is a high heterogeneity and differential spatial effector gene

expression in CAR T cells with respect to the tumor target cells.

Discussion

We describe here the development of a multiplex RNA FISH assay that is coupled with a dnnRRS

algorithm for image segmentation and quantitation. This assay allows to interrogate single cell

expression levels of multiple mRNA of interest and their spatial distribution and can be used to

detect, quantitate and evaluate the frequency and function of CAR T cells and cancer cells in

tissues. The use of specific probes directed against both the wPRE and scFv allows accurate

detection of CAR in tissues. It circumvents the common caveats of immunohistochemistry such

as long lead-time for antibody development and overlap of endogenous epitopes with the chimeric

protein of interest. This also allows to quickly modify this approach for *in situ* trafficking studies

for novel CARs with different target proteins, e.g. CD20, CD22, BMCA or any other genetically

modified cell *(27)* and can be used to study the interplay of CAR T cells, tumor cells and the tumor

microenvironment in solid tumor to develop novel strategies and hypothesis to improve CAR T

cell function in solid tumors; an area where CAR T cells therapies still fall short of the clinical

responses seen in hematologic cancers*(27, 28)*.

In the mouse xenograft samples, we identified 2 distinct CD19t-/CD171+ and CD19t+/CD171+

BE2 cell populations 7 days post-CAR T cells infusion. As our interest in this manuscript is

directed at using the probes to evaluate the structure intersecting between CAR T cells and tumor

tissue, we did not further investigate the underlying molecular mechanism in this artificial tumor

model because CD19 expression in this BE2-tumor model is driven from a lentiviral expression

cassette and the CD19+ BE2 cells in the xenograft sample were purified by magnetic bead

purification (Adam Johnson, Michael C. Jensen, personal communication). Interestingly, we

could not detect PPIB RNA (or other mRNA targets, data not shown) in the CD19+ areas that were

surrounded by IFNγ + and GZMB+ CAR T cells. While this may be merely a RNA preservation

problem in this specific area of the tissue specimen, global RNA degradation is also a hallmark of

apoptosis and the presence of IFNγ+ and GZMB+ CAR T cells in close proximity of the disrupted

PPIB-/CD19+ areas support the notion that they are indeed apoptotic.

Recently, the use of a defined CD4:CD8 ratio was implicated with synergistic antitumor effects

and better persistence *(29, 30)*. Our data indicate that CD4 and CD8 CAR T cells traffic into the

tumor site and probably undergo rapid proliferation. Not unexpectedly, the CD4 and CD8 CAR T

cells localize in close proximity of each other within the tumor at a ratio of ~1:1 at 7 days post-

infusion. This may indicate that the CD4 CAR T cells provide helper-function through secretion

of soluble factors. These observations suggest more detailed evaluation of samples using our

technology may provide insight into CAR T-cell responses and side effects.

The presence of CAR T cells expressing GZMB in close proximity at the border between the solid

tumor and adjacent tissue and the decreased expression in distant, CD19 negative tissues suggests

cell-to-cell contact with target cells is required for CAR T-cell activation. In contrast to the

GZMB+ CAR T cells, the more delocalized expression of IFNγ may indicate that these CAR T

cells migrate in and out from the area of the tumor surface more frequently or express IFNγ for an

extended period of time post-antigen exposure. Furthermore, there is still a considerable number

of CAR T cells close to the residual tumor surface that does not express IFNγ or GZMB and this

functional CAR T cell heterogeneity corroborate the findings of a *in vivo* live cell imaging study

by Cazaux and co-workers in which not every interaction of a CAR T cell with a tumor cell *in vivo*

led to CAR-mediated Ca^{2+}-signaling and killing *(31)*. However, a second study using a similar

approach by Mulazzani and Co-Workers did not report a heterogenous response after CAR T cell

infusion *(32)*. More recently, a third study by Chen and Co-workers reported an

immunohistochemistry and in situ hybridization approach and computational analysis of the anti-

CD19 CAR Axicabtagene ciloleucel in diffuse large B-cell lymphoma in the ZUMA-1 clinical

trial 5 days post-infusion *(12)*. Interestingly, less than 5% of the T cells were CAR T cells, in which

74% expressed IFNγ mRNA and 31% expressed GZMB protein whereas the number of bystander

CD3+ cells that expressed these markers exceeded the CAR T cells by 5-10 fold in absolute

numbers. The authors concluded that the non-CAR cells are the primary source of immune cell

activation and IFNγ production and their data indicate that there is also some degree of functional

heterogeneity of CAR T cells in a clinical setting. In our sample material bystander T cells and

other immune cells are absent and the frequency of GZMB+ and IFNγ + CAR T cells is

substantially lower 10 days post-infusion and warrants further studies into the contribution of

bystander immune cells and the suitability of tumor mouse models. Of note, an important

distinction between our study and the three studies is the costimulatory domain of the CAR. In our

sample tissue, a 41bb costimulatory domain was used whereas a CD28 costimulatory domain was

used in the two mouse studies and the study in clinical specimen*(12, 31, 32)*, which reportedly

have different kinetics and persistence in vivo. Further studies into CAR T cell functional

heterogeneity in xenograft models and human tumor and healthy tissue and how it relates to

clinical response rates will likely contribute to improvement of future CAR T cells interventions.

Methodically, the in vivo two photon microscopy assays published by Cazaux and Co-workers as

well as by Mulazzani and Co-Workers allow to study eGFP-expressing murine CAR T cells in

brain, lung, bone marrow and lymphoid tissue of xenografted mice *(31, 32)* and permit to perform

in vivo live tracking studies over days and weeks. These will be particularly helpful to better

understand the in vivo dynamics of CAR T cells in mice over time, however, the application of

two photon microscopy is still hampered in larger animal models and human because of the

limitations of deep tissue imaging and high immunogenicity of fluorescent proteins in immune-

competent large animal models and humans. The approach to phenotype CAR T cells in situ

reported by Chen and Coworkers is similar to ours in that it uses a computer algorithm to analyze

complex multicolor image datasets derived from tissue sections. Notable differences are that the

dnnRRS algorithm performs semi-supervised machine learning on whole tissue sections and does

the analysis of multicolor fluorescence IHC and FISH in the same run. Chen and coworkers report

a supervised analysis of 3-5 fields of view for the fluorescent IHC images and a separately analysis

for chromogenic RNA FISH*(12)*.

Globally, our studies provide insight on how CAR T cells are acting within in a solid tumor setting.

The IFNγ-expressing CAR T cells may change the local tumor microenvironment such as M2-to-

M1 macrophage differentiation or stimulate incoming tumor-infiltrating T cells and, thus gradually

lead to tumor regression. However, extended IFNγ-expression from tumor-infiltrating T

lymphocytes is associated with increase PD-1L expression and immune evasion in melanoma *(33)*

and neuroblastoma*(34)* and thus, extended IFNγ secretion from CAR T cells may be also

detrimental in a solid tumor setting. Of note, high levels of serum IFNγ are also implicated in an

increased risk of grade 3–5 neurotoxicity in anti-CD19 CAR T cell therapy*(11, 35)*. Further

evaluation of *in situ* cytokine-chemokine expression of CAR T cells in tissue is likely to lead to

increased understanding of the mechanism of CAR T cell tumor killing and toxicity.

Quantitative PCR, flow cytometry and bioluminescence imaging are the main tools to study CAR

T cells in vivo pre-clinical and, to some extent, clinical studies. Even though these assays allow

high sample throughput, macroscopic analysis of the tumor mass and/or provide high statistical

power for quantification, none of them assesses the spatial distribution CAR T cells and the

interplay of CAR T cells and their targets.

The dnnRRS CAR RNA FISH approach described here provides an easy-to-go tool kit for the

CAR T cells studies and promises to provide insights into the trafficking, persistence and

phenotype of CAR T cells in liquid and solid tumors in large animal models and human to better

understand the interplay between the tumor microenvironment and CAR T cell function.

With respect to the application of CAR T cells in the treatment of HIV, we also validated probes

to study anti-HIV CAR T cells in tissues. In this regard, CAR T cell trafficking into germinal

centers and other anatomic sites that harbors the HIV reservoir is considered to be crucial for a

long-term HIV remission *(21, 36, 37)* and the technique described here may provide valuable

insights to improve CAR T cell therapeutic approaches for HIV iteratively. Likewise, dnnRRS

FISH may also be applicable for infectious disease research with the advent of novel RNA and

DNA FISH assays to study persistence of HIV reservoirs in immune-privileged sites in tissues*(21,*

38) and the renewed interest in novel cell-based immunotherapies for HIV*(2, 3)*.

Bibliography

1. L. Liu, B. Patel, M. H. Ghanem, V. Bundoc, Z. Zheng, R. A. Morgan, S. A. Rosenberg, B.

Dey, E. A. Berger, Novel CD4-Based Bispecific Chimeric Antigen Receptor Designed for

Enhanced Anti-HIV Potency and Absence of HIV Entry Receptor Activity, *J. Virol.* **89**, 6685–

6694 (2015).

2. R. S. Leibman, M. W. Richardson, C. T. Ellebrecht, C. R. Maldini, J. A. Glover, A. J. Secreto,

I. Kulikovskaya, S. F. Lacey, S. R. Akkina, Y. Yi, F. Shaheen, J. Wang, K. A. Dufendach, M. C.

Holmes, R. G. Collman, A. S. Payne, J. L. Riley, Supraphysiologic control over HIV-1

replication mediated by CD8 T cells expressing a re-engineered CD4-based chimeric antigen

receptor, *PLoS Pathog.* **13**, 1–30 (2017).

3. M. Hale, T. Mesojednik, G. S. Romano Ibarra, J. Sahni, A. Bernard, K. Sommer, A. M.

Scharenberg, D. J. Rawlings, T. A. Wagner, Engineering HIV-Resistant, Anti-HIV Chimeric

Antigen Receptor T Cells, *Mol. Ther.* **25**, 570–579 (2017).

4. R. A. Morgan, J. C. Yang, M. Kitano, M. E. Dudley, C. M. Laurencot, S. A. Rosenberg, Case

report of a serious adverse event following the administration of T cells transduced with a

chimeric antigen receptor recognizing ERBB2, *Mol. Ther.* **18**, 843–851 (2010).

5. K. Watanabe, S. Kuramitsu, A. D. Posey, C. H. June, Expanding the therapeutic window for

CAR T cell therapy in solid tumors: The knowns and unknowns of CAR T cell biology, *Front.*

Immunol. **9**, 1–12 (2018).

6. M. R. Parkhurst, J. C. Yang, R. C. Langan, M. E. Dudley, D. A. N. Nathan, S. A. Feldman, J.

L. Davis, R. A. Morgan, M. J. Merino, R. M. Sherry, M. S. Hughes, U. S. Kammula, G. Q. Phan,

R. M. Lim, S. A. Wank, N. P. Restifo, P. F. Robbins, C. M. Laurencot, S. A. Rosenberg, T cells

targeting carcinoembryonic antigen can mediate regression of metastatic colorectal cancer but

induce severe transient colitis, *Mol. Ther.* **19**, 620–626 (2011).

7. X. Wang, W. C. Chang, C. L. W. Wong, D. Colcher, M. Sherman, J. R. Ostberg, S. J. Forman,

S. R. Riddell, M. C. Jensen, A transgene-encoded cell surface polypeptide for selection, in vivo

tracking, and ablation of engineered cells, *Blood* **118**, 1255–1263 (2011).

8. A. Taraseviciute, V. Tkachev, R. Ponce, C. J. Turtle, J. M. Snyder, H. D. Liggitt, D. Myerson,

L. Gonzalez-Cuyar, A. Baldessari, C. English, A. Yu, H. Zheng, S. N. Furlan, D. J. Hunt, V.

Hoglund, O. Finney, H. Brakke, B. R. Blazar, C. Berger, S. Riddell, R. Gardner, L. S. Kean, M.

C. Jensen, Chimeric Antigen Receptor T Cell-Mediated Neurotoxicity in Non-Human Primates,

Cancer Discov. **8**, 750–763 (2018).

9. C. J. Turtle, L. Hanafi, C. Berger, T. A. Gooley, S. Cherian, M. Hudecek, D. Sommermeyer,

K. Melville, B. Pender, T. M. Budiarto, E. Robinson, N. N. Steevens, C. Chaney, L. Soma, X.

Chen, C. Yeung, B. Wood, D. Li, J. Cao, S. Heimfeld, M. C. Jensen, S. R. Riddell, D. G.

Maloney, CD19 CAR – T cells of defined CD4+:CD8+ composition in adult B cell ALL

patients, *J Clin Invest* **1**, 1–16 (2016).

10. D. M. O'Rourke, M. P. Nasrallah, A. Desai, J. J. Melenhorst, K. Mansfield, J. J. D.

Morrissette, M. Martinez-Lage, S. Brem, E. Maloney, A. Shen, R. Isaacs, S. Mohan, G. Plesa, S.

F. Lacey, J. M. Navenot, Z. Zheng, B. L. Levine, H. Okada, C. H. June, J. L. Brogdon, M. V.

Maus, A single dose of peripherally infused EGFRvIII-directed CAR T cells mediates antigen

loss and induces adaptive resistance in patients with recurrent glioblastoma, *Sci. Transl. Med.* **9**,

eaaa0984 (2017).

11. A. E. Obstfeld, N. V. Frey, K. Mansfield, S. F. Lacey, C. H. June, D. L. Porter, J. J.

Melenhorst, M. A. Wasik, Cytokine release syndrome associated with chimeric-antigen receptor

T-cell therapy: clinicopathological insights, *Blood* **130**, 2569–2572 (2017).

12. P.-H. Chen, M. Lipschitz, J. L. Weirather, C. Jacobson, P. Armand, K. Wright, F. S. Hodi, Z.

J. Roberts, S. A. Sievers, J. Rossi, A. Bot, W. Y. Go, S. J. Rodig, Activation of CAR and non-

CAR T cells within the tumor microenvironment following CAR T-cell therapy, *JCI Insight* **5**

(2020).

13. L. E. Baum, T. Petrie, G. Soules, N. Weiss, A maximation techique occuring in the statistical

analysis of probablistic functions of markov chains, *Ann. Math. statisics* **41**, 164–171 (1970).

14. J. Li, A. Najmi, R. M. Gray, Image classification by a two-dimensional hidden Markov

model, *IEEE Trans. Signal Process.* **48**, 517–533 (2000).

15. R. Zhang, W. Ouyang, W. Cham, Image multi-scale edge detection using 3-D Hidden

Markov Model based on the non-decimated wavelet, *2009 16th IEEE Int. Conf. Image Process.*

Cairo , 2173–2176 (2009).

16. Y. Zhang, M. Brady, S. Smith, Segmentation of brain MR images through a hidden Markov

random field model and the expectation-maximization algorithm, *IEEE Trans. Med. Imaging* **20**,

45–47 (2001).

17. K. M. Kim, K. Son, G. T. R. Palmore, Neuron image analyzer: Automated and accurate

extraction of neuronal data from low quality images, *Sci. Rep.* **5**, 17062 (2015).

18. A. Li, L. Corey, J. Zhu, Random-reaction-seed method for automated identification of neurite

elongation and branching, *Sci. Rep.* **9**, 2908 (2019).

19. J. Wang, O. W. Press, C. G. Lindgren, P. Greenberg, S. Riddell, X. Qian, C. Laugen, A.

Raubitschek, S. J. Forman, M. C. Jensen, Cellular immunotherapy for follicular lymphoma using

genetically modified CD20-specific CD8+ cytotoxic T lymphocytes, *Mol. Ther.* **9**, 577–586

(2004).

20. A. Kunkele, A. J. Johnson, L. S. Rolczynski, C. A. Chang, V. Hoglund, K. S. Kelly-Spratt,

M. C. Jensen, Functional Tuning of CARs Reveals Signaling Threshold above Which CD8+

CTL Antitumor Potency Is Attenuated due to Cell Fas-FasL-Dependent AICD, *Cancer Immunol.*

Res. **3**, 368–379 (2015).

21. C. Deleage, S. W. Wietgrefe, G. Del Prete, D. R. Morcock, X.-P. Hao, M. Piatak, Jr., J. Bess,

J. L. Anderson, K. Perkey, C. Reilly, J. M. McCune, A. T. Haase, J. D. Lifson, T. W. Schacker,

J. D. Estes, Defining HIV and SIV Reservoirs in Lymphoid Tissues, *Pathog. Immun.* **1**, 68–106

(2016).

22. J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch,

C. Rueden, S. Saalfeld, B. Schmid, J. Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P.

Tomancak, A. Cardona, Fiji: An open-source platform for biological-image analysis, *Nat.*

Methods **9**, 676–682 (2012).

23. O. Ronneberger, P. Fischer, T. Brox, U-net: Convolutional networks for biomedical image

segmentation, *Lect. Notes Comput. Sci. (including Subser. Lect. Notes Artif. Intell. Lect. Notes*

Bioinformatics) **9351**, 234–241 (2015).

24. I. C. Nicholson, K. A. Lenton, D. J. Little, T. Decorso, F. T. Lee, A. M. Scott, H. Zola, A. W.

Hohmann, Construction and characterisation of a functional CD19 specific single chain Fv

fragment for immunotherapy of B lineage leukaemia and lymphoma, *Mol. Immunol.* **34**, 1157–

1165 (1997).

25. R. S. Rudicell, Y. D. Kwon, S.-Y. Ko, A. Pegu, M. K. Louder, I. S. Georgiev, X. Wu, J. Zhu,

J. C. Boyington, X. Chen, W. Shi, Z. -y. Yang, N. A. Doria-Rose, K. McKee, S. O'Dell, S. D.

Schmidt, G.-Y. Chuang, A. Druz, C. Soto, Y. Yang, B. Zhang, T. Zhou, J.-P. Todd, K. E. Lloyd,

J. Eudailey, K. E. Roberts, B. R. Donald, R. T. Bailer, J. Ledgerwood, J. C. Mullikin, L. Shapiro,

R. A. Koup, B. S. Graham, M. C. Nason, M. Connors, B. F. Haynes, S. S. Rao, M. Roederer, P.

D. Kwong, J. R. Mascola, G. J. Nabel, Enhanced Potency of a Broadly Neutralizing HIV-1

Antibody In Vitro Improves Protection against Lentiviral Infection In Vivo, *J. Virol.* **88**, 12669–

12682 (2014).

26. A. Taraseviciute, A. J. Johnson, M. C. Jensen, O. Finney, M. Berger, L. S. Finn, L. S.

Rolczynski, S. Mgebroff, A. Künkele, C. Brown, J. R. Park, C. Berger, C. A. Chang, Preclinical

Assessment of CD171-Directed CAR T-cell Adoptive Therapy for Childhood Neuroblastoma:

CE7 Epitope Target Safety and Product Manufacturing Feasibility, *Clin. Cancer Res.* **23**, 466–

477 (2016).

27. C. H. June, R. S. O'Connor, O. U. Kawalekar, S. Ghassemi, M. C. Milone, CAR T cell

immunotherapy for human cancer, *Science (80-.).* **359**, 1361–1365 (2018).

28. C. S. M. Yong, V. Dardalhon, C. Devaud, N. Taylor, P. K. Darcy, M. H. Kershaw, CAR T-

cell therapy of solid tumors, *Immunol. Cell Biol.* **95**, 356–363 (2017).

29. D. Sommermeyer, M. Hudecek, P. L. Kosasih, T. Gogishvili, D. G. Maloney, C. J. Turtle, S.

R. Riddell, Chimeric antigen receptor-modified T cells derived from de fi ned CD8+ and CD4+

subsets confer superior antitumor reactivity in vivo, *Leukemia* **30**, 492–500 (2016).

30. C. J. Turtle, L. Hanafi, C. Berger, M. Hudecek, B. Pender, E. Robinson, R. Hawkins, C.

Chaney, S. Cherian, X. Chen, L. Soma, B. Wood, D. Li, S. Heimfeld, S. R. Riddell, D. G.

Maloney, Immunotherapy of non-Hodgkin's lymphoma with a defined ratio of CD8+ and CD4+

CD19-specific chimeric antigen receptor – modified T cells, *Sci. Transl. Med.* **8**, 355ra116

(2016).

31. M. Cazaux, C. L. Grandjean, F. Lemaître, Z. Garcia, R. J. Beck, I. Milo, J. Postat, J. B.

Beltman, E. J. Cheadle, P. Bousso, Single-cell imaging of CAR T cell activity in vivo reveals

extensive functional and anatomical heterogeneity, *J. Exp. Med.* **216**, 1038–1049 (2019).

32. M. Mulazzani, S. P. Fräßle, I. von Mücke-Heim, S. Langer, X. Zhou, H. Ishikawa-

Ankerhold, J. Leube, W. Zhang, S. Dötsch, M. Svec, M. Rudelius, M. Dreyling, M. von

Bergwelt-Baildon, A. Straube, V. R. Buchholz, D. H. Busch, L. von Baumgarten, Long-term in

vivo microscopy of CAR T cell dynamics during eradication of CNS lymphoma in mice, *Proc.*

Natl. Acad. Sci. U. S. A. **116**, 24275–24284 (2019).

33. S. L. Topalian, J. M. Taube, A. P. Klein, R. A. Anders, D. M. Pardoll, S. Chen, H. Xu, T. L.

McMiller, L. Chen, R. Sharma, G. D. Young, Colocalization of Inflammatory Response with B7-

H1 Expression in Human Melanocytic Lesions Supports an Adaptive Resistance Mechanism of

Immune Escape, *Sci. Transl. Med.* **4**, 127ra37 (2012).

34. P. Srinivasan, X. Wu, M. Basu, C. Rossi, A. D. Sandler, PD-L1 checkpoint inhibition and

anti-CTLA-4 whole tumor cell vaccination counter adaptive immune resistance: A mouse

neuroblastoma model that mimics human disease, *PLoS Med.* **15**, e1002497 (2018).

35. J. Gust, K. A. Hay, L. A. Hanafi, D. Li, D. Myerson, L. F. Gonzalez-Cuyar, C. Yeung, W. C.

Liles, M. Wurfel, J. A. Lopez, J. Chen, D. Chung, S. Harju-Baker, T. Ozpolat, K. R. Fink, S. R.

Riddell, D. G. Maloney, C. J. Turtle, Endothelial activation and blood–brain barrier disruption in

neurotoxicity after adoptive immunotherapy with CD19 CAR-T cells, *Cancer Discov.* **7**, 1404–

1419 (2017).

36. K. P. Haran, A. Hajduczki, M. S. Pampusch, G. Mwakalundwa, D. A. Vargas-Inchaustegui,

E. G. Rakasz, E. Connick, E. A. Berger, P. J. Skinner, Simian immunodeficiency virus (SIV)-

specific chimeric antigen receptor-T cells engineered to target B cell follicles and suppress SIV

replication, *Front. Immunol.* **9**, 492 (2018).

37. V. I. Ayala, C. Deleage, M. T. Trivett, S. Jain, L. V. Coren, M. W. Breed, J. A. Kramer, J. A.

Thomas, J. D. Estes, J. D. Lifson, D. E. Ott, CXCR5-Dependent Entry of CD8 T Cells into

Rhesus Macaque B-Cell Follicles Achieved through T-Cell Engineering, *J. Virol.* **91**, e02507-16

(2017).

38. J. D. Estes, C. Kityo, F. Ssali, L. Swainson, K. N. Makamdop, G. Q. Del Prete, S. G. Deeks,

P. A. Luciw, J. G. Chipman, G. J. Beilman, T. Hoskuldsson, A. Khoruts, J. Anderson, C.

Deleage, J. Jasurda, T. E. Schmidt, M. Hafertepe, S. P. Callisto, H. Pearson, T. Reimann, J.

Schuster, J. Schoephoerster, P. Southern, K. Perkey, L. Shang, S. W. Wietgrefe, C. V. Fletcher,

J. D. Lifson, D. C. Douek, J. M. McCune, A. T. Haase, T. W. Schacker, Defining total-body

AIDS-virus burden with implications for curative strategies, *Nat. Med.* **23**, 1271–1276 (2017).

Acknowledgements

We thank S. Sun, L. Schroeder, P Guo, J. Vazquez and D. McDonald for technical support and

Mindy Miner for editing the manuscript. We thank Christopher W. Peterson and Hans-Peter Kiem

for providing the VRC07-523 CAR Jurkat cell line.

This independent research was supported by National Institute of Health grant UM1 A126623

and by the Gilead Sciences HIV Cure Grant Program.

This research was supported by the Experimental Histopathology Shared Resource of the Fred

Hutch/University of Washington Cancer Consortium (P30 CA015704).

This research was supported by the Cellular Imaging Shared Resource (CISR) of the Fred

Hutch/University of Washington Cancer Consortium (P30 CA015704).

SUT was a full-time employee of Juno Therapeutics during the period of performing this work.

The remaining authors declare no competing interest.

Code and data availability statement:

Code and raw data are available upon request

Author contributions:

K.E. and L.C. conceived the project. K.E. performed most experiments, analyzed the data

together with A.L. and interpreted the data together with LC.

A.L. developed the algorithm and A. L. and K.E. performed the dnnRRS analysis.

K.D. helped with experiments.

J.Z. helped with scientific advice and valuable expertise

S.T. designed the VRC07-523 CAR and provided scientific advice.

M.J. provided tissue samples and scientific advice.

KE and LC wrote the manuscript with input from all authors.

Figures

Figure 1: CD19 CAR RNA FISH *in vitro* validation.

The probes for CAR mRNA detection are directed against the scFv and wPRE and bridging areas

in the signaling domain of the CAR (A).

Cytospins of actively growing human anti-CD19 CD8 CAR T cells or Jurkat cells were stained

with CAR FMC63 26zz probes or PPIB (+ control) or DapB (- control) probes (B). CD19 CAR T

cells were stained with CAR FMC63 26zz probes-Alexa488 and Texas red-conjugated wheat germ

agglutinin to improve cell segmentation (C). dnnRRS analysis of multicolor raw images includes

nuclei and RNA detection as well random reaction seed detection of plasma membrane staining

(D). Fixed human anti-CD19 CD8 CAR T cells and Jurkat cells were mixed at different ratio,

processed for cytospins and stained with CAR FMC63 26zz probes-Atto550 and DAPI (E).

Correlation of CAR+ cells detected by dnnRRS analysis and flow cytometry (F). Comparison of

RNA dots/cells between different dilution steps. Statistics are calculated by one-way ANOVA

with Tukey's multiple comparison test (G). Staining experiments were performed at least 3 times

with technical duplicates.

Scale bar = 10 μm

Figure 2: in situ detection of anti-CD19-CAR T cells in mouse xenografts of NSG mice after

anti-CD19 CAR T cell infusion.

A BE2 neuroblastoma xenograft with ectopic expression of truncated CD19 and infused with anti-

CAR T cells or control T cells was used for the CAR RNA FISH *in situ* validation.

Staining of CAR+ and CAR- BE2 tumor tissue sections with CAR FMC63 15zz probes (red). The

red dotted line denotes an area that showed specific CAR RNA FISH staining (A). Staining of

CAR+ BE2 tumor with CAR FMC63 15zz probes and Texas Red-conjugated wheat germ

agglutinin improves cell segmentation (B). dnnRRS analysis of multicolor raw images includes

nuclei and CAR RNA detection as well random reaction seed detection of plasma membrane

staining. Data mining can be performed, and differential RNA expression and cell centers can be

visualized with Flowjo. The example picture derives from the area framed in white in B (C). A

thumbnail image and dnnRRS analysis of multiplex fluorescence RNA FISH and

immunohistochemistry staining of CAR RNA FISH and CD19 IHC (n=3) on CAR+ (D) and CAR-

(E) tumor sections shows the global CAR T cells distribution. A thumbnail image and dnnRRS

analysis of multiplex fluorescence RNA FISH and immunohistochemistry staining of PPIB RNA

FISH and CD19 IHC on CAR+ (F) and CAR- (G) tumor sections shows the global RNA integrity.

Figure 3: CD4 and CD8 CAR T cells are present in the tumor and express IFNγ.

CAR T cell lineage phenotyping was done with CAR RNA FISH in combination with CD4 and

CD8α, RNA FISH. A representative image are shown for CAR RNA and CD4 RNA (A) or CD8α

RNA (B) as well as nuclear and plasma membrane counterstaining. Thumbnail image of the

stitched multiplex fluorescence CAR (red) and CD4 (green) RNA FISH raw image channels and

dnnRRS analysis of a CAR+ BE2 tumor sections (n=3) (C). Thumbnail image of the stitched

multiplex fluorescence CAR (red) and CD8α (green) RNA FISH raw image channels and dnnRRS

analysis of a CAR+ BE2 tumor sections (n=3) (D).

Figure 4: GZMB and IFNγ are highly expressed in a minority of CAR+ cells.

CAR T cell functional phenotyping in a BE2 neuroblastoma xenograft was done by IHC for tumor

marker CD19 and CAR RNA FISH in combination with either GZMB (n=5) or IFNγ RNA FISH

(n=5). Confocal images of the combination of CAR RNA and CD19 with either GZMB RNA (A)

or IFNγ RNA in a representative area close to the residual tumor (B). The overview thumbnail

image of CAR RNA, CD19 with either GZMB RNA (C) or IFNγ RNA (D) staining shows the

global distribution of the different markers. dnnRRS analysis of the whole tissue section and the

various plots depict the spatial distribution of CD19+, CAR+ and GZMB+ (E) cells or IFNγ+ cells

(F). Scale bar = 10 μm

Figure 5: Spatial density analysis of the GZMB+ and IFNγ+ CAR T cells in CD19high and

low areas reveals differential spatial effector gene expression

Statistics of the CD19/CAR/GZMB (n=5) and CD19/CAR/IFNγ (n=5) cumulative datasets from

Figure 6 and supplementary Fig. 11 and 12. Total CD19+ cells/mm2 (A) as well as total (B),

frequency (C) and spatial density (D) of CAR+ GZMB-, CAR+ GZMB+, CAR+ IFNγ-, CAR+

IFNγ + are shown. An example shows how tissue sections were separated in CD19high and

CD19low areas denoted by the yellow color and, in this case the CAR+GZMB- cells as red dots

(E). Spatial density of CD19+ cells in the respective areas in the CD19/CAR/GZMB (n=5) (F) and

CD19/CAR/IFNγ (n=4) (G) cumulative datasets. In the CD19/CAR/GZMB cumulative dataset,

total CAR+ cells/mm2 (H), CAR+ GZMB- cells/mm2 (I), CAR+ GZMB+ cells/mm2 (J), ratio of

CAR+ GZMB+ to CAR+ GZMB- cells/mm2 (K) in the CD19high and CD19low area. In the

CD19/CAR/ IFNγ cumulative dataset, total CAR+ cells/mm2 (L), CAR+ IFNγ- cells/mm2 (M),

CAR+ IFNγ+ cells/mm2 (N), ratio of CAR+ IFNγ+ to CAR+ IFNγ- cells/mm2 (O) in the

CD19high and CD19low area. C-D were analyzed by one-way ANOVA. F-O were analyzed by

paired t test.

Supplementary Figure 1: CAR probe table, principle of RNA FISH.

Four zz probes sets were designed for detection of anti-CD19 and anti-HIV CAR T cells. The

probes bind to regions unique to the CAR mRNA that cannot be found in non-genetically

engineered organisms (A).

Supplementary Figure 2: CAR FMC63-15zz probes on human tonsil tissue

Human tonsil sections were stained for IGHG gene (A) or with FMC63-15zz probe with (B) or

without (C) RNAase pretreatment (n=2).

Scale bar = 1000 micrometers

Supplementary Figure 3: purity of primary CAR T cell culture after expansion.

Anti-CD19 CD8 CAR T cells were cultured with γ-irradiated CD19+ TM-LCL cells to stimulate

expansion. On day 12 of cell culture, CAR T cell purity was assessed by flow cytometry for the

cell surface markers CD19-BV786, EGFRt-PE and CD8-FITC (n=1). These cells were processed

for CAR RNA FISH and used for the *in vitro* validation.

Supplementary Figure 4: CAR RNA FISH detects CAR RNA mainly extranuclear.

VRC07-523 CAR Jurkat cells were stained with CAR VRC07-523 42zz probes and nuclear

counterstain to exemplify subcellular localization of CAR RNA. The CAR RNA is mainly detected

extranuclear or on the outline of the nuclear counterstain.

Supplementary Figure 5: VRC07 CAR RNA FISH *in vitro* validation.

Cytospins of VRC07-523 CAR-expressing Jurkat cells or non-transduced Jurkat cells were stained

with VRC07-523 42zz probes (A). Fixed human VRC07-523 CAR-expressing Jurkat cells and

non-transduced Jurkat cells were mixed at different ratio, processed for cytospins and stained with

VRC07-523 42zz probes and DAPI (B). Correlation of dnnRRS algorithm detected CAR+ cells

stained with the VRC07 42zz probes with expected ratio based on EGFRt+-flow cytometry (C).

Comparison of RNA dots/cells between different dilution steps. Statistics are calculated by one-

way ANOVA with Tukey's multiple comparison VRC07-523 CAR Jurkat data set (D).

Staining experiments were performed 3 times with technical duplicates.

Scale bar = 10 μm

Supplementary Figure 6: presence and expression of CAR sequence and wPRE in mouse

xenografts.

ddPCR analysis for the presence of the CAR signaling domain (CD8CD3) and the 3'UTR

containing the wPRE sequence of genomic DNA in xenograft tissue section from CAR T cell-

infused and control mice . (n=2). ddPCR primers and probes were directed against the CD3 CD8

region and wPRE portion to test for CAR and CD19 lentivirus-derived expression cassettes.

Supplementary Figure 7: CAR RNA FISH in situ validation.

Specificity of CAR RNA FISH in situ staining was further validated with RNA FISH staining for

housekeeping gene PPIB (A) and bacterial gene DapB (B) CAR RNA FISH protocol without the

addition of CAR FMZ63 15zz probes (C) or after RNAse treatment (D) (n=3).

Supplementary Figure 8: Manual and dnnRRS cell counting numbers correlate with each

other

We chose 10 images from the CAR+ BE2 tumor dataset that are exemplary for images without

artifacts or that show different imaging artifacts like strong membrane staining or weak nuclei

staining (A). Total cell numbers were counted manually by three operators or with the dnnRRS

algorithm using the nuclei and membrane WGA staining (B). Correlation of the number of cells

counted/images by the three operators and the dnnRRS method (C). The counting time

required/image is shown in (D).

Supplementary Figure 9: Low effector gene expression in CAR T cells that located further

away from the tumor.

CAR T cell functional phenotyping in a BE2 neuroblastoma xenograft was done by IHC for tumor

marker CD19 and CAR RNA FISH in combination with either GZMB (n=5) or IFNγ RNA FISH

(n=5). Confocal images of the combination of CAR RNA and CD19 with either GZMB RNA (A)

or IFNγ RNA (B) in a representative area far from the residual tumor. Exemplary wide field images

that were used to create the overview thumbnail images in Figure 5 image of CAR RNA, CD19

with either GZMB RNA (C) or IFNγ RNA (D).

Supplementary Figure 10: dnnRRS analysis CD19, CAR and GZMB expression of tissue

sections from a BE2 tumor.

CAR T cell functional phenotyping in a BE2 neuroblastoma xenograft was done by IHC for

tumor marker CD19 and CAR RNA FISH in combination with GZMB RNA FISH(n=4).

Supplementary Figure 11: dnnRRS analysis CD19, CAR and IFNγ expression of tissue

sections from a BE2 tumor.

CAR T cell functional phenotyping in a BE2 neuroblastoma xenograft was done by IHC for

tumor marker CD19 and CAR RNA FISH in combination with IFNγ RNA FISH (n=4).

Supplementary Table 1: CAR RNA FISH probes and respective catalog number.

Probe name	Catalog Number
VRC07-CAR-wPRE-O1	300031
FMC63 15ZZ	515321
FMC63 26ZZ	513551
FMC63 29ZZ	exclusive probe
DapB-C1	310043
Hs-PPIB-C2	313901-C2
Hs-PPIB-C1	313901-C1
Hs-GZMB	445971
Hs-GZMB-C2	445971-C2
Hs-CD8A-C3	560391-C3
Hs-CD4-C2	605601-C2
Hs- IFNγ-C1	310501-C1
Hs- IFNγ-C2	310501-C2

Figure 1

Figure 2

Figure 3

Figure 5

	total zz	scFvCAR	core	3'UTR	target tissue/species
FMC63 15zz	15	11	4	0	human and mouse tissue containing different lentiviral constructs
FMC63 27zz	27	11	4	12	human and mouse tissue with only CAR T cells present
FMC63 29zz (JCAR14)	29	11	4	12	human and mouse tissue with only CAR T cells present
VRC07-523 42zz	42	16	13	13	human and nonhuman primate

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

A   B   C   D

DAPI   DAPI   DAPI   DAPI

PPIB   DapB   no probes   CAR FMC63

RNAse

merge   merge   merge   merge

CAR T   +   +   +   +
cells infusion

Supplementary Figure 7

Supplementary Figure 8

Supplementary Figure 10

Supplementary Figure 11