Retinal mechanisms of saccadic suppression and their role in processing sequential stimuli

Saad Idrees[1,2], Matthias-Philipp Baumann[1], Marili Korympidou[1,2,3], Timm Schubert[1,3], Alexandra Kling[4], Katrin Franke[3,5], Ziad M. Hafed[1], Felix Franke[6,7,8]*, Thomas A. Münch[1,3,9]*

[1]Werner Reichardt Centre for Integrative Neuroscience, University of Tübingen, 72076 Tübingen, Germany

[2]International Max Planck Research School, University of Tübingen, 72074 Tübingen, Germany

[3]Institute for Ophthalmic Research, University of Tübingen, 72076 Tübingen, Germany

[4]Department of Neurosurgery, Stanford School of Medicine, Stanford, CA 94305, USA

[5]Bernstein Center for Computational Neuroscience, University of Tübingen, 72076 Tübingen, Germany

[6]Bio Engineering Laboratory, ETH Zürich, 4058 Basel, Switzerland

[7]Institute of Molecular and Clinical Ophthalmology Basel, 4031 Basel, Switzerland

[8]Faculty of Science, University of Basel, 4056 Basel, Switzerland

[9]Lead Contact

*Correspondence: felfranke@googlemail.com (F.F.), thomas.muench@gmail.com (T.A.M.)

Summary

Visual perception remains stable across eye movements such as saccades, despite the concurrent strong disruptive visual flow. This stability is partially associated with a reduction in visual system sensitivity, a phenomenon known as saccadic suppression. Saccadic suppression already starts in the retina, where responses of ganglion cells to flashes are strongly suppressed after saccade-like image shifts. However, the retinal circuit mechanisms giving rise to such suppression remain unknown. Here, we describe these mechanisms using electrophysiology in mouse, pig, and macaque retina, 2-photon calcium imaging, computational modeling, and human psychophysics. We find that several mechanisms support retinal saccadic suppression, including a novel retinal processing motif, "dynamic reversal suppression", which is triggered by sequential stimuli containing contrast reversals. This motif relies on nonlinear transformation of the inherently slow responses of cone photoreceptors. We describe two further components of suppression present in ON ganglion cells which originate in the cells' receptive field surround, highlighting a novel disparity between ON and OFF ganglion cells. Our results increase our understanding of retinal processing of any time-varying stimulus.

Keywords: retina, ganglion cell, cone photoreceptor, saccadic suppression, computational model, sequential stimuli, rapid image shifts, perception, dynamic vision, visual processing

Introduction

Vision appears as a continuous and coherent process. This is a striking achievement of the visual system, considering that the visual flow across the retina is not continuous, but characterized by frequent and sudden changes, irregularities, and disruptions. The most prominent cause for such disruptions are eye movements. For example, as a result of saccades, the number of photons falling onto a given area of the retina can change by several orders of magnitude within tens of milliseconds. These fast eye movements can therefore cause sudden and frequent visual transients of local intensity across the entire retina. At the same time, saccades are critical for efficiently sampling the visual world (Schütz et al., 2011; Tatler et al., 2010; Yarbus, 1967), which is particularly true for species in which high visual resolution is limited to a small fraction of the overall visual space, such as the foveal region in primates.

Perceptually, saccadic disruptions are minimized by reducing the sensitivity of the visual system to new input around the time of saccades - a phenomenon known as saccadic suppression. While this phenomenon has been extensively characterized over the past few decades (Beeler, 1967; Bremmer et al., 2009; Idrees et al., 2020; Krekelberg, 2010; Matin, 1974; Volkmann, 1986; Wurtz, 2008; Zuber and Stark, 1966), its underlying mechanisms still remain unclear. Several electrophysiological studies have shown neural correlates of saccadic suppression throughout the visual system, namely a modulation of neural activity and/or sensitivity around the time of saccades (Bremmer et al., 2009; Chen and Hafed, 2017; Hafed and Krauzlis, 2010; Ibbotson et al., 2008; Kleiser et al., 2004; Krekelberg, 2010; Wurtz, 2008). These observations have often been interpreted to be caused by active suppressive signals originating from (pre-) motor areas, such as corollary discharge signals related to the saccadic eye movement command (Bremmer et al., 2009; Diamond et al., 2000; Duffy and Lombroso, 1968; Ross et al., 2001; Zuber and Stark, 1966). Most studies investigating the mechanisms of saccadic suppression have therefore focused on cortical or subcortical neuronal recordings and/or on behavioral measures of perceptual state, largely neglecting the consequence of visual processing in early visual pathways, for example in the retina.

The retina is an independent signal processing "front end" in the visual system, before visual information is sent along the optic nerve to higher brain areas. Consequently,

image processing triggered by visual transients, such as those that naturally occur
during saccades, could potentially lead to altered retinal output. Retinal signal
processing could therefore contribute to perceptual saccadic suppression. Some
studies have investigated how the retina processes information in the context of
spatio-temporal dynamics that occur during natural visual behavior (Appleby and
Manookin, 2019; Baccus et al., 2008; Berry et al., 1999; Chen et al., 2013; Fried et al.,
2002; Garvert and Gollisch, 2013; Geffen et al., 2007; Gollisch, 2013; Münch et al.,
2009; Tikidji-Hamburyan et al., 2015; Wei, 2018; Zaghloul et al., 2007; Zhang et al.,
2012), including saccades (Amthor et al., 2005; Barlow et al., 1977; Enroth-Cugell and
Jakiela, 1980; Krishnamoorthy et al., 2017; Krueger and Fischer, 1973; Noda and
Adey, 1974; Passaglia et al., 2009; Roska and Werblin, 2003). A retinal neural
correlate of perceptual saccadic suppression has recently been shown by a previous
study from our labs (Idrees et al., 2020). There, we showed that the retinal output is
indeed altered by saccade-like image shifts. In most mouse and pig retinal ganglion
cells (RGCs) that we recorded from, responses to brief probe flashes were suppressed
when preceded by saccade-like image displacements across the retina. This retinal
saccadic suppression had properties consistent with the perceptual suppression of
probe flashes reported by human subjects using similar images, and following either
real or simulated saccades. In fact, we already observed elementary properties of
perceptual saccadic suppression, such as its dependency on background scene
statistics, at the level of the output of the retina, providing strong evidence of a retinal
mechanism directly contributing to perceptual saccadic suppression.

In this study, we describe such a mechanism. We experimentally mimicked the visual
flow resulting from saccades and recorded the neural activity of the output neurons of
the retina (RGCs) from ex vivo retinae of mice, pigs, and macaque monkeys. We found
that retinal saccadic suppression was the result of multiple mechanisms. A major
component originated from a novel visual processing motif within an RGC's receptive
field center, which we call "dynamic reversal suppression". Dynamic reversal
suppression results from a combination of inherent response properties of cone
photoreceptors and non-linear processing of cone output by downstream pathways in
the retina. Two further components of suppression originated from beyond the RGC's
receptive field center, one of them driven by GABAergic inhibition. Interestingly, these

two additional components were observed primarily in ON RGCs, highlighting a novel

disparity between ON and OFF type RGCs. Perhaps one of the most intriguing

outcomes of this study, also consistent with observations of perception (Idrees et al.,

2020), is that the suppressive effects observed in RGCs are not exclusively triggered

by saccades, but occur for many scenarios involving sequential visual stimulation,

which are ever-present during natural vision. Therefore, while the results described

here are crucial for understanding the mechanisms of saccadic suppression, they also

elucidate more general mechanisms of retinal signal processing across any

time-varying visual input over short time scales (10 ms to 1 s).

Results

Experimental Approach

We measured the modulation of retinal ganglion cell (RGC) output following

saccade-like changes of the visual input with a variety of different light stimulation

strategies (Fig. S1). The basic experimental paradigm was similar to that described in

(Idrees et al., 2020). Briefly, we first recorded spiking activity of RGCs in isolated ex

vivo mouse retinae using both high-density and low-density multi-electrode arrays

(MEAs). Each retina was exposed to a background texture having one of several

possible spatial scales that defined its spatial spectrum ("fine" to "coarse", Methods,

Fig. S2). We simulated saccade-like image displacements by rapidly translating the

texture globally across the retina (Methods; Fig. 1a). Most RGCs responded robustly

to such saccade-like texture displacements (see Fig. 1b for responses of example ON

and OFF RGCs). At different times relative to the saccade-like texture displacements

("saccades" from now on), we presented a brief probe flash (Fig. 1c). We then

analyzed how the response to this probe flash (spike rate of the RGC) was influenced

by the preceding saccade, by comparing it to the response to the flash presented in

isolation (baseline). To quantify RGC response modulation, we calculated a

modulation index (Methods) as follows: we first isolated the flash-induced response

component by subtracting the saccade-only response (e.g., Fig. 1b) from the response

to the composite saccade-flash stimulus (e.g., Fig. 1c). Based on this flash-induced

response component (Fig. 1d), we calculated the modulation index as $(r_d - r_b)/(r_d + r_b)$.

Here, r_d is the peak response to the probe flash presented with a delay d relative to

saccade onset, and r_b is the baseline (peak response to the flash presented ~2 s after the saccade). This modulation index is negative when flash-induced responses are suppressed (Fig. 1d shows, on the horizontal dashed line, the modulation index for the responses at each flash-time). In yet further recordings we applied various manipulations to this base paradigm to probe for the mechanisms underlying modulation of RGC responses following saccades. To generalize our findings across other species, we also performed similar analyses of pig and macaque RGC data.

Similarities and differences in retinal saccadic suppression across ON and OFF type RGCs

Suppression was robust across most RGCs that we recorded from, consistent with what we reported previously (Idrees et al., 2020). Here, we more closely inspected functionally different RGCs. Specifically, throughout this study, we divided RGCs into ON and OFF types, i.e. into RGCs responding best to light increments or decrements, respectively (Methods). We always quantified the modulation index defined above for ON RGCs based on their responses to bright probe flashes and for OFF RGCs based on their responses to dark probe flashes (Fig. 1c, d). Flash responses following a saccade were suppressed in both ON and OFF RGCs. Fig. 1d shows example cells, Fig. 1e the temporal profile of the mean population modulation index, and Fig. S3 the underlying population data. Suppression was consistently stronger for coarser background textures (Figs. 1e, S3), for both ON and OFF RGCs. This is consistent with (Idrees et al., 2020), where we showed that this dependency on the texture statistics can be explained by the distinct statistics of luminance and contrast changes when coarse or fine textures move across the RGCs' receptive fields. However, a striking difference existed in suppression recovery times: OFF RGCs recovered by ~350 ms after saccade onset, whereas ON RGCs recovered only by ~1 s. Similar results were obtained under scotopic conditions for coarse textures, while suppression for fine textures was very weak (Fig. S4; all other mouse retina data was recorded at mesopic conditions). In general, the presence of post-saccadic suppression of probe flash responses in both ON and OFF type RGCs suggests a common mechanistic theme across these cell types (Idrees et al., 2020). On the other hand, the different recovery times indicate either additional suppressive mechanisms in ON RGCs or additional recovery mechanisms in OFF RGCs.

Spatial origin of retinal saccadic suppression

Global component of suppression

To probe the mechanisms underlying suppression and its differences across ON and OFF type RGCs, we first examined the spatial origin of suppression. We hypothesized that suppression of flash responses was caused by circuits detecting rapid global shifts across the retina. Typically, these circuits include a lateral network of interneurons, communicating with RGCs even from beyond their classical center-surround receptive field (i.e., from their periphery, or far surround) (Lin and Masland, 2006; Roska and Werblin, 2003). To test whether suppression was caused by such circuits, we modified the spatial layout of the paradigm: we placed a square mask of 1000 x 1000 μm^2 (Fig. 2a, right) to restrict the saccades to the periphery of an RGC's receptive field. Similar to the previous experiments, the probe flash was either a dark or bright flash presented over the entire retina, including the masked region. Figure 2b shows the mean population modulation indices of ON RGCs (top) and OFF RGCs (bottom) from these experiments (Fig. S5 depicts the underlying population data and shows responses of representative ON and OFF RGCs from these experiments). In OFF RGCs, responses to full-field probe flashes were no longer suppressed when saccades were restricted to the periphery. The responses of ON RGCs, on the other hand, were still suppressed in this condition. The resulting suppression was however weaker and short-lived (recovered by 350 ms) than with full-field saccades. These observations (Figs. 2b, S5c) were robust across ON and OFF RGCs whose receptive fields were completely contained within the mask (Fig. S5b).

We will refer to this component of suppression in ON RGCs, which originates from the periphery, as the "global component" from now on. Such spatially far-reaching inhibition is often mediated through GABAergic wide-field amacrine cells. We tested this hypothesis by blocking GABA$_A$ receptors. Indeed, in the periphery saccade condition, the modulation index for most ON RGCs was around 0 in the presence of the GABA$_A$ receptor antagonist SR-95531 (Fig. 2b, S5d). These results suggest that this short-lived global component of suppression is caused by inhibition via GABAergic amacrine cells, perhaps similar to the polyaxonal amacrine cells described previously (Ölveczky et al., 2003; Roska and Werblin, 2003). Thus, suppression is indeed partially caused by circuits detecting global changes across the retina. However, those

circuits seem to act predominantly on ON RGCs, and even there, they do not account

for the total suppression observed with full-field saccades (without mask), which lasts

longer. Other, probably more local sources of suppression must exist that account for

the remaining suppression in ON RGCs and all of the suppression in OFF RGCs.

Local components of suppression

To understand the more local components of suppression, we used different analyses

and manipulations of the main experimental paradigm. As we will see below, the local

components can be subdivided into a "central" and a "surround" component. First, we

eliminated the global component, by repeating our normal full-field saccade paradigm

in the presence of GABA receptor blockers. The suppression profile of both ON and

OFF RGCs was only weakly affected upon blocking $GABA_{A,C}$ receptors (5 μM

SR-95531 and 100 μM Picrotoxin; Fig. 2c, S6a). Since the GABA-block eliminates the

global component of suppression, the remaining more local components did not seem

to rely on GABAergic inhibition. Also, this suggests that the local components

dominate retinal saccadic suppression under full-field conditions. We then also

blocked glycine receptors (1 μM of Strychnine; Fig. 2d, S6b) to test if the local

components of suppression were caused by local inhibition via glycinergic pathways.

Here again, the suppression profiles of both ON and OFF RGCs were only weakly

affected upon blocking glycine receptors in combination with blocking $GABA_{A,C}$

receptors. Therefore, inhibitory synaptic interactions are not the major mechanism

behind the local components of suppression.

Next, we tested whether these local components originated from within the receptive

field center. For this, we modified the spatial layout of our paradigm to exclude

saccades from the very center of the receptive field. Simply reducing the size of our

mask would have severely decreased the number of simultaneously recorded cells

located inside the mask, and we therefore resorted to a different strategy: saccades

and flashes were presented in small square regions spread across the retina,

separated by gaps kept at mean luminance ("checkerboard mask", Fig. 2e). In one

condition (Fig. 2e, left), we presented saccades and flashes in all regions of the

checkerboard mask; in the other condition (Fig. 2e, right), saccades and flashes were

presented in alternate regions. With this arrangement, saccades were excluded from

at most ~300 x 300 μm^2 of a cell's receptive field center, even if that cell was perfectly

centered on a non-saccade region. Flashes were presented in the set of regions that included the square region covering the receptive field center of the analyzed RGC (Fig. S7a).

Probe flash responses following saccades were suppressed in both ON and OFF RGCs when the saccade and flash were presented in all regions (Fig. 2f, thick lines; Fig. S7c; see Fig. S7b for example cells), consistent with the suppression observed after full-field saccades (Figs. 1, S3). When saccades were excluded from the receptive field center, and were presented in alternate regions to the flash, the flash responses were no longer suppressed in OFF RGCs (Fig. 2f, bottom, thin line), even though these cells showed spiking responses to saccades themselves (Fig. S7b). In fact, flash responses were even enhanced. This suggests that the local component of suppression in OFF RGCs arises fully from within the receptive field center ("central component"). This highly localized origin of suppression in OFF RGCs was further confirmed by additional analysis of the large mask experiments (see Figs. S7d, e). In ON RGCs, on the other hand, suppression persisted (Fig. 2f, top; Fig. S7c), even though a loss in suppression was apparent for flashes presented immediately after the saccade, at 117 and 150 ms (marked with an arrow in Fig. 2f). This suggests that in ON RGCs, part of the early suppression originates from the "central component". The leftover suppression during these early time points might be explained by the global component of suppression, described above (Fig. 2b, S5c), which should also be triggered under this experimental setting. However, since the global component also recovers quickly (by 350 ms, Fig. 2b, S5c), the persisting suppression at the later time points (350 ms and beyond) needs to originate from yet another source beyond the receptive field center. We call this the "surround component", and it may originate from the ON RGCs' immediate surround, which also experiences the saccade under this experimental setting. Therefore, in ON RGCs, the local component of suppression can be divided into a central and a surround spatial component.

Summary of retinal saccadic suppression spatial origins
In summary, our data suggests that retinal saccadic suppression is mediated by at least three components with distinct spatial origins and temporal properties (Fig. 3): a central, surround, and global component. Suppression in OFF RGCs is mediated exclusively by the central component, which originates from the cell's receptive field

center and is characterized by fast onset and fast recovery (by 350 ms after saccade onset). In ON RGCs, we most directly observed the global component (Figs. 2b, S5a). It extends into the periphery and its timing is similar to the central component in OFF RGCs. Only this global component is affected by blocking GABA receptors (Figs. 2b, S5b). During full-field saccades, removing this component by blocking GABA receptors has little effect on the overall suppression (Figs. 2c, 2d, S7), suggesting a more dominant role of the remaining components. The central component in ON RGCs can only be observed by the loss in suppression for early flashes when saccades are excluded from the receptive field center (marked with an arrow in Fig. 2f, top). Its full duration and time course are obscured by the concurrently acting global and surround components. However, given the identical pharmacological dependencies and spatial origins, it is plausible that the central component is symmetric across ON and OFF RGCs with a common underlying mechanism. Therefore, the longer suppression in ON RGCs can neither be attributed to the central nor global components. It likely originates from the immediate surround of the receptive field. This surround component is long lasting (recovers by ~1s) and has a slow onset (Figs. 2f, S3).

Suppression is triggered by interaction between consecutive stimuli of opposite polarity

We previously showed (Idrees et al., 2020) that retinal and perceptual saccadic suppression not only occurs after texture displacements, but also after instantaneous texture jumps and structure-free uniform luminance steps. These observations suggested that saccadic suppression is the consequence of rather general mechanisms in which the response to a second stimulus (here: probe flash) gets suppressed by a previous visual transient (caused by saccades or luminance steps). In the following, we apply additional analysis to the luminance-step paradigm (Fig. S1b) dataset of (Idrees et al., 2020), to investigate how the polarity and strength of the visual transients affect the suppression of ON and OFF RGCs.

Similar to all previous experiments, we analyzed the modulation index of ON and OFF RGCs separately, using bright probe flashes to analyze ON RGCs and dark probe flashes for OFF RGCs. Consistent with the suppression after texture displacements (Figs. 1e, S3), responses to flashes after luminance steps were strongly suppressed

in both ON and OFF RGCs, and ON RGC suppression outlasted suppression in OFF

RGCs (Fig. 4a, S8a). The two seemingly different experimental paradigms may

therefore trigger similar mechanisms in the retina.

We hypothesized that the response to a luminance step might strongly activate RGCs,

so that the response to a subsequent probe flash would drive the cells into adaptation

or saturation, effectively resulting in suppressed flash responses. At least for the

central component of suppression, this could be a viable mechanism as suppression

is not caused by inhibitory synaptic interactions. If this was indeed the case, then

positive-contrast luminance steps would suppress responses to bright flashes in ON

RGCs, and negative-contrast luminance steps would suppress responses to dark

flashes in OFF RGCs. To test this, we separately analyzed the effects of positive- and

negative-contrast luminance steps on probe flash responses (Fig. 4b). Surprisingly,

the resulting effects were contrary to our adaptation/saturation hypothesis: the

responses of ON RGCs to bright probe flashes were only weakly suppressed after

positive-contrast luminance steps (Fig. 4b, left), but strongly suppressed following

negative-contrast luminance steps (Fig. 4b, right). Similarly, responses of OFF RGCs

to dark probe flashes were weakly suppressed by negative-contrast luminance steps

(Fig. 4b, right), but strongly suppressed by positive-contrast luminance steps (Fig. 4b,

left). Fig. S8 shows the underlying population data for these experiments, and

Figs. S9a, b show the spiking response of a representative ON and OFF RGC,

respectively. While ON RGCs did show a small component of suppression in support

of our adaptation/saturation hypothesis (Fig. 4b left panel, see Fig. S9a left column

and Fig. S10a for a detailed analysis), the dominant suppressive effect in both ON and

OFF RGCs was caused by luminance steps with the opposite contrast as the

subsequent flash.

Such crossover style of suppression would be consistent with mechanisms involving

crossover inhibition via amacrine cells (Werblin, 2010), where activation of OFF

pathways (here: by the negative-contrast luminance step) would inhibit responses in

the ON pathway (here: to the bright probe flash) and vice versa. However, consistent

with our earlier experiments (Fig. 2c, d), suppression in ON RGCs still persisted upon

blocking $GABA_{A,C}$ and glycine receptors (5 μM SR-95531, 100 μM Picrotoxin and 1 μM

Strychnine) (Fig. 4c). We could not calculate a modulation index for OFF RGCs under

these conditions because they did not respond to brief probe flashes in the presence
of the pharmacological agents, and therefore the modulation index was
mathematically undefined. However, in our texture displacement experiments, the
same pharmacological agents (Fig. 2d) had no substantial effect on OFF RGC
suppression. The crossover-style suppression observed in Fig. 4b was therefore
unlikely to be caused by classical crossover inhibition pathways involving amacrine
cells and $GABA_{A,C}$ or glycine receptors.

Central component of suppression results from cone response kinetics and nonlinearities in downstream retinal pathways

So far our experiments suggest that suppression in OFF RGCs (1) is mediated solely
by the central component of suppression that originates in the receptive field center
(Figs. 2e, S7b-e), (2) is predominantly triggered by the interaction between
consecutive stimuli with opposite polarity (Figs. 4b, S8b), and (3) is not caused by
inhibitory amacrine cells (Figs. 2c, 2d, S6). Similar conclusions can be drawn for the
central component of suppression in ON RGCs. Together, these results restrict the
possible cellular substrates for this central suppressive mechanism to the feed-forward
pathway in the retina, namely photoreceptors - bipolar cells - RGCs. We wondered
whether opposite-polarity stimulus-stimulus interactions could already modulate the
responses of photoreceptors themselves. For this, we recorded the output of cones
with an intensity-based glutamate-sensitive fluorescent reporter (iGluSnFR) (Marvin
et al., 2013; Szatko et al., 2020), predominantly expressed in horizontal cells
postsynaptic to cone terminals (Methods). We presented a shortened version of the
luminance step paradigm in which a homogeneous background alternated between a
brighter and darker gray value (Fig. S1c) to induce positive-contrast (+0.4 Michelson
contrast) and negative-contrast (-0.4 Michelson contrast) luminance steps. Dark or
bright probe flashes (100 ms duration, -0.33 or +0.33 Michelson contrast, respectively)
followed the luminance steps at different delays (50, 250, and 2000 ms), with the flash
at 2000 ms serving as baseline.

The luminance steps caused sustained changes in the cones' glutamate output
(Fig. 5a). The transient responses to the probe flashes were superimposed on these
glutamate modulations (Fig. 5b). This superposition was mostly linear and did not

indicate nonlinear effects such as adaptation or saturation. Therefore, when we
isolated the flash responses by subtracting the step responses, the resulting peak
flash responses were only weakly affected by the preceding luminance step (Fig. 5c).
Thus, at the level of the cone output (Fig. 5b), there was hardly any suppression when
only considering the peak of the probe flash responses (Fig. 5c). How does the
suppression observed at the level of RGC output arise from effectively linear cone
responses? The answer lies in other properties of the cone response, such as its
kinetics, which will be captured by downstream retinal pathways.

To demonstrate this, we used a previously published phenomenological model of
retinal processing. In the model (Drinnenberg et al., 2018), RGC responses are
described by passing a "light stimulus" to model photoreceptors, feeding their output
through a set of linear-nonlinear filters to describe the processing in different bipolar
pathways (Methods), and eventually converting these filter outputs into RGC spiking.
Here, as the first step of analysis, we fitted the parameters of the model cone to reflect
our measured data of cone output. The model faithfully explained the observed cone
responses (Fig. S11) and gave us the opportunity to calculate cone responses to
flashes presented at additional time points. This modeled cone output to step-flash
combinations was fed into the model bipolar cells, finally yielding model RGC
responses (Fig. 6). In the model, different RGC types can be described by varying the
bipolar cell filter properties. We first investigated transient RGC responses (Fig 6a-c)
and calculated a modulation index (Fig. 6d) comparable with the modulation index of
our real RGC data. As a control, we also passed the raw cone output data, instead of
the fitted cone model, to the bipolar cells and found qualitatively the same results
(Fig. S12).

The model's ON and OFF RGCs (Fig. 6d) showed crossover-style suppression that
was consistent with the suppression of real RGCs under similar luminance step
experiments (Fig. 4b): the model's ON RGC showed strong suppression to the bright
flashes presented immediately after the negative-contrast luminance step (Fig. 6d,
right), while flashes after the positive-contrast luminance step were only weakly
affected (Fig. 6d, left). Suppression recovered by 200 ms, consistent with the recovery
time for the central component of suppression in real RGCs. Similarly, the model's
OFF RGC showed strong suppression of its response to dark flashes presented

immediately after the positive-contrast luminance step (Fig. 6d, left); suppression was absent in OFF RGC when the dark flash was presented later or after a negative-contrast luminance step (Fig. 6d). The markers on the curves in Fig. 6d correspond to the time points when the flashes were presented to the cones in the experiments depicted in Fig. 5. Model RGC responses to step-flash combinations and flash-induced responses at these time points are shown in Fig. 6a-c.

What mechanism within the model led to the emergence of the suppressive effect in RGC responses, despite the mostly linear response superposition at the cone output? In the model, the bipolar cells have transient filter properties and are driven predominantly by the instantaneous rate of change in the cone output, i.e. its derivative, rather than by the absolute cone output. The response to a probe flash presented immediately after an opposite-contrast luminance step (50 ms) occurred during the initial phase (ramp) of the cone response to the luminance step (Fig. 5b, columns 2 and 3). This causes a much smaller rate of change in cone output and therefore drives the downstream bipolar cells only weakly, resulting in weak or even completely suppressed model RGC responses (Fig. 6c, columns 2 and 3). On the other hand, flashes presented during the steady state phase of the luminance step response (250 ms and 2000 ms in Fig. 5b, columns 2 and 3), or flashes presented immediately after (50 ms) a same-contrast luminance step (Fig. 5b, columns 1 and 4) caused larger instantaneous changes in the cone output, and therefore resulted in relatively stronger spiking (250 ms and 2000 ms in Fig. 6c; 50 ms in Fig. 6c, columns 1 and 4).

If suppression observed at RGC output did indeed result from the weaker rate of change in the cone output, less transient RGCs should show weaker suppression because they reflect the absolute cone output signal more faithfully. To test this, we varied the linear-nonlinear filter of the model bipolar cell to make the retinal pathway less transient (Methods). Indeed, the resulting crossover-style suppression was weaker in model RGCs with less transient response properties, indicated by the negative modulation index slope in Fig. 6e: see ON RGC suppression after negative-contrast steps (blue lines in row 2) and OFF RGC suppression after positive-contrast steps (red lines in row 1); columns represent different time points sampled. Curious if this effect was also present in our real RGC data, we re-analyzed

the data of Fig. 4 and evaluated suppression as a function of RGC transiency
(Methods). The result was congruent with the model prediction: suppression was
weaker in less transient (more sustained) cells (Fig. 6f).

The analyses above suggest that the central component of retinal saccadic
suppression does not have a single site of origin. Instead, it appears to emerge from
the relatively slow kinetics of cone responses (such that flash responses ride on the
initial rising/falling phase of the cone's step response), combined with temporal
properties and nonlinearities of downstream retinal pathways. The downstream
pathways shape the exact strength and time course of RGC suppression. We term
this novel retinal processing motif "dynamic reversal suppression": "dynamic" because
of the necessarily tight temporal link between responses to the two consecutive stimuli;
and "reversal" because the response to an event gets suppressed when it reverses
the cone output triggered by the first stimulus.

Generalization to other species

Retinal saccadic suppression, at least its central component, was triggered by
stimulus-stimulus interactions (Figs. 4, 6, S8), governed by general retinal signal
processing, without the need for any specialized "saccadic suppression" circuit. It is
likely that such general processing is conserved across species. Indeed, we observed
quantitatively similar retinal saccadic suppression in pig ON and OFF RGCs
(Fig. S13), including the dependency on background texture statistics. Interestingly,
like in mouse, OFF RGCs in pig retina also recovered faster than ON RGCs,
suggesting that the surround component of suppression was also present in pig ON
RGCs. In an additional experiment, we also recorded the activity of RGCs from ex vivo
macaque retina while subjecting it to a shorter version of the luminance step paradigm
(Fig. S1b) with fewer conditions than in the original paradigm. Our results (Fig. 7)
provide an indication that macaque RGC responses to probe flashes, following
luminance steps, are suppressed in a way similar to mouse retina. However, more
data will be required to determine the population trend and for characterizing the
dictionary of response modulations in macaque retina.

Downstream visual areas may modulate retinal saccadic suppression

Given the similarities we previously described between retinal and perceptual saccadic suppression (Idrees et al., 2020), it was tempting to test whether the crossover style of suppression, observed in the retina (Fig. 4b), was also reflected in perception. We therefore conducted human psychophysics experiments where we asked human subjects (N = 5) to maintain saccade-free fixation, while we simply changed the luminance of the homogenous background to a brighter (0.3 to 0.56 Michelson contrast) or darker (-0.3 to -0.56 Michelson contrast) background (Fig. 8a; Methods). At random times relative to the luminance step, we presented a dark (-0.033 Michelson contrast) or bright probe flash (+0.033 Michelson contrast), at one of four locations in the subjects' field of view. At trial end, the subjects were asked to localize the probe flash.

Irrespective of the step → flash combination, subjects were strongly impaired in their ability to localize the probe flashes presented around the time of the luminance step (Fig. 8b). Most interesting in this context was the combination of negative-contrast luminance steps with dark probe flashes. In mouse retina, even though OFF RGCs did show weak suppression to this combination (Fig. S8b, inverted histograms in row 2), this effect was virtually absent at the population level (Fig. 4b, right panel). In human perception, however, this combination led to strong suppression (Fig. 8b, right panel). We cannot exclude that stronger retinal suppression to this specific combination might be present under different light or stimulus conditions. Another possibility is that it might be more pronounced in the retina of humans and other primate species (Fig. 7). Nonetheless, visual mechanisms of suppression certainly exist in higher visual brain areas (Idrees et al., 2020). Perceptual suppression after same-contrast stimulus combinations may arise from processing in these higher visual brain areas, which may modulate and complement retinal saccadic suppression to achieve robust effects at the perceptual level.

Discussion

For most RGCs that we recorded, responses to brief probe flashes were strongly suppressed when presented after saccade-like texture displacements across the retina. In fact, similar suppression occurred when texture displacements were replaced

by sudden uniform changes in background luminance, suggesting that suppression was caused by rather generic mechanisms, triggered by visual transients across the retina, rather than specialized suppression circuits that react to saccadic image motion. We found that the suppression strength depended on four main factors: (1) strength of the visual transients, governed by the statistics of the background texture or the contrast of the luminance step; (2) elapsed time following the visual transient; (3) RGC polarity (ON vs. OFF RGCs); and (4) RGC response properties (RGCs transiency). Stronger visual changes, elicited either by coarser textures or larger luminance step contrasts, caused stronger suppression, peaking approximately 50 ms after the stimulus offset (Figs. 1e, 4). The recovery times depended on RGC polarity: OFF RGCs recovered by 250 - 350 ms whereas suppression in ON RGCs lasted for up to 1 s (Figs. 1e, 4). The suppression was stronger in more transient RGCs (Fig. 6e, f). We identified at least three components of retinal saccadic suppression, with distinct spatial origins, defined as central, surround, and global components (Fig. 3). These components were mediated by different underlying mechanisms.

The central component was the only source of suppression in OFF RGCs, and the dominant source in ON RGCs for time points immediately after a full-field saccade or luminance step. This component was short-lived (~250-350 ms) and originated from a cell's receptive field center. It was triggered by opposite-polarity stimulus-stimulus interactions, which naturally occur during saccades and other forms of sequential visual stimulation. The relatively slow temporal dynamics of the cone photoreceptor responses to two opposite-polarity consecutive stimuli were the basis for subsequent non-linear processing, ultimately suppressing RGC response to the second stimulus. Therefore, the central component of suppression cannot be pinpointed to one specific site but seems to be an emergent property. We call this novel processing motif "dynamic reversal suppression". Such a mechanism, where the cone response itself remains linear, but nonetheless forms the basis for subsequent nonlinear response modulation, is clearly different from adaptation (Clark et al., 2013) or desensitization (Baylor and Hodgkin, 1974) mechanisms within the cones, which would evoke nonlinear responses of the cones themselves.

The suppressive mechanism of the central component also suggests that perceptual saccadic suppression is derived, at least in part, from the inherent response properties

of photoreceptors, the very first cell in the visual processing cascade. In fact, this early implementation could also explain why we observed suppression in most RGCs we recorded from (Figs. S3, S8, S13), covering a wide spectrum of response properties and therefore presumably many RGC cell types (see Supplementary Fig. 6 of Idrees et al., 2020). Still, our results suggest that the suppression initiated at the level of cone photoreceptors is translated differently by the different parallel pathways in the retina, leading to variability in response suppression at the ganglion cell level (such as the stronger suppression in more transient RGCs, Fig. 6e, f). Further cell type classification will be required to relate the degree of modulation with pathway specificity. The type and degree of modulation might also differ across species, even though we see qualitatively similar suppression in mouse, pig, and macaque RGCs.

ON RGCs, in addition to this central component, derive suppression from two more components. First, the global component is a fast but short-lived (~250-350 ms) component, caused by inhibition via GABAergic wide field amacrine cells, triggered by global image changes and carried to the RGC from as far as the cell's periphery (Figs. 2b, S5). This likely belongs to the same class of circuits as described by others in the context of peripheral stimulation (Baccus et al., 2008; Brooks and Holden, 1973; Enroth-Cugell and Jakiela, 1980; Roska and Werblin, 2003; Zhang et al., 2012). The role of this global component in perceptual saccadic suppression might not be so prominent as it is masked by more local components during full-field saccades (Fig. 2c, 2d, S6). It is nevertheless another mechanism to process global motion, in addition to the several others (Wei, 2018). It may, for example, play a role in perceptual modulations during pursuit eye movements (Schütz et al., 2011). Here, the central component of suppression will not be triggered in RGCs whose receptive field centers are locked to the tracked object; but these RGCs will still be suppressed by the global component.

The second additional component is the surround component. It seems to act with a delay of ~200 ms and can last for up to 1000 ms. The spatial origins of this component are not very well understood, but our data indicates that it presumably arises from the immediate surround of a cell's receptive field. Additionally, similar to the central component, it does not rely on $GABA_{A,C}$ or glycine receptors (Fig. 2c, 2d, S6), but the exact mechanisms remain to be explored. Possible mechanisms could involve

negative feedback of horizontal cells onto the cones (Drinnenberg et al., 2018; Kemmler et al., 2014). This slower component most likely contributes to the longer recovery times observed in ON cells. Interestingly, visual masking in cat LGN also lasts longer in ON versus OFF cells (Schiller, 1968), which may be a consequence of the effects we describe here in the retina. While this surround component plays an important role in shaping RGC and downstream neural responses following a visual transient, its contribution to perceptual saccadic suppression can also be disputed. This is because, during real saccades, eye-movement related signals (e.g., corollary discharge) shorten the duration of suppression caused by visual mechanisms (Idrees et al., 2020), such that the long-lasting surround component may not critically shape perception. Yet another additional component of suppression, based on saturation-like mechanisms (Fig. 4b left, S10a), was found only in ON RGCs. It is possible that this component originates at the level of bipolar cells as response saturation has been observed predominantly in ON bipolar cells but not in OFF bipolar cells (Schreyer and Gollisch, 2020). In summary, while Fig. 3 summarizes the three spatial components of suppression and their temporal properties, these components in turn can have further sub-components. An example is the saturation-type of suppression in ON RGCs which acts in addition to the dynamic contrast reversal. Both can be assigned to the central component.

Irrespective of which components of retinal saccadic suppression contribute towards perceptual saccadic suppression, our results show that retinal responses to stimuli following visual transients are modulated concurrently by several mechanisms (Fig. 3). Additional mechanisms might exist under different stimulus conditions. From the perspective of retinal visual feature processing, this would be consistent with how multiple mechanisms concurrently process other visual features in the retina, such as motion (Wei, 2018). Consequently, our study is not only relevant from the perspective of saccadic suppression, but also contributes new insights into retinal signal processing during the dynamics of natural vision where local and global visual transients are frequent. Our visual system might exploit this retinal signal processing to enhance visual performance during high-acuity tasks. For example, while threading a needle, microsaccades will jitter the needle and its background across the retina. RGCs exposed to the background will experience stronger visual transients than

RGCs exposed to the needle. As a result, RGCs 'seeing' the background might show
suppressed responses following microsaccades, perceptually blurring out the
background. This relative enhancement of the fixated object by background
suppression could be valuable for visual tasks such as object recognition.

Suppression during different paradigms of sequential visual stimulation has regularly
been observed in downstream visual areas, e.g. when studying saccadic suppression
(Idrees et al., 2020), visual masking (Breitmeyer, 2007) or neural adaptation (Mayo
and Sommer, 2008). The similarities between the observations there and the
suppression we observe in the retina suggest that the retina may be a common neural
substrate of these seemingly different types of suppression, unifying their underlying
mechanisms. Indeed, it is remarkable that an elementary property of retinal saccadic
suppression, i.e. its dependence on the scene statistics (Fig. 1e), is clearly preserved
all the way to perception (Idrees et al., 2020). On the other hand, retinal saccadic
suppression alone cannot explain all perceptual suppressive effects. For example,
suppression before a saccade (or a luminance step) is observed perceptually (even in
the absence of eye movements, Fig. 8), but not in the retinal output (Idrees et al.,
2020), and therefore most likely originates from visual processing beyond the retina.
Similarly, not all properties of retinal saccadic suppression were preserved in
perception. The crossover style of suppression observed in the retinal output of mice
(Fig. 4b) was not observed perceptually in humans (Fig. 8). The more far-reaching
perceptual suppression may be the result of additional visual processing beyond the
retina (Shapley and Enroth-Cugell, 1984). Alternatively, our results, despite their
robustness, may not capture the full array of retinal processing. For one, our stimulus
conditions in mouse retina experiments may not have been adequate to capture all
retinal suppressive effects. Further, the retinal output of humans and other primate
species (Fig. 7) might differ from mouse retina in this respect.

Retinal saccadic suppression, or rather sequential-stimuli suppression, is only one
way the retina alters its output during dynamic vision. Other forms certainly co-exist,
such as brief changes in RGC polarity following peripheral shifts (Geffen et al., 2007),
or sensitization of some RGC types following a change in background luminance
(Appleby and Manookin, 2019). These and several other studies (Amthor et al., 2005;
Barlow et al., 1977; Enroth-Cugell and Jakiela, 1980; Krishnamoorthy et al., 2017;

Noda and Adey, 1974; Passaglia et al., 2009; Roska and Werblin, 2003), together with ours, demonstrate the complex image processing capabilities in the retina to facilitate downstream visual processing for the ultimate service of perception during natural vision. Looking forward, the detailed characterization of retinal output provided here paves the way to investigate the interactions between retinal and extra-retinal (visual and non-visual) mechanisms of saccadic suppression, to further our understanding of how the visual system maintains stability in the face of constant disruptions.

Acknowledgements

We thank Elisabeth Gustafsson for providing technical support; Andreas Hierlemann for providing the HiDens CMOS MEA system and helping establish our high-density MEA recordings; Roland Diggelmann for helping in setting up the pipeline (including providing code) for automatic spike sorting of high-density MEA recordings; Martin Schenk at the Department of Experimental Surgery, Tübingen, for providing us domestic pig eyes. Our work was supported by funds of the Deutsche Forschungsgemeinschaft (DFG) to the Werner Reichardt Center for Integrative Neuroscience (EXC 307) and to T.A.M. (MU3792/1-1 and MU3792/3-1). T.A.M. received support from the Tistou and Charlotte Kerstan Foundation. S.I., T.A.M. and Z.M.H. were also supported by an intra-mural funding program (Project 2013-05) of the Werner Reichardt Center for Integrative Neuroscience. F.F. was supported by a Swiss National Science Foundation Ambizione grant (PZ00P3_167989) and Swiss National Science Foundation Eccellenza grant (PCEFP3_187001). M.P.B. and Z.M.H. were further funded by the SFB 1233 on "Robust Vision" (DFG) - project number 276693517. K.F. was supported by the Bundesministerium für Bildung und Forschung (BMBF, 01GQ1002) and the Max Planck Gesellschaft (MPG, M.FE.A.KYBE0004).

Author contributions

TAM conceptualized the study; SI and TAM designed the overall study; FF developed and supervised computational modeling; SI, MPB, AK, KF, ZMH, FF and TAM designed experiments; SI performed ex vivo mouse and pig retina experiments; AK performed ex vivo macaque retina experiment; MPB performed human psychophysics experiments; KF performed viral injections; MK performed cone imaging experiments;

640   SI and FF implemented computational model; All authors analyzed and interpreted the
641   data and wrote the manuscript.

Declaration of interests

643   The authors declare no competing interests.

Figures

Figure 1

a Saccade paradigm

b Response to saccades-alone

c Response to saccades followed by probe flashes

d Flash-induced responses

e Modulation index (population)

Figure 1 Similarities and differences in retinal saccadic suppression across ON and OFF RGCs.

a. RGC action potentials were recorded from ex vivo retinae placed on multielectrode arrays. Saccades were mimicked by displacing a texture projected onto the retina (blue arrows in left panel indicate texture displacement paths). The texture remained static for 2 s and was then displaced over 100 ms (blue outlines) followed by a brief probe flash (here bright probe flash is depicted, orange outline). Each trial consisted of 39 such successive saccade-flash sequences (Fig. S1a).

b,c. Average activity (firing rate) of an example ON RGC (left column) and OFF RGC (right column) to 39 saccade sequences not followed by a probe flash (**b**), and to 39 saccade sequences followed by probe flashes at different delays after saccade onset (**c**). Blue window: timing of saccades; orange markers: timing of probe flashes.

d. Isolated flash-induced responses (firing rate) of the same RGCs obtained by subtracting responses to saccades-alone (**b**) from responses to saccades followed by probe flashes (**c**). Lines connecting the response peaks highlight the time courses of retinal saccadic suppression relative to baseline flash-induced responses. Numbers above each response peak represent the modulation index which quantifies how much the probe flash response is modulated by the preceding saccade (Methods, negative modulation indices correspond to suppressed flash-induced responses).

e. Population modulation index (mean ± s.e.m.) of ON (light gray) and OFF (dark gray) RGCs, for different background textures with different spatial scales (left to right: fine to coarse). The number of ON and OFF RGCs in the population varied between 68 and 574 for different flash times and textures (see Fig. S3 for exact numbers and relevant statistics). Hash symbols: significant modulation difference between ON and OFF RGCs ($p < 10^{-4}$, two-tailed Wilcoxon rank-sum test).

Figure 2

Significant difference in modulation across groups (p < 0.01)

Figure 2 Spatial origins of retinal saccadic suppression.

a. Spatial layout of the visual stimulation paradigm used in experiments to probe the global component of suppression. Saccades were presented either full-field (left; same as in Fig. 1) or in the periphery (right), where a 1000 x1000 μm^2 mask (intensity: mean luminance of texture) covered at least 2-σ of the 2D Gaussian fit to the RGC receptive fields (Fig. S5b).

b. Population modulation index (mean ± s.e.m.) of ON (top) and OFF (bottom) RGCs for full-field saccades condition (thick gray lines, same as Fig. 1e rightmost panel; N = 68 to 574 RGCs (see Fig. S3 for exact numbers)); periphery saccades condition (thin gray lines; N = 91 ON RGCs, N = 56 OFF RGCs); and periphery saccades condition in the presence of GABA$_A$ receptor blocker (5μM SR-95531; green lines; N = 62 ON RGCs, N = 35 OFF RGCs). Blue window shows the timing of the saccade. In these experiments, we used a coarse background texture (300 μm spatial scale). Timing of probe flashes: 50 and 150 (only for full-field saccade), 117, 200, 350, 600 and 2100 ms (baseline) after saccade onset.

c,d. Population modulation index (mean ± s.e.m.) of ON (top) and OFF (bottom) RGCs for full-field saccades without any pharmacological agents (gray lines; N = 82 ON RGCs, N = 30 OFF RGCs) and with GABA$_{A,C}$ receptor blockers 5 μM SR-95531 + 100 μM Picrotoxin (**c**; green lines), and for a subset of RGCs where we additionally

blocked glycine receptors using 1 μM Strychnine (**d**; green lines; N = 51 ON RGCs, N = 13 OFF RGCs). In these experiments, we used a coarse background texture (150 μm spatial scale). Probe flashes were presented at 117 ms, 150, 200, 350, 600, 1100 and 2100 (baseline) after saccade onset.

e. Spatial layout of the visual stimulation paradigm used in experiments to probe local components of suppression. Saccades and flashes were presented in 100 x 100 μm^2 square regions, separated by 100 μm gaps with mean overall luminance. **Left:** Saccades and flashes were presented in all regions. **Right:** Saccades and flashes were presented in alternate regions; only cells with receptive fields (RFs) in the non-saccade regions (orange) were analyzed (black ellipse: 1-σ of the 2D Gaussian fit to an example RGC receptive field). Consequently, saccades were excluded from at most ~300 x 300 μm^2 of a cell's RF center. In these experiments, we used a coarse background texture (150 μm spatial scale).

f. Population modulation index (mean ± s.e.m.) of ON (top; N = 32) and OFF (bottom; N = 38) RGCs for saccades and flashes in all regions (thick lines) or saccades excluded from RGC RF center (thin lines). Red arrow indicates significant loss in suppression in ON RGCs for early flashes at 117 and 150 ms upon excluding saccades from RF center (p = 0.0016 and p = 0.002 respectively; two-tailed Wilcoxon rank-sum test).

In all panels, hash symbols indicate statistically significant difference between groups (p < 0.01, two-tailed Wilcoxon rank-sum test).

 Figure 3

Figure 3 Schematic summarizing the spatial origins of retinal saccadic suppression.
Saccadic suppression in OFF RGCs (right) is mediated primarily by the central component of suppression (blue) that originates from the cells' receptive field center. ON RGCs (left) get suppressed from two additional components: First, the fast but short-lived global component (red), mediated by GABAergic inhibition, that originates from as far as the cells' periphery. This global component has a similar temporal profile as the central component. It is however weaker than the central component and acts in parallel to it, indicated by the red arrows parallel to blue arrows in the total suppression schematic. Second, the delayed but long-lasting surround component (green), which might originate from the cell's immediate surround. The central component and surround component do not depend on classical GABAergic or glycinergic inhibitory pathways. The differences in the suppression recovery time in ON and OFF RGCs was mainly due to this surround component acting on ON RGCs. Inset shows the legend for arrow schematics. Length of the arrows represent suppression strength; spread of the arrows show the temporal profile of suppression.

Figure 4

Figure 4 Suppression following luminance steps.

a. Population modulation index (mean ± s.e.m.) of ON (light gray, N = 259) and OFF (dark gray, N = 107) RGCs for probe flashes following luminance steps (blue line). Modulation index for each RGC was based on its average response to 56 or 156 luminance step sequences (Fig. S1b) spanning a contrast range of -0.5 to +0.5 Michelson contrast (Methods). Probe flashes were presented at 17 ms, 33, 50, 100, 250, 500, 1000 and 2000 (baseline) after luminance steps. Probe flash responses were suppressed in both ON and OFF RGCs, with similar time course and recovery as in the saccade paradigm with textures (Fig. 1e). Error bars are not visible due to small s.e.m.

b. Same as in **a**, except that the modulation index for each RGC was separately based on average responses to probe flashes after positive-contrast luminance steps (left panel; +0.03 to +0.5 Michelson contrast), and after negative-contrast luminance steps (right panel; -0.03 to -0.5 Michelson contrast). Underlying population data is shown in Fig. S8.

c. Same as in **b**, for a subset of ON RGCs (N = 115) in control conditions (light gray lines) and with $GABA_{A,C}$ and glycine receptors blocked (green lines; cocktail of 5μM SR-95531, 100 μM Picrotoxin and 1μM Strychnine). Hash symbol: significant difference between modulation of ON and OFF RGCs in **a** or between ON RGCs without and with pharmacological blockers in **c** ($p < 10^{-4}$, two-tailed Wilcoxon rank-sum test).

Figure 5

Figure 5 Cone output in response to probe flashes following luminance steps.
a,b. Cone responses (baseline normalized iGluSnFR indicator fluorescence signal) to positive- and negative-contrast (+0.4 and -0.4 on Michelson scale) luminance steps alone **(a)** and to luminance steps followed by probe flashes at 17, 250 and 2000 ms (orange bars below the intensity bar shows timing of probe flashes) **(b)**. Probe flashes were either bright or dark (+0.33 or -0.33 Michelson contrast respectively; 100 ms long). In **b**, responses to step-alone (blue) and individual step → flash pairs (dark gray) are overlaid. Dashed blue lines: timing of luminance step; orange circles: peak cone response to flashes; horizontal dashed line: cone level prior to the luminance step.
c. Flash-induced responses, isolated by subtracting luminance step alone responses (blue) from individual composite luminance step-probe flash responses (dark gray) in **b**. Lines connecting the response peaks highlight the time courses of suppression relative to baseline flash-induced (2000 ms) responses.

Figure 6

Figure 6 Model RGC responses to probe flashes following luminance steps.

a,b. Spiking response of model ON (columns 1-2) and OFF (columns 3-4) RGCs to luminance steps alone (**a**, blue) and to luminance steps followed by probe flashes (**b**, black) at 50, 250 and 2000 ms (different rows; analogous to real RGCs in Fig. 1b, c). Luminance steps are depicted by intensity bars in **a**. First column in each cell type: responses following a positive-contrast luminance step; second column: responses following a negative-contrast luminance step. Vertical blue lines: timing of luminance step; orange bars: timing of probe flashes. Note the ON RGC and OFF RGC did not spike in response to negative-contrast (column 2) and positive-contrast (column 3) luminance steps respectively.

c. Flash-induced responses, after subtracting **a** from **b**, overlaid to show the modulation of probe flash responses at different times (analogous to real RGCs in Fig. 1d). Lines connecting the response peaks highlight the time courses of suppression relative to baseline flash-induced responses (2000 ms).

d. Modulation indices for probe flashes in ON (light gray) and OFF model RGCs (dark gray), following positive-contrast (left panel) and negative-contrast (right panel) luminance steps. Modulation indices were calculated based on model responses to probe flashes presented at 10 ms intervals after luminance steps, and baseline as response to a flash at 2000 ms. Circle markers indicate modulation indices based on probe flashes at 50 and 250 ms shown in **b, c**. Cyan and red arrows highlight the suppression of opposite-contrast flashes at 50 ms in ON and OFF RGCs, respectively.

e. Modulation indices of model ON (cyan) and OFF (red) RGCs plotted as a function of RGC transiency. Individual panels correspond to different flash times (chosen from real-RGC recordings, Fig. 4) after positive-contrast (top row) and negative-contrast (bottom row) luminance steps. In **a-d**, transiency parameter was set to 1. Arrows highlight the same data as in **d**.

f. Same as in **e** but for real ON (light gray circles; N = 228; cyan line: linear regression fit) and OFF RGCs (dark gray circles; N = 92; red line: linear regression fit). These RGCs are a subset of the population data shown in Fig. 4b for which we could compute a transiency index (Methods). Consistent with model predictions, suppression after negative-contrast steps was weaker in less transient ON RGCs (bottom row, blue regression line has negative slope) and, after positive-contrast steps, suppression was weaker in less transient OFF RGCs (top row, red regression line has negative slope). Numbers in each panel indicate the slope of the fits and asterisk symbol indicates statistically significant slope (slope ≠ 0, p < 0.01, two-tailed t-test).

Fig. S12 shows model RGC responses based on real cone data of Fig. 5 instead of model cone responses.

    Figure 7

Figure 7 Retinal saccadic suppression in Macaque RGCs.
a. Average activity of an example macaque ON RGC (top) and OFF RGC (bottom) to positive-contrast (left column) and negative-contrast (right column) luminance steps alone (blue traces) and luminance steps followed by probe flashes (black traces). ON RGCs were analyzed for bright probe flashes and OFF RGCs for dark probe flashes. Responses were averaged across the different positive-contrast (0.05 to 0.5 Michelson contrast, N = 10 sequences) and negative-contrast luminance steps (-0.05 to -0.5 Michelson contrast, N = 10 sequences).
b. Median modulation index (thick lines) of macaque ON (light gray) and OFF (dark gray) RGCs for probe flashes presented after positive-contrast (left panel; N = 13 ON RGCs, N = 1 OFF RGC) and negative-contrast luminance steps (right panel; N = 7 ON RGCs, N = 2 OFF RGC). Circles represent modulation indices of individual RGCs. This experiment was performed under scotopic light conditions.

Figure 8

Figure 8 Perceptual suppression following luminance steps.

a. Example visual task trial. Subjects fixated a small spot on a uniform background with a set luminance (pre-step luminance) for a random duration (800-1700 ms). Background luminance then increased (like shown here) or decreased (positive or negative-contrast luminance step, respectively). At one of 5 times relative to the luminance step (-24, -12, 36, 72, or 108 ms), a luminance pedestal (probe flash, 147.8 x 147.8 min arc) was applied for ~12 ms at one of four locations relative to the fixation spot: 7 deg above (shown here), below, to the right, or to the left. The probe flash was brighter (shown here) or darker than the current screen luminance. The background remained at the post-step luminance until the subject responded with the perceived location of the flash, plus an additional 500-1000 ms but without the fixation spot, allowing the subject to relax. The current luminance was the pre-step luminance of the consecutive trial.

b. Performance of human subjects (mean ± s.e.m., N = 5 subjects), to correctly localize a dark (dark gray) or a bright (light gray) probe flash presented at different times relative to positive-contrast (left panel) and negative-contrast (right panel) luminance steps (blue line). Each subject's responses were averaged across the different positive-contrast (0.3 to 0.56 Michelson contrast) and negative-contrast (-0.3 to -0.56 Michelson contrast) luminance steps. Perceptual performance was reduced around the time of luminance steps, reflecting suppression, irrespective of the combination of luminance step polarity and flash polarity. There were no statistically significant differences in suppression of dark and bright probe flashes (two-tailed Wilcoxon rank-sum test). Note that in the right panel, the suppression profile for bright probe flashes almost completely overlaps the suppression profile of dark probe flashes.

 STAR METHODS

 Key resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
AAV2.7m8.hSyn.iGluSnFR	Dalkara lab (Institut de la Vision), (Dalkara et al., 2013)	N/A
Chemicals, Peptides, and Recombinant Proteins		
SR-95531	Sigma	N/A
Picrotoxin	Sigma	N/A
Strychnine	Sigma	N/A
Experimental Models: Organisms/Strains		
Mouse: PV-Cre (strain: B6;129P2-Pvalb$^{tm1(cre)Arbr}$/J)	Jackson laboratories	RRID: IMSR_JAX:008069
Mouse: Thy-S-Y (strain: C57BL/6-tg (ThystopYFPJS))	FMI Basel	N/A
Mouse: BL6 (C57BL/6)	Jackson laboratories	RRID: IMSR_JAX:000664
Pigs: domestic pigs	Department of Experimental Surgery at the Medical Faculty of the University of Tübingen	N/A
Software and Algorithms		
MATLAB	MathWorks	RRID: SCR_001622
Spike sorting algorithm for high-density microelectrode array recordings	(Diggelmann et al., 2018)	N/A
Spike sorting post hoc processing: UnitBrowser	(Idrees et al., 2016)	N/A
Other		
Low density multielectrode array setup: 60pMEA200/30ir-Ti-gt	Multichannel Systems	N/A
High density multielectrode array setup: HiDens CMOS	Hierlemann Lab, (Frey et al., 2009)	N/A
High density multielectrode array setup: MaxOne	MaxWell Biosystems, (Müller et al., 2015)	N/A
High density multielectrode array setup	Chichilnisky et al (Chichilnisky and Baylor, 1999)	N/A

MOM-type two-photon microscope setup	Designed by W. Denk, MPI, Heidelberg; purchased from Sutter Instruments / Science Products	

Contact for reagent and resource sharing

858 Further information and requests for resources should be directed to and will be
859 fulfilled by the Lead Contact, Thomas A. Münch (thomas.muench@gmail.com).

Experimental model and subject details

Animals

We performed electrophysiological experiments on ex vivo mouse, pig and macaque
retinae; and imaging experiments on ex vivo mouse retinae.

Mouse and pig ex vivo retinae experiments were performed in Tübingen, in
accordance with German and European regulations, and animal experiments were
approved by the Regierungspräsidium Tübingen. Macaque ex vivo retina experiment
was performed at Stanford University. Eyes were removed from a terminally
anesthetized macaque rhesus monkey used by other laboratories in the course of their
experiments, in accordance with the Institutional Animal Care and Use Committee
guidelines of Stanford University.

For mouse retina electrophysiology, we used 47 retinae from 15 male and 30 female
PV-Cre x Thy-S-Y mice (*B6;129P2-Pvalb$^{tm1(cre)Arbr}$/J × C57BL/6-tg (ThystopYFPJS))*,
3-12 months old, which are functionally wild type (Farrow et al., 2013; Münch et al.,
2009; Tikidji-Hamburyan et al., 2015). Additionally, we recorded the cone output from
4 retinae obtained from two C57BL/6 male mice, 9 to 10 weeks old. We housed mice
on a 12/12 h light/dark cycle, in ambient temperatures between 20-22 °C and humidity
levels of 40%.

We also replicated experiments on 9 pig retinae obtained from domestic female pigs
after they had been sacrificed during independent studies at the Department of
Experimental Surgery at the Medical Faculty of the University of Tübingen. Pigs were

anesthetized using atropine, azaperone, benzodiazepine (midazolam), and ketamine,
and then sacrificed with embutramide (T61). Before embutramide administration,
heparin was injected.

One experiment was conducted with a retina extracted from a macaque rhesus
monkey.

Humans

Human psychophysics experiments were performed in Tübingen. Human subjects
provided written, informed consent, and they were paid 10 Euros per session of 60
minutes each, for three sessions. Human experiments were approved by ethics
committees at the Medical Faculty of Tübingen University, and they were in
accordance with the Declaration of Helsinki.

In total, we collected data from 5 subjects (24-29 years old; one female).

Method details

Retina electrophysiology (mouse and pig): procedure and laboratory setup

Mice were dark adapted for 4-16 h before experiments. We then sacrificed them under
dim red light, removed the eyes, and placed eyecups in Ringer solution (in mM: 110
NaCl, 2.5 KCl, 1 $CaCl_2$, 1.6 $MgCl_2$, 10 D-glucose, and 22 $NaHCO_3$) bubbled with 5%
CO_2 and 95% O_2. We removed the retina from the pigment epithelium and sclera while
in Ringer solution.

Pigs were dark-adapted for 15-20 min before sacrifice. Immediately after
veterinary-confirmed sacrifice, the eyes were enucleated under dim red light, and the
cornea, lens, and vitreous were removed. Eyecups were kept in CO_2-independent
culture medium (Gibco) and protected from light. We transported eyecups to our
laboratory and cut pieces from mid-peripheral or peripheral retinae. Only those retinae
which showed ganglion cell responses to light stimuli were used in our experiments.

We recorded mouse and pig retinal ganglion cell (RGC) activity using either low- or
high-density multi-electrode arrays (MEAs). The low-density setup consisted of a

perforated 60-electrode MEA (60pMEA200/30ir-Ti-gt, Multichannel Systems (MCS),

Reutlingen, Germany) having a square grid arrangement and 200 μm inter-electrode

distance. We whole mounted an isolated retina on a nitrocellulose filter (Millipore) with

a central 2 x 2 mm hole. The mounted retina was placed with the RGC side down into

the recording chamber, and good electrode contact was achieved by negative

pressure through the MEA perforation. We superfused the tissue with Ringer solution

at 30-34 °C during recordings, and we recorded extracellular activity at 25 kHz using

a USB-MEA-system (USB-MEA 1060, Multichannel Systems) or a memory-card

based system (MEA1060, Multichannel Systems). Data was acquired using MC Rack

version 4.6.2 (Multichannel Systems). A detailed step-by-step approach is provided in

(Reinhard et al., 2014).

The high-density MEA setup consisted of either a HiDens CMOS MEA (Frey et al.,

2009) (developed by the lab of Andreas Hierlemann, Basel, Switzerland) or a MaxOne

system (Müller et al., 2015) (Maxwell Biosystems, Basel, Switzerland). The HiDens

CMOS MEA featured 11,011 metal electrodes with inter-electrode (center-to-center)

spacing of 18 μm placed in a honeycomb pattern over an area of 2 x 1.75 mm. Any

combination of 126 electrodes could be selected for simultaneous recording. The

MaxOne MEA featured 26,400 metal electrodes with center-to-center spacing of 17.5

μm in a grid-like arrangement over an area of 3.85 x 2.1 mm. In this system, up to

1024 electrodes could be selected for simultaneous recordings. For each experiment,

a piece of isolated retina covering almost the entire electrode array was cut and placed

RGC-side down in the recording chamber. We achieved good electrode contact by

applying pressure on the photoreceptor side of the retina by carefully lowering a

transparent permeable membrane (Corning Transwell polyester membrane, 10 μm

thick, 0.4 μm pore diameter) with the aid of a micromanipulator. The membrane was

drilled with 200 μm holes, with center-center distance of 400 μm, to improve access

of the Ringer solution to the retina. We recorded extracellular activity at 20 kHz using

FPGA signal processing hardware. In the case of the HiDens CMOS MEA, data were

acquired using custom data acquisition software, called MEA 1k Scope (developed by

the lab of Andreas Hierlemann, Basel, Switzerland). In the case of the MaxOne MEA,

data were acquired using MaxLab software provided by Maxwell Biosystems, Basel,

Switzerland.

In total, we performed 59 recordings, 47 from mouse and 12 from pig retinae. 24 of the 59 recordings were done using low-density MEAs. Once a basic experimental protocol was established, we shifted to HiDens CMOS MEA providing much higher throughput. 12 experiments were done using this setup. We upgraded to the MaxOne MEA for even higher throughput and did 23 recordings using this setup. A subset of the data collected from 32 of the 59 recordings (20 from mouse and 12 from pig retinae), was also used in our previous study (Idrees et al., 2020). Here, we show further in-depth analysis of that data.

We presented light stimuli to the retinal piece that was placed on the MEA using a DLP projector running at 60 Hz (Acer K11 for low-density MEA experiments and Lightcrafter 4500 from EKB Technologies Ltd. with internal red, green and blue light-emitting diodes, for high-density MEA experiments). 60 Hz is above the flicker fusion frequency of both mouse and pig retinae; therefore, the framerate of these projectors was adequate for our purposes. The Acer K11 projector had a resolution of 800 x 600 pixels covering 3 x 2.25 mm on the retinal surface. Lightcrafter 4500 had a resolution of 1280 x 800 pixels, extending 3.072 x 1.92 mm on the retinal surface. We focused images onto the photoreceptors using a condenser (low-density MEA recordings, illumination from below) or a 5x objective (high-density MEAs, illumination from above). In each case, the light path contained a shutter and two motorized filter wheels with a set of neutral density (ND) filters (Thorlabs NE10B-A to NE50B-A), having optical densities from 1 (ND1) to 5 (ND5). The filters allowed us to adjust the absolute light level of the stimulation.

We measured the spectral intensity profile (in $\mu W \, cm^{-2} \, nm^{-1}$) of our light stimuli with a calibrated USB2000+ spectrophotometer (Ocean Optics) and converted the physical intensity into a biological equivalent of photoisomerizations per rod photoreceptor per second ($R^* rod^{-1} s^{-1}$), as described before (Tikidji-Hamburyan et al., 2015). Light intensities of the projector output covered a range of 3 log units (i.e. 1000-fold difference between black and white pixels, over an 8-bit range). We linearized the projector output, and we used only grayscale images of limited contrast, spanning at most the range from 0 to 120 in the 8-bit range of the projector (see stimulus description below for details). Absolute light intensities were set to the mesopic level, where a stimulus intensity of '30' in our 8-bit DLP projector scale (0-255) corresponded

to 225 to 1000 $R^*rod^{-1}s^{-1}$, depending on the experimental rig used for the experiment

(i.e. different DLP projectors and MEAs). We pooled all data from the different rigs as

separate individual analyses from the individual setups revealed no effects of

recording conditions in the different setups. For experiments of Fig. S4, we also

recorded at scotopic light levels where a stimulus intensity of '30', corresponded to 23

$R^*rod^{-1}s^{-1}$ at scotopic level.

Retina electrophysiology (macaque): procedure and laboratory setup

In one experiment, we recorded the activity of macaque retinal ganglion cells. For this

experiment we used a high-density MEA, as described previously (Chichilnisky and

Baylor, 1999; Field et al., 2007). Following enucleation, the anterior portion of the eye

and vitreous were removed. The eye was stored in a dark container in oxygenated

Ames' solution (Sigma, St. Louis, MO) at 33°C, pH 7.4. Under infrared illumination, a

small piece of retina approximately 1x1 mm, from a retinal region with eccentricity

around 12 mm (4.0-17mm temporal equivalent eccentricity; (Chichilnisky and Kalmar,

2002)), was dissected and placed ganglion cell side down on a MEA for recording.

The retina pigment epithelium remained attached during the recording; the retina was

perfused with oxygenated Ames' solution. A custom planar large-scale MEA (Field et

al., 2007; Litke et al., 2004) with a hexagonal outline of 519 electrodes at 30 μm pitch

was used. Recorded voltages were band-pass filtered, amplified, and digitized at 20

kHz using custom electronics (Litke et al., 2004). Spike sorting process was described

previously (Field et al., 2007).

Visual stimulation was performed with the optically reduced image of a

gamma-corrected OLED microdisplay (eMagin) refreshing at 60.35 Hz focused on the

photoreceptor outer segments. The visual stimulus was delivered through the

mostly-transparent electrode array. The power of each display primary was measured

at the preparation with a calibrated photodiode (UDT Instruments). At the mean

background illumination level, the photoisomerization rates for the rods and the L, M,

and S cones were approximately 29, 9, 9, and 2 $P^*receptor^{-1}s^{-1}$, respectively (see Li

et al., 2014), placing the retina in scotopic regime.

Retina electrophysiology: pharmacology

In several MEA experiments, we used pharmacological agents to block specific receptors in the mouse retina. To block $GABA_A$ receptors selectively, we used 5 μM SR-95531 (gabazine, an antagonist of $GABA_A$ receptors; Sigma). To block both $GABA_A$ and $GABA_C$ receptors, we used 100 μM picrotoxin (an antagonist of $GABA_A$ and $GABA_C$ receptors; Sigma). To block glycine receptors, we used 1 μM Strychnine (antagonist of Glycine receptors).

We first prepared a 1000x stock solution of these pharmacological blockers as follows: SR-95531 was dissolved in water at a concentration of 5 mM; picrotoxin was dissolved in DMSO at a concentration of 100 mM; Strychnine was dissolved in Chloroform at a concentration of 1mM. During the experiments, we pipetted the stock solution to the Ringer solution in a 1:1000 ratio. Wash-in was performed for 20 min.

Cone photoreceptors imaging: laboratory setup

To record the output of cone photoreceptors in the mouse retina, we measured the glutamate release using an intensity based glutamate-sensitive fluorescent reporter, iGluSnFR (Marvin et al., 2013) expressed in horizontal cell processes post-synaptic to cone terminals, using a viral approach. We recorded the cone output from 4 retinae obtained from two C57BL/6 male mice, 9 to 10 weeks old. Below, we reproduce the methods, previously described in Szatko et al (Szatko et al., 2020).

We dark-adapted the mice for ≥1 h before the experiments. They were then anaesthetized using isoflurane (Baxter) and sacrificed by cervical dislocation. The eyes were enucleated and hemisected in carboxygenated (95% O_2 and 5% CO_2) artificial cerebrospinal fluid (ACSF) solution containing (in mM): 125 NaCl, 2.5 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 1.25 NaH_2PO_4, 26 $NaHCO_3$, 20 glucose, and 0.5 L-glutamine (pH 7.4). We then moved the tissue to the recording chamber where it was continuously perfused with carboxygenated ACSF at ~36 °C. In these experiments, ACSF contained ~0.1 μM Sulforhodamine-101 (SR101, Invitrogen) to reveal blood vessels and any damaged cells in the red fluorescence channel (Euler et al., 2019). All procedures were carried out under very dim red (>650 nm) light.

iGluSnFR was expressed in the retina by viral transduction of AAV2.7m8.hSyn.iGluSnFR, generated in the Dalkara lab (Institut de la Vision) as described in (Dalkara et al., 2013; Khabou et al., 2016). The iGluSnFR plasmid construct was provided by J. Marvin and L. Looger (Janelia Research Campus, USA). A volume of 1 μL of the viral construct was injected into the vitreous humour of the mice, anaesthetized with 10% Ketamine (Bela-Pharm GmbH & Co. KG) and 2% xylazine (Rompun, Bayer Vital GmbH) in 0.9% NaCl (Fresenius). For the injections, we used a micromanipulator (World Precision Instruments) and a Hamilton injection system (syringe: 7634-01, needles: 207434, point style 3, length 51 mm, Hamilton Messtechnik GmbH). Imaging experiments were performed 3-4 weeks after injection. In the outer retina, iGluSnFR was predominantly expressed in horizontal cells. As the expression tended to be weaker in the central retina, most scan fields were acquired in the medial to peripheral ventral or dorsal retina.

To record the iGluSnFR signal, we used a MOM-type two-photon microscope setup (designed by W. Denk, MPI, Heidelberg; purchased from Sutter Instruments / Science Products). The design and procedures have been previously described in (Euler et al., 2009, 2019). In brief, the system was equipped with a mode-locked Ti:Sapphire laser (MaiTai-HP DeepSee, Newport Spectra-Physics), two fluorescence detection channels for iGluSnFR (HQ 510/84, AHF/Chroma) and SR101 (HQ 630/60, AHF), and a water immersion objective (W Plan-Apochromat 20x /1.0 DIC M27, Zeiss). The laser was tuned to 927 nm for imaging iGluSnFR. For image acquisition, we used custom made software (ScanM by M. Müller and T. Euler) running under IGOR Pro 6.3 for Windows (Wavemetrics), taking time-lapsed 128 x 128 pixel image scans at 3.9 Hz in the outer plexiform layer (OPL).

For light stimulation in cone imaging experiments, we used the Lightcrafter (LCr; DPM-E4500UVBGMKII), a DLP projector from EKB Technologies Ltd. with internal UV and green light-emitting diodes (LEDs). The light from the DLP projector was focused through the objective. To optimize spectral separation of mouse M- and S- opsins, LEDs were band-pass filtered (390/576 Dualband, F59-003, AHF/Chroma). LEDs of the DLP projector were synchronized with the microscope's scan retrace. Stimulus intensity (as isomerization rate, $P^*cone^{-1}s^{-1}$) was calibrated to range from ~500 (black image) to ~20,000 for M- and S-opsins. In addition, a steady illumination component

of ~10^4 P*cone^{-1}s^{-1} was present during the recordings because of two-photon

excitation of photopigments. The overall light intensity falling on to the retina was

therefore in the low photopic regime. The light stimulus was centered to the recording

field before every experiment. For all experiments, the retinal tissue was kept at a

constant mean stimulator intensity level for at least 15 s after the laser scanning started

and before stimuli were presented.

Human psychophysics: laboratory setup

We used a similar laboratory setup as described previously (Idrees et al., 2020).

Briefly, subjects sat in a dark room 57 cm in front of a CRT monitor (85 Hz refresh

rate; 41 pixels per deg resolution) spanning 34.1 x 25.6 deg (horizontal x vertical).

Head fixation was achieved with a custom head, forehead, and chin rest (Hafed,

2013), and we tracked eye movements of the left eye at 1 kHz using a video-based

eye tracker (EyeLink 1000, SR Research Ltd, Canada). Gray backgrounds in the

luminance step experiment (Fig. 8) were always presented at an average luminance

of 49.84 cd m^{-2}, and the monitor was linearized (8-bit resolution) such that equal

luminance increments and decrements for luminance steps were possible around this

average. In total, we collected data from 5 subjects (24-29 years old; one female). A

subset of the data from 4 subjects was used in our previous study (Idrees et al., 2020).

Here, we perform novel analyses of the complete dataset, in addition to one new

subject.

Visual stimuli: Retina electrophysiology (Figs. 1-4, 7, S3-S10, S13)

In retina electrophysiology experiments, we used two broad visual stimulation

paradigms: a saccade (texture displacements) paradigm (Fig. S1a), and a luminance

step paradigm (Fig. S1b), described in detail below. In different experiments we used

different spatial and/or pharmacological manipulations of these two paradigms.

Saccade (texture displacements) paradigm

<u>Background textures</u>

We created background textures (Fig. S2a) by convolving a random binary (i.e. white or black) pixel image with a two-dimensional Gaussian blurring filter (Schwartz et al., 2012) defined by the kernel

$$G(x,y) = e^{\frac{-(x^2+y^2)}{2\sigma^2}}$$ (Equation 1)

The parameter σ of the kernel influenced the amount of blurring. This resulted in textures having effectively low-pass spectral content (Fig. S2b) with a cutoff frequency depending on σ. For easier interpretation, we define the spectral content of these textures by a spatial scale. Intuitively, the spatial scale approximates the size of the smallest dark and bright image blobs of the texture (Fig. S2a). Quantitatively, the spatial scale is defined as the $2*\sigma$ parameter of the Gaussian blurring filter. We generated textures with four different spatial scales: 25, 50, 150 and 300 μm, that resulted in dark and bright image blobs approximating a range of receptive field sizes between bipolar cells (texture with spatial scale 25 μm, see (Zhang et al., 2012)) and RGCs (textures with spatial scale 150 and 300 μm). In other words, coarser textures matched the resolution of RGCs, and finer textures matched the resolution of one processing stage earlier, the retinal bipolar cells. Calculating power spectra for the textures (Fig. S2b) confirmed that the spatial scale and hence the cutoff frequencies were consistent with this design aim. In different experiments, we used textures of all or a subset of the different spatial scales.

We normalized the pixel intensities in the textures to have uniform variations in luminance around a given mean. We used pixel intensities (from our 8-bit resolution scale) ranging from 0 to 60 around a mean of 30, or ranging from 30 to 90 around a mean of 60 (see sub-section *Saccades and probe flashes* for when each paradigm was used).

<u>Saccades and probe flashes</u>

To simulate saccades in our ex vivo retina electrophysiology experiments, we displaced the texture across the retina in 6 display frames (100 ms at 60 Hz refresh rate). For easier readability, we usually refer to these saccade-like texture

displacements as "saccades". The textures were displaced in each frame by a constant distance along a linear trajectory. While each "saccade" lasted 100 ms, displacement direction was varied randomly for each "saccade" (uniformly distributed across all possible directions), and "saccade" amplitude could range from 310 μm to 930 μm (corresponding to a velocity range of 3100-9300 μm s^{-1} on the retinal surface). In visual degrees, this corresponds to a velocity range of 100-300 deg s^{-1} and displacement range of 10-30 deg in mice, well in the range of observed mouse saccade amplitudes (Sakatani and Isa, 2007). Similar to primates, mice also have oculomotor behavior, even under cortical control (Itokazu et al., 2018). For example, they make, on average, 7.5 saccade-like rapid eye movements per minute when their head is fixed (Sakatani and Isa, 2007) (humans make several saccades per second). We used the same retinal displacement range of 310 μm to 930 μm for pig retinae. To the best of our knowledge, pig oculomotor behavior has not been documented in the literature. However, with their larger eyeball sizes, our translations of the retinal image would correspond to slower saccades (e.g. small saccades in humans and monkeys), which are also associated with saccadic suppression. Moreover, retinal "saccadic suppression" is not critically dependent on the details of movement kinematics, as it is triggered by visual transients (Fig. 4, also see Figs. 4, 5 in Idrees et al., 2020).

Each trial consisted of successive sequences (Fig. 1a, S1a) that combined a "saccade" with a probe flash, as follows: there was first a "pre-saccade" fixation of 2 seconds, where the texture remained static over the retina, then a 100 ms "saccade", followed by "post-saccade" fixation where the texture again remained static over the retina but now with a shifted texture. At a certain time from "saccade" onset (delay d, range: 50 ms to 2100 ms), we presented a probe flash (see below). Following the probe flash, the texture remained static at the post-saccade fixation position for another 2 seconds before the next saccade of the successive sequence occurred. The post-probe-flash fixation of one sequence was therefore also the pre-saccade fixation of the next sequence. This way the texture remained visible during the entire trial, being translated during saccades of the successive sequences. In a single trial, 39 such sequences occurred. In each successive sequence, the direction and amplitude of the saccade was pseudo-randomly determined by the range of allowed saccade amplitudes and directions. The texture always landed at unique locations within a trial.

The end result was that, within a single trial, RGCs experienced a wide spectrum of

saccade amplitudes, directions and contrasts across these 39 saccades. As such, by

analyzing the average effects of the 39 saccades on RGC responses to probe flashes,

we captured a wide range of saccade-induced kinematics and luminance changes

over the RGC receptive fields.

In most cases, the probe flash had a duration of 2 frames (~33 ms). We used 1 frame

(~16 ms) in a subset of earlier experiments (mouse: 709 of 1616 cells; pig: 116 of 228

cells). Results were pooled across these paradigms as the effects were

indistinguishable. The probe flash was a full-screen positive ("bright") or negative

("dark") stimulus transient.

Bright or dark probe flashes could happen in two different ways across our

experiments. The results were indistinguishable between the two ways, so we pooled

results across them. Briefly, in one manipulation, the probe flash replaced the texture

with a homogeneous bright (pixel intensity of 60 in our 8-bit projectors) or dark (pixel

intensity of 0) full-screen (in these experiments, the textures themselves had

intensities ranging from 0 to 60 pixel intensity; see *Background textures* above). This

way, the flash contrast from the underlying background luminance was variable across

space (e.g. a bright flash on a bright portion of a texture had lower contrast from the

underlying texture than the same flash over a dark portion of the texture). In the second

manipulation, the bright and dark flashes were simply luminance increments or

decrements (by pixel values of 30 on our 8-bit projectors) over the existing textures.

This way, spatial contrast relationships in the background textures were maintained.

In these experiments, the textures themselves had a range of 30-90 pixel intensities

and a mean pixel value of 60 (on our 8-bit projectors). Out of the 1616 RGCs that we

analyzed for saccadic suppression across all experiments where texture

displacements were used as saccades (irrespective of the spatial or pharmacological

manipulations), 1129 RGCs experienced such probe flashes, whereas the rest (487

RGCs) experienced the homogenous probe flash. For pig retina recordings, we always

used the homogenous framework. However, in the subset of pig experiments where

the 2-frame probe flash was employed (112 of 228 RGCs), we used a high-contrast

probe flash such that a bright flash would be achieved by first going to 0 in the first

frame of the flash then going to 60 (on our 8-bit projectors) in the next frame (and vice

versa for a dark flash). Again, all data were pooled across these different paradigms
because their outcomes were indistinguishable.

The number of trials required during a physiology experiment depended on the number
of conditions that we ran on a specific day. For example, testing 7 different flash delays
required 15 trials (7 with bright probe flashes, 7 with dark probe flashes, and 1 without
probes). In a given experiment, we always interleaved all the conditions; that is, in any
one of the 15 necessary trials, each of the 39 saccades could be followed by a bright
or a dark probe at any of the 7 delays, or no probe flash at all (Fig. S1a shows
schematic of one such trial). Moreover, we repeated the total number of conditions
(e.g. the interleaved 15 trials) 4 times per session, and we averaged responses across
those repetitions. Since one trial typically lasted for 2 minutes, the example of 15 trials
repeated 4 times lasted for approximately 2 hours. This was usually combined with
additional conditions (e.g. other background texture scales). Therefore, the total
number of saccades shown in any given experiment could be computed by #trials x 39
saccades per trial x #textures x #repetitions. A typical experiment lasted 10-12 hours.
If the combination of conditions would have required even longer recordings in a given
experiment, we reduced the number of conditions (e.g. we presented flashes at fewer
delays, or used fewer texture scales).

<u>Full-field saccades</u>
In the full-field saccades experiments, saccades and probe flashes occurred over the
entire retina. This was our main experimental paradigm that we used for characterizing
how saccades modulate RGC responses to probe flashes. This paradigm was also
used as a control in experiments in which we applied different spatial or
pharmacological manipulations of this paradigm to probe for the spatial origins and
mechanisms of saccadic suppression. Further, results from this paradigm served as a
baseline standard across different experimental rigs. This paradigm was used in a total
of 32 retinal recordings (32 retinae from 30 mice). In different recordings, we used a
subset of the texture scales and probe flash delays. This explains the different values
of N seen for different conditions in, for example, Figs. 1e, S3. However, to ensure
comparison, some conditions always overlapped across different recordings. This
paradigm was also used in 12 recordings with retinae from 6 pigs.

 Periphery saccades (global component of suppression)

In this manipulation, we restricted saccades to the RGC's receptive fields periphery

(i.e. its far surround). This spatial manipulation was used to investigate the spatial

origins of the global component of suppression (Figs. 2a-b, S5). We performed 13

recordings (13 retinae from 13 mice) with this paradigm, always with the high-density

multielectrode array system (MaxOne by MaxWell), as it provided a large electrode

area (~2 x 4 mm^2) for the retina to be placed on. The recording region was typically

either a high density block of electrodes (inter-electrode spacing: ~17.5 μm) or a block

with one-electrode spacing (inter-electrode spacing: ~35 μm). The recording region

was selected close to the center of the electrode array. We centered a large square

mask (1000 x 1000 μm^2) over the recording region to restrict the texture and saccade

presentation to the periphery of RGC receptive fields (Fig. S5b). The mask had a

homogenous intensity corresponding to the mean luminance of the texture. At different

times relative to texture displacements, full-field probe flashes were presented, similar

to experiments with full-field texture displacements. The intensity of each pixel of the

stimulus (both the mask and the texture regions) was adjusted for the probe flashes,

either decreased or increased by a pixel value of 30 (on our 8-bit projectors) for dark

and bright probe flashes, respectively. In all periphery saccade experiments, we used

probe flash duration of ~33 ms, and a coarse texture background of spatial scale 300

μm.

Checkerboard mask paradigm (local component of suppression)

In this spatial manipulation we presented saccades and flashes in small square

regions spread equidistantly over the entire retina. Each square region measured

100 x 100 μm^2, separated from adjacent squares by an edge-edge gap of 100 μm.

The gap was kept at mean luminance throughout the experiment. Saccades and

flashes could either be presented in all the regions (similar to full-field saccades,

except for the gap), or in alternate regions (Fig. 2e, S7a), arranged like in a

checkerboard. This paradigm was used to investigate the origins of the local

component of suppression (Figs. 2f, S7a-c). We performed 4 recordings (4 retinae

from 3 mice) with this paradigm, always with the low density MCS MEA rig. In all

experiments with this spatial manipulation, we used probe flash duration of ~17 ms,

and a coarse texture background of spatial scale of 150 μm.

Luminance step paradigm

In this paradigm (Fig. S1b), we used no textures at all. The screen was always a homogenous gray field, and the visual event of a "saccade" was replaced by an instantaneous step to a different gray value. The gray backgrounds had intensities between 30 and 90 (on our 8-bit projector). The instantaneous step in intensity caused either a positive contrast luminance step (in the range of +0.03 to +0.50 Michelson contrast) or a negative contrast luminance step (-0.03 to -0.50 Michelson contrast). This paradigm was used to characterize the stimulus-stimulus interactions that ultimately trigger retinal saccadic suppression (Figs. 4, 7, S8-10). We performed a total of 4 recordings (4 retinae from 4 mice) with this paradigm, always with the high density MaxOne MEA rig. A trial consisted of either 56 or 156 successive sequences (Fig. S1b) that each combined a luminance step with a probe flash, as follows: there was first a "pre-step" fixation of 2 seconds where the retina was exposed to a fixed gray level (analogous to "pre-saccade" fixation in texture displacements), then an instantaneous switch to "post-step" fixation (analogous to "post-saccade"). At a certain time from the luminance step (delay: 17, 33, 50, 100, 250, 500, 1000 or 2000 ms), we presented a 2-frame (~33 ms) dark (-0.33 Michelson contrast) or bright (+0.33 Michelson contrast) probe flash. Some sequences contained no probe flash, the next luminance step then happened 4 seconds after the previous one. In a given experiment, we had 17 trials representing the 17 conditions: 8 flash delays x 2 probe flash polarities + 1 condition with no probe flash. Similar to the saccade paradigm, we always interleaved all conditions; that is, in any one of the 17 necessary trials, each luminance step could be followed by a bright or a dark probe at any of the 8 delays, or no probe flash. Moreover, we repeated the 17 trials at least 4 times.

A shorter version of this paradigm was used in our macaque retina recording (Fig. 7). Here, a trial consisted of 20 successive sequences. The 20 luminance steps induced contrasts in the range -0.5 to +0.5. Flashes of ~33 ms were presented with a delay of 17, 50, 100, 250, 500 and 2000 ms after the luminance step.

Other stimuli

Finally, we used other stimuli unrelated to the main experiments to help us characterize RGC properties (e.g. response polarity, latency, transiency, and spatial receptive fields). These stimuli had the same mean intensities and intensity ranges as

the textures or luminance steps used in each experiment. Below, we describe these stimuli for the condition in which the texture intensities ranged from 0 to 60 pixel intensity (represented as grayscale RGB values in the units of our 8-bit projects). In experiments in which the textures ranged in intensity from 30 to 90, or the luminance step experiment, all intensities reported below were shifted upward by 30. (1) Full-field contrast steps. ON steps: stepping from 0 to 30 (+1 Michelson contrast) and from 30 to 60 (+0.33) for 2 s. OFF steps: stepping from 60 to 30 (-0.33) and from 30 to 0 (-1) for 2 s. (2) Full-field Gaussian flicker, 1 minute. Screen brightness was updated every frame and was drawn from a Gaussian distribution with mean 30 and standard deviation 9. This stimulus was used to calculate linear filters representing the temporal receptive fields of RGCs through reverse correlation (spike-triggered averaging of the stimulus history). (3) Binary checkerboard flicker, 10-15 minutes. The screen was divided into a checkerboard pattern; each checker either covered an area of 55 x 55 μm, 60 x 60 μm, or 65 x 65 μm depending on the recording rig. The intensity of each checker was updated independently from the other checkers and randomly switched between 10 and 50 or 0 and 120. This stimulus also allowed us to calculate linear filters representing the spatial receptive fields of RGCs.

Visual stimuli: Cone photoreceptors imaging (Figs. 5, 6)

For cone imaging experiments, we used a minimalistic version of the luminance step paradigm used in retina electrophysiology. A homogeneous background alternated between a darker (pixel intensity 50 on 8-bit projector) and brighter gray value (pixel intensity 120 on 8-bit projector); the transitions between these two background values represented positive and negative contrast of 0.4 Michelson contrast. At various times after the luminance step (50, 250, and 2000 ms) we presented a probe flash (100 ms duration, +0.33 or -0.33 Michelson contrast). The probe flash at 2000 ms served as the baseline. The next background transition always happened 2 s after the preceding probe flash. The combination of the two luminance steps and the two probe flash polarities yielded a total of four combinations: negative-contrast luminance step followed by dark flash; negative-contrast luminance step followed by bright flash; positive-contrast luminance step followed by dark flash; and positive-contrast luminance step followed by bright flash. A single trial (Fig. S1c) was composed of the four step-flash combinations occurring three times (for the three delays with which the

flash was presented); and the negative- and positive-contrast luminance step without a flash. Within a trial, these conditions were randomized. A trial was repeated three times. The luminance steps and the flashes were presented within a 700 μm disc region centered over the scan field.

Visual stimuli: Human psychophysics (Fig. 8)

In the human psychophysics experiment (Fig. 8), we mimicked the retinal luminance step experiments of Fig. 4. The paradigm (Fig. 8a) was similar to the one described in (Idrees et al., 2020). Subjects fixated a central fixation spot over a gray background that remained there for the entire duration of a trial. The background had one of 8 luminances (22.4, 30.24, 38.08, 45.92, 53.76, 61.6, 69.44, 77.28 cd m^{-2}). After a random initial fixation duration (800-1700 ms after fixation spot onset), the luminance of the background was changed suddenly (in one display frame update) to one of the remaining 7 luminances, inducing a positive-contrast luminance step or a negative-contrast luminance step. In our analysis, we used the luminance steps that induced contrasts in the range +0.3 to +0.56 Michelson contrast and -0.3 to -0.56 Michelson contrast. At one of 5 different possible times relative to the time of the luminance step (-24, -12, 36, 72, or 108 ms), a luminance pedestal (probe flash) was applied briefly for one display frame (~12 ms) at one of four locations relative to display center (7 deg above, below, to the right of, or to the left of center). Note that because the display was rasterized (that is, drawn by the computer graphics board from the top left corner in rows of pixels), the exact flash time and duration depended on the location of the flash on the display (but in a manner like other psychophysical experiments studying the same phenomenon, and also in a manner that is unlikely to affect our results). The luminance pedestal consisted of a square of 147.8 x 147.8 min arc, in which we added or subtracted a value to represent bright and dark probe flashes. We ensured that the contrast of the flash (relative to the currently displayed background luminance) was always the same across all trials: +0.033 for a bright flash, and -0.033 for a dark flash. Following each trial, the fixation spot was removed from the background to allow the subjects to relax. This inter trial period lasted for 500-1000 ms. The next trial happened consecutively, in a way that the current luminance of the background was used as the pre-step luminance. Subjects maintained fixation

1339 throughout all trials (except the inter trial period) and simply reported the locations of
1340 the brief flashes. Each subject performed 3 sessions, with 1120 trials per session.

Quantification and statistical analysis

Data analysis: Retina electrophysiology

MEA recordings preprocessing

Low-density MEA recordings were high-pass filtered at a 500 Hz cutoff frequency
using a tenth-order Butterworth filter. We extracted spike waveforms and times using
thresholding, and we semi-manually sorted spikes using custom software. For
high-density MEA recordings, we performed spike sorting by an offline automatic
algorithm (Diggelmann et al., 2018) and assessed the sorted units using a custom
developed tool, the UnitBrowser (Idrees et al., 2016). We judged the quality of all units
using inter-spike intervals and spike shape variation. Low quality units, such as ones
with high inter-spike intervals, missing spikes, or contamination, were discarded. All
firing rate analyses were based on spike times of individual units. In total, we extracted
3,510 high quality units after the spike sorting (referred to as RGCs from now on), from
recordings of mouse retina. From pig retina recordings, we extracted 376 RGCs and
from macaque retina we extracted 57 RGCs after the spike sorting. However, as we
mention below, only a subset of these could be analyzed for saccadic suppression.

RGCs characterization: Receptive fields, ON-OFF index, Transiency index

We first characterized the properties of RGCs. We calculated linear filters in response
to full-field Gaussian flicker and binary checkerboard flicker by summing the 500-ms
stimulus history before each spike. The linear filters allowed determining cell polarity.
Specifically, the amplitude of the first peak of the filter was used: If the peak was
positively deflected, the cell was categorized as an ON cell; if negatively deflected, the
cell was an OFF cell. ON cells were later always analyzed with respect to their
responses to bright probe flashes, and OFF cells were analyzed with dark probe
flashes. We determined the spatial receptive fields of RGCs by calculating the linear
filters for each region (checker) defined by the binary checkerboard flickering stimulus.
The modulation strength of each linear filter, measured as the standard deviation (s.d.)
along the 500 ms temporal kernel, is an estimate for how strongly that region drives

ganglion cell responses. We fitted the resulting 2D-map of s.d. values with a two dimensional Gaussian and took the 2-σ ellipse (long axis) as the receptive field diameter. For all other figures and analyses, we converted spike times to estimates of firing rate by convolving these times with a Gaussian of σ = 10 ms standard deviation and amplitude 0.25 $\sigma^{-1}e^{1/2}$.

For each RGC, we used responses to full-field contrast steps to calculate an ON-OFF index, a transiency index, and a response latency index. These indices were used to characterize the properties of RGCs that we included in our analyses. The ON-OFF index was calculated by dividing the difference between ON and OFF step peak response by their sum. The resulting index values ranged between -1 (OFF) and +1 (ON) and were then scaled to span between 0 (OFF) and +1 (ON). The transiency index was defined as the ratio of the response area within the first 400 ms and the total response area spanning 2000 ms. The resulting index had a value of 1 for pure transient cells. Response latency was calculated as the time from stimulus onset to 90% of peak response. This value was normalized to the maximum response latency in our dataset to create the response latency index.

Modulation index

To quantify retinal saccadic suppression, we first determined a baseline response, defined as the response to a probe flash approximately 2 s after texture displacement onset or 2 s after luminance step (delay between 1967 to 2100 ms, depending on the specific flash times used in a specific experiment). This baseline response was compared to responses of the same cell to the same flash when it occurred at an earlier time (i.e. closer in time to the "saccade"). Usually, the saccade-like texture displacements themselves caused significant neural responses (saccade-alone response, e.g. Fig. 1b), and the responses to the flashes were superimposed on these "saccade-responses". We therefore first isolated the component of the responses caused by the flashes by subtracting the saccade-alone responses from the composite saccade and flash responses. We refer to this isolated component as the flash-induced responses.

To get a robust estimate of the response to saccades-alone (i.e. without any flashes), we averaged spike rate from before saccade onset up until the next saccade onset for

conditions in which no flash was presented, or until just before the flash onset for conditions in which a post-saccade flash was presented. This was done for each of the 39 successive saccades in a given trial.

We then computed a neural modulation index, ranging from -1 to +1. A value of -1 represents complete suppression of flash-induced responses, whereas +1 indicates "complete enhancement" of flash-induced responses (that is, there was only a response to a flash after saccades, but not to a flash in isolation). A modulation index of 0 meant no change in flash-induced response relative to the "baseline" response. The modulation index of an RGC for a given flash delay d after saccade onset was calculated as $(r_d - r_b)/(r_d + r_b)$ where r_d is the peak firing rate for the flash-component of the response (see above for how we isolated this from the composite saccade+flash response) and r_b is the peak firing rate for the baseline flash response (i.e. the same flash but occurring ~2 s away from any "saccade"; see above). In all cases, peak firing rate was estimated after averaging responses from all repetitions of a given condition (delay d or baseline) for a given RGC, across all saccades. For ON cells, the modulation index was based only on responses to bright flashes, and for OFF cells, it was based on responses to dark flashes. To quantify the modulation at a population level, we averaged the modulation indices of the individual RGCs in that population. For some analyses, we also calculated modulation indices of RGCs for each of the 39 individual "saccades" using the same procedure.

In some cells, individual saccades from the sequence of 39 were discarded. For example, imagine that "saccade No 3" gets discarded. This would happen when the baseline response strength (response to the probe flash with 2 s delay) after saccade No 3 is weak (specifically: peak amplitude less than 60% of the median of all 39 baseline response strengths). We did this to ensure that our modulation indices were not marred by a denominator approaching zero (e.g. if both flash and baseline responses were weak). We did, however, re-include some sequences. For example, if the probe flash with delay 100 ms after saccade No 3 triggered a strong response (specifically: peak amplitude larger than the median baseline response peak across the 39 saccades), then saccade No 3 would be re-included for the condition "100 ms delay". This was done in order to re-include sequences (if discarded by the first step) for which the baseline flash response was weak but a flash after saccades nonetheless

gave a robust response. For example, this could happen if a cell did not respond to

the baseline flash but the saccade enhanced the response to a flash following it.

Finally, to perform statistics, we applied tests both at the level of individual cells and

at the level of the population. At the individual cell level, we determined whether the

modulation index for a probe flash presented at a certain delay was significantly

different from 0 (i.e. "Is the response of this cell modulated by the 'saccade'?"). For

this, we performed a one-tailed sign test of the null hypothesis that the 39 individual

modulation indices (or its subset in case weak sequences were discarded, as

described above) came from a distribution with zero median. The alternative

hypothesis was that the median was below (for negative modulation index) or above

(for positive modulation index) zero. The modulation index was considered significant

(i.e. the flash response was modulated by the saccade) at $p < 0.05$. However, we did

not consider cells significantly modulated if the test had a power $(1-\beta)$ smaller than

0.8, which could happen if we previously had to exclude too many sequences ($N \leq 39$).

At the population level, we determined whether the retinal output as a whole was

modulated by saccades. For this, we performed a two-tailed Wilcoxon signed rank test

of the null hypothesis that the median of the distribution of modulation indices did not

differ from 0. Lastly, we tested whether the population modulation index was

significantly different across populations of ON and OFF RGCs or across different

paradigms. For this, we performed a two-tailed Wilcoxon rank-sum test of the null

hypothesis that the median of the distribution of modulation indices did not differ

across the two populations being tested.

Since the modulation index was based on responses to the brief probe flashes, it could

only be computed for RGCs that did respond to these brief flash stimuli. In our analysis,

we included all such RGCs. Of the spike sorted RGCs across all paradigms, we

included: 2002 of 3510 in mice; 228 of 376 in pigs; and 15 of 57 in macaque.

Saccade (texture displacement) paradigm

<u>Full-field saccades</u>

We analyzed 1010 mouse RGCs (633 ON; 377 OFF) and 228 pig RGCs (197 ON; 31

OFF) for saccadic suppression using the full-field version of the saccade paradigm. A

subset of data from 688 of the 1010 mouse RGCs and a subset of data from all the

228 pig RGCs was presented previously (Idrees et al., 2020). Here, we perform novel

analyses on the complete datasets from the RGCs recorded previously, in addition to

analyzing the newly recorded RGCs. For each RGC, we quantified the modulation

index for a full-field probe flash presented at different times from saccade onset.

A subset of these RGCs were also tested for saccadic suppression while blocking

GABAergic and glycinergic inhibition. For 82 ON and 30 OFF RGCs, we had a direct

comparison with and without $GABA_{A,C}$ blockers (5 μM SR-9553 + 100 μM Picrotoxin)

(Fig. 2c, S6a). For 51 ON and 13 OFF RGCs, we had a direct comparison with and

without $GABA_{A,C}$ blockers in addition to glycine blocker (1μM Strychnine; Fig. 2d, S6b).

In yet another subset of RGCs (72 ON; 49 OFF), we also analyzed saccadic

suppression at scotopic light level (Fig. S4), which was 1 log unit dimmer than the light

level at which all other recordings were performed. For these cells, we had a direct

comparison of responses at scotopic and mesopic levels.

<u>Periphery saccades (global component of suppression)</u>

For each recorded RGC, we computed a masking factor (post hoc) to quantify how

well its receptive field was covered by the 1000 x 1000 μm^2 mask. We first determined

the spatial receptive field center of each RGC (described above in *RGCs*

characterization). The masking factor was defined as the multiple of σ of the two

dimensional Gaussian fit for which the ellipse just touched the mask boundary

(Fig. S7d). Cells with receptive field centers within the mask were defined to have a

positive masking factor while those lying outside were given a negative masking factor.

The magnitude of the factor increased with distance from the edge of the mask. This

way, cells for which the mask covered their receptive field centers and immediate

surround had masking factors > 2 (these were the cells included in analysis shown in

Figs. 2b, S5); cells with a mask covering only the receptive field center had masking

factors between 1 and 1.5. Cells located close to the mask's edge, with masking

factors between -1 to +1, had their receptive field centers partially exposed to the

saccade. Finally, cells lying outside the mask where the receptive field center always

experienced saccades had masking factors < -1. A total of 642 RGCs (401 ON; 241

OFF) from 13 experiments were recorded with this spatial layout of the saccade

paradigm. Cells for which clear receptive fields could not be calculated were excluded

from any further analysis. The exact number of cells for different conditions within this

paradigm are reported in the results section. For each RGC, we calculated the

modulation index for flashes presented at different times from saccade onset, in the

same manner as described under the heading *Modulation index*. Fig. S7e shows the

modulation index of individual RGCs as a function of its masking factor. Only a subset

of RGCs with masking factors in the range -3 to +5 were included in this analysis. The

median modulation index was calculated by taking the median of modulation indices

of RGCs within a 1.2 masking factor window, sampled at intervals of 0.1 (running

median).

In a subset of these experiments, we analyzed the effects of blocking GABAergic

inhibition (using 5 μM SR-95531) on the modulation of probe flash responses in RGCs

with masking factors > 2, i.e. with receptive field centers and immediate surround

effectively masked. This included 62 ON RGCs and 35 OFF RGCs with robust

responses to brief probe flashes with the pharmacological agent; only these RGCs

were analyzed for saccadic suppression under the presence of SR-95531 (Fig. 2b). In

a subset of these RGCs (29 ON; 25 OFF), we had a direct comparison of modulation

indices with the control condition, i.e. periphery saccades in the absence of SR-95531

(Fig. S5d).

Checkerboard mask regions saccades

In experiments where we investigated the local component of suppression

(Figs. 2e-f, S7b, c), we had three stimulus presentation settings (Fig. S7a): (1)

saccade and flash presented in all squares regions; (2, 3) saccades and flashes in

alternate regions of a hypothetical binary checkerboard. For each recorded RGC we

calculated the modulation index for the following two scenarios: saccades and flashes

in all regions (Fig. S7a presentation setting 1); and saccades excluded from the

receptive field center (Fig. S7a presentation setting 1 or 2 depending on the RGCs

spatial position relative to the presentation area). For this, we first calculated the

spatial receptive field center (see heading *RGCs characterization*; receptive field

center was defined as the 1-σ ellipse of the 2D Gaussian fit) for each RGC. We then

calculated the Euclidean distance between the receptive field center and the closest

flash region for both presentation settings 2 and 3. For further analysis, we used the

presentation setting with the shortest distance between a flash region and the

receptive field center. In case this flash region is perfectly centered over the receptive
field center, saccades will be excluded from a region of at most 300 μm diameter
centered over the receptive field center. In most cases, the RGCs were indeed well
centered within a flash region since the center of each region coincided with electrodes
in the low density MEA. Nonetheless, for each RGC we calculated the intersection
between its receptive field area and saccade regions in pixels (where each pixel
corresponded to 3.75 μm on the retinal surface). RGCs for which more than 15% of
their receptive field area intersected with the saccade regions were excluded from
further analysis. In the end, a total of 51 RGCs (32 ON; 38 OFF) from 4 retinae
recordings were used for further analysis of saccadic suppression.

Luminance step paradigm
To quantify retinal "saccadic suppression" with the luminance-step paradigm (Fig. 4,
Fig. 7), we used the same analyses and statistical procedures to those described
above for the saccade (texture displacement) paradigm. The only difference was that
instead of 39 successive sequences in a trial, we now had either 56 or 156 successive
sequences (or 20 in case of macaque retina experiment), spanning a contrast range
of ±0.03 to ±0.5 Michelson contrast. Similar to the texture displacement paradigms,
the modulation index was based on responses to brief probe flashes (~33 ms flash
duration), and it could therefore only be computed for cells that did respond to these
flash stimuli (N = 366 of 668 spike sorted RGCs from 4 mouse retinae; N = 15 of 57
spike sorted RGCs from a macaque retina). The modulation index for ON RGCs
(N = 259 mouse RGCs; N = 13 macaque RGCs) was calculated from responses to
bright probe flashes, and that for OFF RGCs (N = 107 mouse RGCs; N = 2 macaque
RGCs) was calculated from responses to dark flashes. A subset of data from all mouse
RGCs was presented previously (Idrees et al., 2020). Here, we perform novel analyses
on the complete dataset of the same RGCs.

In a subset of luminance step experiments (2 mouse retinae), we analyzed the effects
of blocking GABAergic and glycinergic inhibition on the modulation of probe flash
responses. 115 ON RGCs showed robust responses to the brief probe flashes with
and without the pharmacological blockers. However, none of the OFF RGCs
responded robustly to the baseline probe flash in the presence of pharmacological

blockers and therefore OFF RGCs were excluded in quantification of saccadic
suppression in the presence of pharmacological blockers.

We also analyzed if the modulation of flash-induced responses depended on the
strength of the response to the preceding luminance step. This analysis was done to
establish whether suppression of flash responses resulted from saccade-induced
saturation or adaptation of ganglion cell responses (Fig. S10). For each RGC, we
calculated an association index which quantified the monotonic relationship between
response to the luminance step and response to subsequent flashes. We first binned
the responses across the 56 or 156 step-flash sequences (Fig. S1b) based on the
contrast induced by the luminance step in each sequence. Bin width was set to 0.025
Michelson contrast. Then, within each bin, we averaged the responses to luminance
steps alone (Fig. S9a, b, top row) and to luminance steps followed by probe flashes.
For each probe flash delay and contrast bin, we quantified the strength of the response
induced by the luminance step preceding each probe flash (we integrated the average
response to the luminance steps followed by probe flashes, up until the response to
that probe flash). The modulation index was calculated as usual. We then calculated
the Spearman's rank correlation coefficient (R) between the modulation index and the
response strength induced by the luminance step, across all the contrast bins. This
can be visualized from the insets in Fig. S9. For ease in interpretation of the results,
we termed the resulting correlation coefficient as the association index. Intuitively, this
association index describes the monotonic relationship between the step response
strength and the strength of suppression and can be interpreted as follows: the larger
the magnitude of the association index, the stronger is the monotonic relation between
the two quantities. A negative value indicates that stronger step responses are
associated with decreasing (more negative) modulation indices (i.e. weaker flash
responses → suppression). A positive value indicates that stronger step responses
are associated with increasing modulation indices (i.e. stronger flash responses →
less suppression or even enhancement). In the example cell of Fig. S9a, the
association index has large negative values for flashes immediately after the positive
luminance step, suggesting that a stronger step response is indeed strongly correlated
with stronger suppression of subsequent flashes. A robust calculation of association
index was only possible for luminance steps that activated the RGC (i.e.

positive-contrast luminance steps in ON RGCs, negative-contrast luminance steps in OFF RGCs). Fig. S10 shows the association index for each RGC and flash time as a function of the cell's modulation index.

In Fig. 6f, we plot modulation index as a function of RGC transiency index (see heading *RGCs characterization* for details on transiency index). The RGCs shown in this sub-figure were a subset of the RGCs analyzed with the luminance step paradigm for which we could also compute a transiency index. The relation between RGC transiency and modulation index for each condition was modeled using a linear regression least-squares fit through the ON and OFF RGC population. To determine if the slope of the resulting line was statistically significant non-zero, we conducted a t-test of this slope.

Data analysis: Cone photoreceptors

Quantifying cone responses

We analyzed data from 11 scan fields recorded from four retinae (2 mice). Each scan field was 128 x 128 pixels, which on the retinal surface was 94 x 94 μm^2, 110 x 110 μm^2, or 132 x 132 μm^2, depending on the zoom factor used. In each scan field, we identified regions of interest (ROIs) as a group of neighboring pixels with correlated fluorescence signals in time. Only ROIs with diameters corresponding to the cone axon terminal diameter (3-7 μm) were considered for further analysis. The output signal of the ROIs (baseline normalized iGluSnFR indicator fluorescence signal), represented the changes in glutamate release at the cone terminals. A total of 931 ROIs were extracted from the 11 scan fields. Identifying the ROIs and extracting their output signal were automated using custom IGOR Pro scripts.

Within a scan field, each ROI was sampled every 256 ms (3.9 Hz sampling rate). This interval was greater than the duration of probe flashes (100 ms) in these experiments. Therefore, the measured signal of many ROIs might not capture the peak response to the probe flashes. A conventional upsampling method, such as interpolation, could also underestimate the peak response in this case. However, since all ROIs (within and across scan fields) experienced the same visual stimulus (Fig. S1c), but were sampled at different points in time, we temporally 'stitched' the output from these ROIs.

The resulting signal had a sampling interval of 2 ms, where the signal in a specific time bin was computed from ROIs sampled within that specific time window. In this 'stitching' approach, we first baseline-adjusted the output signal of each ROI for a trial by subtracting the baseline activity (calculated as the average ROI output across 1 s epoch prior to the first luminance step in the trial). Then, for each 2-ms time bin, we averaged the response across those ROIs that were sampled within that time window. This resulted in an output vector of the same duration as the trial but with a sampling interval of 2 ms. The output vector was empty for time bins where no ROIs were sampled. We therefore convolved this output vector with a moving average filter of size 80 ms to fill in the empty time bins and to also smooth out the stitching boundaries (boundaries between time bins filled with output from different ROIs). This method gave a much better temporal resolution than conventional upsampling techniques, robustly capturing the peak for the 100 ms duration probe flash, as shown in Fig. 5b.

The 'stitched' signal was then cut into snippets that captured the relevant responses to our stimulus (e.g. step followed by probe flash). For this, we used stimulation trigger signals that marked the presentation of each luminance step. Each snippet was then baseline-adjusted by subtracting the average response over 800 ms prior to the luminance step. This way the output signal was 0 prior to a luminance step (Fig. 5). The output was averaged across the three repetitions of the trial. The normalized and averaged snippets represented the cone response to a particular stimulus sequence (for example Fig. 5a) and were used to fit parameters for model cones as described in the next section.

RGC model

To describe cone responses (Fig. 5) and RGC responses (Fig. 7) to our luminance step paradigm, we used a phenomenological model of the retina, previously published in Drinnenberg et al., 2018.

The original model related light intensity to retinal ganglion spiking activity by three layers of processing: first, the "light stimulus" was passed to model cone photoreceptors. Their activity was modulated by negative feedback from model horizontal cells. Second, the output of the model cones was passed to six inner retina pathways describing retinal processing by three different ON and three different OFF

1650 bipolar cells (fast, intermediate and slow pathways). Third, the output from the model

1651 pathways were then fed into model RGCs to yield RGC spiking activity. This cascade

1652 modeled RGC spiking in response to the "light stimulus" passed to model cones.

1653 The cone responses were describes as

1654
$$r(t) = \frac{\alpha_c y(t)}{[1 + \beta_c z(t)]} - h(t)$$
(Equation 2)

1655 where

1656
$$r(t) = V(t) - V_{dark}$$
(Equation 3)

1657 $V(t)$ and V_{dark} were the instantaneous and dark membrane potentials of the cone,

1658 respectively, $h(t)$ was the feedback signal from the horizontal cell, and α_c and β_c were

1659 numerical factors. The time-varying functions $y(t)$ and $z(t)$, were related to light input

1660 through linear convolution, as

1661
$$y(t) = \int_{-\infty}^{t} dt' K_y(t - t') I(t')$$
(Equation 4)

1662

1663

1664
$$z(t) = \int_{-\infty}^{t} dt' K_z(t - t') I(t')$$
(Equation 5)

1665

1666

1667 where $I(t)$ was the incident light intensity (or, more precisely, R*/s). The kernels

1668 describing the cone response were given by

1669
$$K_y(t) = \frac{t}{\tau_y} \frac{e^{-\frac{t}{\tau_y}}}{\tau_y}$$
(Equation 6)

1670 and

$$K_z(t) = \gamma K_y(t) + (1 - \gamma) \frac{t}{\tau_z} \frac{e^{-\frac{t}{\tau_z}}}{\tau_z} \qquad \text{(Equation 7)}$$

where τ_z was larger than τ_y, and $0 \leq \gamma \leq 1$ ensured proper normalization. Note that $\int_0^\infty dt' K_y(t - t') = 1$ for all filters. The response of the horizontal cell was described by

$$h(t) = \alpha_c \int_{-\infty}^{t} dt' K_h(t - t') r(t') \qquad \text{(Equation 8)}$$

with

$$K_h(t) = \frac{t}{\tau_h} \frac{e^{-\frac{t}{\tau_h}}}{\tau_h} \qquad \text{(Equation 9)}$$

Here, instead of the cone model parameters used in the published model (Drinnenberg et al., 2018), we re-fitted the parameters of the model cone to reflect our measured data of cone output (Fig. 5b) which yielded faithful fits (Fig. S12) to our experimentally measured cone responses.

All the parameters of the outer retina component (Equations 12 and 13) of our circuit model were fit once to cone responses (Fig. 5b, S11) and then kept unchanged for all simulations reported in Fig. 6. The fitted values are given as follows (original values (Drinnenberg et al., 2018) are reported in brackets):

$\alpha_c = -3.342 * 10^{-5}$ $(-9.602 * 10^{-6})$, $\beta_c = -1.273 * 10^{-6}$ $(-1.148 * 10^{-5})$, $\gamma = 0.842$ (0.764), $\alpha_h = 0.016$ (0.177), $\tau_y = 48.98 \, ms$ (50.64), $\tau_z = 200ms$ (576.9), $\tau_h = 1.232 * 10^3 ms$ (371).

The small α_h suggested that in our cone recordings, horizontal cell feedback had only a minor effect.

In (Drinnenberg et al., 2018) three different retinal pathways were modelled according to

$$b_{p,k}(t) = \left\lfloor -1^k \left(\int_{-\infty}^{t} dt' K_p(t-t')V(t') - \theta_{p,k} \right) \right\rfloor \qquad \text{(Equation 10)}$$

1692

where $p = 1,2,3$ labeled the pathway based on its response properties (1 = fast, 2 = intermediate, 3 = slow), $k = 0$ for OFF pathways and $k = 1$ for ON pathways.

1693
1694

$$\lfloor x \rfloor = \begin{cases} 0, & x < 0 \\ x, & x \geq 0 \end{cases} \qquad \text{(Equation 11)}$$

1695

was a thresholding non-linearity, and $\theta_{p,k}$ acted as a threshold.

1696

The main difference between the pathways was the temporal characteristics of the filters K_p. In the current study we wanted to smoothly vary the transiency of the model ganglion cells. To this end, we based our bipolar pathway on the fast bipolar pathway $(p = 1)$ and modified its temporal characteristics to make it less transient. K_1 represented a high-pass filter which took the derivative of the cone potential on the order of 1 ms. We obtained K_1 by convolving the high-pass filter of the form

$$G(t) = \sin\left(\frac{\pi t}{\mu}\right) \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\left(\frac{t-\mu}{\sigma}\right)^2}, with \; \mu = 3 \; ms, \sigma = 1 \; ms \qquad \text{(Equation 12)}$$

1703

with an exponential function.

1704

$$K_1(t) = \int_{-\infty}^{t} dt' \left(e^{-\frac{t-t'}{\tau_d}} \right) G(t') \qquad \text{(Equation 13)}$$

1705

Higher values of the time constant, τ_d, of the exponential function decreased the pathway transiency. We set $\tau_d = 0.5$ as the default transient pathway (Fig. 6a-d). The 'transiency parameter' shown in In Fig. 6e was obtained by normalizing $15 \geq \tau_d \geq 0.5$ in the range 0 to 1, with 0 being less transient. In the original model (Drinnenberg et al., 2018), $K_1(t) = G(t)$.

The threshold $\theta_{1,k}$ was set to $-1^k \cdot 0.1$.

1711

$b_{1,k}(t)$, the output of fast inner retina models was used as the input to the model RGCs used in this study. The spiking rate of the model RGC was obtained as the thresholded input and a temporally coarse version of the input's derivative,

1712
1713
1714

$$1715 \qquad R_{1,k}(t) = \left| (1-\alpha)b_{1,k}(t) + \alpha\left(\int_{-\infty}^{t} dt'\, K(t-t')\, b_{1,k}(t')\right) - \theta \right| \qquad \text{(Equation 14)}$$

1716

where $K(t)$ was a biphasic filter similar in its form to $G(t)$. The threshold, θ, was a

multiple of the peak response to any given input. We used the same parameters for

the inner retina component as the ones described in the published model (Drinnenberg

et al., 2018):

(i) $\qquad Transient\ OFF,\ R_{1,0}(t)\colon \alpha = 0,\ \theta = 0;$

(ii) $\qquad Transient\ ON, R_{1,1}(t)\colon \alpha = 0,\ \theta = 0;$

All simulations were computed with a 1 ms sampling interval.

Using the above cascade, we calculated model RGC's spike rate in response to the

recorded cone output when subjected to "luminance steps alone" and "luminance

steps followed by probe flash" stimuli (Fig. 6a, b). Similar to real RGC analysis, we

calculated the "flash-induced responses" (Fig. 6c) by subtracting "response to

luminance steps alone" (Fig. 6a) from "response to luminance steps followed by probe

flash" (Fig. 6b). We then calculated the modulation index also in the same way as for

the real RGCs: $(r_d - r_b)/(r_d + r_b)$ where r_d was the peak spiking rate of the flash-induced

response for the flash presented with delay d from the luminance step, and r_b was the

peak firing rate for the baseline flash-induced response (flash at 2000 ms).

As a control, we replaced the model cone responses with the experimentally acquired

cone responses (Fig. 5) thus forming a hybrid model. Before passing the cone

response to the downstream model pathways, we passed it through a low pass filter

to further smooth the fluctuations at the stitching boundaries in order to avoid

discontinuities in calculation of its temporal derivative. For this smoothing, we

convolved the cone output with a moving average filter of size 40 ms. Here, the model

ON RGC responses (Fig. S12a, b, columns 1-2) were calculated using the cone

responses to bright probe flashes (Fig. 5a, b, columns 1-2), and OFF RGC responses

(Fig. S12a, b, columns 3-4) were calculated using cone responses to dark probe

flashes (Fig. 5a, b, columns 3-4). The resulting hybrid model RGC responses (Fig. S12) were consistent with the pure model responses shown in Fig. 6.

Data analysis: Human psychophysics

We analyzed eye movements in all trials and detected saccades using established methods (Chen and Hafed, 2013). We excluded from analysis trials in which a saccade or microsaccade happened anywhere in the interval from 200 ms before to 50 ms after a probe flash. At each flash time, we calculated the proportion of correct trials to obtain time courses of this perceptual measure. We obtained time course curves for each subject individually, and then averaged it across trials and different contrasts of the luminance steps. Reduced proportion of correct trials at any flash time indicated perceptual saccadic suppression. A subset of data from 4 of the 5 subjects was used in our previous study (Idrees et al., 2020). Here we perform novel analyses of the complete dataset.

We applied a two-tailed Wilcoxon rank-sum test to determine if the suppression after luminance steps differed across bright and dark probe flashes.

All data analyses were performed in MATLAB (The MathWorks Inc).

Data and software availability

Requests for data should be directed to and will be fulfilled by the Lead Contact, Thomas A. Münch (thomas.muench@gmail.com).

References

Amthor, F.R., Tootle, J.S., and Gawne, T.J. (2005). Retinal ganglion cell coding in simulated active vision. Vis. Neurosci. *22*, 789–806.

Appleby, T.R., and Manookin, M.B. (2019). Neural sensitization improves encoding fidelity in the primate retina. Nat. Commun. *10*, 1–15.

Baccus, S.A., Olveczky, B.P., Manu, M., and Meister, M. (2008). A Retinal Circuit That Computes Object Motion. J. Neurosci. *28*, 6807–6817.

Barlow, H.B., Derrington, A.M., Harris, L.R., and Lennie, P. (1977). The effects of remote retinal stimulation on the responses of cat retinal ganglion cells. J. Physiol. *269*, 177–194.

Baylor, D.A., and Hodgkin, A.L. (1974). Changes in time scale and sensitivity in turtle photoreceptors. J. Physiol. *242*, 729–758.

Beeler, G.W. (1967). Visual threshold changes resulting from spontaneous saccadic eye movements. Vision Res. *7*, 769–775.

Berry, M., Brivanlou, I., Jordan, T., and Meister, M. (1999). Anticipation of moving stimuli by the retina. Nature *398*.

Breitmeyer, B. (2007). Visual masking: Past accomplishments, present status, future developments. Adv. Cogn. Psychol. *3*, 9–20.

Bremmer, F., Kubischik, M., Hoffmann, K.P., and Krekelberg, B. (2009). Neural dynamics of saccadic suppression. J. Neurosci. *29*, 12374–12383.

Brooks, B., and Holden, A.L. (1973). Suppression of visual signals by rapid image displacement in the pigeon retina: A possible mechanism for "saccadic" suppression. Vision Res. *13*, 1387–1390.

Chen, C.-Y., and Hafed, Z.M. (2013). Postmicrosaccadic enhancement of slow eye movements. J. Neurosci. *33*, 5375–5386.

Chen, C.-Y., and Hafed, Z.M. (2017). A neural locus for spatial-frequency specific saccadic suppression in visual-motor neurons of the primate superior colliculus. J.

Neurophysiol. *117*, 1657–1673.

Chen, E.Y., Marre, O., Fisher, C., Schwartz, G., Levy, J., da Silviera, R.A., and Berry, M.J. (2013). Alert response to motion onset in the retina. J. Neurosci. *33*, 120–132.

Chichilnisky, E.J., and Baylor, D.A. (1999). Receptive-field microstructure of blue-yellow ganglion cells in primate retina. Nat. Neurosci. *2*, 889–893.

Chichilnisky, E.J., and Kalmar, R.S. (2002). Functional asymmetries in ON and OFF ganglion cells of primate retina. J. Neurosci. *22*, 2737–2747.

Clark, D.A., Benichou, R., Meister, M., and Azeredo da Silveira, R. (2013). Dynamical Adaptation in Photoreceptors. PLoS Comput. Biol. *9*.

Dalkara, D., Byrne, L.C., Klimczak, R.R., Visel, M., Yin, L., Merigan, W.H., Flannery, J.G., and Schaffer, D. V. (2013). In vivo-directed evolution of a new adeno-associated virus for therapeutic outer retinal gene delivery from the vitreous. Sci. Transl. Med. *5*.

Diamond, M.R., Ross, J., and Morrone, M.C. (2000). Extraretinal control of saccadic suppression. J. Neurosci. *20*, 3449–3455.

Diggelmann, R., Fiscella, M., Hierlemann, A., and Franke, F. (2018). Automatic spike sorting for high-density microelectrode arrays. J. Neurophysiol. *120*, 3155–3171.

Drinnenberg, A., Franke, F., Morikawa, R.K., Jüttner, J., Hillier, D., Hantz, P., Hierlemann, A., Azeredo da Silveira, R., and Roska, B. (2018). How Diverse Retinal Functions Arise from Feedback at the First Visual Synapse. Neuron 1–18.

Duffy, F.H., and Lombroso, C.T. (1968). Electrophysiological evidence for visual suppression prior to the onset of a voluntary saccadic eye movement. Nature *218*, 1074–1075.

Enroth-Cugell, C., and Jakiela, H.G. (1980). Suppression of cat retinal ganglion cell responses by moving patterns. J. Physiol. *302*, 49–72.

Euler, T., Hausselt, S.E., Margolis, D.J., Breuninger, T., Castell, X., Detwiler, P.B., and Denk, W. (2009). Eyecup scope—optical recordings of light stimulus-evoked

fluorescence signals in the retina. Pflügers Arch. - Eur. J. Physiol. *457*, 1393–1414.

Euler, T., Franke, K., and Baden, T. (2019). Studying a light sensor with light:
Multiphoton imaging in the retina. Neuromethods *148*, 225–250.

Farrow, K., Teixeira, M., Szikra, T., Viney, T.J., Balint, K., Yonehara, K., and Roska,
B. (2013). Ambient illumination toggles a neuronal circuit switch in the retina and
visual perception at cone threshold. Neuron *78*, 325–338.

Field, G.D., Sher, A., Gauthier, J.L., Greschner, M., Shlens, J., Litke, A.M., and
Chichilnisky, E.J. (2007). Spatial properties and functional organization of small
bistratified ganglion cells in primate retina. J. Neurosci. *27*, 13261–13272.

Frey, U., Egert, U., Heer, F., Hafizovic, S., and Hierlemann, A. (2009).
Microelectronic system for high-resolution mapping of extracellular electric fields
applied to brain slices. Biosens. Bioelectron. *24*, 2191–2198.

Fried, S., Münch, T., and Werblin, F. (2002). Mechanisms and circuitry underlying
directional selectivity in the retina. Nature 411–414.

Garvert, M.M., and Gollisch, T. (2013). Local and global contrast adaptation in retinal
ganglion cells. Neuron *77*, 915–928.

Geffen, M.N., De Vries, S.E.J., and Meister, M. (2007). Retinal ganglion cells can
rapidly change polarity from off to on. PLoS Biol. *5*, 0640–0650.

Gollisch, T. (2013). Features and functions of nonlinear spatial integration by retinal
ganglion cells. J. Physiol. Paris *107*, 338–348.

Hafed, Z.M. (2013). Alteration of Visual Perception prior to Microsaccades. Neuron
77, 775–786.

Hafed, Z.M., and Krauzlis, R.J. (2010). Microsaccadic suppression of visual bursts in
the primate superior colliculus. J. Neurosci. *30*, 9542–9547.

Ibbotson, M.R., Crowder, N.A., Cloherty, S.L., Price, N.S.C., and Mustari, M.J.
(2008). Saccadic modulation of neural responses: possible roles in saccadic
suppression, enhancement, and time compression. J. Neurosci. *28*, 10952–10960.

Idrees, S., Franke, F., Diggelmann, R., Hierlemann, A., and Münch, T.A. (2016).
UnitBrowser - A tool to evaluate and post-process units sorted by automatic spike
sorting algorithms. Front. Neurosci. *10*.

Idrees, S., Baumann, M.P., Franke, F., Münch, T.A., and Hafed, Z.M. (2020).
Perceptual saccadic suppression starts in the retina. Nat. Commun. *11*, 1977.

Itokazu, T., Hasegawa, M., Kimura, R., Osaki, H., Albrecht, U.-R., Sohya, K.,
Chakrabarti, S., Itoh, H., Ito, T., Sato, T.K., et al. (2018). Streamlined sensory motor
communication through cortical reciprocal connectivity in a visually guided eye
movement task. Nat. Commun. *9*, 338.

Kemmler, R., Schultz, K., Dedek, K., Euler, T., and Schubert, T. (2014). Differential
regulation of cone calcium signals by different horizontal cell feedback mechanisms
in the mouse retina. J. Neurosci. *34*, 11826–11843.

Khabou, H., Desrosiers, M., Winckler, C., Fouquet, S., Auregan, G., Bemelmans,
A.P., Sahel, J.A., and Dalkara, D. (2016). Insight into the mechanisms of enhanced
retinal transduction by the engineered AAV2 capsid variant -7m8. Biotechnol.
Bioeng. *113*, 2712–2724.

Kleiser, R., Seitz, R.J., and Krekelberg, B. (2004). Neural correlates of saccadic
suppression in humans. Curr. Biol. *14*, 386–390.

Krekelberg, B. (2010). Saccadic suppression. Curr. Biol. *20*, R228–R229.

Krishnamoorthy, V., Weick, M., and Gollisch, T. (2017). Sensitivity to image
recurrence across eye-movement-like image transitions through local serial inhibition
in the retina. Elife e22431.

Krueger, J., and Fischer, B. (1973). Strong periphery effect in cat retinal ganglion
cells. Excitatory responses in ON- and OFF-center neurones to single grid
displacements. Exp. Brain Res. *18*, 316–318.

Li, P.H., Field, G.D., Greschner, M., Ahn, D., Gunning, D.E., Mathieson, K., Sher, A.,
Litke, A.M., and Chichilnisky, E.J. (2014). Retinal Representation of the Elementary
Visual Signal. Neuron *81*, 130–139.

Lin, B., and Masland, R.H. (2006). Populations of wide-field amacrine cells in the mouse retina. J. Comp. Neurol. *499*, 797–809.

Litke, A.M., Bezayiff, N., Chichilnisky, E.J., Cunningham, W., Dabrowski, W., Grillo, A. a., Grivich, M., Grybos, P., Hottowy, P., Kachiguine, S., et al. (2004). What does the eye tell the brain?: Development of a system for the large-scale recording of retinal output activity. IEEE Trans. Nucl. Sci. *51*, 1434–1440.

Marvin, J.S., Borghuis, B.G., Tian, L., Cichon, J., Harnett, M.T., Akerboom, J., Gordus, A., Renninger, S.L., Chen, T.W., Bargmann, C.I., et al. (2013). An optimized fluorescent probe for visualizing glutamate neurotransmission. Nat. Methods *10*, 162–170.

Matin, E. (1974). Saccadic suppression: a review and an analysis. Psychol. Bull. *81*, 899–917.

Mayo, J.P., and Sommer, M.A. (2008). Neuronal Adaptation Caused by Sequential Visual Stimulation in the Frontal Eye Field. J. Neurophysiol. *100*, 1923–1935.

Müller, J., Ballini, M., Livi, P., Chen, Y., Radivojevic, M., Shadmani, A., Viswam, V., Jones, I.L., Fiscella, M., Diggelmann, R., et al. (2015). High-resolution CMOS MEA platform to study neurons at subcellular, cellular, and network levels. Lab Chip *15*, 2767–2780.

Münch, T.A., da Silveira, R.A., Siegert, S., Viney, T.J., Awatramani, G.B., and Roska, B. (2009). Approach sensitivity in the retina processed by a multifunctional neural circuit. Nat. Neurosci. *12*, 1308–1316.

Noda, H., and Adey, W.R. (1974). Excitability Changes in Cat Lateral Geniculate Cells during Saccadic Eye Movements. Science (80-.). *183*, 543–545.

Ölveczky, B.P., Baccus, S.A., and Meister, M. (2003). Segregation of object and background motion in the retina. Nature *423*, 401–408.

Passaglia, C.L., Freeman, D.K., and Troy, J.B. (2009). Effects of remote stimulation on the modulated activity of cat retinal ganglion cells. J. Neurosci. *29*, 2467–2476.

Reinhard, K., Tikidji-Hamburyan, A., Seitter, H., Idrees, S., Mutter, M., Benkner, B.,

and Münch, T.A. (2014). Step-By-Step instructions for retina recordings with

perforated multi electrode arrays. PLoS One *9*, e106148.

Roska, B., and Werblin, F. (2003). Rapid global shifts in natural scenes block spiking

in specific ganglion cell types. Nat. Neurosci. *6*, 600–608.

Ross, J., Morrone, M.C., Goldberg, M.E., and Burr, D.C. (2001). Changes in visual

perception at the time of saccades. Trends Neurosci. *24*, 113–121.

Sakatani, T., and Isa, T. (2007). Quantitative analysis of spontaneous saccade-like

rapid eye movements in C57BL/6 mice. Neurosci. Res. *58*, 324–331.

Schiller, P.H. (1968). Single unit analysis of backward visual masking and

metacontrast in the cat lateral geniculate nucleus. Vision Res. *8*, 855–866.

Schreyer, H.M., and Gollisch, T. (2020). Nonlinearities in retinal bipolar cells shape

the encoding of artificial and natural stimuli. BioRxiv 1–40.

Schütz, A., Braun, D., and Gegenfurtner, K. (2011). Eye movements and perception:

A selective review. J. Vis. *11*, 1–30.

Schwartz, G.W., Okawa, H., Dunn, F. a, Morgan, J.L., Kerschensteiner, D., Wong,

R.O., and Rieke, F. (2012). The spatial structure of a nonlinear receptive field. Nat.

Neurosci. *15*, 1572–1580.

Shapley, R., and Enroth-Cugell, C. (1984). Chapter 9 Visual adaptation and retinal

gain controls. Prog. Retin. Res. *3*, 263–346.

Szatko, K.P., Korympidou, M.M., Ran, Y., Berens, P., Dalkara, D., Schubert, T.,

Euler, T., and Franke, K. (2020). Neural circuits in the mouse retina support color

vision in the upper visual field. Nat. Commun. *11*, 3481.

Tatler, B.W., Wade, N.J., Kwan, H., Findlay, J.M., and Velichkovsky, B.M. (2010).

Yarbus, Eye Movements, and Vision. Iperception. *1*, 7–27.

Tikidji-Hamburyan, A., Reinhard, K., Seitter, H., Hovhannisyan, A., Procyk, C.A.,

Allen, A.E., Schenk, M., Lucas, R.J., and Münch, T.A. (2015). Retinal output

changes qualitatively with every change in ambient illuminance. Nat. Neurosci. *18*,

66–74.

Volkmann, F.C. (1986). Human visual suppression. Vision Res. *26*, 1401–1416.

Wei, W. (2018). Neural mechanisms of motion processing in the mammalian retina.
Annu. Rev. Vis. Sci. *4*, 165–192.

Werblin, F. (2010). Six different roles for crossover inhibition in the retina: correcting
the nonlinearities of synaptic transmission. Vis. Neurosci. *27*, 1–8.

Wurtz, R.H. (2008). Neuronal mechanisms of visual stability. Vision Res. *48*, 2070–
2089.

Yarbus, A.L. (1967). Eye Movements and Vision (Boston, MA: Springer US).

Zaghloul, K.A., Manookin, M.B., Borghuis, B.G., Boahen, K., and Demb, J.B. (2007).
Functional circuitry for peripheral suppression in Mammalian Y-type retinal ganglion
cells. J Neurophysiol *97*, 4327–4340.

Zhang, Y., Kim, I.-J., Sanes, J.R., and Meister, M. (2012). The most numerous
ganglion cell type of the mouse retina is a selective feature detector. Proc. Natl.
Acad. Sci. *109*, E2391–E2398.

Zuber, B.L., and Stark, L. (1966). Saccadic suppression: Elevation of visual
threshold associated with saccadic eye movements. Exp. Neurol. *16*, 65–79.

1943