Discrepancy between repetition suppression and pattern analysis provides new insights into the role of M1 and parietal areas in sequence learning

Eva Berlot[1,*], Nicola J. Popp[1], Scott T. Grafton[2-3], & Jörn Diedrichsen[1,4,5, *]

[1] The Brain and Mind Institute, University of Western Ontario, Canada
[2] Department of Psychological and Brain Sciences, University of California, Santa Barbara, CA, USA
[3] Institute for Collaborative Biotechnologies, University of California, Santa Barbara, CA, USA
[4] Department of Statistical and Actuarial Sciences, University of Western Ontario, Canada
[5] Department of Computer Science, University of Western Ontario, Canada

Corresponding authors:
eva.berlot@gmail.com and jdiedric@uwo.ca

Abstract

How does the brain change during skill learning? We previously conducted a longitudinal fMRI motor sequence learning study and, using multivariate techniques, found learning-related changes in premotor and parietal areas, but not in the primary motor cortex (M1) (1). However, a study using repetition suppression (RS) had previously suggested that M1 represents learned sequences. Here we replicate this discrepancy in a single experiment, allowing us to investigate the differences between RS and multivariate pattern analysis in detail. We found that the RS effect in M1 and parietal areas reflect fundamentally different processes. M1's activity represents the starting finger of the sequence, an effect that vanishes with repetition. In contrast, activity patterns in parietal areas exhibit sequence dependency, which persists with repetition. These findings demonstrate that combining RS and pattern analysis can provide novel functional insights, here specifically into the relative contribution of cortical motor areas to sequence production.

Introduction

The ability to learn and produce complex sequences of movements is essential for many everyday activities, from tying laces to riding a bicycle and playing instruments. Searching for where these acquired skills are represented in the brain has been one of the central questions in motor neuroscience (2). One central issue in this debate is whether skilful sequence execution relies on representations in premotor and supplementary motor areas, or whether the sequences are represented in the primary motor cortex (M1) itself (3–11; for reviews see 12,13). We recently conducted a systematic longitudinal 5-week training study (1) employing functional magnetic resonance imaging (fMRI) to assess brain changes with motor learning. We observed no overall activity changes with learning in M1, and no changes in the sequence-specific activity patterns in this region. In contrast, clear learning-related changes in both overall activity and fine-grained activity patterns were observed in premotor and parietal areas, suggesting learning-related changes occur outside of M1. Other recent studies suggest that activity patterns in M1 reflect individual movement elements, but do not vary with the sequential context (9,11,14). This suggests that M1 does not represent learnt motor sequences, but must rely on inputs from other areas to initiate the next movement element.

Using the technique of repetition suppression, however, Wymbs and Grafton (15) provided evidence for skilled sequence representation in M1. Repetition suppression (RS) refers to the observation that a stimulus repetition evokes reduced neuronal activity compared to its initial presentation (16). This technique has been commonly used as a tool for investigating brain representation (17–24) following the logic that if, on average, regional activation reduces upon repetition, then the underlying neuronal population must represent at least one of the repeated aspects of the stimulus. Based on this, stronger RS effects are commonly interpreted as more robust representations (18). Wymbs and Grafton (15) found learning-related changes in RS across several regions,

including M1, suggesting that M1 might represent trained sequences. This stands in stark contrast to the above-mentioned studies that used pattern dissimilarity analyses and found no evidence of sequential representation in M1.

We reasoned that this discrepancy may reflect the fact that RS can be caused by a range of changes in the neuronal population code, and hence may reflect different changes in brain computation (18,25). To understand RS effects in more detail, we need to know what aspects of the underlying representations reduce from the first to the second repetition of a stimulus or movement. To address this question, we here performed a set of novel multivariate and RS analyses on the data from the longitudinal fMRI learning study (1). In the study, participants performed all sequences, trained and untrained, twice in a row, allowing us to conduct both pattern and RS analysis on the same data. We first replicate the discrepancy in the literature, with RS indicating that M1 develops a sequence representation with training, and multivariate analysis indicating the opposite. Using pattern analysis, we then decompose the activation patterns in the first and second repetition to determine which representational aspects underlie the RS effects in the different regions.

Results

Learning paradigm

We trained 26 participants to execute six 9-digit finger sequences with their right hand (**Fig 1a**), over a period of 5 weeks (**Fig 1b**). Participants also underwent scanning (using 7T MRI) while executing the 6 trained and 6 untrained sequences. In this paper we focus on results from week 1, i.e., before training, and week 5, i.e., after training (**Fig. 1c**; see **Methods** for details). The trials in the scanning sessions were structured so that the same sequence was always executed twice in a row (**Fig 1d**). This design choice

86 allowed us to contrast brain responses during the first execution to that of the second

87 execution to examine repetition effects.

Figure 1. Experimental paradigm. a) Experimental setup – finger sequences composed of 9 digits were executed on a keyboard device. Participants received visual feedback on correctness of their presses – digits turned green for correct presses, red for incorrect presses. **b)** Group-averaged performance on trained sequences over the 5-week behavioural training protocol. Red shade indicates the standard error of the group mean. **c)** Group-averaged performance during the pre-training and post-training scanning sessions. Trained sequences are in red, untrained in blue. Dark colour indicates first execution, light second execution. White bars indicate the group mean performance. **d)** Experimental paradigm inside the scanner. Each sequence was presented twice in a row. Trials started with a 1s preparation time in which the 9-digit sequence was presented, followed by a 3.5s period of sequence execution and 0.5s of inter-trial interval (ITI). Periods of rest were intermixed between trials to allow for an estimate of resting baseline activation. Yellow and purple indicate two different motor sequences. Darker shades indicate first execution, lighter shades second execution.

Changes in repetition suppression and pattern dissimilarities with learning

To examine learning-related changes in repetition suppression and pattern analysis, we calculated both metrics on fMRI activation patterns in the pre-learning session (i.e. week 1), and in the post-learning session (week 5). Relative to rest, sequence execution activated the primary motor cortex (M1), the primary somatosensory cortex (S1), premotor dorsal and ventral regions, supplementary motor cortex and the superior parietal lobules (SPL; **Fig 2a-b**). Activity was higher for the first than for the second execution. The amount of repetition suppression was calculated as the difference in the elicited activation between the two executions of the same sequence (exe 2 – exe 1). Negative values indicate a relative reduction in activation with repetition, i.e., repetition suppression (RS). Already in week 1, prior to learning, RS was observed in nearly all regions displaying task-evoked activation (**Fig 2b-c**). In regions that showed de-activation during task performance (blue shades in **Fig 2a**), we observed positive difference values between the executions (areas in red shades in **Fig 2c**). This indicates that, both the amount of activation, as well as the amount of deactivation reduced with repetition.

Next, we examined whether the regions that exhibited RS also displayed distinct activity patterns for different sequences (i.e., sequence-specific representations). Pattern dissimilarities were calculated as the average crossvalidated Mahalanobis dissimilarity (i.e., Crossnobis dissimilarity) between activation patterns of all possible sequence pairs (see **Methods** for details). Overall, regions with dissimilar activity patterns for the different sequences corresponded to regions which also exhibited RS effects (**Fig 2c-e**). Additionally, both metrics (RS and pattern dissimilarities) increased with from week 1 to week 5, with the effect particularly pronounced in the parietal cortex (**Fig 2c, e**). Thus, based on visual inspection, RS and pattern dissimilarity metrics seem to provide consistent evidence for the development of sequence-specific representations with learning in an overlapping set of regions.

However, a more detailed inspection reveals some important differences between

these two metrics. To show this, we focussed on a set of predefined regions of interest:

the anterior SPL (SPLa), the dorsal premotor cortex (PMd), and M1. In SPLa and PMd,

pattern dissimilarities increased specifically for the trained sequences (**Fig 2g**), as

quantified by a significant interaction in a week x sequence type ANOVA (SPLa:

$F_{(1,25)}$=4.80; p=.038, PMd: $F_{(1,25)}$=5.29, p=.030). The RS effects in the same regions

paralleled those of pattern dissimilarities, with bigger increase in RS for trained than

untrained sequences (**Fig 2f**; SPLa: $F_{(1,25)}$=17.44; p=3.1e^{-4}, PMd: $F_{(1,25)}$=7.27, p=1.1e^{-6}).

Furthermore, the observed change in RS across weeks replicated a previously reported

finding (15; see Figure 2–figure supplement 1). Thus, in PMd and SPLa, trained

sequences exhibited more dissimilar activity patterns and a stronger RS, together

suggesting these regions develop a dedicated representation of the trained sequences

with learning.

In contrast, the two metrics diverged in M1. Here, pattern dissimilarities did not

change with learning, i.e., the week by sequence type interaction was not significant (**Fig**

2g; $F_{(1,25)}$=2.13, p=.16). In contrast, RS increased more for trained than untrained

sequences with learning (**Fig 2f**; $F_{(1,25)}$=25.09; p=3.6e^{-4}). These results, thus, replicate

the previously reported discrepancy in findings showing increased RS in M1 with

learning (15), yet no change in pattern dissimilarities (1), within the same dataset.

Figure 2. Changes in repetition suppression and dissimilarities with learning. a) Group-averaged evoked activation, measured as percent signal change over resting baseline in week 1, averaged across all sequences and projected to an inflated representation of the left hemisphere. **b)** Group-averaged activation for each execution (Exe1, Exe2), in the baseline session (Week 1) and after training (Week 5) represented on a flattened representation of the left hemisphere. CS stands for the central sulcus. **c)** The difference in evoked activation between the two executions. Blue represents relative suppression of activation on the second, relative to the first, execution. Regions of interest: primary motor cortex (M1), dorsal premotor cortex (PMd), anterior superior parietal lobule (SPLa). **d)** Average dissimilarity between evoked patterns for all pairs of sequences, in week 1, averaged across the group. Pattern dissimilarity was computed using a searchlight approach, by calculating the average Crossnobis dissimilarity of activation patterns between all sequence pairs in each searchlight. The mean dissimilarity of a searchlight

was assigned to its centre (see **Methods**). **e)** Average dissimilarity between activation patterns of different sequence pairs in weeks 1 and 4. **f)** Repetition suppression in the predefined regions of interest. Repetition suppression is calculated as the difference in group-averaged activation between the two executions for trained (red) and untrained (blue) sequences. Error bars reflect the standard error of the group. More negative values indicate more suppression during second execution, relative to the first. * signals $p<.05$. **g)** Dissimilarities between trained (red) and untrained (blue) sequence patterns, across weeks 1 and 5. Error bars reflect the standard error of the group. * signals $p<.05$.

Pattern dissimilarities reduce with repetition

Pattern analysis and RS therefore provide conflicting evidence on the nature of learning-related change in M1. While the increase in pattern dissimilarities (**Fig 2f**), as well as direct evidence for cross-session pattern changes (1), clearly argue that sequence-specific learning occurs in premotor and parietal areas and not in M1, RS provides evidence for the development of sequence-specific representations in both parietal / premotor regions and M1.

How can this discrepancy be explained? To resolve this question, we need to understand how the representations in each area during skilled performance change from the first to the second execution. We first inspected pattern dissimilarities for each of the two executions separately (execution 1, execution 2) in the trained state (Week 5). We observed that, on average, pattern dissimilarities in week 5 decreased with repetition in most cortical regions (**Fig 3a**). This decrease was particularly pronounced in M1 (**Fig 3b**).

The relationship between the reduction in overall activity with repetition and the reduction of pattern dissimilarity provides a way of distinguishing three different accounts of RS (**Fig 3c**; 18). According to one hypothesis (the scaling account), all of the involved voxels scale their activation with repetition, proportional to the initial activity in the first instance. This account predicts that dissimilarities reduce proportionally, i.e.

scale, to the decrease in activation. Alternatively (following the sharpening account),

representation could sharpen with repetition – this would occur if the neuronal population

most sensitive to different motor sequences would still be active on repetition, but the

less selective neurons (i.e., background noise) would decrease their activation. As a

third option (the fatigue account), the most active and selective neurons would decrease

their firing the most on repetition, resulting in a representational fatigue, which predicts

that dissimilarities reduce more than the reduction in overall activity.

Given this set of competing accounts, we first investigated how overall

dissimilarities within a region change across the two executions relative to changes in

activation within the same region. We quantified the amount of activation as the length

of the average activity vector relative to rest, separately for each execution (length of

lines in **Fig 3c**, see **Methods** for details). We then calculated the ratio of the activity

during the second execution over the activity during the first. This ratio was applied to

the observed dissimilarities on the first execution (dashed line in **Fig 3d**), yielding a

prediction of what dissimilarities would be expected for the second execution, if

representation scaled with activation. Relative to this expected dissimilarity, the

observed dissimilarities of M1 patterns were significantly lower during the second

execution (**Fig 3d**; $t_{(25)}$=3.87, p=6.9e^{-4}). This indicates that M1's representational change

is more in line with the representational fatigue account, which proposes that the

neuronal populations most tuned to performed sequences on the first execution reduce

their firing the most with repetition.

In contrast to M1, PMd and SPLa's dissimilarities on the second execution were

closer to the expected dissimilarities under scaling (**Fig 3e**; also see figure 3-figure

supplement 1 for a depiction of distance deviation from scaling across the cortical

sheet). To quantify whether the representational nature of repetition differed qualitatively

across regions, we performed a region x sequence type ANOVA. The main effect of

region was significant ($F_{(2,50)}$=7.42, p=1.5e^{-3}). Post-hoc t-tests revealed that this was driven by a significantly larger deviation from scaling in M1 as compared to SPLa ($t_{(25)}$=3.55, p=1.56e^{-3}). M1 and PMd did not differ from one another ($t_{(25)}$=1.25, p=.22). There was a significant difference between PMd and SPLa ($t_{(25)}$=2.65, p=.013), indicating a more 'scaling-like' representation in SPLa. There was no main effect of sequence type ($F_{(1,25)}$=1.35, p=.26), nor a region x sequence type interaction ($F_{(2,50)}$=2.10, p=.14). Altogether this indicates that representational change with repetition differed across regions: proportional scaling of representation in parietal regions, and much more pronounced decrease of dissimilarities in M1, similar to the fatigue account.

Figure 3. Representational change with repetition related to different hypothesized accounts of repetition suppression. a) Dissimilarities between pairs of sequences in week 5 (post-training), split by first and second executions. **b)** Difference in pattern dissimilarities between executions 1 and 2. Blue hues reflect relatively lower dissimilarities on the second execution. **c)** Different possible accounts of representational change with repetition. Yellow and purple indicate two different conditions, i.e. motor sequences in our paradigm. Darker shades

indicate first execution, lighter shades second execution. *Scaling*: dissimilarities scale

proportionally to activation reduction with repetition. *Sharpening*: dissimilarities on the second

execution are relatively higher than during the first execution, after accounting for activation

reduction. *Fatigue*: Dissimilarities reduce on repetition more than expected from the reduction in

activation. **d)** Average Crossnobis dissimilarity between pairs of evoked sequence-specific

patterns in week 5 in the primary motor cortex (M1), split by execution. Dashed line depicts

expected dissimilarities if distances scaled proportionally with reduction of activation on

repetition. The difference between dissimilarities on execution 2 and dashed line is plotted in e).

* denotes $p<.05$ for comparison between the observed dissimilarities on execution 2 and the

expected ones under scaling. **e)** Difference between the expected dissimilarity under scaling of

activation and the observed dissimilarities on the second for trained (red) and untrained (blue)

activity patterns. Regions: M1, dorsal premotor cortex (PMd), anterior superior parietal lobule

(SPLa). Zero indicates proportional scaling with activation. * denotes $p<.05$.

Decomposing representations across executions 1 and 2

Analysis of average dissimilarities across executions revealed bigger compression of

representation in M1 than in parietal regions. This analysis, however, does not reveal

which aspects of the representations are responsible for this regional difference. To

investigate exactly how the representation changed, we decomposed the

representations during each execution into several underlying representational

components. Differences in activity patterns can reflect differences in various

characteristics, or features, of the performed sequences (**Fig 4a**). Specifically, based on

previous results (11,14), we hypothesized that the covariance (or similarity) between

activity patterns can be explained with the following 5 components (**Fig 4b**, see **Methods**

for details): *1) first finger:* the pattern is determined by the starting finger, *2) all fingers*:

the pattern reflects a linear mixture of finger-specific patterns, *3) sequence type*: trained

and untrained sequences have different average patterns, *4) trained sequence identity*:

the trained sequences differ amongst each other, *5) untrained sequence identity*: the

untrained sequences (7-12) differ amongst each other. Using pattern component

modelling framework (26), we constructed a model family out of these components,

which consisted of all possible combinations of those 5 components, totalling 2^5 models

(see Figure 4-figure supplement 1a). These models were then fit to the observed regional

covariance structure (second moment matrices; **Fig 4c**), separately for executions 1 and

2. The crossvalidated log Bayes factor relative to null model (i.e. no component present),

and relative to the noise ceiling, is shown in Figure 4-figure supplement 1 for each of the

combination models. To integrate the results across models, we then used Bayesian

model averaging and estimated how much pattern variance was accounted for by each

of these components (see **Methods** for details).

In M1, the regional representation on the first execution was accounted for by the

individual movement elements (fingers), with especially high weight on the first finger

(**Fig 4d**). This replicates the recent finding that M1's representation during sequence

production tasks can be fully explained by the starting finger, but does not reflect

differences later in the sequence (11,14). In the current study, we did not match the

sequences in terms of the number of times each of the five fingers was pressed. Thus,

subsequent finger presses, encoded in the 'all finger' component, also accounted for

substantial variance, independent of the exact ordering of these movements.

To statistically quantify these effects, we calculated component Bayes factors for

individual components. The Bayes factors were significant for both first and all finger

factors (first finger: BF=13.3, $t_{(25)}$=3.1, p=4.8e^{-3}; all fingers: BF=19.6, $t_{(25)}$=4.1, p=3.4e^{-4}).

In contrast, the component Bayes factors were not significant for any sequence-related

feature – neither sequence type (BF=3.2, t=1.9, p=.07), nor sequence identity: of trained

sequences (BF=1.6, $t_{(25)}$=1.5, p=.16) or untrained sequences (BF=0, $t_{(25)}$=-0.2, p=.85).

Thus, the pattern analysis clearly shows that activity patterns during the first execution

in M1 can be explained by a superposition of individual movements, without clear

evidence of a sequence representation.

In SPLa and PMd, the variance explained during the first execution was well

accounted for by sequence type (SPLa: BF=48.0, $t_{(25)}$=5.85, p=4.22e^{-6}, PMd: BF=55.2,

$t_{(25)}$=4.97, p=4.0e^{-4}), and trained sequence identity (SPLa: BF=7.2, $t_{(25)}$=3.03, p=5.6e^{-3};

PMd: BF=6.0, $t_{(25)}$=2.5, p=.019). There was no significant evidence for representation of

untrained sequence identity in either of the regions (SPLa: BF=0.8, PMd: BF=0;

$t_{(25)}$<=1.1, p>=.28). In comparison to M1, the variance related on individual movements

– either the first finger or all fingers were weaker across PMd and M1. In PMd the first

finger still accounted for some variance (BF=4.1), but this was further reduced in SPLa

(BF=0.5).

With repetition, the pattern component related to the first finger drastically

reduced, in M1 by 93% (**Fig 4d**). The reduction in variance explained by the first finger

component was larger than for the all finger component, which reduced by 75% (paired

t-test: $t_{(25)}$=9.03, p=2.4e^{-9}). This indicates that the drastic reduction of average

dissimilarities in M1 with repetition is mostly due a pronounced first-finger effect during

the first execution that almost vanishes on the second execution. Large reductions in

first finger effect were also observed in PMd (by 81%) and SPLa (by 83%). In contrast,

the representation of sequence type and trained sequence identity in these areas clearly

reduced less (PMd: sequence type: 44%, trained sequence: 64%; SPLa: sequence type:

49%, trained sequence: 55%). The statistical significance of these effects was confirmed

using a paired t-test, comparing the % reduction of the first finger and trained sequence

factors within each region (PMd: $t_{(25)}$=7.96, p=2.6^{-8}; SPLa: $t_{(25)}$=12.8, p=1.7e^{-12}).

In summary, SPLa's regional activation patterns were better accounted for by

components related to the sequence identity than to the first finger, which also reduced

much less with repetition. This likely explains why the average dissimilarities did not

compress with repetition in SPLa regions as much as in M1. PMd's representation was

in-between those of M1 and SPLa – more variance was accounted for by the first finger

than in SPLa, but less than in M1. With repetition, the proportion of different components

to overall regional representation remained relatively stable in PMd and SPLa (**Fig 4e**),

but changed substantially in M1 in that the dominant first-finger representation on the

first execution nearly disappeared on the second execution.

Figure 4. Component decomposition of regional representation across executions 1 and

2. a) Executed 9-digit sequences. Trained sequences of group 1 (presented here) were

untrained for group 2, and vice versa. **b)** Candidate component models used to assess regional

representations across first and second executions. Each row and column indicate a specific

sequence, and values in the matrices reflect the correspondence across sequences on that

component, with yellow indicating higher correspondence. **c)** Regional representations during

the first execution of sequences, as assessed by the crossvalidated second moment matrix,

averaged across subjects of group 1. Similar as for models, each row and column reflect an activation pattern for an individual sequence. Regions: primary motor cortex (M1) and anterior superior parietal lobule (SPLa). **d)** Variance explained by candidate model components on executions 1 (black) and 2 (grey) in M1, PMd (dorsal premotor cortex) and SPLa. See figure supplement for performance of all models in model family. **e)** Relative contribution of variance explained in d) across the different components. The total variance explained across the different components (i.e. sum of the bars in d) was normalized across the two executions to display the relative shift of importance of different representational components.

Discussion

In the present study, we examined how brain representations of motor sequences evolve with training using a combination of pattern analysis and repetition suppression. Both pattern analysis and RS are commonly used to examine brain representations (18,27–30). The two methods often provide convergent results about the representational content of brain regions (18,24) – but sometimes there are discrepancies between the two. Specifically, in the context of motor sequence learning, previous studies using pattern analysis have reported no evidence of M1 representing trained motor sequences with learning (1,10,11). In contrast, one study (15) has reported learning-related increases in RS even for M1, which suggests a representation of the trained sequences in this region.

We first replicated the discrepancy between these two metrics in M1 in the same experiment. Closer examination of this discrepancy revealed that, with repetition, pattern dissimilarities decreased, particularly in M1. When decomposing regional representations into different underlying components, we observed that M1 mainly represents the first finger in a sequence. This component diminishes dramatically on a repetition. In contrast, the representation of sequence type and identity, which accounted for most of the variance in premotor and parietal areas, remained more stable across the two executions. This suggests that RS acts differently on different

components of neuronal representations; thus it's interpretation also varies across regions which differ in their regional code.

Our findings suggest a new model of the interaction between parietal/premotor areas and M1 during the execution of skilled motor sequences. During the first execution, premotor and parietal regions contain information about the specific sequence that needs to be executed (**Fig 5**). Additionally, premotor regions also reflect the starting finger of the sequence. These regions send signals to M1, pre-activating the neural circuits for the movement of the first finger. This replicates a previous finding that the difference between M1's activation patterns is explained by the starting finger, rather than reflecting true sequence representation (11). The finding is also consistent with results from neurophysiology (31) and magneto-encephalography (MEG; 32), showing the first action in a sequence is most highly activated in premotor and motor areas during the preparatory period. Since M1 does not contain representation of sequence identity, continuous signals from higher-order regions provide it with the sequential context needed throughout the sequence production (9,33).

Upon repetition of the same sequence, activation reduces across all regions (**Fig 5**). This effect is already present before training, but its strength increases with training. The global reduction in activation likely indicates that the initial volley of inter-regional communication is established the first time, and does not need to be re-initialised (18). The analysis of the regional representations in terms of their components indicates that this reduction is most pronounced in the pre-activation of the first finger. In contrast, the information about sequence identity in premotor and parietal regions reduces only moderately, but otherwise remains similar to the first execution. This suggests that the sequence representation is always necessary for successfully guiding M1 through the correct sequences of actions, whereas the first finger does not need to be pre-activated for a successful second execution.

Figure 5. Conceptual depiction of changes in representation across regions and with

repetition. Different dots represent activation patterns for different finger sequences. Regions:

anterior superior parietal lobule (SPLa), dorsal premotor cortex (PMd), primary motor cortex

(M1). Arrows depict communication between higher-order areas and M1. Activation levels of

three hypothetical voxels are indicated across the 3 axes.

Overall, our results suggest that M1 does not represent individual trained

sequences with learning, despite increased RS. Instead, its representation is

characterized by combinations of finger presses. If this is true, why was RS in M1

stronger for trained than for untrained sequences? Since fMRI activation reflects the

driving input into the area (34), this effect may reflect changes in the communication

between higher-order areas and M1, which may become more efficient with repetition of

trained sequences. Some support for this idea comes from a recent study demonstrating

layer-specific effects in M1 (35). By measuring changes in cerebral blood volume across

layers, the authors demonstrated that superficial M1 layers (which reflect inputs into M1)

show RS, whereas deep layers' activation (which is more indicative of M1's outputs) is

enhanced during repetition. Since the blood-oxygenated level depend signal (BOLD) is

biased towards the superficial vascular signals, our activation results more likely reflect

inputs into M1.

An alternative explanation for M1's increased RS for trained sequences is that M1

does contain sequence information, albeit at a spatial scale which cannot be detected

by pattern analyses. In line with this idea, a prior electrophysiology study reported differential M1's responses to trained relative to random sequences (8). This study, however, only reflected an overall difference between trained and random sequences, but not M1's differential activation for different trained sequences, thus no sequence representation as defined here. More recent electrophysiological studies have shown that M1 only represents the ongoing movement, and does not represent the sequential context (9,33). Altogether, this makes M1 representing individual trained sequences a less likely explanation of our results.

There are a number of limitations of the current study which merit future experimentation. First, our study did not differentiate between repetition suppression and expectation suppression (36,37). Expectation of repetition has been shown to modulate RS. We suspect that our observed effects would likely be weaker had the sequence repetition been less predictable, although we would not expect qualitatively different patterns of results. Second, the RS accounts tested here are by no means an exhaustive list. Other studies have examined additional models of RS (25,38). Some of those can only be tested with measurements providing better temporal resolution than that afforded by fMRI. For instance, a facilitation model predicts a faster start and finish of neuronal responses with repetition (22,39), which we are unable to assess using the sluggish temporal resolution of fMRI. Similarly, differences in neural synchronization across regions have been discussed as a mechanism of RS (40). MEG (41,42) or electrocorticography approaches (43) are more appropriate to test those accounts, and may provide important insights into the dynamics of parietal-premotor-M1 interactions.

Nevertheless, with the RS accounts employed here, we demonstrated that RS can reflect very different representational changes from the first to the second execution depending on the regions. During the repetition of skilled motor sequences, associative cortical regions represent the sequence itself in a stable fashion, while the activation in M1 changes qualitatively, with a reduced activation of the beginning of the sequence.

435　These results emphasize that employing RS only using the average regional activation

436　sometimes provides incomplete, and possibly misleading, insights into regional

437　representation. Instead, RS should be combined with pattern analyses to determine how

438　the representations change with repetition, to ultimately provide a deeper understanding

439　of brain circuits and their function.

Materials and Methods

Participants

Twenty-seven participants took part in the experiment. Data of one participant were excluded because their fieldmap was distorted in one of the four scans, resulting in 26 participants whose data was analysed (17 females, 9 males). Their mean age was 22.2 years (SD = 3.3 years). Criteria for study inclusion were right-handedness and no prior history of psychiatric or neurological disorders. They provided written informed consent to all procedures and data usage before the study started. The experimental procedures were approved by the Ethics Committee at Western University.

Apparatus

Finger sequences were performed using a right-hand MRI-compatible keyboard device (**Fig 1a**). The keys of the device had a groove for each fingertip, with keys numbered 1-5 for thumb-little finger. The keys were not depressible, so participants performed isometric finger presses. The force of the presses was measured by the force transducers underneath each finger groove (FSG-15N1A, Sensing and Control, Honeywell; dynamic range 0-25 N; update rate 2 ms; sampling 200 Hz). For the key to be recognized as pressed, the applied force had to exceed 1 N.

Learning paradigm

Participants were trained over a five-week time period to perform six 9-digit finger sequences. They were split into two groups, with trained sequences of one group being the untrained sequences of the second group, and vice versa (see **Fig 4b** for all of the chosen sequences). The chosen sequences for both groups were matched as closely as possible on several features: starting finger, number of repetitions per finger, and first-order finger transitions. The decision to split participants into two groups was made

to ensure that none of the observed effects could be due to the specific set of sequences

chosen.

On day 1 of the study, participants were acquainted with the apparatus and the

task performed in the scanner. To ensure no sequence-specific learning would take

place prior to scan 1, we used finger sequences different from the trained and untrained

sets which participants did not encounter at any later stage of the experiment.

During the behavioral training sessions, participants were trained to perform the

six sequences. Each training session was split into several blocks with 24 trials in each

block (i.e. 4 repetitions of each of the 6 sequences). Participants received visual

feedback on the correctness of their presses online with digits turning green for correct

presses and red for incorrect presses (**Fig 1a**). They were instructed to perform the

sequences as fast as possible, while keeping the overall accuracy within each block

<15% error rate. To encourage fast and accurate performance throughout the training,

participants received 0, 1 or 3 points after each trial completion based on correctness

and movement time (MT) – time from the first press until the last release in the sequence.

Specifically, they received 1 point for correct presses performed at their median MT, 3

points for correct presses executed at least 20% faster than their median speed until

that point. If they made a mistake on any of the presses or the MT in the trial was lower

than their median MT, they received 0 points. The median MT threshold was dynamically

adjusted across blocks every time a block was completed with an error rate <15% and

the median MT faster than previous threshold. Overall, their MT decreased across the

five weeks of training (**Fig 1b**).

We included several other design features in the behavioural training, such as

behavioural test sessions comparing participants' performance of trained sequences to

random sequences, and their performance using the left hand. Those sessions were not

assessed for this paper, and have been described elsewhere (1).

Experimental design during scanning

Participants underwent a total of 4 MRI scanning sessions (**Fig 1c**) while executing trained and untrained sequences. The first session served as a baseline prior to the start of the training protocol (in week 1), where the "trained" and "untrained" sequences were both untrained and seen for equivalent amount. The second session was conducted in week 2, and the last two after training protocol was completed – in week 5. In scanning sessions 1-3, participants' performance inside the scanner was paced with a metronome, whereas in session 4, they performed as fast as possible. For the purpose of this paper, data of scanning session 1 (prior to training) and 4 (fully skilled performance) were analysed (**Fig 1c**). Session 4 allows for the closest comparison to the previous RS study (15) which also employed a full-speed performance design. Pattern analyses of sessions 2 and 3 are reported in (1).

Each scanning session consisted of eight functional runs with event-related design randomly intermixing trials containing the 6 trained and the 6 untrained sequences (totalling 72 trials per functional run). Each sequence was executed for two trials in a row (**Fig 1d**). The trial started with a 1 second preparation time with nine digits of the sequence presented on the screen. A 'go' signal was presented afterwards. In scans 1-3, a pink line appeared underneath the sequence and started expanding, indicating the pace at which participants were to press. In scan 4, participants executed the sequence as fast as possible after the go cue. After execution, they received feedback on their overall performance – 3 points for correct and 0 for incorrect performance. Each trial lasted for 5 s total (**Fig 1d**), with a 0.5 s inter-trial interval. Five periods of 10 s rests were added throughout each functional run to provide a better estimate of baseline activation. These rests were added randomly, but never between the first and second execution of the same sequence. In total, each scanning session lasted for approximately 75 minutes.

Image acquisition

Data were acquired on a 7-Tesla Siemens Magnetom MRI scanner with a 32-receive channel head coil (8-channel parallel transmit). At the beginning of the first scan, we acquired anatomical T1-weighted scan for each participant. This was obtained using a magnetization-prepared rapid gradient echo sequence (MPRAGE) with voxel size of 0.75x0.75x0.75 mm isotropic (field of view = 208 x 157 x 110 mm [A-P; R-L; F-H], encoding direction coronal). Data during functional runs were acquired using the following sequence parameters: GRAPPA 3, multi-band acceleration factor 2, repetition time [TR] = 1.0 s, echo time [TE] = 20 ms, flip angle [FA] = 30 deg, slice number: 44, voxel size: 2x2x2 mm isotropic. To estimate magnetic field inhomogeneities, we acquired a gradient echo field map with the following parameters: transversal orientation, field of view: 210 x 210 x 160 mm, 64 slices, 2.5 mm thickness, TR = 475 ms, TE = 4.08 ms, FA = 35 deg.

Preprocessing and first level analysis

Data preprocessing was carried out using SPM12 and custom MATLAB code. Preprocessing of functional data included correcting for geometric distortions using the acquired fieldmap data, and head motion correction (3 translations: x, y, z; 3 rotations: pitch, roll yaw). The data across sessions were all aligned to the first run of the first session, and then co-registered to the anatomical scan.

Preprocessed data were analysed using a general linear model (GLM; 44). We defined a regressor for each of the performed 12 sequences (6 trained, 6 untrained), separately for their first and second execution – resulting in a total of 24 regressors per run. The regressor was a boxcar function defined for each trial, and convolved with a two-gamma canonical hemodynamic response function (time to peak: 5.5 s, time to undershoot: 12.5 s). All instances of sequence execution were included into estimating

regressors, regardless of whether the execution was correct or erroneous. This analysis

choice was also taken by (15), thus allowing a more direct comparison of repetition

suppression results. Even when the error trials were excluded (i.e. removing all error

trials as well as second execution trials when the first execution was erroneous), our

results remained unchanged. Ultimately, the first level analysis resulted in activation

images (beta maps) for each of the 24 conditions per run, for each of the four scanning

sessions.

Surface reconstruction and regions of interest

Individual subject's cortical surfaces were reconstructed using FreeSurfer (45), and

aligned to the FreeSurfer's Left-Right symmetric template (Workbench's 164 nodes

template) via spherical registration. For our regions of interest (ROI), we defined areas

covering the primary motor cortex and secondary associative regions. The primary motor

cortex (M1) was defined using probabilistic cytoarchitectonic map (46) by including

nodes with the highest probability of belonging to Brodmann area (BA) 4 which in

addition corresponded to the hand knob area (47). The dorsal premotor cortex (PMd)

was included as the lateral part of the middle frontal gyrus. The anterior part of the

superior parietal lobule (SPLa) was defined to include anterior, medial and ventral

intraparietal sulcus.

Evoked activation and repetition suppression

We calculated the percent signal change for execution of each sequence relative to the

baseline activation for each voxel. The calculation was split between the first and second

execution. To calculate repetition suppression, the activation during the first execution

was subtracted from the elicited activation during the second execution. Thus, negative

values of this difference contrast represented relative suppression of activation on the

second execution, i.e. repetition suppression. For most subsequent analyses, the

obtained values of activation and repetition suppression were averaged separately for

trained and the untrained sequences. For ROI analysis, the volume maps were averaged

across the predefined regions (M1, PMd, SPLa) in the native volume space of each

subject. Additionally, for visualization the volume maps were projected to the surface for

each subject, and averaged across the group in Workbench space.

Dissimilarities between activity patterns for different sequences

To evaluate which regions displayed sequence-specific representation, we calculated

Crossnobis dissimilarities between the evoked beta patterns of individual sequences. To

do so, we first multivariately prewhitened the beta values – i.e. we standardized them by

voxels' residuals and weighted by the voxel noise covariance matrix. Prewhitening has

previously been found to increase the reliability of dissimilarity estimates (48). Next, we

calculated the crossvalidated squared Mahalanobis dissimilarities (i.e. the Crossnobis

dissimilarities) between evoked regional patterns of different pairs of sequences,

resulting in a total of 66 dissimilarities. This was performed twice: once by combining the

activation patterns across the two executions and second time by separately obtaining

dissimilarities between evoked patterns split per execution. The obtained dissimilarities

were then averaged overall, as well as separately within the pairs of trained sequences,

and the untrained sequences.

The multivariate analysis was conducted separately for each ROI and then

repeated using a surface searchlight approach (49). For each surface node, a circular

region of 120 voxels was defined and Crossnobis dissimilarities were calculated

amongst those voxels, with the resulting dissimilarities values assigned to the centre of

the searchlight. The searchlight was then moved across the cortical sheet in a

continuous manner.

Assessing accounts of repetition suppression

We examined several proposed accounts of RS – scaling, sharpening and fatigue (**Fig 3c**), by relating the observed changes in pattern dissimilarities with repetition to activation changes with repetition. Activation pattern for each sequence can be characterized as a point in a high-dimensional space, with each axis referring to the activation of a voxel. As a measure of the overall activation, we used the length of the activity vector from the origin (rest), and as dissimilarities the lengths of the vectors between different dots (**Fig 3c**). Unbiased estimates of the length of activity vectors relative to rest were derived from the crossvalidated second-moment matrix. The square root of each diagonal element (variance of evoked pattern) indicates the length of the activity vector, relative to rest. The square root of Crossnobis dissimilarity (variance – covariance between patterns) is the length of the vector between the two patterns (50). These measures were averaged across the 6 trained, and 6 untrained conditions, separately for executions 1 and 2.

For each subject and region, we then computed the decrease in activation as activity length for 2^{nd} execution divided by activity length for the 1^{st} execution. If dissimilarities decreased proportional to changes in activation, we would expect the ratio of dissimilarities between the two executions to equal the ratio obtained from activity vector lengths. Likewise, the dissimilarities expected under the scaling account can be computed as: $\text{diss}_{\text{scaling}} = \frac{\text{act}_{\text{exe2}}}{\text{act}_{\text{exe1}}} \text{ x diss}_{\text{exe1}}$. This expected dissimilarity value was compared to the observed dissimilarities on the second execution. Higher value of the observed relative to expected dissimilarity would be in line with the 'sharpening' account, lower value would be in support of the 'fatigue' account, while a value that wouldn't differ from the expected dissimilarities would support the 'scaling' account of RS for that region.

Pattern component analyses: modelling representational components

The analysis on dissimilarities between sequence-specific activity patterns is sensitive

to any possible difference between patterns of different sequences. To decompose what

specific features of stimuli might be driving these differences, we used pattern

component modelling (PCM; 26,51). PCM models the covariance structure (second

moment matrix) of regional activity patterns according to different representational

hypotheses. In our experiment based on presented sequences, we defined five

representational components.

1) First finger

Both trained and untrained sequences started with 3 possible fingers: thumb, middle

finger and pinky finger. The first finger feature characterizes activity pattern as different

for sequences starting with different first finger, and indistinguishable for sequences

starting with the same first finger. The model component was weighted by the natural

statistics of hand movement (52).

2) All fingers

The chosen sequences had slight differences in terms of how many times different

fingers were executed in the sequence. Thus, the 'all fingers' feature characterized

number of repetitions across fingers and how correspondent different sequences were

in finger repetitions. Also this model component was weighted by the natural statistics

of hand movement (52).

3) Sequence type

This component split the performed sequences based on whether they were trained or

not, predicting different regional representations between trained and untrained

sequences, but indistinguishable patterns within each of the two groups.

4) Trained sequence identity

This component assumes that each of the trained sequences is associated with unique

activity patterns.

5) Untrained sequence identity

Similar as the trained sequence identity component, this component assumes unique

activity patterns for different untrained sequences.

The overall predicted second moment matrix (G) was then a convex combination of the

component matrices (G_c):

658
$$G = \sum_c \exp(\Theta_c) \, G_c$$

The construction of the models was split between the two groups of participants, as

different sequences constituted 'trained' or 'untrained' sequences for the two groups.

The subsequent steps of model fitting and evaluation were carried together for all

subjects.

We formulated a family of models containing all possible combinations of the five

chosen components. This resulted in 32 combinations (Figure 4-figure supplement 1),

also containing the 'null' model without any component, which predicted no differences

amongst any of the sequence patterns. We then evaluated all of the 32 models using a

crossvalidated leave-one-subject-out scheme, estimating the likelihood of the observed

regional second moment data under each model. The resultant cross-validated

likelihoods were used as evidence of the model. The difference between the log-

likelihood of each model and the null model results in a log-Bayes factor representing

the relative model evidence over a null model.

In addition to the model family of the chosen components, we also fit a 'noise-

ceiling' model to assess the maximal explainable information present in each region's

representation across participants (26,53). For each of the two groups, we predicted the

second moment matrix of a left-out subject based on n-1 subjects in the same group.

This metric of inter-subject consistency was then combined across the subjects of the

two groups.

To integrate the results across models, we used model averaging to determine the amount of pattern variance accounted for by each component in the context of all the other components. Prior to fitting, we normalized the trace of each model component to be 12 (number of conditions). Thus, the fitted component weight $\exp(\Theta_i)$ can be taken as the amount of variance accounted for by the component in the context of that model. Assuming a uniform prior probability across models, we compute a posterior probability of each model i (from 1 to 32) directly from their Bayes factors:

$$posterior_i = \frac{\exp(logBF_i)}{\sum_{j=1}^{c} \exp(logBF_j)}$$

The model-averaged amount of variance accounted for by each component c is then given by:

$$variance_c = \sum_{i=1}^{32} posterior_i \; x \; \exp(\Theta_c)$$

Important to note is that this variance calculation always results in positive estimates.

Data availability

The dataset used for these analyses is publicly available on OpenNeuro (doi: 10.18112/openneuro.ds002776.v1.0.2).

Acknowledgments

The work was supported by an Ontario Trillium Scholarship to EB, an NSERC Discovery Grant (RGPIN-2016-04890) to JD, and the Canada First Research Excellence Fund (BrainsCAN). We thank Giacomo Ariani for helpful comments on an earlier version of the manuscript.

 Supplementary Figures

Figure 2-figure supplement 1. Learning-related changes in repetition suppression.
Activation elicited on the first (x-axis) and second (y-axis) execution for trained (red) and
untrained (blue) sequences across weeks 1, 2 and 5, in sessions using paced performance.
Regions: anterior superior parietal lobule (SPLa), dorsal premotor cortex (PMd), primary motor
cortex (M1). Ellipses reflect the standard error of the group. Dots below the diagonal line indicate
activation was lower on second execution.

crossnobis dissimilarity from 'scaling'

-0.02   0   0.02

Figure 3-figure supplement 1. Difference in the Crossnobis dissimilarity on second execution from the 'scaling' account prediction. This was calculated for each of the regularly tessellated regions. Blue depicts lower dissimilarities than expected under the scaling account, yellow higher dissimilarities than predicted under scaling of activation.

716
717

Figure 4-figure supplement 1. Pattern component modelling – formulation and model fits.

a) Construction of a model family from the five components. All possible combinations of five components were used to construct models, resulting in 32 (2^5) models. The first model is a 'null' model (no component) and the last is a full model (all components). White squares indicate presence of the component in each model. **b)** Evidence for all models across regional activation, separately examined for the first and second sequence execution (left, right column). Regions: primary motor cortex (M1), dorsal premotor cortex (PMd), anterior superior parietal lobule (SPLa). The 32 evaluated models are presented on the x-axis. Model fits were assessed with a type-II log-likelihood. The difference in log-likelihood between each model and the null model was calculated; this can be interpreted as a log-Bayes factor (Log-BF) and is plotted on the y-axis. The grey shaded area represents the mean noise ceiling ± standard error of the group mean. The bar with a star is the best performing model. All models in red performed within the noise ceiling and did not differ from one another. Bars in dark blue depict models which hit the noise ceiling, but performed significantly worse than the best performing model. Bars in light blue did not reach the noise ceiling performance.

References

1. Berlot E, Popp NJ, Diedrichsen J. A critical re-evaluation of fMRI signatures of motor sequence learning. Elife. 2020;9(e55241):1–24.

2. Lashley K. In search of the engram. 1950.

3. Floyer-Lea A, Matthews PM. Distinguishable brain activation networks for short- and long-term motor skill learning. J Neurophysiol. 2005;94(1):512–8.

4. Karni A, Meyer G, Rey-Hipolito C, Jezzard P, Adams MM, Turner R, et al. The acquisition of skilled motor performance: fast and slow experience-driven changes in primary motor cortex. Proc Natl Acad Sci U S A. 1998;95(3):861–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/9448252%5Cnhttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC33809

5. Karni A, Meyer G, Jezzard P, Adams MM, Turner R, Ungerleider LG. Functional MRI evidence for adult motor cortex plasticity during motor skill learning. Nature. 1995;377(6545):155–8. Available from: http://www.nature.com/doifinder/10.1038/377155a0%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/7675082

6. Lehéricy S, Benali H, Van de Moortele P-F, Pélégrini-Issac M, Waechter T, Ugurbil K, et al. Distinct basal ganglia territories are engaged in early and advanced motor sequence learning. Proc Natl Acad Sci U S A. 2005;102(35):12566–71. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1194910&tool=pmcentrez&rendertype=abstract

7. Penhune VB, Doyon J. Dynamic cortical and subcortical networks in learning and delayed recall of timed motor sequences. J Neurosci. 2002;22(4):1397–406. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11850466

8. Matsuzaka Y, Picard N, Strick PL, Barnes TD, Mao J, Hu D, et al. Skill Representation in the Primary Motor Cortex After Long-Term Practice Skill Representation in the Primary Motor Cortex After Long-Term Practice. J Neurophysiol. 2011;(December 2006):1819–32.

9. Russo AA, Khajeh R, Bittner SR, Perkins SM, Cunningham JP, Abbott LF, et al. Neural trajectories in the supplementary motor area and primary motor cortex exhibit distinct geometries, compatible with different classes of computation. Neuron. 2020;107(1–14).

10. Wiestler T, Diedrichsen J. Skill learning strengthens cortical representations of motor sequences. Elife. 2013;2013(2):1–20.

11. Yokoi A, Arbuckle SA, Diedrichsen J. The role of human primary motor cortex in the production of skilled finger sequences. J Neurosci. 2018;38(6):1430–42.

12. Dayan E, Cohen LG. Neuroplasticity subserving motor skill learning. Neuron. 2011;72(3):443–54. Available from: http://dx.doi.org/10.1016/j.neuron.2011.10.008

13. Berlot E, Popp NJ, Diedrichsen J. In search of the engram, 2017. Curr Opin Behav Sci. 2018;20.

14. Yokoi A, Diedrichsen J. Neural Organization of Hierarchical Motor Sequence Representations in the Human Neocortex. Neuron. 2019;103(6):1178–1190.e7.

15. Wymbs NF, Grafton ST. The human motor system supports sequence-specific representations over multiple training-dependent timescales. Cereb Cortex. 2015;25(11):4213–25.

16. Gross, C.G., Schiller, P.H., Wells, C., Gerstein GL. Single-unit activity in temporal association cortex of the monkey. J Neurophysiol. 1967;30(4):833–43.

17. Weiner KS, Sayres R, Vinberg J, Grill-Spector K. fMRI-adaptation and category selectivity in human ventral temporal cortex: Regional differences across time scales. J Neurophysiol. 2010;103(6):3349–65.

18. Grill-Spector K, Henson R, Martin A. Repetition and the brain: Neural models of stimulus-

specific effects. Trends Cogn Sci. 2006;10(1):14–23.

19.   Grafton ST, Hamilton AFDC. Evidence for a distributed hierarchy of action representation
in the brain. 2007;26:590–616.

20.   Shmuelof L, Zohary E. Dissociation between Ventral and Dorsal fMRI. 2005;47:457–70.

21.   Buckner RL, Goodman J, Burock M, Rotte M, Koutstaal W, Schacter D, et al. Functional-
Anatomic Correlates of Object Priming in Humans Revealed by Rapid Presentation
Event-Related fMRI. 1998;20:285–96.

22.   Henson RN, Ganel T, Otten LJ. Electrophysiological and Haemodynamic Correlates of
Face Perception , Recognition and Priming. 2003;793–805.

23.   Noppeney U, Price CJ. An fMRI Study of Syntactic Adaptation. J Cogn Neurosci.
2004;16(4):702–13.

24.   Barron HC, Vogels TP, Emir UE, Makin TR, O'Shea J, Clare S, et al. Unmasking Latent
Inhibitory Connections in Human Cortex to Reveal Dormant Cortical Memories. Neuron.
2016;90(1):191–203. Available from: http://dx.doi.org/10.1016/j.neuron.2016.02.031

25.   Alink A, Abdulrahman H, Henson RN. From neurons to voxels - repetition suppression is
best modelled by local neural scaling. Nat Commun. 2018;9(3854):1–10. Available from:
https://www.biorxiv.org/content/early/2017/07/31/170498

26.   Diedrichsen J, Yokoi A, Arbuckle SA. Pattern component modeling: A flexible approach
for understanding the representational structure of brain activity patterns. Neuroimage.
2017;

27.   Barron HC, Garvert MM, Behrens TEJ. Repetition suppression: a means to index neural
representations using BOLD? Philos Trans R Soc B Biol Sci. 2016;371(1705):20150355.
Available from: http://rstb.royalsocietypublishing.org/lookup/doi/10.1098/rstb.2015.0355

28.   Segaert K, Weber K, de Lange FP, Petersson KM, Hagoort P. The suppression of
repetition enhancement: A review of fMRI studies. Neuropsychologia. 2013;51(1):59–66.
Available from: http://dx.doi.org/10.1016/j.neuropsychologia.2012.11.006

29.   Walther A, Ejaz N, Kriegeskorte N, Diedrichsen J. Representational fMRI analysis: an
introductory tutorial. 2014;

30.   Kriegeskorte N, Kievit RA. Representational geometry: Integrating cognition,
computation, and the brain. Trends Cogn Sci. 2013;17(8):401–12.

31.   Averbeck BB, Chafee M V, Crowe DA, Georgopoulos AP. Parallel processing of serial
movements in prefrontal cortex. PNAS. 2002;99(20):13172–7.

32.   Kornysheva K, Bush D, Meyer SS, Sadnicka A, Barnes G, Burgess N. Neural
Competitive Queuing of Ordinal Structure Underlies Skilled Sequential Action. Neuron.
2019;101(6):1166–1180.e3.

33.   Zimnik AJ, Churchland MM. Generation of Rapid Sequences by Motor Cortex. bioRxiv.
2020;

34.   Picard N, Matsuzaka Y, Strick PL. Extended practice of a motor skill is associated with
reduced metabolic activity in M1. Nat Neurosci. 2013;16(9):1340–7. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3757119&tool=pmcentrez&re
ndertype=abstract

35.   Persichetti AS, Avery JA, Huber L, Merriam EP, Martin A. Layer-Specific Contributions to
Imagined and Executed Hand Movements in Human Primary Motor Cortex. SSRN
Electron J. 2019;1–5.

36.   Summerfield C, Trittschuh EH, Monti JM, Mesulam MM, Egner T. Neural repetition
suppression reflects fulfilled perceptual expectations. Nat Neurosci. 2008;11(9):1004–6.

37.   Kok P, Jehee JFM, de Lange FP. Less Is More: Expectation Sharpens Representations in
the Primary Visual Cortex. Neuron. 2012;75(2):265–70.

38.   Garrido MI, Kilner JM, Kiebel SJ, Stephan KE, Friston KJ. Repetition suppression and
plasticity in the human brain. Neuroimage. 2010;48(1):269–79.

39.   James TW, Gauthier I. Repetition-Induced Changes in BOLD Response Reflect

835   Accumulation of Neural Activity. 2006;46:37–46.

836 40. Gotts SJ, Chow CC, Martin A. Repetition priming and repetition suppression : A case for
837   enhanced efficiency through neural synchronization. Cogn Neurosci. 2012;3(May).

838 41. Ghuman AS, Bar M, Dobbins IG, Schnyer DM. The effects of priming on frontal-temporal
839   communication. PNAS. 2008;105(24):8405–9.

840 42. Gilbert JR, Gotts SJ, Carver FW, Martin A. Object repetition leads to local increases in
841   the temporal coordination of neural responses. Front Hum Neurosci. 2010;4(April):1–12.

842 43. Korzeniewska A, Wang Y, Benz HL, Fifer MS, Collard M, Milsap G, et al. Changes in
843   human brain dynamics during behavioral priming and repetition suppression. Prog
844   Neurobiol. 2020;189(May 2019).

845 44. Friston KJ, Worsley KJ, Poline J-PB, Frith CD, Frackowiak RSJ, Holmes a. P, et al.
846   Statistical parametric maps in functional imaging: A general linear approach. Hum Brain
847   Mapp. 1994;2(4):189–210. Available from: http://doi.wiley.com/10.1002/hbm.460020402

848 45. Dale AM, Fischl B, Sereno MI. Cortical Surface-Based Analysis. Neuroimage.
849   1999;9(2):179–94. Available from:
850   http://www.ncbi.nlm.nih.gov/pubmed/9931268%0Ahttp://linkinghub.elsevier.com/retrieve
851   /pii/S1053811998903950

852 46. Fischl B, Rajendran N, Busa E, Augustinack J, Hinds O, Yeo BTT, et al. Cortical folding
853   patterns and predicting cytoarchitecture. Cereb Cortex. 2008;18(8):1973–80.

854 47. Yousry TA, Schmid UD, Alkadhi H, Schmidt D, Peraud A, Buettner A, et al. Localization
855   of the motor hand area to a knob on the precentral gyrus. A new landmark. Brain.
856   1997;120(1):141–57.

857 48. Walther A, Nili H, Ejaz N, Alink A, Kriegeskorte N, Diedrichsen J. Reliability of
858   dissimilarity measures for multi-voxel pattern analysis. Neuroimage. 2016;137:188–200.
859   Available from: http://dx.doi.org/10.1016/j.neuroimage.2015.12.012

860 49. Oosterhof NN, Wiestler T, Downing PE, Diedrichsen J. A comparison of volume-based
861   and surface-based multi-voxel pattern analysis. Neuroimage. 2011;56(2):593–600.
862   Available from: http://dx.doi.org/10.1016/j.neuroimage.2010.04.270

863 50. Diedrichsen J, Provost S, Zareamoghaddam H. On the distribution of cross-validated
864   Mahalanobis distances. arXiv Prepr arXiv. 2016;1607.01371. Available from:
865   http://arxiv.org/abs/1607.01371

866 51. Diedrichsen J, Ridgway GR, Friston KJ, Wiestler T. Comparing the similarity and spatial
867   structure of neural representations: A pattern-component model. Neuroimage.
868   2011;55(4):1665–78. Available from: http://dx.doi.org/10.1016/j.neuroimage.2011.01.044

869 52. Ejaz N, Hamada M, Diedrichsen J ouml rn. Hand use predicts the structure of
870   representations in sensorimotor cortex. Nat Neurosci. 2015;18(7):1034–40. Available
871   from: http://dx.doi.org/10.1038/nn.4038%5Cnpapers3://publication/doi/10.1038/nn.4038

872 53. Nili H, Wingfield C, Walther A, Su L, Marslen-Wilson W, Kriegeskorte N. A Toolbox for
873   Representational Similarity Analysis. PLoS Comput Biol. 2014;10(4):e1003553.