**Analysis of paralogs in target enrichment data pinpoints multiple ancient polyploidy events**

**in *Alchemilla* s.l. (Rosaceae).**

Diego F. Morales-Briones[1,2,*], Berit Gehrke[3], Chien-Hsun Huang[4], Aaron Liston[5], Hong Ma[6],

Hannah E. Marx[7], David C. Tank[2], Ya Yang[1]

[1]Department of Plant and Microbial Biology, University of Minnesota-Twin Cities, 1445 Gortner

Avenue, St. Paul, MN 55108, USA

[2]Department of Biological Sciences and Institute for Bioinformatics and Evolutionary Studies,

University of Idaho, 875 Perimeter Drive MS 3051, Moscow, ID 83844, USA

[3]University Gardens, University Museum, University of Bergen, Mildeveien 240, 5259

Hjellestad, Norway

[4]State Key Laboratory of Genetic Engineering and Collaborative Innovation Center of Genetics

and Development, Ministry of Education Key Laboratory of Biodiversity and Ecological

Engineering, Institute of Plant Biology, Center of Evolutionary Biology, School of Life Sciences,

Fudan University, Shanghai 200433, China

[5]Department of Botany and Plant Pathology, Oregon State University, 2082 Cordley Hall,

Corvallis, OR 97331, USA

[6]Department of Biology, the Huck Institute of the Life Sciences, the Pennsylvania State

University, 510D Mueller Laboratory, University Park, PA 16802 USA

[7]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI

48109-1048, USA

* Correspondence to be sent to: Diego F. Morales-Briones. Department of Plant and Microbial

Biology, University of Minnesota, 1445 Gortner Avenue, St. Paul, MN 55108, USA, Email:

dfmoralesb@gmail.com

***Abstract.*** —Target enrichment is becoming increasingly popular for phylogenomic studies.

Although baits for enrichment are typically designed to target single-copy genes, paralogs are

often recovered with increased sequencing depth, sometimes from a significant proportion of

loci. Common approaches for processing paralogs in target enrichment datasets include removal,

random selection, and manual pruning of loci that show evidence of paralogy. These approaches

can introduce errors in orthology inference, and sometimes significantly reduce the number of

loci, especially in groups experiencing whole-genome duplication (WGD) events. Here we used

an automated approach for paralog processing in a target enrichment dataset of 68 species of

*Alchemilla* s.l. (Rosaceae), a widely distributed clade of plants primarily from temperate climate

regions. Previous molecular phylogenetic studies and chromosome numbers both suggested the

polyploid origin of the group. However, putative parental lineages remain unknown. By taking

paralogs into consideration, we identified four nodes in the backbone of *Alchemilla* s.l. with an

elevated proportion of gene duplication. Furthermore, using a gene-tree reconciliation approach

we established the autopolyploidy origin of the entire *Alchemilla* s.l. and the nested

allopolyploidy origin of four clades within the group. Here we showed the utility of automated

orthology methods, commonly used in genomic or transcriptomic datasets, to study complex

scenarios of polyploidy and reticulate evolution from target enrichment datasets.

**Keywords:** *Alchemilla*; allopolyploidy; autopolyploidy; gene tree discordance; orthology

inference; paralogs; Rosaceae; target enrichment; whole genome duplication.

Polyploidy, or whole genome duplication (WGD), is prevalent throughout the evolutionary

history of plants (Cui et al. 2006; Jiao et al. 2011; Jiao et al. 2012; Leebens-Mack et al. 2019). As

a result, plant genomes often contain large numbers of paralogous genes from recurrent gene and

genome duplication events. With very few nuclear genes being truly single- or low-copy, careful

evaluation of orthology is critical for phylogenetic analyses (Fitch 1970). Orthology inference

has received much attention in the phylogenomic era with multiple pipelines available for this

task (e.g., Li et al. 2003; Dunn et al. 2013; Kocot et al. 2013; Yang and Smith 2014; Emms and

Kelly 2019, also see Glover et. al 2019 and Fernández et al. 2020 for recent reviews). All of

these approaches were developed for genome and transcriptome datasets. So far, few studies

have employed automated, phylogeny-aware orthology inference in target enrichment datasets.

The most common approach for resolving paralogy in target enrichment datasets is removing

loci that show evidence of potential paralogy (e.g., Nicholls et al. 2015; Jones et al. 2019;

Andermann et al. 2020; but see Moore et al. 2018). Removal of paralogous genes might be

appropriate in target enrichment datasets in which only a small number of loci show evidence of

paralogy (e.g., Larridon et al. 2020), but in some datasets this could result in a significant

reduction of the number of loci (e.g., Montes et al. 2019). Given the prevalence of WGD in

plants and the continuous increase of sequencing depth in typical target enrichment studies, the

latter scenario is becoming increasingly frequent, calling for automated, tree-based approaches

for processing paralogs in target enrichment data.

The ability to explicitly process paralogs opens the door for using target enrichment data

for inferring gene duplication events and pinpointing the phylogenetic locations of putative

WGDs. In the past, the phylogenetic placement of WGD events have most often been carried out

using genome and transcriptome sequencing data (e.g., Li et al. 2015; Huang et al. 2016; McKain

et al. 2016; Yang et al. 2018) using either the synonymous distance between of paralogs gene

pairs (Ks; Lynch and Conery 2000) or tree-based reconciliation methods (e.g., Jiao et al. 2011;

Li et al. 2015; Yang et al. 2015; Huang et al. 2016; Xiang et al. 2017; Leebens-Mack et al.

2019). Recently, Viruel et al. (2019) showed that allele frequency information from target

enrichment data can be informative in inferring recent WGDs. Target enrichment methods (e.g.,

Mandel et al. 2014; Weitemier et al. 2014; Buddenhagen et al. 2016) have been widely adopted

to collect hundreds to over a thousand nuclear loci for plant systematics, allowing studies at

different evolutionary scales (e.g., Villaverde et al. 2018), and the use of museum-preserved

collections (e.g., Forrest et al. 2019). This creates new opportunities to adopt tree-reconciliation

based methods to explore WGD patterns in groups for which genomic and transcriptomic

resources are not available or feasible.

With >1,100 species worldwide, *Alchemilla* in the broad sense has been a challenging

group to study due to the presence of reticulate evolution, polyploidy, and apomixis (Table S1).

Based on previous phylogenetic analyses, *Alchemilla* s.l. contains four clades: Afromilla,

*Aphanes,* Eualchemilla (or *Alchemilla* s.s.), and *Lachemilla*. Together they form a well-supported

clade nested in the subtribe Fragariinae, which also includes the cultivated strawberries (Gehrke

et al. 2008). Unlike the more commonly recognized members of the rose family (Rosaceae),

*Alchemilla* s.l. is characterized by small flowers with no petals, and a reduced number [1–4(–5)]

of stamens that have anthers with one elliptic theca on the ventral side of the connective that

opens by one transverse split (Perry 1929; Soják 2008). Gehrke et al. (2008) presented the first

phylogeny of *Alchemilla* s.l. and established the paraphyly of traditional *Alchemilla* s.s. into a

primarily African clade, Afromilla, and a Eurasian clade, Eualchemilla. Gehrke et al. (2008) also

suggested treating Afromilla and Eualchemilla, along with *Aphanes* and *Lachemilla* as a single

genus based on nomenclatural stability and the lack of morphological characters to distinguish

between Afromilla and Eualchemilla. The four clades within *Alchemilla* s.l. are mainly defined

by geographic distribution, as well as the number and insertion of the stamens on the disk lining

the hypanthium (Table 1). Phylogenetic analyses using at least one nuclear and one chloroplast

marker (Gehrke et al. 2008; 2016) found significant cytonuclear discordance regarding the

relationships among the four major clades. Similar patterns, often attributed to hybridization and

allopolyploidy, have been detected among most genera in Fragariinae (Lundberg 2009; Eriksson

et al. 2015; Gehrke et al. 2016, Kamneva et al. 2017; Morales-Briones et al. 2018a), leaving the

phylogenetic relationships of *Alchemilla* s.l. to the rest of Fragariinae unresolved. Recent

phylogenomic analyses using target enrichment that included 32 species of *Lachemilla* detected a

putative WGD event (or multiple events) shared by all *Alchemilla* s.l., based on high frequency

of paralogs (Morales-Briones et al. 2018b). However, sampling was limited to one species each

of Eualchemilla, Afromilla, and *Aphanes*, and the location of this shared WGD remains

unresolved.

In this study we increased sampling to 68 species across the major clades of *Alchemilla*

s.l., and included 11 additional closely related species in Fragariinae, which allowed us to 1) test

for polyploid events in the origin of *Alchemilla* s.l., and 2) explore the reticulate evolution within

the clade using a target enrichment dataset. Given the prevalence of polyploidy and reticulate

history within *Alchemilla* s.l., this is an excellent group to explore the utility of tree-based

methods for (1) processing paralogs, and (2) detecting and placing WGDs using target

enrichment datasets.

**MATERIALS AND METHODS**

*Taxon sampling and data collection*

We sampled 68 species representing the four major clades of *Alchemilla* s.l. (sensu Gehrke et al.

2008), and 11 species to represent all other genera in Fragariinae (except *Chamaecallis*; sensu

Morales-Briones and Tank 2019). Additionally, we sampled one species each of *Potentilla*,

*Sanguisorba*, and *Rosa* as outgroups. Voucher information is provided in Table S2. We used a

Hyb-Seq approach (Weitemier et al. 2014), that combines target enrichment and genome

skimming, to target 257 putatively single-copy orthologous genes (in 1,419 exons) using baits

designed for *Fragaria vesca* (strawberries, also member of Fragariinae; Kamneva et al. 2017).

125            Of the 82 total species, only sequences for *Fragaria vesca*, were from a reference

genome (Shulaev et. al 2010). Twenty-two were from a previously published Hyb-Seq dataset

using the same bait set as this study (Morales-Briones et al. 2018b; Table S2), including 19

species of *Lachemilla* that did not show evidence of hybridization within *Lachemilla,* and one

species each of Eualchemilla, Afromilla, and *Aphanes*. Newly generated sequence data for 55

species (Table S2) were collected as follows. Total genomic DNA was isolated from silica-dried

or herbarium material with a modified CTAB method (Doyle and Doyle 1987). Probe synthesis,

library preparation, capture enrichment, and high-throughput sequencing (HiSeq2000 instrument,

$2 \times 101$ bp) were carried out at Rapid Genomics LLC (Gainesville, FL, USA). Data for the

remaining four species, *Drymocallis glandulosa*, *Potentilla indica*, *Rosa woodsii*, and

*Sanguisorba menziesii* were collected as described in Kamneva et al. (2017).

139                         *Read processing and assembly*

We removed sequencing adaptors and trimmed low-quality bases (Phred scores < 20) from raw

reads with SeqyClean v.1.10.07 (Zhbannikov et al. 2017) using default settings. Plastomes were

assembled using Alignreads v.2.5.2 (Straub et al. 2011) and 12 closely related plastome

references (with one Inverted Repeat removed; Table S3). Plastome assemblies were annotated

using *Fragaria vesca* as a reference in Geneious v.11.1.5 (Kearse et al. 2012). Assembly of

nuclear loci was carried out with HybPiper v.1.3.1 (Johnson et al. 2016) using exons of *F. vesca*

as references. Given the large number of paralogs detected in *Lachemilla,* Eualchemilla, and

Afromilla, multi-exon gene assemblies resulted in chimeric sequences of exons from distinct

paralogs (Morales-Briones et al. 2018b). To avoid chimeric sequences that can affect paralog

processing and phylogenetic analyses, assemblies were performed on each exon separately. Only

exons with a reference length of ≥ 150 bp were assembled (939 exons from 257 genes). Paralog

detection was carried out for all exons with the 'paralog_investigator' option in HybPiper. This

option flags loci with potential paralogs when multiple contigs cover at least 85% of the

reference sequence length. Exon assemblies that included flagged paralogs were extracted using

the 'paralog_retriever' command of HybPiper and used for orthology inference.

156                         *Orthology inference for nuclear exons*

To infer orthologs for phylogenetic analyses, all exons were processed as follows (Fig. 1).

Individual exons were aligned using MACSE v.2.03 (Ranwez et al. 2018) with default

parameters. Codons with frameshifts (labeled with '!' by MACSE) were replaced with gaps, and

aligned columns with more than 90% missing data were removed using Phyx (Brown et al.

2017). Initial homolog trees were built using RAxML v.8.2.11 (Stamatakis 2014) with a

GTRCAT model and clade support assessed with 100 rapid bootstrap (BS) replicates.

Monophyletic and paraphyletic grades that belonged to the same taxon were masked by keeping

only the tip with the highest number of characters in the trimmed alignment following Yang and

Smith (2014). To obtain the final homolog trees, outlier tips with unusually long branches were

detected and removed by maximally reducing the tree diameter with TreeShrink v.1.3.2 (Mai and

Mirarab 2018). Orthology inference was carried out using two outgroup-aware strategies from

Yang and Smith (2014). First, we used the 'monophyletic outgroup' (MO) approach keeping

only ortholog groups with at least 25 ingroup taxa. The MO approach filters for homolog trees

with outgroup taxa being monophyletic and single-copy, and therefore filters for single- and low-

copy genes. The second approach used was the 'rooted tree' (RT), with at least 25 ingroup taxa.

The RT approach searches for the subtree with the highest number of ingroup taxa and cuts it out

as a rooted tree. Both approaches root the gene tree by the outgroups, traverse the rooted tree

from root to tip, and remove the side with fewer taxa (MO) or keep both sides (RT) when gene

duplication is detected at any given node. The RT approach maximizes the number of orthologs

compared to MO, but removes outgroups. To add outgroups back to the RT orthologs for

downstream analyses, we kept only RT orthologs from homologs that had a MO ortholog. Then

we used the outgroups of the MO ortholog for all the RT orthologs of the same homolog (Fig 1).

Scripts for orthology inference can be found at

https://bitbucket.org/dfmoralesb/target_enrichment_orthology.

**Figure 1.** Paralog processing workflow and orthology inference methods used in *Alchemilla* s.l. homolog

trees. a) Flow chart of paralog processing and homolog tree inference. b) Outgroup present and

monophyletic. Monophyletic outgroups (MO) will prune single-copy genes keeping clades with at least a

user-defined minimum number of ingroup taxa. Rooted ingroups (RT) will keep all subtrees with at least

a user-defined minimum number of ingroups taxa. If the homolog trees can be pruned using both MO and

RT, then RT orthologs are added to the same root. c) Outgroup present but non-monophyletic. MO

orthologs cannot be obtained from this kind of homologs. RT orthologs can be pruned from this kind

homologs but do not have a root. These RT orthologs can be used for phylogenetic inference (e.g., species

tree inference), but were not used here as some downstream analyses require rooted orthologs.

## *Phylogenetic analyses*

We used concatenation and coalescent-based methods to reconstruct the phylogeny of *Alchemilla*

s.l. Analyses were carried out in the two sets of final orthologs, MO and RT, separately. Each

ortholog exon was aligned using MACSE v.2.03 with default parameters. Codons with

frameshifts were replaced with gaps, and aligned columns with more than 90% missing data

were removed using Phyx, and alignments with at least 150 characters and 25 taxa were retained.

We first estimated a maximum likelihood (ML) tree from the concatenated matrices with

RAxML using a partition by gene scheme with a GTRGAMMA model for each partition. Clade

support was assessed with 100 rapid bootstrap (BS) replicates. To estimate a species tree that is

statistically consistent with the multi-species coalescent (MSC), we first inferred individual ML

gene trees using RAxML with a GTRGAMMA model, and 100 BS replicates to assess clade

support. Individual gene trees were then used to estimate a species tree using ASTRAL-III

v.5.6.3 (Zhang et al. 2018) using local posterior probabilities (LPP; Sayyari and Mirarab 2016) to

assess clade support.

To evaluate nuclear gene tree discordance, we calculated the internode certainty all (ICA)

value to quantify the degree of conflict on each node of the species tree given individual gene

trees (Salichos et al. 2014). Also, we calculated the number of conflicting and concordant

bipartitions on each node of the species trees. We calculated both the ICA scores and the number

of conflicting/concordant bipartitions with Phyparts (Smith et al. 2015) using individual gene

trees with BS support of at least 50% for the corresponding node. Additionally, to distinguish

strong conflict from weakly supported branches, we evaluated tree conflict and branch support

with Quartet Sampling (QS; Pease et al. 2018) using 1,000 replicates. Quartet Sampling

subsamples quartets from the input species tree and concatenated alignment to assess the

confidence, consistency, and informativeness of each internal branch by the relative frequency of

the three possible quartet topologies at each node (Pease et al. 2018).

In addition to species tree construction using inferred orthologs, we used a recently

developed quartet-based species tree method (ASTRAL-Pro; Zhang et al. 2019) to estimate the

phylogeny of *Alchemillla* s.l. ASTRAL-Pro directly uses multi-labeled gene trees while

accounting for gene duplications and losses to estimate a species tree that is statistically

consistent with the MSC and birth-death gene duplication and loss model. We used all 923 final

homolog trees as input for ASTRAL-Pro, ignoring trees with less than 20 taxa, and estimated

LPP to assess clade support. Additionally, we calculated ICA scores and the number of

conflicting/concordant bipartitions with Phyparts using homolog trees with BS support of at least

50% for the corresponding nodes.

For the plastome phylogenetic analyses, 74 partial plastome assemblies and eight

reference plastome sequences were included (Table S3). Plastome sequences were aligned using

the automatic alignment strategy in MAFFT v.7.307 (Katoh and Standley 2013), and aligned

columns with more than 70% missing data were removed with Phyx. We estimated an ML tree

of the plastome alignment with RAxML using a partition by coding (CDS) and noncoding

regions (introns and intergenic spacers) scheme, with a GTRGAMMA model for each partition

and clade support assessed with 100 rapid BS replicates and QS using 1,000 replicates, to detect

potential plastome conflict in the backbone of *Alchemilla* s.l. as recently reported in other groups

(e.g., Gonçalves et al. 2019; Walker et al. 2019; Morales-Briones et al. 2020; Zhang et al. 2020).

237                              *Mapping whole genome duplications*

We took two alternative approaches for detecting WGD events by mapping gene duplication

events from gene trees onto the species tree. The first approach begins by extracting orthogroups

from the final homolog trees. Orthogroups are rooted ingroup lineages separated by outgroups

that include the complete set of genes in a lineage from a single copy in their common ancestor.

We extracted orthogroups requiring at least 50 out of 79 species in Fragariinae. Gene duplication

events were then recorded on the most recent common ancestor (MRCA) on the species tree

when two or more species overlapped between the two daughter clades (Yang et al. 2018). Each

node on a species tree can be counted only once from each gene tree to avoid nested gene

duplications inflating the number of recorded duplications (Yang et al. 2018). We mapped

duplication events onto both the MO and RT species trees using orthogroups from all 923 final

homologs while filtering orthogroups using an BS average of at least 50%. We carried out the

mapping using all exons and using only the longest exon per gene to avoid inflating the counts in

multi-exon genes.

For the second strategy we explicitly tested for polyploidy mode using GRAMPA

(Thomas et al. 2017). GRAMPA uses MRCA reconciliation with multi-labeled gene trees to

compare allo-or autopolyploid scenarios in singly- or multi-labeled species trees. To reduce the

computational burden of searching all possible reconciliations, we constrained searches to only

among crown nodes of major clades of *Alchemilla* s.l. (including the 'dissected' and 'lobed' clades

of Eualchemilla; see results) and genera within Fragariinae. We ran reconciliation searches using

all 923 final homologs, as well as using only the longest exon per gene (256), against either the

MO or RT species tree. We expected multiple WGD events within *Alchemilla* s.l. (see results),

but GRAMPA can only infer one WGD at a time. To disentangle nested duplication events, we

also carried out similar GRAMPA reconciliations using the MO species tree and sequentially

excluding major groups of *Alchemilla* s.l. that were identified as a polyploidy clade. We only

used the MO species tree as the only difference with the RT species trees is the location of the

'lobed' clade that was first clade identified as allopolyploid (see results). The removal of the

'lobed' clade from either the MO or RT species trees resulted in the same backbone for the

remaining *Alchemilla* s.l. Finally, to test for a polyploid origin of *Alchemilla* s.l., we carried out

searches among the constrained crown node of *Alchemilla* s.l. and the rest of the genera within

Fragariinae using the MO and cpDNA trees. The backbone of Fragariinae differed between the

MO (same as RT) species tree and the cpDNA tree. Thus, we tested how this affected the

inference of the polyploidy origin of *Alchemilla* s.l. We also carried out similar searches but

using each of the five major clades individually.

272                *Distribution of synonymous distance among gene pairs (Ks plots)*

To verify WGD events inferred from target enrichment, we analyzed the distribution of

synonymous distances (Ks) from RNA-seq data of four species of *Alchemilla* s.l. and nine

species of Fragariinae (Table S4). Read processing and transcriptome assembly followed

Morales-Briones et al. (2020). For each of the four species, a Ks plot of within-taxon paralog

pairs based on BLASTP hits was done following Yang et al. (2015). Significant Ks peaks were

identified using a mixture model as implemented in mixtools v.1.2.0 (Benaglia et al. 2009). The

optimal number of mixing components was estimated using parametric bootstrap replicates of

the likelihood ratio test statistic (McLachlan and Peel 2000). We tested up to five components

using 500 bootstrap replicates in mixtools.

283                                **RESULTS**

284                          *Assembly and orthology inference*

The number of assembled exons per species (with > 25% of the target length) ranged from 692

(*Alchemilla fissa*) to 939 (*Farinopsis salesoviana*) out of 939 single-copy exon references from

*F. vesca*, with an average of 902 exons (Table S5). The number of exons with paralog warnings

ranged from 10 in *Drymocallis glandulosa* to 746 in *Alchemilla mollis* (Table S5). The number

of exon alignments with ≥ 25 species was 923 from 256 genes. The orthology inference resulted

in 914 MO orthologs (Table S6), and 1,906 RT orthologs (Table S6). The trimmed alignments of

the MO orthologs ranged from 136 to 5,740 characters with a mean of 425 characters (median =

268). The concatenated alignment of the MO orthologs, with at least 150 aligned characters and

25 species for each exon, included 910 exons and 387,042 characters with a matrix occupancy of

66%. The trimmed alignments of the RT orthologs ranged from 136 to 5,740 characters with a

mean of 394 characters (median = 259). The concatenated alignment of the RT orthologs, with at

least 150 aligned characters and 25 species, included 1,894 exons and 746,562 characters with a

matrix occupancy of 54%. The chloroplast alignment included 124,079 characters.

299                                        *Nuclear phylogenetic analyses*

All nuclear analyses recovered the monophyly of *Alchemilla* s.l. with maximum support (i.e.,

bootstrap percentage [BS] = 100, local posterior probabilities [LPP] = 1.0; Fig. 2; Fig. S1), most

gene trees being concordant (858 out of 863 for MO; 977/984 for RT; 912/932 for ASTRAL-

Pro; ICA = 0.95), and full QS support (1.0/–/1.0; i.e., all sampled quartets supported that node).

Five major clades were identified within *Alchemilla* s.l.: Afromilla, *Aphanes*, Eualchemilla-

'dissected', Eualchemilla-'lobed,' and *Lachemilla*. Moreover, the relationships among these

clades showed high levels of discordance and varied among the MO and RT species trees.

307            Analyses of the MO ortholog exons using ASTRAL and concatenated ML approaches

resulted in similar topologies for the backbone of *Alchemilla* s.l. (Fig. 2). The monophyly of the

five major clades each received maximum support (BS = 100; LPP = 1.0) and had most gene

trees being concordant (except for the two clades of Eualchemilla). Eualchemilla was

paraphyletic and split into the 'dissected' and 'lobed' clades. The 'dissected' clade was supported

by 73 out of 420 informative gene trees (ICA = 0.06) and strong QS score (0.61/0.98/0.99),

while the 'lobed' clade was supported by 113 out of 316 informative gene trees (ICA = 0.06) and

strong QS score (0.87/0.34/1). In both cases, the 'dissected' and 'lobed' clades had a relatively

small percentage of supporting gene trees, but the conflict analysis and QS score did not reveal

any well-supported alternative topology. *Aphanes* was recovered as sister to the Eualchemilla-

'lobed' clade with relatively low support (BS = 90, LPP = 0.62), 60 concordant gene trees (out of

430 informative gene trees; ICA = 0.08), and weak QS score (0.016/0.95/0.98) with similar

frequencies for all three possible alternative topologies. The Eualchemilla-'dissected' clade was

recovered as sister to Eualchemilla-'lobed' + *Aphanes* with maximum support, 279 concordant

gene trees (out of 482 informative gene trees; ICA = 0.29), and full QS score. Afromilla was

recovered as sister to the clade formed of Eualchemilla ('dissected and 'lobed') and *Aphanes*

with high to low support (BS = 100, LPP = 0.88), only 146 concordant gene trees (out of 413

informative gene trees; ICA = 0.22), and weak QS support (0.2/0.44/0.99) with a skew in

discordance suggesting a possible alternative topology. Lastly, *Lachemilla* was recovered as the

sister to the rest of *Alchemilla* s.l.

Analysis of the RT ortholog exons using ASTRAL and concatenated ML approaches

both recovered the same major clades, but they differed in the relationship among these five

clades (Fig 2a; Fig. S1). In both analyses, *Lachemilla*, Afromilla, and *Aphanes* had maximum

support (BS = 100; LPP = 1.0) and had most gene trees being concordant. Eualchemilla was

recovered as monophyletic and composed of the 'dissected' and 'lobed' clades. Eualchemilla had

high to low support (BS = 99, LPP = 0.63), only 231 concordant gene trees (out of 819

informative gene trees; ICA = 0.07), and weak QS support (0.023/0.87/0.98) with similar

frequencies for all three possible alternative topologies. As in the MO analyses, the 'dissected'

and 'lobed' clades had the maximum support, low number of concordant gene trees (218 out of

557 [ICA = 0.19] and 136 out of 707 [ICA = 0.08], respectively), and strong QS support

(0.98/0/1 and 0.62/0.17/0.99, respectively). Both the 'dissected' and 'lobed' clades had a reduced

number of supporting gene trees. Still, in the case of 'dissected' clade, the conflict analysis and

QS score did not reveal any well-supported alternative topology. In contrast, in the 'lobed' clade,

only the QS score showed the possible presence of a supported secondary topology.

Eualchemilla was recovered as sister of *Aphanes* with maximum support (BS = 100), 348

concordant gene trees (out of 728 informative gene trees; ICA = 0.29) and full QS support. The

ML concatenated tree (Fig 2a; Fig. S1) placed Afromilla as sister to the clade formed of

Eualchemilla and *Aphanes* with maximum support (BS = 100), 212 concordant gene trees (out of

771 informative gene trees; ICA = 0.27), and weak QS support (0.18/0.66/0.99) with a small

skew in discordance suggesting a possible alternative topology. *Lachemilla* was placed as sister

to the rest of *Alchemilla* s.l. The ASTRAL tree in turn (Fig S1a), retrieved *Lachemilla* as sister to

the clade formed of Eualchemilla and *Aphanes* with no support (LPP = 0.01), 247 concordant

gene trees (out of 953 informative gene trees; ICA = 0.19), and QS counter-support

(−0.21/0.29/0.99), showing that the majority of the quartets supported one of the alternatives

topologies. In this case, Afromilla was placed as sister to the rest of *Alchemilla* s.l.

The ASTRAL-Pro analysis using multi-labeled homolog trees recovered the same

backbone topology of *Alchemilla* s.l. as the concatenated ML analysis from the RT orthologs

(Fig. 2b; Fig. S3). All five major clades had the maximum support (LPP = 1.0). Eualchemilla,

composed of the 'dissected' and 'lobed' clades, had high to low support (LPP = 0.76) and only

502 concordant trees (out of 1,274 informative trees; ICA = 0.09). The 'dissected' and 'lobed'

clades had low numbers of concordant trees (402 out of 1,157 [ICA = 0.13] and 97 out of 975

[ICA = 0.05], respectively), but did not showed signal of any well-supported alternative

topology. *Aphanes* was placed as the sister of Eualchemilla with maximum support (LPP = 1.0),

577 concordant trees (out of 1,329 trees; ICA = 0.16), and no support for any well-supported

alternative topology. Afromilla was recovered as sister to the clade formed of Eualchemilla and

*Aphanes* with low support (LPP = 0.52), 649 concordant trees (out of 1,415 trees; ICA = 0.20),

and no support for any well-supported alternative topology.

365                                 *Chloroplast phylogenetic analyses*

The chloroplast ML tree (Fig 2c; Fig S4) recovered a well-supported backbone *Alchemilla* s.l.

where the monophyly of *Aphanes,* Afromilla, and *Lachemilla,* had maximum or near maximum

support (i.e., bootstrap percentage [BS] = 100, QS support = [1.0/–/1.0]). Eualchemilla,

composed of the 'dissected' and 'lobed' clades, also had the maximum support. The 'dissected'

clade had strong support (BS = 75), but the QS score (0.8/0.43/0.88) showed a small skew in

discordance suggesting a possible alternative topology. Similarly, the 'lobed' clade also had

strong support (BS = 100) and a QS score (0.95/0.25/0.92) that suggested a possible alternative

topology. *Aphanes* and Eualchemilla formed, with maximum support, a clade as in the nuclear

analyses. In turn, Afromilla and *Lachemilla* were recovered as sister clades with maximum

support, which differed from the nuclear analyses.

**Figure 2.** Maximum likelihood phylogeny of *Alchemilla* s.l. inferred from RAxML analysis of the concatenated 910-nuclear gene supermatrix from the 'monophyletic outgroup' (MO) orthologs. Bootstrap support (BS) and Local posterior probability (LLP) are shown above branches. Nodes with full support (BS= 100/LLP= 100) are noted with an asterisk (*). Em dashes (—) denoted alternative topology compared to the ASTRAL species tree. Quartet Sampling (QS) scores for major clades are shown below branches. QS scores in blue indicate strong support and red scores indicate weak support. QS scores: Quartet concordance/Quartet differential/Quartet informativeness. Pie charts for major clades represent

the proportion of exon ortholog trees that support that clade (blue), the proportion that support the main

alternative bifurcation (green), the proportion that support the remaining alternatives (red), and the

proportion (conflict or support) that have < 50% bootstrap support plus missing data (gray). Branch

lengths are in number of substitutions per site (scale bar on the bottom). Inset: a) Summary Maximum

likelihood phylogeny inferred from RAxML analysis of the concatenated 1,894-nuclear gene supermatrix

from the 'rooted ingroup' orthologs (RT). BS and LLP are shown above branches and QS scores below

the branches. Branch lengths are in number of substitutions per site; b) Summary ASTRAL-Pro species

trees inferred from 923 multi-labeled homolog trees. LLP are shown next to nodes. Branch lengths are in

coalescent units. c) Summary Maximum likelihood phylogeny inferred from RAxML analysis of partial

plastomes. BS and LLP are shown above branches and QS scores below the branches. Branch lengths are

in number of substitutions per site.

397                                    *Mapping whole genome duplications*

By mapping the most recent common ancestor (MRCA) of gene duplication events from rooted

orthogroup trees onto the species tree, we found four nodes in *Alchemilla* s.l. that each had an

elevated proportion of gene duplications (Fig. 3a–b). This trend was consistent regardless of

using all 923 homologs exons (826 after orthogroup inference and BS filtering) or using only the

256 longest exons for each gene (250 after orthogroup inference and BS filtering; Fig. S5).

Therefore, here we describe the results only for the latter. These four clades include (Fig. 3a; Fig.

S5): 1) the MRCA of *Alchemilla* s.l. (86.0% of the 250 genes show evidence of duplication), 2)

the MRCA of Eualchemilla , *Aphanes*, and Afromilla (34.4%), 3) the MRCA of Eualchemilla +

*Aphanes* (MO: 18.4%; RT:15.6% ), and 4) the MRCA of *Lachemilla* (18.4%). These four nodes

have an elevated proportion of gene duplications compared to all other nodes in Fragariinae (Fig.

3b) and it is consistent with the number of paralogs counted from the final homolog trees (Fig.

3c). Interestingly, *Aphanes* had a lower number of paralogs than the rest of *Alchemilla* s.l.,

resembling the other members of Fragariinae (Fig. 3c).

**Figure 3.** Orthogroup gene duplication mapping results. a) Summarized cladogram of *Alchemilla* s.l.
from the 'monophyletic outgroup' (MO) orthologs species trees. Percentages next to nodes denote the
proportion of duplicated genes when using orthogroups from only one exon per gene. Nodes with
elevated proportions of gene duplications are numbered 1–4 as referenced in the main text. See Fig. S6
for the full tree. b) Histogram of percentages of gene duplication per branch. c) Number of paralogs per
taxa in the final homolog trees. Each locus is represented by the longest exon per gene.

Similar to the results of MRCA mapping, the GRAMPA analyses recovered the same

results when using all 923 homolog exons or only one exon per gene (256). GRAMPA

reconciliations using all major clades of *Alchemilla* s.l. recovered multi-labeled trees with the

best score where the 'lobed' clade of Eualchemilla was of an allopolyploid origin, but the putative

parental lineages varied between the MO and RT ortholog species trees. The reconciliations

using the MO species tree (Fig. 4a; Fig. S6) showed that the 'lobed' clade was of allopolyploid

origin between an unsampled or extinct lineage sister to *Aphanes* and an unsampled or extinct

lineage ('lineage' for short hereafter) sister to 'dissected' + *Aphanes*. In turn, the reconciliations

using the RT species tree (Fig. S6) showed that the 'lobed' clade was of allopolyploid origin

between a 'lineage' sister to the 'dissected' clade, and also a 'lineage' sister to 'dissected' +

*Aphanes*.

The GRAMPA reconciliations performed on the MO species trees with removal of major

clades of *Alchemilla* s.l. inferred as allopolyploid resulted in the identification of additional

polyploidy events (Fig. 4b–d ). First, we removed the 'lobed' clade that resulted in the recovery

of Afromilla as an allopolyploid clade. Afromilla parental lineages were a 'lineage' sister to

*Aphanes* + the 'dissected' clade, and a 'lineage' sister to all remaining *Alchemilla* s.l. (Fig. 4b).

The further removal of Afromilla resulted in recovery of the 'dissected' clade as allopolyploid.

The 'lobed' clade had parental lineages to a 'lineage' sister to *Aphanes* and to a 'lineage' sister to

all remaining *Alchemilla* s.l. (Fig. 4c). Finally, the removal of the 'dissected' clade resulted in the

*Lachemilla* being recovered also as an allopolyploid clade. The parental lineages of *Lachemilla*

were a 'lineage' sister to *Aphanes* and a 'lineage' sister to all remaining *Alchemilla* s.l. (Fig. 4d).

The GRAMPA results from the analyses with constrained searches on the crown node of

*Alchemilla* s.l. recovered different modes of polyploidy when using the MO species tree versus

the cpDNA tree. The MO species tree had *Farinopsis*, *Sibbaldianthe* + *Sibbaldia*, *Comarum*, and

*Fragaria* forming a grade sister to *Alchemilla* s.l., while *Drymocallis*, *Chamaerhodos*, *Potaninia*,

and *Dasiphora* form a clade that is sister to all other Fragariinae (Fig. 2). The reconciliations on

the MO tree resulted in an allopolyploid event for the clade composed of *Alchemilla* s.l.,

*Farinopsis, Sibbaldianthe*, and *Sibbaldia* (Fig. S7)*. The parental lineages of this clade were a

'lineage' sister to *Comarum*, and a 'lineage' sister to the grade formed of *Comarum* and

*Fragaria* (Fig. S7). The reconciliations using individual major clades of *Alchemilla* s.l. resulted

in identical patterns as in the full constrained analysis (Fig. S8). The cpDNA tree had *Alchemilla*

s.l. as sister to the grade formed of *Farinopsis*, *Comarum*, and *Sibbaldianthe + Sibbaldia*, while

*Fragaria* was recovered as sister to the clade composed of *Drymocallis*, *Chamaerhodos*,

*Potaninia*, and *Dasiphora,* which is sister to all other Fragariinae (Fig. S6). The reconciliations

on the cpDNA tree recovered *Alchemilla* s.l. as of autopolyploid origin (Fig. 4e; Fig. S7). The

analyses using individual major clades of *Alchemilla* s.l. recovered identical patterns as in the

full constrained analysis, except for *Aphanes* that resulted in a singly-labeled tree (Fig. S8).

The distribution of synonymous distances in the four species of Eualchemilla (one

'dissected' and three 'lobed') shared three optimal mixing components with a Ks mean at

approximately 0.1, 0.34, and 1.67, respectively (Fig. S9). All nine species from other genera in

Fragariinae had two optimal mixing components with a shared Ks peak at 1.44–1.86. The only

two polyploid species from the other genera of Fragariinae, *Sibbaldianthe bifurca* ($2n=28$) and

*Comarum palustre* ($2n=64$), had KS peaks at 0.11 and 0.46, respectively (Fig. S9).

**Figure 4.** Summarized optimal multi-labeled tree (MUL-tree) inferred from GRAMPA analyses. a) MUL-tree for the complete 'monophyletic outgroup' (MO) orthologs species tree. Red branches denote the allopolyploid origin of the 'lobed' clade of Eualchemilla. b) MUL-tree for MO species trees after the removal of the 'lobed' clade of Eualchemilla. Green branches denote the allopolyploid origin of Afromilla. c) MUL-tree for MO species trees after the removal of the 'lobed' clade and Afromilla. Blue branches denote the allopolyploid origin of the 'dissected' clade of Eualchemilla. d). MUL-tree for MO species trees after the removal of the 'lobed' and 'dissected' clades, and Afromilla. Green lines denote the allopolyploid origin of *Lachemilla*. e) MUL-tree using constrained searches the crown node of *Alchemilla* s.l. on the cpDNA tree. Orange branches denote the autopolyploid origin *Alchemilla* s.l. f) Putative summary network of all reticulation events in *Alchemilla* s.l. Colored curved branches denote different polyploid events as in a–e. Dashed curved lines represent the maternal lineage (cpDNA) in allopolyploid events.

## DISCUSSION

### *Processing paralogs in target enrichment datasets*

The increased use of target enrichment methods in combination with reduced sequencing costs

and higher read coverage have facilitated the recovery of paralogs in such datasets. Paralogy is

viewed as a nuisance for phylogenetic reconstruction and is commonly aimed to be reduced in

early stages of experimental design, by targeting only single- or low-copy genes during the

selection of loci (e.g., Chamala et al. 2015; Nicholls et al. 2015; Gardner et al. 2016; Kamneva et

al. 2017). Still, the recovery of paralogs is inevitable when working with groups where WGD is

prevalent, especially in plants, leading to various strategies to remove them prior to phylogenetic

analyses. Commonly used target enrichment assembly pipelines (e.g., Faircloth 2016; Johnson et

al. 2016; Andermann et al. 2018) use different criteria to flag assembled loci with putative

paralogs that are later filtered or sorted prior to phylogenetic analysis. The most common

approach for processing paralogs in target enrichment datasets is removing the entire loci that

show any signal of potential paralogy (e.g., Crowl et al. 2017; Montes et al. 2019; Bagley et al.

2020). Other approaches either retain or remove contigs based on the distinction between

putative allelic variation (flagged sequences form monophyletic groups) or putative paralogs

(non-monophyletic) in combination with study-specific threshold or random selection (e.g.,

Villaverde et al; 2018; Liu et al. 2019; Stubbs et al. 2019), or manual processing (e.g., Garcia et

al. 2017; Karimi et al. 2019). Using these approaches, especially the removal of entire loci, can

significantly reduce the size of target enrichment datasets. As dataset size increases, manual

processing becomes prohibitive.

The presence of WGDs also poses some challenges for locus assembly. Target

enrichment design commonly includes multi-contig targets that assembly pipelines attempt to

assemble into single contigs (e.g. Faircloth 2016) or 'supercontigs' composed of multiple exons

and partially assembled introns (e.g., Johnson et al. 2016). In groups like *Alchemilla* s.l., where

multiple, nested WGD events led to a prevalence of closely-related paralogs, 'supercontigs'

produces chimeric assemblies (Morales-Briones et al. 2018b). Instead, we assembled the exons

individually to minimize chimeric loci, at the cost of generating short exons that may produce

poorly resolved gene trees, which can affect orthology inference and downstream phylogenetic

analyses. Therefore, it is important to take this into consideration during target enrichment

experimental design, and to preferentially target long exons when possible. An alternative

strategy to avoid chimeric supercontig when gene duplications are frequent is to perform a

preliminary orthology inference in single exon-based trees and then use the inferred orthologs as

a reference to reassemble the loci into 'supercontigs' (e.g., Gardner et al. 2020; Karimi et al.

2020).

The utility of paralogs for phylogenetic reconstruction in target enrichment datasets is

gaining more attention (e.g., Johnson et al. 2016; Gardner et al. 2020). A few studies have

considered tree-based orthology inference to process affected loci (e.g., Garcia et al. 2017;

Moore et al. 2018, Morales-Briones et al. 2018b), but in some cases the orthology approaches

used cannot be applied to other groups. Here we demonstrated the utility of automated, tree-

based orthology inference methods (Yang and Smith 2014), originally designed for genomic or

transcriptomic datasets, to infer orthology from paralog-flagged loci in a target enrichment

dataset. Our approach facilitates the automated inference of orthologs while maximizing the

number of loci retained for downstream analyses. It is suitable for any type of target enrichment

dataset (e.g., anchored phylogenomics, exon capture; Hyb-Seq, ultraconserved elements) as long

as individual gene trees are informative.

Orthology inference methods used here (Yang and Smith et al. 2014) are a powerful tool

for target enrichment datasets. In the case of allopolyploidy, however, these methods can

introduce bias in the distribution of ortholog trees inferred. In the case of MO, each time a gene

duplication event is detected, the side with a smaller number of taxa is removed. When

allopolyploidy occurs, MO will likely result in preserving one subgenome instead of the other.

One could alternatively preserve a random side each time a gene duplication event is identified.

However, in practice, the side with a smaller number of taxa often contains misassembled or

misplaced sequences. The RT method of separating duplicated gene copies, on the other hand,

keeps any subtree with sufficient number of taxa, but removes outgroups, and worked best when

hierarchical outgroups were included in the taxon sampling. Therefore, both MO and RT lose

information, especially in cases with complex, nested polyploidy. Recently developed methods

based on quartet similarity (Zhang et al. 2019) or Robinson-Foulds distances (Molloy and

Warnow 2020) can directly estimate species trees from multi-labeled trees that are consistent

with the MSC and gene duplication and loss without inferring orthologs. However, their

behavior on complex datasets using archival materials is yet to be explored. For example, both

methods do not define ingroup-outgroup relationships a priori, and correctly inferring the root of

homolog trees can be challenging with missing data, or when WGD occurs near the root. In

addition, none of these above methods were designed to explicitly handle reticulate relationships.

This can result in species tree topology that is an "average" between subgenomes. Depending on

the topological distance of subgenomes, the resulting species tree may not represent any

subgenome history. Finally, most current methods for evaluating node support still require

orthologous gene trees as input. In such cases tools like Phyparts can still be used to visualize

gene tree discordance and calculate ICA scores using multi-labeled trees.

548                                *Phylogenetic implications in Alchemilla s.l.*

Previous phylogenetic studies established the monophyly *Alchemilla* s.l. and four major clades of

the group (Gehrke et al. 2008; 2016), but the relationship among them and the placement of

*Alchemilla* s.l. within Fragariinae remain unresolved. Our nuclear and plastid analyses both

confirmed the monophyly of *Alchemilla* s.l. and its sister relationship to *Farinopsis*, as

previously shown by Morales-Briones and Tank (2019) based on plastome sequences only.

Gehrke et al. (2008) identified two well supported clades within Eualchemilla that were

distinguished by leaf shape, namely the 'dissected' and 'lobed' clades. Most species of

Eualchemilla have a leaf shape consistent with their clade placement, but some had different leaf

shapes that were attributed to their hybrid origin between the two clades (Gehrke et al. 2008).

More recently, Gehrke et al. (2016) and Morales-Briones and Tank (2019) found that

Eualchemilla is not monophyletic in analyses that included the external transcribed spacer (ETS)

of the nuclear ribosomal DNA (nrDNA) cistron. Both studies found *Aphanes* nested between the

'dissected' and 'lobed' clades of Eualchemilla. Our analyses of the nuclear loci supported the

monophyly of 'dissected' and 'lobed' clades, but the monophyly of Eualchemilla had low support

(Fig. 2; Fig. S1). The analysis using only the MO orthologs even weakly supported the 'lobed'

clade as sister of *Aphanes* (Fig. 2). In contrast, our plastome analysis recovered a well-supported,

monophyletic Eualchemilla, as well as well-supported 'dissected' and 'lobed' clades. Both nuclear

and plastid analyses strongly supported the clade composed of *Aphanes* and both clades of

Eualchemilla (Fig. 2; Fig. S4), a relationship that is consistent with previous nuclear and plastid

analyses (Gehrke et al. 2008; 2016; Morales-Briones and Tank 2019). Given the revealed

hybridization in the evolution and early divergence within *Alchemilla* s.l., the non-monophyly of

Eualchemilla could be explained by ancient gene flow or an allopolyploid origin of the

'dissected' and 'lobed' clades (Fig. 4a,c; see below). Besides the well supported relationship of

Eualchemilla + *Aphanes*, our nuclear analysis showed high levels of conflict among the other

clades (Fig. 2; Fig. S1) which could also be explained by additional ancient allopolyploid events

within *Alchemilla* s.l. (Fig. 4; see below).

576                                    *Ancient allopolyploidy in Alchemilla s.l.*

Whole-genome duplications are frequent across Rosaceae (Dickinson et al. 2007; Xiang et al.

2017), and allopolyploidy has been suggested as the primary source for the cytonuclear

discordance in Fragariinae (Lundberg et al. 2009; Gehrke et al. 2016; Morales-Briones and Tank

2019). We recovered four nodes in *Alchemilla* s.l. with a high percentage of gene duplications

(Fig. 3a; Fig. S5). One of the nodes showing a high percentage of gene duplication (18.4%) was

the MRCA of *Aphanes* and Eualchemilla (node 3 in Fig. 3a; Fig. S5). This duplication event

agreed with the MRCA of the ancestral lineages inferred with GRAMPA for the allopolyploid

origin of the 'lobed' clade of Eualchemilla (Fig. 4a). Moreover, the GRAMPA reconciliations

after the removal of the 'lobed' clade and Afromilla (Fig. 4c) inferred a scenario where the

'dissected' clade is of allopolyploid origin. Ks plots of all species of the 'dissected' and 'lobed'

clades shared two peaks that are missing from taxa outside of *Alchemilla* s.l. (Fig. S10)

suggested that at least two WGD events have happened between the stem lineage of *Alchemilla*

s.l. to the crown node of the dissected and the lobed clades. Given the elevated ploidy level of

the sampled species ($2n$=102–156) and that the sister relationship of the 'dissected' and 'lobed'

clades are not supported by nuclear genes, the two clades of Eualchemilla seem to have had

independent allopolyploid origins, while sharing the same or a closely related maternal lineage

(cpDNA; Fig. 4f).

594            The next node with high percentage of gene duplication (34.4%) was placed at the

MRCA of the clade composed of Afromilla, Eualchemilla, and *Aphanes* (node 2 in Fig. 3a). The

GRAMPA reconciliation (after the removal of the 'lobed' clades) recovered an allopolyploid

origin of Afromilla (Fig. 4b), with the MRCA of inferred parental lineages being the base of

*Alchemilla* s.l., consistent with the high percentage of gene duplication mapped to that node

(86.0% at node 1; Fig. 3a). After removing both Afromilla and Eualchemilla, *Lachemilla* was

recovered as allopolyploid, again with the MRCA of parental lineages mapped to the base of

*Alchemilla* s.l. (Fig. 4d)

602            Finally, the node with the highest percentage of duplicated genes (86%) was placed at the

MRCA of *Alchemilla* s.l. (node 1 in Fig. 3a). GRAMPA analysis using the MO species tree

showed an allopolyploid event for the clade that included *Alchemilla* s.l., *Farinopsis salesoviana*,

*Sibbaldia*, and *Sibbaldianthe* (Fig. S7). However, an allopolyploid origin of *Farinopsis*

*salesoviana*, *Sibbaldia*, and *Sibbaldianthe* is not supported by chromosome numbers, orthogroup

gene duplication counts, or Ks plots. All members in Fragariinae, with the exception of

*Alchemilla* s.l. mainly consists of diploid species and base chromosome number of seven ($n$ =7),

including *Sibbaldia* and *Sibbaldianthe.* On the other hand, *Alchemilla* s.l. has a base number of

eight ($n$ =8), and contains mostly species with high ploidy levels (octoploid to 24-ploid), with the

exception of most species of *Aphanes* ($2n$=16) and one species of *Lachemilla* (*L. mandoniana,*

$2n$=16). Also, our duplication counts show low percentages (1.6%) of gene duplication for the

MRCA of the GRAMPA-inferred allopolyploid clade or the MRCA (3.6%) of the inferred

parental lineages (Fig. 3). Furthermore, previous phylotranscriptomic analyses of Rosaceae

(Xiang et al. 2017) that included one species each of the 'dissected' and 'lobed' clades of

Eualchemilla, found 33.21% of duplicated genes for the MRCA of these two clades, but did not

recover any other node with elevated gene duplications within Fragariinae. The Ks plots of the

four species of *Alchemilla* s.l. all showed peaks with similar Ks means among them, but these

largely differed from Ks plots for diploid and polyploid species of *Sibbaldia* and *Sibadianthe*,

and do not share any peak. Instead, *Sibbaldia* and *Sibadianthe* showed Ks patterns similar to

other genera across Fragariinae (Fig. S9). Although the chromosome number and Ks data for

*Farinopsis salesoviana* is not available, all these suggest an unlikely allopolyploid origin of

*Farinopsis, Sibbaldia*, and *Sibadianthe* along with *Alchemilla* sl. In turn, the GRAMPA

reconciliations using the cpDNA tree resulted in an optimal multi-labeled tree where *Alchemilla.*

s.l. had an autopolyploid origin (Fig. 4e). This scenario is compatible with the high percentage of

gene duplication at the MRCA of *Alchemilla* s.l. and the low percentage of gene duplication in

the rest of Fragariinae. Another compatible scenario is an allopolyploid origin of *Alchemilla* s.l.

where both parental lineages are missing or extinct, but this scenario is indistinguishable from

autopolyploidy. The atypical high proportion of gene duplication in the crown of *Alchemilla* s.l.

can be explained by the inferred autopolyploid event. However, this high proportion of gene

duplication could also be the results of gene tree estimation error (e.g., uninformative genes),

incomplete lineage sorting (ILS), allopolyploid events among major clades of *Alchemilla* s.l.,

and/or homoeologous exchanges among subgenomes (Edger et al. 2018; McKain et al. 2018).

Altogether, our results showed evidence that the entire *Alchemilla* s.l. is the product of an ancient

autopolyploidy event, and that Afromilla, Eualchemilla ('lobed' and 'dissected' clades), and

*Lachemilla* originated from subsequent and nested ancient allopolyploid events (Fig. 4f).

While GRAMPA has been shown to be useful to identify multiple polyploidy events in

the same tree (e.g., Thomas et al. 2017; Koenen et al. 2020), this tree-based approach can also

be sensitive to gene tree estimation error or ILS (Thomas et al. 2017). Methods to infer species

networks in the presence of ILS (e.g., Solís-Lemus and Ané 2016; Wen et al. 2018) could also be

used to explore the prevalence of ancient hybridization in *Alchemilla* s.l. Although these methods

are under constant development and improvement, they are still only tractable in simple

scenarios with few reticulation events (Hejase and Liu 2016; Kamneva and Rosenberg 2017).

Similarly, the signal of the *D*-Statistic (Green et al. 2010; Durand et al. 2011), commonly used to

detect introgression, can be lost or distorted in presence of multiple reticulations (Elworth et al.

2018). Complex reticulate scenarios like *Alchemilla* s.l. are likely to face these problems and

have phylogenetic network and *D*-statistic identifiability issues as seen in other groups (e.g.,

Morales-Briones et al. 2020).

650                                                      *Conclusions*

In this study, we have shown the potential of target enrichment datasets in combination

with tree-based methods for paralog processing and WGD investigation. We have used

*Alchemilla* s.l., a group showing nested WGDs, to highlight the potential of these methods in

investigating cases of complex scenarios. Our study shows the need in selecting long exons that

produce informative gene trees for target enrichment studies. As research continues to deepen in

other clades across the Tree of Life using similar methods, we expect that complex patterns of

duplications and reticulations, as those shown here in *Alchemilla* s.l. are more common than we

previously thought.

660　　　　　　　　　　　　　　　SUPPLEMENTARY MATERIAL

661　Data available from the Dryad Digital Repository: http://dx.doi.org/10.5061/.[NNNN]

663　　　　　　　　　　　　　　　　ACKNOWLEDGMENTS

664　This work was supported by the University of Minnesota, Graduate Student Research Grants

665　from the Botanical Society of America, American Society of Plant Taxonomists, International

666　Association of Plant Taxonomists, the University of Idaho Stillinger Herbarium Expedition

667　Funds to D.F.M.-B., and a National Science Foundation Doctoral Dissertation Improvement

668　Grant to D.C.T. for D.F.M.-B. (DEB-1502049).

670　　　　　　　　　　　　　　　　　REFERENCES

Andermann T., Cano Á., Zizka A., Bacon C., Antonelli A. 2018. SECAPR—a bioinformatics

pipeline for the rapid and user-friendly processing of targeted enriched Illumina

sequences, from raw reads to alignments. PeerJ. 6:e5175.

Andermann T., Torres Jiménez M.F., Matos-Maraví P., Batista R., Blanco-Pastor J.L.,

Gustafsson A.L.S., Kistler L., Liberal I.M., Oxelman B., Bacon C.D., Antonelli A. 2020.

A Guide to Carrying Out a Phylogenomic Target Sequence Capture Project. Front.

Genet. 10:1407.

Bagley J.C., Uribe-Convers S., Carlsen M.M., Muchhala N. 2020. Utility of targeted sequence

capture for phylogenomics in rapid, recent angiosperm radiations: Neotropical

Burmeistera bellflowers as a case study. Mol. Phylogenet. Evol.:106769.

Benaglia T., Chauveau D., Hunter D.R., Young D. 2009. mixtools : An R Package for Analyzing

Finite Mixture Models. J. Stat. Softw. 32:1–29.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx - phylogenetic tools for unix. Bioinformatics.

684             33:1886–1888.

685     Buddenhagen C., Lemmon A.R., Lemmon E.M., Bruhl J., Cappa J., Clement W.L., Donoghue

686             M.J., Edwards E.J., Hipp A.L., Kortyna M., Mitchell N., Moore A., Prychid C.J.,

687             Segovia-Salcedo M.C., Simmons M.P., Soltis P.S., Wanke S., Mast A. 2016. Anchored

688             Phylogenomics of Angiosperms I: Assessing the Robustness of Phylogenetic Estimates.

689             bioRxiv.:086298.

690     Chamala S., García N., Godden G.T., Krishnakumar V., Jordon-Thaden I.E., De Smet R.,

691             Barbazuk W.B., Soltis D.E., Soltis P.S. 2015. MarkerMiner 1.0: A new application for

692             phylogenetic marker development using angiosperm transcriptomes. Appl. Plant Sci.

693             3:1400115.

694     Crowl A.A., Manos P.S., McVay J.D., Lemmon A.R., Lemmon E.M., Hipp A.L. 2019.

695             Uncovering the genomic signature of ancient introgression between white oak lineages

696             (Quercus). New Phytol.:nph.15842.

697     Dickinson T.A., Lo E., Talent N. 2007. Polyploidy, reproductive biology, and Rosaceae:

698             understanding evolution and making classifications. Plant Syst. Evol. 266:59–78.

699     Doyle J.J., Doyle J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf

700             tissue. Phytochem. Bull. 19:11–15.

701     Dunn C.W., Howison M., Zapata F. 2013. Agalma: an automated phylogenomics workflow.

702             BMC Bioinformatics. 14:330.

703     Durand E.Y., Patterson N., Reich D., Slatkin M. 2011. Testing for Ancient Admixture between

704             Closely Related Populations. Mol. Biol. Evol. 28:2239–2252.

705    Edger P.P., McKain M.R., Bird K.A., VanBuren R. 2018. Subgenome assignment in

706            allopolyploids: challenges and future directions. Curr. Opin. Plant Biol. 42:76–80.

707    Elworth R.A.L., Allen C., Benedict T., Dulworth P., Nakhleh L.K. 2018. DGEN: A Test Statistic

708            for Detection of General Introgression Scenarios. WABI.

709    Emms D.M., Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative

710            genomics. Genome Biol. 20:238.

711    Eriksson T., Lundberg M., Töpel M., Östensson P., Smedmark J.E.E. 2015. Sibbaldia: a

712            molecular phylogenetic study of a remarkably polyphyletic genus in Rosaceae. Plant

713            Syst. Evol. 301:171–184.

714    Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic

715            loci. Bioinforma. Oxf. Engl. 32:786–788.

716    Fernández R., Gabaldon T., Dessimoz C. 2020. Orthology: Definitions, Prediction, and Impact

717            on Species Phylogeny Inference. In: Scornavacca C., Delsuc F., Galtier N., editors.

718            Phylogenetics in the Genomic Era. No commercial publisher | Authors open access

719            book. p. 2.4:1--2.4:14.

720    Fitch W.M. 1970. Distinguishing Homologous from Analogous Proteins. Syst. Biol. 19:99–113.

721    Forrest L.L., Hart M.L., Hughes M., Wilson H.P., Chung K.-F., Tseng Y.-H., Kidner C.A. 2019.

722            The Limits of Hyb-Seq for Herbarium Specimens: Impact of Preservation Techniques.

723            Front. Ecol. Evol. 7:439.

724    García N., Folk R.A., Meerow A.W., Chamala S., Gitzendanner M.A., Oliveira R.S. de, Soltis

725            D.E., Soltis P.S. 2017. Deep reticulation and incomplete lineage sorting obscure the

726            diploid phylogeny of rain-lilies and allies (Amaryllidaceae tribe Hippeastreae). Mol.

727            Phylogenet. Evol. 111:231–247.

Gardner E.M., Johnson M.G., Pereira J.T., Ahmad Puad A.S., Arifiani D., Sahromi, Wickett

N.J., Zerega N.J.C. 2020. Paralogs and off-target sequences improve phylogenetic

resolution in a densely-sampled study of the breadfruit genus (Artocarpus, Moraceae).

bioRxiv.:854232.

Gardner E.M., Johnson M.G., Ragone D., Wickett N.J., Zerega N.J.C. 2016. Low-coverage,

whole-genome sequencing of Artocarpus camansi (Moraceae) for phylogenetic marker

development and gene discovery. Appl. Plant Sci. 4:1600017.

Gehrke B., Bräuchler C., Romoleroux K., Lundberg M., Heubl G., Eriksson T. 2008. Molecular

phylogenetics of Alchemilla, Aphanes and Lachemilla (Rosaceae) inferred  from plastid

and nuclear intron and spacer DNA sequences, with comments on generic classification.

Mol. Phylogenet. Evol. 47:1030–1044.

Gehrke B., Kandziora M., Pirie M.D. 2016. The evolution of dwarf shrubs in alpine

environments: a case study of Alchemilla in Africa. Ann. Bot. 117:121–131.

Glover N., Dessimoz C., Ebersberger I., Forslund S.K., Gabaldón T., Huerta-Cepas J., Martin

M.-J., Muffato M., Patricio M., Pereira C., da Silva A.S., Wang Y., Sonnhammer E.,

Thomas P.D. 2019. Advances and Applications in the Quest for Orthologs. Mol. Biol.

Evol. 36:2157–2164.

Gonçalves D.J.P., Simpson B.B., Ortiz E.M., Shimizu G.H., Jansen R.K. 2019. Incongruence

between gene trees and species trees and phylogenetic signal variation in plastid genes.

Mol. Phylogenet. Evol. 138:219–232.

Green R.E., Krause J., Briggs A.W., Maricic T., Stenzel U., Kircher M., Patterson N., Li H., Zhai

W., Fritz M.H.Y., Hansen N.F., Durand E.Y., Malaspinas A.S., Jensen J.D., Marques-

Bonet T., Alkan C., Prufer K., Meyer M., Burbano H.A., Good J.M., Schultz R., Aximu-

Petri A., Butthof A., Hober B., Hoffner B., Siegemund M., Weihmann A., Nusbaum C.,

Lander E.S., Russ C., Novod N., Affourtit J., Egholm M., Verna C., Rudan P., Brajkovic

D., Kucan Z., Gusic I., Doronichev V.B., Golovanova L.V., Lalueza-Fox C., de la

Rasilla M., Fortea J., Rosas A., Schmitz R.W., Johnson P.L.F., Eichler E.E., Falush D.,

Birney E., Mullikin J.C., Slatkin M., Nielsen R., Kelso J., Lachmann M., Reich D.,

Paabo S. 2010. A Draft Sequence of the Neandertal Genome. Science. 328:710–722.

Hejase H.A., Liu K.J. 2016. A scalability study of phylogenetic network inference methods using

empirical datasets and simulations involving a single reticulation. BMC Bioinformatics.

17:422.

Huang C.-H., Zhang C., Liu M., Hu Y., Gao T., Qi J., Ma H. 2016. Multiple Polyploidization

Events across Asteraceae with Two Nested Events in the Early History Revealed by

Nuclear Phylogenomics. Mol. Biol. Evol. 33:2820–2835.

Jiao Y., Leebens-Mack J., Ayyampalayam S., Bowers J.E., McKain M.R., McNeal J., Rolf M.,

Ruzicka D.R., Wafula E., Wickett N.J., Wu X., Zhang Y., Wang J., Zhang Y., Carpenter

E.J., Deyholos M.K., Kutchan T.M., Chanderbali A.S., Soltis P.S., Stevenson D.W.,

McCombie R., Pires J., Wong G., Soltis D.E., dePamphilis C.W. 2012. A genome

triplication associated with early diversification of the core eudicots. Genome Biol.

13:R3.

Jiao Y., Wickett N.J., Ayyampalayam S., Chanderbali A.S., Landherr L., Ralph P.E., Tomsho

L.P., Hu Y., Liang H., Soltis P.S., Soltis D.E., Clifton S.W., Schlarbaum S.E., Schuster

S.C., Ma H., Leebens-Mack J., dePamphilis C.W. 2011. Ancestral polyploidy in seed

plants and angiosperms. Nature. 473:97–100.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett

N.J. 2016. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics from

High-Throughput Sequencing Reads Using Target Enrichment. Appl. Plant Sci.

4:1600016.

Jones K.E., Fér T., Schmickl R.E., Dikow R.B., Funk V.A., Herrando-Moraira S., Johnston P.R.,

Kilian N., Siniscalchi C.M., Susanna A., Slovák M., Thapa R., Watson L.E., Mandel

J.R. 2019. An empirical assessment of a single family-wide hybrid capture locus set at

multiple  evolutionary timescales in Asteraceae. Appl. Plant Sci. 7:e11295.

Kamneva O.K., Rosenberg N.A. 2017. Simulation-Based Evaluation of Hybridization Network

Reconstruction Methods in the Presence of Incomplete Lineage Sorting. Evol.

Bioinforma. 13:117693431769193.

Kamneva O.K., Syring J., Liston A., Rosenberg N.A. 2017. Evaluating allopolyploid origins in

strawberries (Fragaria) using haplotypes generated from target capture sequencing.

BMC Evol. Biol. 17:401.

Karimi N., Grover C.E., Gallagher J.P., Wendel J.F., Ané C., Baum D.A. 2020. Reticulate

Evolution Helps Explain Apparent Homoplasy in Floral Biology and Pollination in

Baobabs (Adansonia; Bombacoideae; Malvaceae). Syst. Biol. 69:462–478.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780.

Kearse M., Moir R., Wilson A., Stones-Havas S., Cheung M., Sturrock S., Buxton S., Cooper A.,

Markowitz S., Duran C., Thierer T., Ashton B., Meintjes P., Drummond A. 2012.

Geneious Basic: An integrated and extendable desktop software platform for the

organization and analysis of sequence data. Bioinformatics. 28:1647–1649.

Kocot K.M., Citarella M.R., Moroz L.L., Halanych K.M. 2013. PhyloTreePruner: A

Phylogenetic Tree-Based Approach for Selection of Orthologous Sequences for

Phylogenomics. Evol. Bioinforma. Online. 9:429–435.

Koenen E.J.M., Ojeda D.I., Bakker F.T., Wieringa J.J., Kidner C., Hardy O.J., Pennington R.T.,

Herendeen P.S., Bruneau A., Hughes C.E. 2020. The Origin of the Legumes is a

Complex Paleopolyploid Phylogenomic Tangle closely associated with the Cretaceous-

Paleogene (K-Pg) Mass Extinction Event. Syst. Biol. syaa041.

Larridon I., Villaverde T., Zuntini A.R., Pokorny L., Brewer G.E., Epitawalage N., Fairlie I.,

Hahn M., Kim J., Maguilla E., Maurin O., Xanthos M., Hipp A.L., Forest F., Baker W.J.

2020. Tackling Rapid Radiations With Targeted Sequencing. Front. Plant Sci. 10:1655.

Leebens-Mack J.H., Barker M.S., Carpenter E.J., Deyholos M.K., Gitzendanner M.A., Graham

S.W., Grosse I., Li Z., Melkonian M., Mirarab S., Porsch M., Quint M., Rensing S.A.,

Soltis D.E., Soltis P.S., Stevenson D.W., Ullrich K.K., Wickett N.J., DeGironimo L.,

Edger P.P., Jordon-Thaden I.E., Joya S., Liu T., Melkonian B., Miles N.W., Pokorny L.,

Quigley C., Thomas P., Villarreal J.C., Augustin M.M., Barrett M.D., Baucom R.S.,

Beerling D.J., Benstein R.M., Biffin E., Brockington S.F., Burge D.O., Burris J.N.,

Burris K.P., Burtet-Sarramegna V., Caicedo A.L., Cannon S.B., Çebi Z., Chang Y.,

Chater C., Cheeseman J.M., Chen T., Clarke N.D., Clayton H., Covshoff S., Crandall-

Stotler B.J., Cross H., dePamphilis C.W., Der J.P., Determann R., Dickson R.C., Di

Stilio V.S., Ellis S., Fast E., Feja N., Field K.J., Filatov D.A., Finnegan P.M., Floyd

S.K., Fogliani B., García N., Gâteblé G., Godden G.T., Goh F. (Qi Y., Greiner S.,

Harkess A., Heaney J.M., Helliwell K.E., Heyduk K., Hibberd J.M., Hodel R.G.J.,

Hollingsworth P.M., Johnson M.T.J., Jost R., Joyce B., Kapralov M.V., Kazamia E.,

Kellogg E.A., Koch M.A., Von Konrat M., Könyves K., Kutchan T.M., Lam V., Larsson

A., Leitch A.R., Lentz R., Li F.-W., Lowe A.J., Ludwig M., Manos P.S., Mavrodiev E.,

McCormick M.K., McKain M., McLellan T., McNeal J.R., Miller R.E., Nelson M.N.,

Peng Y., Ralph P., Real D., Riggins C.W., Ruhsam M., Sage R.F., Sakai A.K.,

Scascitella M., Schilling E.E., Schlösser E.-M., Sederoff H., Servick S., Sessa E.B.,

Shaw A.J., Shaw S.W., Sigel E.M., Skema C., Smith A.G., Smithson A., Stewart C.N.,

Stinchcombe J.R., Szövényi P., Tate J.A., Tiebel H., Trapnell D., Villegente M., Wang

C.-N., Weller S.G., Wenzel M., Weststrand S., Westwood J.H., Whigham D.F., Wu S.,

Wulff A.S., Yang Y., Zhu D., Zhuang C., Zuidof J., Chase M.W., Pires J.C., Rothfels

C.J., Yu J., Chen C., Chen L., Cheng S., Li J., Li R., Li X., Lu H., Ou Y., Sun X., Tan

X., Tang J., Tian Z., Wang F., Wang J., Wei X., Xu X., Yan Z., Yang F., Zhong X.,

Zhou F., Zhu Y., Zhang Y., Ayyampalayam S., Barkman T.J., Nguyen N., Matasci N.,

Nelson D.R., Sayyari E., Wafula E.K., Walls R.L., Warnow T., An H., Arrigo N.,

Baniaga A.E., Galuska S., Jorgensen S.A., Kidder T.I., Kong H., Lu-Irving P., Marx

H.E., Qi X., Reardon C.R., Sutherland B.L., Tiley G.P., Welles S.R., Yu R., Zhan S.,

Gramzow L., Theißen G., Wong G.K.-S., One Thousand Plant Transcriptomes Initiative.

2019. One thousand plant transcriptomes and the phylogenomics of green plants. Nature.

574:679–685.

Li L. 2003. OrthoMCL: Identification of Ortholog Groups for Eukaryotic Genomes. Genome

Res. 13:2178–2189.

Li Z., Baniaga A.E., Sessa E.B., Scascitelli M., Graham S.W., Rieseberg L.H., Barker M.S.

2015. Early genome duplications in conifers and other seed plants. Sci. Adv.

1:e1501084.

Liu Y., Johnson M.G., Cox C.J., Medina R., Devos N., Vanderpoorten A., Hedenäs L., Bell N.E.,

Shevock J.R., Aguero B., Quandt D., Wickett N.J., Shaw A.J., Goffinet B. 2019.

Resolution of the ordinal phylogeny of mosses using targeted exons from organellar and

nuclear genomes. Nat. Commun. 10:1485.

Lundberg M., Töpel M., Eriksen B., Nylander J.A.A., Eriksson T. 2009. Allopolyploidy in

Fragariinae (Rosaceae): Comparing four DNA sequence regions, with comments on

classification. Mol. Phylogenet. Evol. 51:269–280.

Lynch M., Conery J.S. 2000. The Evolutionary Fate and Consequences of Duplicate Genes.

Science. 290:1151–1155.

Mai U., Mirarab S. 2018. TreeShrink: fast and accurate detection of outlier long branches in

collections of phylogenetic trees. BMC Genomics. 19:272.

Mandel J.R., Dikow R.B., Funk V.A., Masalia R.R., Staton S.E., Kozik A., Michelmore R.W.,

Rieseberg L.H., Burke J.M. 2014. A Target Enrichment Method for Gathering

Phylogenetic Information from Hundreds of Loci: An Example from the Compositae.

Appl. Plant Sci. 2:1300085.

McKain M.R., Estep M.C., Pasquet R., Layton D.J., Vela Díaz D.M., Zhong J., Hodge J.G.,

Malcomber S.T., Chipabika G., Pallangyo B., Kellogg E.A. 2018. Ancestry of the two

subgenomes of maize. bioRxiv.:352351.

McKain M.R., Tang H., McNeal J.R., Ayyampalayam S., Davis J.I., dePamphilis C.W., Givnish

T.J., Pires J.C., Stevenson D.W., Leebens-Mack J.H. 2016. A Phylogenomic Assessment

of Ancient Polyploidy and Genome Evolution across the Poales. Genome Biol. Evol.

8:1150–1164.

McLachlan G., Peel D. 2000. Finite Mixture Models. New York: Wiley.

Molloy E.K., Warnow T. 2020. FastMulRFS: fast and accurate species tree estimation under

generic gene duplication and loss models. Bioinformatics. 36:i57–i65.

Montes J.R., Peláez P., Willyard A., Moreno-Letelier A., Piñero D., Gernandt D.S. 2019.

Phylogenetics of Pinus Subsection Cembroides Engelm. (Pinaceae) Inferred from Low-

Copy Nuclear Gene Sequences. Syst. Bot. 44:501–518.

Moore A.J., Vos J.M.D., Hancock L.P., Goolsby E., Edwards E.J. 2018. Targeted Enrichment of

Large Gene Families for Phylogenetic Inference: Phylogeny and Molecular Evolution of

Photosynthesis Genes in the Portullugo Clade (Caryophyllales). Syst. Biol. 67:367–383.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington S.F.,

Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2020. Disentangling Sources of Gene

Tree Discordance in Phylogenomic Datasets: Testing Ancient Hybridizations in

Amaranthaceae s.l. Syst. Biol. syaa066

Morales-Briones D.F., Romoleroux K., Kolář F., Tank D.C. 2018a. Phylogeny and Evolution of

the Neotropical Radiation of *Lachemilla* (Rosaceae): Uncovering a History of Reticulate

Evolution and Implications for Infrageneric Classification. Syst. Bot. 43:17–34.

Morales-Briones D.F., Liston A., Tank D.C. 2018b. Phylogenomic analyses reveal a deep history

of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae). New

Phytol. 218:1668–1684.

Morales-Briones D.F., Tank D.C. 2019. Extensive allopolyploidy in the neotropical genus

*Lachemilla* (Rosaceae) revealed by. Am. J. Bot. 106:415–437.

Nicholls J.A., Pennington R.T., Koenen E.J.M., Hughes C.E., Hearn J., Bunnefeld L., Dexter

K.G., Stone G.N., Kidner C.A. 2015. Using targeted enrichment of nuclear genes to

increase phylogenetic resolution in the neotropical rain forest genus Inga (Leguminosae:

Mimosoideae). Front. Plant Sci. 6:710.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling

distinguishes lack of support from conflicting support in the green plant tree of life. Am.

J. Bot. 105:385–403.

Perry L.M. 1929. A Tentative Revision of Alchemilla § Lachemilla. Contrib. Gray Herb. Harv.

Univ.:1–57.

Ranwez V., Douzery E.J.P., Cambon C., Chantret N., Delsuc F. 2018. MACSE v2: Toolkit for

the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. Mol.

Biol. Evol. 35:2582–2584.

Salichos L., Stamatakis A., Rokas A. 2014. Novel Information Theory-Based Measures for

Quantifying Incongruence among Phylogenetic Trees. Mol. Biol. Evol. 31:1261–1271.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Shulaev V., Sargent D.J., Crowhurst R.N., Mockler T.C., Folkerts O., Delcher A.L., Jaiswal P.,

Mockaitis K., Liston A., Mane S.P., Burns P., Davis T.M., Slovin J.P., Bassil N.,

Hellens R.P., Evans C., Harkins T., Kodira C., Desany B., Crasta O.R., Jensen R.V.,

Allan A.C., Michael T.P., Setubal J.C., Celton J.-M., Rees D.J.G., Williams K.P., Holt

S.H., Rojas J.J.R., Chatterjee M., Liu B., Silva H., Meisel L., Adato A., Filichkin S.A.,

Troggio M., Viola R., Ashman T.-L., Wang H., Dharmawardhana P., Elser J., Raja R.,

Priest H.D., Bryant D.W., Fox S.E., Givan S.A., Wilhelm L.J., Naithani S., Christoffels

A., Salama D.Y., Carter J., Girona E.L., Zdepski A., Wang W., Kerstetter R.A., Schwab

W., Korban S.S., Davik J., Monfort A., Denoyes-Rothan B., Arus P., Mittler R., Flinn

B., Aharoni A., Bennetzen J.L., Salzberg S.L., Dickerman A.W., Velasco R.,

Borodovsky M., Veilleux R.E., Folta K.M. 2011. The genome of woodland strawberry

(Fragaria vesca). Nat. Genet. 43:109–116.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals

conflict, concordance, and gene duplications with examples from animals and plants.

BMC Evol. Biol. 15:150.

Soják J. 2008. Notes on Potentilla XXI. A new division of the tribe Potentilleae (Rosaceae) and

notes on generic delimitations. Bot. Jahrb. Für Syst. Pflanzengesch. Pflanzengeogr.

127:349–358.

Solís-Lemus C., Ané C. 2016. Inferring Phylogenetic Networks with Maximum

Pseudolikelihood under Incomplete Lineage Sorting. PLoS Genet. 12:e1005896–21.

Stamatakis A. 2014. RAxML version 8 - a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 30:1312–1313.

Straub S.C., Fishbein M., Livshultz T., Foster Z., Parks M., Weitemier K., Cronn R.C., Liston A.

2011. Building a model: developing genomic resources for common milkweed

(Asclepias syriaca) with low coverage genome sequencing. BMC Genomics. 12:211.

Stubbs R.L., Folk R.A., Xiang C.-L., Chen S., Soltis D.E., Cellinese N. 2020. A Phylogenomic

Perspective on Evolution and Discordance in the Alpine-Arctic Plant Clade Micranthes

(Saxifragaceae). Front. Plant Sci. 10:1773.

Thomas G.W.C., Ather S.H., Hahn M.W. 2017. Gene-Tree Reconciliation with MUL-Trees to

Resolve Polyploidy Events. Syst. Biol. 66:1007–1018.

Villaverde T., Pokorny L., Olsson S., Rincón-Barrado M., Johnson M.G., Gardner E.M., Wickett

N.J., Molero J., Riina R., Sanmartín I. 2018. Bridging the micro- and macroevolutionary

levels in phylogenomics: Hyb-Seq solves relationships from populations to species and

above. New Phytol. 220:636–650.

Viruel J., Conejero M., Hidalgo O., Pokorny L., Powell R.F., Forest F., Kantar M.B., Soto

Gomez M., Graham S.W., Gravendeel B., Wilkin P., Leitch I.J. 2019. A Target Capture-

Based Method to Estimate Ploidy From Herbarium Specimens. Front. Plant Sci. 10:937.

Walker J.F., Walker-Hale N., Vargas O.M., Larson D.A., Stull G.W. 2019. Characterizing gene

tree conflict in plastome-inferred phylogenies. PeerJ. 7:e7747.

Weitemier K., Straub S.C.K., Cronn R.C., Fishbein M., Schmickl R., McDonnell A., Liston A.

2014. Hyb-Seq: Combining Target Enrichment and Genome Skimming for Plant

Phylogenomics. Appl. Plant Sci. 2:1400042.

Wen D., Yu Y., Zhu J., Nakhleh L. 2018. Inferring Phylogenetic Networks Using PhyloNet.

Syst. Biol. 67:735–740.

Xiang Y., Huang C.-H., Hu Y., Wen J., Li S., Yi T., Chen H., Xiang J., Ma H. 2017. Evolution

of Rosaceae Fruit Types Based on Nuclear Phylogeny in the Context of Geological

Times and Genome Duplication. Mol. Biol. Evol. 34:262–281.

Yang Y., Moore M.J., Brockington S.F., Mikenas J., Olivieri J., Walker J.F., Smith S.A. 2018.

Improved transcriptome sampling pinpoints 26 ancient and more recent polyploidy

events in Caryophyllales, including two allopolyploidy events. New Phytol. 217:855–

870.

Yang Y., Moore M.J., Brockington S.F., Soltis D.E., Wong G.K.-S., Carpenter E.J., Zhang Y.,

Chen L., Yan Z., Xie Y., Sage R.F., Covshoff S., Hibberd J.M., Nelson M.N., Smith

S.A. 2015. Dissecting Molecular Evolution in the Highly Diverse Plant Clade

Caryophyllales Using Transcriptome Sequencing. Mol. Biol. Evol. 32:2001–2014.

Yang Y., Smith S.A. 2014. Orthology Inference in Nonmodel Organisms Using Transcriptomes

and Low-Coverage Genomes: Improving Accuracy and Matrix Occupancy for

Phylogenomics. Mol. Biol. Evol. 31:3081–3092.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

Zhang C., Scornavacca C., Molloy E.K., Mirarab S. 2019. ASTRAL-Pro: quartet-based species

tree inference despite paralogy. bioRxiv.:2019.12.12.874727.

Zhang R., Wang Y.-H., Jin J.-J., Stull G.W., Bruneau A., Cardoso D., De Queiroz L.P., Moore

M.J., Zhang S.-D., Chen S.-Y., Wang J., Li D.-Z., Yi T.-S. 2020. Exploration of Plastid

Phylogenomic Conflict Yields New Insights into the Deep Relationships of

Leguminosae. Syst. Biol. 69:613–622.

Zhbannikov I.Y., Hunter S.S., Foster J.A., Settles M.L. 2017. SeqyClean: A Pipeline for High-

throughput Sequence Data Preprocessing. Proc. 8th ACM Int. Conf. Bioinforma.

Comput. Biol. Health Inform.:407–416.

970