Analysis of paralogs in target enrichment data pinpoints multiple ancient polyploidy events

in *Alchemilla* s.l. (Rosaceae)

Diego F. Morales-Briones[1,2,*], Berit Gehrke[3], Chien-Hsun Huang[4], Aaron Liston[5], Hong Ma[6],

Hannah E. Marx[7,8], David C. Tank[2], Ya Yang[1]

[1]Department of Plant and Microbial Biology, University of Minnesota-Twin Cities, 1445 Gortner

Avenue, St. Paul, MN 55108, USA

[2]Department of Biological Sciences and Institute for Bioinformatics and Evolutionary Studies,

University of Idaho, 875 Perimeter Drive MS 3051, Moscow, ID 83844, USA

[3]University Gardens, University Museum, University of Bergen, Mildeveien 240, 5259

Hjellestad, Norway

[4]State Key Laboratory of Genetic Engineering and Collaborative Innovation Center of Genetics

and Development, Ministry of Education Key Laboratory of Biodiversity and Ecological

Engineering, Institute of Plant Biology, Center of Evolutionary Biology, School of Life Sciences,

Fudan University, Shanghai 200433, China

[5]Department of Botany and Plant Pathology, Oregon State University, 2082 Cordley Hall,

Corvallis, OR 97331, USA

[6]Department of Biology, the Huck Institute of the Life Sciences, the Pennsylvania State

University, 510D Mueller Laboratory, University Park, PA 16802 USA

[7]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI

48109-1048, USA

[8]Museum of Southwestern Biology and Department of Biology, University of New Mexico,

Albuquerque, NM 87131, USA

* Correspondence to be sent to: Diego F. Morales-Briones. Department of Plant and Microbial

Biology, University of Minnesota, 1445 Gortner Avenue, St. Paul, MN 55108, USA, Email:

dfmoralesb@gmail.com

Abstract. —Target enrichment is becoming increasingly popular for phylogenomic studies. Although baits for enrichment are typically designed to target single-copy genes, paralogs are often recovered with increased sequencing depth, sometimes from a significant proportion of loci, especially in groups experiencing whole-genome duplication (WGD) events. Common approaches for processing paralogs in target enrichment datasets include random selection, manual pruning, and mainly, the removal of entire genes that show any evidence of paralogy. These approaches are prone to errors in orthology inference or removing large numbers of genes. By removing entire genes, valuable information that could be used to detect and place WGD events is discarded. Here we use an automated approach for orthology inference in a target enrichment dataset of 68 species of *Alchemilla* s.l. (Rosaceae), a widely distributed clade of plants primarily from temperate climate regions. Previous molecular phylogenetic studies and chromosome numbers both suggested ancient WGDs in the group. However, both the phylogenetic location and putative parental lineages of these WGD events remain unknown. By taking paralogs into consideration, we identified four nodes in the backbone of *Alchemilla* s.l. with an elevated proportion of gene duplication. Furthermore, using a gene-tree reconciliation approach we established the autopolyploid origin of the entire *Alchemilla* s.l. and the nested allopolyploid origin of four major clades within the group. Here we showed the utility of automated tree-based orthology inference methods, previously designed for genomic or transcriptomic datasets, to study complex scenarios of polyploidy and reticulate evolution from target enrichment datasets.

Keywords: *Alchemilla*; allopolyploidy; autopolyploidy; gene tree discordance; orthology inference; paralogs; Rosaceae; target enrichment; whole genome duplication.

Polyploidy, or whole genome duplication (WGD), is prevalent throughout the evolutionary

history of plants (Cui et al. 2006; Jiao et al. 2011; Jiao et al. 2012; Leebens-Mack et al. 2019). As

a result, plant genomes often contain large numbers of paralogous genes from recurrent gene and

genome duplication events (Lynch and Conery 2000; Panchy et al. 2016). Paralogs are defined as

homologous genes that share a common ancestor as the product of gene duplication (Fitch 1970),

either from small scale duplications or WGD. One special case of WGD is allopolyploidy, where

genome doubling is accompanied by hybridization between two different species. The duplicated

genes in allopolyploids are not paralogs in the traditional sense and are referred to as

homoeologs, which are expected to be sister to the orthologous in the parental taxa, rather than to

each other (Smedmark et al., 2003). For practical purposes, however, we refer to the product of

any kind of duplications found in gene trees hereafter as paralog, as homoeologs are

indistinguishable from paralogs until diagnosed as resulting from allopolyploidy. With very few

nuclear genes being truly single- or low-copy, careful evaluation of orthology is critical for

phylogenetic analyses (Fitch 1970). Orthology inference has received much attention in the

phylogenomic era with multiple pipelines available for this task (e.g., Li et al. 2003; Dunn et al.

2013; Kocot et al. 2013; Yang and Smith 2014; Emms and Kelly 2019, also see Glover et. al

2019 and Fernández et al. 2020 for recent reviews). But these approaches have been mainly

applied to genomic or transcriptomic data sets. So far, few studies have employed automated,

phylogeny-aware orthology inference in target enrichment datasets. The most common approach

for dealing with paralogy in target enrichment datasets is removing entire genes that show any

evidence of potential paralogy (e.g., Nicholls et al. 2015; Jones et al. 2019; Andermann et al.

2020; but see Moore et al. 2018). Removal of entire genes might seem appropriate in target

enrichment datasets in which only a small number of genes show evidence of paralogy (e.g.,

Larridon et al. 2020), but in some datasets this could result in a significant reduction of the

number of loci (e.g., Montes et al. 2019). More importantly, dealing with paralogy only by

removal of entire genes assumes that target enrichment assembly pipelines (e.g., Faircloth 2016;

Johnson et al. 2016; Andermann et al. 2018), have flagged all genes with paralogs. It also

assumes that if no sequence in a gene is flagged, all sequences are all single-copy and

orthologous. On the other hand, this approach also removes genes that show allelic variation

instead of paralogs. Given the prevalence of WGD and reticulations these assumptions can lead

to errors in orthology inference. As paralogous genes are prevalent in plants, more appropriate

orthology inference methods need to be applied in target enrichment data. The same automated

approaches used for genome and transcriptome datasets can be applied for target enrichment, as

these are tree-based and agnostic to the data source for tree inference.

The ability to explicitly process paralogs opens the door for using target enrichment data

for inferring gene duplication events and pinpointing the phylogenetic locations of putative

WGDs. In the past, the phylogenetic placement of WGD events have most often been carried out

using genome and transcriptome sequencing data (e.g., Li et al. 2015; Huang et al. 2016; McKain

et al. 2016; Yang et al. 2018) using either the synonymous distance between paralog gene pairs

(Ks; Lynch and Conery 2000) or tree-based reconciliation methods (e.g., Jiao et al. 2011; Li et

al. 2015; Yang et al. 2015; Huang et al. 2016; Xiang et al. 2017; Leebens-Mack et al. 2019).

Similar to orthology inference, tree-based methods used to investigate WGDs in genome and

transcriptome datasets should be useful in target enrichment data. Target enrichment methods

(e.g., Mandel et al. 2014; Weitemier et al. 2014; Buddenhagen et al. 2016) have been widely

adopted to collect hundreds to over a thousand nuclear loci for plant systematics, allowing

studies at different evolutionary scales (e.g., Villaverde et al. 2018), and the use of museum-

preserved collections (e.g., Forrest et al. 2019). This creates new opportunities to adopt tree-

based reconciliation methods to explore WGD patterns in groups for which genomic and

transcriptomic resources are not available or feasible.

With at least 350 (–1,100) species worldwide, *Alchemilla* in the broad sense has been a

challenging group to study due to the presence of reticulate evolution, polyploidy, and apomixis .

Based on previous phylogenetic analyses, *Alchemilla* s.l. contains four clades: Afromilla,

Aphanes, Eualchemilla, and *Lachemilla* (Table S1). Together they form a well-supported clade

nested in the subtribe Fragariinae, which also includes the cultivated strawberries (Gehrke et al.

2008). Unlike the more commonly recognized members of the rose family (Rosaceae),

Alchemilla s.l. is characterized by small flowers with no petals, and a reduced number (1–4[–5])

of stamens that have anthers with one elliptic theca on the ventral side of the connective that

opens by one transverse split (Perry 1929; Soják 2008). Gehrke et al. (2008) presented the first

phylogeny of *Alchemilla* s.l. and established the paraphyly of traditional *Alchemilla* s.s. as

consisted of a primarily African clade, Afromilla, and a Eurasian clade, Eualchemilla. Gehrke et

al. (2008) also suggested treating Afromilla and Eualchemilla, along with *Aphanes* and

Lachemilla as a single genus based on nomenclatural stability and the lack of morphological

characters to distinguish between Afromilla and Eualchemilla. The four clades within *Alchemilla*

s.l. are mainly defined by geographic distribution, as well as the number and insertion of the

stamens on the disk lining the hypanthium (Table S1). Phylogenetic analyses using at least one

nuclear and one chloroplast marker (Gehrke et al. 2008; 2016) found significant cytonuclear

discordance regarding the relationships among the four major clades. Similar patterns, often

attributed to hybridization and allopolyploidy, have been detected in other genera of Fragariinae

(Lundberg 2009; Eriksson et al. 2015; Gehrke et al. 2016, Kamneva et al. 2017; Morales-Briones

et al. 2018a), leaving the phylogenetic relationships of *Alchemilla* s.l. to the rest of Fragariinae

unresolved. Unlike most members of Fragariinae that have predominantly diploid species,

Alchemilla s.l. is known for high rates of polyploidy. The base chromosome number of

Alchemilla s.l. is eight ($x = 8$), which differs from all other members in Fragariinae that have a

base of number of seven ($x = 7$; Dickinson et al. 2007; Lundberg et al., 2009). Ploidy levels have

been well documented in Eualchemilla that shows only polyploid species ($2n = 64$ to 220–224;

octoploid to 28-ploid; e.g., Turesson 1943; Izmailow 1981; Walters and Bozman, 1967;

Hayirhoğlu-Ayaz et al. 2006). *Aphanes* has mainly diploid species ($2n = 16$), with the exception

of *Aphanes arvensis* that is an hexaploid ($2n = 48$; Montgomery et al. 1997). *Lachemilla* has

mostly polyploid members ($2n = 24$ to 96; triploid to 12-ploid) with a single species reported to

have diploid ($2n = 16$) and triploid ($2n = 24$) populations (Morales-Briones et al. 2018a). Lastly,

little is known about ploidy levels in Afromilla, but so far, the two species reported were both

polyploids ($2n = 64$ to 80; octoploid and decaploid; Hjelmqvist 1956; Morton 1993). A recent

phylogenomic analysis focused on *Lachemilla* using target enrichment and 32 species of the

group detected a high frequency of paralogs shared with Eualchemilla and Afromilla (Morales-

Briones et al. 2018b). This paralog frequency suggested a possible ancient WGD event; however,

the sampling was limited to one species each of Eualchemilla and Afromilla, and the location

and mode of this putative WGD remained uncertain.

In this study we sampled 68 species across the major clades of *Alchemilla* s.l., and

included 11 additional closely related species in Fragariinae, which allowed us to 1) test for

polyploid events in the origin of *Alchemilla* s.l., and 2) explore the reticulate evolution among

major clades of *Alchemilla* s.l. using a target enrichment dataset. Given the prevalence of

polyploidy and reticulate history within *Alchemilla* s.l., this is an excellent group to explore the

utility of tree-based methods for (1) processing paralogs, and (2) detecting and placing WGDs

using target enrichment datasets.

147   **MATERIALS AND METHODS**

148   *Taxon sampling and data collection*

We sampled 68 species representing the four major clades of *Alchemilla* s.l. (sensu Gehrke et al.

2008), and 11 species to represent all other genera in Fragariinae (except *Chamaecallis*; sensu

Dobeš et al. 2015; Morales-Briones and Tank 2019). Additionally, we sampled one species each

of *Potentilla*, *Sanguisorba*, and *Rosa* as outgroups. Voucher information is provided in Table S2.

We used a Hyb-Seq approach (Weitemier et al. 2014), that combines target enrichment and

genome skimming, to capture nuclear exon sequences and off-target cpDNA. We used baits

designed for *Fragaria vesca* (strawberry, also a member of Fragariinae) to target 1,419 exons in

257 genes (Kamneva et al. 2017). These genes were identified as single-copy orthologs among

the apple (*Malus domestica*), peach (*Prunus persica*), and strawberry genomes based on

reciprocal nucleotide similarity comparisons. The 257 genes resulted from first retaining only

genes >960 bp long and with >85% similarity in pairwise comparisons among the three

genomes. The remaining genes were further filtered by removing exons <80 bp long, with GC

content <30% or >70%, and with >90% sequence similarity to annotated repetitive DNA in the

genome, followed by removing exons with any paralogs with >90% sequence similarity in the

same genome (Kamneva et al. 2017).

164   Of the 82 total species, only sequences for *Fragaria vesca*, were from a reference

genome (Shulaev et. al 2010). Twenty-two were from a previously published Hyb-Seq dataset

using the same bait set as this study (Morales-Briones et al. 2018b; Table S2), including 19

species of *Lachemilla* that did not show evidence of hybridization within *Lachemilla,* and one

species each of Eualchemilla, Afromilla, and *Aphanes*. Newly generated sequence data for 55

species (Table S2) were collected as follows. Total genomic DNA was isolated from silica-dried

or herbarium material with a modified CTAB method (Doyle and Doyle 1987). Probe synthesis,

library preparation, capture enrichment, and high-throughput sequencing (HiSeq2000 instrument,

2 × 101 bp) were carried out at Rapid Genomics LLC (Gainesville, FL, USA). Data for the

remaining four species, *Drymocallis glandulosa*, *Potentilla indica*, *Rosa woodsii*, and

Sanguisorba menziesii were collected as described in Weitemier et al. (2014).

176   *Read processing and assembly*

We removed sequencing adaptors and trimmed low-quality bases (Phred scores < 20) from raw

reads with SeqyClean v.1.10.07 (Zhbannikov et al. 2017) using default settings. Plastomes were

assembled using Alignreads v.2.5.2 (Straub et al. 2011) and 12 closely related plastome

references (with one Inverted Repeat removed; Table S3). Plastome assemblies were annotated

using *Fragaria vesca* as a reference in Geneious v.11.1.5 (Kearse et al. 2012). Assembly of

nuclear loci was carried out with HybPiper v.1.3.1 (Johnson et al. 2016) using exons of *F. vesca*

as references. Given the large number of paralogs detected in *Lachemilla,* Eualchemilla, and

Afromilla, multi-exon gene assemblies resulted in chimeric sequences of exons from distinct

paralogs (Morales-Briones et al. 2018b). To avoid chimeric sequences that can affect orthology

inference and phylogenetic analyses, assemblies were performed on each exon separately. Only

exons with a reference length of ≥ 150 bp were assembled (939 exons from 257 genes). Paralog

detection was carried out for all exons with the 'paralog_investigator' option in HybPiper. This

option flags loci with potential paralogs when multiple contigs cover at least 85% of the

reference sequence length. Exon assemblies that included flagged paralogs were extracted using

the 'paralog_retriever' command of HybPiper and used for orthology inference.

193   *Orthology inference for nuclear exons*

To infer orthologs for phylogenetic analyses, all exons were processed as follows (Fig. 1a).

Individual exons were aligned using MACSE v.2.03 (Ranwez et al. 2018) with default

parameters. Codons with frameshifts (labeled with '!' by MACSE) were replaced with gaps and

aligned columns with more than 90% missing data were removed using Phyx (Brown et al.

2017). Initial homolog trees were built using RAxML v.8.2.11 (Stamatakis 2014) with a

GTRCAT model and clade support assessed with 100 rapid bootstrap (BS) replicates. Clades and

paraphyletic grades that belonged to the same taxon were pruned by keeping only the tip with the

highest number of characters in the trimmed alignment following Yang and Smith (2014). To

obtain the final homolog trees, outlier tips with unusually long branches were detected and

removed by maximally reducing the tree diameter with TreeShrink v.1.3.2 (Mai and Mirarab

2018). Orthology inference was carried out using two outgroup-aware strategies from Yang and

Smith (2014). We set *Potentilla*, *Sanguisorba*, and *Rosa* as outgroups and all members of

Fragariinae as ingroups. First, we used the 'monophyletic outgroup' (MO) approach keeping

only ortholog groups with at least 25 ingroup taxa. The MO approach filters for homolog trees

with outgroup taxa being monophyletic and single-copy, and therefore filters for single- and low-

copy genes. The second approach used was the 'rooted ingroup' (RT), with at least 25 ingroup

taxa. The RT approach iteratively searches subtrees of ingroup taxa and cuts them out as rooted

trees. Both approaches root the gene tree by the outgroups, traverse the rooted tree from root to

tip, and remove the side with fewer taxa (MO) or keep both sides (RT) when gene duplication is

detected at any given node. In the case of MO, homolog trees with non-monophyletic outgroups

or duplicated taxa in the outgroups are discarded. If no taxon duplication is detected in a

homolog tree, the MO approach outputs a one-to-one ortholog. The RT approach maximizes the

number of orthologs compared to MO while not requiring monophyletic outgroups and allowing

for duplicated taxa in the outgroups but removes outgroups from all orthologs. To add outgroups

back to the RT orthologs for downstream analyses, we kept only RT orthologs from homologs

that had a MO ortholog (i.e., using only homolog trees with monophyletic and non-duplicated

outgroups for both MO and RT). Then we used the outgroups of the MO ortholog for all the RT

orthologs of the same homolog (Fig. 1b). Scripts for orthology inference can be found at

https://bitbucket.org/dfmoralesb/target_enrichment_orthology.

Figure 1. Paralog processing workflow and orthology inference methods used in *Alchemilla* s.l.
homolog trees. a) Flow chart of paralog processing and homolog tree inference. b) Only
homologs with outgroup present and monophyletic were used for orthology inference.
Monophyletic outgroups (MO) will prune single-copy genes keeping clades with at least a user-
defined minimum number of ingroup taxa. Rooted ingroups (RT) will keep all subtrees with at
least a user-defined minimum number of ingroups taxa. If the homolog trees can be pruned using
both MO and RT, then RT orthologs are added to the same root. Homologs that lack
monophyletic outgroups were excluded from further consideration.

Phylogenetic analyses

We used concatenation and coalescent-based methods to reconstruct the phylogeny of *Alchemilla*
s.l. Analyses were carried out in the two sets of final orthologs, MO and RT, separately. Each
ortholog was aligned using MACSE v.2.03 with default parameters. Codons with frameshifts

were replaced with gaps, aligned columns with more than 90% missing data were removed using

Phyx, and alignments with at least 150 characters and 25 taxa were retained. We first estimated a

maximum likelihood (ML) tree from the concatenated matrices with RAxML using a partition by

gene scheme with a GTRGAMMA model for each partition. Clade support was assessed with

100 rapid bootstrap (BS) replicates. To estimate a species tree that is statistically consistent with

the multi-species coalescent (MSC), we first inferred individual ML gene trees using RAxML

with a GTRGAMMA model, and 100 BS replicates to assess clade support. Individual gene trees

were then used to estimate a species tree using ASTRAL-III v.5.6.3 (Zhang et al. 2018) using

local posterior probabilities (LPP; Sayyari and Mirarab 2016) to assess clade support.

To evaluate nuclear gene tree discordance, we calculated the internode certainty all (ICA)

value to quantify the degree of conflict on each node of the map tree (e.g., species tree) given

individual gene trees (Salichos et al. 2014). Also, we calculated the number of conflicting and

concordant bipartitions on each node of the map tree. We calculated both the ICA scores and the

number of conflicting/concordant bipartitions with Phyparts (Smith et al. 2015) using individual

ortholog trees with BS support of at least 50% for the corresponding node. Additionally, to

distinguish strong conflict from weakly supported branches, we evaluated tree conflict and

branch support with Quartet Sampling (QS; Pease et al. 2018) using 1,000 replicates. Quartet

Sampling subsamples quartets from the input map tree (e.g., species tree) and concatenated

alignment to assess the confidence, consistency, and informativeness of each internal branch by

the relative frequency of the three possible quartet topologies at each node (Pease et al. 2018).

In addition to species tree construction using inferred orthologs, we used a recently

developed quartet-based species tree method (ASTRAL-Pro; Zhang et al. 2020a) to estimate the

phylogeny of *Alchemilla* s.l. ASTRAL-Pro directly uses multi-labeled gene trees while

accounting for gene duplications and losses to estimate a species tree that is statistically

consistent with the MSC and birth-death gene duplication and loss model. We used all 923 final

homolog trees as input for ASTRAL-Pro, ignoring trees with less than 20 taxa, and estimated

LPP to assess clade support. Additionally, we calculated ICA scores and the number of

conflicting/concordant bipartitions with Phyparts using homolog trees with BS support of at least

50% for the corresponding nodes.

268   For the plastome phylogenetic analyses, 74 partial plastome assemblies and eight

reference plastome sequences were included (Table S3). Contiguous plastome sequences were

aligned using the default settings in MAFFT v.7.307 (Katoh and Standley 2013) and aligned

columns with more than 70% missing data were removed with Phyx. We estimated an ML tree

of the plastome alignment with RAxML using a partition by coding (CDS) and noncoding

regions (introns and intergenic spacers) scheme, with a GTRGAMMA model for each partition

and clade support assessed with 100 rapid BS replicates and QS using 1,000 replicates, to detect

potential within-plastome conflict in the backbone of *Alchemilla* s.l. as recently reported in other

groups (e.g., Gonçalves et al. 2019; Walker et al. 2019; Zhang et al. 2020b; Morales-Briones et

al. 2021).

279   *Mapping whole genome duplications*

We took two alternative approaches for detecting WGD events by mapping gene duplication

events from gene trees onto a map tree (e.g., species tree). The first approach begins by

extracting orthogroups from the final homolog trees. Orthogroups are rooted ingroup lineages

separated by outgroups that include the complete set of genes in a lineage from a single copy in

their common ancestor. We extracted orthogroups requiring at least 50 out of 79 species in

Fragariinae. Gene duplication events were then recorded on the most recent common ancestor

(MRCA) on the map tree when two or more species overlapped between the two daughter clades

Each node on a map tree can be counted only once from each gene tree to avoid nested gene

duplications inflating the number of recorded duplications (Yang et al. 2018;

https://bitbucket.org/blackrim/clustering, 'extract_clades.py' and 'map_dups_mrca.py'). We

mapped duplication events onto both the MO and RT trees using orthogroups from all 923 final

homologs, filtering orthogroups using an average BS of at least 50%. We carried out the

mapping using two sets of orthogroups, one from all homologs, and the from the longest

homologs (the single longest aligned exon per gene) to avoid inflating the counts in multi-exon

genes.

For the second strategy of WGD mapping, we explicitly tested for polyploidy mode using

GRAMPA (Thomas et al. 2017). GRAMPA uses MRCA reconciliation with multi-labeled gene

trees to compare allo-or autopolyploid scenarios in singly- or multi-labeled map trees (e.g.,

species tree). To reduce the computational burden of searching all possible reconciliations, we

constrained searches to only among crown nodes of major clades of *Alchemilla* s.l., which all are

well supported (including the 'dissected' and 'lobed' clades of Eualchemilla; see results) and

genera within Fragariinae. We ran reconciliation searches using all 923 final homologs, as well

as using only the longest homologs (the single longest aligned exon per gene; 256), against either

the MO or RT tree. We expected multiple WGD events within *Alchemilla* s.l. (see results), but

GRAMPA can only infer one WGD at a time. To disentangle nested duplication events, we also

carried out similar GRAMPA reconciliations using the MO tree and sequentially excluding

major groups of *Alchemilla* s.l. that were identified as a polyploid clade. We only used the MO

tree as it differs from the RT tree only by the location of the 'lobed' clade, which was the first

clade identified as allopolyploid (see results) and was removed for subsequent GRAMPA

analyses. Finally, to test for a polyploid origin of *Alchemilla* s.l., we carried out searches among

the constrained crown node of *Alchemilla* s.l. and the rest of the genera within Fragariinae using

the MO and cpDNA trees. The backbone of Fragariinae differed between the MO (same as RT)

tree and the cpDNA tree. Thus, we tested how this affected the inference of the polyploid origin

of *Alchemilla* s.l. We also carried out similar searches but using each of the five major clades

individually.

Both approaches used here to detect WGD events use final homolog trees and as any

other tree-based method they may be sensitive to tree informativeness. To explore node support

across homologs, we run a conflict analysis with Phyparts using individual final homologs trees

with a BS support filter of at least 50% for each node. We used both the MO and RT trees as

map trees and ran the analysis using all homolog exons as well as only the longest homolog exon

per gene.

322   *Distribution of synonymous distance among gene pairs (Ks plots)*

To obtain further evidence for WGD events and compare them to those inferred from gene

duplication events from target enrichment, we analyzed the distribution of synonymous distances

(Ks) from RNA-seq data of four species of *Alchemilla* s.l. and nine species of Fragariinae (Table

S4). Read processing and transcriptome assembly followed Morales-Briones et al. (2020). For

each of the four species, a Ks plot of within-species paralog pairs based on BLASTP hits was

done following Yang et al. (2018; https://bitbucket.org/blackrim/clustering; 'ks_plots.py'). Ks

peaks were identified using a mixture model as implemented in mixtools v.1.2.0 (Benaglia et al.

2009). The optimal number of mixing components was estimated using parametric bootstrap

replicates of the likelihood ratio test statistic (McLachlan and Peel 2000). We tested up to five

components using 500 bootstrap replicates in mixtools. Additionally, we used between-species

Ks plots to determine the relative timing of the split between two species and compare it to that

of WGD events inferred with within-species Ks plots. Ks plots of between-species also followed

Yang et al. (2018; 'ks_between_taxa_cds.py'). Lastly, we also attempted to build Ks plots using

raw homologs from target enrichment, but the relatively low number of genes (256) failed to

produce a meaningful distribution (not shown).

339   **RESULTS**

340   *Assembly and orthology inference*

The number of assembled exons per species (with > 75% of the target length) ranged from 632

(*Alchemilla fissa*) to 934 (*Dasiphora fruticosa*) out of 939 single-copy exon references from *F.*

vesca, with an average of 873 exons (Table S5). The number of exons with paralog warnings

ranged from 10 in *Drymocallis glandulosa* to 746 in *Alchemilla mollis* (Table S5). The number

of exon alignments with ≥ 25 species was 923 from 256 genes. The orthology inference resulted

in 914 MO orthologs (Table S6), and 1,906 RT orthologs (Table S6). The trimmed alignments of

the MO orthologs ranged from 136 to 5,740 characters with a mean of 425 characters (median =

268). The concatenated alignment of the MO orthologs, with at least 150 aligned characters and

25 species for each exon, included 910 exons and 387,042 characters with a matrix occupancy of

66%. The trimmed alignments of the RT orthologs ranged from 136 to 5,740 characters with a

mean of 394 characters (median = 259). The concatenated alignment of the RT orthologs, with at

least 150 aligned characters and 25 species, included 1,894 exons and 746,562 characters with a

matrix occupancy of 54%. The chloroplast alignment included 124,079 characters with a matrix

occupancy of 77%.

356   *Nuclear phylogenetic analyses*

All nuclear analyses recovered the monophyly of *Alchemilla* s.l. with maximum support (i.e.,

bootstrap percentage [BS] = 100, local posterior probabilities [LPP] = 1.0; Fig. 2; Fig. S1), most

informative gene trees being concordant with this node (858 out of 863 for MO; 977/984 for RT;

912/932 for ASTRAL-Pro; ICA = 0.95), and full QS support (1.0/–/1.0; i.e., all sampled quartets

supported that node). Five major clades were identified within *Alchemilla* s.l.: Afromilla,

Aphanes, Eualchemilla-'dissected', Eualchemilla-'lobed,' and *Lachemilla*. Moreover, the

relationships among these clades showed high levels of discordance and varied among the MO

and RT trees.

365   Analyses of the MO orthologs using ASTRAL and concatenated ML approaches resulted

in similar topologies for the backbone of *Alchemilla* s.l. (Fig. 2). The monophyly of the five

major clades each received maximum support (BS = 100; LPP = 1.0) and had most trees being

concordant (except for the two clades of Eualchemilla). Eualchemilla was paraphyletic and split

into the 'dissected' and 'lobed' clades. Monophyly of the 'dissected' clade was supported by 118

out of 429 informative trees (ICA = 0.08) and strong QS score (0.87/0.34/1), while the 'lobed'

clade was supported by 73 out of 420 informative trees (ICA = 0.06) and strong QS score

(0.61/0.98/0.99). In both cases, the 'dissected' and 'lobed' clades had a relatively small

percentage of supporting trees, but the conflict analysis and QS score did not reveal any well-

supported alternative topology. *Aphanes* was recovered as sister to the Eualchemilla-'lobed'

clade with relatively low support (BS = 90, LPP = 0.62), 60 concordant trees (out of 430

informative gene trees; ICA = 0.08), and weak QS score (0.016/0.95/0.98) with similar

frequencies for the two discordant alternative topologies. The Eualchemilla-'dissected' clade was

recovered as sister to Eualchemilla-'lobed' + *Aphanes* with maximum support, 279 concordant

trees (out of 482 informative gene trees; ICA = 0.29), and full QS score. Afromilla was

recovered as sister to the clade consisted of Eualchemilla ('dissected and 'lobed') and *Aphanes*

with high to low support (BS = 100, LPP = 0.88), only 146 concordant trees (out of 413

informative gene trees; ICA = 0.22), and weak QS support (0.2/0.44/0.99) with a skew in

discordance suggesting a possible alternative topology (*Lachemilla* sister to Eualchemilla +

Aphanes). Lastly, *Lachemilla* was recovered as the sister to the rest of *Alchemilla* s.l.

Analysis of the RT orthologs using ASTRAL and concatenated ML approaches both

recovered the same major clades, but they differed in the relationship among these five clades

(Fig. 2a; Fig. S1). In both analyses, *Lachemilla*, Afromilla*, and *Aphanes* had maximum support

(BS = 100; LPP = 1.0) and had most trees being concordant. Eualchemilla was recovered as

monophyletic and composed of the 'dissected' and 'lobed' clades. The monophyly of

Eualchemilla had high to low support (BS = 99, LPP = 0.63), only 231 concordant trees (out of

819 informative gene trees; ICA = 0.12), and weak QS support (0.023/0.87/0.98) with similar

frequencies for the two discordant alternative topologies. Similar to the MO analyses, the

'dissected' and 'lobed' clades each had low number of concordant trees (218 out of 557 [ICA =

0.19] and 136 out of 707 [ICA = 0.08], respectively), and strong QS support (0.98/0/1 and

0.62/0.17/0.99, respectively). Eualchemilla was recovered as sister of *Aphanes* with maximum

support (BS = 100), 348 concordant trees (out of 728 informative trees; ICA = 0.29) and full QS

support. The ML concatenated tree (Fig. 2a; Fig. S1) placed Afromilla as sister to the clade

formed of Eualchemilla and *Aphanes* with maximum support (BS = 100), 212 concordant gene

trees (out of 771 informative trees; ICA = 0.27), and weak QS support (0.18/0.66/0.99) with no

significant skew between the two discordant alternatives. *Lachemilla* was placed as sister to the

rest of *Alchemilla* s.l. The ASTRAL tree in turn (Fig. S1a), retrieved *Lachemilla* as sister to the

clade formed of Eualchemilla and *Aphanes* with no support (LPP = 0.01), 247 concordant trees

(out of 953 informative trees; ICA = 0.19), and QS counter-support (-0.21/0.29/0.99), showing

that the majority of the quartets supported one alternative topology (Afromilla sister to

Eualchemilla + *Aphanes*). In this case, Afromilla was placed as sister to the rest of *Alchemilla* s.l.

406   The ASTRAL-Pro analysis using multi-labeled homolog trees recovered the same

backbone topology of *Alchemilla* s.l. as the concatenated ML analysis from the RT orthologs

(Fig. 2b; Figs S2–S3). All five major clades had the maximum support (LPP = 1.0).

Eualchemilla, composed of the 'dissected' and 'lobed' clades, had moderate support (LPP = 0.76)

and only 415 concordant trees (out of 1106 informative trees; ICA = 0.17). The 'dissected' and

'lobed' clades had low numbers of concordant trees (379 out of 941 [ICA = 0.23] and 65 out of

824 [ICA = 0.09], respectively), but did not show signal of any alternative topology. *Aphanes*

was placed as the sister of Eualchemilla with maximum support (LPP = 1.0), 426 concordant

trees (out of 952 trees; ICA = 0.34), and no support for any major alternative topology. Afromilla

was recovered as sister to the clade formed of Eualchemilla and *Aphanes* with low support (LPP

= 0.52), 492 concordant trees (out of 953 trees; ICA = 0.42), and no support for any alternative

topology.

419   *Chloroplast phylogenetic analyses*

The chloroplast ML tree (Fig. 2c; Fig. S4) recovered a well-supported backbone *Alchemilla* s.l.

where the monophyly of *Aphanes,* Afromilla, and *Lachemilla,* had maximum or near maximum

support (i.e., bootstrap percentage [BS] = 100, QS support = [1.0/–/1.0]). Eualchemilla,

composed of the 'dissected' and 'lobed' clades, also had the maximum support. The 'dissected'

and 'lobed' clades had strong support (BS = 75, QS = 0.8/0.43/0.88 and BS = 100, QS =

0.95/0.25/0.92, respectively). *Aphanes* and Eualchemilla formed, with maximum support, a clade

as in the nuclear analyses. In turn, Afromilla and *Lachemilla* were recovered as sister clades with

maximum support, which differed from the nuclear analyses.

Figure 2. Maximum likelihood phylogeny of *Alchemilla* s.l. inferred from RAxML analysis of
the concatenated 910-nuclear exon supermatrix from the 'monophyletic outgroup' (MO)
orthologs. Bootstrap support (BS) and Local posterior probability (LLP) are shown above
branches. Nodes with full support (BS= 100/LLP= 1) are noted with an asterisk (*). Em dashes
(—) denoted alternative topology compared to the ASTRAL tree. Quartet Sampling (QS) scores
for major clades are shown below branches. QS scores in blue indicate strong support and red
scores indicate weak support. QS scores: Quartet concordance/Quartet differential/Quartet

informativeness. Pie charts for major clades represent the proportion of exon ortholog trees that

support that clade (blue), the proportion that support the main alternative bifurcation (green), the

proportion that support the remaining alternatives (red), and the proportion (conflict or support)

that have < 50% bootstrap support (gray). Gene trees with missing data that were uninformative

for the node were ignored. Branch lengths are in number of substitutions per site (scale bar on

the bottom). Inset: a) Summary Maximum likelihood phylogeny inferred from RAxML analysis

of the concatenated 1,894-nuclear exon supermatrix from the 'rooted ingroup' orthologs (RT).

BS and LLP are shown above branches and QS scores below the branches. Branch lengths are in

number of substitutions per site; b) Summary ASTRAL-Pro tree inferred from 923 multi-labeled

exon homolog trees. LLP are shown next to nodes. Branch lengths are in coalescent units. c)

Summary Maximum likelihood phylogeny inferred from RAxML analysis of concatenated

partial plastomes. BS and LLP are shown above branches and QS scores below the branches.

Branch lengths are in number of substitutions per site.

450   *Mapping whole genome duplications*

By mapping the most recent common ancestor (MRCA) of gene duplication events from

orthogroup trees onto the MO and RT trees, we found four nodes in *Alchemilla* s.l. that each had

an elevated proportion of gene duplications (Fig. 3a–b). This trend was consistent regardless of

using all 923 homolog exons (868 after orthogroup inference and BS filtering) or using only the

256 longest homolog exons per gene (250 after orthogroup inference and BS filtering; Fig. S5).

Therefore, here we describe the results only for the latter. These four clades include (Fig. 3a; Fig.

S5): 1) the MRCA of *Alchemilla* s.l. (86.0% of the 250 genes show evidence of duplication), 2)

the MRCA of Eualchemilla , *Aphanes*, and Afromilla (34.4%), 3) the MRCA of Eualchemilla +

Aphanes (MO: 18.4%; RT:15.6%), and 4) the MRCA of *Lachemilla* (18.4%). These four nodes

have an elevated proportion of gene duplications compared to all other nodes in Fragariinae (Fig.

3b) and it is consistent with the number of paralogs counted from the final homolog trees (after

pruning of clades or paraphyletic grades of same species; Fig. 3c). Interestingly, although deeply

nested in *Alchemilla* s.l., *Aphanes* had a lower number of paralogs than the rest of *Alchemilla* s.l.,

resembling the other members of Fragariinae (Fig. 3c).

Figure 3. Orthogroup gene duplication mapping results. a) Summarized cladogram of *Alchemilla*

s.l. from the 'monophyletic outgroup' (MO) ortholog tree. Percentages next to nodes denote the

proportion of duplicated genes when using orthogroups from the longest homologs (250 after

orthogroup inference and filtering). Nodes with elevated proportions of gene duplications are

numbered 1–4 as referenced in the main text. See Fig. S5 for the full tree. b) Histogram of

percentages of gene duplication per branch. c) Number of paralogs per taxa in the final homolog

trees. In final homologs clades and paraphyletic grades of the same species were pruned leaving

only one tip per species. Each locus is represented by the longest homolog (the single longest

aligned exon per gene).

Bootstrap support for exon homologs were informative (BS ≥ 50%) at most nodes,

especially regarding the relationship among the major clades of *Alchemilla* s.l. (Fig. S6).

Therefore, uninformative homolog trees were unlikely to affect the results from WGD detection

analysis overall. The proportion of uninformative nodes (BS < 50%) were at most 30% in the

worst case (Eualchemilla + *Aphanes* + Afromilla) when using all homolog exons. This

proportion reduces significantly when using only the longest homolog exons (Fig. S6).

Similar to the results of MRCA mapping, the GRAMPA analyses recovered the same

results when using all 923 homologs or only the longest homologs (256). GRAMPA

reconciliations using all major clades of *Alchemilla* s.l. recovered optimal multi-labeled trees

with the best score (i.e., lowest reconciliation score; Fig. S7) where the 'lobed' clade of

Eualchemilla was of an allopolyploid origin, but the putative parental lineages varied between

the MO and RT trees. The reconciliations using the MO tree (reconciliation score [RS] = 70,250;

Fig. 4a; Fig. S8) showed that the 'lobed' clade was of allopolyploid origin between an unsampled

or extinct lineage sister to *Aphanes* and an unsampled or extinct lineage ('lineage' for short

hereafter) sister to 'dissected' + *Aphanes*. In turn, the reconciliations using the RT tree (RS =

70,721; Fig. S8) showed that the 'lobed' clade was of allopolyploid origin between a 'lineage'

sister to the 'dissected' clade, and also a 'lineage' sister to 'dissected' + *Aphanes*. Alternative

multi-labeled trees had higher (worse) RSs (70,482 for MO and 70,739 for RT; Fig. S7). The

GRAMPA reconciliations performed on the MO tree with removal of major clades of *Alchemilla*

s.l. inferred as allopolyploid resulted in the identification of additional polyploidy events (Fig.

4b–d). First, we removed the 'lobed' clade, and this resulted in the recovery of Afromilla as an

allopolyploid clade (RS = 127,836). Afromilla parental lineages were a 'lineage' sister to

Aphanes + the 'dissected' clade, and a 'lineage' sister to all remaining *Alchemilla* s.l. (Fig. 4b).

Alternative multi-labeled trees reconciliations had scores starting at 127,869 (Fig. S7). The

further removal of Afromilla resulted in recovery of the 'dissected' clade as allopolyploid (RS =

167,545). The 'dissected' clade had as parental lineages the 'lineage' sister to *Aphanes* and the

'lineage' sister to all remaining *Alchemilla* s.l. except for *Lachemilla* (Fig. 4c). Other

reconciliation alternatives had scores starting at 167,612 (Fig. S7). Finally, the removal of the

'dissected' clade resulted in the *Lachemilla* being recovered also as an allopolyploid clade (RS =

181,302). The parental lineages of *Lachemilla* were a 'lineage' sister to *Aphanes* and a 'lineage'

sister to all remaining *Alchemilla* s.l. (Fig. 4d). Alternative multi-labeled trees reconciliations had

scores starting at 181,564 (Fig. S7).

509   The GRAMPA results from the analyses with constrained searches on the crown node of

Alchemilla s.l. recovered different modes of polyploidy when using the MO tree or the cpDNA

tree. The MO tree had *Farinopsis*, *Sibbaldianthe* + *Sibbaldia*, *Comarum*, and *Fragaria* forming a

grade sister to *Alchemilla* s.l., while *Drymocallis*, *Chamaerhodos*, *Potaninia*, and *Dasiphora*

form a clade that is sister to all other Fragariinae (Fig. 2). The reconciliations using the MO tree

resulted in an allopolyploid event for the clade composed of *Alchemilla* s.l., *Farinopsis,*

Sibbaldianthe, and *Sibbaldia* (RS = 339,755; Fig. S9). The parental lineages of this clade were a

'lineage' sister to *Comarum*, and a 'lineage' sister to the grade formed of *Comarum* and

Fragaria (Fig. S9). Alternative multi-labeled trees had scores starting at 340,053 (Fig. S7). The

reconciliations using individual major clades of *Alchemilla* s.l. resulted in identical patterns as in

the full constrained analysis (Fig. S10). The cpDNA tree had *Alchemilla* s.l. as part of a grade

formed along with *Farinopsis*, *Comarum*, and *Sibbaldianthe* + *Sibbaldia*, while *Fragaria* was

recovered as sister to the clade composed of *Drymocallis*, *Chamaerhodos*, *Potaninia*, and

Dasiphora, which is sister to all other Fragariinae (Fig. S4). The reconciliations on the cpDNA

tree recovered *Alchemilla* s.l. as of autopolyploid origin (RS = 364,594; Fig. 4e; Fig. S9).

Alternative multi-labeled trees had scores starting at 363,987 (Fig. S7). The analyses using

individual major clades of *Alchemilla* s.l. recovered identical patterns as in the full constrained

analysis, except for *Aphanes* that resulted in a singly-labeled tree (Fig. S10).

To further explore WGD events using alternative data sources, we analyzed Ks plots

from genomes and transcriptomes across Fragariinae. The distribution of synonymous distances

in the transcriptomes of four species of Eualchemilla (one 'dissected' and three 'lobed') shared

three optimal mixing components with a Ks mean at approximately 0.1, 0.34, and 1.67,

respectively (Fig. S11). The first two components partially overlapped and corresponded to at

least two WGD events in all four sampled species of Eualchemilla, that happened before the

splits between the lobed vs. the dissected clades of Eualchemilla (Ks ~ 0.02; Fig. S12). The third

shared component corresponds to a whole genome triplication event early in the core eudicots

(Jiao et al. 2012; Fig. S11). All nine species from other genera in Fragariinae had two optimal

mixing components. One component is a Ks peak at 1.61–1.78 corresponding to the whole

genome triplication event early in eudicot (Fig. S11). In the case of the diploid species, the

second component represents a small and very young (~ 0.05) peak, most likely the product of

small-scale recent gene duplications. The only two polyploid species from the other genera in

Fragariinae, *Comarum palustre* (2n=28–64) and *Sibbaldianthe bifurca* (2n=28), had a single

additional significant component at 0.11 and 0.08, respectively (Fig. S11). The Ks plots between

species of Eualchemilla and Fragariinae species outside of *Alchemilla* s.l., and between species

of Fragariinae showed that the WGD events detected in Eualchemilla were not shared with other

genera outside of *Alchemilla* s.l. Likewise, the WGD events in *Comarum palustre* and

Sibbaldianthe bifurca occurred after the split of the two species (Fig. S12).

Figure 4. Summary of optimal multi-labeled tree (MUL-tree) inferred from GRAMPA analyses.

a) MUL-tree based on reconciling homologs against the species tree inferred from 'monophyletic

outgroup' (MO) orthologs including all taxa. Red branches denote the allopolyploid origin of the

'lobed' clade of Eualchemilla. b) MUL-tree after removing the 'lobed' clade of Eualchemilla as in

a). Green branches denote the allopolyploid origin of Afromilla. c) MUL-tree after removing

Afromilla as in b). Blue branches denote the allopolyploid origin of the 'dissected' clade of

Eualchemilla. d). MUL-tree after further removing the 'dissected' clade as in c). Yellow lines

denote the allopolyploid origin of *Lachemilla*. e) MUL-tree using constrained searches of the

crown node of *Alchemilla* s.l. on the cpDNA tree. Orange branches denote the autopolyploid

origin *Alchemilla* s.l. f) Putative summary network of all reticulation events in *Alchemilla* s.l.

Colored curved branches denote different polyploid events as in a–e. Dashed curved lines

represent the maternal lineage (cpDNA) in allopolyploid events.

561   DISCUSSION

562   *Processing paralogs in target enrichment datasets*

The increased use of target enrichment methods in combination with reduced sequencing costs

and higher read coverage have facilitated the recovery of paralogs in such datasets. Paralogy is

sometimes viewed as a nuisance for phylogenetic reconstruction and is commonly aimed to be

reduced in early stages of experimental design, by targeting only single- or low-copy genes

during the selection of loci (e.g., Chamala et al. 2015; Nicholls et al. 2015; Gardner et al. 2016;

Kamneva et al. 2017). Still, the recovery of paralogs is inevitable when working with groups

where WGD is prevalent, especially in plants, leading to various strategies to remove them prior

to phylogenetic analyses. Commonly used target enrichment assembly pipelines (e.g., Faircloth

2016; Johnson et al. 2016; Andermann et al. 2018) use different criteria to flag assembled loci

with putative paralogs that are later filtered or processed prior to phylogenetic analysis. The most

used common approach for dealing with paralogous loci in target enrichment datasets is

removing the entire locus that show any signal of potential paralogy (e.g., Crowl et al. 2017;

Montes et al. 2019; Bagley et al. 2020). The removal of paralogous loci can significantly reduce

the size of target enrichment datasets and most often do not take in consideration the reason why

a locus was flagged for putative paralogy (i.e., allelic variation or gene duplication). Orthology

inference should be carried for all loci in target enrichment data, as relying on settings in

assembly pipelines does not guarantee that non-removed or non-flagged loci are orthologous.

Furthermore, removing paralogs before phylogenetic inference eliminates valuable information

that could have been used to detect and place WGD events using target enrichment data. Other

approaches either retain or remove contigs based on the distinction being putative allelic

variation (flagged sequences from monophyletic conspecific groups) or putative paralogs

(paralogs from the same species are non-monophyletic) in combination with study-specific

threshold or random selection (e.g., Villaverde et al; 2018; Liu et al. 2019; Stubbs et al. 2019), or

manual processing (e.g., Garcia et al. 2017; Karimi et al. 2019). As dataset size increases,

manual processing becomes prohibitive.

The presence of WGDs also poses some challenges for locus assembly. Target

enrichment design commonly includes multi-contig targets that assembly pipelines attempt to

assemble into single contigs (e.g., Faircloth 2016) or 'supercontigs' composed of multiple exons

and partially assembled introns (e.g., Johnson et al. 2016). In groups like *Alchemilla* s.l., where

multiple, nested WGD events led to a prevalence of paralogs, 'supercontigs' can produce

chimeric assemblies (Morales-Briones et al. 2018b). Instead, we assembled the exons

individually to minimize chimeric loci, at the cost of working with some short exons that

contribute little phylogenetic information, which can affect orthology inference and downstream

analyses. Therefore, it is important to take this into consideration during target enrichment

experimental design, and to preferentially target long exons when possible in groups where

WGD is expected. An alternative strategy to avoid chimeric supercontigs when gene duplications

are frequent is to perform a preliminary orthology inference in single exon-based trees and then

use the inferred orthologs as a reference to reassemble the loci into 'supercontigs' (e.g., Gardner

et al. 2020; Karimi et al. 2020). Another aspect to take in consideration during or right after

assembly is allele phasing. While phasing heterozygous loci, from population or individual

variation, have been shown to have minimal impact in phylogenetic reconstruction in target

enrichment data (e.g., Kates et al. 2018), the effect on unphased or merged loci in cases of WGD

can be larger and be a source of gene tree error. Here we were interested in ancient WGD in

Alchemilla s.l. and relied on enough sequencing coverage and sequence dissimilarity to assemble

separate paralogs (homoeologs in the case of allopolyploidy) that can be flagged as such by

HybPiper. While we obtained a large number of deep paralogs across *Alchemilla* s.l. (Fig. 3c;

Table S5), there is still the possibility of some locus included merged sequences from paralogs

with high sequence similarity. Paralog merger should be more problematic in cases of recent

allopolyploidy or neo-allopolyploidy taxa. To this end, recently developed tools have been

designed to phase gene copies into polyploid subgenomes using phylogenetic and similarity

approaches (e.g., Freyman et al. 2020, Nauheimer et al. 2020).

The utility of paralogs for phylogenetic reconstruction in target enrichment datasets is

gaining more attention (e.g., Johnson et al. 2016; Gardner et al. 2020). A few studies have

considered tree-based orthology inference to process affected loci (e.g., Garcia et al. 2017;

Moore et al. 2018, Morales-Briones et al. 2018b), but in some cases the orthology approaches

used cannot be applied to other groups. Here we demonstrated the utility of automated, tree-

based orthology inference methods (Yang and Smith 2014), originally designed for genomic or

transcriptomic datasets, to infer orthology from paralog-flagged loci in a target enrichment

dataset. Our approach facilitates the automated inference of orthologs while maximizing the

number of loci retained for downstream analyses. These methods are agnostic of the data source

and should work for any type of target enrichment dataset (e.g., anchored phylogenomics, exon

capture; Hyb-Seq, ultraconserved elements).

Orthology inference methods used here (Yang and Smith et al. 2014) are a powerful tool

for target enrichment datasets. In the case of allopolyploidy, however, these methods can

introduce bias in the distribution of ortholog trees inferred. In the case of MO, each time a gene

duplication event is detected, the side with a smaller number of taxa is removed. When

allopolyploidy occurs, MO may bias towards one subgenome due to 1) bias in gene loss between

subgenomes. Even if the submissive subgenome is present in some parts of the genome, it is

differentially lost in a higher number of loci, 2) bias in bait design. In the case of *Alchemilla* s.l.,

this is less likely as baits are designed in outgroups. If baits are designed according to ingroup

taxa, depending on which taxa were used it can have higher affinity to one subgenome instead of

another, and 3) unequal sampling of parental lineages. If one parental lineage is more densely

sampled than the other or one parental lineage is unsampled, the two subgenomes will be in

species-rich versus species-poor clades respectively in gene trees. One could alternatively

preserve a random side each time a gene duplication event is identified. However, in practice, the

side with a smaller number of taxa often contains misassembled or misplaced sequences. The RT

method of separating duplicated gene copies, on the other hand, keeps any subtree with sufficient

number of taxa, but removes outgroups, and worked best when hierarchical outgroups were

included in the taxon sampling. Therefore, both MO and RT lose information, especially in cases

with complex, nested polyploidy. Recently developed methods based on quartet similarity

(Zhang et al. 2020a) or Robinson-Foulds distances (Molloy and Warnow 2020) can directly

estimate species trees from multi-labeled trees that are consistent with the MSC and gene

duplication and loss without inferring orthologs (for a recent review see Smith and Hahn 2020).

However, their behavior on complex datasets using archival materials is yet to be explored. For

example, both methods do not define ingroup-outgroup relationships a priori, and correctly

inferring the root of homolog trees can be challenging with missing data, or when WGD occurs

near the root. In addition, none of these above species tree reconstruction methods (Molloy and

Warnow 2020; Zhang et al. 2020a) were designed to handle reticulate relationships. This can

result in species tree topology that is an "average" between subgenomes. Depending on the

topological distance of subgenomes, the resulting species tree may not represent any subgenome

history. Finally, most current methods for evaluating node support still require orthologous gene

trees as input. In such cases tools like Phyparts can still be used to visualize gene tree

discordance and calculate ICA scores using multi-labeled trees.

657   *Phylogenetic implications in Alchemilla s.l.*

Previous phylogenetic studies established the monophyly *Alchemilla* s.l. and four major clades of

the group (Gehrke et al. 2008; 2016), but the relationship among them and the placement of

Alchemilla s.l. within Fragariinae remain unresolved. Our nuclear and plastid analyses both

confirmed the monophyly of *Alchemilla* s.l. and its sister relationship to *Farinopsis*, as

previously shown by Morales-Briones and Tank (2019) based on plastome sequences only.

Gehrke et al. (2008) identified two well supported clades within Eualchemilla that were

distinguished by leaf shape, namely the 'dissected' and 'lobed' clades. Most species of

Eualchemilla have a leaf shape consistent with their clade placement, but some had different leaf

shapes that were attributed to their hybrid origin between the two clades (Gehrke et al. 2008).

More recently, Gehrke et al. (2016) and Morales-Briones and Tank (2019) found that

Eualchemilla is not monophyletic in analyses that included the external transcribed spacer (ETS)

of the nuclear ribosomal DNA (nrDNA) cistron. Both studies found *Aphanes* nested between the

'dissected' and 'lobed' clades of Eualchemilla. Our analyses of the nuclear loci supported the

monophyly of 'dissected' and 'lobed' clades, but the monophyly of Eualchemilla had low support

(Fig. 2; Fig. S1). The analysis using only the MO orthologs even weakly supported the 'lobed'

clade as sister of *Aphanes* (Fig. 2). In contrast, our plastome analysis recovered a well-supported,

monophyletic Eualchemilla, as well as well-supported 'dissected' and 'lobed' clades. Both nuclear

and plastid analyses strongly supported the clade composed of *Aphanes* and both clades of

Eualchemilla (Fig. 2; Fig. S4), a relationship that is consistent with previous nuclear and plastid

analyses (Gehrke et al. 2008; 2016; Morales-Briones and Tank 2019). Given the revealed

hybridization in the evolution and early divergence within *Alchemilla* s.l., the non-monophyly of

Eualchemilla could be explained by ancient gene flow or an allopolyploid origin of the

'dissected' and 'lobed' clades (Fig. 4a,c; see below). Besides the well supported relationship of

Eualchemilla + *Aphanes*, our nuclear analysis showed high levels of conflict among other major

clades in *Alchemilla* s.l. (Fig. 2; Fig. S1) which could also be explained by additional ancient

allopolyploid events (Fig. 4; see below).

685   *Ancient polyploidy in Alchemilla s.l.*

Whole-genome duplications are frequent across Rosaceae (Dickinson et al. 2007; Xiang et al.

2017), and allopolyploidy has been suggested as the primary source for the cytonuclear

discordance in Fragariinae (Lundberg et al. 2009; Gehrke et al. 2016; Morales-Briones and Tank

2019). We recovered four nodes in *Alchemilla* s.l. with a high percentage of gene duplications

(Fig. 3a; Fig. S5). One of the nodes showing a high percentage of gene duplication (18.4%) was

the MRCA of *Aphanes* and both clades of Eualchemilla (node 3 in Fig. 3a; Fig. S5). This

duplication event agreed with the MRCA of the ancestral lineages inferred with GRAMPA for

the allopolyploid origin of the 'lobed' clade of Eualchemilla (Fig. 4a). Moreover, the GRAMPA

reconciliations after the removal of the 'lobed' clade and Afromilla inferred a scenario where the

'dissected' clade is of allopolyploid origin with one of the parental lineages as sister to *Aphanes*

(Fig. 4c). Although there is some uncertainty about the placement of the parental lineage of

'dissected' clade, due to the removal of major clades for the GRAMPA analyses, the cpDNA tree

suggest that it is likely sister to the parental lineage of 'lobed' clade that is also sister to *Aphanes*.

Ks plots of all species of the 'dissected' and 'lobed' clades had two peaks that are not shared with

members of Fragariinae (Figs S11–S12), suggesting that at least two WGD events have

happened between the stem lineage of *Alchemilla* s.l. to the crown node of the 'dissected' and the

'lobed' clades. The between-species Ks plots between 'dissected' and 'lobed' (Fig. S12), showed

that the split between these two groups is more recent than the WGD events, suggesting a single

origin (or very close in time) of both clades. Still, the sister relationship of the 'dissected' and

'lobed' clades is not supported by nuclear genes, suggesting that the two clades of Eualchemilla

might had independent allopolyploid origins, while sharing the same or a closely related

maternal lineage (cpDNA; Fig. 4f).

The GRAMPA reconciliation, after the removal of the 'lobed' clade, recovered an

allopolyploid origin of Afromilla (Fig. 4b) with a MRCA of the ancestral lineages at the crown

of the remaining *Alchemilla* s.l. Similarly, the further removal of both Afromilla and the

'dissected' clade recovered *Lachemilla* as allopolyploid, with the MRCA of parental lineages

mapped to the crown of the remaining *Alchemilla* s.l. (Fig. 4d). In the case of *Lachemilla,*

because of the removal of major clades for the GRAMPA analyses, there is also no certainty in

the placement of its parental lineages. Still, Afromilla and *Lachemilla* are sisters in the cpDNA

tree (Fig 1C.), suggesting these two share the same or a closely related maternal lineage (Fig. 4f).

The high percentage of gene duplication (34.4%) placed at the MRCA of the clade composed of

Afromilla, Eualchemilla, and *Aphanes* (node 2 in Fig. 3a), could be explained in part by the

allopolyploid origin of Afromilla.

Finally, the node with the highest percentage of duplicated genes (86%) was placed at the

MRCA of *Alchemilla* s.l. (node 1 in Fig. 3a). The GRAMPA analysis using the MO tree showed

an allopolyploid event for the clade that included *Alchemilla* s.l., *Farinopsis salesoviana*,

Sibbaldia, and *Sibbaldianthe* (Fig. S9). However, an allopolyploid origin of *Farinopsis*

salesoviana, *Sibbaldia*, and *Sibbaldianthe* is not supported by chromosome numbers, orthogroup

gene duplication counts, or Ks plots. All members in Fragariinae, with the exception of

Alchemilla s.l. mainly consists of diploid species and base chromosome number of seven ($x = 7$),

including *Sibbaldia* and *Sibbaldianthe.* On the other hand, *Alchemilla* s.l. has a base number of

eight ($x = 8$) and contains mostly species with high ploidy levels (octoploid to 24-ploid), with the

exception of most species of *Aphanes* ($2n=16$) and one species of *Lachemilla* (*L. mandoniana,*

$2n=16$). Also, our gene duplication counts show low percentages (1.6%) of gene duplication for

the MRCA of the GRAMPA-inferred allopolyploid clade or the MRCA (3.6%) of the inferred

parental lineages (Fig. 3). Previous phylotranscriptomic analyses of Rosaceae (Xiang et al. 2017)

that included one species each of the 'dissected' and 'lobed' clades of Eualchemilla, found

33.21% of duplicated genes for the MRCA of these two clades, but did not recover any other

node with elevated gene duplications within Fragariinae. The Ks plots of the four species of

Alchemilla s.l. all showed peaks with similar Ks means, but these peaks were not shared with

species of *Sibbaldia* and *Sibbadianthe* (Fig. S11). Furthermore, the between-species Ks plots

showed that the WGD events detected in *Alchemilla* were more recent than the split with

members of Fragariinae (Fig. S12). Although the chromosome number and Ks data for

Farinopsis salesoviana are not available, all the above evidence suggest an unlikely

allopolyploid origin of the clade consisting of *Farinopsis, Sibbaldia*, *Sibadianthe*, and *Alchemilla*

sl. On the other hand, the GRAMPA reconciliations using the cpDNA tree resulted in an optimal

multi-labeled tree where *Alchemilla* s.l. had an autopolyploid origin (Fig. 4e). This scenario is

compatible with the high percentage of gene duplication at the MRCA of *Alchemilla* s.l. and the

low percentage of gene duplication in the backbone of the rest of Fragariinae. Another

compatible scenario is an allopolyploid origin of *Alchemilla* s.l. where both parental lineages are

missing or extinct, but this scenario is indistinguishable from autopolyploidy. The atypical high

proportion of gene duplication at the base of *Alchemilla* s.l. can be explained by the

autopolyploid event at this branch. In addition, given the short branch lengths among major

clades within *Alchemilla* s.l., gene tree estimation error (e.g., uninformative genes), incomplete

lineage sorting (ILS), allopolyploid events among major clades of *Alchemilla* s.l., and/or

homoeologous exchanges among subgenomes (Edger et al. 2018; McKain et al. 2018) can all

contribute to additional gene duplication events mapped to the MRCA of *Alchemilla* s.l.

Although deeply nested in *Alchemilla* s.l., remarkably, *Aphanes* showed a significantly

lower number of paralogs than the rest of *Alchemilla* s.l. (Fig. 3). The relatively low number of

paralogs, its diploid species being mainly diploid, and the best GRAMPA reconciliation resulting

in a singly-labeled tree (Fig. S10), suggesting that *Aphanes* is a functional diploid clade. One

plausible scenario is that post-polyploid diploidization (reviewed in Mandáková and Lysak 2018)

occurred after the autopolyploidy event at the base of *Alchemilla* s.l. After diploidization,

Afromilla, Eualchemilla ('lobed' and 'dissected' clades), and *Lachemilla* originated from

allopolyploid events (Fig. 4f). On the other hand, *Aphanes* seems to descend from a diploidized

ancestor that did not duplicate further. The orthogroup gene duplication mapping showed

Aphanes as part of a clade that had nested high proportions of gene duplication in the orthogroup

mapping (Fig. 3a-b, nodes 2–3). But this does not necessarily mean that *Aphanes* should show

the same duplication pattern, or neither does it contradict its diploid condition, as a duplication

event does not affect or include all descendants of the mapped MRCA in the map tree (e.g.,

species trees).

GRAMPA has been shown to be useful to identify multiple polyploidy events in the same

tree (e.g., Thomas et al. 2017; Guo et al. 2020; Koenen et al. 2020), but a tree-based approach

can also be sensitive to gene tree estimation error or ILS (Thomas et al. 2017). Methods to infer

species networks in the presence of ILS (e.g., Solís-Lemus and Ané 2016; Wen et al. 2018) could

also be used to explore the prevalence of ancient hybridization in *Alchemilla* s.l. Although these

methods are under constant development and improvement, they are still only tractable in simple

scenarios with few reticulation events (Hejase and Liu 2016; Kamneva and Rosenberg 2017).

Similarly, the signal of the *D*-Statistic (Green et al. 2010; Durand et al. 2011), commonly used to

detect introgression, can be lost or distorted in presence of multiple reticulations (Elworth et al.

2018). Complex reticulate scenarios like *Alchemilla* s.l. are likely to face these problems and

have phylogenetic network and *D*-statistic identifiability issues as seen in other groups (e.g.,

Morales-Briones et al. 2021).

780   *Conclusions*

In this study, we have shown the utility of target enrichment datasets in combination with

tree-based methods for orthology inference and WGD investigation. Here, we used *Alchemilla*

s.l. to highlight the importance of processing paralogs, rather than discarding them before

phylogenetic analysis, to shed light on the complex polyploidy histories. We showed evidence

that the entire *Alchemilla* s.l. is the product of an ancient autopolyploidy event, and that

Afromilla, Eualchemilla ('lobed' and 'dissected' clades), and *Lachemilla* originated from

subsequent and nested ancient allopolyploid events. Our results from analyzing target enrichment

data corroborated with previously published chromosome numbers and distribution of Ks values

from transcriptomes. Our analyses has several important implications for future target

enrichment projects, including 1) design baits to obtain a relatively large number of loci as this is

required for accurate species tree and networks estimation in complex scenarios (e.g., higher

levels of ILS; Solís-Lemus and Ané 2016; Nute et al. 2018), 2) increase the length of individual

loci to improve the information content of individual gene trees for proper tree-based orthology

inference and identifying gene duplication events, and 3) design baits to minimize lineage-

specific and paralog-specific capture efficiency and missing data. Furthermore, in target

enrichment, unlike genome or transcriptome data, only a few hundreds of genes are typically

recovered with levels of missing data that varies by lineage and are non-random. This limits the

utility of target enrichment for generating Ks plots, and creates the need to carefully scrutinize

the variation in percentage of gene duplications among nodes. In the end, even with these

limitations, target enrichment is an overall valuable and cost-effective approach of genomic

subsampling to explore patterns of reticulation and WGD, especially in groups for which whole

genome or transcriptome data are not possible to generate, including from museum/herbarium

specimens. As research continues to deepen in other clades across the Tree of Life using similar

target enrichment methods, we expect that other complex patterns of duplication and reticulation,

as those shown here in *Alchemilla* s.l. will continue to emerge.

807   SUPPLEMENTARY MATERIAL

Data available from the Dryad Digital Repository: http://dx.doi.org/10.5061/.[NNNN]

ACKNOWLEDGMENTS

This work was supported by the University of Minnesota, Graduate Student Research Grants

from the Botanical Society of America, American Society of Plant Taxonomists, International

Association of Plant Taxonomists, the University of Idaho Stillinger Herbarium Expedition

Funds to D.F.M.-B., and a National Science Foundation Doctoral Dissertation Improvement

Grant to D.C.T. for D.F.M.-B. (DEB-1502049).

REFERENCES

Andermann T., Cano Á., Zizka A., Bacon C., Antonelli A. 2018. SECAPR—a bioinformatics

820   pipeline for the rapid and user-friendly processing of targeted enriched Illumina

821   sequences, from raw reads to alignments. PeerJ. 6:e5175.

822   Andermann T., Torres Jiménez M.F., Matos-Maraví P., Batista R., Blanco-Pastor J.L.,

823   Gustafsson A.L.S., Kistler L., Liberal I.M., Oxelman B., Bacon C.D., Antonelli A. 2020.

824   A Guide to Carrying Out a Phylogenomic Target Sequence Capture Project. Front.

825   Genet. 10:1407.

826   Bagley J.C., Uribe-Convers S., Carlsen M.M., Muchhala N. 2020. Utility of targeted sequence

827   capture for phylogenomics in rapid, recent angiosperm radiations: Neotropical

828   Burmeistera bellflowers as a case study. Mol. Phylogenet. Evol.:106769.

829   Benaglia T., Chauveau D., Hunter D.R., Young D. 2009. mixtools : An R Package for Analyzing

830   Finite Mixture Models. J. Stat. Softw. 32:1–29.

831   Brown J.W., Walker J.F., Smith S.A. 2017. Phyx - phylogenetic tools for unix. Bioinformatics.

832   33:1886–1888.

Buddenhagen C., Lemmon A.R., Lemmon E.M., Bruhl J., Cappa J., Clement W.L., Donoghue

M.J., Edwards E.J., Hipp A.L., Kortyna M., Mitchell N., Moore A., Prychid C.J.,

Segovia-Salcedo M.C., Simmons M.P., Soltis P.S., Wanke S., Mast A. 2016. Anchored

Phylogenomics of Angiosperms I: Assessing the Robustness of Phylogenetic Estimates.

bioRxiv.:086298.

Chamala S., García N., Godden G.T., Krishnakumar V., Jordon-Thaden I.E., De Smet R.,

Barbazuk W.B., Soltis D.E., Soltis P.S. 2015. MarkerMiner 1.0: A new application for

phylogenetic marker development using angiosperm transcriptomes. Appl. Plant Sci.

3:1400115.

Crowl A.A., Manos P.S., McVay J.D., Lemmon A.R., Lemmon E.M., Hipp A.L. 2019.

Uncovering the genomic signature of ancient introgression between white oak lineages

(Quercus). New Phytol.:nph.15842.

Dickinson T.A., Lo E., Talent N. 2007. Polyploidy, reproductive biology, and Rosaceae:

understanding evolution and making classifications. Plant Syst. Evol. 266:59–78.

Dobeš C., Lückl A., Kausche L., Scheffknecht S., Prohaska D., Sykora C., Paule J. 2015. Parallel

origins of apomixis in two diverged evolutionary lineages in tribe Potentilleae

(Rosaceae): Origin of Apomixis in Potentilleae. Bot. J. Linn. Soc. 177:214–229.

Doyle J.J., Doyle J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf

tissue. Phytochem. Bull. 19:11–15.

Dunn C.W., Howison M., Zapata F. 2013. Agalma: an automated phylogenomics workflow.

BMC Bioinformatics. 14:330.

Durand E.Y., Patterson N., Reich D., Slatkin M. 2011. Testing for Ancient Admixture between

Closely Related Populations. Mol. Biol. Evol. 28:2239–2252.

Edger P.P., McKain M.R., Bird K.A., VanBuren R. 2018. Subgenome assignment in

857   allopolyploids: challenges and future directions. Curr. Opin. Plant Biol. 42:76–80.

858   Elworth R.A.L., Allen C., Benedict T., Dulworth P., Nakhleh L.K. 2018. DGEN: A Test Statistic

859   for Detection of General Introgression Scenarios. WABI.

860   Emms D.M., Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative

861   genomics. Genome Biol. 20:238.

862   Eriksson T., Lundberg M., Töpel M., Östensson P., Smedmark J.E.E. 2015. Sibbaldia: a

863   molecular phylogenetic study of a remarkably polyphyletic genus in Rosaceae. Plant

864   Syst. Evol. 301:171–184.

865   Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic

866   loci. Bioinforma. Oxf. Engl. 32:786–788.

867   Fernández R., Gabaldon T., Dessimoz C. 2020. Orthology: Definitions, Prediction, and Impact

868   on Species Phylogeny Inference. In: Scornavacca C., Delsuc F., Galtier N., editors.

869   Phylogenetics in the Genomic Era. No commercial publisher | Authors open access

870   book. p. 2.4:1--2.4:14.

871   Fitch W.M. 1970. Distinguishing Homologous from Analogous Proteins. Syst. Biol. 19:99–113.

872   Forrest L.L., Hart M.L., Hughes M., Wilson H.P., Chung K.-F., Tseng Y.-H., Kidner C.A. 2019.

873   The Limits of Hyb-Seq for Herbarium Specimens: Impact of Preservation Techniques.

874   Front. Ecol. Evol. 7:439.

875   Freyman W.A., Johnson M.G., Rothfels C.J. 2020. homologizer: Phylogenetic phasing of gene

876   copies into polyploid subgenomes. bioRxiv 2020.10.22.351486

877   García N., Folk R.A., Meerow A.W., Chamala S., Gitzendanner M.A., Oliveira R.S. de, Soltis

878   D.E., Soltis P.S. 2017. Deep reticulation and incomplete lineage sorting obscure the

diploid phylogeny of rain-lilies and allies (Amaryllidaceae tribe Hippeastreae). Mol.

Phylogenet. Evol. 111:231–247.

Gardner E.M., Johnson M.G., Pereira J.T., Ahmad Puad A.S., Arifiani D., Sahromi, Wickett

N.J., Zerega N.J.C. 2020. Paralogs and off-target sequences improve phylogenetic

resolution in a densely-sampled study of the breadfruit genus (Artocarpus, Moraceae).

Syst. Bio. syaa073.

Gardner E.M., Johnson M.G., Ragone D., Wickett N.J., Zerega N.J.C. 2016. Low-coverage,

whole-genome sequencing of Artocarpus camansi (Moraceae) for phylogenetic marker

development and gene discovery. Appl. Plant Sci. 4:1600017.

Gehrke B., Bräuchler C., Romoleroux K., Lundberg M., Heubl G., Eriksson T. 2008. Molecular

phylogenetics of Alchemilla, Aphanes and Lachemilla (Rosaceae) inferred from plastid

and nuclear intron and spacer DNA sequences, with comments on generic classification.

Mol. Phylogenet. Evol. 47:1030–1044.

Gehrke B., Kandziora M., Pirie M.D. 2016. The evolution of dwarf shrubs in alpine

environments: a case study of Alchemilla in Africa. Ann. Bot. 117:121–131.

Glover N., Dessimoz C., Ebersberger I., Forslund S.K., Gabaldón T., Huerta-Cepas J., Martin

M.-J., Muffato M., Patricio M., Pereira C., da Silva A.S., Wang Y., Sonnhammer E.,

Thomas P.D. 2019. Advances and Applications in the Quest for Orthologs. Mol. Biol.

Evol. 36:2157–2164.

Gonçalves D.J.P., Simpson B.B., Ortiz E.M., Shimizu G.H., Jansen R.K. 2019. Incongruence

between gene trees and species trees and phylogenetic signal variation in plastid genes.

Mol. Phylogenet. Evol. 138:219–232.

Green R.E., Krause J., Briggs A.W., Maricic T., Stenzel U., Kircher M., Patterson N., Li H., Zhai

W., Fritz M.H.Y., Hansen N.F., Durand E.Y., Malaspinas A.S., Jensen J.D., Marques-

Bonet T., Alkan C., Prufer K., Meyer M., Burbano H.A., Good J.M., Schultz R., Aximu-

Petri A., Butthof A., Hober B., Hoffner B., Siegemund M., Weihmann A., Nusbaum C.,

Lander E.S., Russ C., Novod N., Affourtit J., Egholm M., Verna C., Rudan P., Brajkovic

D., Kucan Z., Gusic I., Doronichev V.B., Golovanova L.V., Lalueza-Fox C., de la

Rasilla M., Fortea J., Rosas A., Schmitz R.W., Johnson P.L.F., Eichler E.E., Falush D.,

Birney E., Mullikin J.C., Slatkin M., Nielsen R., Kelso J., Lachmann M., Reich D.,

Paabo S. 2010. A Draft Sequence of the Neandertal Genome. Science. 328:710–722.

Guo X., Mandáková T., Trachtová K., Özüdoğru B., Liu J., Lysak M.A. 2020. Linked by

ancestral bonds: multiple whole-genome duplications and reticulate evolution in a

Brassicaceae tribe. Mol. Biol. Evol. msaa327.

Hayirhoğlu-Ayaz S., İnceer H., Frost-Olsen P. 2006. Chromosome counts in the genus

Alchemilla (Rosaceae) from SW Europe. Folia Geobot. 41:335–344.

Hejase H.A., Liu K.J. 2016. A scalability study of phylogenetic network inference methods using

empirical datasets and simulations involving a single reticulation. BMC Bioinformatics.

17:422.

Hjelmqvist H. 1956. The embryology of some African *Alchemilla* species. Bot. Not. 109:21–32.

Huang C.-H., Zhang C., Liu M., Hu Y., Gao T., Qi J., Ma H. 2016. Multiple Polyploidization

Events across Asteraceae with Two Nested Events in the Early History Revealed by

Nuclear Phylogenomics. Mol. Biol. Evol. 33:2820–2835.

Izmailow R. Karyological studies in species of *Alchemilla* L. from the series *Calycinae* Bus.

(section *Brevicaulon* Rothm.). Acta Biol. Cracoviensia Ser. Bot. 23:117–130.

Jiao Y., Wickett N.J., Ayyampalayam S., Chanderbali A.S., Landherr L., Ralph P.E.,

Tomsho L.P., Hu Y., Liang H., Soltis P.S., Soltis D.E., Clifton S.W., Schlarbaum S.E.,

Schuster S.C., Ma H., Leebens-Mack J., dePamphilis C.W. 2011. Ancestral polyploidy

in seed plants and angiosperms. Nature. 473:97–100.

Jiao Y., Leebens-Mack J., Ayyampalayam S., Bowers J.E., McKain M.R., McNeal J., Rolf M.,

Ruzicka D.R., Wafula E., Wickett N.J., Wu X., Zhang Y., Wang J., Zhang Y., Carpenter

E.J., Deyholos M.K., Kutchan T.M., Chanderbali A.S., Soltis P.S., Stevenson D.W.,

McCombie R., Pires J., Wong G., Soltis D.E., dePamphilis C.W. 2012. A genome

triplication associated with early diversification of the core eudicots. Genome Biol.

13:R3.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett

N.J. 2016. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics from

High-Throughput Sequencing Reads Using Target Enrichment. Appl. Plant Sci.

4:1600016.

Jones K.E., Fér T., Schmickl R.E., Dikow R.B., Funk V.A., Herrando-Moraira S., Johnston P.R.,

Kilian N., Siniscalchi C.M., Susanna A., Slovák M., Thapa R., Watson L.E., Mandel

J.R. 2019. An empirical assessment of a single family-wide hybrid capture locus set at

multiple evolutionary timescales in Asteraceae. Appl. Plant Sci. 7:e11295.

Kamneva O.K., Rosenberg N.A. 2017. Simulation-Based Evaluation of Hybridization Network

Reconstruction Methods in the Presence of Incomplete Lineage Sorting. Evol.

Bioinforma. 13:117693431769193.

Kamneva O.K., Syring J., Liston A., Rosenberg N.A. 2017. Evaluating allopolyploid origins in

strawberries (Fragaria) using haplotypes generated from target capture sequencing.

BMC Evol. Biol. 17:401.

Karimi N., Grover C.E., Gallagher J.P., Wendel J.F., Ané C., Baum D.A. 2020. Reticulate

Evolution Helps Explain Apparent Homoplasy in Floral Biology and Pollination in

Baobabs (Adansonia; Bombacoideae; Malvaceae). Syst. Biol. 69:462–478.

Katoh K., Standley D.M. 2013. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780.

Kates H.R., Johnson M.G., Gardner E.M., Zerega N.J.C., Wickett N.J. 2018. Allele phasing has

minimal impact on phylogenetic reconstruction from targeted nuclear gene sequences in

a case study of Artocarpus. Am. J. Bot. 105:404–416.

Kearse M., Moir R., Wilson A., Stones-Havas S., Cheung M., Sturrock S., Buxton S., Cooper A.,

Markowitz S., Duran C., Thierer T., Ashton B., Meintjes P., Drummond A. 2012.

Geneious Basic: An integrated and extendable desktop software platform for the

organization and analysis of sequence data. Bioinformatics. 28:1647–1649.

Kocot K.M., Citarella M.R., Moroz L.L., Halanych K.M. 2013. PhyloTreePruner: A

Phylogenetic Tree-Based Approach for Selection of Orthologous Sequences for

Phylogenomics. Evol. Bioinforma. Online. 9:429–435.

Koenen E.J.M., Ojeda D.I., Bakker F.T., Wieringa J.J., Kidner C., Hardy O.J., Pennington R.T.,

Herendeen P.S., Bruneau A., Hughes C.E. 2020. The Origin of the Legumes is a

Complex Paleopolyploid Phylogenomic Tangle closely associated with the Cretaceous-

Paleogene (K-Pg) Mass Extinction Event. Syst. Biol. syaa041.

Larridon I., Villaverde T., Zuntini A.R., Pokorny L., Brewer G.E., Epitawalage N., Fairlie I.,

Hahn M., Kim J., Maguilla E., Maurin O., Xanthos M., Hipp A.L., Forest F., Baker W.J.

2020. Tackling Rapid Radiations With Targeted Sequencing. Front. Plant Sci. 10:1655.

Leebens-Mack J.H., Barker M.S., Carpenter E.J., Deyholos M.K., Gitzendanner M.A., Graham

S.W., Grosse I., Li Z., Melkonian M., Mirarab S., Porsch M., Quint M., Rensing S.A.,

Soltis D.E., Soltis P.S., Stevenson D.W., Ullrich K.K., Wickett N.J., DeGironimo L.,

Edger P.P., Jordon-Thaden I.E., Joya S., Liu T., Melkonian B., Miles N.W., Pokorny L.,

Quigley C., Thomas P., Villarreal J.C., Augustin M.M., Barrett M.D., Baucom R.S.,

Beerling D.J., Benstein R.M., Biffin E., Brockington S.F., Burge D.O., Burris J.N.,

Burris K.P., Burtet-Sarramegna V., Caicedo A.L., Cannon S.B., Çebi Z., Chang Y.,

Chater C., Cheeseman J.M., Chen T., Clarke N.D., Clayton H., Covshoff S., Crandall-

Stotler B.J., Cross H., dePamphilis C.W., Der J.P., Determann R., Dickson R.C., Di

Stilio V.S., Ellis S., Fast E., Feja N., Field K.J., Filatov D.A., Finnegan P.M., Floyd

S.K., Fogliani B., García N., Gâteblé G., Godden G.T., Goh F. (Qi Y., Greiner S.,

Harkess A., Heaney J.M., Helliwell K.E., Heyduk K., Hibberd J.M., Hodel R.G.J.,

Hollingsworth P.M., Johnson M.T.J., Jost R., Joyce B., Kapralov M.V., Kazamia E.,

Kellogg E.A., Koch M.A., Von Konrat M., Könyves K., Kutchan T.M., Lam V., Larsson

A., Leitch A.R., Lentz R., Li F.-W., Lowe A.J., Ludwig M., Manos P.S., Mavrodiev E.,

McCormick M.K., McKain M., McLellan T., McNeal J.R., Miller R.E., Nelson M.N.,

Peng Y., Ralph P., Real D., Riggins C.W., Ruhsam M., Sage R.F., Sakai A.K.,

Scascitella M., Schilling E.E., Schlösser E.-M., Sederoff H., Servick S., Sessa E.B.,

Shaw A.J., Shaw S.W., Sigel E.M., Skema C., Smith A.G., Smithson A., Stewart C.N.,

Stinchcombe J.R., Szövényi P., Tate J.A., Tiebel H., Trapnell D., Villegente M., Wang

C.-N., Weller S.G., Wenzel M., Weststrand S., Westwood J.H., Whigham D.F., Wu S.,

Wulff A.S., Yang Y., Zhu D., Zhuang C., Zuidof J., Chase M.W., Pires J.C., Rothfels

C.J., Yu J., Chen C., Chen L., Cheng S., Li J., Li R., Li X., Lu H., Ou Y., Sun X., Tan

X., Tang J., Tian Z., Wang F., Wang J., Wei X., Xu X., Yan Z., Yang F., Zhong X.,

Zhou F., Zhu Y., Zhang Y., Ayyampalayam S., Barkman T.J., Nguyen N., Matasci N.,

Nelson D.R., Sayyari E., Wafula E.K., Walls R.L., Warnow T., An H., Arrigo N.,

Baniaga A.E., Galuska S., Jorgensen S.A., Kidder T.I., Kong H., Lu-Irving P., Marx

H.E., Qi X., Reardon C.R., Sutherland B.L., Tiley G.P., Welles S.R., Yu R., Zhan S.,

Gramzow L., Theißen G., Wong G.K.-S., One Thousand Plant Transcriptomes Initiative.

2019. One thousand plant transcriptomes and the phylogenomics of green plants. Nature.

574:679–685.

Li L. 2003. OrthoMCL: Identification of Ortholog Groups for Eukaryotic Genomes. Genome

Res. 13:2178–2189.

Li Z., Baniaga A.E., Sessa E.B., Scascitelli M., Graham S.W., Rieseberg L.H., Barker M.S.

2015. Early genome duplications in conifers and other seed plants. Sci. Adv.

1:e1501084.

Liu Y., Johnson M.G., Cox C.J., Medina R., Devos N., Vanderpoorten A., Hedenäs L., Bell N.E.,

Shevock J.R., Aguero B., Quandt D., Wickett N.J., Shaw A.J., Goffinet B. 2019.

Resolution of the ordinal phylogeny of mosses using targeted exons from organellar and

nuclear genomes. Nat. Commun. 10:1485.

Lundberg M., Töpel M., Eriksen B., Nylander J.A.A., Eriksson T. 2009. Allopolyploidy in

Fragariinae (Rosaceae): Comparing four DNA sequence regions, with comments on

classification. Mol. Phylogenet. Evol. 51:269–280.

Lynch M., Conery J.S. 2000. The Evolutionary Fate and Consequences of Duplicate Genes.

1014   Science. 290:1151–1155.

1015   Mai U., Mirarab S. 2018. TreeShrink: fast and accurate detection of outlier long branches in

1016   collections of phylogenetic trees. BMC Genomics. 19:272.

1017   Mandáková T., Lysak M.A. 2018. Post-polyploid diploidization and diversification through

1018   dysploid changes. Curr. Opin. Plant Biol. 42:55–65.

1019   Mandel J.R., Dikow R.B., Funk V.A., Masalia R.R., Staton S.E., Kozik A., Michelmore R.W.,

1020   Rieseberg L.H., Burke J.M. 2014. A Target Enrichment Method for Gathering

1021   Phylogenetic Information from Hundreds of Loci: An Example from the Compositae.

1022   Appl. Plant Sci. 2:1300085.

1023   Nauheimer L., Weigner N., Joyce E., Crayn D., Clarke C., Nargar K. 2020. HybPhaser: a

1024   workflow for the detection and phasing of hybrids in target capture datasets. bioRxiv

1025   2020.10.27.354589

1026   McKain M.R., Estep M.C., Pasquet R., Layton D.J., Vela Díaz D.M., Zhong J., Hodge J.G.,

1027   Malcomber S.T., Chipabika G., Pallangyo B., Kellogg E.A. 2018. Ancestry of the two

1028   subgenomes of maize. bioRxiv.:352351.

1029   McKain M.R., Tang H., McNeal J.R., Ayyampalayam S., Davis J.I., dePamphilis C.W., Givnish

1030   T.J., Pires J.C., Stevenson D.W., Leebens-Mack J.H. 2016. A Phylogenomic Assessment

1031   of Ancient Polyploidy and Genome Evolution across the Poales. Genome Biol. Evol.

1032   8:1150–1164.

1033   McLachlan G., Peel D. 2000. Finite Mixture Models. New York: Wiley.

1034   Molloy E.K., Warnow T. 2020. FastMulRFS: fast and accurate species tree estimation under

1035   generic gene duplication and loss models. Bioinformatics. 36:i57–i65.

1036   Montes J.R., Peláez P., Willyard A., Moreno-Letelier A., Piñero D., Gernandt D.S. 2019.

1037   Phylogenetics of Pinus Subsection Cembroides Engelm. (Pinaceae) Inferred from Low-

1038   Copy Nuclear Gene Sequences. Syst. Bot. 44:501–518.

1039   Montgomery L. 1997. Contributions to a cytological catalogue of the British and Irish flora, 5.

1040   Watsonia. 21:365–368.

1041   Moore A.J., Vos J.M.D., Hancock L.P., Goolsby E., Edwards E.J. 2018. Targeted Enrichment of

1042   Large Gene Families for Phylogenetic Inference: Phylogeny and Molecular Evolution of

1043   Photosynthesis Genes in the Portullugo Clade (Caryophyllales). Syst. Biol. 67:367–383.

1044

1045   Morales-Briones D.F., Romoleroux K., Kolář F., Tank D.C. 2018a. Phylogeny and Evolution of

1046   the Neotropical Radiation of *Lachemilla* (Rosaceae): Uncovering a History of Reticulate

1047   Evolution and Implications for Infrageneric Classification. Syst. Bot. 43:17–34.

1048   Morales-Briones D.F., Liston A., Tank D.C. 2018b. Phylogenomic analyses reveal a deep history

1049   of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae). New

1050   Phytol. 218:1668–1684.

1051   Morales-Briones D.F., Tank D.C. 2019. Extensive allopolyploidy in the neotropical genus

1052   *Lachemilla* (Rosaceae) revealed by. Am. J. Bot. 106:415–437.

1053   Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A.,

1054   Brockington S.F., Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021. Disentangling

1055   Sources of Gene Tree Discordance in Phylogenomic Datasets: Testing Ancient

1056   Hybridizations in Amaranthaceae s.l. Syst. Biol. 70:219–235

1057   Morton J. 1993. Chromosome numbers and polyploidy in the flora of Cameroons Mountain.

1058   Opera Bot. 121:159–172.

Nicholls J.A., Pennington R.T., Koenen E.J.M., Hughes C.E., Hearn J., Bunnefeld L., Dexter

K.G., Stone G.N., Kidner C.A. 2015. Using targeted enrichment of nuclear genes to

increase phylogenetic resolution in the neotropical rain forest genus Inga (Leguminosae:

Mimosoideae). Front. Plant Sci. 6:710.

Nute M., Chou J., Molloy E.K., Warnow T. 2018. The performance of coalescent-based species

tree estimation methods under models of missing data. BMC Genomics. 19:286.

Panchy N., Lehti-Shiu M., Shiu S.-H. 2016. Evolution of Gene Duplication in Plants. Plant

Physiol. 171:2294–2316.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling

distinguishes lack of support from conflicting support in the green plant tree of life. Am.

J. Bot. 105:385–403.

Perry L.M. 1929. A Tentative Revision of Alchemilla § Lachemilla. Contrib. Gray Herb. Harv.

Univ.:1–57.

Ranwez V., Douzery E.J.P., Cambon C., Chantret N., Delsuc F. 2018. MACSE v2: Toolkit for

the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. Mol.

Biol. Evol. 35:2582–2584.

Salichos L., Stamatakis A., Rokas A. 2014. Novel Information Theory-Based Measures for

Quantifying Incongruence among Phylogenetic Trees. Mol. Biol. Evol. 31:1261–1271.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Shulaev V., Sargent D.J., Crowhurst R.N., Mockler T.C., Folkerts O., Delcher A.L., Jaiswal P.,

Mockaitis K., Liston A., Mane S.P., Burns P., Davis T.M., Slovin J.P., Bassil N.,

Hellens R.P., Evans C., Harkins T., Kodira C., Desany B., Crasta O.R., Jensen R.V.,

Allan A.C., Michael T.P., Setubal J.C., Celton J.-M., Rees D.J.G., Williams K.P., Holt

S.H., Rojas J.J.R., Chatterjee M., Liu B., Silva H., Meisel L., Adato A., Filichkin S.A.,

Troggio M., Viola R., Ashman T.-L., Wang H., Dharmawardhana P., Elser J., Raja R.,

Priest H.D., Bryant D.W., Fox S.E., Givan S.A., Wilhelm L.J., Naithani S., Christoffels

A., Salama D.Y., Carter J., Girona E.L., Zdepski A., Wang W., Kerstetter R.A., Schwab

W., Korban S.S., Davik J., Monfort A., Denoyes-Rothan B., Arus P., Mittler R., Flinn

B., Aharoni A., Bennetzen J.L., Salzberg S.L., Dickerman A.W., Velasco R.,

Borodovsky M., Veilleux R.E., Folta K.M. 2011. The genome of woodland strawberry

(Fragaria vesca). Nat. Genet. 43:109–116.

Smedmark J.E.E., Eriksson T., Evans R.C., Campbell C.S. 2003. Ancient Allopolyploid

Speciation in Geinae (Rosaceae): Evidence from Nuclear Granule-Bound Starch

Synthase (GBSSI) Gene Sequences. Syst. Biol. 52:374–385.

Smith M.L., Hahn M.W. 2020. New Approaches for Inferring Phylogenies in the Presence of

Paralogs. Trends Genet.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals

conflict, concordance, and gene duplications with examples from animals and plants.

BMC Evol. Biol. 15:150.

Soják J. 2008. Notes on Potentilla XXI. A new division of the tribe Potentilleae (Rosaceae) and

notes on generic delimitations. Bot. Jahrb. Für Syst. Pflanzengesch. Pflanzengeogr.

127:349–358.

Solís-Lemus C., Ané C. 2016. Inferring Phylogenetic Networks with Maximum

Pseudolikelihood under Incomplete Lineage Sorting. PLoS Genet. 12:e1005896–21.

Stamatakis A. 2014. RAxML version 8 - a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 30:1312–1313.

Straub S.C., Fishbein M., Livshultz T., Foster Z., Parks M., Weitemier K., Cronn R.C., Liston A.

2011. Building a model: developing genomic resources for common milkweed

(Asclepias syriaca) with low coverage genome sequencing. BMC Genomics. 12:211.

Stubbs R.L., Folk R.A., Xiang C.-L., Chen S., Soltis D.E., Cellinese N. 2020. A Phylogenomic

Perspective on Evolution and Discordance in the Alpine-Arctic Plant Clade Micranthes

(Saxifragaceae). Front. Plant Sci. 10:1773.

Thomas G.W.C., Ather S.H., Hahn M.W. 2017. Gene-Tree Reconciliation with MUL-Trees to

Resolve Polyploidy Events. Syst. Biol. 66:1007–1018.

Villaverde T., Pokorny L., Olsson S., Rincón-Barrado M., Johnson M.G., Gardner E.M., Wickett

N.J., Molero J., Riina R., Sanmartín I. 2018. Bridging the micro- and macroevolutionary

levels in phylogenomics: Hyb-Seq solves relationships from populations to species and

above. New Phytol. 220:636–650.

Walker J.F., Walker-Hale N., Vargas O.M., Larson D.A., Stull G.W. 2019. Characterizing gene

tree conflict in plastome-inferred phylogenies. PeerJ. 7:e7747.

Walters S., Boznan V. *Alchemilla faeroensis* (Lange) Buser and *A. alpina* L. Proc. Bot. Soc. Br.

Isles. 7:83.

Weitemier K., Straub S.C.K., Cronn R.C., Fishbein M., Schmickl R., McDonnell A., Liston A.

2014. Hyb-Seq: Combining Target Enrichment and Genome Skimming for Plant

Phylogenomics. Appl. Plant Sci. 2:1400042.

Wen D., Yu Y., Zhu J., Nakhleh L. 2018. Inferring Phylogenetic Networks Using PhyloNet.

Syst. Biol. 67:735–740.

Xiang Y., Huang C.-H., Hu Y., Wen J., Li S., Yi T., Chen H., Xiang J., Ma H. 2017. Evolution

of Rosaceae Fruit Types Based on Nuclear Phylogeny in the Context of Geological

Times and Genome Duplication. Mol. Biol. Evol. 34:262–281.

Yang Y., Moore M.J., Brockington S.F., Mikenas J., Olivieri J., Walker J.F., Smith S.A. 2018.

Improved transcriptome sampling pinpoints 26 ancient and more recent polyploidy

events in Caryophyllales, including two allopolyploidy events. New Phytol. 217:855–

870.

Yang Y., Moore M.J., Brockington S.F., Soltis D.E., Wong G.K.-S., Carpenter E.J., Zhang Y.,

Chen L., Yan Z., Xie Y., Sage R.F., Covshoff S., Hibberd J.M., Nelson M.N., Smith

S.A. 2015. Dissecting Molecular Evolution in the Highly Diverse Plant Clade

Caryophyllales Using Transcriptome Sequencing. Mol. Biol. Evol. 32:2001–2014.

Yang Y., Smith S.A. 2014. Orthology Inference in Nonmodel Organisms Using Transcriptomes

and Low-Coverage Genomes: Improving Accuracy and Matrix Occupancy for

Phylogenomics. Mol. Biol. Evol. 31:3081–3092.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

Zhang C., Scornavacca C., Molloy E.K., Mirarab S. 2020a. ASTRAL-Pro: Quartet-Based

Species-Tree Inference despite Paralogy. Mol. Biol. Evol. 37:3292–3307.

Zhang R., Wang Y.-H., Jin J.-J., Stull G.W., Bruneau A., Cardoso D., De Queiroz L.P., Moore

M.J., Zhang S.-D., Chen S.-Y., Wang J., Li D.-Z., Yi T.-S. 2020b. Exploration of Plastid

Phylogenomic Conflict Yields New Insights into the Deep Relationships of

Leguminosae. Syst. Biol. 69:613–622.

Zhbannikov I.Y., Hunter S.S., Foster J.A., Settles M.L. 2017. SeqyClean: A Pipeline for High-

throughput Sequence Data Preprocessing. Proc. 8th ACM Int. Conf. Bioinforma.

Comput. Biol. Health Inform.:407–416.

1152