Recombinant SARS-CoV-2 RBD molecule with a T helper epitope as a built in adjuvant induces strong neutralization antibody response

Qiudong Su[1, #], Yening Zou[2, #], Yao Yi[1, #], Liping Shen[1, #], Xiangzhong Ye[3, #], Yang Zhang[4], Hui Wang[4], Hong Ke[4], Jingdong Song[1], Keping Hu[5], Bolin Cheng[6], Feng Qiu[1], Pengcheng Yu[1], Wenting Zhou[1], Lei Cao[1], Shengli Bi[1, *], Guizhen Wu[1, *], George Fu Gao[1, *], Jerry Zheng [4, *]

[1]National Institute For Viral Disease Control and Prevention, Chinese Center For Disease Control and Prevention, Beijing 102206, China

[2]Sinovac Biotech Co., LTD, Beijing 100085, China

[3]Beijing WanTai Biological Pharmacy Enterprise Co., LTD, Beijing 102206, China

[4]Artron Bioresearch Inc., Burnaby, BC V5A1M6, Canada

[5]Institute of Medicinal Plant Development, Chinese Academy of Medical Science and Peking Union Medical College, Beijing 100193, China

[6] Jianxuan Business Center, Tianjin 300110, China

[#] QD Su, YN Zou, Y Yi, LP Shen and XZ Ye contributed equally.

[*] SL Bi (bisl@ivdc.chinacdc.cn), GZ Wu (wugz@ivdc.chinacdc.cn), GF Gao (gaof@im.ac.cn) and J Zheng (jzheng@artronbio.com) were co-corresponding authors.

Abstract

Without approved vaccines and specific treatment, severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) is spreading around the world with above 20 million

COVID-19 cases and approximately 700 thousand deaths until now. An efficacious and

affordable vaccine is urgently needed. The Val308 – Gly548 of Spike protein of SARS-

CoV-2 linked with Gln830 – Glu843 of Tetanus toxoid (TT peptide) (designated as S1-4)

and without TT peptide (designated as S1-5), and prokaryotic expression,

chromatography purification and the rational renaturation of the protein were performed.

The antigenicity and immunogenicity of S1-4 protein was evaluated by Western Blotting

(WB) in vitro and immune responses in mice, respectively. The protective efficiency of it

was measured by virus neutralization test in Vero E6 cells with SARS-CoV-2. S1-4

protein was prepared to high homogeneity and purity by prokaryotic expression and

chromatography purification. Adjuvanted with Alum, S1-4 protein stimulated a strong

antibody response in immunized mice and caused a major Th2-type cellular immunity

compared with S1-5 protein. Furthermore, the immunized sera could protect the Vero E6

cells from SARS-CoV-2 infection with neutralization antibody GMT 256. The candidate

subunit vaccine molecule could stimulate strong humoral and Th1 and Th2-type cellular

immune response in mice, giving us solid evidence that S1-4 protein could be a

promising subunit vaccine candidate.

Keywords: SARS-CoV-2; RBD; TT peptide; subunit vaccine

Introduction

A highly virulent and pathogenic Coronavirus disease (COVID-19) viral infection

with incubation period ranging from two to fourteen days, transmitted by breathing of

infected droplets or contact with infected droplets, was spreading all over the world[1].

Until 12 August 2020, 20,162,474 COVID-19 cases were reported with 737,417 deaths

globally[2].

COVID-19 is caused by Severe Acute Respiratory Syndrome coronavirus 2 (SARS-

CoV-2)[3]. Coronaviruses named after their morphology as spherical virions with a core

shell and surface projections resembling a solar corona, are enveloped, positive single-

stranded large RNA viruses that infect humans, but also a wide range of animals[4]. SARS-

CoV-2 closely related to the SARS-CoV virus[5] belongs to the B lineage of the beta-

coronaviruses[6]. There are more than 100 candidate vaccines in development worldwide,

while among the vaccine technologies under evaluation are whole virus vaccines,

recombinant protein subunit vaccines, and nucleic acid vaccines[7]. Based on previous

experiences with SARS-CoV vaccines, SARS-CoV-2 vaccines should refrain from

immunopotentiation that could lead to increased infectivity or eosinophilic infiltration[7].

Therefore, subunit vaccine might be one of the better candidate vaccine for SARS-CoV-2

for its simpler in composition and less uncontrollable risk factors[8]. There are four major

structural proteins including the nucleocapsid protein (N), the spike protein (S), a small

membrane protein (SM) and the membrane glycoprotein (M) with an additional

membrane glycoprotein (HE) in the HCoV-OC43 and HKU1 beta-coronaviruses[9]. Like

other coronaviruses, SARS-CoV-2 infects lung alveolar epithelial cells by receptor-

mediated endocytosis with Angiotensin-Converting Enzyme II (ACE2) as the cell

receptor[5]. SARS-CoV-2 binds ACE2 via the receptor binding domain (RBD) of S

protein[10, 11]. Full-length S (S1 and S2) or S1 which contains RBD could induce

neutralization antibodies that prevent viral entry[12]. RBD structurally and functionally

distinct from the remainder of the S1 domain are highly stable and held together by four

disulfide bonds[8]. Numerous researchers have demonstrated that RBD contains abundant

T cell and B cell epitopes including neutralization epitope and has the potential quality as

a subunit vaccine[8, 13-15]. Subunit vaccines possessed high safety profile, consistent

production and could induce cellular and humoral immune responses, but need

appropriate adjuvants[12].

Tetanus toxoid (TT) as a protein carrier could enhance effectively humoral and

cellular immune responses of antigenic epitopes[16]. TT exerts the requisite immunological

enhancement and promotes production of high levels of antibodies through traditional

and effective T-B cell reaction[16]. TT has been proposed because of its safe and also has

been commercially applied in human vaccine[17]. Since virtually all people have

endogenous antibodies against TT, such commonly occurring antibodies are promising

candidates to utilize for immune modulation[18] through antibody complexed antigen.

Gln830 – Glu843 of TT (TT peptide) as a promiscuous T-helper epitope has a strong

binding capacity to DR3 allele[19, 20] and is frequently used as T cell stimulator to enhance

the immunogenicity of exogenous epitopes[19, 21, 22].

Without specific prevention measures, current effective managements mainly focus

on the enforcement of quarantine, isolation and physical distancing[23]. An efficacious and

affordable vaccine is urgently needed. In this study, we prepared a novel molecule of

RBD linked TT peptide (designated as S1-4) and evaluated its antigenicity,

immunogenicity and protective efficiency as potential subunit vaccine.

Results

S1-4 protein was prepared to high homogeneity and purity by prokaryotic expression

and chromatography purification:

Upon induction, in small-expression, the expected S1-4 protein of 29.0 kDa was

observed (Figure 1A) suggesting that S1-4 protein could be appropriately expressed in

E.coli. Through large-scale expression (4 L LB medium), S1-4 protein accounted for

approximately 26.38% in the total cell lysates, and increased up to 70.13% in the total

proteins from the inclusion bodies (IBs) (Figure 1B). Therefore, S1-4 protein expressed

and existed mainly in the IBs form in *E.coli*. S1-4 protein was treated with 8mol/l urea

(pH8.0) and purified by IEX, S1-4 protein existed mainly in the unbound fraction (Figure

1C). The purity of S1-4 protein can reach 96.9% at 2.64 mg/mL through HIC (Figure 1D).

Renaturated by stepwise dialysis, the concentration and purity of S1-4 protein were 0.84

mg/mL and 98.71%, respectively (Figure 1D).

Figure 1. The preparation of S1-4 protein

(A). Small-scale expression of S1-4 protein. (B). Large-scale expression of S1-4 protein.

(C). S1-4 protein purified by ion-exchange chromatography (IEX). (D). S1-4 protein

purified by hydrophobic interaction chromatography (HIC) and renaturation.

1. Negative control; 2. Non-induced expression bacteria; 3 – 4, Induced expression

bacteria; 5. Total bacterial proteins after large-scale expression; 6. Soluble fraction of

bacterial sonication; 7. Insoluble aggregates of bacterial sonication; 8. S1-4 protein in

Urea solution; 9. Unbound fraction eluted by IEX; 10. S1-4 protein fraction eluted by

HIC; 11. Re-natured S1-4 protein after stepwise dialysis.

　　　After diluted with normal saline, S1-4 protein was mixed with aluminum adjuvant

and adjusted antigen concentration in the final adsorption product was 800 μg/mL.

S1-4 protein possessed antigenicity in vitro

The antigenicity of S1-4 protein was demonstrated by WB analysis with a

commercial recombinant RBD protein (Sino Biological, Beijing, China) as positive

control (Figure 2). Whether it was convalescent serum of COVID-19 patients (Figure 2B),

Rabbit monoclonal antibody against RBD (Figure 2C) or Rat polyclonal antibody against

inactivated SARS-CoV-2 (Figure 2D) as the primary antibody, there was an obvious

band at about 30 kDa, suggesting the high antigenicity of S1-4 protein.

Figure 2. Western Blotting analysis of S1-4 protein

(A). Silver staining after SDS-PAGE. (B). Convalescent serum of COVID-19 patients as

the primary antibody. (C). Rabbit monoclonal antibody against RBD as the primary

antibody. (D). Rat polyclonal antibody against inactivated SARS-CoV-2 as the primary

antibody.

1. S1-4 protein; 2. Commercial recombiant RBD protein.

SARS-CoV-2 subunit vaccine displayed strong immunogenicity in immunized mice and

caused a major Th2-type cellular immunity

To analyze the immunogenicity of SARS-CoV-2 candidate subunit vaccine, we

determined the anti-SARS-CoV-2 antibody titers in sera of immunized mice by Enzyme

Linked Immunosorbent Assay (ELISA) (Figure 3A). The total antibody GMT in mice

immunized with S1-4 protein/Alum came to 100 one week after the first injection, while

it climbed to 1481 one week after the second injection. However, the total antibody GMT

in mice immunized with S1-5 protein/Alum was only 24 after the first injection and 192

after the second injection. Finally, one week after the third injection, the total antibody

GMT in mice immunized with S1-4 protein/Alum was 1728, while that of S1-5

protein/Alum was 456. Obviously, S1-4 conjugated with TT peptide tail were more

immunogenic than S1-5 without TT peptide.

Enzyme-linked Immunospot (ELISpot) assay was applied to enumerate Spot-forming

cells (SFCs) for IL-4 and IFN-γ in splenocytes of immunized mice (Figure 3B). More IL-

4 SFCs existed in splencytes of mice immunized with S1-4 protein/Alum comparing with

mice immunized with S1-5 protein/Alum ($P < 0.05$). Meanwhile, there were more IFN-γ

SFCs in splencytes of mice immunized with S1-4 protein/Alum than that of S1-5

protein/Alum ($P < 0.05$). These results indicated that not only Th1 (IFN-γ) but also Th2

(IL-4) SFCs were involved in the immune response against S1-4 protein/Alum and S1-5

protein/Alum. Remarkably, both IL-4 and IFN-γ SFCs were more in mice immunized

with S1-4 protein/Alum than those in mice immunized with S1-5 protein/Alum.

Figure 3. Kinetics of anti-SARS-CoV-2 antibody and neutralization antibody in sera

(A) and IL-4 & IFN-γ spot-forming cells (SFCs) in splenocytes of immunized mice

(B)

S1-4 Immunized sera could protect the Vero E6 cells from SARS-CoV-2 infection

The virus neutralizing activity of SARS-CoV-2 subunit vaccine immunized mice was

examined using live SARS-CoV-2 operating within Biosafety Level 3 (BSL-3)

Laboratory (Figure 3A). The neutralization antibody GMT were 192 on Day28 and

reached 256 on Day35 in the sera of mice immunized with S1-4 protein/Alum. However,

the neutralization antibody GMTs of sera of mice immunized with S1-5 protein/Alum

were only 48 and 64 on Day28 and Day35, respectively. Furthermore, neutralization

antibody GMT increased along with total antibody GMT, whether in S1-4 or S1-5 protein

immunized mice ($P < 0.05$). Those results indicated that TT peptide could not only

enhance the total antibody response, but also the neutralization antibody.

Discussion

The independence of structure and function made RBD the ideal choice of subunit

vaccine. Several expression systems such as CHO-, baculovirus-, and yeast had been

used to express recombinant RBD protein[11, 24-26]. Considering its efficiency, time and cost

saving and the safety consideration, nothing else could match the *E. coli* expression

system, especially in the circumstance that SARS-CoV-2 vaccine is urgently needed. Xia

had developed successfully the Hepatitis E virus (HEV) vaccine[27] and Human

Papillomavirus (HPV) vaccine[28] by prokaryotic expression system. Furthermore, both of

the vaccine had been licenced by National Medical Products Administration of China. By

the approach of prokaryotic expression system, the annual productivity could reach

billions of vaccine dosages after the candidate vaccine was approved, which would make

it possible to prevent and eliminate COVID-19 on a global scale. However, two key

questions, glycosylation and renaturation, bear the brunt, if prokaryotic expression

system is applied. First of all, there are only two predictive N-linked (Asn331 and

Asn343) and O-linked (Thr323 and Thr385) glycosylation sites within Val308 –

Gly548[29]. It was presumed that glycosylations at those sites of RBD were not crucial for

maintance of antigenicity and immunogenicity of it[8]. In this study, S1-4 protein without

glycosylation displayed ideal antigenicity (Figure 2) and immunogenicity (Figure 3).

Moreover, glycosylation and mature structure might hinder the recognition of related

epitopes by immune cells. We found that anti-S1-4 protein antibody titers (unpublished)

were much higher than anti-SARS-CoV-2 antibody titers. Secondly, the existence of four

pairs of cysteine brings great difficulties to the reconstruction of natural structure[8].

Fortunately, our previous experience of SARS-CoV RBD renaturation facilitated the

current work[30].

Humoral immune response, especially the neutralization antibody response, plays an

important role by limiting infection at later phase and prevents reinfection in the future[12].

S1-4 protein can induce high titer neutralizing antibody. One week after the third

immunization, anti-SARS-CoV-2 antibody GMT detected by ELISA reached 1728 while

neutralization antibody GMT came to 256 measured by micro-neutralization assay

(Figure 3A). In contrast, S1-5 protein without TT peptide was barely satisfactory.

SARS-CoV-2 maybe induce delayed type I IFN and miss the best time for viral

control in an early phase of infection[12]. In general, Th1-type immune response is crucial

in an adaptive immunity to eliminate viral infection[12]. The Th1 cells produce cytokines

that promote cell-mediated immune responses such as IFN-γ whilst the Th2 cells induce

cytokines such as IL-4, IL-5 and IL-13, which will activate humoral immune responses[31].

Current evidences strongly indicated that Th1 type response is a key for successful

control of SARS-CoV and MERS-CoV and probably is true for SARS-CoV-2 as well[12].

According to the ELISpot assay, S1-4 protein/Alum could stimulate Th1 and Th2 – type

immune responses simultaneously in mice (Figure 3B). Furthermore, TT peptide played

an important role in promoting cytokine secretion including Th1 and Th2-type cytokines

(Figure 3B).

Protective efficacy is crucial for a vaccine. Neutralization activity of immune serum

is often the key to judge whether a candidate vaccine will be successful or not. The serum

from mice immunized with S1-4 protein/Alum could protect Vero E6 cells from SARS-

CoV-2 infection with neutralization antibody GMT 256 on Day35 (Figure 3A).

In conclusion, our preparation of trunked RBD protein fused TT peptide could

stimulate humoral and Th1 and Th2- type cell-mediated immune response in mice and

the immunized sera could protect Vero E6 cells from SARS-CoV-2 infection. This would

make this subunit protein ideal vaccine candidate.

Experimental Protocol

Preparation of SARS-CoV-2 subunit vaccine

The SARS-CoV subunit antigen, designated as S1-4, consisted of the truncated Spike

protein Val308 – Gly548 (GenBank No. YP009724390) and TT peptide Gln830 –

Glu843 with GGG as the linker (Figure 4). The gene encoding S1-4 protein was codon-

optimized and sub-cloned into pET28a (+) vector using restriction endonucleases (*Nco*I

and *Xho*I). The positive plasmids were confirmed by restriction endonuclease analysis

and sequencing, while the single colony of freshly transformed BL21(DE3) was chosen

depending on the SDS-PAGE analysis of small-scale expression.

Figure 4. Schematic diagram of the gene of SARS-CoV-2 subunit antigen (S1-4 and

S1-5) in the pET28a expression vector

Small-scale expression cultures inoculated with BL21(DE3) carrying S1-4 plasmids

were induced by IPTG (1 mmol/L) at 37°C for 2 h. A large-scale preparation was

performed in 4-L LB medium induced by IPTG (1 mmol/L) at 32°C for 4 h. The cell

harvested by centrifugation (4,000 g, 10 min, 4°C) were resuspended with Buffer I (10

mmol/L Tris-HCl, 1 mmol/L EDTA, 0.1% Triton X-100, pH8.0) before sonic disruption

(300 W, 20 s, 20 s, and 30 times). The pellets retained by centrifugation (17,300 g, 20

min, 4°C) were washed twice with Buffer I. After the pellets were dissolved with Buffer

II (10 mmol/L Tris-HCl, 8 mol/L urea, pH8.0), S1-4 protein in the denatured solution was

purified by DEAE ion-exchange chromatography (IEX) then followed with Octyl

Sepharose 4 Fast Flow hydrophobic interaction chromatography (HIC). After being

appropriately diluted, S1-4 protein was renatured by gradually reducing the urea content

with 20 mmol/L PB (pH8.0) at 4°C for 4 h each cycle. After refolding, S1-4 protein

solution was centrifuged to remove the precipitate and sterile filtered by 0.2 μm filter.

Protein analysis was conducted by SDS polyacrylamide gel electrophoresis (SDS-PAGE)

and Western Blotting (WB) using convalescent serum of COVID-19 patients, Rabbit

monoclonal antibody against RBD and related HRP –conjugated secondary antibodies.

After diluted with normal saline, S1-4 protein was mixed with aluminum adjuvant in a

certain proportion for adsorption, and the antigen concentration in the final adsorption

product was adjusted to 800 μg/mL. The mixture is designated as SARS-CoV-2 subunit

vaccine.

The prokaryotic expression and chromatography purification of S1-5 protein were

performed according to those of S1-4 protein. The MW of S1-5 was approximately 27.4

kDa. Similarly, S1-5 was adjuvanted with Aluminum as a control.

Preliminary analysis of immunogenicity of SARS-CoV-2 subunit vaccine

All animal procedures were reviewed and approved by the Animal Care and Welfare

Committee at the National Institute for Viral Disease Control and Prevention, Chinese

Center for Disease Control and Prevention (No. 20200201003). The neutralization tests

were performed in a BSL-3 laboratory. Ten specific pathogen-free, female 4~6 weeks old

BALB/c mice (Vital River Laboratories, Beijing, China) each group were immunized

i.m. with 50 μL of S1-4/Aluminum, S1-5/ Aluminum or Aluminum alone respectively,

with two booster shots at Day 7 and 14. Orbital venous blood was sampled at regular

intervals (Day 0, 14, 28 and 35) and sera was collected by centrifugation (1,000 g, 10 min,

4°C) after coagulation at room temperature for 2 h.

ELISA was applied to evaluate the antibody response according to the previous

study[32]. Anti-SARS-CoV-2 antibody titers were determined by an endpoint dilution

ELISA on micro-well plates coated with inactivated SARS-CoV-2 viral culture (30

ng/well). Briefly, sera serially diluted in phosphate-buffered saline (PBS) were added into

micro-well plates (100 μL/well) and incubated at 37°C for 1 h. Goat anti-mouse IgG

antibody (HRP conjugate) (Sigma-Aldrich, SL, USA) was used as the detection antibody

(100 μL/well). Color reaction was developed with 3, 3', 5, 5'-tetramethylbenzidine (TMB)

and terminated with H_2SO_4 solution. The absorbance was read at 450 nm with an ELISA

reader (Thermo Fisher, Mannheim, Finland). Antibody titers were determined as the

highest dilution at which the mean absorbance of the sample was 2.1-fold greater than

that of the control serum.

Evaluation of protective efficiency of SARS-CoV-2 subunit vaccine

The neutralization antibody titers were detected by a traditional virus neutralization

assay using SARS-CoV-2. The assay was performed in quadruplicate, and a series of

eight two-fold serial dilutions of the plasma or serum were assessed. Briefly, 100 tissue

culture infective dose 50 ($TCID_{50}$) units of SARS-CoV-2 were added to two-fold serial

dilutions of heat-inactivated serum (60°C, 30 min), and incubated for 1 h at 37°C. The

virus – serum mixture was added to Vero E6 cells (3×10^5/well) grown in an ELISpot

plate and incubated for 3 d. The endpoint of the micro-neutralization assay was

designated as the highest serum dilution at which all three, or two of three, wells were not

protected from virus infection, as assessed by visual examination. The neutralizing titers

of the dilutions of sera were measured with the maximum dilution achieving the median

neutralization[33].

Investigation of cell-mediated immune response by IFN-γ and IL-4 ELISpot assay

Mouse IFN-γ and IL-4 ELISpot assay were conducted according to the

manufacturer's guidelines (DaKeWe, Shenzhen, China). Briefly, splenocytes (3×10^5 per

well, in triplicate) of immunized mice at Day35 were plated into ELISpot plate and

stimulated with PHA (Positive control) or S1-4 protein. After incubation for 16 h in a

37°C humidified incubator with 5% CO_2, the cells were removed before adding the

detection antibody (biotin conjugate). Two hours later at room temperature (RT),

Streptavidin-HRP was added after washing plate. One hour later at RT, TMB substrate

was applied until distinct spots emerged. Count spots in an ELISpot reader after stopping

color development with deionized water.

Statistical analysis

Antibody titers were log-transformed to establish geometric mean titers (GMT) for

analysis. The number of specific IFN-γ- or IL-4-secreting T cells were expressed as spots

forming cells (SFCs) per 3×10^5 cells. The related data were statistically analyzed by

Mann-Whitney U test for non-parametric data and the paired t test for parametric data.

SPSS software (22.0, Chicago, IL, USA) and GraphPad Prism software (8.00, San Diego,

CA, USA) was applied to analyze the data and draw graphics, respectively. Significance

was indicated when value of $P < 0.05$.

Acknowledgements

This work is supported by a project funded by Denogen (Beijing) Biotechnology Co., Ltd.

We thank Mrs Jianrong Cao especially for the funding. The funder has no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

S.B., G.Z., G.F.G. and J.Z. conceived, designed and supervised the research. Q.S., Y. Z.,

Y.Y., L.S., X.Y., J.S., F.Q., P.Y., W.Z. and L.C. performed the experiments. L.S. and Y.

Z analyzed the data. H.W., H.K., K.H. and B.C. contributed with discussion of the results.

Q.S. and S.B. wrote the paper. All authors read, revised and approved the manuscript.

Competing interests

The authors declare that they have no known competing financial interests or personal

relationships that could have influenced the work reported in this paper.

References

1. Madabhavi, I., Sarkar, M. & Kadakol, N. COVID-19: a review. *Monaldi archives*

for chest disease = Archivio Monaldi per le malattie del torace **90** (2020).

2. World Health Organization. Coronavirus disease (COVID-19) Situation Report -

205, https://www.who.int/docs/default-source/coronavirus/situation-

reports/20200812-covid-19-sitrep-205.pdf?sfvrsn=627c9aa8_2; 2020 [accessed

12 August 2020].

3. Zhu, N. et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019.

The New England journal of medicine **382**, 727-733 (2020).

4. Velavan, T.P. & Meyer, C.G. The COVID-19 epidemic. *Tropical medicine &*

international health : TM & IH **25**, 278-280 (2020).

5. Zhou, P. et al. A pneumonia outbreak associated with a new coronavirus of

probable bat origin. *Nature* **579**, 270-273 (2020).

6. Lu, R. et al. Genomic characterisation and epidemiology of 2019 novel

coronavirus: implications for virus origins and receptor binding. *Lancet (London,*

England) **395**, 565-574 (2020).

7. Chen, W.H., Strych, U., Hotez, P.J. & Bottazzi, M.E. The SARS-CoV-2 Vaccine

Pipeline: an Overview. *Current tropical medicine reports*, 1-4 (2020).

8. Su, Q.D. et al. The biological characteristics of SARS-CoV-2 spike protein

Pro330-Leu650. *Vaccine* **38**, 5071-5075 (2020).

9. Ludwig, S. & Zarbock, A. Coronaviruses and SARS-CoV-2: A Brief Overview.

Anesthesia and analgesia **131**, 93-96 (2020).

10.   Hoffmann, M. et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2

and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e278

(2020).

11.   Shang, J. et al. Structural basis of receptor recognition by SARS-CoV-2. *Nature*

581, 221-224 (2020).

12.   Prompetchara, E., Ketloy, C. & Palaga, T. Immune responses in COVID-19 and

potential vaccines: Lessons learned from SARS and MERS epidemic. *Asian*

Pacific journal of allergy and immunology **38**, 1-9 (2020).

13.   Sarkar, B., Ullah, M.A., Johora, F.T., Taniya, M.A. & Araf, Y.

Immunoinformatics-guided designing of epitope-based subunit vaccines against

the SARS Coronavirus-2 (SARS-CoV-2). *Immunobiology* **225**, 151955 (2020).

14.   Wang, D. et al. Immunoinformatic Analysis of T- and B-Cell Epitopes for SARS-

CoV-2 Vaccine Design. *Vaccines* **8** (2020).

15.   Noorimotlagh, Z. et al. Immune and bioinformatics identification of T cell and B

cell epitopes in the protein structure of SARS-CoV-2: A systematic review.

International immunopharmacology **86**, 106738 (2020).

16.   He, Q. et al. Immunogenicity and safety of a novel tetanus toxoid-conjugated anti-

gastrin vaccine in BALB/c mice. *Vaccine* **36**, 847-852 (2018).

17.   Lise, L.D. et al. Enhanced epitopic response to a synthetic human malarial peptide

by preimmunization with tetanus toxoid carrier. *Infection and immunity* **55**, 2658-

2661 (1987).

18.   Mangsbo, S.M. et al. Linking T cell epitopes to a common linear B cell epitope: A

2   targeting and adjuvant strategy to improve T cell responses. *Molecular*

3   *immunology* **93**, 115-124 (2018).

4   19.   Moriyama, H. et al. Induction and acceleration of insulitis/diabetes in mice with a

5   viral mimic (polyinosinic-polycytidylic acid) and an insulin self-peptide.

6   *Proceedings of the National Academy of Sciences of the United States of America*

7   **99**, 5539-5544 (2002).

8   20.   Cai, H. et al. Self-adjuvanting synthetic antitumor vaccines from MUC1

9   glycopeptides conjugated to T-cell epitopes from tetanus toxoid. *Angewandte*

10   *Chemie (International ed. in English)* **52**, 6106-6110 (2013).

11   21.   Abu-Zeid, Y.A. et al. Seroprevalence of antibodies to repetitive domains of

12   Plasmodium vivax circumsporozoite protein in United Arab Emirates children.

13   *Transactions of the Royal Society of Tropical Medicine and Hygiene* **96**, 560-564

14   (2002).

15   22.   Boitel, B. et al. Strong similarities in antigen fine specificity among DRB1* 1302-

16   restricted tetanus toxin tt830-843-specific TCRs in spite of highly heterogeneous

17   CDR3. *Journal of immunology (Baltimore, Md. : 1950)* **154**, 3245-3255 (1995).

18   23.   Zhu, F.C. et al. Safety, tolerability, and immunogenicity of a recombinant

19   adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label,

20   non-randomised, first-in-human trial. *Lancet (London, England)* **395**, 1845-1854

21   (2020).

24. Chen, W.H. et al. Yeast-Expressed SARS-CoV Recombinant Receptor-Binding Domain (RBD219-N1) Formulated with Alum Induces Protective Immunity and Reduces Immune Enhancement. *bioRxiv : the preprint server for biology* (2020).

25. Yang, J. et al. A vaccine targeting the RBD of the S protein of SARS-CoV-2 induces protective immunity. *Nature* (2020).

26. Yuan, M. et al. A highly conserved cryptic epitope in the receptor binding domains of SARS-CoV-2 and SARS-CoV. *Science (New York, N.Y.)* **368**, 630-633 (2020).

27. Zhang, J. et al. Long-term efficacy of a hepatitis E vaccine. *The New England journal of medicine* **372**, 914-922 (2015).

28. Hu, Y.M. et al. Safety of an Escherichia coli-expressed bivalent human papillomavirus (types 16 and 18) L1 virus-like particle vaccine: an open-label phase I clinical trial. *Human vaccines & immunotherapeutics* **10**, 469-475 (2014).

29. Vankadari, N. & Wilce, J.A. Emerging WuHan (COVID-19) coronavirus: glycan shield and structure prediction of spike glycoprotein and its interaction with human CD26. *Emerging microbes & infections* **9**, 601-604 (2020).

30. Yi, Y. et al. Cloning and expression of SARS coronavirus spike gene fragment and its application. *Chinese Journal of Virology* **19**, 267-268 (2003).

31. Mosmann, T.R. & Coffman, R.L. TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annual review of immunology* **7**, 145-173 (1989).

32. Su, Q.D. et al. Intranasal vaccination with ebola virus GP amino acids 258-601 protects mice against lethal challenge. *Vaccine* **36**, 6053-6060 (2018).

1 33.   He, Y. et al. Receptor-binding domain of SARS-CoV spike protein induces highly

2   potent neutralizing antibodies: implication for developing subunit vaccine.

3   *Biochemical and biophysical research communications* **324**, 773-781 (2004).

Figure Legends

Figure 1. The preparation of S1-4 protein

(A). Small-scale expression of S1-4 protein. (B). Large-scale expression of S1-4 protein. (C). S1-

4 protein purified by ion-exchange chromatography (IEX). (D). S1-4 protein purified by

hydrophobic interaction chromatography (HIC) and renaturation.

1. Negative control; 2. Non-induced expression bacteria; 3 – 4, Induced expression bacteria; 5.

Total bacterial proteins after large-scale expression; 6. Soluble fraction of bacterial sonication; 7.

Insoluble aggregates of bacterial sonication; 8. S1-4 protein in Urea solution; 9. Unbound

fraction eluted by IEX; 10. S1-4 protein fraction eluted by HIC; 11. Re-natured S1-4 protein after

stepwise dialysis.

Figure 2. Western Blotting analysis of S1-4 protein

(A). Silver staining after SDS-PAGE. (B). Convalescent serum of COVID-19 patients as the

primary antibody. (C). Rabbit monoclonal antibody against RBD as the primary antibody. (D).

Rat polyclonal antibody against inactivated SARS-CoV-2 as the primary antibody.

1. S1-4 protein; 2. Commercial recombiant RBD protein.

Figure 3. Kinetics of anti-SARS-CoV-2 antibody and neutralization antibody in sera (A)

and IL-4 & IFN-γ spot-forming cells (SFCs) in splenocytes of immunized mice (B)

Figure 4. Schematic diagram of the gene of SARS-CoV-2 subunit antigen (S1-4 and S1-5) in

the pET28a expression vector

SARS-CoV-2 subunit candidate vaccine