DNA extraction and host depletion methods significantly impact and potentially bias bacterial detection in a biological fluid

Erika Ganda[a,b#*], Kristen L. Beck[b,c], Niina Haiminen[b,d], Justin D. Silverman[e,f,g], Ban Kawas[b,c], Brittany Cronk[h], Renee R. Anderson[h], Laura B. Goodman[h], and Martin Wiedmann[a,b]

[a] Department of Food Science, Cornell University, Ithaca, NY, USA

[b] Consortium for Sequencing the Food Supply Chain, San Jose, CA, USA

[c] IBM Almaden Research Center, San Jose, CA, USA

[d] IBM T.J. Watson Research Center, Yorktown Heights, NY, USA

[e] College of Information Sciences and Technology, Penn State University, University Park, PA, USA

[f] Department of Statistics, Penn State University, University Park, PA, USA

[g] Department of Medicine, Penn State College of Medicine, Hershey, PA, USA

[h] Department of Population Medicine and Diagnostic Sciences, College of Veterinary Medicine, Cornell University, Ithaca, NY, USA

Running Head: Nucleic acid extraction and host depletion in milk

[#] Address correspondence to Erika Ganda, ganda@psu.edu.
[*] Present address: Department of Animal Science, The Pennsylvania State University, State College, PA, USA

Abstract word count: Abstract: **250**; Importance: **147**

Text word count: 4180 (excluding, methods, references, table footnotes, and figure legends)

ABSTRACT

Abstract: Untargeted sequencing of nucleic acids present in food can inform the detection of food safety and origin, as well as product tampering and mislabeling issues. The application of such technologies to food analysis could reveal valuable insights that are simply unobtainable by targeted testing, leading to the efforts of applying such technologies in the food industry. However, before these approaches can be applied, it is imperative to verify that the most appropriate methods are used at every step of the process: gathering primary material, laboratory methods, data analysis, and interpretation.

The focus of this study is in gathering the primary material, in this case, DNA. We used bovine milk as a model to 1) evaluate commercially available kits for their ability to extract nucleic acids from inoculated bovine milk; 2) evaluate host DNA depletion methods for use with milk, and 3) develop and evaluate a selective lysis-PMA based protocol for host DNA depletion in milk.

Our results suggest that magnetic-based nucleic acid extraction methods are best for nucleic acid isolation of bovine milk. Removal of host DNA remains a challenge for untargeted sequencing of milk, highlighting that the individual matrix characteristics should always be considered in food testing. Some reported methods introduce bias against specific types of microbes, which may be particularly problematic in food safety where the detection of Gram-negative pathogens and indicators is essential. Continuous efforts are needed to develop and validate new approaches for untargeted metagenomics in samples with large amounts of DNA from a single host.

Importance: Tracking the bacterial communities present in our food has the

potential to inform food safety and product origin. To do so, the entire genetic material

present in a sample is extracted using chemical methods or commercially available kits

and sequenced using next-generation platforms to provide a snapshot of what the

relative composition looks like. Because the genetic material of higher organisms

present in food (e.g., cow in milk or beef, wheat in flour) is around one thousand times

larger than the bacterial content, challenges exist in gathering the information of

interest. Additionally, specific bacterial characteristics can make them easier or harder

to detect, adding another layer of complexity to this issue. In this study, we demonstrate

the impact of using different methods in the ability of detecting specific bacteria and

highlight the need to ensure that the most appropriate methods are being used for each

particular sample.

Keywords: Host Depletion, Shotgun Metagenomics, Milk, DNA, RNA, Biases,

Propidium Monoazide, Dairy, Low Biomass, Food Microbiome

INTRODUCTION

In the past decade, recent developments in molecular methods, including high throughput sequencing (**HTS**) technologies, have demonstrated the feasibility of sequencing-based analysis of various foods and food-associated environments, and the potential for application in informing food safety practices (1), product processing methods (2) and ingredient authentication (3, 4). This has spurred the efforts to translate the use of such technologies to the industry setting with the objective of moving food safety testing to the next frontier (5). However, before such refined approaches can be reliably applied in the industry, it is imperative that appropriate methods are employed at every step of the process: gathering primary material, laboratory methods, data analysis, and interpretation. In this study, we highlight the importance of considering the food matrix characteristics as well as the laboratory methodologies applied. Bovine milk is used as a model with to demonstrate the several challenges associated with developing HTS methods for use in food matrices.

Milk, similar to many other foods, is a chemically complex biological fluid. It contains several compounds that can hamper the chemistry involved in DNA and RNA extraction (6, 7) and act as PCR inhibitors, such as calcium ions, fats, and proteins (8, 9). Additionally, lactoferrin, an enzyme present in bovine milk, has recently been described to have both DNase and RNase activity (10). Another challenge for untargeted sequencing applications of bovine milk is the presence of bovine somatic cells. The bovine genome is one thousand times larger than an average bacterial genome (Bovine: 2.7 Gb, Bacteria: average 3.6 Mb, median 3.4 Mb (11, 12)). Thus, even when present in much smaller amounts than bacterial cells, bovine somatic cells

introduce an enormous quantity of typically unwanted host nucleic acids in untargeted

HTS studies. Realistically, however, high-quality raw milk may contain around 200,000

bovine somatic cells and 20,000 or less bacterial cells per mL, leading to a 10,000-fold

higher abundance of bovine DNA in high quality raw milk.

Despite the challenges associated with nucleic acid extraction from milk and the

amount of host DNA present, several investigations have successfully used milk (13–

15) or other dairy products (16) as their sample of interest in targeted and untargeted

HTS studies, highlighting the potential for application of HTS technologies in food

production settings. Total RNA sequencing can be more informative than untargeted

DNA sequencing as it has the potential to provide gene expression (e.g. toxin

production) in addition to taxonomic relative abundance in a community of food-

associated microorganisms (4, 17). When compared to DNA extraction, RNA extraction

is a more complex and challenging process, given the short half-life of RNA compared

to DNA, its inherent susceptibility to degradation, and the known presence of RNases in

milk (10). Nevertheless, a number of studies have been successful in extracting RNA

from dairy products highlighting the potential of RNA based techniques for food safety

and quality surveillance (18–20).

When the ultimate objective is to use HTS to create a tool that can be used to

inform food safety and quality in industry settings, understanding the impact and biases

introduced by using protocols that have not yet been tested or optimized for a given

food matrix is of great importance. In addition, to be potentially adopted in industry

settings, laboratory protocols should be performed in a timely manner. It is important to

characterize the entire food sample, and most protocols available for nucleic acid

extraction in milk begin with a centrifugation and fat-removal step, which can in itself

introduce significant bias to the final result. For example, bacterial spores have been

described to aggregate in the fat layer of milk samples subjected to gravity separation

(21, 22), and RNA yield has been shown to be variable among different milk fractions

(23). Unfortunately, these potential introductions of bias are overlooked in most

investigations.

While a number of studies have investigated commercially available protocols for

nucleic acid extraction of milk in terms of DNA concentration, quality, and ability to

detect specific pathogens (24–26), their main objective was not the assessment of

differential DNA extraction nor biases in the representation of diverse bacterial

populations, which would require the inclusion of mock bacterial communities of interest

into a milk sample and comparison of several protocols.

Host DNA contamination is a challenge not exclusive to milk and other food

samples, as mammalian DNA has been shown to dominate the number of sequencing

reads in cerebrospinal fluid, skin, vaginal, and oral metagenomes in humans(27–29). To

tackle the host DNA issue, enzymatic and immunomagnetic protocols aimed at

decreasing host DNA contamination became commercially available and have been

tested in select sample types (27, 28, 30). In addition, a number of "homebrew"

methods have been tested to allow for successful depletion of host DNA (27, 31, 32).

Nevertheless, these methods are not guaranteed to work in every sample type, and to

the best of our knowledge, no host DNA depletion methods have been evaluated for

their applicability in milk.

The goals of this study were 1) to evaluate commercially available kits for their

ability to extract nucleic acids from bovine milk; 2) to evaluate host DNA depletion

methods for use with bovine milk, and 3) to develop and evaluate a selective lysis-PMA

based protocol for host DNA depletion in milk. Our overarching hypothesis was that

methodologies would differ regarding the efficacy of nucleic acid extraction, and

potential biases would be observed. Experiments were thus performed on raw bovine

milk inoculated with mock bacterial communities, which included Gram-negative

(*Salmonella enterica*), Gram-positive (*Listeria monocytogenes*), *Mycobacteria* (*M.*

smegmatis) as well as spores representing aerobic sporeformers (*Bacillus wiedmannii*).

METHODS

Comparison of Nucleic Acid Extraction Protocols in Spiked Milk

Strain Selection

We selected a Gram-positive (*L. monocytogenes*), a Gram-negative (*S. enterica*), and a spore former (*B. wiedmanii*) previously isolated from milk or the dairy environment to create a mock microbial community that would be inoculated into raw milk. We included an additional Gram-positive bacterium, a member of the *Mycobacteriaceae* family (*M. smegmatis*), because of the potential public health implications and the uniqueness of the cell structure of *Mycobacteria*. Milk samples were specifically inoculated with *Salmonella enterica*, *Listeria monocytogenes,* and *Mycobacterium smegmatis* vegetative cells grown to stationary phase*,* and *Bacillus wiedmannii* spores. Specific strain information is available in Table 1 and in the Food Microbe Tracker Database (www.foodmicrobetracker.com). Bacterial load of the inoculated milk sample was chosen to represent the largest bacterial concentration allowed for raw milk in the United States, which is 300,000 colony forming units **(CFUs)** per mL, in an attempt to simulate the highest legal bacterial load of incoming milk in a dairy processing plant(33).

Inoculum Preparation

Bacillus wiedmannii spore suspension was prepared according to (34). Briefly, the bacterial isolate was streaked from frozen culture into Brain Heart Infusion (**BHI**) agar (Becton, Dickinson and Co., Sparks, MD) and incubated for 24h at 37°C. Following incubation, a single colony was selected to inoculate a tube containing 5 mL of BHI broth followed by incubation at 37°C for 72h. Next, 100 µL of inoculated BHI was spread plated in duplicate on sporulating media AK Agar #2 (Becton, Dickinson and Co.), which

was incubated for 120h at 37°C. Sporulation was confirmed via microscopy with a 7.5%

malachite green endospore stain (J.T. Baker, Phillipsburg, NJ) as detailed in (35).

Spores were harvested by flooding the agar surface with 10 mL of phosphate-buffered

saline (**PBS**) (Weber Scientific, Hamilton, NJ) and scraping the bacterial culture with a

cell scraper. Harvested cells were transferred to a sterile centrifuge tube and washed

with 10 mL of sterile water three times by centrifugation at 10,500 rpm for 15 minutes

and resuspension of the pellet. Following the third wash, 5 mL of sterile water and 5 mL

of 100% ethanol (Decon Labs, King of Prussia, PA) were added to the tube and the

pellet was resuspended by vortexing. The bacterial pellet resuspended in 50% ethanol

was incubated for 12h at 4°C in a rotating platform to eliminate any remaining

vegetative cells. After ethanol treatment, the spore suspension was washed another

three times,with 10 mL of sterile water as described above. The final spore suspension

was kept in sterile water at 4°C until used for spiking experiments.

Mycobacterium smegmatis cells were streaked from frozen stocks into BHI

agar, followed by incubation at 37°C for 48h. A single colony was used to inoculate a 5

mL tube containing BHI broth with 1% Tween 80 (Fisher Scientific, Hampton, NH, USA)

followed by incubation at 37°C for 48h. The final inoculum was prepared by inoculating

100 µL of liquid culture onto 100 mL of pre-warmed BHI broth with 1% Tween 80,

followed by incubation at 37°C for 72h. The resulting stationary phase culture was kept

at 4°C until spiking.

Listeria monocytogenes and *Salmonella enterica* cells were streaked

separately from frozen stocks into BHI agar, followed by incubation at 37°C for 24h. For

each strain, a single colony was used to inoculate a 5 mL tube containing BHI broth

followed by incubation at 37°C for 24h. The final inoculums were prepared by

inoculating 100 µL of each liquid culture onto 100 mL of pre-warmed BHI broth that was

subsequently incubated at 37°C for 12h.

At harvesting, bacteria were spiral plated on agar using an Eddy Jet 2W Spiral

plater (IUL micro, Barcelona, Spain) at various dilutions to determine bacterial

concentrations. Bacterial liquid cultures were kept at 4°C until bacterial enumeration (*B.*

wiedmanii, *L. monocytogenes*, and *Salmonella* sp. were kept for 24h and *M. smegmatis*

was kept for 48h at 4°C). Inoculum volumes were calculated based on CFUs.

Inoculation of Raw Milk

Raw milk was collected from the Cornell University Ruminant Center (CURC –

Hartford, NY) bulk tank into sterile 10 oz lock tab containers (Capitol plastics,

Amsterdam, NY) and transported on ice to the Milk Quality Improvement Program

laboratory in Ithaca, NY. Samples were combined into a single sterile 1000 mL glass

bottle and homogenized by inverting the container 50 times. From that bottle, one

aliquot (1 mL) was used to make dilutions and determine the initial bacterial count using

Standard Plate Count (**SPC**) agar (Milipore Sigma, Burlington, MA), that was incubated

for 48h at 32°C. Bacterial enumeration was performed using an automated colony

counter (SphereFlash® – Automatic Colony Counter, IUL micro, Barcelona, Spain). A

second aliquot (60 mL) was transported on ice to the DairyOne laboratory (DairyOne,

Ithaca, NY) for determination of somatic cell counts (**SCC**) and milk contents (fat,

protein, lactose, and total solids). The remaining milk was used for inoculation with the

strains described above. To allow for interaction between inoculated bacteria and milk

components, and to mimic conditions similar to when raw milk is stored in dairy silos,

inoculated milk was held at 4°C for 24h prior to use as starting sample for nucleic acid

extraction comparisons.

Bacterial Enumeration of Inoculated Milk Samples

Total bacterial enumeration was performed through serial dilutions of each milk

sample in PBS that was spiral plated in duplicate on standard plate count agar as

described above and incubated for 48h at 32°C.

Nucleic Acids Extraction

Raw (uninoculated) milk samples were processed in parallel with inoculated milk

samples for all kits. Additionally, a no template nucleic acid extraction was carried out

as a negative control to assess cross-contamination. As a positive control, a mock

bacterial community was created in PBS with the same bacteria described above at the

same concentration as the inoculated milk samples; this was included on each

extraction plate or run as a control. Inoculated milk samples were extracted in three

technical replicates in each of three independent biological replicates, which were

performed on separate days with a different raw milk sample used for each biological

replicate.

Kits, manufacturers, and protocol details are described in Table 2. For each of

the nine protocols evaluated, samples were processed in parallel (uninoculated milk,

inoculated milk triplicates, mock bacterial community in PBS, and a negative control

without a starting sample "kit buffers only") according to the manufacturer's instructions.

Exact versions of protocols followed can be found in a GitHub repository as

supplementary materials (https://github.com/ErikaGanda/MilkDNA). Extracted nucleic

acids were frozen at -80°C until quantification and qPCR assays were performed.

Nucleic Acids Quantification

Nucleic acid quantification was performed with both a spectrophotometer

(Nanodrop 2000, ThermoFisher Scientific) and a fluorescence-based method.

Absorbance was measured at 280, 260, and 230 nm for DNA and RNA. Total DNA was

also measured with a Quant-iT™ dsDNA High-Sensitivity (HS) Assay Kit (ThermoFisher

Scientific); fluorescence measurements were performed using a Synergy H1 plate

reader (BioTek Instruments, Winooski, VT, USA) with wavelengths of 490 nm for

excitation and 535 nm for emission.

qPCR Primer Development and Assay Conditions

For each bacterial strain used to inoculate milk samples, qPCR primers were

designed to target the RNA polymerase subunit beta gene (*rpoB*) because it is a single-

copy gene and allows for a more accurate comparison between bacterial numbers than

16S rDNA. Primers were also designed to target a conserved region of the 16S gene,

and calculated 16S copy numbers were used as a proxy for total bacterial numbers.

Primer details are described in Table 3. Reactions were carried out in duplicate using 2

μL of extracted DNA. The final qPCR reaction volumes totaled 25 μL containing 12.5 μL

SYBR green master mix (ThermoFisher Scientific), 2 μL extracted DNA, 0.5 μM forward

primer, 0.5 μM reverse primer, and 9.5 μL nuclease-free water. Reactions were carried

out in a Quantstudio 6 instrument (ThermoFisher Scientific), with the following cycling

conditions: 95°C for 10 minutes, followed by 40 cycles of 95°C for 15 seconds and 60°C

for 1 minute, and a melting curve: 95°C for 15 seconds, 60°C for 1 minute, and 95°C for

15 seconds.

Bovine genome copy numbers were calculated using a commercial TaqMan

assay targeting the UXT gene (ThermoFisher Scientific). The final qPCR reaction

totaled 20 µL, and included 1 µL of gene expression assay, 10 µL TaqMan Fast

Advanced Master Mix (ThermoFisher Scientific), 7 µL of nuclease-free water, and 2 µL

of template. Reactions were carried out in a Quantstudio 6 instrument, with the following

cycling conditions: 95°C for 20 seconds, followed by 40 cycles of 95°C for 1 second and

60°C for 20 seconds. For each of the three biological replicates, samples extracted with

all kits were amplified in a single PCR plate and compared using the same standard

curve.

qPCR Data Analysis

Amplification data were exported from QuantStudio Real-Time PCR Software

(ThermoFisher Scientific) into Excel (version 16.0.11325.20156, Microsoft Corp.,

Redmond, WA). Standard curves were built with serial dilutions of purified bacterial

DNA that was quantified using a Nanodrop spectrophotometer (ThermoFisher) and

genome equivalents in each reaction were calculated as described in (36). Standard

curves with an R2 < 0.9 and efficiency < 70% were discarded and reactions were

repeated. For 2 out of 24 reaction plates, one outlier was removed based on visual

inspection of deviating standard curve data points.

Copies per µL of DNA input were calculated for each reaction. Melt curves were

visually inspected and cycle threshold (**CT**) values of samples in which melt curves did

not match standard samples' peaks were not used for final copy number calculations.

Because different kits required varying amounts of sample input and nucleic acid output

(Table 2), data was normalized to allow for comparison between kit protocols. We chose

to simulate copy numbers in 1 mL of milk input and 100 µL DNA output for each kit,

using the following equations:

$$\text{MilkEquivalents}=\frac{input\ milk\ sample\ (µL)}{eluted\ DNA\ (µL)}$$

$$\text{Copies/mLofMilk}=\frac{copies\_µL\_DNA*1000}{MilkEquivalents}$$

Normalized copy number data were log-transformed prior to statistical analysis.

The final database was cross-referenced and checked with the open-source software

openrefine (https://github.com/OpenRefine). All statistical analyses with inoculated milk

PCR data were performed in R (version 3.4.3; R project, Vienna, Austria).

Statistical Model Selection

Linear models were fit using the "lm" function in R to compare log copy numbers

between kits. For each assay, three linear models were compared; including (i)

experiment number (SpikeSet) only, (ii) including qPCR efficiency only, and (iii) both (i)

and (ii) as covariates. Given that only three experiments were performed, experiment

number was treated as a fixed effect rather than a random effect. To account for

different variances observed across kits, models not assuming homoscedasticity were

also fit using generalized least-squares using the nlme (37) package in R. Models were

inspected by Q-Q plots and residual vs. predicted plots. The model with best fit as

determined by smallest AIC was used as the final model. Estimated marginal means of

each model were calculated using the emmeans (38) package and are depicted in

graphs overlaid to scatterplots of original datapoints. Raw data and code are available

on a GitHub repository: https://github.com/ErikaGanda/MilkDNA.

Host Depletion Protocols

Selective Osmotic Lysis of Host Cells and Host DNA Depletion through Propidium

Monoazide (PMA) Treatment of Uninoculated Raw Milk

An osmotic lysis-based host DNA depletion protocol was adapted from (27).

Briefly, 500 µL of uninoculated milk were centrifuged at 10,000g for 8 minutes, whey

was discarded while fat and pellet were kept in the microcentrifuge tube. Five hundred

microliters of sterile double distilled water (**ddH$_2$O**) were added to the pellet. The tube

was vortexed until pellet dissolution followed by incubation at room temperature for 5

minutes to allow for osmotic lysis of mammalian cells. Four different concentrations of

PMA (Biotium, Hayward, CA, catalog # 40019) were compared: 10, 20, 40, and 50µM.

Untreated milk (milk that was not centrifuged or exposed to osmotic lysis) and milk

exposed to osmotic lysis but not exposed to PMA were also included in comparisons.

To account for biological variation and assess repeatability, experiments were

performed in three biological replicates performed on three different days, using a

different raw milk sample for each biological replicate.

After incubation at room temperature, the appropriate volume of PMA was added

to each tube to achieve desired concentrations, followed by briefly mixing, and

incubation in the dark (in an aluminum-foil wrapped box) for 5 minutes on a rotating

platform.

Following PMA incubation in the dark, PMA was inactivated by light exposure.

Samples were placed horizontally on ice, < 20 cm from a 500W halogen light source

(Woods Halogen Work Light, Southwire Company LLC, Carrollton, GS, USA). Samples

were exposed to light on a rotating platform for 5 minutes, and frozen at -80°C until DNA

extraction with a magnetic-based method (MagMAX CORE Nucleic Acid Purification Kit

298 – ThermoFisher). Quantitative PCR reactions and data analysis were performed as

299 described above, with the exception that only bovine DNA and total bacterial copy

300 numbers were quantified, as no bacteria was inoculated in the milk samples used.

301 Linear models were fitted to assess the effect of selective lysis in non-PMA treated

302 samples, (NTC versus 0µM) and the effect of PMA concentration on copy numbers.

Selective Enzymatic Lysis of Host Cells and Host DNA depletion through PMA

Treatment in Inoculated Raw Milk

Because milk has more fat, protein, and minerals then saliva we hypothesized

that the 1:1 osmotic lysis included in the protocol adapted from (27) was not optimal for

lysing bovine cells in milk compared to human cells in saliva.

We thus also evaluated a combination of a mild lysis solution followed by

incubation with two concentrations of subtilisin, a protease from *Bacillus licheniformis*

(Krackeler Scientific, cat# 45-P5380-25MG). The lysis solution contained 7.6 g/L sodium

carbonate, 8.8 g/L sodium bicarbonate, 2.43 g/L disodium EDTA, and 2.71 g/L

tetrasodium EDTA. Reagents were solubilized in 1L of sterile double distilled water

using a stir plate with a magnetic rotating bar for 6 hours, prior to pH measurement and

titration to 9.5, and filter-sterilization with a 0.22 µm filter (Corning Disposable Vacuum

Filter – Fisher Scientific, Waltham, MA).

Experiments were performed in two separate days with a newly collected milk

sample in each day. To access potential lysis biases, we spiked milk with a Gram-

positive and a Gram-negative bacterium as described above (*L. monocytogenes* and

Salmonella sp.). Spiked milk samples were treated prior to DNA extraction as follows:

Four hundred microliters of inoculated milk were added to 1.6 mL of lysis solution and

incubated at room temperature for 5 minutes prior to the addition of subtilisin. Two

concentrations of subtilisin were tested: 10 mg/mL (10x) and 1 mg/mL (1x) – to achieve

final concentrations of 0.73 and 0.073 units per reaction, respectively. After adding the

enzyme, samples were incubated at 50°C for 5 minutes and placed on ice after

incubation. Samples were then centrifuged at 10,000g for 8 minutes. The supernatant

containing lysis solution and enzyme was discarded, and the pellet was resuspended

with 400µL of PBS. Treated samples were exposed to PMA at 20 µM as described

above, photo-activated using two light sources (Halogen lamp described above or BLU-

V System – Qiagen) and stored at -80°C until DNA extraction with a magnetic-based

method (MagMAX CORE extraction kit – ThermoFisher).

Quantitative PCR reactions and data analysis were performed as described

above, with the exception that Bovine DNA, Total bacterial copy numbers, *L.*

monocytogenes, and *Salmonella* were quantified, as we hypothesized that the lysis

solution could affect Gram-negative bacteria differently than Gram-positive bacteria.

Comparison of Host DNA Depletion Methods in Uninoculated Raw Milk

Based on initial experiments with various PMA concentrations we decided to use

20µM to include in a comparison with three commercial host DNA depletion kits. Kits,

manufacturers, and protocol details are described in Table 4. Samples were processed

according to the manufacturer's instructions and frozen at -80°C.

For the NEB host depletion method, DNA was extracted with a magnetic-based

extraction and quantified using a qubit fluorometer (ThermoFisher) prior to methylated

host DNA capture reactions. Final microbial enriched DNA cleanup was performed

using AMPure magnetic beads (Beckman Coulter). Exact versions of protocols followed

can be found in our GitHub repository https://github.com/ErikaGanda/MilkDNA. Three

biological replicates (each with two technical replicates) were prepared using a new

sample of raw milk for each biological replicate (with the exception of MolZym kit, which

was only performed with duplicate milk samples in one day due to limitation in available

reagent amounts).

Quantitative PCR reactions and data analysis were performed as described

above, with the exception that only bovine DNA and total bacterial copy numbers were

quantified, as no bacteria was inoculated in the milk samples used.

Sequencing

In addition to qPCR, we performed deep untargeted sequencing of a subset of

nine samples at the Cornell Biotechnology Resource Center. Briefly, DNA quality control

was performed in a fragment analyzer and sequencing libraries were constructed using

a Nextera DNA Flex kit (now renamed to Illumina DNA Prep, Illumina, San Diego, CA)

and 2x150 BP paired-end sequencing was performed in an Illumina NextSeq500.

Sequencing data quality control and in silico host signal removal was performed as

previously described (3, 4), and microbial data was processed as described in Beck et

al. 2020 (4). Briefly, adapter removal and quality trimming was performed with

TrimGalore (39) and trimmed reads of at least 50bp in length were classified using

Kraken v0.3 (40) against a multi-eukaryotic database of 31 common food ingredients

and contaminants including the bovine reference genome (GCF_000003205.7,

Btau_5.0.1) for bioinformatic removal of the host signal. Then taxonomic profiling of

samples was completed using Kraken v0.3 against microbial RefSeq genomes as

described in Beck et. al., 2020 (4). Principal Component Analysis (PCA) and Robust

PCA (41) were performed in pseudocounted Isometric Log-Ratio transformed microbial

reads (4).

Data availability

Raw data and code are available on our GitHub repository:

https://github.com/ErikaGanda/MilkDNA. Raw sequencing reads have been deposited

into SRA under the submission number SUB7938697.

RESULTS

Extraction Method Significantly Impacts Bacterial Quantification through qPCR

We assessed seven commercially available DNA extraction methods on their ability to isolate DNA from milk samples inoculated with a mock bacterial community (for milk sample characteristics see Supplementary Table S1). Nucleic acid quantification and quality measurements (Supplementary Table S2) need to be interpreted with caution due to the low overall DNA yield (<10 ng/µL for all samples), which placed readings below the linear range for most samples. However, all kits yielded sufficient DNA to allow for qPCR-based quantification of the members of the bacterial mock community (i.e., *Bacillus wiedmanii*, *Listeria monocytogenes*, *Mycobacterium smegmatis*, and *Salmonella* sp.) as well as total bacterial 16S rDNA copies and bovine DNA. Quantitative PCR data (Figure 1) demonstrated that the two magnetic-based DNA extraction methods (COREDNA and Mastitis, the two leftmost methods in Fig. 1) always yielded numerically higher copy numbers (for all members of the mock community and for total bacterial 16S rDNA) than the other five kits; these values were generally significantly higher (except for *L. monocytogenes*, where values for one non-magnetic kit did not differ significantly from the two magnetic kits). Bacterial copy numbers for the two magnetic-based kits also had lower variability between technical replicates than column-based kits, with E.Z.N.A. Food DNA kit showing particularly large variability between technical replicates; DNA yield for this kit were also so low that no qPCR amplification was observed in the *L. monocytogenes* qPCR targeting *rpoB*.

Among the column-based extraction methods, Power Food, Power Viral, and Zymo were comparable with regard to (i) their ability to recover DNA of the members of

the Mock community and (ii) variability within technical replicates (see Fig. 1; Table 5

provides detailed descriptive statistics). The Power Soil kit (another column-based kit),

generally provided lower bacterial DNA yields for each member of the mock community

than these three kits, even though the differences in log10 copies per ml of milk were

only significantly for some organisms (e.g., *Mycobacterium*).

We also evaluated the ability to isolate bacterial RNA for (i) two kits designed for

isolation of RNA only and for (ii) two kits designed for isolation of both RNA and DNA

(see Table 2). Similar to the DNA detection data detailed above, all kits yielded very low

total nucleic acid concentrations based on spectrophotometry. Reverse transcriptase

qPCR amplification of gene targets for *Salmonella* sp. and *L. monocytogenes* as well as

bovine RNA did not yield amplification on samples treated with DNase (these initial tests

were performed on two technical replicates for each of the four kits). Control

amplifications on nucleic acids before DNAse treatment however, yielded amplification

with CT values that did not differ between the RT-qPCR and the control qPCR targeting

DNA, suggesting the presence of residual DNA in the extracted nucleic acids.

Osmotic Lysis followed by PMA Treatment Decreases Host DNA but also Impacts

Bacterial DNA

We compared protocols that combined osmotic lysis and treatment with multiple

concentrations of PMA for their ability to reduce host DNA, compared to bacterial DNA

copy numbers. To assess if osmotic lysis alone impacted copy-numbers, we compared

non-treated control (NTC) samples to samples that underwent centrifugation and

osmotic lysis with ddH_2O, but no PMA addition (0µM PMA); this comparison revealed

significantly lower bovine and bacterial DNA copy numbers in 0µM PMA samples,

suggesting significant effects of sample processing (including lysis) independent of PMA

addition (Figure 2, NTC vs. 0µM comparisons). We also observed that PMA

concentration significantly impacted bovine and bacterial DNA copy numbers in a dose-

dependent manner: as PMA concentration increased, a sharp decrease was observed

in bovine copy-numbers, whereas a less pronounced, but still noticeable decrease was

observed in bacterial copy-numbers (Figure 2). This trend was confirmed in the model

estimates, which showed a significant and negative effect of PMA in both linear models,

with a greater estimate for bovine copy numbers (-0.022) versus bacterial copy numbers

(-0.012). Treatment with 20 µM PMA was the concentration deemed optimal as it

yielded the greatest decrease in host DNA without critically compromising bacterial DNA

recovery, and thus was chosen for subsequent comparisons with commercially

available host DNA depletion protocols.

We also selected a subset of milk samples from each of the three biological

replicates to undergo deep untargeted sequencing, including one sample each of the

three biological replicates representing (i) untreated samples, (ii) samples treated with

20 µM PMA and (iii) samples that underwent osmotic lysis but were not treated with

PMA (underwent centrifugation, selective lysis with ddH$_2$O but were not treated with

PMA); (iii) was included to allow us to determine if differences observed would be due to

PMA treatment or sample processing prior to PMA addition.

Sequencing had an average read depth of 51,563,707 reads per sample (range

25,488,728 to 127,203,413). Overall less than 1% of all reads that passed quality

control were determined to be of microbial origin regardless of PMA treatment (Table 6,

Figure 3), and around 3% remained unclassified.

While we observed a pronounced increase in the number of bacterial reads per

one million sequenced reads with the PMA treatment (Figure 4A), attempting to deplete

host DNA with PMA also appeared to influence the resulting microbial profiles of

samples, leading to visible differences between replicates of the same biological

samples (Figure 4B). For example, PMA treatment of Bio1 sample resulted in greater

relative abundances of *Cutibacterium* when compared to aliquots of the same sample

(Bio1) that were not treated (NTC) or underwent selective lysis with ddH$_2$O but no PMA

treatment (0µM). Similar differences between NTC, 0µM, and 20µM were observed for

the other two samples sequenced (Bio2 and Bio3, Figure 4B).

The effect of PMA treatment was also seen when the first 3 principal components

were plotted in PCA (Figure 4C) or robust PCA (Figure 4D). We observed that PMA

treated samples separated from other samples (all have negative PC_2) instead of

samples clustering by biological sample origin regardless of PMA treatment. Robust

PCA(42) was used in an effort to alleviate the effects of PMA treatment; however, it was

not able to effectively improve the signal-to-noise ratio introduced by PMA treatment.

Host Depletion Methods Successfully Decrease Host DNA, But Not to the Extent

Needed to Make a Significant Impact for Use in Milk Sequencing Studies

We compared host DNA depletion methods and osmotic lysis followed by PMA

treatment in three independent biological replicates. In each biological replicate, a

freshly collected milk sample was homogenized and divided into five separate aliquots

that were processed in parallel for each of the five methods (e.g., PMA method, Molzym

kit, NEB kit, Qiagen kit, and negative controls [NTC]). Means by which host DNA

depletion was accomplished varied between methods: some kits depended on host

DNA degradation followed by treatment with different enzymes (e.g., Qiagen and

Molzym kits), while the NEB kit was based on the capture of methylated host DNA (for

details on kits and methods, see Table 4, for descriptive statistics, see Table 7). Overall,

host DNA removal treatments significantly decreased bovine copy numbers when

compared to a negative control (Figure 5A). However, we also observed a significant

loss of bacterial copy numbers across methods (Figure 5B). Notably, the Molzym kit

caused the largest reduction of both bovine and bacterial DNA copy numbers to levels

that likely would be insufficient as inputs for shotgun metagenomics sequencing.

Treatment of raw milk with Molzym reagents specifically resulted in a pellet that was

challenging to bring back to solution, which likely led to considerable nucleic acid losses

and renders this method unreliable for use with milk.

As all host DNA depletion methods also led to the depletion of bacterial DNA, we

choose to use the calculated log difference between bovine and bacterial DNA copy

numbers as key metrics for evaluating the different host DNA depletion methods.

Untreated (NTC) samples showed an average of 0.77 log higher bacterial DNA copy

number as compared to bovine DNA copy numbers (Figure 5C). By comparison, the log

difference after different host depletion methods ranged from 1.22 log (PMA method) to

2.09 log (Molzym kit). If treated samples underwent untargeted sequencing, this

decrease in bovine copies would translate into negligible differences in terms of relative

abundances of bacterial reads versus bovine reads given the approximately thousand-

fold difference between the bovine genome size and average bacterial genome sizes.

Enzymatic Selective Lysis of Inoculated Milk followed by PMA Treatment

Differentially Affects Detection of Gram-positive and Gram-negative Bacteria

through qPCR

In a final attempt to optimize a host depletion protocol that would be applicable to

raw milk, we tested a lysis protocol that included a mild protease, with the objective of

more efficiently permeabilizing mammalian cells while keeping bacterial cells intact.

Because we were aware of potential biases that could be introduced by adding an

enzymatic lysis step prior to PMA treatment, we decided to inoculate the samples tested

with Gram-positive and Gram-negative bacteria to address potential differential bacterial

permeabilization by subtilisin that would result in DNA inactivation by PMA binding. We

also investigated if the light source would have an effect on the efficiency of PMA

binding by processing samples in parallel and exposing PMA-treated sample duplicates

to either a halogen-light source or a commercial apparatus designed for use with PMA

treated samples (BLU-V, Qiagen). We prepared two biological replicates with two

technical replicates for each comparison, from which duplicate qPCR reactions were

done.

Treatment with subtilisin decreased the number of culturable bacteria as

assessed through CFU plate counts of milk that had been treated with lysis solution

prior to PMA exposure (Figure 6A). However, we did not observe a difference in CFU in

milk samples treated with different enzyme concentrations. The light source did not

affect copy numbers (*P=0.74*, data not shown); therefore qPCR comparisons were

performed in combined data from both halogen light source and BLU-V apparatus for

Figure 6B. As expected, we observed a decrease in bovine copy numbers in subtilisin-

PMA treated samples compared to negative control to a much greater extent than what

we observed with osmotic lysis (Figure 6B, first panel). However, we also observed a

dose-dependent decrease in Gram-negative copy numbers in treated samples (Figure

6B, third panel), while no differences were observed in Gram-positive copy numbers

(Figure 6B, second panel). This trend was confirmed by a less steep but still noticeable

decrease in total bacterial copy numbers when compared to Gram-negative copy

numbers (Figure 6B, fourth panel). These data led us to conclude that enzymatic

treatment followed by PMA inactivation differentially affects Gram-negative bacteria

copy numbers, leading to biased results if used to prepare DNA for sequencing

purposes.

DISCUSSION

In this study we comprehensively evaluated both nucleic acid extraction and host depletion protocols in bovine raw milk. The rationale of using raw milk as a model included the recent publication of several studies highlighting the variability of raw milk microbiota (43–45); the potential influence of raw milk microbial quality on processed dairy products (46–48); and the fact that patterns of the milk microbiome represent potential "biomarkers" that could be tracked (19, 20). All of these studies highlight milk as a potential candidate for using HTS as part of quality assurance and risk assessment in the food industry in the future. Our results indicate that (i) magnetic-based DNA extraction methods are superior for extracting DNA from milk, (ii) although host DNA depletion methods decrease the amount of bovine DNA in a given sample, the reduction would not be sufficient for effectively depleting bovine DNA in HTS studies of raw milk, and (iii) we observed biases introduced by certain protocols; both on the overall microbial profile as well as selective bias against Gram-negative bacteria.

Magnetic-based DNA extraction is superior to column-based methods for raw milk samples. We compared several nucleic acid extraction protocols on bovine raw milk. While all protocols evaluated for extraction of DNA were able to successfully extract (albeit low amounts of) total DNA, we observed higher variability in replicates in some protocols when compared to others. The low DNA concentration from milk extracts is in agreement with previous reports (24, 25, 49). One particular aspect of this study is that we inoculated different types of bacteria that would be of interest in a dairy processing environment into raw milk and performed targeted qPCR to assess differential extraction of Gram-positive and Gram-negative bacteria. While most

protocols were able to successfully extract DNA from the bacteria inoculated, magnetic-

based DNA extraction approaches had the best recovery and the lowest inter-replicate

variability as evaluated by qPCR. It must be noted that the differences observed in the

recovery of bacterial DNA are a characteristic of processing steps rather than a flaw or

inefficiency of a given protocol (i.e. while some extraction methods carry the entire

sample lysate from one step to the following, some protocols require fixed or maximum

volumes to be transferred to the next step due to volume limitations in reaction tubes

used). It is important to highlight that different DNA extraction methods have

advantages and drawbacks, which can also vary according to the material they are

applied to. As no general gold-standard method exists to date for this application, the

selection of an extraction method should be based on the objectives of each particular

study (50). Nevertheless, a growing consensus exists on the importance of mechanical

disruption such as bead-beating for microbiome applications (51), and magnetic-based

extractions have been demonstrated to be particularly effective in the diagnosis of

tuberculosis in serum and plasma of patients (52) and the extraction of algae DNA for

NGS studies (53), highlighting magnetic methods as a common theme around diverse

nucleic-acid based applications.

Extracting bacterial RNA from high-quality raw milk is challenging. The

methods used here co-purify both DNA and RNA, or each nucleic acid separately, and

based on inoculation of bacterial loads in some cases exceeding regulatory standards

for raw milk; we expected that bacterial RNA should be detectable. We observed very

low concentrations of extracted DNA and were unable to successfully detect bacterial

RNA from our inoculated milk samples. This was a surprising finding, as literature on

the milk metatranscriptome is available (18). However, the protocols previously used to

successfully isolate RNA from milk involved extensive centrifugation of large sample

volumes followed by several wash steps (54), as well as samples with much greater

bacterial loads, including dairy products such as cheese, which contain high levels of

organisms responsible for fermentation of the products (18, 20).

Current methods for host DNA depletion are not suitable for application in

untargeted HTS studies of milk. Because the bovine cells are present in milk along

with the microbes of interest, and due to the striking difference between bovine and

bacterial genome sizes, developing efficient yet unbiased host-DNA depletion methods

is critical for the adoption of HTS technologies in food safety. While we observed a

decrease in the number of bovine copies when compared to bacterial copies as

assessed through qPCR in all protocols evaluated, this decrease would not be sufficient

to effectively change the relative abundance of reads being assigned to the bovine

genome in HTS studies. This was confirmed when we performed deep untargeted

sequencing in a subset of samples that had demonstrated promising results in qPCR

but still detected over 99% of reads mapping to the bovine genome independent of host

DNA depletion.

Among the other host DNA depletion protocols evaluated were

immunoprecipitation of methylated eukaryotic DNA and selective lysis of mammalian

cells followed by DNAse treatment. One reason for the lack of efficiency observed in the

treatment of milk with either of these methods could be the challenges associated with

extracting DNA from milk in the first place. Such protocols require large amounts of

good quality DNA, which is practically impossible to obtain from milk samples without

extensive centrifugation and pellet washes, potentially proving impractical in industry

settings and representing added opportunities for sample contamination or unintentional

bias of the microbiota for lipid- or protein-bound microorganisms. The characteristics of

milk in itself can also pose a challenge (presence of fats, proteins, and ions) (8, 9).

Challenges specific to nucleic acid extraction from milk have been discussed in the

literature to some extent, specifically Metzger et al. reported challenges in amplifying

bacterial DNA from DNA extracted from milk (55).

To the best of our knowledge, this was the first study to evaluate host DNA

depletion methods in a food matrix. Limited literature is available trying to address host

contamination in clinical sequencing applications for pathogen detection, with reported

improvements in the detection of malaria (56) and pathogens in infected tissue samples

(57) and sputum (31). Reports are also available describing attempts of using host DNA

depletion in cerebrospinal (29), and arthroplasty (28) fluids, as well as human saliva

(27). The common theme around these studies was the varying efficiencies across

methods and sample types, highlighting the need for individual assessment of host

depletion methods using the desired sample type.

Enzymatic-based selective lysis followed by PMA treatment differentially

affects Gram-negative bacteria in inoculated raw milk. Propidium monoazide has

been extensively used to differentiate between live and dead bacterial cells due to its

inability to penetrate intact cell membranes and has recently been indicated to be an

effective method for host DNA depletion for use in HTS studies (27, 58, 59). We thus

developed an enzymatic-based method for host DNA depletion (which utilized a mild

protease) combined with subsequent PMA treatment in an effort to better lyse

mammalian cells while retaining intact bacterial cells.

While we detected an effect of the enzymatic treatment on total bacterial

numbers as measured through plate counts that appeared to be independent of

enzymatic concentration, we did not observe a decrease in Gram-positive bacterial

numbers as measured through qPCR after PMA exposure. Nevertheless, we observed

an enzyme dose-dependent decrease of Gram-negative copy numbers in samples that

underwent selective lysis followed by PMA exposure. These data led us to conclude

that enzymatic treatment might have differentially permeabilized the membrane of

Gram-negative cells, allowing PMA to bind to DNA without observable differences in

overall bacterial survival. Taken together, these observations suggest that potential

biases could occur against Gram-negative bacteria detection through qPCR in this

method.

It is imperative to optimize protocols for samples with different

characteristics. The generalizability of this study lies on the importance of

standardization and validation of methods for each specific food matrix. It is also

important to highlight the challenges associated with low bacterial biomass samples and

the fact that various foods have a wide range of biomass, from very low in raw milk, to

relatively high microbial content in some cultured dairy products. Efforts to standardize

and recommend best practices in HTS studies have recently begun, particularly

pertaining to low biomass samples (60–62) and must be continued.

CONCLUSION

Our results suggest that magnetic-based extraction methods are superior for

bacterial nucleic acid isolation from bovine milk. Host DNA remains a challenge for

untargeted sequencing of milk, highlighting that the food matrix characteristics should

always be considered whenever planning HTS studies. Enzymatic-lysis based PMA

host depletion introduced dose-dependent biases against Gram-negative bacteria,

suggesting that selective lysis permeabilized Gram-negative to PMA, which

subsequently hindered our ability to detect Gram-negative bacteria through qPCR

without affecting live bacterial counts. While it is not possible to test all methods

available at any given time, we focused on kits and protocols that have been most

widely used for the extraction of nucleic acids from milk. As procedures are improved, or

new methods are developed, reevaluation of protocols available would prove useful, as

the development of HTS-based tools to aid and improve quality assurance and food

safety programs continue to hold great promise.

DECLARATIONS

Ethics Approval

All experimental procedures were carried out at Cornell University, because no animal sampling was necessary for this study, an approved research protocol by the Cornell University Institutional Animal Care and Use Committee was not necessary.

Consent for publication

Not applicable.

Availability of data and material

The datasets and R code used and/or analyzed during the current study are available on https://github.com/ErikaGanda/MilkDNA. Raw sequencing reads have been deposited into SRA under the submission number SUB7938697.

Competing interests

Authors are part of the Consortium for Sequencing the Food Supply Chain. The authors declare that they have no conflict of interest.

Funding

This work was supported by the New York State Dairy Promotion Advisory Board (Albany, NY; OSP no. 88975), dairy farmers dedicated to the manufacture of high-quality dairy products.

Authors' contributions

Conceived and designed the experiments: **EKG, MW** Performed the experiments: **EKG** Performed automated DNA extraction: **BC, RRA** Analyzed the data:

EKG, KLB, NH, BK Wrote the paper: **EKG** Revised the manuscript: **EKG, KLB, NH,**

BK, LG, MW. All authors reviewed the manuscript.

Acknowledgements

We thank members of CURC for allowing us to sample, Dr. VanDerVen Lab for

providing us *Mycobacterium smegmatis* MC2-155 strain, and Dr. Mann Lab for

providing bovine RNA used as a standard.

REFERENCES

1. Tan X, Chung T, Chen Y, Macarisin D, Laborde L, Kovac J. 2019. The occurrence
    of Listeria monocytogenes is associated with built environment microbiota in three
    tree fruit processing facilities. Microbiome https://doi.org/10.1186/s40168-019-
    0726-2.

2. Guidone A, Zotta T, Matera A, Ricciardi A, De Filippis F, Ercolini D, Parente E.
    2016. The microbiota of high-moisture mozzarella cheese produced with different
    acidification methods. Int J Food Microbiol
    https://doi.org/10.1016/j.ijfoodmicro.2015.09.002.

3. Haiminen N, Edlund S, Chambliss D, Kunitomi M, Weimer BC, Ganesan B, ...
    2019. Food authentication from shotgun sequencing reads with an application on
    high protein powders. npj Sci Food in press.

4. Beck KL, Haiminen N, Chambliss D, Edlund S, Kunitomi M, Carol Huang B, Kong
    N, Ganesan B, Baker R, Kawas B, Davis M, Prill RJ, Krishnareddy H, Marlowe
    CH, Pierre S, Quintanar A, Parida L, Kaufman J, Weimer BC. 2020. Monitoring
    the microbiome for food safety and quality using deep 1 shotgun sequencing 2 3
    Running Title: Monitoring the microbiome for food safety and quality 4 5. bioRxiv
    2020.05.18.102574.

5. Jagadeesan B, Gerner-Smidt P, Allard MW, Leuillet S, Winkler A, Xiao Y,
    Chaffron S, Van Der Vossen J, Tang S, Katase M, McClure P, Kimura B, Ching
    Chai L, Chapman J, Grant K. 2019. The use of next generation sequencing for
    improving food safety: Translation into practice. Food Microbiol.

6.   Foroutan A, Guo AC, Vazquez-Fresno R, Lipfert M, Zhang L, Zheng J, Badran H,

Budinski Z, Mandal R, Ametaj BN, Wishart DS. 2019. Chemical Composition of

Commercial Cow's Milk. J Agric Food Chem

https://doi.org/10.1021/acs.jafc.9b00204.

7.   Demeke T, Jenkins GR. 2010. Influence of DNA extraction methods, PCR

inhibitors and quantification methods on real-time PCR assay of biotechnology-

derived traits. Anal Bioanal Chem.

8.   Schrader C, Schielke A, Ellerbroek L, Johne R. 2012. PCR inhibitors -

occurrence, properties and removal. J Appl Microbiol.

9.   Bickley J, Short JK, McDowell DG, Parkes HC. 1996. Polymerase chain reaction

(PCR) detection of Listeria monocytogenes in diluted milk and reversal of PCR

inhibition caused by calcium ions. Lett Appl Microbiol

https://doi.org/10.1111/j.1472-765X.1996.tb01131.x.

10. Soboleva SE, Zakharova OD, Sedykh SE, Ivanisenko N V, Buneva VN, Nevinsky

GA. 2019. DNase and RNase activities of fresh cow milk lactoferrin. J Mol

Recognit2019/02/15. e2777.

11. DiCenzo GC, Finan TM. 2017. The Divided Bacterial Genome. Microbiol Mol Biol

Rev https://doi.org/10.1128/ MMBR.00019-17.

12. Zhou S, Goldstein S, Place M, Bechner M, Patino D, Potamousis K, Ravindran P,

Pape L, Rincon G, Hernandez-Ortiz J, Medrano JF, Schwartz DC. 2015. A clone-

free, single molecule map of the domestic cow (Bos taurus) genome. BMC

Genomics https://doi.org/10.1186/s12864-015-1823-7.

13. Bhatt VD, Ahir VB, Koringa PG, Jakhesara SJ, Rank DN, Nauriyal DS, Kunjadia

AP, Joshi CG. 2012. Milk microbiome signatures of subclinical mastitis-affected

cattle analysed by shotgun sequencing. J Appl Microbiol2012/01/27. 112:639–

650.

14.   Kuehn JS, Gorden PJ, Munro D, Rong R, Dong Q, Plummer PJ, Wang C, Phillips

GJ. 2013. Bacterial community profiling of milk samples as a means to

understand culture-negative bovine clinical mastitis. PLoS One 8:e61959.

15.   Ganda EK, Gaeta N, Sipka A, Pomeroy B, Oikonomou G, Schukken YH, Bicalho

RC. 2017. Normal milk microbiome is reestablished following experimental

infection with Escherichia coli independent of intramammary antibiotic treatment

with a third-generation cephalosporin in bovines. Microbiome 5.

16.   Walsh AM, Crispie F, Daari K, O'Sullivan O, Martin JC, Arthur CT, Claesson MJ,

Scott KP, Cotter PD. 2017. Strain-level metagenomic analysis of the fermented

dairy beverage nunu highlights potential food safety risks. Appl Environ Microbiol

https://doi.org/10.1128/AEM.01144-17.

17.   Addis MF, Tanca A, Uzzau S, Oikonomou G, Bicalho RC, Moroni P. 2016. The

bovine milk microbiota: Insights and perspectives from -omics studies. Mol

Biosyst 12:2359–2372.

18.   Richards VP, Choi SC, Pavinski Bitar PD, Gurjar AA, Stanhope MJ. 2013.

Transcriptomic and genomic evidence for Streptococcus agalactiae adaptation to

the bovine environment. BMC Genomics 14:920.

19.   Wolfe BE, Button JE, Santarelli M, Dutton RJ. 2014. Cheese rind communities

provide tractable systems for in situ and in vitro studies of microbial diversity. Cell

https://doi.org/10.1016/j.cell.2014.05.041.

20. De Filippis F, Genovese A, Ferranti P, Gilbert JA, Ercolini D. 2016.

Metatranscriptomics reveals temperature-driven functional changes in microbiome

impacting cheese maturation rate. Sci Rep 6:21871.

21. Geer SR, Barbano DM. 2014. The effect of immunoglobulins and somatic cells on

the gravity separation of fat, bacteria, and spores in pasteurized whole milk. J

Dairy Sci 97:2027–2038.

22. Caplan Z, Melilli C, Barbano DM. 2013. Gravity separation of fat, somatic cells,

and bacteria in raw and pasteurized milks1. J Dairy Sci 96:2011–2019.

23. Kirchner B, Pfaffl MW, Dumpler J, Von Mutius E, Ege MJ. 2016. MicroRNA in

native and processed cow's milk and its implication for the farm milk effect on

asthma. J Allergy Clin Immunol 137:1893-1895.e13.

24. Volk H, Piskernik S, Kurinčič M, Klančnik A, Toplak N, Jeršek B. 2014. Evaluation

of different methods for DNA extraction from milk. J Food Nutr Res.

25. Quigley L, O'Sullivan O, Beresford TP, Paul Ross R, Fitzgerald GF, Cotter PD.

2012. A comparison of methods used to extract bacterial DNA from raw milk and

raw milk cheese. J Appl Microbiol2012/03/29. 113:96–105.

26. Lima SF, Bicalho MLS, Bicalho RC. 2018. Evaluation of milk sample fractions for

characterization of milk microbiota from healthy and clinical mastitis cows. PLoS

One2018/03/22. 13:e0193671.

27. Marotz CA, Sanders JG, Zuniga C, Zaramela LS, Knight R, Zengler K. 2018.

Improving saliva shotgun metagenomics by chemical host DNA depletion.

Microbiome2018/02/28. 6:42.

28. Thoendel M, Jeraldo PR, Greenwood-Quaintance KE, Yao JZ, Chia N, Hanssen

765  AD, Abdel MP, Patel R. 2016. Comparison of microbial DNA enrichment tools for

766  metagenomic whole genome sequencing. J Microbiol Methods

767  https://doi.org/10.1016/j.mimet.2016.05.022.

768 29. Hasan MR, Rawat A, Tang P, Jithesh P V, Thomas E, Tan R, Tilley P. 2016.

769  Depletion of Human DNA in Spiked Clinical Specimens for Improvement of

770  Sensitivity of Pathogen Detection by Next-Generation Sequencing. J Clin

771  Microbiol 54:919–927.

772 30. Oechslin CP, Lenz N, Liechti N, Ryter S, Agyeman P, Bruggmann R, Leib SL,

773  Beuret CM. 2018. Limited Correlation of Shotgun Metagenomics Following Host

774  Depletion and Routine Diagnostics for Viruses and Bacteria in Low Concentrated

775  Surrogate and Clinical Samples. Front Cell Infect Microbiol2018/11/09. 8:375.

776 31. Nelson MT, Pope CE, Marsh RL, Miller SI, Borenstein E, Hoffman

777  Correspondence LR. 2019. Human and Extracellular DNA Depletion for

778  Metagenomic Analysis of Complex Clinical Infection Samples Yields Optimized

779  Viable Microbiome Profiles. CellReports 26:2227-2240.e5.

780 32. Chiu CY, Miller SA. 2019. Clinical metagenomics. Nat Rev Genet. Nature

781  Publishing Group.

782 33. FDA. 2017. Grade "A" Pasteurized Milk Ordinance.

783 34. Buehler AJ, Martin NH, Boor KJ, Wiedmann M. 2018. Psychrotolerant spore-

784  former growth characterization for the development of a dairy spoilage predictive

785  model. J Dairy Sci2018/05/29. 101:6964–6981.

786 35. Gaillard S, Leguerinel I, Mafart P. 1998. Model for Combined Effects of

787  Temperature, pH and Water Activity on Thermal Inactivation of Bacillus cereus

Spores. J Food Sci 63:887–889.

36.   Brankatschk R, Bodenhausen N, Zeyer J, Burgmann H. 2012. Simple absolute

quantification method correcting for quantitative PCR efficiency variations for

microbial community samples. Appl Env Microbiol 78:4481–4489.

37.   Pinheiro J, Bates D, DebRoy S, Sarkar D RCT. 2020. nlme: Linear and Nonlinear

Mixed Effects Models. R package version 3.1-148.

38.   Russell Lenth, Henrik Singmann, Jonathon Love, Paul Buerkner MH. 2020.

emmeans: Estimated marginal means. R package verison 1.4.8.

39.   Krueger F. TrimGalore.

40.   Wood DE, Salzberg SL. 2014. Kraken: Ultrafast metagenomic sequence

classification using exact alignments. Genome Biol 15:R46.

41.   Candès EJ, Li X, Ma Y, Wright J. 2011. Robust principal component

analysis?Journal of the ACM.

42.   Chen X, Zhang B, Wang T, Wang T, Bonni A, Zhao G. 2020. Robust principal

component analysis for accurate outlier sample detection in RNA-Seq data. BMC

Bioinformatics 21:269.

43.   Quigley L, O'Sullivan O, Stanton C, Beresford TP, Ross RP, Fitzgerald GF, Cotter

PD. 2013. The complex microbiota of raw milk. FEMS Microbiol Rev 37:664–698.

44.   Kable ME, Srisengfa Y, Laird M, Zaragoza J, McLeod J, Heidenreich J, Marco

ML. 2016. The Core and Seasonal Microbiota of Raw Bovine Milk in Tanker

Trucks and the Impact of Transfer to a Milk Processing Facility. MBio2016/08/25.

7.

45.   Doyle CJ, Gleeson D, O'Toole PW, Cotter PD. 2017. Impacts of Seasonal

Housing and Teat Preparation on Raw Milk Microbiota: a High-Throughput

Sequencing Study. Appl Env Microbiol 83.

46. Skeie SB, Haland M, Thorsen IM, Narvhus J, Porcellato D. 2019. Bulk tank raw

milk microbiota differs within and between farms: A moving goalpost challenging

quality control. J Dairy Sci2019/01/15. 102:1959–1971.

47. Porcellato D, Aspholm M, Skeie SB, Monshaugen M, Brendehaug J, Mellegård H.

2018. Microbial diversity of consumption milk during processing and storage. Int J

Food Microbiol https://doi.org/10.1016/j.ijfoodmicro.2017.11.004.

48. De Filippis F, Parente E, Ercolini D. 2018. Recent Past, Present, and Future of

the Food Microbiome. Annu Rev Food Sci Technol 9:589–608.

49. Sun L, Dicksved J, Priyashantha H, Lundh, Johansson M. 2019. Distribution of

bacteria between different milk fractions, investigated using culture-dependent

methods and molecular-based and fluorescent microscopy approaches. J Appl

Microbiol https://doi.org/10.1111/jam.14377.

50. Ali N, De Cássia R, Rampazzo P, Dias Tavares Costa A, Krieger MA. 2017.

Current Nucleic Acid Extraction Methods and Their Implications to Point-of-Care

Diagnostics https://doi.org/10.1155/2017/9306564.

51. Lim MY, Song EJ, Kim SH, Lee J, Nam Y Do. 2018. Comparison of DNA

extraction methods for human gut microbial community profiling. Syst Appl

Microbiol 41:151–157.

52. Pan S, Gu B, Wang H, Yan Z, Wang P, Pei H, Xie W, Chen D, Liu G. 2013.

Comparison of four DNA extraction methods for detecting Mycobacterium

tuberculosis by real-time PCR and its clinical application in pulmonary

tuberculosis. J Thorac Dis 5:251–257.

53.   Fort A, Guiry MD, Sulpice R. 2018. Magnetic beads, a particularly effective novel

method for extraction of NGS-ready DNA from macroalgae. Algal Res

https://doi.org/10.1016/j.algal.2018.04.015.

54.   García-Nogales P, Serrano A, Secchi S, Gutiérrez S, Arís A. 2010. Comparison of

commercially-available RNA extraction methods for effective bacterial RNA

isolation from milk spiked samples. Electron J Biotechnol 13.

55.   Metzger SA, Hernandez LL, Skarlupka JH, Suen G, Walker TM, Ruegg PL. 2018.

Influence of sampling technique and bedding type on the milk microbiota: Results

of a pilot study. J Dairy Sci2018/04/24. 101:6346–6356.

56.   Feehery GR, Yigit E, Oyola SO, Langhorst BW, Schmidt VT, Stewart FJ,

Dimalanta ET, Amaral-Zettler LA, Davis T, Quail MA, Pradhan S. 2013. A Method

for Selectively Enriching Microbial DNA from Contaminating Vertebrate Host DNA.

PLoS One 8:e76096.

57.   Heravi FS, Zakrzewski M, Vickery K, Hu H. 2020. Host DNA depletion efficiency

of microbiome DNA enrichment methods in infected tissue samples. J Microbiol

Methods 170.

58.   Fittipaldi M, Nocker A, Codony F. 2012. Progress in understanding preferential

detection of live cells using viability dyes in combination with DNA amplification. J

Microbiol Methods.

59.   Nocker A, Camper AK. 2009. Novel approaches toward preferential detection of

viable cells using nucleic acid amplification techniques. FEMS Microbiol Lett

291:137–142.

60.   Sinha R, Abu-Ali G, Vogtmann E, Fodor AA, Ren B, Amir A, Schwager E,

858   Crabtree J, Ma S, Abnet CC, Knight R, White O, Huttenhower C. 2017.

859   Assessment of variation in microbial community amplicon sequencing by the

860   Microbiome Quality Control (MBQC) project consortium. Nat Biotechnol

861   https://doi.org/10.1038/nbt.3981.

862   61.   Minich JJ, Sanders JG, Amir A, Humphrey G, Gilbert JA, Knight R. 2019.

863   Quantifying and Understanding Well-to-Well Contamination in Microbiome

864   Research. mSystems https://doi.org/10.1128/msystems.00186-19.

865   62.   Eisenhofer R, Minich JJ, Marotz C, Cooper A, Knight R, Weyrich LS. 2019.

866   Contamination in Low Microbial Biomass Microbiome Studies: Issues and

867   Recommendations. Trends Microbiol.

869 **Table 1 – Organisms used in this study**

Target	Isolate	Source	Accession*
Total Bacterial DNA/RNA	Pooled Culture		NC_000913
Bacillus wiedmanii	FSL H7-0344	Pasteurized 2% milk	NC_004722
Listeria monocytogenes	FSL A5-0145	Raw Milk	NC_003210
Mycobacterium smegmatis	FSL X3-0054	VanDerVen Lab – MC2155	NC_008596
Salmonella enterica	FSL A5-0218	Raw milk	NC_003197
Bovine DNA	Bovine Blood	Cornell University Dairy Farm	NM_001037471.2

870   *NCBI accession number of genomes used for primer design; primers for

871 detection of total bacterial DNA target 16S rDNA and were designed based on

872 conserved regions of the 16S gene.

873 **Table 2– Nucleic Acid Extraction Kit Characteristics**

Abbreviations	Kit	Sample Input (µL)	Nucleic Acid Output (µL)	Nucleic Acid	Processing	Capture	Catalog Number	Manufacturer
PviralDNA PviralRNA	AllPrep PowerViral DNA/RNA kit	200	100	DNA/RNA	Manual	Column-based	28000-50	Qiagen
COREDNA CORERNA	MagMAX CORE Nucleic Acid Isolation kit	200	90	DNA/RNA	Automated	Magnetic-based	A32700	ThermoFisher
ZymoDNA ZymoRNA	ZymoBIOMICs DNA/RNA mini kit	250	100	DNA/RNA	Manual	Column-based	R2002	Zymo
Mastitis	MagAttract Mastitis	400	100	DNA	Automated	Magnetic-based	947757	Qiagen
Pfood	DNeasy PowerFood	1800	100	DNA	Manual	Column-based	21000-100	Qiagen
PSoilP	Power Soil Pro	500	100	DNA	Manual	Column-based	47014	Qiagen
EZFood	E.Z.N.A. ® Food DNA kit	500	100	DNA	Manual	Column-based	D4616-00	Omega
RNeasy	RNeasy Protect Bacteria Mini Kit	1,500	50	RNA	Manual	Column-based	74524	Qiagen
EZNARNA	E.Z.N.A.® HP Total RNA Kit	500	50	RNA	Manual	Column-based	R6812-00	Omega

875 **Table 3 – Primers Used in this Study**

Target	Primer	Sequence	Fragment Size	Source
Total Bacterial DNA/RNA	EKG43-16-TotalBact16S-F2	GTAGCGGTAAATGCGTAGA	120	This study
	EKG43-4-TotalBact16S-R	GACTACCAGGGTATCTAATC		This study
Bacillus wiedmannii	JK2739-3rpoB_F	AACGTGCTTGTTGGCTTCAT	*152*	This study
	JK2739-4rpoB_R	TCTTCTGGTCCAAGCTTCGT		This study
Listeria monocytogenes	VGO-23-rpoB-RT-F	TCGTCGTCTTCGTTCTGTTG	221	(Liu et al., 2017)
	VGO-24-rpoB-RT-R	GTTCGCCAAGTGGATTTGTT		(Liu et al., 2017)
Mycobacterium smegmatis	EKG43-15-Mycobacterium-rpoB-F2	TCGGTGAGCTGATCCAGAAC	156	This study
	EKG43-14-Mycobacterium-rpoB-R	TGCCGAAGAACTCCTTGATC		This study
Salmonella sp.	EKG43-7-Salmonella-rpoB-F	GTACCGTCGTGTGGTTGATG	170	This study
	EKG43-8-Salmonella-rpoB-R	GGCTGAACAAGCTGGATTCG		This study
Bovine RNA	Bt03229278_m1	bovine ubiquitously-expressed transcript	89	Commercial assay
Bovine DNA	Bt03229276_g1	bovine ubiquitously-expressed transcript	88	Commercial assay

878   **Table 4 - Host DNA depletion Kit Characteristics**

Abbreviations	Kit	Method	Catalog Number
NEB	NEBNext Microbiome DNA Enrichment	Methylated Host DNA capture	E2612
Qia	QIAamp DNA Microbiome	Host DNA degradation with Benzonase	51704
Molz	Molzym Ultra-Deep Microbiome Prep	Host DNA degradation with MolDNaseB	G-020-025
PMA[1]	Propidium Monoazide	Covalent attachment to free DNA	40019

879   [1]Not a commercial kit, in-house protocol combining selective lysis and exposure

880   to PMA and light.

881 **Table 5 – Descriptive Statistics of Log Copies per mL of Milk – DNA extraction**

Organism	Mean log10 copies per mL of milk as detected via qPCR (StdDev), N						
	COREDNA	Mastitis	EZFood	PFood	PSoilP	PviralDNA	ZymoDNA
Total Bacterial DNA	6.99 (0.96),18	7.51 (0.54),18	4.30 (1.72),12	6.71 (0.57),18	6.31 (0.48),18	6.96 (0.59),18	6.78 (0.40),18
Bacillus wiedmanii	6.53 (0.39),18	6.67 (0.25),18	4.31 (0.77),12	5.79 (0.67),18	5.61 (0.45),18	6.09 (0.49),18	6.02 (0.32),18
Listeria monocytogenes	6.21 (0.89),18	6.58 (0.48),18	NA (NA)[a],12	5.93 (0.49),18	5.46 (0.68),18	6.84 (0.09),18	6.00 (0.59),18
Mycobacterium smegmatis	6.81 (0.31),18	6.70 (0.42),18	3.97 (1.39),12	6.29 (0.34),18	5.44 (0.82),18	6.25 (0.38),18	6.30 (0.58),18
Salmonella sp.	6.07 (0.26),18	5.80 (0.61),18	3.83 (1.12),12	5.24 (0.72),18	4.13 (0.77),18	5.77 (0.19),18	5.34 (0.61),18
Bovine DNA	5.45 (0.08),18	5.37 (0.05),18	4.71 (0.59),12	4.86 (0.08),18	4.05 (0.24),18	5.20 (0.11),18	4.81 (0.06),18

882 [a] Not applicable, this method did not yield detectable signal in qPCR for *Listeria monocytogenes*.

884 **Table 6 – Descriptive statistics of shotgun metagenomics sequencing results assessing the effect of PMA**

885 **treatment**

Parameter[1]	Mean (StdDev), N		
	No Treatment	0 µM PMA (Wash only)	20 µM PMA
Bovine (%)	93.87(0.54),3	93.83(0.366),3	92.46(0.866),3
Microbial (%)	0.010(0.002),3	0.011(0.005),3	0.109(0.027),3
Unclassified (%)	2.933(0.122),3	3.066(0.04),3	3.113(0.11),3
Low quality (QC filtered) (%)	3.18(0.441),3	3.09(0.384),3	4.31(0.851),3
Average Input Reads[2]	67,983,431	45,846,439	40,861,252

887 [1] Indicates which class (e.g., eukariotic, bacterial, viral, archaea) sequencing reads were assigned to using methods

888 described in Beck et al., 2020.

889 [2] Average across three samples tested in each category

Table 7 – Descriptive Statistics of Log Copies per mL of Milk – Host Depletion

Target	Mean (StdDev), N				
	NTC	20µM PMA	Molz	NEB	Qia
Total Bacterial DNA	6.18 (0.23) 12	5.25 (0.09) 12	4.42 (0.40) 8	5.99 (0.07) 12	5.04 (0.17) 12
Bovine DNA	5.41 (0.23) 12	4.03 (0.28) 12	2.32 (0.03) 8	4.56 (0.08) 12	3.55 (0.34) 12

Supplementary Table S1 – Milk Sample Characteristics

Sample	Experiment	CFU[1]	Log10 CFU	SCC[2]	Fat (%)	Protein (%)	Lactose (%)	Solids (%)
Uninoculated Milk – Incoming	1	7,100	3.85	153,000	3.69	3.1	4.77	12.54
Uninoculated Milk – DNA extraction	1	22,875	4.36	152,000	3.68	3.08	4.78	12.51
Inoculated Milk – DNA extraction	1	309,500	5.49	138,000	3.67	3.11	4.73	12.47
Uninoculated Milk – Incoming	2	3,020	3.48	149,000	4.03	2.97	4.85	12.83
Uninoculated Milk – DNA extraction	2	7,630	3.88	143,000	4.03	2.97	4.81	12.79
Inoculated Milk – DNA extraction	2	453,000	5.66	147,000	4.03	2.98	4.86	12.81
Uninoculated Milk – Incoming	3	19,850	4.30	116,000	3.85	2.88	4.86	12.52
Uninoculated Milk – DNA extraction	3	12,070	4.08	109,000	3.85	2.88	4.88	12.51
Inoculated Milk – DNA extraction	3	2,617,500	6.42	122,000	3.82	2.88	4.84	12.45

895 [1] CFU – Colony Forming Units

896 [2] SCC – Somatic Cell Count

897 **Supplementary Table S2 – Descriptive Statistics of Nucleic Acid Quantification and Quality Measurements**

Concentration (Fluorescence)	Mean ng/µL (StdDev), N						
	COREDNA	Mastitis	EZFood	PFood	PSoilP	PviralDNA	ZymoDNA
Uninoculated Milk	1.21 (0.42) 3	2.07 (0.82) 3	1.93 (1.59) 2	2.88 (2.94) 3	1.03 (1.20) 3	0.57 (0.17) 3	0.52 (0.27) 3
Inoculated Milk	1.22 (0.30) 9	2.32 (0.69) 9	1.10 (0.72) 6	2.68 (1.85) 9	2.03 (2.50) 9	0.63 (0.19) 9	1.22 (0.30) 9
Mock Community	0.08 (0.05) 3	0.18 (0.17) 3	0.21 (0.23) 2	0.17 (0.18) 3	0.19 (0.18) 3	0.05 (0.03) 3	0.16 (0.18) 3
No Template Control	0.03 (0.02) 3	0.15 (0.19) 3	0.19 (0.25) 2	0.25 (0.13) 3	0.15 (0.19) 3	0.03 (0.02) 3	0.15 (0.19) 3

Concentration (Absorbance)	Mean ng/µL (StdDev), N						
	COREDNA	Mastitis	EZFood	PFood	PSoilP	PviralDNA	ZymoDNA
Uninoculated Milk	6.46 (0.35) 3	7.80 (2.29) 3	12.3 (2.75) 2	6.10 (5.60) 3	-0.8 (2.11) 3	1.03 (1.22) 3	15.3 (2.97) 3
Inoculated Milk	8.25 (5.15) 9	8.55 (2.49) 9	10.7 (4.89) 6	6.38 (0.95) 9	-1.5 (1.68) 9	1.03 (1.20) 9	11.0 (4.19) 9
Mock Community	2.00 (0.34) 3	5.13 (2.62) 3	5.75 (3.46) 2	2.63 (2.87) 3	1.5 (1.83) 3	1.60 (0.55) 3	12.1 (2.73) 3
No Template Control	1.86 (0.60) 3	4.43 (2.39) 3	4.45 (0.21) 2	3.9 (5.56) 3	1.63 (3.07) 3	1.13 (1.33) 3	9.36 (5.76) 3

260/280 Ratios (Absorbance)	Mean (StdDev), N						
	COREDNA	Mastitis	EZFood	PFood	PSoilP	PviralDNA	ZymoDNA
Uninoculated Milk	2.22 (0.30) 3	3.23 (1.61) 3	1.62 (0.08) 2	2.31 (0.47) 3	-2.60 (6.58) 3	0.70 (1.69) 3	1.26 (0.11) 3

Inoculated Milk	2.32 (0.46) 9	2.78 (0.96) 9	1.57 (0.12) 6	2.04 (0.12) 9	0.44 (1.68) 9	1.56 (0.72) 9	1.19 (0.11) 9
Mock Community	3.29 (0.80) 3	10.7 (3.47) 3	1.97 (0.02) 2	0.95 (0.86) 3	2.19 (0.90) 3	1.76 (0.95) 3	1.16 (0.19) 3
No Template Control	-2.30 (8.94) 3	11.0 (12.5) 3	2.22 (0.86) 2	1.66 (0.20) 3	2.25 (0.85) 3	1.15 (1.12) 3	1.11 (0.13) 3

Figure Legends

Figure 1

Scatterplots of raw data of normalized log copy numbers per mL of milk obtained with different DNA extraction kits. Bars overlaid represent model estimated marginal means and standard deviations. Different letters represent significantly different means at the 0.05 level after Tukey multiple comparison adjustment. Scatterplot shapes represent each of three independent biological replicates (circles - first; triangles - second; and squares - third., whereas colors represent kits evaluated. Each panel depicts results of a single gene target evaluated.

Figure 2

Effect of osmotic lysis of raw milk followed by treatment with various PMA concentrations on host and bacterial DNA counts as determined by qPCR. Boxplots represent normalized log copy numbers per mL of milk. NTC: untreated sample, 0µM: milk that underwent osmotic lysis but was not treated with PMA. Asterisks represent significant differences at $p < 0.05$ in linear model comparisons. Osmotic lysis significantly decreases Log10 copies in non-PMA treated samples (NTC vs 0µM) and increasing PMA concentrations significantly decreases Log10 Copies in a dose-dependent manner.

Figure 3

Effect of host DNA depletion with PMA on percentage of sequencing reads assigned to the bovine genome or microbial genomes. Raw milk was collected in three separate

days and divided into aliquots that underwent each of the processing methods, including

(i) NTC: untreated sample, (ii) 0μM: milk that underwent osmotic lysis but was not

treated with PMA, and (iii) 20μM: milk that underwent osmotic lysis and PMA treatment

at 20 μM.

Figure 4

Microbial profile of samples exposed to osmotic lysis and PMA treatment. (A) Microbial

reads per million reads by biological sample and treatment. (B) Relative abundance

within microbial reads by biological sample and treatment. (C) Principal Component

Analysis (PCA) and (D) Robust PCA performed in Isometric Log-Ratio transformed

microbial reads.

NTC: untreated sample, 0μM: milk that underwent osmotic lysis but was not treated with

PMA, 20μM: milk that underwent osmotic lysis and PMA treatment at 20 μM.

Figure 5

Comparisons of host depletion methods in uninoculated milk. Scatterplots of raw data of

normalized log copy numbers per mL of milk. Bars overlaid represent model estimated

marginal means and standard deviations for comparisons performed on Total Bacterial

copy numbers (A) and Bovine copy numbers (B) and. Different letters represent

significant Tukey-adjusted differences at the 0.05 level. Scatterplot shapes represent

each independent experiment, whereas colors represent kits evaluated. Paired boxplots

(C) represent normalized log copy numbers per mL of milk of two independent

experiments with two technical replicates each, with exception of Molzym, that only had

technical replicates performed in one day. Numbers represent the log difference

between mean log Bacterial copy numbers and mean log Bovine copy numbers.

⁂Denotes that technical replicates were only performed in one day due to sample

becoming an insoluble pellet during processing.

Figure 6

Comparison of enzymatic lysis prior to host DNA depletion with 20 µM of PMA

inoculated milk. Aerobic standard plate count results (A) and qPCR results (B). Different

colors correspond to two independent experiments with two technical replicates each,

from with duplicate qPCR reactions were done. CT values of samples in which melt

curves did not match standard samples' peaks were not used.