Age effects aggressive behavior: RNA-seq analysis in cattle with implications for studying

neoteny under domestication

Paulina G. Eusebi[1], Natalia Sevane[1], Thomas O´Rourke[2,3], Manuel Pizarro[1], Cedric Boeckx[2,3,4],

Susana Dunner[1]

[1]Universidad Complutense de Madrid, Avenida Puerta de Hierro, s/n, 28040, Madrid, Spain.

[2] Universitat de Barcelona, Gran Vía de les Corts Catalanes 585, 08007, Barcelona, Spain.

[3]UBICS, Carrer Martí Franqués 1, 08028 Barcelona, Spain.

[4]ICREA, Passeig Lluís Companys 23, 08010 Barcelona, Spain.

Corresponding author. Paulig01@ucm.es

Abstract

Aggressiveness is one of the most basic behaviors, characterized by targeted intentional actions

oriented to cause harm. The reactive type of aggression is regulated mostly by the brain's

prefrontal cortex; however, the molecular changes underlying aggressiveness in adults have not

been fully characterized. Here we used an RNA-seq approach to investigate differential gene

expression in the prefrontal cortex of bovines from the aggressive Lidia breed at different age

stages: young three-year old and adult four-year-old bulls. A total of 50 up and 193 down-

regulated genes in the adult group were identified. Furthermore, a cross-species comparative

analysis retrieved 29 genes in common with previous studies on aggressive behaviors,

representing an above-chance overlap with the differentially expressed genes in adult bulls.

Particularly, we detected changes in the regulation of networks such as synaptogenesis, involved

in maintenance and refinement of synapses, and the glutamate receptor pathway, which acts as

excitatory driver in aggressive responses. Our results provide insights into candidate genes and

networks involved in the molecular mechanisms leading to the maturation of the brain. The

reduced reactive aggression typical of domestication has been proposed to form part of a

retention of juvenile traits as adults (neoteny). The significant age-associated differential

expression of genes implicated in aggressive behaviors and concomitant increase in Lidia cattle

aggression validates this species as a novel model comparator to explore the impact of behavioral

neoteny under domestication.

Keywords gene expression, prefrontal cortex, cattle, aggressiveness, behavior genomics

Introduction

Aggression in animals, including humans, serves important purposes in securing mates, territory

and food and, therefore, is one of our most basic behaviors. We understand aggression as an

overt behavior directed at an object or a subject with an intention to cause harm or damage (

Gannon et al. 2009). This definition encapsulates a heterogeneous and multifaceted construct,

including the important distinction between two different subtypes of aggression, reactive and

proactive (Smeets et al. 2017). Reactive aggression is the impulsive response to provocations,

frustrations or threats from the environment. Proactive aggression, is conscious and goal-

directed, planned with a specific outcome in mind (e.g. attainment of resources). While both

subtypes can co-occur, different neurobiological structures have been shown to underlie reactive

and proactive subtypes (Blair, 2013).

Animal studies have shown that reactive aggression is mediated by the limbic system, through a

circuit that runs from the amygdala to the hypothalamus, and from there to the periaqueductal

grey. This circuitry is regulated by frontal cortical regions, particularly by the prefrontal cortex

(PFC) and the anterior cingulate cortex (Blair, 2013). There is strong evidence that reactive

aggression often appears earlier in life than proactive aggression, and that heightened levels of

reactive aggression at early stages are associated with increased levels of anxiety in adulthood. In

humans, childhood reactive aggression can be found in different neuropsychiatric that can lead to

conduct disorders during adulthood (Fite et al., 2010). Therefore, a comparison of gene

expression profiles in PFC tissues at different stages of development may be relevant to reveal

some of the molecular characteristics and mechanisms involved in aggressive-related

perturbations that characterize early stages of aggression and its progression into adulthood.

Research studies across different fields suggest that modern humans have undergone positive

selection for reduced reactive aggression relative to our archaic counterparts. Behavioral

comparisons with our closest extant ape relatives have shown that instances of reactive

aggression in our species are between two and three orders of magnitude lower (Wrangham,

2019). Modern humans also have marked reductions in hypothalamic-pituitary-adrenal (HPA)

axis activity (which regulates stress responses) compared to other primates (Chrousos et al .,

1982). Furthermore, evidence of convergent phenotypical changes in anatomically modern

humans and domesticated species strongly suggest that such reductions in aggressive behaviors

have continued in our recent evolutionary history, marking an important contributor to the

modern—archaic split: domesticated species ubiquitously display reductions in reactive

aggression when compared to their wild counterparts, often accompanied by broad phenotypical

changes known as the ``domestication syndrome'' (reductions in cranial, brain and tooth size,

shortening of muzzles, depigmentation, the development of floppy ears, and attenuated HPA

activity) (Wilkins et al., 2014). Experimentally controlled selection for reduced reactive

aggression in farm-bred foxes brought about many of these broader changes (Dugatkin and Trut,

2017). Modern humans exhibit craniofacial changes relative to Neanderthals reminiscent of those

in domesticated species, including reductions in tooth size, contraction of the skull (in our case

giving a more globularized shape), and flattening of the face (Sánchez-Villagra & van Schaik,

2019; Theofanopoulou et al., 2017). Genes identified within selective-sweep regions in modern

humans have been shown to have above-chance intersection with those implicated in domesticate

evolution, including disproportionate targeting glutamate receptor genes associated with

aggressive behaviors and implicated in regulation of the HPA axis (O'Rourke & Boeckx, 2020;

Theofanopoulou et al., 2017). This evidence for (self-) domestication in ours and domesticated

species has long been considered an instance of retained juvenile (neotenic) traits (see e.g. 5).

This includes behavioral traits such as diminished reactive aggression. A corollary of this

proposal is that Lidia cattle, for which reactive aggression increases with maturity, should have

non-juvenile gene expression patterns.

The Lidia cattle breed is a promising and natural animal model for studying reactive aggression,

given that the breed's agonistic responses have been exacerbated through long-term selection of

aggressive behaviors. In relation to other domesticated cattle breeds, the selection regime that

Lidia cattle have undergone is analogous to that practiced in the farm-fox experiment, where

distinct fox lineages have been bred for tame versus aggressive behaviors. Although not perhaps

the most obvious models of human evolution, genes targeted by selective sweeps in tame cattle

(like dogs and unlike cats and horses) show above-chance intersection with genes falling within

selective sweep regions in modern humans (Theofanopolou et al., 2017). The Lidia is a primitive

breed, raised for centuries with the exclusive purpose of taking part in socio-cultural and

anachronistic events grouped under the term of "*tauromaquias*" (Eusebi et al., 2018). At these

festivities, male bovines are placed on a bullring in solitary with a human subject, resulting in a

series of aggressive reactions that measures the animal´s fighting capacities by means of reactive

responses, such as fighting, chasing and ramming. Different types of *tauromaquia* take place at

the bullrings, perhaps the most popular are those known as major festivities: the "*corrida*" and

the "*novillada*" (Domecq, 2009). The difference between these two events is the age of the

bovines. In a "*corrida*", the protagonists are 4-year-old bulls, while a "*novillada*" uses 3-year-old

bovines (Maudet, 2010). This age discrimination between events is not random; 3-year-old

animals are smaller, less strong and their reactive responses tend to be less explosive and intense

than those of 4 year-old bovines.

Our previous studies using Lidia bovines as a model to study the neurobiological basis of

aggressive behaviors revealed widespread changes in PFC gene expression compared with tamed

Wagyu bovines, a meat-production breed known for their docile-temperament. Thus, we aimed

to provide insights into potential molecular and cortical substrates of aggressiveness (Eusebi et

al., 2020). Furthermore, genome scans for selection signatures in the Lidia cattle breed have

uncovered the existence of several genes strongly associated with aggressive behaviors

previously reported in humans and other animal models (Eusebi et al., 2019). Particularly, we

have associated a novel polymorphism in the promoter of the monoamone oxidase A (*MAOA*)

with different levels of aggressiveness among cattle breeds, a gene that has been clearly linked to

aggression in humans (Brunner et al., 1993; Eusebi et al., 2019).

Although studies characterizing gene expression differences between juveniles and adults in the

PFC have shown promising results in mice (Agoglia et al., 2017; Lander et al., 2017), to our

knowledge, there is a lack of such studies using cattle as a model. Thus, the aim of the present

study is to explore differences in gene expression at two age stages of Lidia cattle, "young"

three-year-old and "adult" four-year-old bovines, given the marked differences in the intensity of

their agonistic responses. For this purpose, gene expression profiles from the PFC tissue of eight

adult and eight young Lidia bovines were compared. The results of the present study will give

insights into the molecular modulation involved in the complex mechanisms leading to the

maturation of the brain, with a focus on aggressive behaviors. From an evolutionary perspective,

comparisons identifying developmental changes in gene expression can inform as to the possible

influences of transcriptional neoteny on the retention of juvenile behaviors into adulthood

(Somel et al., 2009). Domestic animals such as dogs retain playful juvenile behaviors as adults,

and the reduction of reactive aggression under domestication may also result from a more

juvenile neurotranscriptome. Increased aggression of four-year-old Lidia cattle may be expected

to rely on an expression profile involving genes not typically implicated in domestication.

Material and Methods

Ethics statement

No special permits were required to conduct the research. All animals were sacrificed for reasons

other than their participation in this study. Harvesting of bovines´ brains at the cutting room

followed standard procedures approved by the Spanish legislation applied to abattoirs (BOE,

1997).

Animals

Post-mortem PFC tissue samples were retrieved from 16 non-castrated bovine males of the Lidia

breed, eight three-year-old and eight four-year-old, for the young and adult groups respectively.

In standard RNA-seq studies, this level of replication is 2.6-fold higher than the minimum

required (3 individuals/group). Similarly to a classic resident-intruder laboratory test, where an

unfamiliar animal (intruder) is placed in the territory of another animal (resident) resulting

frequently in an aggressive conflict (Koolhaas et al., 2013), each bovine was placed in a bullring

and incited to develop reactive aggressive responses for approximately thirty minutes prior to its

sacrifice, to measure their fighting abilities (Domecq, 2009). Samples of PFC were collected less

than an hour post-mortem and immediately submerged in RNA-later™ (Thermo Fisher

Scientific, Madrid, Spain), followed by 24 hours' storage at 5°C and long-term conservation at -

80°C.

RNA isolation and sequencing

Total RNA was extracted from PFC samples using the RNeasy Lipid Tissue Mini Kit (QIAGEN,

Madrid, Spain), and total RNA concentration and purity were assessed with a Nanodrop ND-

1000 spectophotometer (Thermo Fisher Scientific, Spain). To guarantee their preservation, RNA

samples were treated with RNAstable (Sigma-Aldrich, Spain), and shipped at ambient

temperature to DNA-link Inc. sequencing laboratory (Seoul, Korea) to perform high throughput

sequencing. For quality check, the OD 260/280 ratio was determined to be between 1.87 and 2.0.

All these procedures were conducted according to the respective manufacturers´ protocols.

Individual libraries for each of the analyzed bovines (N=16) were processed using the TruSeq

Stranded mRNA Preparation kit (Illumina, San Diego, CA, USA) according to manufacturer´s

instructions. Each library was pair-end sequenced (2x100 bp). Individual reads were de-

multiplexed using the CASAVA pipeline (Illumina v1.8.2), obtaining the FASTQ files used for

downstream bioinformatics analysis. Illumina reads generated from all samples have been

deposited in the NCBI GEO / bioproject browser database (Accession Number: GSE149676).

Bioinformatics analyses

Quality control and pre-processing of genomic datasets was carried out with the PRINSEQ v.

0.20.4 software (Schmeider and Edwards, 2011). To improve the downstream analysis, low

quality ($Q < 20$) and ambiguous bases (N) were first trimmed from both ends of the reads and,

then, the trimmed reads were filtered by a Phred quality score ($Q \geq 20$ for all bases) and read

length of ≥ 68 bp. The obtained FASTQ files were mapped to the bovine reference genome (Bos

taurus ARS.UCD 1.2) with the STAR Alignment v.2.7.3a software (Dobin et al., 2013), using

default parameters for pair-end reads and including the Ensembl *Bos taurus* ARS-UCD 1.2 as

reference annotation. Once reads were mapped, the expression of each mRNA and their relative

abundance in fragments per kilobase of exon per million fragments mapped (FKPM) were

measured with Cufflinks v.2.2.1 (Dobin et al., 2013). Only expressed transcripts with FKPM

values <0 were used for downstream gene expression analyses, leaving a total of 14 animals

(eight young and six adults).

The analysis of differential gene expression between young and adult groups was performed

using Cufflinks software (Trapnell et al., 2010). The assembled transcripts from all samples were

merged using the Cuffmerge command. Cuffdiff was then used applying default parameters and

the option "fr-firststrand" to define pair-end reads. A Benjamini-Hochberg False Discovery Rate

(FDR), which defines the significance of the Cuffdiff output, was set as threshold for statistically

significant values of the Differentially Expressed Genes (DEG). The results of the Cufdiff RNA-

seq analysis were visualized with the R software application CummeRbund v.2.28.0 (Goff et al.,

2012).

To study the relationship between differences in PFC gene expression among groups and its

biological function, we separated the results of DEG in two independent gene-lists according to

their Log_2 Fold Change (FC) in the adult group: up-regulated for those transcripts displaying a

$Log_2FC \geq 0.1$; and down-regulated for those with a $Log_2FC \leq 0.1$.

Cross Species Comparative Analysis (CSCA)

We conducted a comparison between our DEG and a compendium of genes associated with

aggressiveness and previously identified in different genomic studies in humans, rodents, foxes,

dogs and bovines, as proposed by Zhang-James et al. (Zhang-James et al., 2019). This list is

based on four categories of genomic evidence: a) genome wide association studies (GWAS) in

humans of different age groups, children and adults (Fernández del Catillo and Cormand, 2016);

b) genes showing selection signatures previously identified in the Lidia breed (Eusebi et al.,

2018; , Eusebi et al., 2019); c) RNA-seq studies in rodents (Clinton et al., 2011; ,Malki et al.,

2014) and silver foxes (Kukekova et al., 2011, Kukekova et al., 2018); and d) genes identified in

studies with causal evidence of the Online Mendelian Inheritance in Man (OMIM) and the

Online Mendelian Inheritance in Animals (OMIA) databases, and a report of knockout (KO)

studies in mice (Våge et al., 2010; Veroude et al., 2106; Zhang-James et al., 2019). The gene-list

and details of the different studies are described in Supplementary Table 1. Bovine official gene

names were converted to their human orthologues using biomaRt (Durinck et al., 2005) in order

to homogenize our DEG with the cross-species gene list. We assigned a weight value (weighted

ranking, WR) with the same conditions proposed by Zhang-James et al. (2019). For statistical

analysis of the intersection between DEGs and genes identified in studies of aggression, we

cross-referenced each gene list using the PANTHER (www.pantherdb.org), NCBI

HomoloGene,(www.ncbi.nlm.nih.gov/homologene) and Ensembl orthologue databases *Bos*

taurus ARS-UCD 1.2 and Human reference (GRCh38.p13) genomes. If no human–bovine one-

to-one orthologues were found in any database, we removed the relevant genes for statistical

analysis.

Ingenuity pathway analysis

Up and down-regulated DEG lists were imported into the Ingenuity Pathway Analysis (IPA)

(QIAGEN, www.qiagen.com/ingenuity) software to assess Gene Ontologies (GOs) and

canonical pathways enrichment scores. Additionally, up and down-regulated DEGs in common

with the CSCA were imported together to build biological networks and identify upstream

regulators.

For the canonical pathway analyses, a right-tailed Fisher's exact test was performed for the

enrichment of the DEGs in IPA´s hand-curated canonical pathway database. Here, the P-value

calculated for a pathway measures the probability of being randomly selected from all of the

curated pathways. To control the error rate in the analysis, IPA also provide the corrected P-

values to identify the most significant results in IPA's canonical pathways based on the

Benjamini-Hochberg method (Benjamini and Hochberg, 1995). This tool allowed us to identify

the signaling pathways in which the DEGs were enriched. We used the predetermined cut-off of

the corrected P-value < 0.05 to define the significant pathways.

For network enrichment and detection of upstream regulators, the genes in common with the

CSCA were overlaid onto a global molecular network (GMN) developed based on the Ingenuity

Pathways Knowledge Base, in which functional relationships such as activation, chemical-

protein interaction, expression, inhibition and regulation of binding were manually searched.

Sub-networks of genes were then extracted from the GMN based on their connectivity using the

algorithm developed by IPA. The sub-networks were visualized using IPA´s Path Designer tool.

Results

Sequencing and read assembly

For the 16 individually sequenced samples, we generated an average of 61.6 million pairs of

100-bp paired-end reads per sample after filtering by quality score. For each sample, an average

of 91.6% reads were uniquely mapped to the bovine reference genome, ranging from 86.8 to

95.2% (Supplementary Table 2). The transcripts expressed in both the young and adult groups

reaching FKPM values < 0 (N=8 young and N=6 adult) were included in the differential

expression and subsequent IPA analyses (Supplementary Figure 1).

Gene expression differences revealed a total of 27,153 DEGs between young and adult groups.

Of those genes, only 243 were statistically significant; 50 up-regulated ($\log_2$FC $\geq$ 0.1) and 193

down-regulated ($\log_2$FC $\geq$ -0.1) (Figure 1A and B). For the detailed list of up and down-

regulated DEGs see Supplementary Table 3.

A) Up-regulated DEG   **B) Down-regulated DEG**

$\log_{10}$ FPKM + 1

Young   Adult   Young   Adult
(Three-year-old) (Four-year-old)   (Three-year-old) (Four-year-old)

Figure 1. Heatmap of up-regulated (A) and down-regulated (B) differentially expressed genes

(DEG) in the adult group.

Genes in common with the cross-species comparative analysis (CSCA)

The cross-reference analysis between our up and down-regulated DEGs in the adult group and

the gene-list associated with aggressive behavior, overlapped at numerous loci. Notably, the

down-regulated DEGs shared 21 genes, while the up-regulated displayed only 8 genes in

common with the gene reference list (Table 1). The subset of DEGs included in the CSCA are

shown in the expression bar plot of Figure 2, which details their standard deviation of FPKM

values. As well as these 29 genes, a many-to-many orthologous match was found between the

bovine non-classical major histocompatibility complex class I antigen (BOLA-NC1) and the

human equivalents (HLA-A, HLA-B, HLA-C, and HLA-F).

Table 1. Up and down regulated DEGs in four-year-old bulls in common with the cross-species

comparative analysis (CSCA).

Up-regulated DEGs			Down-regulated DEGs		
Gene symbol	**Gene Name**	**Weighted Ranking (WR)**	**Gene symbol**	**Gene Name**	**Weighted Ranking (WR)**
NPAS4	Neuronal PAS Domain Protein 4	1.5	*IGF2*	Insulin Like Growth Factor 2	1
PENK	Proenkephalin	1.5	*PAMR1*	Peptidase Domain Containing Associated With Muscle Regeneration 1	1
FMN1	Formin 1	1.5	*ROBO1*	Roundabout Guidance Receptor 1	1
SLC24A2	Solute Carrier Family 24 member 2	1	*SCARA5*	Scavenger Receptor Class A Member 5	1
SEL1L3	SEL1L Family Member 3	1	*SPARC*	Secreted Protein Acidic And Cysteine Rich	1
VGF	VGF Nerve Growth Factor Inducible	1	*DACT2*	Dishevelled Binding Antagonist Of Beta Catenin 2	1
LGI2	Leucine Rich Repeat LGI Family Member 2	0.5	*ALCAM*	Activated Leukocyte Cell Adhesion Molecule	0.5
BRINP1	BMP/Retinoic Acid Inducible Neural Specific 1	0.5	*KCNG1*	Potassium Voltage-Gated Channel Modifier Subfamily G Member 1	0.5
			TIMP3	TIMP Metallopeptidase Inhibitor 3	0.5
			GJA1	Gap Junction Protein Alpha 1	0.5
			ADGRA2	Adhesion G Protein-Coupled Receptor A2	0.5

TP53I11	Tumor Protein P53 Inducible Protein 11	0.5
DCN	Decorin	0.5
HES5	Hes Family BHLH Transcription Factor 5	0.5
HTR2C	5-Hydroxytryptamine Receptor 2C	0.5
FRZB	Frizzled Related Protein	0.5
HACD4	3-Hydroxyacyl-CoA Dehydratase 4	0.5
NUPR1	Nuclear Protein 1, Transcriptional Regulator	0.5
CRIM1	Cysteine Rich Transmembrane BMP Regulator 1	0.5
ZFP36L1	ZFP36 Ring Finger Protein Like 1	0.5
CRISPLD1	Cysteine Rich Secretory Protein LCCL Domain Containing 1	0.5

Figure 2. Bar chart of up and down regulated DEGs in common with the cross-species

comparative analysis (CSCA). Gene abundance is represented in fragments per kilobase of exon

per million fragments mapped (FPKM) of the adult (4-year-old) and young (3-year-old) groups.

Statistical analysis of aggression-associated differentially expressed genes (DEG)

In order to test whether the 29 adult Lidia DEGs identified in the CSCA represent a statistically

significant association between maturation and aggression, we calculated the cumulative

hypergeometric probability of this overlap occurring. Following removal of non-orthologous

genes, 1701 aggression-associated genes across species and 238 Lidia DEGs remained. Taking

the estimated 22,000 genes in the bovine genome (Elsik et al., 2009), an overlap of at least 29

between these two gene samples was significantly unlikely to occur (p-value: 0.0099). We

carried out the same analysis taking only aggression-associated genes that were identified in

brain-expression studies across the different species (totaling 1151). Of the 238 Lidia DEGs, 21

were also identified in these studies (p-value: 0.0137). Under a more restrictive analysis,

whereby only cortically expressed genes are taken as potential comparators ─ estimated at 85%

of all protein-coding genes in both humans and pigs (Sjöstedt et al., 2020) ─ , the intersection of

21 was slightly more likely to have occurred by chance (p-value: 0.0617).

Ontological analysis of the differentially expressed genes (DEG)

The up and down-regulated DEGs in the 4-year-old adult group were subjected to a Fisher's

exact test within IPA to obtain Ingenuity ontology annotations. IPA annotations follow the GO

annotation principle, but are based on a proprietary knowledge base of over a million protein-

protein interactions. Significant results are summarized in Supplementary table 4. The most

relevant results for the up-regulated DEGs were obtained under the *physiological system*

development and the *disease and disorders* categories. Within the first of these categories, the

top of the list gathered terms related with *behavior* (highest p-value range of 5.51E-08 and 15

DEG) and *nervous system development and function* (highest p-value range of 7.30E-07 and 24

DEG), while within the category of *diseases*, terms related to *cancer* (highest p-value range of

1.21E-07 and 44 DEGs), *neurological diseases* (highest p-value range of 8.07E-07 and 21

DEGs) and *psychological disorders* (highest p-value range of 3.35E-06 and 12 DEGs) occupied

the top positions.

Additionally, we explored the overrepresentation of specific enriched functions within the

category of *behavior* and its relationship with diseases or function annotations, finding high

overall significance levels of association between up-regulated DEGs and diverse behavioral

conditions (p-values $< 10^{-5}$) (Table 2, Figure 3).

Table 2. Overrepresentation of enriched specific functions within the category of *behavior*. N:

Number of up-regulated DEGs in the adult group.

Disease / function annotation	P-value	N	Genes
Emotional behavior	5.51E-08	9	*BRINP1,CBLN4,GRIN2A,PENK,SCN1B,SLC17A6, SCN4B,SCN1B,SCN1A,VGF*
Learning	3.27E-06	9	*ABCC8,BRINP1,DKK1,GRIN2A,NPAS4,SCN1A,SLC4A2,TAFA2,VGF*
Memory	7.70E-07	8	*ABCC8,BRINP1,DKK1,GRIN2A,NPAS4,SCN1A,SLC4A2,TAFA2*
Anxiety	1.75E-05	6	*BRINP1,CBLN4,GRIN2A,PENK,SCN1A,TAFA2*
Spatial learning	7.09E-05	5	*ABCC8,GRIN2A,SLC24A2,TAFA2,VGF*
Irritable behavior	3.35E-06	4	*GRIN2A,SCN1A,,SCN1B,SCN4B*

Figure 3. Regulator effects of the IPA package applied to the up-regulated DEG and diverse

behavioral conditions (p-values $< 10^{-5}$). In the lower tier, the expected behavioral consequences

of the up-regulated DEG changes are shown by considering the Ingenuity Knowledge Base. In

orange color are marked those functions predicted to be activated and in blue those predicted to

be suppressed.

The Core Analysis of the down-regulated DEGs in the 4-year-old group also revealed genes

related to the *physiological system development* and the *disease and disorders* categories. The

top enriched genes in the first category were associated with *organismal* (highest p-value range

of 2.38E-17 and 63 DEGs), *embryonic* (highest p-value range of 2.38E-17 and 58 DEGs) and

cardiovascular system (highest p-value range of 1.08E-16 and 43 DEGs) development functions.

Furthermore, at the *diseases and disorders* category the down-regulated DEGs were shown to be

involved with *cardiovascular disease* (highest p-value range of 3.05E-11 and 34 DEGs), *cancer*

(highest p-value range of 1.73E-10 and 84 DEGs), and *organismal injury and abnormalities*

(highest p-value range of 1.73E-10 and 84 DEGs) (Supplementary Table 4).

Metabolic pathways affected by the up and down-regulated DEGs in the adult group

Within the up-regulated DEGs in the adult 4-year-old group, four pathways surpassed the

significance threshold (p-values < 0.05, Supplementary Figure 2). The *synaptogenesis* and the

Wnt/β- catenin signaling are among the most significant pathways (p-value = 3.08E-03).

Besides, the *glutamate receptor pathway*, which mediates synaptic signaling by this primary

excitatory neurotransmitter in the central nervous system (CNS), resulted also highly significant

(p-value = 5.74E-03).

The analysis of down-regulated pathways revealed the involvement of the retrieved DEGs on 53

metabolic routes (Supplementary Table 5). Amongst the most significant, the *intrinsic*

prothrombin activation pathway (p-value =1-08E-08) and, noteworthy, again the *Wnt/β- catenin*

signaling (p-values = 4.76E-05) route, are highlighted within the down-regulated DEGs in the

adult group (Supplementary Figure 1).

Gene networks and upstream activator analyses of the genes in common with the CSCA

We analyzed the gene transactivation networks and upstream activators of the 29 DEGs with

one-to-one orthologous matches in the CSCA, finding three highly interconnected networks

related to organ morphology, organismal development, behavior, cancer, neurological disease

and psychological disorders (Figure 4).

Finally, the upstream analysis tool of the IPA package was employed to identify potential

upstream regulators that may explain the differential patterns of expression between young and

adult groups. We explored the activation/inhibition states of the top upstream regulators by using

IPA´s activation z-score tool, identifying four main upstream regulators: KRAS proto-oncogene

(*KRAS),* Brain Derived Neurotrophic Factor *(BNDF),* CAMP Responsive Element Binding

Protein 1 *(CREB1)* and SRY-Box Transcription Factor 2 *(SOX2).* As shown in the graphic

network in Figure 5, these genes are involved in an array of diverse biological functions related

with behavioral development.

Figure 4. Network analysis of the 29 DEGs in the adult group in common with the CSCA.

Genes highlighted in color correspond to the DEGs in our study; in red the up-regulated and in

green the down-regulated DEGs.

Figure 5. Network visualization of the four major upstream regulators of the DEGs in common

with the CSCA. In orange, upstream regulators predicted to be activated; in red and green, genes

whose expression increases or decreases in response to the activation of upstream regulation,

respectively. The shapes of the nodes reflect the functional class of each gene product, as shown

in the legend. The symbols marked in grey define functions. Solid and dashed lines between

genes represent known direct and indirect interactions, respectively, with orange lines leading to

activation and blue to inhibition.

Discussion

The structural and molecular changes occurring in the PFC of diverse species at different age

stages have been previously explored (Counotte et al., 2010; , Moczulska et al., 2014; ,

González-Lozano et al., 2016). The majority of these analyses conducted in rodents focus on the

study of the molecular changes occurring in post-natal and puberty-adolescence stages.

However, studies on the specific changes of genetic expression during adulthood are still scarce

(12, Agoglia et al., 2017, Lander et al., 2017). Here, we used RNA-seq to identify genes and

gene networks that show differential expression in the PFC of adult bulls when compared to

young bovines of the aggressive Lidia breed.

A total of 143 significant DEGs, 50 up and 193 down-regulated in the adult group were

identified. The GO analysis of up-regulated DEGs in the adult group rendered highly significant

results for behavioral and nervous system development features, while the down-regulated genes

were mostly implicated in systemic developmental functions (organismal, embryonic and

cardiovascular system). We detected a modest coincidence between these results and what has

been published in previous analysis using high-throughput molecular data in PFC. For instance,

Agoglia et al. (2017) used an unbiased proteomics approach to analyze the differential

expression of proteins in adolescent compared to adult PFC in mouse. There is a concordance

between the proportion of up and down-regulated DEGs in the adult groups reported by both, us

and Agoglia et al. (2017), as well as results from the GO analyses showing down-regulation of

organismal growth and development processes in adults. This suggests that, both technologies,

proteomics and RNA-seq are, as awaited, correlated, detecting similar sets of DE molecules (40).

Canonical pathway analyses also revealed that metabolic routes involved in cellular signaling

and neurotransmission impact in the adult PFC during the aging process (Figure 2). We detected

an over representation of the synaptogenesis and the glutamate signaling reception pathways in

the four-year-old bulls, both involved in brain´s synaptic plasticity; the synaptogenesis pathway

is implicated in developmental processes of formation, maintenance and refinement of synapses

(Cohen-Cory, 2002), while the glutamate signaling receptor acts as excitatory driver of the

Hypothalamus-Pituitary-Adrenal (HPA) system, a complex system that triggers reactive

aggression responses (Herman et al., 2004; Evanson and Herman, 2015). Similar results were

obtained by Lander et al. (2017); they detected an over expression in adults of glutamate markers

comparing the cortical mRNA expression of mid-adolescent and adult mice.

Most of the genes making up the glutamate receptor signaling pathway are implicated in NMDA

and AMPA-mediated excitation driving long term potentiation at mature synapses. This contrasts

with findings that metabotropic and kainate receptor genes ─ which tend to down-regulate

glutamanergic excitation at synapses, including in limbic circuits that control HPA activity ─

are disproportionately implicated in domestication and modern human evolution (O'Rourke &

Boeckx, 2020).

In our recent comparison between the Lidia and Wagyu we noted evidence for down-regulated

glutamate-to-GABA conversion in the more aggressive breed, suggesting heightened excitatory

signaling (Eusebi et al., 2020). In previous studies of selective sweeps distinguishing the Lidia

from tamer breeds, we also noted differential signals of selection on the domestication-associated

glutamate receptor gene *GRIK3*, further suggesting that heightened excitatory signaling may play

a role in the increased reactive aggression of this breed (Eusebi et al., 2018). In future studies it

may be worthwhile to compare the age-dependent PFC expression profile of tame cattle breeds

with that of the Lidia. Under the hypothesis that increased excitatory signaling contributes to

driving aggressive behaviors, one may expect that genes tending to contribute to a relative down-

regulation of glutamatergic signaling or delayed maturation of excitatory synapses may be

targeted in tamer breeds.

Interestingly, we detected differences in the expression of genes included in the Wnt//β- catenin

signaling pathway in both up and down-regulated DEGs in the adult group. The complex

interactions of genes, with actors triggering the expression of the Wnt//β- catenin signaling

pathway while others are needed to silence routes leading to its suppression, justifies the

inclusion of genes from this network in the up and down-regulated cohorts. This pathway is

considered one of the most strongly conserved routes among species and is involved in synaptic

transmission activities (Maguschak and Ressler, 2012). Malfunction of many components of the

Wnt//β- catenin signaling pathway in the adult brain can result in altered behavior and cognitive

disorders (Svenningsson et al., 2002).

In addition, the prothrombin activation pathway acts as signaling route at the CNS, inducing the

decrease of cyclic AMP (CAMP) and the activation of protein kinase C (PKC), both enzymes

linked to catecholamine and serotonin receptors (David et al., 2004). It has been proposed that,

during encounters between individuals, a de-synchronization of the top-down control at these

receptors influence a lack of control of emotional reactivity (Kretshcmer et al., 2014). As

suggested by Agoglia et al. (2017), the increased expression on these routes observed in the 4-

year-old PFC may contribute to the fine-tuning of synaptic transmission, resulting in a superior

control of the PFC executive functioning in mature brains, intensifying the addressed aggressive

behaviors.

The significant overlap between the age-associated DEGs we identified and the genes associated

with aggression in other species strongly suggests an important influence of maturation on the

development of aggressive traits in Lidia cattle. This contrasts with the retention of a docile

temperament from adolescence to adulthood in other cattle breeds and marks the Lidia as a

viable model for research into the neurobiological basis of behavioral neoteny under

domestication. Complementarily to these results, the IPA network tool applied to the 29 DEGs

with one-to-one orthologues in the CSCA, retrieved three inter-connected networks associated

with organismal morphology and development, behavior, cancer, neurological disease and

psychological disorders (Figure 3). Particularly, the Mitogen Activated Protein Kinase (*MAPK*)

and two Extracellular Signal Regulated Kinases (*ERK*) appear to be involved in diverse

molecular mechanisms underlying aggression. Similar results were obtained in previous studies

on PFC gene expression (Zhang-James et al., 2019; Eusebi et al., 2020), identifying a differential

expression of the ERK/MAPK signaling pathway in their comparisons between aggressive and

non-aggressive individuals.

At the regulatory level, three of the four genes predicted to be activated display clear roles in

aggressive behaviors, including *BDNF* (Kretshcmer et ak., 2014;,Vigers et al., 2012), *CREB*

(David et al., 2004) and *SOX2* (Gatewood et al., 2006). The behavioral functions affected by

these regulators include learning and memory cognitive processes, along with behavioral

conditions and neuropsychiatric disorders, such as fear, addictive behaviors, schizophrenia and

bipolar disorder, all displaying aggressiveness (Figure 4). In this regard, there is evidence of

increased memory encoding during early adulthood (Rubin et al., 1998) and also, it has been

observed that adults maintain elevated plasma levels of molecules associated with abnormal

aggressive behaviors (i.e. corticosterone) longer than adolescents (Romeo et al., 2006).

Furthermore, some neural receptors at adult PFC have shown heightened expression levels,

conferring them higher risk of addictive and impulsive behaviors (Sonntag et al., 2014). This

evidence suggest that behavioral and physiological functional processes of adults differs from

that of adolescents, which may reflect the ongoing maturation of the limbic structures that

regulate reactive aggression responses in adulthood.

Furthermore, the presence of *KRAS* gene as upstream regulator was also observed. Although

KRAS is a proto-oncogene associated principally with cancer (Zhu et al., 2008), at the

transcriptional level it acts as a short-term inductor to astrocytes in response to stimulus and as a

sensor that adapts cells to metabolic needs and oxidative stress at the brain (Messina et al.,

2017). The reactive aggression responses of the bulls during the *corrida* may trigger the

mechanisms implicated in oxidative stress states, which have been observed to increase with age

due to a disruption in the oxidant and anti-oxidant balance (Sies, 1991). Besides, in

neuropsychiatric disorders such as bipolar disorder, elevated markers in blood of oxidative stress

levels has been reported among adults, relating oxidative stress as mediator on neuropathological

processes of adulthood (Andreazza et al., 2008).

Finally, it is worth also highlighting two genes implicated in functional development functions:

the *NPAS4* gene, which participates in the structural and functional plasticity of neurons acting

as regulator of the excitatory-inhibitory neural balance (Jaehne et al., 2015); and the *HTR2C*

gene, modulator of the neural activity by functioning as down-stream effector of insulin

sensitivity and glucose homeostasis (Stam et al., 1994). Both genes have also been candidates in

studies of human pathophysiology in mental disorders, including schizophrenia, bipolar and

cognitive disorders (Iwamoto et al., 2009), and are up (*NPAS4*) and down-regulated (*HTR2C*) in

the adult group of bulls (Table 2).

Interestingly, whereas Agoglia et al. (2017) reported enhanced protein alterations in networks

that regulate neuronal signaling, anxiety-related behavior and neurological disease in the

adolescent PFC when compared to adult mice, our gene expression results found similar

associations but only in adults. This different outcome may be due to differences in the age-stage

comparison between studies. While in mice the biological transition of adolescence to adulthood

is well studied, defining maturation stages in cattle has proved more complex, as its lifespan

depends on the production purpose of each breed, among other factors (Essl, 1998). Given that

the average life of a Lidia sire is approximately 15 years, and the average generation sire interval

pathway (sire-offspring) is 7,5 years (Fernández-Salcedo, 1990), perhaps here the comparison

between early and middle age adulthood would be a more correct approximation.

The present findings suggest that the cortical gene expression in 4-year-old Lidia cattle is

characterized by an enhancement of molecular mechanisms involved in the increase of

aggressive behaviors and neurophysiological disorders. However, despite our results suggesting

cortical enhanced genetic expression associated with reactive aggression in adults, this may not

be representative of other brain regions within the limbic system reported also as important

actors in this process. Future experiments to evaluate the role of these other brain regions,

particularly in the context of aggressive responses, will shed further light on this age transition

period.

Acknowledgements

The authors are grateful to the enterprise of Las Ventas, bullring "*Plaza 1*" and its veterinarian's

team for providing post-mortem Lidia breed brain tissue samples for this study.

References

• Agoglia A. E., Holstein, S. E., Small, A. T., Spanos, M., Burrus, B. M., & Hodge, C. W.

2017. Comparison of the adolescent and adult mouse prefrontal cortex proteome. PloS

one, 12(6). https://doi.org/10.1371/journal.pone.0178391

• Andreazza A. C., Kauer-Sant'Anna M., Frey B. N., Bond D. J., Kapczinski F., Young L. T., &

Yatham, L. N. 2008. Oxidative stress markers in bipolar disorder: a meta-analysis. *J affect*

disord, 111(2-3), 135-144. https://doi.org/10.1016/j.jad.2008.04.013

• Benjamini Y., & Hochberg, Y. 1995. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. J R stat Methodol 57(1); 289-300.

https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

• Blair R. J. R. 2013. The neurobiology of psychopathic traits in youths. Nat Rev Neurosci.

14(11), 786-799. https://doi.org/10.1038/nrn3577.

• Boletin Oficial del Estado BOE-A-1997-3081 ley 10/1997. 1997. Reglamento general de

mataderos. Boletin Oficial del estado.

• Brunner H. G., Nelen M., Breakefield X. O., Ropers H. H., & Van Oost B. A. 1993.

Abnormal behavior associated with a point mutation in the structural gene for monoamine

oxidase. Anim Sci. 262(5133): 578-580. DOI: 10.1126/science.8211186

• Chrousos G. P., Renquist D., Brandon D., Eil C., Pugeat M., Vigersky R., Cutler Jr G.B.,

Loriaux D.L. & Lipsett, M. B. 1982. Glucocorticoid hormone resistance during primate

evolution: receptor-mediated mechanisms. Proc Natl Academ Sci, 79(6), 2036-2040.

https://doi.org/10.1073/pnas.79.6.2036

• Clinton S.M., Stead J.D., Miller S., Watson S.J. &Akil H. 2011. Developmental

underpinnings of differences in rodent novelty-seeking and emotional reactivity. Eur J

Neurosci. 34:994–1005. https://doi.org/10.1111/j.1460-9568.2011.07811.x

• Cohen-Cory S. 2002. The developing synapse: construction and modulation of synaptic

structures and circuits. *Science*. *298*(5594), 770-776. https://doi.org/10.1186/1471-2164-13-

629

• Counotte D. S., Li, K. W., Wortel, J., Gouwenberg, Y., Van Der Schors, R. C., Smit, A. B., &

Spijker, S. 2010. Changes in molecular composition of rat medial prefrontal cortex synapses

during adolescent development. Europ J Neurosci, 32(9), 1452-1460.

https://doi.org/10.1111/j.1460-9568.2010.07404.x

• David J. T., Cervantes, M. C., Trosky, K. A., Salinas, J. A., & Delville, Y. 2004. A neural

network underlying individual differences in emotion and aggression in male golden

hamsters. *Neurosci*, *126*(3), 567-578. https://doi.org/10.1016/j.neuroscience.2004.04.031

• Dobin, A, Davis C.A., Schlesinger, F., Drenkow J., Zaleski C., Jha S., Chaisson M., &

Gingeras T.R. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*. 29(1):15–21

https://doi.org/10.1093/bioinformatics/bts635

• Domecq J. P. 2009. Del toreo a la bravura. Madrid, Spain: Alianza Editorial.

• Dugatkin L. A. & Trut L. N. 2017. *How to tame a fox (and build a dog): visionary scientists*

and a Siberian tale of jump-started evolution. University of Chicago Press.

• Durinck S., Moreau Y., Kasprzyk A., Davis S., de Moor B., Brazma A. & Huber W. 2005.

BioMart and bioconductor: a powerful link between biological databases and microarray data

analysis. Bioinformatics. 21:3439–40.

• Essl A. 1998. Longevity in dairy cattle breeding: a review. *Livestock Production Science*.

57(1), 79-89. https://doi.org/10.1016/S0301-6226(98)00160-2

• Elsik, C. G., Tellam, R. L., & Worley, K. C. (2009). The genome sequence of taurine cattle: a

window to ruminant biology and evolution. Science, 324(5926), 522-528. DOI:

10.1126/science.1169588

• Eusebi P.G., Cortés O., Dunner S. & Cañón J. (2018) Detection of selection signatures for

agonistic behavior in cattle. J Anim Breed Genet. 135:170-177. DOI: 10.1111/jbg.12325

• Eusebi P.G, Sevane N. Pizarro M. &Dunner S. 2020. Gene expression profiles underlying

aggressive behavior in the prefrontal cortex of cattle. Biorxiv.

https://doi.org/10.1101/2020.07.09.194704

• Eusebi P.G., Sevane N., Cortés O., Contreras E., Cañon J. & Dunner S. 2019. Aggressive

behavior in cattle is associated with a polymorphism in the MAOA gene promoter. Anim

Genet. 51:14-21. https://doi.org/10.1111/age.12867

• Evanson N .K., & Herman J.P. 2015. Role of paraventricular nucleus glutamate signaling in

regulation of HPA axis stress responses. Interdiscip inf sci. 21(3); 253-260.

https://doi.org/10.4036/iis.2015.B.10

• Fernández del Castillo N, Cormand B. 2016. Aggressive behavior in humans: genes and

pathways identified through association studies. Am J Med Genet part B Neuropsych Genet.

171:676–96.   https://doi.org/10.1002/ajmg.b.32419

• Fernández-Salcedo Luis. 1990. Veinte toros de Martínez. Madrid, Spain. Egartorre editorial.

• Fite P. J., Raine A., Stouthamer-Loeber M., Loeber,R., & Pardini D. A. 2010. Reactive and

proactive aggression in adolescent males: Examining differential outcomes 10 years later in

early adulthood. Crim Justice Behav. 37(2), 141-157.

https://doi.org/10.1177/0093854809353051

• Gannon T. A., Ward T., Beech A. R., & Fisher D. 2009. Aggressive Offenders' Cognition:

Theory, research, and practice (Vol. 43). John Wiley & Sons.

• Gatewood J. D. et al. 2006. Sex chromosome complement and gonadal sex influence

aggressive and parental behaviors in mice. *J Neurosci*. *26*(8), 2335-2342. DOI:

https://doi.org/10.1523/JNEUROSCI.3743-05.2006

• Goff L., Trapnel C. & Kelley, D. 2012. cummeRbund: Analysis, exploration, manipulation,

and visualization of Cufflinks high-throughput sequencing data. R package version 2.6.1.

• Gonzalez-Lozano M. A., Klemmer P., Gebuis T. Hassan Ch., Nierop P., van Kesteren R.E.,

Smit A.B. & Li K.W. 2016. Dynamics of the mouse brain cortical synaptic proteome during

postnatal brain development. Sci rep, *6*, 35456. https://doi.org/10.1038/srep35456

• Herman J.P., Mueller N.K. & Figueiredo H. 2004. Role of GABA and glutamate circuitry in

hypothalamo-pituitary-adrenocortical stress integration. Ann NY Acad Sci. 1018:35-

45. https://doi.org/10.1196/annals.1296.004

• Iwamoto K., Bundo M., & Kato T. 2009. Serotonin receptor 2C and mental disorders:

genetic, expression, and RNA editing studies. *RNA biology*, *6*(3), 248-253.

https://doi.org/10.4161/rna.6.3.8370

• Jaehne E. J., Klarić T. S., Koblar S. A., Baune B. T., & Lewis M. D. 2015. Effects of Npas4

deficiency on anxiety, depression-like, cognition and sociability behaviour. *Behav b res 281*,

276-282. https://doi.org/10.1016/j.bbr.2014.12.044

• Kogenaru S., Qing Y., Guo Y. & Wang N. 2012. RNA-seq and microarray complement each

other in transcriptome profiling. BMC Genomics. 13, 629. https://doi.org/10.1186/1471-2164-

13-629

• Koolhaas J. M., Coppens C. M., de Boer S. F., Buwald, B., Meerlo P., & Timmermans P. J.

2013. The resident-intruder paradigm: a standardized test for aggression, violence and social

stress. J Vis Exps. 77: e4367. doi: 10.3791/4367

• Kretschmer T, Vitaro F, Barker ED. 2014. The Association Between Peer and own

Aggression is Moderated by the Val-met Polymorphism. *J Res Adolesc*.; 24(1):177–85

https://doi.org/10.1111/jora.12050

• Kukekova A. V. et al. 2011. Sequence comparison of prefrontal cortical brain transcriptome

from a tame and an aggressive silver fox (Vulpes vulpes). BMC genomics.12(1), 482.

https://doi.org/10.1186/1471-2164-12-482

• Kukekova A. V. et al.   2018. Red fox genome assembly identifies genomic regions associated

with tame and aggressive behaviours. Nat Ecol & Evol. 2(9), 1479-1491.

https://doi.org/10.1038/s41559-018-0611-6

• Lander S. S., Linder-Shacham D., & Gaisler-Salomon, I. 2017. Differential effects of social

isolation in adolescent and adult mice on behavior and cortical gene expression. Behav brain

Res. 316, 245-254. https://doi.org/10.1016/j.bbr.2016.09.005

• Maguschak K. A., & Ressler K. J. 2012. A role for WNT/β-catenin signaling in the neural

mechanisms of behavior. J Neuroimmune Pharm 7(4), 763-773.

https://doi.org/10.1007/s11481-012-9350-7

• Malki K. Pain O, Du Rietz E., Tosto M.G., Paya-Cano J., Sandnabba K.N., de Boer S.,

Scjalkwyk L.C. & Sluyter F., 2014. Genes and gene networks implicated in aggression related

behaviour. Neurogenet. ;15: 255–66. DOI 10.1007/s10048-014-0417-x

• Maudet J.B. 2010. Terres de taureaux: les jeux taurins de LʹEurope à LʹAmerique. Madrid:

Casa de Velázquez.

• Messina S., Di Zazzo E., & Moncharmont B. 2017. Early and late induction of KRAS and

HRAS proto-oncogenes by reactive oxygen species in primary astrocytes. *Antioxidants 6*(3),

48. https://doi.org/10.3390/antiox6030048

• Moczulska K. E., Pichler, P., Schutzbier M., Schleiffer A., Rumpel S., & Mechtler K. 2014.

Deep and precise quantification of the mouse synaptosomal proteome reveals substantial

remodeling during postnatal maturation. J proteome res. 13(10), 4310-4324.

https://doi.org/10.1021/pr500456t

• O'Rourke T., & Boeckx, C. (2020). Glutamate receptors in domestication and modern human

evolution. Neurosci Biobehav Rev. 108, 341-357.

https://doi.org/10.1371/journal.pone.0185306

• Romeo R. D., Bellani R., Karatsoreos I. N. et al. 2006. Stress history and pubertal

development interact to shape hypothalamic-pituitary-adrenal axis

• Rubin D. C., Rahhal T. A., & Poon L. W. 1998. Things learned in early adulthood are

remembered best. *Memory & cognition*, *26*(1), 3-19. https://doi.org/10.3758/BF03211366

• Sánchez-Villagra M. R., & van Schaik C. P. 2019. Evaluating the self-domestication

hypothesis of human evolution. Evol Anthropol, 28(3), 133-

143. https://doi.org/10.1002/evan.21777

• Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets.

Bioinformatics. 2011:27: 863–864. https://doi.org/10.1093/bioinformatics/btr026

• Smeets K. C., Oostermeijer S., Lappenschaar M., Cohn M., Van der Meer J.M.J., Popma A.,

Jansen L.M.C., Rommelse N.N.J., Scheepers F.E. & Buitelaar J.K. 2017. Are proactive and

reactive aggression meaningful distinctions in adolescents? A variable-and person-based

approach. J. Abnor Child Psychol. 2017;45(1): 1-14. (doi:10.1007/s10802-016-0149-5).

• Sies H. 1991. Oxidative stress: from basic research to clinical application. *Am j med*, 91(3),

S31-S38. https://doi.org/10.1016/0002-9343(91)90281-2

• Sjöstedt E, Zhong W, et al., 2020. An atlas of the protein-coding genes in the human, pig,

and mouse brain. Science 367(6482) DOI: 10.1126/science.aay5947

- Somel, M. et al . 2009. Transcriptional neoteny in the human brain. *Proc Natl Academ Sci*, *106*(14), 5743-5748. https://doi.org/10.1073/pnas.0900544106

- Sonntag K. C., Brenhouse H. C., Freund N., Thompson B. S., Puhl M., & Andersen S. L. 2014. Viral over-expression of D1 dopamine receptors in the prefrontal cortex increase high-risk behaviors in adults: comparison with adolescents. *Psychopharmacology*. *231*(8), 1615-1626. https://doi.org/10.1007/s00213-013-3399-8

- Stam N. J., Vanderheyden P., van Alebeek C., Klomp J., de Boer T., van Delft A.M.L & Olijve W. 1994. Genomic organisation and functional expression of the gene encoding the human serotonin 5-HT2C receptor. *Eur J Pharmacol: Mol Pharmacol*.*269*(3), 339-348. https://doi.org/10.1016/0922-4106(94)90042-6

- Svenningsson P. Tzavara, E. T., Liu, F. Allen A., Fienber A.A., Nomikos G.G. & Greengard P.2002. DARPP-32 mediates serotonergic neurotransmission in the forebrain. Proc Natl Acad Sci. 99(5): 3188-3193. https://doi.org/10.1073/pnas.052712699

- Theofanopoulou C., Gastaldon S., O'Rourke T., Samuels B. D., Messner A., Martins P. T., Delogu F., Alamri S., & Boeckx C. 2017. Self-domestication in Homo sapiens: Insights from comparative genomics. *PloS one*, *12*(10), e0185306. https://doi.org/10.1371/journal.pone.0185306

- Trapnell C., Williams B.A., Pertea et al. 2010. Transcript assembly andquantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. Nat Biotechnol. 28: 511–515. doi: 10.1038/nbt.1621

- Våge J., Bønsdorff T. B., Arnet E., Tverdal A., & Lingaas F. 2010. Differential gene expression in brain tissues of aggressive and non-aggressive dogs. BMC vet res. 6(1), 3. https://doi.org/10.1186/1746-6148-6-34

• Veroude K, Zhang-James Y, Fernandez-Castillo N, Bakker MJ, Cormand B, Faraone SV.

2016. Genetics of aggressive behavior: an overview. Am J Med Genet pat B Neuropsych

Genet. 1:3–43. https://doi.org/10.1002/ajmg.b.32364

• Vigers A.J., Amin D.S., Talley-Farnham T., Gorski J.A., Xu B. & Jones K.R. 2012. Sustained

expression of brain-derived neurotrophic factor is required for maintenance of dendritic spines

and normal behavior. *Neurosci*. 212:1–18. https://doi.org/10.1016/j.neuroscience.2004.04.031

• Wilkins A. S., Wrangham R. W., & Fitch W. T. 2014. The "domestication syndrome" in

mammals: a unified explanation based on neural crest cell behavior and

genetics. Genetics, 197(3), 795-808. https://doi.org/10.1534/genetics.114.165423

1. Wrangham, R. 2019. *The goodness paradox: the strange relationship between virtue and*

violence in human evolution. Vintage plasticity. *Endocrinology*, 147(4), 1664-1674.

https://doi.org/10.1210/en.2005-1432

• Zhang-James Y. et al. 2019. An integrated analysis of genes and functional pathways for

aggression in human and rodent models. Molecular Psychiatry. 24(11), 1655-1667.

https://doi.org/10.1038/s41380-018-0068-7

• Zhu C. Q. et al. 2008. Role of KRAS and EGFR as biomarkers of response to erlotinib in

National Cancer Institute of Canada Clinical Trials Group Study BR. 21. Journal of clinical

oncology. 26(26), 4268-4275. DOI: 10.1200/JCO.2007.14.8924