Preexisting memory CD4 T cells in naïve individuals confer robust immunity upon vaccination

Running title: Preexisting memory T cells confer immunity to vaccination

George Elias [1,2]%, Pieter Meysman [2,3,4]%, Esther Bartholomeus [2,5]%, Nicolas De Neuter [2,3,4], Nina Keersmaekers [2,6], Arvid Suls [2,5], Hilde Jansens [7], Aisha Souquette [8], Hans De Reu [1,9], Evelien Smits [1,9], Eva Lion[1,9], Paul G. Thomas [8], Geert Mortier [2,5], Pierre Van Damme [2,10], Philippe Beutels [2,6], Kris Laukens [2,3,4]%, Viggo Van Tendeloo [1,11]%, Benson Ogunjimi [2,6,12,13]%

% = equal contribution

[1]Laboratory of Experimental Hematology (LEH), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, Belgium.

[2]Antwerp Unit for Data Analysis and Computation in Immunology and Sequencing (AUDACIS), Antwerp, Belgium.

[3]Adrem Data Lab, Department of Mathematics and Computer Science, University of Antwerp, Antwerp, Belgium.

[4]Biomedical Informatics Research Network Antwerp (biomina), University of Antwerp, Antwerp, Belgium.

[5]Department of Medical Genetics, University of Antwerp/Antwerp University Hospital, Edegem, Belgium

[6]Centre for Health Economics Research & Modeling Infectious Diseases (CHERMID), Vaccine & Infectious Disease Institute (VAXINFECTIO), University of Antwerp, Antwerp, Belgium

[7]Department of Clinical Microbiology, Antwerp University Hospital

[8]Department of Immunology, St. Jude Children's Research Hospital, Memphis, Tennessee, USA

[9]Center for Cell Therapy and Regenerative Medicine, Antwerp University Hospital, Edegem, Belgium

[10]Centre for the Evaluation of Vaccination (CEV), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, Belgium.

[11]Janssen Research & Development, Immunosciences WWDA, Johnson & Johnson, Beerse, Belgium

[12]Antwerp Center for Translational Immunology and Virology (ACTIV), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, Belgium

[13]Department of Paediatrics, Antwerp University Hospital, Antwerp, Belgium.

28   Corresponding authors: George Elias, Benson Ogunjimi

29   **E-mail address:**

30   george.elias@student.uantwerpen.be, benson.ogunjimi@uantwerp.be

31   Laboratory of Experimental Hematology (LEH), Vaccine and Infectious Disease Institute,

32   University of Antwerp, Universiteitsplein 1, 2610 Antwerp, Belgium

Footnotes

This work was funded by

1. University of Antwerp [BOF Concerted Research Action (PS ID 30730) Methusalem

funding; Industrial Research Fund SBO].

2. Research Foundation Flanders (1861219N grant to B. Ogunjimi and FWO SB grant to N.

De Neuter.

3. ALSAC at St. Jude Children's Research Hospital and by HSN272201400006C and

R01AI107625 from the National Institute of Allergy and Infectious Disease.

45
46

Abstract

Antigen recognition through the T cell receptor (TCR) αβ heterodimer is one of the primary determinants of the adaptive immune response. Vaccines activate naïve T cells with high specificity to expand and differentiate into memory T cells that allow for a quick and robust T cell response upon exposure to the pathogen or re-exposure to the vaccine antigen. This is why the induction of memory T cells is a key feature in vaccine development. However, it has become increasingly evident that antigen-specific memory CD4 and CD8 T cells exist in unexposed antigen-naïve hosts and it is likely that exposure to one antigen might alter the TCR repertoire of memory T cells to a different unrelated antigen. In this study, we utilize high-throughput sequencing to profile the memory CD4 TCRβ repertoire and track vaccine-specific TCRβ clonotypes following the de novo administration of hepatitis B (HepB) vaccine in healthy HepB-naïve individuals and show that vaccinees with preexisting vaccine-specific memory CD4 T cell clonotypes elicited earlier and higher antibody concentrations and mounted a more robust CD4 T cell response to the vaccine. We further identify vaccine-specific TCRβ sequence patterns that can be used to predict which individuals will have an early and more vigorous vaccine-elicited immunity to HepB vaccine. Moreover, we find that an expansion of 4-1BB+ memory T$_{REG}$ is a prominent feature in individuals with delayed and modest vaccine-induced immunity. Our approach shows that modeling pre-vaccination TCRβ repertoire enables prediction of both antibody and CD4 responses to vaccines.

Keywords

CD4 T cell, T cell receptor, immune repertoire sequencing, vaccination

Introduction

Antigen recognition through the T cell receptor (TCR) is one of the key determinants of the adaptive immune response [1]. Antigen presentation via major histocompatibility complex (MHC) (encoded by HLA genes), together with the right costimulatory and cytokine signals, are responsible for T cell activation [2,3]. In this system, every T cell receptor (TCR) αβ heterodimer imparts specificity for a peptide-MHC (pMHC) complex. A highly diverse TCR repertoire ensures that an effective T cell response can be mounted against pathogen-derived peptides [4]. High TCRαβ diversity is generated through V(D)J recombination at the complementary-determining region 3 (CDR3) of TCRα and TCRβ chains, accompanied with junctional deletions and insertions of nucleotides, further adding to the diversity [5].

Vaccines activate naïve T cells with high specificity to vaccine-derived peptides and induce their expansion and differentiation into effective and multifunctional T cells. This is followed by a contraction phase from which surviving cells constitute a long-lived memory T cell pool that allows for a quick and robust T cell response upon a second exposure to the pathogen [6]. However, recent work has shown that a prior pathogen encounter is not a prerequisite for the formation of memory T cells and that CD4 T cells with a memory phenotype can be found in antigen-naïve individuals [7]. The existence of memory-like CD4 T cells in naïve individuals [8] can be explained by molecular mimicry, as the encounter with environmentally-derived peptides activates cross-reactive T cells due to the highly degenerate nature of the CD4 T cell recognition of peptide-MHC complex [9]. Indeed, work that attempted to replicate the history of human pathogen exposure in mice has shown that sequential infections altered the immunological profile and remodeled the immune response to vaccination [10]. The existence of memory CD4 T cells specific to vaccine-derived peptides in unexposed individuals might confer an advantage in

vaccine-induced immunity. In the present study we used high-throughput sequencing to profile

the memory CD4 TCRβ repertoire of healthy adults before and after administration of a hepatitis

B vaccine to investigate the impact of preexisting memory CD4 T cells on the immune response

to the vaccine. Based on anti-hepatitis B surface (anti-HBs) antibody titers over 365 days,

vaccinees were grouped into early, late and non-converters. Our data reveals that individuals

with preexisting vaccine-specific CD4 T cell clonotypes in the memory CD4 compartment had

earlier emergence of antibodies and mounted a more vigorous CD4 T cell response to the

vaccine. Moreover, we identify a set of vaccine-specific TCRβ sequence patterns which can be

used to predict which individuals will have an early and more vigorous response to hepatitis B

vaccine.

Methods

Human study design and clinical samples. A total of 34 healthy individuals (20-29y: 10, 30-39y: 7, 40-49y: 16, 50+y: 1) without a history of HBV infection or previous hepatitis B vaccination were recruited in this study after obtaining written informed consent. Individuals were vaccinated with a hepatitis B vaccine by intramuscular (m. deltoideus) injection (Engerix-B® containing 20 µg dose of alum-adjuvanted hepatitis B surface antigen, GlaxoSmithKline) on days 0 and 30 (and on day 365). At days 0 (pre-vaccination), 60, 180 and 365, peripheral blood samples were collected on spray-coated lithium heparin tubes, spray-coated K2EDTA (dipotassium ethylenediamine tetra-acetic acid) tubes and serum tubes (Becton Dickinson, NJ, USA). The study was approved by the Ethics Committee of Antwerp University Hospital and University of Antwerp (Antwerp, Belgium).

Peripheral blood mononuclear cells. Peripheral blood mononuclear cells (PBMC) were isolated by Ficoll-Paque Plus gradient separation (GE Healthcare, Chicago, IL, USA). Cells were stored in 10% dimethyl sulfoxide in fetal bovine serum (Thermo Fisher Scientific, Waltham, MA, USA). After thawing and washing cryopreserved PBMC, cells were cultured in AIM-V medium that contained L-glutamine, streptomycin sulfate at 50 µg/ml, and gentamicin sulfate at 10 µg/ml (Thermo Fisher Scientific, Waltham, MA, USA) and supplemented with 5% human serum (One Lambda, Canoga Park, CA, USA).

Serology and complete blood count. Serum was separated and stored immediately at – 80°C until time of analysis. Anti-HBs antibody was titrated in serum from day 0, 60, 180 and 365

using Roche Elecsys® Anti-HBs antibody assay on an Elecsys® 2010 analyzer (Roche, Basel,

Switzerland). An anti-HBs titer above 10 IU/ml was considered protective [11].

Serum IgG antibodies to Cytomegalovirus (CMV), Epstein–Barr virus viral-capsid antigen

(EBV-VCA), and Herpes Simplex virus (HSV)-1 and 2 were determined using commercially

available sandwich ELISA kits in accordance with the manufacturer's instructions.

A complete blood count including leukocyte differential was run on a hematology analyzer

(ABX MICROS 60, Horiba, Kyoto, Japan).

Sorting of memory CD4 T cells. Total CD4 T cells were isolated by positive selection using

CD4 magnetic microbeads (Miltenyi Biotech, Bergisch Gladbach, Germany). Memory CD4 T

cells were sorted after gating on single viable CD3+CD4+CD8–CD45RO+ cells. The following

fluorochrome-labeled monoclonal antibodies were used for staining: CD3-PerCP (BW264/56)

(Miltenyi Biotech), CD4-APC (RPA-T4) and CD45RO-PE (UCHT1) (both from Becton

Dickinson, Franklin Lakes, NJ, USA) and CD8-Pacific Orange (3B5) (from Thermo Fisher

Scientific, Waltham, MA, USA). Cells were stained at room temperature for 20 min and sorted

with FACSAria II (Becton Dickinson). Sytox blue (Thermo Fisher Scientific) was used to

exclude non-viable cells.

Single peptides, matrix peptide pools and epitope mapping. A set of 15-mers peptides with an

11-amino acid overlap spanning the 226 amino acids along the small S protein of hepatitis B

(HB) surface antigen (HBsAg), also designated as small HBs (SHBs) [12], were synthesized by

JPT Peptide Technologies (Berlin, Germany). The set, composed of 54 **single peptides** (See

supplementary table 1), was used in a matrix-based strategy to map epitopes against which the

immune response is directed [13]. The matrix layout enables efficient identification of epitopes

within the antigen using a minimal number of cells. For this purpose, a matrix of 15 pools, 7

rows and 8 columns, referred to as **matrix peptide pool**, was designed so that each peptide is in

exactly one row-pool and one column-pool, thereby allowing for the identification of positive

peptides at the intersection of positive pools. Matrix peptide pools that induced a CD4 T cell

response (as determined by CD40L/CD154 assay described below) which meets the threshold

criteria for a positive response were considered in the deconvolution process. Top six single

peptides were considered for peptide-specific T cell expansion and sorting. A **master peptide**

pool is composed of all of the 54 single peptides and was used to identify and sort total vaccine-

specific CD4 T cells. Each peptide was used at a final concentration of 2 µg/ml.

Ex vivo **T cell stimulation (CD40L/CD154 assay).** Thawed PBMC from each vaccinee were

cultured in AIM-V medium that contained L-glutamine, streptomycin sulphate at 50 µg/ml, and

gentamicin sulphate at 10 µg/ml. (GIBCO, Grand Island, NY) and supplemented with 5% human

serum (One Lambda, Canoga Park, CA, USA). Cells were stimulated for 6 hours with 2 µg/ml of

each of the 15 matrix peptide pools in the presence of 1 µg/ml anti-CD40 antibody (HB14)

(purchased from Miltenyi Biotec, Bergisch Gladbach, Germany) and 1 µg/ml anti-CD28

antibody (CD28.2) (purchased from BD Biosciences, Franklin Lakes, NJ, USA).

Cells were stained using the following fluorochrome-labelled monoclonal antibodies: CD3-

PerCP (BW264/56), CD4-APC (REA623), CD8-VioGreen (REA734) and CD40L-PE (5C8)

(purchased from Miltenyi Biotec, Bergisch Gladbach, Germany). Viability dye Sytox blue from

Invitrogen (Thermo Fisher Scientific, Waltham, MA, USA) was used to exclude non-viable cells.

Data was acquired on FACSAria II using Diva Software, both from BD Biosciences (Franklin

Lakes, NJ, USA), and analyzed on FlowJo software version 10.5.3 (Tree Star, Inc., Ashland, OR,

USA). Fluorescence-minus-one controls were performed in pilot studies. Gates for CD40L+CD4

T cells were set using cells left unstimulated.

***In vitro* T cell expansion and cell sorting.** Thawed PBMC were labelled with

carboxyfluorescein succinimidyl ester (CFSE) (Invitrogen, Carlsbad, CA, USA) and cultured in

AIM-V medium that contained L-glutamine, streptomycin sulphate at 50 µg/ml, and gentamicin

sulphate at 10 µg/ml. (GIBCO, Grand Island, NY) and supplemented with 5% human serum

(One Lambda, Canoga Park, CA, USA). Cells were stimulated for 7 days with 2 µg/ml of

selected single peptides in addition to the master peptides pool. Cells were stained using the

following fluorochrome-labelled monoclonal antibodies: CD3-PerCP (BW264/56), CD4-APC

(REA623) and CD8-VioGreen (REA734) (purchased from Miltenyi Biotec, Bergisch Gladbach,

Germany). Viability dye Sytox blue from Invitrogen (Thermo Fisher Scientific, Waltham, MA,

USA) was used to exclude non-viable cells. Single viable CFSE$_{low}$ CD3+ CD8– CD4+ T cells

were sorted into 96-well PCR plates containing DNA/RNA Shield (Zymo Research, Irvine, CA,

USA) using FACSAria II and Diva Software (BD Biosciences, Franklin Lakes, NJ, USA). For

each of the selected single peptides, 500 cells were sorted in two technical replicates. For the

master peptide pool, 1000 cells were sorted in two technical replicates. Plates were immediately

centrifuged and kept at – 20°C before TCR cDNA library preparation and sequencing.

TCRβ cDNA Library Preparation and Sequencing of memory CD4 T cells. DNA was

extracted from sorted memory CD4 T cells using Quick-DNA Microprep kit (Zymo Research,

Irvine, CA, USA). ImmunoSEQ hsTCRB sequencing kit (Adaptive Biotechnologies, Seattle,

WA, USA) was used to profile TCRβrepertoire following the manufacturer's protocol.

After quality control using Fragment Analyzer (Agilent, Santa Clara, CA, USA), libraries were

pooled with equal volumes. The concentration of the final pool was measured with the Qubit™

dsDNA HS Assay kit (Thermo Fisher Scientific, Waltham, MA, USA). The final pool was

processed to be sequenced on the Miseq and NextSeq platforms (Illumina, San Diego, CA,

USA).

TCR cDNA Library Preparation and Sequencing of CFSElow CD4 T cells. RNA was

extracted from each of the two technical replicates of sorted CFSElow CD4 T cells using Quick-

RNA Microprep kit (Zymo Research, Irvine, CA, USA). Without measuring the resulted RNA

concentration, an RNA-based library preparation was used. The QIAseq Immune Repertoire

RNA Library kit (Qiagen, Venlo, Netherlands) amplifies TCR alpha, beta, gamma and delta

chains. After quality control using Fragment Analyzer (Agilent, Santa Clara, CA, USA),

concentration was measured with the Qubit™ dsDNA HS Assay kit (Thermo Fisher Scientific,

Waltham, MA, USA) and pools were equimolarly pooled and prepared for sequencing on the

Nextseq platform (Illumina, San Diego, CA, USA).

TCRβ Sequence Analysis. TCRβ clonotypes were identified as previously described [14] where a

unique TCRβ clonotype is defined as a unique combination of a V gene, CDR3 amino acid

sequence, and J gene. All memory CD4 T cell DNA-based TCRβ sequencing reads were

annotated using the immunoSEQ analyzer (v2) from Adaptive Biotechnologies. All small bulk

RNA-based TCR sequencing reads were annotated using the MiXCR tool (v3.0.7) from the

FASTQ files. As all RNA-based TCR sequencing experiments featured two technical replicates,

only those TCR sequences that occurred in both replicates were retained and their counts were

summed. Tracking of vaccine-specific TCRβ clonotypes is based on exact TCRβ CDR3 amino

acid matches to remove any bias introduced by the different VDJ annotation pipelines. Non-

HBsAg TCR annotations were done with the TCRex web tool [15] on the 24th of July, 2019 using

version 0.3.0. Inference of similar epitope binding between two TCR sequences is defined

according to the Hamming distance (d) calculated on the CDR3 amino acid sequence with a

cutoff c, as supported by Meysman et al. [16]. All scripts used in this analysis are available via

github (https://github.com/pmeysman/HepBTCR).

Predictive HBs-response model. From the single peptide data generated in the matrix peptide

pool experiments, we aimed to create a predictive model to enumerate the HBs response from

full TCRβ repertoire data. This approach allows for predictions that are epitope-specific rather

than simply vaccine-specific. This model was applied in a leave-one-out cross validation so that

vaccine-specific TCRβ sequences from a vaccinee are not used to make predictions for the same

vaccinee. While the predictive model is derived from epitope-specific data, it cannot be

guaranteed that some of the expanded CD4 T cells detected in the in vitro assay are not due to

bystander activation. Vaccine-specific TCRβ sequences of vaccinees who did not respond to the

vaccine at day 60 (late-converters and non-converters) are expected to be more enriched in cells

triggered to expand due to bystander activation. Indeed, running the set of vaccine-specific

TCRβ sequences through the TCRex webtool [15] reveals that there is strong enrichment for the

CMV NLVPMVATV epitope (P value = 1.16e-71) and the Mart-1 variant ELAGIGILTV

epitope (P value = 2.16e-60), which supports the notion that some of these TCRβ sequences

might not be specific to HBsAg. This set of vaccine-specific TCRβ sequences can thus be used

to make predictions about possible TCRβ sequences due to bystander activation of CD4 T cells,

i.e. common TCRβ sequences that might be present as false positives. The final output of the

model is thus a ratio R_{hbs} for any repertoire rep_i describing a set of TCRβ sequences t_{repi}:

256
$$R_{hbs}(t_{repi}) = \frac{\sum_{pep=1}^{54}|\{t_{repi}|d(t_{repi}, t_{pep}) < c\}|/|t_{pep}|}{|\{t_{repi}|d(t_{repi}, t_{bystander}) < c\}|/|t_{bystander}|}$$

with t_{pep} as the set of TCRβ sequences occurring in both biological replicates for a single sample

and a single peptide (*pep*) from the HBsAg matrix peptide pool experiment, and $t_{bystander}$ as the

set of TCRβ sequences occurring in both biological replicates of the master peptide pool in any

of the non-responding samples. Thus the ratio signifies the number of TCR clonotypes predicted

to be reactive against one of the HBsAg peptides, normalized by a count of putative false

positive predictions from bystander T-cells.

***Ex vivo* T cell phenotyping of vaccine-specific T cells.** Thawed PBMC from each vaccinee

were cultured in AIM-V medium that contained L-glutamine, streptomycin sulphate at 50 µg/ml,

and gentamicin sulphate at 10 µg/ml. (GIBCO, Grand Island, NY) and supplemented with 5%

human serum (One Lambda, Canoga Park, CA, USA). Cells were stimulated for 6 hours with 2

µg/ml of a master peptide pool representing the full length of the small surface envelope protein

of hepatitis B, in the presence of 1 µg/ml anti-CD40 antibody (HB14) (purchased from Miltenyi

Biotec, Bergisch Gladbach, Germany) and 1 µg/ml anti-CD28 antibody (CD28.2) (purchased

from BD Biosciences, Franklin Lakes, NJ, USA). Cells were stained using the following

fluorochrome-labelled monoclonal antibodies:

CD3-BV510 (SK7), CD4-PerCP/Cy5.5 (RPA-T4), CD8-APC/Cy7 (SK1), CD45RA-AF488

(HI100), CD25-BV421 (M-A251), CD127-BV785 (A019D5) and CD137-PE (4-1BB)

(purchased from BioLegend, San Diego, CA, USA), CXCR5 (CD185)-PE-eFluor 610

(MU5UBEE) (from eBioscience, Thermo Fisher Scientific, Waltham, MA, USA) and CD40L-

APC (5C8)   (purchased from Miltenyi Biotec, Bergisch Gladbach, Germany). Fixable viability

dye Zombie NIR™ from BioLegend (San Diego, CA, USA) was used to exclude non-viable

cells. Data was acquired on FACSAria II using Diva Software, both from BD Biosciences

(Franklin Lakes, NJ, USA), and analyzed on FlowJo software version 10.5.3 (Tree Star, Inc.,

Ashland, OR, USA) using gating strategy shown in **Fig. S1a**. Fluorescence-minus-one controls

were performed in pilot studies. Gates for CD40L+ and 4-1BB+ CD4 T cells **(Fig. S1b)** were set

using cells left unstimulated (negative control contained DMSO at the same concentration used

to solve peptide pools). In order to account for background expression of CD40L and 4-1BB on

CD4 T cells, responses in cells left unstimulated were subtracted from the responses to peptides,

and when peptides-specific CD40L+ or 4-1BB+ CD4 T cells were not significantly higher than

those detected for cells left unstimulated (using one-sided Fisher's exact test), values were

mutated to zero.

Statistical analysis and data visualization. The two-sided Fisher's exact test was used to

evaluate the significance of relationship between early/late-converters and CMV, EBV or HSV

seropositivity. For the visualization of marker expression, TCRβ counts and cell frequencies

between time points or groups of vaccinees, ggplot2 (V3.3.2) and ggpubr (V0.2.5) packages in R

were used. The Wilcoxon signed-rank test was used to compare two or more groups, with

unpaired and paired analysis as necessary. The nonparametric Spearman's rank-order correlation

was used to test for correlation. We used the following convention for symbols indicating

statistical significance; ns $P > 0.05$, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$, **** $P \leq 0.0001$.

Statistical analysis. The two-sided Fisher's exact test was used to evaluate the significance of

relationship between early/late-converters and CMV, EBV or HSV seropositivity. For the

visualization of marker expression, TCRβ counts and cell frequencies between time points or

groups of vaccinees, ggpubr (V0.2.5) package in R was used. The Wilcoxon signed-rank test was

used to compare two or more groups, with unpaired and paired analysis as necessary. The

nonparametric Spearman's rank-order correlation was used to test for correlation. We used the

following convention for symbols indicating statistical significance; ns $P > 0.05$, * $P \leq 0.05$, ** P

≤ 0.01, *** $P \leq 0.001$, **** $P \leq 0.0001$.

Data availability

The sequencing data that support the findings of this study have been deposited on Zenodo

(https://doi.org/10.5281/zenodo.3989144). Flow Cytometry Standard (FCS) data files with

associated FlowJo workspaces are deposited at flowrepository.org [17] under the following

experiment names: epitope mapping: https://flowrepository.org/id/FR-FCM-Z2TN; *in vitro* T

cell expansion: https://flowrepository.org/id/FR-FCM-Z2TM; *ex vivo* CD4 T cell assay:

https://flowrepository.org/id/FR-FCM-Z2TL.

Results:

Vaccinee cohort can be classified into three groups

Out of 34 vaccinees, 21 vaccinees seroconverted (an anti-HBs titer above 10 IU/ml was considered protective [11]) at day 60 and were classified as early-converters; 9 vaccinees seroconverted at day 180 or day 365 and were classified as late-converters; remaining 4 vaccinees had an anti-HBs antibody titer lower than 10 IU/ml at all time points following vaccination and were classified as non-converters **(Fig. 1 and Fig. S2a)**.

Members of *Herpesviridae* family might alter immune responses to vaccines [18]. We found no significant differences in CMV, EBV or HSV seropositivity between the three groups in our cohort (**Fig. S2b**). Early-converters were slightly younger than late-converters and non-converters were notably younger than both early and late converters (**Fig. S2c**).

Figure 1. Hepatitis B vaccination (Engerix-B®) study design.

a Hepatitis B (Engerix-B®) vaccination and experimental design. (Top) Timeline of vaccination
and blood collection. (Bottom) Memory CD4 T cells were magnetically enriched and FACS-
sorted from two time points (day 0 and day 60) for TCRβ repertoire sequencing. Matrix peptide
pools were used to map CD4 T cell epitopes of the vaccine from PBMCs collected at day 60 and
to select single peptides. After 7 days of *in vitro* expansion, single peptide-specific and master
peptide pool-specific CFSE_low CD4 T cells from PBMCs collected at day 60 were FACS-sorted
in two technical replicates for TCRβ repertoire sequencing. PBMCs collected at days 0, 60, 180,
and 365 were stimulated with the master peptide pool (HBsAg) and assessed for converse
expression of 4-1BB and CD40L by flow cytometry.
b Vaccinee cohort can be classified into three groups as determined by anti-Hepatitis B surface
(anti-HBs) titer over four times points.
Early-converters seroconverted at day 60, late-converters seroconverted at day 180 or day 365
and non–converters did not have an anti-HBs titer higher than 10 IU/ml at any of the time points.

Memory CD4 T cell repertoire decreases in clonality following vaccination in early-converters

A genomic DNA-based TCRβ sequence dataset of memory CD4 T cells isolated from peripheral blood was generated from a cohort of 34 healthy vaccinees right before vaccination (day 0) and 60 days after administration of the first dose of hepatitis B vaccine (30 days after administration of the second vaccine dose).

Between 7.32×10^4 and 4.41×10^5 productive TCRβ sequence reads were obtained for each vaccinee at each time point **(Fig. S3a)**. Between 30,000 and 90,000 unique TCRβ sequences were sequenced for each vaccinee at each time point **(Fig. S3b)**. As expected, considering the extremely diverse memory CD4 T cell repertoire [19], only an average of 25% of the TCRβ sequences can be detected at both time points for each vaccinee **(Fig. S3c)**.

The diversity of the memory CD4 T cell repertoire of each vaccinee at the two time points was explored. Even though the number of unique clones in the memory CD4 TCRβ repertoire remained stable in between the two time points, we detected a significant increase in the TCRβ repertoire Shannon's entropy for early-converters **(Fig. 2a)** (P value = 0.042), but not late-converters, suggesting that the memory CD4 T cell repertoires of early-converters have become less clonal, despite the number of distinct TCRβ sequences not changing significantly.

Unique vaccine-specific TCRβ sequences are trackable within memory CD4 T cell repertoire and increase following vaccination

Peripheral blood mononuclear cells from day 60 were labeled with carboxyfluorescein succinimidyl ester (CFSE) and stimulated with a pool of peptides spanning hepatitis B (HB)

surface antigen (HBsAg). At day 7 of *in vitro* expansion, we sorted CFSElow CD4 T cells [20] and

extracted mRNA for quantitative assessment of vaccine-specific TCRβ clonotypes by sequencing

(see *Methods* for details), allowing for the tracking of vaccine-specific TCRβ within memory

CD4 T cell repertoire over the two times points, based on CDR3β amino acid sequence mapping.

We detected a significant increase in the frequency of unique vaccine-specific TCRβ sequences

at day 60 post-vaccination compared to pre-vaccination (mean increase = 96.5%, 95% CI = 56.7

- 170%) **(Fig. 2b)**. Moreover, this increase was larger for early-converters (mean = 132.1%, 95%

CI = 76.4 - 238.2%) than late-converters (mean = 22.1%, 95% CI = 5.9 – 50.1%). For non-

converters the mean was 81.6% (95% CI: [42.7% - 110.6%]). A Wilcoxon test shows that the

difference between the increase for the early converters and late converters had a P value of

0.04909.

As vaccine-specific TCRβ sequences were already detected in the memory CD4 T cell repertoire

prior to vaccination, we sought to determine whether the vaccination results in an expansion of

those sequences. However, using the abundance of vaccine-specific TCRβ sequences within the

memory CD4 T cell repertoire, the data does not support a vaccine-induced expansion of

preexisting vaccine-specific TCRβ sequences **(Fig. S3d)**. Thus, although we see a rise in the

number of vaccine-specific TCRβ clonotypes from day 0 to day 60, this cannot be attributed to

an expansion of preexisting TCRβ clonotypes but rather the recruitment of new TCRβ

clonotypes (presumably from the naïve T cell compartment), as visualized for one vaccinee in

Fig. 2c.

It makes sense to not only look at the difference in vaccine-specific TCRβ sequences between

time points, but also explore whether there are differences in the proportion of HBsAg-specific

clones in the memory repertoire between early-converters, late-converters and non-converters

after vaccination. In this case, as we aim for a between-vaccinees comparison (in contrast to the

within-vaccinees timepoint comparison), we normalize by the number of HBsAg-specific TCRβ

found for each vaccinee. Thus, the values are different from those reported before. From this

analysis, it can be concluded that there is a difference in HBsAg-specific TCRβ at day 60

between the three groups **(Fig. 2d)** (ANOVA P value = 0.00238). A Wilcoxon test between

early-converters and other vaccinees shows a significant P value of 0.000473, thereby implying

that early-converters have a higher relative frequency of vaccine-specific TCRβ sequences

present in their memory repertoire at day 60 compared to the rest.

It is notable that some of the vaccine-specific T cell clonotypes can be detected in the memory

repertoire of vaccinees prior to vaccination **(Fig. 2e)**. However, the relative frequency was not

statistically significant different between the categories (ANOVA P value = 0.238).

Figure 2. CD4 T cell memory TCRβ repertoire and vaccine-specific TCRβ clonotypes.

a Repertoire diversity and entropy between the two time points

b Frequency of unique vaccine-specific TCRβ sequences out of total sequenced TCRβ sequences between two time points for all vaccinees colored by group

c Sequenced CD4+ TCR memory repertoire of vaccinee H35 at day 60. Each TCR clonotype is represented by a node. TCRs are connected by an edge if their Hamming distance is one. Only clusters with at least three TCRs are shown. TCR clonotypes in red are the vaccine-specific TCRβ sequences that were not present prior to vaccination.

d Frequency of vaccine-specific TCRβ sequences within memory CD4 T cell repertoire normalized by amount of HBsAg-specific TCRβ found for each vaccinee at time point 60 and (**e**) at time point 0.

HBsAg single peptide-specific TCRβ identification allows predictive modelling of early converters prior to vaccination

To quantify the T-cell response at the level of individual peptides that make up the HBsAg, a matrix peptide pool covering 54 overlapping peptides of the HBsAg was used to extract peptide-specific T-cells using a CD40L/CD154 activation-induced marker (AIM) assay (see *Methods* for details, **Fig. S4**). The top 6 peptides for each individual were selected for TCR sequencing after a CFSE assay **(Supplementary Table 2)**. In this manner, TCRβ sequences were identified for T-cells reactive against 44 individual HBsAg peptides. These were not uniformly distributed across the HBsAg amino acid sequence, with the most prominent epitopes covering the regions 1-15, 129-144, 149-164, 161-176, 181-200, 213-228. For each of those regions, more than 10 individuals had a strong T-cell response and more than 150 unique TCRβ sequences could be identified **(Fig. 3a)**.

These peptide-specific TCRβ sequences can be utilized in a peptide-TCR interaction classifier to identify other TCRβ that are likely to react against the same HBsAg epitopes, as it has been shown that similar TCRβ sequences tend to target the same epitopes [16,21]. These classifications were integrated into a model which outputs a ratio R_{hbs} for any TCRβ repertoire representing the amount of HBsAg peptide-specific clonotypes. The ratio R_{hbs} is based on the frequency of putative peptide-specific TCRβ divided by a normalization term for putative false positive predictions due to bystander activations in the training data set. This model applied to the memory repertoire at day 60 shows that early-converters tend to have a higher frequency of putative HBsAg peptide-specific TCRβ, while late-converters tend to have relatively more false positive hits **(Fig. 3b)**. Thus, the defined ratio R_{hbs} shows significant difference between the early-converters and the late-converters at day 60 (one-sided Wilcoxon-test P value= 0.0313,

Fig. 3c). Furthermore, calculating the R_{hbs} on the memory repertoires prior to vaccination (day 0)

shows a similar difference (one-sided Wilcoxon-test P value= 0.0010, **Fig. 3d**). In this manner,

R_{hbs} has predictive potential and can be used as a classifier to distinguish early from late-

converters prior to vaccination **(Fig. 3e)**, with an AUC of 0.825 (95% CI: 0.657 – 0.994) in a

leave-one-out cross validation setting.

While R_{hbs} is able to differentiate between early-and late-converters, it seems to be worse at

distinguishing non-converters. This is mainly due to a single non-converter vaccinee (H21) with

a high R_{hbs}, signifying a high number of putative HBsAg peptide-specific TCRβ in their memory

repertoire.

Predictive capacity of TCRβ repertoire holds true for humoral and cellular immune

response

Response groups used thus far were established based on anti-HBs titers at different time points.

However, response groups can be defined differently if one considers the results from the

CD40L assay. For the most part, only those with a high Ab titer at day 60, had a positive value

for the CD40L assay **(Fig S3e)**. The only exception was H21, who had a low measurable Ab titer

but had a measurable CD40L value. Thus, this vaccinee seems to have had a T-cell response

against the vaccine, without developing an antibody response. It is interesting to note that this

vaccinee is classified as a non-responder in the data set but it is predicted as a possible strong

early-converter by the R_{hbs} predictive model in the leave-one-out cross validation. Redoing the

analysis with R_{hbs} ratio to predict the CD40L assay at day 60, shows that the R_{hbs} ratio is a good

classifier in a leave-one-out cross validation **(Fig. 3f)**. We tested the predictive capacity of R_{hbs}

ratio at different time points and cell types, and found good AUC values for both 4-

1BB–CD40L+ (T$_{CON}$) and 4-1BB+CD40L– (T$_{REG}$, see further) at day 60, but to a lesser extent at

days 180 and 365, thereby indicating that R_{hbs} predictive model is currently limited to predicting

mid-term responses following vaccination **(Fig S3f and g)**.

Figure 3

a. Overview of found HBsAg epitope-specific TCRβ sequences. Each bar corresponds to unique
TCRβ sequences found against a single 15mer HBsAg peptide, with 11 amino acid overlap to
each subsequent peptide. Bars in blue denote those epitopes for which 10 or more volunteers had
a strong T-cell reaction. Motif logos on top of bars denote a sampling of the most common
TCRβ amino acid sequence motifs for those epitopes.
b. Scatter plot with the percentage predicted epitope-specific and bystander TCRβ sequences.
Predictions done as a leave-one-out cross-validation. Each dot represents a vaccinee with the
color denoting the responder status (blue: early-converter, yellow: late-converter, red: non-
converter)
c. HBsAg-predictive ratio R(hbs) when calculated on the memory repertoires at day 60
d. HBsAg-predictive ratio R(hbs) when calculated on the memory repertoires at day 0
e. ROC curve of using epitope-specific / bystander ratio to differentiate between early-converters
and late-converters in a leave-one-out cross validation at day 0. Diagonal line denotes a random
classifier. Reported is the area under the curve (AUC) and its 95% confidence interval
f. ROC curve of using epitope-specific / bystander ratio to differentiate between samples with a
positive CD4L value (> 0) in a leave-one-out cross validation. Diagonal line denotes a random
classifier. Reported is the area under the curve (AUC) and its 95% confidence interval. Note that
the classifier is worse as several samples (EC, LC and NC) had NA values for the CD154 assay,
which were all set to zero.
499
500

Vaccine-specific conventional and regulatory memory CD4 T cells induced in early-converters

After showing evidence for the existence of vaccine-specific TCRβ sequences pre-vaccination and that individuals with a higher number of HBsAg peptide-specific clonotypes had earlier seroconversion, we attempted to link this observation to differences in vaccine-specific CD4 T cells responses using CD4 T cell assays. As T_REG cells might suppress vaccine-induced immune responses [22], we used activation markers CD40L (CD154) and 4-1BB (CD137) to help delineate the conventional (T_CON) and regulatory (T_REG) phenotypes of activated CD4 T cells [23,24]. In this scheme, after 6 hours of antigen stimulation, CD40L+4-1BB− can be used as a signature for antigen-specific CD4 T_CON cells, as opposed to CD40L−4-1BB+ signature for antigen-specific CD4 T_REG cells.

Additionally, we added CD25 and CD127 to better identify T_REG cells [25,26] and CXCR5 to further distinguish circulatory T follicular helper cells (cT_FH) and circulatory T follicular regulatory cells (cT_FR) [27,28].

Using the converse expression of CD40L and 4-1BB, CD40L+4-1BB− and CD40L−4-1BB+ CD4 T cells had a T_CON and T_REG phenotype, respectively, as shown by the expression of CD25 and CD127 **(Fig. S5a and b)**, and validate their use for the distinction of activated T_CON and T_REG cells.

We detected a significant increase in the frequency of CD40L+4-1BB− and CD40L−4-1BB+ memory CD4 T cells at day 60 in our cohort **(Fig. 4a)** that correlated positively with the increase in antibody titer between day 0 and day 365 **(Fig. 4b and Fig. S6)**. Upon a closer look, the induction of both signatures of vaccine-specific memory CD4 T cells was only true for early-converters (**Fig. 4c**, see **Fig. S7a** for non-converters and **Fig. S7b** for vaccine-specific CD4 T

cells) while late-converters did not show a detectable memory CD4 T cell response. Although a

subset of both early and late-converters had detectable memory CD4 T cell responses prior to

vaccination, we detected no significant difference in the frequency of $CD40L_+4\text{-}1BB_-$ and

$CD40L_-4\text{-}1BB_+$ memory CD4 T cells between the two groups at day 0 **(Fig. 4d)**.

Collectively, flow cytometry data reveal that the expression of CD40L and 4-1BB in our ex vivo

assay is consistent with our serological data and reflects the lack of seroconversion at day 60 in

late-converters. However, it does not support the existence of more vaccine-specific memory

CD4 T cells in early-converters prior to vaccination.

Figure 4 Hepatitis B vaccine induces a vaccine-specific CD40L+4-1BB− and CD40L−4-1BB+ memory CD4 T cell response in early-converter vaccinees.

PBMCs from vaccinees were stimulated with 2 µg/ml of the master peptide pool (HBsAg) and assessed for converse expression of 4-1BB and CD40L by flow cytometry on days 0, 60, 180, and 365. Shown is number of of vaccine-specific memory CD4 T cells out of 1×10^6 memory CD4 T cells after subtraction of responses in negative control.
a Aggregate analysis from vaccinees (including early, late and non-converters) showing a peak of vaccine-specific CD40L+4-1BB− and CD40L−4-1BB+ memory CD4 T cell at day 60 (day 60 after 1_{st} dose of the vaccine and day 30 after 2_{nd} dose), declining thereafter. Shown are proportions of vaccine-specific memory CD4 T cells out of total memory CD4 t cells.
b Correlation between the difference in antibody titer between day 365 and day 0 and vaccine-specific CD40L+4-1BB− and CD40L−4-1BB+ memory CD4 T cell at day 60
c Aggregate analysis from early and late-converter vaccinees showing a significant induction of vaccine-specific CD40L+4-1BB− and CD40L−4-1BB+ memory CD4 T cell in early-converters and lack thereof in late-converters.
d Aggregate analysis from early and late-converter vaccinees showing no significant differences in vaccine-specific CD40L+4-1BB− and CD40L−4-1BB+ memory CD4 T cell at day 0.
Wilcoxon signed-rank with unpaired and paired analysis as necessary; statistical significance was indicated with ns P > 0.05, * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, **** P ≤ 0.0001
rs, Spearman correlation coefficient, −1 ≤ *r*s ≤ 1; *rs* and *p* value by Spearman's correlation test

An expanded subset of 4-1BB+CD45RA– T_{REG} cells is a prominent feature of late-converters

In order to detect any distinct signatures of early and late-converters, we analyzed pre-vaccination flow cytometry data to examine major CD4 T cell subsets; T_H, T_{REG}, cT_{FH} and cT_{FR} cells. Using manual gating in which regulatory T cells (T_{REG}) were defined as viable CD3+CD4+CD8–CD25+CD127–CXCR5– and were further divided into CD45RA+ and CD45RA– T_{REG} cells, we identified a significantly higher frequency of 4-1BB+ CD45RA– T_{REG} cells in late-converters compared to early-converters **(Fig. 5a and Fig. S8)**.

T_{REG} cells showed higher 4-1BB expression compared to T_H, cT_{FH} and cT_{FR} cells **(Fig. 5b)** and within T_{REG} subset, CD45RA– T_{REG} cells showed significantly higher expression of 4-1BB, accompanied with a higher expression of CD25, compared to CD45RA+ T_{REG} cells **(Fig. 5c)**. In this scheme, CD45RA– T_{REG} can be divided into 4-1BB+CD25$_{high}$ and 4-1BB-CD25$_{int}$ subsets. It is worth noting here that no differences were detected in the frequency of CD45RA– or CD45RA+ T_{REG} cells within CD4 T cell compartment between the two groups **(Fig. 5d)**, and that the composition of T_{REG} compartment that is distinct between the two groups **(Fig. 5e)**.

In summary, an expanded subset of 4-1BB+CD45RA– T_{REG} cells pre-vaccination is a prominent feature of a delayed and modest immune response to hepatitis B vaccine in our cohort.

Figure 5 An expanded 4-1BB+CD45RA– T_{REG} cells within T_{REG} compartment is is a prominent feature in late-converters prior to vaccination.

PBMCs from vaccinees at day 0 (prior to vaccination) were phenotyped for expression of markers of T_{REG}.
a Aggregate analysis of 4-1BB+CD45RA– T_{REG} within CD45RA– T_{REG} CD4 T cells in early and late and non-converter vaccinees before vaccination
b Aggregate analysis of the median fluorescence intensity of 4-1BB in T_H, cT_{FH}, T_{REG} and cT_{FR} cells before vaccination.
c Aggregate analysis of the median fluorescence intensity of 4-1BB (left panel) and CD25 (right panel) in CD45RA– T_{REG} and CD45RA+ T_{REG} cells before vaccination.
d Frequency of T_{REG}, CD45RA– T_{REG} and CD45RA+ T_{REG} cells within total CD4 T cells in early, late and non-converter vaccinees before vaccination.
e Composition of T_{REG} compartment as determined by expression of 4-1BB and CD45RA in early, late and non-converter vaccinees before vaccination.
Wilcoxon signed-rank with unpaired and paired analysis as necessary; statistical significance was indicated with ns P > 0.05, * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, **** P ≤ 0.0001

Discussion

In this study, we used high-throughput TCRβ repertoire profiling and *ex vivo* T cell assays to characterize memory CD4 T cell repertoires before and after immunization with hepatitis B vaccine, an adjuvanted subunit vaccine, and tracked vaccine-specific TCRβ clonotypes over two time points. As antigen-naïve adults were found to have an unexpected abundance of memory-phenotype CD4 T cells specific to viral antigens [7,29], we sought to investigate the influence that preexisting memory CD4 T cells can have on vaccine-induced immunity.

Commercially available HBV vaccines produces a robust and long-lasting anti-HBs response, and protection is provided by induction of an anti-HBs (antibody against HBV surface antigen) titer higher than 10 mIU/mL after a complete immunization schedule of 3 doses [30]. However, 5-10% of healthy adult vaccinees fail to produce protective titers of anti-HBs and can be classified as non-responders [30]. In our cohort, 13 vaccinees did not seroconvert by day 60 (30 days following administration of the second vaccine dose), as determined by antibody titer. Out of this group, 9 vaccinees seroconverted by day 180 or day 365, referred to here as late-converters, and 4 vaccinees did not seroconvert, referred to here as non-converters.

A hallmark of adaptive immunity is a potential for memory immune responses to increase in both magnitude and quality upon repeated exposure to the antigen [31]. Our systems immunology data supports the theory that preexisting memory CD4 T cell TCRβ sequences specific to HBsAg, the antigenic component of the current hepatitis B vaccine, predict which individuals will mount an early and more vigorous immune response to the vaccine as evidenced by a higher fold change in anti-HBs antibody titer and a more significant induction of antigen-specific CD4 T cells. It is postulated that preexisting memory CD4 T cell clonotypes are generated due to the highly degenerate nature of T cell recognition of antigen/MHC and are cross-reactive to

environmental antigens [8]. For example, preexisting memory CD4 T cells are well-established in

unexposed HIV-seronegative individuals, although at a significantly lower magnitude than HIV-

exposed seronegative individuals [32,33], and were likely primed by exposure to environmental

triggers or the human microbiome.

We and others have shown before that the TCRβ repertoire of CD4 T cells encodes the antigen

exposure history of each individual and that antigen-specific TCRβ sequences could serve to

automatically annotate the infection or exposure history [14,34,35]. In this study, we show that

similar principles can be used to study vaccine responsiveness. Specifically, the recruitment of

novel vaccine-specific T-cell clonotypes into memory compartment following vaccination can be

tracked by examining the CD4 memory TCRβ repertoire over time. While we observed no

increase in the frequency of the vaccine-specific memory T-cells, as the time point may have

missed the peak of the clonal expansion of effector CD4 T cells as was reported before [36–38], a

significant rise in the number of unique vaccine-specific T-cell clonotypes was detected.

Nevertheless, this observation is consistent with earlier T cell immune repertoire sequencing

studies that showed that antigen-specific TCRβ sequences do not always overlap with those

TCRβ sequences that increase in frequency after infection or vaccination [39]. More interestingly,

individuals with the earlier and more robust response against the vaccine, had a telltale antigen-

specific signature in their memory TCRβ repertoire prior to vaccination, despite the lack of

HBsAg antibodies or prior vaccination history.

Detection of this vaccine-specific signature was possible due to the development of a novel

predictive model that used epitope-specific TCRβ sequences from one set of individuals to make

predictions about another. A correction factor was needed to account for the occurrence of

bystander activated T-cells within the original epitope-specific TCRβ sequences. Indeed, in those

vaccinees without a positive antibody titer at day 60, any putative vaccine-specific T-cells may

be considered as bystander vaccination. This was supported by predictions using the TCRex tool

[15], which matched these TCR sequences to common viral or other epitopes. It is of note that

these TCRβ sequences are matched with CD8 T cell epitopes, while they originate from isolated

CD4 T cells. This is likely due to the great similarity between the TCRβ sequences of CD4 and

CD8 T cells as noted in prior research [16].

However, our in vitro antigen-specificity data, using an assay that enables discrimination of

T_{CON} and T_{REG} cells using the converse expression of the activation markers CD40L and 4-1BB

[23,40], failed to show a significant difference in preexisting antigen-specific CD4 T cells between

early and late-converters prior to vaccine administration. It is plausible that the signal is below

the detection limit of the assay and that more sensitive assays that require pre-enrichment of

$CD40L_+$ and $4-1BB_+$ T cells (using magnetic beads) [41] or cultured ELISpot assay [42] are needed to

capture preexisting vaccine-specific memory CD4 T cells directly from human peripheral blood.

Another plausible explanation is that our activation proteins, CD40L and 4-1BB, might be

unsuitable to detect preexisting memory CD4 T cells but this is unlikely as both proteins have

been used successfully in similar studies [43,44]. A different explanation may be that the diversity of

preexisting antigen-specific CD4 T clonotypes as determined by TCRβ sequencing is not

reflected in the quantitative measurements of the fractional cell counts.

TREG cells represent about 5 – 10% of human CD4 T cell compartment and are identified by the

constitutive surface expression of CD25, also known as IL-2 receptor α subunit (IL-2Ra), and the

nuclear expression of forkhead family transcription factor 3 (Foxp3), a lineage specification

factor of TREG cells [45]. Regulatory memory T cells play a role in the mitigation of tissue damage

induced by effector memory T cells during protective immune responses, resulting in a selective

advantage against pathogen-induced immunopathology [46–49]. Several studies have identified CD4

TREG cells with specificity to pathogen-derived peptides in murine models and showed evidence

for an induced expansion of TREG cells followed by an emergence and a long-term persistence of

TREG cells with a memory phenotype and potent immunosuppressive properties [47,50]. Blom et al.

reported a significant and transient activation of TREG cells (identified by upregulation of CD38

and Ki67) in humans 10 days after administration of live attenuated yellow fever virus 17D

vaccine [37]. The induction of vaccine-specific TREG cells in our cohort is unexpected and the role

it might play in vaccine-induced immunity warrants further investigation.

The association of an expanded 4-1BB+ CD45RA− TREG subset with a delayed immune response

to hepatitis B vaccine was not described before. Miyara et al. showed that blood contains two

distinct subsets of stable and suppressive TREG cells: resting TREG, identified as

FOXP3lowCD45RA+ CD4 T cells, and activated TREG, identified as FOXP3highCD45RA− CD4 T

cells. They further noted that activated TREG cells constitute a minority subset within cord blood

TREG cells and increase gradually with age [51]. As activated TREG cells were shown to have an

increased expression of proteins indicative of activation, including ICOS and HLA-DR [51–55], it

might be the case that an upregulation of 4-1BB is one more feature of this population or a subset

thereof. Moreover, TREG cells in mice were shown to modulate TFH formation and GC B cell

responses and to diminish antibody production in a CTLA-4 mediated suppression [56]. 682 Interestingly, CD45RA− TREG cells were shown to be more rich in preformed CTLA-4 stored in 683 intracellular vesicles compared to CD45RA+ TREG cells [51].

4-1BB was shown to be constitutively expressed by TREG cells [57] and that 4-1BB+ TREG cells are 685 functionally superior to 4-1BB− TREG cells in both contact-dependent and contact-independent 686 immunosuppression [58]. 4-1BB+ TREG cells are the major producers of the alternatively-spliced 687 and soluble isoform of 4-1BB among T cells [58]. 4-1BB was shown before to be preferentially 688 expressed on TREG cells compared with other non-regulatory CD4 T cell subsets [57] and that 4-689 1BB-costimualtion induces the expansion of TREG cells both in vitro and in vivo [59]. Moreover, 690 agonistic anti-4-1BB mAbs have been shown to abrogate T cell-dependent antibody responses in 691 vivo [60] and to ameliorate experimental autoimmune encephalomyelitis by skewing the balance 692 against TH17 differentiation in favor of TREG differentiation [61]. It is plausible that the expansion 693 4-1BB+CD45RA− TREG cells in late-converters is involved in the suppression of GC vaccine-694 specific TFH cells and the ensuing antibody response in our cohort, but this remains speculative 695 and further research is warranted.

697 It is enticing to speculate that the preexisting memory CD4 T cells result from the complex 698 interplay between cellular immunity and the human microbiome. A role for the microbiota in 699 modulating immunity to viral infection was suggested in 1960's [62], and since then we gained 700 better understanding of the impact of the various components of the microbiota including 701 bacteria, fungi, protozoa, archaea and viruses on the murine and human immune systems [63]. 702 Viral clearance of hepatitis B virus infection depends on the age of exposure and neonates and 703 young children are less likely to spontaneously clear the virus [64]. Han-Hsuan et al. have shown

evidence in mice that this age-dependency is mediated by gut microbiota that prepare the liver

immunity system to clear HBV, possibly via a TLR4 signaling pathway [65]. In this study, young

mice that have not reached an equilibrium in the gut microbiota, exhibited prolonged HBsAg

persistence, impaired anti-HBs antibody production, and limited Hepatitis B core antigen

(HBcAg)-specific IFNγ+ splenocytes. More recently, Tingxin et al. provided evidence for a

critical role of the commensal microbiota in supporting the differentiation of GC B cells, through

follicular T helper (TFH) cells, to promote the anti-HBV humoral immunity [66].

Our study bears some intrinsic limitations. A major drawback is the restricted number of days at

which TCRβ repertoire was profiled, as vaccine-specific perturbations within the repertoire may

occur at different time points for early, late and non-converters. Additionally, more in-depth

characterization and functional studies on 4-1BB+CD45RA− TREG cells could have helped shed

more light on the role they play in vaccine-induced immunity. Future studies in larger cohorts

and with a more comprehensive TCRβ repertoire profiling and CD4 T cells immunophenotyping

are required to validate our findings.

In conclusion, our analysis of the memory CD4 T cell repertoire has uncovered a role for

preexisting memory CD4 T cells in naïve individuals in mounting an earlier and more vigorous

immune response to hepatitis B vaccine and argue for the utility of pre-vaccination TCRβ

repertoire in the prediction of vaccine-induced immunity. Moreover, we identify a subset of 4-

1BB+ memory TREG cells that is expanded in individuals with delayed immune response to the

vaccine, which might further explain the heterogeneity of response to hepatitis B vaccine.

AUTHOR CONTRIBUTIONS

Conception: PM, BO

Design: GE, PM, EB, AS, GM, PVD, PB, KL, VVT, BO

Experiments: GE, PM, EB, NDN, HJ, AS

Data-analysis: GE, PM, NDN, BO

Supervision: HDR, EL, PT, GM, PVD, PB, KL, VVT, BO

First draft: GE, PM, BO

Contributed to the paper: all authors

COMPETING FINANCIAL INTERESTS

Parts of the contents of this manuscript form the topic of patent EPO 19159931.5.

VVT is an employee of Johnson & Johnson since 1/11/2019 and remains currently employed at the University of Antwerp.

References

743   1.   Rudolph, M. G., Stanfield, R. L. & Wilson, I. A. HOW TCRS BIND MHCS, PEPTIDES, AND CORECEPTORS. *Annu. Rev. Immunol.* **24**, 419–466 (2006).

2.   Esensten, J. H., Helou, Y. A., Chopra, G., Weiss, A. & Bluestone, J. A. CD28 Costimulation: From Mechanism to Therapy. *Immunity* **44**, 973–988 (2016).

3.   Curtsinger, J. M. & Mescher, M. F. Inflammatory cytokines as a third signal for T cell activation. *Curr. Opin. Immunol.* **22**, 333–40 (2010).

4.   Turner, S. J., La Gruta, N. L., Kedzierska, K., Thomas, P. G. & Doherty, P. C. Functional implications of T cell receptor diversity. *Current Opinion in Immunology* **21**, 286–290 (2009).

5.   Krangel, M. S. Mechanics of T cell receptor gene rearrangement. *Current Opinion in Immunology* **21**, 133–139 (2009).

6.   Farber, D. L., Yudanin, N. A. & Restifo, N. P. Human memory T cells: Generation, compartmentalization and homeostasis. *Nature Reviews Immunology* **14**, 24–35 (2014).

7.   Su, L. F., Kidd, B. A., Han, A., Kotzin, J. J. & Davis, M. M. Virus-Specific CD4+ Memory-Phenotype T Cells Are Abundant in Unexposed Adults. *Immunity* **38**, 373–383 (2013).

8.   Sewell, A. K. Why must T cells be cross-reactive? *Nature Reviews Immunology* **12**, 669–677 (2012).

9.   Wilson, D. B. *et al.* Specificity and degeneracy of T cells. *Mol. Immunol.* **40**, 1047–1055 (2004).

10. Reese, T. A. *et al.* Sequential Infection with Common Pathogens Promotes Human-like Immune Gene Expression and Altered Vaccine Response. *Cell Host Microbe* **19**, 713–719 (2016).

11. Keating, G. M. & Noble, S. Recombinant Hepatitis B Vaccine (Engerix-B). *Drugs* **63**, 1021–1051 (2003).

12. Shouval, D. Hepatitis B vaccines. *J. Hepatol.* **39**, 70–76 (2003).

13. Precopio, M. L. *et al.* Optimizing peptide matrices for identifying T-cell antigens. *Cytom. Part A* **73**, 1071–1078 (2008).

14. de Neuter, N. *et al.* Memory CD4+T cell receptor repertoire data mining as a tool for identifying cytomegalovirus serostatus. *Genes Immun.* (2018). doi:10.1038/s41435-018-0035-y

15. Gielis, S. *et al.* Detection of Enriched T Cell Epitope Specificity in Full T Cell Sequence Repertoires. *Front. Immunol.* **10**, (2019).

16. Meysman, P. *et al.* On the viability of unsupervised T-cell receptor sequence clustering for epitope preference. *Bioinformatics* **35**, 1461–1468 (2018).

17. Spidlen, J., Breuer, K., Rosenberg, C., Kotecha, N. & Brinkman, R. R. FlowRepository: A resource of annotated flow cytometry datasets associated with peer-reviewed publications. *Cytom. Part A* **81 A**, 727–731 (2012).

18. Furman, D. *et al.* Cytomegalovirus infection enhances the immune response to influenza. *Sci. Transl. Med.* **7**, 281ra43 (2015).

19. Klarenbeek, P. L. *et al.* Human T-cell memory consists mainly of unexpanded clones. *Immunol. Lett.* **133**, 42–48 (2010).

20. Becattini, S. *et al.* Functional heterogeneity of human memory CD4+ T cell clones primed by pathogens or vaccines. *Science (80-.).* **347**, 400–406 (2015).

21. De Neuter, N. *et al.* On the feasibility of mining CD8+ T cell receptor patterns underlying immunogenic peptide recognition. *Immunogenetics* **70**, 159–168 (2018).

22. Brezar, V. *et al.* T-regulatory cells and vaccination "pay attention and do not neglect them": Lessons from HIV and cancer vaccine trials. *Vaccines* **4**, (2016).

23. Schoenbrunn, A. *et al.* A converse 4-1BB and CD40 ligand expression pattern delineates activated regulatory T cells (Treg) and conventional T cells enabling direct isolation of alloantigen-reactive natural Foxp3+ Treg. *J. Immunol.* **189**, 5985–94 (2012).

24. Elias, G., Ogunjimi, B. & Van Tendeloo, V. Activation-induced surface proteins in the identification of antigen-responsive CD4 T cells. *Immunology Letters* **219**, 1–7 (2020).

25. Liu, W. *et al.* CD127 expression inversely correlates with FoxP3 and suppressive function of human CD4+ T reg cells. *J. Exp. Med.* **203**, 1701–1711 (2006).

26. Seddiki, N. *et al.* Expression of interleukin (IL)-2 and IL-7 receptors discriminates between human regulatory and activated T cells. *J. Exp. Med.* **203**, 1693–1700 (2006).

27. Bentebibel, S.-E., Schmitt, N., Banchereau, J. & Ueno, H. Human tonsil B-cell lymphoma 6 (BCL6)-expressing CD4+ T-cell subset specialized for B-cell help outside germinal centers. *Proc. Natl. Acad. Sci.* **108**, E488–E497 (2011).

28. Fonseca, V. R. *et al.* Human blood Tfr cells are indicators of ongoing humoral activity not fully licensed with suppressive function. *Sci. Immunol.* **2**, (2017).

29. Su, L. F. & Davis, M. M. Antiviral memory phenotype T cells in unexposed adults. *Immunol. Rev.* **255**, 95–109 (2013).

30. Meireles, L. C., Marinho, R. T. & Van Damme, P. Three decades of hepatitis B control with vaccination. *World J. Hepatol.* **7**, 2127–32 (2015).

31. Sallusto, F., Lanzavecchia, A., Araki, K. & Ahmed, R. From vaccines to memory and back. *Immunity* **33**, 451–463 (2010).

32. Ritchie, A. J. *et al.* Differences in HIV-Specific T Cell Responses between HIV-Exposed and -Unexposed HIV-Seronegative Individuals. *J. Virol.* **85**, 3507–3516 (2011).

33. Campion, S. L. *et al.* Proteome-wide analysis of HIV-specific naive and memory CD4+ T cells in unexposed blood donors. *J. Exp. Med.* **211**, 1273–1280 (2014).

34. Emerson, R. O. *et al.* Immunosequencing identifies signatures of cytomegalovirus exposure history and HLA-mediated effects on the T cell repertoire. *Nat. Genet.* **49**, 659–665 (2017).

35. DeWitt, W. S. *et al.* Human T cell receptor occurrence patterns encode immune history, genetic background, and receptor specificity. *Elife* **7**, (2018).

36. Kohler, S. *et al.* The early cellular signatures of protective immunity induced by live viral vaccination. *Eur. J. Immunol.* **42**, 2363–2373 (2012).

37. Blom, K. *et al.* Temporal Dynamics of the Primary Human T Cell Response to Yellow Fever Virus 17D As It Matures from an Effector- to a Memory-Type Response. *J. Immunol.* **190**, 2150–2158 (2013).

38. Pogorelyy, M. V. *et al.* Precise tracking of vaccine-responding T cell clones reveals convergent and personalized response in identical twins. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 12704–12709 (2018).

39. DeWitt, W. S. *et al.* Dynamics of the Cytotoxic T Cell Response to a Model of Acute Viral Infection. *J. Virol.* **89**, 4517–4526 (2015).
40. Frentsch, M. *et al.* Direct access to CD4+ T cells specific for defined antigens according to CD154 expression. *Nat. Med.* **11**, 1118–
1124 (2005).
41. Bacher, P. *et al.* Antigen-reactive T cell enrichment for direct, high-resolution analysis of the human naive and memory Th cell
repertoire. *J. Immunol.* **190**, 3967–76 (2013).
42. Reece, W. H. H. *et al.* A CD4+ T-cell immune response to a conserved epitope in the circumsporozoite protein correlates with
protection from natural Plasmodium falciparum infection and disease. *Nat. Med.* **10**, 406–410 (2004).
43. Bacher, P. *et al.* Identification of Immunogenic Antigens from Aspergillus fumigatus by Direct Multiparameter Characterization of
Specific Conventional and Regulatory CD4+ T Cells. *J. Immunol.* **193**, 3332–3343 (2014).
44. Bacher, P. *et al.* Antigen-specific expansion of human regulatory T cells as a major tolerance mechanism against mucosal fungi.
Mucosal Immunol. **7**, 916–928 (2014).
45. Rudensky, A. Y. Regulatory T cells and Foxp3. *Immunol. Rev.* **241**, 260–268 (2011).
46. Lovelace, P. & Maecker, H. T. Multiparameter Intracellular Cytokine Staining. in *Methods in molecular biology (Clifton, N.J.)* **1678**,
151–166 (2018).
47. Lin, P. H. *et al.* Vaccine-induced antigen-specific regulatory T cells attenuate the antiviral immunity against acute influenza virus
infection. *Mucosal Immunol.* **11**, 1239–1253 (2018).
48. Garner-Spitzer, E. *et al.* Tick-Borne Encephalitis (TBE) and Hepatitis B Nonresponders Feature Different Immunologic Mechanisms in
Response to TBE and Influenza Vaccination with Involvement of Regulatory T and B Cells and IL-10. *J. Immunol.* **191**, 2426–2436
(2013).
49. Lanteri, M. C. *et al.* Tregs control the development of symptomatic West Nile virus infection in humans and mice. *J. Clin. Invest.* **119**,
3266–3277 (2009).
50. Sanchez, A. M., Zhu, J., Huang, X. & Yang, Y. The Development and Function of Memory Regulatory T Cells after Acute Viral
Infections. *J. Immunol.* **189**, 2805–2814 (2012).
51. Miyara, M. *et al.* Functional Delineation and Differentiation Dynamics of Human CD4+ T Cells Expressing the FoxP3 Transcription
Factor. *Immunity* **30**, 899–911 (2009).
52. Booth, N. J. *et al.* Different Proliferative Potential and Migratory Characteristics of Human CD4 + Regulatory T Cells That Express
either CD45RA or CD45RO. *J. Immunol.* **184**, 4317–4326 (2010).
53. Ito, T. *et al.* Two Functional Subsets of FOXP3+ Regulatory T Cells in Human Thymus and Periphery. *Immunity* **28**, 870–880 (2008).
54. Mason, G. M. *et al.* Phenotypic Complexity of the Human Regulatory T Cell Compartment Revealed by Mass Cytometry. *J. Immunol.*
195, 2030–2037 (2015).
55. Mohr, A., Malhotra, R., Mayer, G., Gorochov, G. & Miyara, M. Human FOXP3+ T regulatory cell heterogeneity. *Clinical and*
Translational Immunology **7**, (2018).
56. Wing, J. B., Ise, W., Kurosaki, T. & Sakaguchi, S. Regulatory T cells control antigen-specific expansion of Tfh cell number and
humoral immune responses via the coreceptor CTLA-4. *Immunity* **41**, 1013–1025 (2014).
57. McHugh, R. S. *et al.* CD4(+)CD25(+) immunoregulatory T cells: gene expression analysis reveals a functional role for the
glucocorticoid-induced TNF receptor. *Immunity* **16**, 311–23 (2002).
58. Kachapati, K. *et al.* The B10 Idd9.3 locus mediates accumulation of functionally superior CD137(+) regulatory T cells in the nonobese
diabetic type 1 diabetes model. *J. Immunol.* **189**, 5001–15 (2012).
59. Zheng, G., Wang, B. & Chen, A. The 4-1BB costimulation augments the proliferation of CD4+CD25+ regulatory T cells. *J. Immunol.*
173, 2428–34 (2004).
60. Mittler, R. S., Bailey, T. S., Klussman, K., Trailsmith, M. D. & Hoffmann, M. K. Anti-4-1BB monoclonal antibodies abrogate T cell-
dependent humoral immune responses in vivo through the induction of helper T cell anergy. *J. Exp. Med.* **190**, 1535–40 (1999).
61. Kim, Y. H. *et al.* 4-1BB Triggering Ameliorates Experimental Autoimmune Encephalomyelitis by Modulating the Balance between
Th17 and Regulatory T Cells. *J. Immunol.* **187**, 1120–1128 (2011).
62. Robinson, C. M. & Pfeiffer, J. K. Viruses and the Microbiota. *Annu. Rev. Virol.* **1**, 55–69 (2014).
63. Winkler, E. S. & Thackray, L. B. A long-distance relationship: the commensal gut microbiota and systemic viruses. *Current Opinion in*
Virology **37**, 44–51 (2019).
64. Yuen, M.-F. *et al.* Hepatitis B virus infection. *Nat. Rev. Dis. Prim.* **4**, 18035 (2018).
65. Chou, H. H. *et al.* Age-related immune clearance of hepatitis B virus infection requires the establishment of gut microbiota. *Proc. Natl.*
Acad. Sci. U. S. A. **112**, 2175–2180 (2015).
66. Wu, T. *et al.* CD4+ T cells play a critical role in microbiota-maintained Anti-HBV immunity in a mouse model. *Front. Immunol.* **10**,
(2019).
861
862

Supplemental tables

Supplementary *Table 1.* List of 54 single peptides, each 15 AA long with an 11-amino acid overlap spanning the 226 amino acids along the small S protein of hepatitis B (HB) surface antigen (HBsAg)

Peptide No.	15 AA with 11 AA overlap	No. of AA at which the peptide start
1	MENITSGFLGPLLVL	1
2	TSGFLGPLLVLQAGF	5
3	LGPLLVLQAGFFLLT	9
4	LVLQAGFFLLTRILT	13
5	AGFFLLTRILTIPQS	17
6	LLTRILTIPQSLDSW	21
7	ILTIPQSLDSWWTSL	25
8	PQSLDSWWTSLNFLG	29
9	DSWWTSLNFLGGSPV	33
10	TSLNFLGGSPVCLGQ	37
11	FLGGSPVCLGQNSQS	41
12	SPVCLGQNSQSPTSN	45
13	LGQNSQSPTSNHSPT	49
14	SQSPTSNHSPTSCPP	53
15	TSNHSPTSCPPICPG	57
16	SPTSCPPICPGYRWM	61
17	CPPICPGYRWMCLRR	65
18	CPGYRWMCLRRFIIF	69
19	RWMCLRRFIIFLFIL	73
20	LRRFIIFLFILLLCL	77
21	IIFLFILLLCLIFLL	81
22	FILLLCLIFLLVLLD	85
23	LCLIFLLVLLDYQGM	89
24	FLLVLLDYQGMLPVC	93
25	LLDYQGMLPVCPLIP	97
26	QGMLPVCPLIPGSTT	101
27	PVCPLIPGSTTTNTG	105
28	LIPGSTTTNTGPCKT	109
29	STTTNTGPCKTCTTP	113
30	NTGPCKTCTTPAQGN	117
31	CKTCTTPAQGNSMFP	121

32	TTPAQGNSMFPSCCC	125
33	QGNSMFPSCCCTKPT	129
34	MFPSCCCTKPTDGNC	133
35	CCCTKPTDGNCTCIP	137
36	KPTDGNCTCIPIPSS	141
37	GNCTCIPIPSSWAFA	145
38	CIPIPSSWAFAKYLW	149
39	PSSWAFAKYLWEWAS	153
40	AFAKYLWEWASVRFS	157
41	YLWEWASVRFSWLSL	161
42	WASVRFSWLSLLVPF	165
43	RFSWLSLLVPFVQWF	169
44	LSLLVPFVQWFVGLS	173
45	VPFVQWFVGLSPTVW	177
46	QWFVGLSPTVWLSAI	181
47	GLSPTVWLSAIWMMW	185
48	TVWLSAIWMMWYWGP	189
49	SAIWMMWYWGPSLYS	193
50	MMWYWGPSLYSIVSP	197
51	WGPSLYSIVSPFIPL	201
52	LYSIVSPFIPLLPIF	205
53	VSPFIPLLPIFFCLW	209
54	IPLLPIFFCLWVYI	213

Supplementary Table 2. Overview of the single peptides tested for each vaccinee in the CFSE assay

Vaccinee	Gender	Age	Status	peptides_number	single peptides
2	F	43.6	Late-converter	6	51, 35, 50, 34, 54, 38
3	F	48.5	Late-converter	6	53, 5, 37, 49, 1, 33
4	M	47.1	Late-converter	6	21, 20, 5, 53, 4, 52
6	F	44.3	Late-converter	6	23, 39, 7, 21, 37, 5
7	M	38.3	Early-converter	2	17, 33
8	M	43.4	Early-converter	3	47, 41, 42
10	F	33.2	Early-converter	5	49, 1, 33, 41, 9
11	F	39	Early-converter	6	53, 50, 5, 2, 8, 21
13	F	45.9	Early-converter	3	24, 8, 16
14	M	36.3	Late-converter	2	45, 47
17	F	26.5	Early-converter	6	10, 42, 9, 41, 14, 46
18	M	43.3	Early-converter	4	6, 38, 5, 37
19	M	41.3	Early-converter	3	40, 16, 8
20	F	48.7	Early-converter	2	6, 1
21	F	27.2	Non-converter	6	2, 1, 3, 42, 41, 43
22	F	22.3	Early-converter	4	10, 13, 12, 16
23	F	43.9	Late-converter	6	51, 54, 48, 43, 47, 46
24	F	47.7	Late-converter	6	22, 24, 23, 20, 18, 46
25	F	22.6	Non-converter	6	29, 30, 28, 21, 22, 20
26	F	41.2	Late-converter	6	23, 47, 17, 41, 7, 49
28	F	41.7	Early-converter	6	54, 38, 52, 36, 40, 39
29	M	39.5	Early-converter	6	7, 47, 6, 1, 23, 46
30	F	45.1	Late-converter	6	22, 46, 54, 19, 20, 43
31	M	32.4	Early-converter	6	54, 38, 49, 33, 53, 37
32	F	31.4	Early-converter	6	7, 4, 1, 6, 31, 28
33	F	41	Early-converter	6	1, 41, 7, 5, 47, 45
34	M	22.5	Early-converter	5	7, 4, 5, 2, 6
35	F	50.2	Early-converter	6	46, 45, 47, 22, 21, 23
36	F	23.2	Early-converter	6	54, 47, 46, 39, 48, 38
38	F	23.2	Early-converter	6	37, 34, 33, 35, 5, 2
39	M	45.8	Early-converter	6	51, 50, 53, 49, 35, 34
40	F	21.3	Early-converter	5	33, 38, 37, 39, 40
41	M	21.6	Non-converter	6	23, 22, 7, 6, 17, 1
42	M	22.7	Non-converter	4	1, 33, 25, 41

	CD45RA	CXCR5	CD25	CD127
Naive T$_H$	+	—	—	low + high
Memory T$_H$	—	—	—	low + high
Naive T$_{FH}$	+	+	—	low + high
Memory T$_{FH}$	—	+	—	low + high
CD45RA$^+$ T$_{REG}$	+	—	+	low
CD45RA$^-$ T$_{REG}$	—	—	+	low
CD45RA$^+$ T$_{FR}$	+	+	+	low
CD45RA$^-$ T$_{FR}$	—	+	+	low

Figure S1. Gating strategy of ex vivo T cell phenotyping of vaccine-specific T cells.

a Gating strategy started by a lymphocyte gate, followed by gating on viable CD3+CD8– T cells. Doublets were excluded using
doublet discrimination (area against the height of forward scatter pulse) before gating on CD4+ T
cells. Next, CD45RA, CXCR5, CD25 and CD127 were used to identify main subsets of CD4 T
cells using Boolean gates as specified in the accompanying table.
b Shown an example of gating for CD154 (CD40L) and CD137 (4-1BB) for cells left
unstimulated (left) and cells stimulated with a master peptide pool (right) for an early-converter
vaccinee at day 60.

883
Figure S2.

a Anti-Hepatitis B surface (anti-HBs) titer of vaccinees over four times points, facetted by
groups of early, late and non-converters. An anti-HBs titer above 10 IU/ml was considered
protective. Early-converters seroconverted at day 60, late-converters seroconverted at day 180 or
day 365 and non–converters did not have an anti-HBs titer higher than 10 IU/ml at any of the
time points. **b** CMV, EBV and HSV seropositivity in the three groups of the cohort as
determined by serum IgG antibodies to CMV, EBV-VCA, and HSV-1 and 2 using sandwich
ELISA. **c** Age of vaccinees per group.
Wilcoxon signed-rank with unpaired and paired analysis as necessary; statistical significance
was indicated with ns P > 0.05, * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, **** P ≤ 0.0001
894

895
Figure S3

a Scatter plot of the DNA-based TCRβ reads for each vaccinee at each time point

b Scatter plot of number of unique TCRβ amino acid sequences for each vaccinee at each time

point, where the shape denotes the response as based on antibody titer.

c Overview of unique TCRβ amino acid sequences in the memory CD4 T cell repertoire of each

vaccinee. The bottom blue bar denotes those TCR sequences that were found at both time points.

The green and red bars denote the number of unique TCR sequences at each time point. The total

bar height thus represents the total number of unique memory CD4 T cell clonotypes sequences

for a specific vaccinee.

d Change in frequency of those HBV-specific T-cells present at both time points. The (ns) mark

denotes a non-significant paired Wilcoxon signed-rank test (P value = 0.7577).

e Antibody-defined categories (EC, LC, NC)) plotted against the values from the CD40L assay

f ROC curve for the Rhbs predictive model from day 0 data in a leave-one-out cross-validation

compared to CD40L+4-1BB- (TCON) memory CD4 T cell values for each vaccinee at time

points 60 (AUC = 0.84), 180 (AUC = 0.56) and 365 (AUC = 0.57).

g ROC curve for the Rhbs predictive model from day 0 data in a leave-one-out cross-validation

compared to CD40L-4-1BB+ (TREG) memory CD4 T cell values for each vaccinee at time

points 60 (AUC = 0.62), 180 (AUC = 0.56) and 365 (AUC = 0.52).

In vitro expansion of CD4 T cells (7 days)

915
916 **Figure S4. Overview of the in vitro expansion experiments for all vaccinees, vaccinees per**
917 **group and for each vaccinee.**

918
919 **Figure S5. CD40L+4-1BB– and CD40L–4-1BB+ CD4 T cells have a TCON and TREG**
920 **phenotype, respectively.**

Correlation of vaccine-specific memory CD4 T cells at day 60 and antibody titer difference between day 365 and day 0

Figure S6

Correlation between the difference in antibody titer between day 365 and day 0 and vaccine-specific CD40L+4-1BB– and CD40L–4-1BB+ memory CD4 T cell at day 60 colored by vaccinee group and labeled with vaccinee ID.

rs, Spearman correlation coefficient, $-1 \leq rs \leq 1$; rs and p value by Spearman's correlation test

Figure S7 Hepatitis B vaccine induces a vaccine-specific CD4 T cell response in early-converter vaccinees.

PBMCs from vaccinees were stimulated with 2 μg/ml of a pool of peptides of HBsAg and assessed for converse expression of 4-1BB and CD40L by flow cytometry on days 0, 60, 180, and 365.

a Aggregate analysis from early, late and non-converter vaccinees showing a significant induction of vaccine-specific CD40L+4-1BB− and CD40L−4-1BB+ memory CD4 T cell in early-converters and lack thereof in late and non-converters. Shown is number of of vaccine-specific memory CD4 T cells out of $1x10^6$ memory CD4 T cells after subtraction of responses in negative control (see Methods for details).

b Aggregate analysis from early, late and non-converter vaccinees showing a significant induction of vaccine-specific CD40L+4-1BB− and CD40L−4-1BB+ CD4 T cell in early-converters and lack thereof in late and non-converters. Shown is number of of vaccine-specific CD4 T cells out of $1x10^6$ CD4 T cells after subtraction of responses in negative control (see Methods for details).

945
Figure S8

947 Aggregate analysis of 4-1BB+CD45RA– T$_{REG}$ within CD45RA– T$_{REG}$ CD4 T cells in early, late
948 and non-converter vaccinees at days 0, 60, 180 and 365.
949 Wilcoxon signed-rank with unpaired and paired analysis as necessary; statistical significance
950 was indicated with ns P > 0.05, * P ≤ 0.05, ** P ≤ 0.01, *** P ≤ 0.001, **** P ≤ 0.0001
951