1   **Value Certainty in Diffusion Decision Models**

2   Douglas Lee[1] & Marius Usher[2]

3   [1] California Institute of Technology, Division of the Humanities and Social Sciences

4   [2] Tel Aviv University, School of Psychological Sciences

11   E-mail for correspondence:

12   DouglasGLee@gmail.com

14   Keywords:

15   Word count:

Abstract

Decision models such as the drift-diffusion model (DDM) are a widely used and broadly

accepted tool that performs remarkably well in accounting for binary choices and their response

time distributions, as a function of the option values. Such models are built on an evidence

accumulation to bound concept, where a decision maker repeatedly samples a mental

representation of the values of the options on offer until satisfied that there is enough evidence

in favor of one option over the other. The value estimates that drive the DDM evidence are

derived from the relative strength of value signals that are not stable across time, so that

repeated sequential samples are necessary to average out noise. The DDM, however, typically

does not allow for different options to have different levels of variability in their value

representations. However, studies have shown that a decision maker often reports levels of

certainty regarding value estimates that vary across options. We thus propose that future

versions of DDM should include an option-specific value certainty component. We present four

different versions of such a model and validate them against empirical data from four previous

studies. The data show that a model built around a sort of signal-to-noise ratio for each option

(rather than a pure signal that randomly fluctuates) performs best, accounting for the positive

impact of value certainty on choice consistency and the negative impact of value certainty on

response time.

The drift-diffusion model (DDM) is ubiquitous in the contemporary literature on decision

making, including research spanning the fields of psychology, neuroscience, economics, and

consumer behavior. The DDM is a parsimonious mechanism that explains normative decisions by

averaging out noise in information processing and implementing an optimal stopping rule

(optimizing response time (RT) for a specified accuracy). This model accounts well for the

dependency of the RT distribution on the values of the choice options, and for the speed-accuracy

tradeoff—the oft-observed phenomenon that faster choices are typically less accurate (see

Ratcliff et al, 2016; Gold and Shadlen, 2007 for reviews). While initially used in the domain of

perceptual decision-making, the DDM has since become a widespread tool in studies of

preferential (i.e., value-based) decision-making as well (Busemeyer et al, 2019; Tajima,

Drugowitsch, & Pouget, 2016; Polania et al, 2015; Philiastides & Ratcliff, 2013; Milosavljevic et al,

2010; Krajbich, Armel, & Rangel, 2010; Basten et al, 2010). In brief, in its application to

preferential choice, the DDM assumes that the value of each option is represented by a

probability distribution whose mean is the true value of the option, and whose variance

corresponds to processing noise. This noise can be interpreted as imprecision in the value

representations themselves, or as a stochastic distortion of the value signals as they are relayed

through the decision system by populations of neurons (whose firing patterns are known to be

stochastic). Either way, the momentary signal about the relative value of the options (the so-

called evidence for one option over the other) fluctuates randomly around a fixed trajectory (the

so-called drift). In order to average out this processing noise, evidence signals are thought to

accumulate over time until a sufficient amount of evidence has been acquired to allow for a

choice to be made. The DDM thus includes thresholds for each option (typically symmetric) that

trigger a choice once reached by the evidence accumulator. So, the fundamental components of

the drift-diffusion process are the drift rate (the difference in the option values), the diffusion

coefficient (the degree of stochasticity of the system), and the choice boundaries (the minimum

required evidence threshold; see Figure 1).

Figure 1: An illustration of the classic basic DDM. Evidence accumulates across time, following a

fixed drift trajectory (black dashed arrow, corresponding to the value difference of choice

options), corrupted by white processing noise. Here, the accumulated evidence reaches the

upper threshold after 3.9 seconds, and a choice for option 1 is recorded.

The standard DDM implicitly assumes that processing noise is independent of the identity

of the options contained in a particular choice set. That is to say, there is no option-specific noise

in the DDM (but see Ratcliff, Voskuilen, & Teodorescu, 2018; Teodorescu, Moran, & Usher, 2016

for DDM variants in which the noise increases with task difficulty). Most of the DDM applications

to preference-based decisions have assumed no option-specific noise. Since the brain encodes

not only the subjective value of options, but also the subjective certainty about the value

(Lebreton et al, 2015), it is possible to suggest that the representations of value that the brain

uses to inform the decision process fluctuate, and that the degree of imprecision (or uncertainty)

is not the same for all choice options. Indeed, it has been shown that decision makers hold highly

variable beliefs about the certainty of their value estimates for different options (Lee & Coricelli,

2020; Lee & Daunizeau, 2020a, 2020b; Gwinn & Krajbich, 2020; Polania, Woodford, & Ruff, 2019).

It has further been shown that the variability in value (un)certainty has a clear impact on both

choice and RT (Lee & Coricelli, 2020; Lee & Daunizeau, 2020a, 2020b). Specifically, judged value

certainty, C, correlates positively with choice consistency and negatively with RT (see Figure 2).

This provides a qualitative benchmark that any DDM variant that includes option-specific

uncertainty should be able to account for.

Figure 2: Previous results (pooled across four studies, n=152) demonstrate the relationships

between value difference (dV), value certainty (C= .5 (c1+c2)), and certainty difference (dC= c1

−c2) with choice consistency and log(RT). Here we show the beta weights from a logistic

regression on choice (upper left) and a linear regression on log(RT) (upper right; bar heights

represent population means, error bars represent s.e.m.). Note: dV, C, and dC were all

simultaneously included as independent variables in each regression model. Lower plots show

the isolated relationships between log(RT) and C (left) and C1 - C2 (right). Note: the distribution

of C was highly skewed, so we show the data binned by quantiles. (Lee & Daunizeau, 2020a,

2020b; Lee & Coricelli, 2020; Gwinn & Krajbich, 2020)

As we show below, the most straightforward way to include option-specific noise in the

preferential DDM – by assuming that noise increases with value uncertainty – leads to the wrong

qualitative predictions, with regards to the RT x certainty benchmark. In particular, as certainty

increases, noise decreases, resulting in lower RT. Thus, unlike what we see in experimental data,

such a model would predict that people would speed up when they are less certain of the options'

values (all else equal).

The aim of this paper is to examine a number of DDM variants, which could potentially

rise to the challenge of accounting for the behavioral value certainty, choice, and RT data. In

particular, we will first present a variety of derivations of the standard DDM, each of which

incorporates the concept of option-specific value certainty in a unique and realistic way, starting

from the default DDM that has no option-specific noise, and progressing to signal-to-noise type

of models. We then fit each of these models to experimental data from a variety of empirical

datasets. Finally, we quantitatively compare the performance of each model across the different

datasets and suggest the best recommended approach for future studies to incorporate option-

specific value certainty in models derived from the DDM. To anticipate our results, we find that

a signal-to-noise DDM variant provides the best fit to the data and accounts for all qualitative

benchmarks.

METHODS

Computational models

In each of the models described below, we consider decisions between two alternatives,

with values, v1 and v2, and with uncertainties, sigma1, sigma2, respectively. Evidence of these

values is integrated over deliberation time, subject to noise. The evidence accumulator for each

decision is initialized at a neutral starting point (i.e., zero evidence or default bias in favor of

either option), and evolves until reaching one of two symmetric boundaries (see Figure 1 above).

For each decision, the output of the model is: which boundary is reached, or the choice (ch = {0,

1}) and the number of integration time steps elapsed when that boundary is reached (RT).

Model 1

As a baseline default model for comparison (without any option-specific certainty term),

we first consider the classic basic DDM. In this model, the equations that govern the evidence

accumulation process are:

$$E_{t+1} = E_t + d\Delta$$

$$\Delta \ N(\mu_1 - \mu_2, \sigma^2)$$

where E represents the cumulative balance of evidence that one option is better than the

other, t represents deliberation time, Δ represents the incremental evidence in favor of one

option over the other, d is a scalar, μ_i is the value estimate of option i, and σ^2 represents

processing noise in the evidence accumulation and/or comparator systems. In a standard basic

DDM, choice probability and expected response time can be analytically calculated using the

following equations:

$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{2\theta d}{\sigma^2}(\mu_1 - \mu_2)\right)}}$$

$$RT = \frac{\theta(2 * p(ch = 1) - 1)}{d(\mu_1 - \mu_2)}$$

where θ is the height of the evidence accumulation threshold where a decision is triggered

(shown here as the upper threshold, for a choice of option 1), p(ch=1) is the probability that the

upper threshold will be reached (rather than the lower threshold), and RT is the expected time

at which the accumulation process will end (by reaching the upper threshold). Because this

system of equations is over-parameterized, one of the free parameters must be fixed for a

practical application of the equations. In this work, we will fix the threshold θ to a value of 1

when fitting the models, for simplicity. Choice probability and RT will thus be functions of the

drift rate d and the noise σ^2 (see Figure 3). As expected, accuracy increases and RT decreases with

the drift. The noise, on the other hand, reduces both accuracy and RT.

Figure 3: With all other parameters fixed, an increase in drift rate leads to a sigmoidal increase

in the probability of choosing option 1 from 0.5 (random guess) to 1, and a sigmoidal decrease

in RT (left plots in blue). With all other parameters fixed, an increase in processing noise will

lead to a sigmoidal decrease in the probability of choosing option 1, and a parallel sigmoidal

decrease in RT (right plots in red).

Model 2

The simplest and most obvious solution to incorporate option-specific uncertainty into

the DDM would be to model the process as:

$$E_{t+1} = E_t + d\Delta$$

$$\Delta \ N(\mu_1 - \mu_2, \sigma_1^2 + \sigma_2^2)$$

which is simply the standard DDM equation capturing the evolution of the accumulated evidence

across time, but with σ_i^2 representing the uncertainty about the value estimate of option i. The

only difference between this formulation and the standard one is that here the variance of Δ is

specific to the options in the current choice set, whereas in the standard DDM it is fixed across

choices (for an individual decision maker). A direct result of this reformulation is that choices

between options with greater value uncertainty (lower C) will be more stochastic and take less

time (on average), as can be seen by examining the (revised) DDM equations for choice

probability and expected response time (see also Fig 3, red lines):

169
$$p(ch = 1) = \cfrac{1}{1 + e^{\left(\frac{2\theta d(\mu_1 - \mu_2)}{\sigma_1^2 + \sigma_2^2}\right)}}$$

170
$$RT = \frac{\theta(2 * p(ch = 1) - 1)}{d(\mu_1 - \mu_2)}$$

Model 3

An alternative way in which the concept of option-specific value (un)certainty could be

incorporated into the DDM would be through a signal-to-noise dependency in the drift rate. The

drift rate in the DDM symbolizes the accumulation of evidence for one option over the other,

equal to the value estimate of one option minus the value estimate of the other option (scaled

by a fixed term). The accumulator variable is referred to as "evidence" because the probability

distributions controlling it (or the neural activity driving it) are thought to provide reliable signal

that will accurately inform the decision. If the value representations of different options can have

different levels of uncertainty, it stands to reason that the reliability of the "evidence" that these

signals provide about the correct decision will also be different. As such, evidence drawn from a

more reliable source (i.e., one with higher certainty) should be weighted more heavily. Under

this framework, the equation governing the DDM process would be:

183
$$E_{t+1} = E_t + d\Delta$$

184
$$\Delta\ N\left(\frac{\mu_1 - \mu_2}{\sigma_1^2 + \sigma_2^2}, \sigma^2\right)$$

where σ² (without a subscript) is the noise in the system unrelated to specific choice options. The

only difference between this formulation and the standard one is that here the mean of Δ is

divided by the variance of the choice options. A direct result of this reformulation is that choices

between options with greater value uncertainty will be more stochastic and also take *more* time

(on average), as can be seen by examining the (revised) DDM equations for choice probability

and expected response time:

191
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{2\theta d(\mu_1 - \mu_2)}{\sigma^2(\sigma_1^2 + \sigma_2^2)}\right)}}$$

192
$$RT = \frac{\theta(2 * p(ch = 1) - 1)}{\frac{d(\mu_1 - \mu_2)}{\sigma_1^2 + \sigma_2^2}}$$

Here the impact of option-specific uncertainty on RT is more complex. First, greater

uncertainty decreases RT through its effect on choice stochasticity (as before). Second, greater

uncertainty directly increases RT by diminishing the slope of the drift rate. The second effect

dominates.

Model 4

A variant of the DDM in which the drift rate is altered by the option-specific value

certainty could be one in which the evidence in favor of each option is scaled by its own precision

term before the evidence for the different options is compared. The drift rate would thus become

the accumulation of adjusted evidence for one option over the other, equal to the precision-

weighted value estimate of one option minus the precision-weighted value estimate of the other

option (scaled by a fixed term). Here, the evidence drawn from a more reliable source (i.e., one

with higher certainty) will be weighted more heavily (as in Model 3) but prior to comparison

between the alternative options. Under this framework, the equation governing the DDM

process would be:

207
$$E_{t+1} = E_t + d\Delta$$

208
$$\Delta\; N\left(\frac{\mu_1}{\sigma_1^2} - \frac{\mu_2}{\sigma_2^2}, \sigma^2\right)$$

The only difference between this formulation and the standard one is that here the mean

of Δ is adjusted by the variance of the choice options. Because the evidence in favor of each

option (prior to comparison) will be scaled by its own specific (and importantly, potentially

different) precision term, the impact on both choice stochasticity and response time could go in

either direction. This can be seen by examining the (revised) DDM equations for choice

probability and expected response time:

215
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{2\theta d}{\sigma^2}\left(\frac{\mu_1}{\sigma_1^2} - \frac{\mu_2}{\sigma_2^2}\right)\right)}}$$

216
$$RT = \frac{\theta(2 * p(ch = 1) - 1)}{d\left(\frac{\mu_1}{\sigma_1^2} - \frac{\mu_2}{\sigma_2^2}\right)}$$

Here the impact of option-specific uncertainty on both choice and RT is more complex

than in the other models presented above. If the evidence stream for one option is has both a

larger mean and a smaller variance, relative to the other option, the effective drift rate will be

higher than in a standard DDM (e.g., choices will be less stochastic and faster). On the other

hand, if the option with the larger mean evidence stream is different from the option with the

more reliable evidence stream, the effective drift rate will be lower than in a standard DDM (e.g.,

choices will be more stochastic and slower).

Model 5

Yet another alternative way in which the DDM could include option-specific value

certainty is in the form of an independent evidence accumulator -- a secondary drift, with a rate

proportional to the difference in certainty between the choice options. In this way, the decision

about which option to choose would be influenced both by which value estimate was higher (via

the primary, standard drift) and by which value estimate was more certain (via the secondary,

novel drift). The secondary drift might represent an aversion (or attraction) to risk or ambiguity,

where the deliberation process would be both slowed (or hastened) by the risk/ambiguity and

pulled towards the less (or more) uncertain option.   Under this framework, the equation

governing the DDM process would be:

234
$$E_{t+1} = E_t + d_v \Delta_v + d_c \Delta_c$$

235
$$\Delta_v \; N(\mu_1 - \mu_2, \sigma^2)$$

236
$$\Delta_c \; N(\sigma_1^2 - \sigma_2^2, \sigma^2)$$

where Δ_v and Δ_c are the incremental evidence for value and certainty, respectively, and d_v and d_c

are scalars (it is assumed that d_v will always be positive, but d_c could take either sign). The

inclusion of a secondary drift rate (in essence, a parallel evidence stream that monitors value

certainty rather than value itself) can have either a positive or a negative impact on both choice

and RT, depending on whether the option with the higher mean evidence stream is the same as

or different than the option with the higher evidence reliability (as well as on the sign of d_c). This

can be seen by examining the (revised) DDM equations for choice probability and expected

response time:

245
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{2\theta}{\sigma^2}[d_v(\mu_1 - \mu_2) + d_c(\sigma_1^2 - \sigma_2^2)]\right)}}$$

246
$$RT = \frac{\theta(2 * p(ch = 1) - 1)}{d_v(\mu_1 - \mu_2) + d_c(\sigma_1^2 - \sigma_2^2)}$$

Here, the secondary drift rate will result in more consistent and faster choices if the sign

of $\mu_1-\mu_2$ is the same as that of $\sigma_1^2-\sigma_2^2$, but it will result in less consistent and slower choices

otherwise. This, of course, is under the assumption that the decision maker is risk/ambiguity

averse (i.e., $d_c>0$). If the decision maker were risk/ambiguity seeking (i.e., $d_c<0$), the opposite

predictions would hold.

Materials and Design

Using a variety of different datasets from previous studies, one at a time, we fit

experimental data to each of the models that we described above. We then performed Bayesian

model comparison to determine which of the models (if any) performed better than the others

across the population of participants. For this model fitting and comparison exercise, we relied

on the Variational Bayesian Analysis toolbox (VBA, available freely at https://mbb-

team.github.io/VBA-toolbox/; Daunizeau, Adam, & Rigoux, 2014) for Matlab R2020a. We used

the VBA_NLStateSpaceModel function to fit the data for each participant individually, followed

by the VBA_groupBMC function to compare the results of the model fitting across models for the

full group of participants. The input for the model inversion was a series of two-alternative forced

choice (2AFC) data, including measures of value estimate and value estimate certainty for each

choice option, the chosen option (left or right) for each trial, and response time (RT) for each

trial. Some datasets also included choice confidence reports for each trial, which were included

in supplementary analyses as will be described below. The parameters to be fitted included all

of the d and σ^2 terms described above, plus additional parameters for an affine transformation

of experimental certainty measures into theoretical ones. This is necessary because in the

experimental data, the measures of value and those of value certainty reside on the same scale,

but this is likely untrue for the cognitive variables that are meant to drive the models.

Dataset 1

The first dataset we examined was from Lee & Daunizeau (2020a). In this study,

participants made choices between various snack food options based on their personal subjective

preferences. Value estimates for each option were provided in a separate rating task prior to the

choice task. Participants used a slider scale to respond to the question, "Does this please you?"

After each rating, participants used a separate slider scale to respond to the question, "Are you

sure?" This provided a measure of value estimate certainty for each item. During the choice

task, participants were presented with pairs of snack food images and asked, "What do you

prefer?" After each choice, participants used a slider scale to respond to the question, "Are you

sure about your choice?" to provide a subjective report of choice confidence. This dataset

contained 51 subjects, each of whom were faced with 54 choice trials.

Dataset 2

The second dataset we examined was from Lee & Daunizeau (2020b). In this study,

participants made choices between various snack food options based on their personal subjective

preferences. Value estimates for each option were provided in a separate rating task prior to the

choice task. Participants used a slider scale to respond to the question, "How much do you like

this item?" After each rating, participants used the same slider scale to respond to the question,

"How certain are you about the item's value?" by indicating a zone in which they believed the

value of the item surely fell. This provided a measure of value estimate certainty for each item.

During the choice task, participants were presented with pairs of snack food images and asked,

"Which do you prefer?" After each choice, participants used a slider scale to respond to the

question, "Are you sure about your choice?" to provide a subjective report of choice confidence.

This dataset contained 32 subjects, each of whom were faced with 74 choice trials.

Dataset 3

The third dataset we examined was from Lee & Coricelli (2020). In this study, participants

made choices between various snack food options based on their personal subjective

preferences. Value estimates for each option were provided in a separate rating task prior to the

choice task. Participants used a slider scale to respond to the question, "How pleased would you

be to eat this?" After each rating, participants used a six-point descriptive scale to respond to

the question, "How sure are you about that?" This provided a measure of value estimate

certainty for each item. During the choice task, participants were presented with pairs of snack

food images and asked, "Which would you prefer to eat?" After each choice, participants used a

slider scale to respond to the question, "How sure are you about your choice?" to provide a

subjective report of choice confidence. This dataset contained 47 subjects, each of whom were

faced with 55 choice trials.

Dataset 4

The fourth dataset we examined was from Gwinn & Krajbich (2020). In this study,

participants made choices between various snack food options based on their personal subjective

preferences. Value estimates for each option were provided in a separate rating task prior to the

choice task. Participants used a 10-point numerical scale to respond to the prompt, "Please

indicate how much you want to eat this item." After each rating, participants used a seven-point

numerical scale to respond to the prompt, "Please indicate how confident you are in your rating

of this item." This provided a measure of value estimate certainty for each item. During the

choice task, participants were presented with pairs of snack food images and instructed to choose

the one that they preferred to eat. Choice confidence was not measured in this study. This

dataset contained 36 subjects, each of whom were faced with 200 choice trials.

RESULTS

Model comparison

The classic basic DDM, our Model 1, has been validated countless times for its ability to

account for two-alternative forced choice responses and response time distributions. The other

models we described above, Models 2-5, are new and have therefore never been tested with

empirical data. Thus, we start our model comparison exercise with one-on-one competitions

between Model 1 and each of Models 2-5, separately. This serves as a simple test of whether

the addition of the option-specific value estimate certainty term, as suggested in each of the four

different manners described in Models 2-5, improves the fit of the classic DDM. We then perform

a comparison across all five models simultaneously, and test whether any of them dominates the

others in terms of best fit to the data.

Before we present the model-fitting results, we first show the qualitative predictions that

each model makes with respect to the effects of value difference, value certainty, and certainty

difference on choice consistency, RT, and choice confidence (see Figure 4). We first simulated

10^7 trials for each model, based on random input values with similar distributions as in our

experimental data. The model-parameters we used were in line with the best fit parameters to

the experimental data. We then performed GLM regressions: dV, C, and dC on choice (binomial)

and on RT (linear). (Note: we coded the data such that option 1 always had the higher value.) A

preliminary inspection of the results suggests that Model 4 is the only model that accounts for all

of the qualitative benchmarks of the certainty-RT correlations, in particular the decrease in RT

with both average certainty and certainty difference. As expected, Model 2 makes the wrong

qualitative prediction (higher RT with value certainty), while Model 3 and Model 5 fail to account

for the dependency of RT on either certainty difference (dC) or average certainty (C), respectively.

Figure 4: Qualitative predictions of the effects of value difference, value certainty, and

certainty difference on choice consistency and RT (shown for models 2-5; top row shows

experimental data). (Bar heights represent mean GLM beta weights based on 10^8 simulated

trials for each model.)

Below, we present the quantitative results of the model-fit comparison. The results are

summarized in Figure 5.

Figure 5: Model comparison results. For the five models we examined, we show here the

model attributions (top row) for each of the studies we examined, at the subject level; each cell

represents the probability that the model (column) best represents the behavior of the subject

(row). We also show the exceedance probability for each model being the best in each study

(bottom row).

For Dataset 1, Model 1 dominated Model 2, with an exceedance probability of 1 and an

estimated model frequency (across the participant population) of 0.972. Model 1 dominated

Model 3, with an exceedance probability of 1 and an estimated model frequency of 0.990. Model

4 dominated Model 1, with an exceedance probability of 1 and an estimated model frequency of

0.779. Model 5 dominated Model 1, with an exceedance probability of 1 and an estimated model

frequency of 0.757. When comparing all models simultaneously, Model 4 dominated, with an

exceedance probability of 0.956 and an estimated model frequency of 0.541. Models 1, 2, 3, and

5 had estimated model frequencies of 0.127, 0.004, 0.004, and 0.324, respectively. Because

Models 4 and 5 each performed better than Model 1, we ran the comparison again including

Model 6, which was a combination of Models 4 and 5. Model 4 again dominated, with Model 6

receiving no support.

For Dataset 2, Model 1 dominated Model 2, with an exceedance probability of 1 and an

estimated model frequency of 0.984. Model 1 dominated Model 3, with an exceedance

probability of 1 and an estimated model frequency of 0.983. Model 4 dominated Model 1, with

an exceedance probability of 1 and an estimated model frequency of 0.897. Model 5 dominated

Model 1, with an exceedance probability of 1 and an estimated model frequency of 0.889. When

comparing all models simultaneously, Model 4 outperformed the others, with an exceedance

probability of 0.742 and an estimated model frequency of 0.546. Models 1, 2, 3, and 5 had

estimated model frequencies of 0.127, 0.004, 0.004, and 0.324, respectively. Because Models 4

and 5 each performed better than Model 1, we ran the comparison again including Model 6,

which was a combination of Models 4 and 5. Model 4 again outperformed the others, with Model

6 receiving no support.

For Dataset 3, Model 1 dominated Model 2, with an exceedance probability of 1 and an

estimated model frequency of 0.987. Model 1 dominated Model 3, with an exceedance

probability of 1 and an estimated model frequency of 0.985. Model 4 dominated Model 1, with

an exceedance probability of 1 and an estimated model frequency of 0.860. Model 5 slightly

outperformed Model 1, with an exceedance probability of 0.561 and an estimated model

frequency of 0.511. When comparing all models simultaneously, Model 4 dominated, with an

exceedance probability of 1 and an estimated model frequency of 0.865. Models 1, 2, 3, and 5

had estimated model frequencies of 0.122, 0.004, 0.004, and 0.004, respectively. Because

Models 4 and 5 each performed better than Model 1, we ran the comparison again including

Model 6, which was a combination of Models 4 and 5. Model 4 again dominated, with Model 6

receiving no support.

For Dataset 4, Model 1 dominated Model 2, with an exceedance probability of 1 and an

estimated model frequency of 0.955. Model 1 dominated Model 3, with an exceedance

probability of 1 and an estimated model frequency 0.982. Model 4 dominated Model 1, with an

exceedance probability of 1 and an estimated model frequency of 0.986. Model 5 dominated

Model 1, with an exceedance probability of 1 and an estimated model frequency of 0.986. When

comparing all models simultaneously, Model 4 dominated, with an exceedance probability of 1

and an estimated model frequency of 0.808. Models 1, 2, 3, and 5 had estimated model

frequencies of 0.005, 0.005, 0.005, and 0.175, respectively. Because Models 4 and 5 each

performed better than Model 1, we ran the comparison again including Model 6, which was a

combination of Models 4 and 5. Model 4 again dominated, with Model 6 receiving no support.

DISCUSSION

The aim of this study was to examine a number of variants of drift-diffusion model for preferential choice, and to probe them in their ability to account for benchmark data on the dependency of choice and RT on value uncertainty. As illustrated in Figure 2, the experimental data that we examined show that value certainty has a clear impact on both choice and RT (thus extending beyond the default DDM without option-specific noise), and also provides strong constraints on the way one can introduce option-specific noise into the model. As we have shown, the simplest DDM extension, in which the noise increases with value uncertainty, produces the wrong qualitative prediction: RT increases with certainty (certainty reduces the noise in the system, which slows down RT; see Figure 3, right panel).

We have examined and tested three additional DDM variants. The first two (Models 3-4) were based on signal-to-noise principles, while the last one (Model 5) included an independent and additive diffusion process based on certainty. While each of these models was able to account for some of the relationships in the data, only Model 4 accounted for all of them. In this model, the drift rate of the diffusion process is not simply the fluctuating difference in the values of the options (Tajima, Drugowitsch, & Pouget, 2016), but rather a difference between the ratios of the values and their corresponding value uncertainties. This mechanism has a normative flavor, as it penalizes values that are associated with uncertain alternatives. Some similar type of signal-to-noise models have also been supported by data in perceptual choice tasks. For example, de Gardelle and Summerfield (2011) examined choices in which an array of eight visual patches of variable color or shape are compared (in binary choice) to a reference (color or shape). By independently varying the set mean distance (in the relevant dimension) from the reference as

well as the set variance, they found that both independently affect choice accuracy and RT. In

particular, set variance (which is the analog of our value uncertainty) reduces choice accuracy

and increases RT. As shown by de Gardelle and Summerfield (2011), a signal-to-noise model can

account for this dependency. Indeed, the random dot motion task that is widely used alongside

the DDM in perceptual decision making studies provides a signal-to-noise ratio as input for the

drift rate (e.g., Gold & Shadlen, 2007). With this task, drift rate is typically determined by the

motion coherence, which is composed of the number of dots moving in the same direction

(signal) as well as the number of dots moving randomly (noise). Future work is needed to examine

the neural mechanism that extracts the drift from fluctuating values (sampled from memory or

prospective imagination; Bakkour et al, 2019; Poldrack et al, 2001; Schacter, Addis, & Buckner,

2007) and that reduces the drift of strongly fluctuating items. Future research is also needed to

examine if the effects of value uncertainty on choice correlate with risk-aversion at the level of

individual participants, and to integrate this type of model with dynamical attentional affects as

in the attentional drift-diffusion model (aDDM; Krajbich et al, 2010; Sepulveda et al, 2020).

While we have focused here on how value certainty affects choice and RT, the

experimental data also importantly show a marked and systematic effect of value certainty on

choice confidence. In particular, higher average value certainty (C) and certainty difference (dC)

both lead to higher choice confidence. This pattern raises a further challenge for most

accumulation-to-bound style choice models that aim to account for both RT and choice

confidence. For example, in the balance of evidence (BOE) type models (Vickers & Packer, 1982;

De Martino et al, 2013), confidence corresponds to the difference in the activation of two

accumulators that independently race to a decision boundary. If we were to naively introduce

option-specific noise in such models, they would predict, contrary to the data, that the

confidence becomes larger for options with more value uncertainty (as the noise increases the

BOE; see Lee & Daunizeau, 2020b). Similarly, if we were to model confidence using a DDM with

collapsing boundaries (e.g., Tajima et al, 2016), with confidence corresponding to the height of

the boundary at the time the choice is made, naively introducing option-specific noise would

once again provide us with a prediction opposite from what we see in the data. For uncertain

alternatives, there would be more noise in the evidence accumulation process, resulting in faster

choices and therefore higher boundaries, and thus higher confidence (in fact, this would be true

for any model that assumes that confidence decreases with RT; Kiani & Shadlen, 2009).

There are very few value-based choice studies that simultaneously examined value

certainty and choice confidence (but see Lee & Daunizeau, 2020a, 2020b; Lee & Coricelli, 2020;

De Martino et al, 2013). We have not modeled choice confidence here, as there are many

potential ways to do this, with substantial divergence among them (Vickers & Packer, 1982; Kiani

& Shadlen, 2009; Pleskac & Busemeyer, 2010; De Martino et al, 2013; Moran, Teodorescu, &

Usher, 2015; Calder-Travis, Bogacz, & Yeung, 2020; see Calder-Travis et al, 2020). Nevertheless,

all of these models strive to predict a strong negative correlation between RT and choice

confidence, as has been demonstrated in a plethora of experimental data. We note that in the

data we examined, the impact of value certainty on choice confidence was essentially the reverse

of its effect on RT (see Supplementary Material, Figure S1). While we did not explore this further,

it suggests that a signal-to-noise DDM can also capture the dependency of choice confidence on

value certainty. Future work is needed to determine how signal detection style DDM variants

might be extended towards an optimal unified account of choice, RT, and choice confidence.

BIBLIOGRAPHY

Bakkour, A., Palombo, D. J., Zylberberg, A., Kang, Y. H., Reid, A., Verfaellie, M., ... & Shohamy, D. (2019). The hippocampus supports deliberation during value-based decisions. elife, 8, e46080.

Basten, U., Biele, G., Heekeren, H. R., & Fiebach, C. J. (2010). How the brain integrates costs and benefits during decision making. Proceedings of the National Academy of Sciences, 107(50), 21767-21772.

Boldt, A., & Yeung, N. (2015). Shared neural markers of decision confidence and error detection. *Journal of Neuroscience*, *35*(8), 3478-3484.

Busemeyer, J. R., Gluth, S., Rieskamp, J., & Turner, B. M. (2019). Cognitive and neural bases of multi-attribute, multi-alternative, value-based decisions. Trends in cognitive sciences, 23(3), 251-263.

Calder-Travis, J., Bogacz, R., & Yeung, N. (2020). Bayesian confidence for drift diffusion observers in dynamic stimuli tasks. bioRxiv.

Calder-Travis, J., Charles, L., Bogacz, R., & Yeung, N. (2020). Bayesian confidence in optimal decisions. psyArXiv.

Daunizeau, J., Adam, V., & Rigoux, L. (2014). VBA: a probabilistic treatment of nonlinear models for neurobiological and behavioural data. *PLoS Comput Biol*, *10*(1), e1003441.

De Gardelle, V., & Summerfield, C. (2011). Robust averaging during perceptual judgment. Proceedings of the National Academy of Sciences, 108(32), 13341-13346.

De Martino, B., Fleming, S. M., Garrett, N., & Dolan, R. J. (2013). Confidence in value-based choice. *Nature neuroscience*, *16*(1), 105-110.

Gold, J. I., & Shadlen, M. N. (2007). The neural basis of decision making. *Annual review of neuroscience*, *30*.

Gwinn, R., & Krajbich, I. (2020). Attitudes and attention. *Journal of Experimental Social*
Psychology, *86*, 103892.

Kiani, R., & Shadlen, M. N. (2009). Representation of confidence associated with a decision by
neurons in the parietal cortex. science, 324(5928), 759-764.

Krajbich, I., Armel, C., & Rangel, A. (2010). Visual fixations and the computation and
comparison of value in simple choice. *Nature neuroscience*, *13*(10), 1292-1298.

Lebreton, M., Abitbol, R., Daunizeau, J., & Pessiglione, M. (2015). Automatic integration of
confidence in the brain valuation signal. *Nature neuroscience*, *18*(8), 1159-1167.

Lee, D., & Coricelli, G. (2020). An empirical test of the role of value certainty in decision
making. *bioRxiv,* 2020.06.16.155234.

Lee, D., & Daunizeau, J. (2020a). Choosing what we like vs liking what we choose: How choice-
induced preference change might actually be instrumental to decision-making. *PloS*
one, *15*(5), e0231081.

Lee, D., & Daunizeau, J. (2020b). Trading Mental Effort for Confidence: The Metacognitive
Control of Value-Based Decision-Making. *bioRxiv*, 837054.

Milosavljevic, M., Malmaud, J., Huth, A., Koch, C., & Rangel, A. (2010). The drift diffusion model
can account for the accuracy and reaction time of value-based choices under high and low
time pressure. Judgment and Decision Making, 5(6), 437.

Moran, R., Teodorescu, A. R., & Usher, M. (2015). Post choice information integration as a
causal determinant of confidence: Novel data and a computational account. Cognitive
psychology, 78, 99-147.

Philiastides, M. G., & Ratcliff, R. (2013). Influence of branding on preference-based decision
making. Psychological science, 24(7), 1208-1215.

Pleskac, T. J., & Busemeyer, J. R. (2010). Two-stage dynamic signal detection: a theory of
choice, decision time, and confidence. Psychological review, 117(3), 864.

Polania, R., Woodford, M., & Ruff, C. C. (2019). Efficient coding of subjective value. *Nature*
neuroscience, *22*(1), 134-142.

Polanía, R., Moisa, M., Opitz, A., Grueschow, M., & Ruff, C. C. (2015). The precision of value-
based choices depends causally on fronto-parietal phase coupling. Nature
communications, 6(1), 1-10.

Poldrack, R. A., Clark, J., Paré-Blagoev, E. A., Shohamy, D., Moyano, J. C., Myers, C., & Gluck,
M. A. (2001). Interactive memory systems in the human brain. Nature, 414(6863), 546-550.

Ratcliff, R., Smith, P. L., Brown, S. D., & McKoon, G. (2016). Diffusion decision model: Current
issues and history. *Trends in cognitive sciences*, *20*(4), 260-281.

Ratcliff, R., Voskuilen, C., & Teodorescu, A. (2018). Modeling 2-alternative forced-choice tasks:
Accounting for both magnitude and difference effects. Cognitive psychology, 103, 1-22.

Schacter, D. L., Addis, D. R., & Buckner, R. L. (2007). Remembering the past to imagine the
future: the prospective brain. Nature reviews neuroscience, 8(9), 657-661.

Sepulveda, P., Usher, M., Davies, N., Benson, A., Ortoleva, P., & De Martino, B. (2020). Visual
attention modulates the integration of goal-relevant evidence and not value. bioRxiv.

Tajima, S., Drugowitsch, J., & Pouget, A. (2016). Optimal policy for value-based decision-
making. *Nature communications*, *7*(1), 1-12.

Teodorescu, A. R., Moran, R., & Usher, M. (2016). Absolutely relative or relatively absolute:
violations of value invariance in human decision making. Psychonomic bulletin & review,
23(1), 22-38.

Vickers, D., & Packer, J. (1982). Effects of alternating set for speed or accuracy on response
time, accuracy and confidence in a unidimensional discrimination task. Acta psychologica,
50(2), 179-197.

Supplementary Materials

Choice Confidence

In this study, we chose not to include choice confidence in our model predictions, as there

is not currently an agreed upon standard for doing so. Nevertheless, we did briefly examine this

variable in those datasets that contained it (Studies 1-3; Lee & Daunizeau, 2020a, 2020b; Lee &

Coricelli, 2020). In general, choice confidence exhibited patterns qualitatively opposite to those

exhibited by RT. Specifically, regression beta weights for dV, C, and dC were of similar magnitude

as those for RT, but were all positive (whereas for RT, they were all negative). (see Figure S1)

Figure S1: Beta weights from a linear regression of dV, C, and dC on choice confidence (left) and

log(RT) (right; n=124; bar heights represent population means, error bars represent s.e.m.).

554