1 **Value Certainty in Drift-Diffusion Models of Preferential Choice**

2 Douglas Lee[1]   Marius Usher[2]

3 [1] Institute of Cognitive Sciences and Technologies, National Research Council of Italy

4 [2] Tel Aviv University, School of Psychological Sciences & Sagol School of Neuroscience

11 E-mail for correspondence:

12 DouglasGLee@gmail.com

Abstract

The *drift-diffusion model* (DDM) is widely used and broadly accepted for its ability to account for binary choices (in both the perceptual and preferential domains) and for their response times (RT), as a function of the stimulus or the option values. The DDM is built on an evidence accumulation to bound concept, where, in the value domain, a decision maker repeatedly samples the mental representations of the values of the options until satisfied that there is enough evidence in favor of one option over the other. As the signals that drive the evidence are derived from value estimates that are not known with certainty, repeated sequential samples are necessary to average out noise. The classic DDM does not allow for different options to have different levels of variability in their value representations. However, recent studies have shown that decision makers often report levels of certainty regarding value estimates that vary across choice options. There is therefore a need to extend the DDM to include an option-specific value certainty component. We present several such DDM extensions and validate them against empirical data from four previous studies. The data support best a DDM version in which the drift of the accumulation is based on a sort of signal-to-noise ratio of value for each option (rather than a mere accumulation of samples from the corresponding value distributions). This DDM variant accounts for the positive impact of value certainty on choice consistency and for the negative impact of value certainty on decision time, both of which are present in the empirical data.

The *drift-diffusion model* (DDM; Bogacz et al, 2006; Gold & Shadlen, 2007; Ratcliff & Rouder, 1998) is ubiquitous in the contemporary literature on decision making, including research spanning the fields of psychology, neuroscience, economics, and consumer behavior. The DDM is a parsimonious mechanism that explains normative decisions by averaging out noise in value comparisons (e.g., stochastic fluctuations in value signals that may result from variability in the firing of neural populations, from variability in attention or in memory retrieval of the values associated with the choice alternatives, or from variability intrinsic in the environment when the alternatives are characterized by variable/stochastic values) and implementing an optimal stopping rule (optimizing response time (RT) for a specified accuracy; Wald, 1948; Gold & Shadlen, 2007; Bogacz et al, 2006). This model accounts well for the dependency of the RT distribution on the values (or stimulus strength) of the choice options (Ratcliff & Rouder, 1998), and for the *speed-accuracy* tradeoff — the often-observed phenomenon that decision makers are able to improve their choice accuracy by taking longer to decide (on average; Wickelgren, 1977) — which can be accounted for by an expansion of the DDM response boundaries.

While initially used in the domain of perceptual (Ratcliff & Rouder, 1998) or memory-based decisions (Ratcliff, 1978), the DDM core principles of sequential sampling and integration to boundary have since become a central component of models of preference-based decisions (Busemeyer & Townsend, 1993; Busemeyer & Diederich, 2002; Busemeyer et al, 2019; Fudenberg, Strack, & Strzalecki, 2018; Tajima, Drugowitsch, & Pouget, 2016; Polania et al, 2015; Philiastides & Ratcliff, 2013; Milosavljevic et al, 2010; Krajbich, Armel, & Rangel, 2010; Basten et al, 2010; Roe, Busemeyer & Townsend, 2001; Turner, Schley, Muller, & Tsetsos (2018); Usher & McClelland, 2004). For example, in the *Decision Field Theory* (DFT) model (Busemeyer & Townsend, 1993; Busemeyer & Diederich, 2002; Busemeyer et al, 2019; Roe et al., 2001) — one of the first models that applied sequential sampling principles to preferential choice — it is assumed that the variability in the value integration corresponds to fluctuations of attention between the relevant attributes that characterize the alternatives. In a more recent application of the DDM to preferential choice between food items, the drift rate is assumed to be modulated by visual attention across the alternatives (thus this model was labeled the *attentional drift-diffusion model* or aDDM), which explains why people tend to choose more often the alternatives that they look longer at (Krajbich, Armel, & Rangel, 2010). In two more recent studies (Tajima et al; 2016; Fudenberg, Strack, & Strzalecki, 2018), the authors carried out normative analyses of the optimal

decision policy in preferential choice. Both of these studies were based on principles of Bayesian updating of value estimates and optimal stopping policies, given the cost of accumulating evidence. Tajima et al (2016) converged on the conclusion that "similar to perceptual decisions, drift diffusion models implement the optimal strategy for value-based decisions." An important assumption for the derivation of this optimal policy is that "the value of each option is represented by a probability distribution whose mean is the true (subjective) value of the option, and whose variance corresponds to sampling noise or uncertainty about the true value." This noise could be interpreted as imprecision in the value representations themselves. Similarly, the model of Fudenberg et al (2018) includes a common variance term meant to represent the uncertainty about the option value estimates (prior to accumulating evidence, which is reduced as more evidence is accumulated). Thus, the momentary signal about the relative value of the options (the so-called evidence for one option over the other) fluctuates randomly around a fixed value (the so-called drift rate, which corresponds to the difference in the means of the two value distributions). In order to average out this noise, the values of the alternatives are repeatedly sampled (possibly from memory; Bakkour et al, 2019) and accumulated over time until a sufficient amount of evidence has been acquired to allow for a choice to be made. Accordingly, the DDM includes response boundaries for each option (typically symmetric) that trigger a choice once reached by the evidence accumulator. Thus, the fundamental components of the drift-diffusion process are the drift rate (proportional to the difference in the option values), the diffusion coefficient (the degree of stochasticity of the system), and the choice boundaries (the threshold of minimum required evidence for a given level of caution, which controls the speed-accuracy tradeoff; see Figure 1).[1]

---

[1] According to Tajima et al (2016) and Fudenberg et al (2018), the optimal policy requires a collapsing rather than fixed boundary. The collapse of the boundary results, however, from auxiliary assumptions about the cost of deliberation time to the subject. Since these assumptions can be debated, and since there is an intensive debate on whether experimental data best support DDM with fixed or collapsing boundaries (Evans, Trueblood, & Holmes, 2020; Hawkins et al, 2015; Palestro et al, 2018), we consider both cases in this study. We will start with the simple fixed boundary DDM, which is the most prominent version in the literature, and we will then show that our conclusions are unchanged for the collapsing boundary variant (see Figure S1 in the Supplementary Material).

Figure 1: *An illustration of the classic* DDM. *Evidence accumulates across time, following a fixed drift trajectory (black dashed arrow, whose slope corresponds to the value difference of the choice options multiplied by a scalar coefficient), corrupted by white processing or sampling noise. Here, the accumulated evidence reaches the upper boundary after 3.9 seconds, and a choice for option 1 is recorded.*

The classic DDM implicitly assumes that processing (or sampling) noise is independent of the identity of the options contained in a particular choice set. That is to say, the sampling noise in the DDM is not option-specific (but see Ratcliff, Voskuilen, & Teodorescu, 2018; Teodorescu, Moran, & Usher, 2016 for DDM variants in which the noise increases with task difficulty; and Ratcliff & McKoon, 2018, for an application of option-specific variability in decisions about numerosity). Most of the DDM applications to preference-based decisions have assumed no option-specific noise. However, the brain is known to encode not only the subjective value of options, but also the subjective certainty about the value (Lebreton et al, 2015). Efficient coding principles also suggest that the values of certain options will be represented with greater precision when those options are encountered more frequently (Heng, Woodford, & Polania, 2020). It is thus reasonable to suggest that the representations of value that the brain uses to inform the decision process vary (Tajima et al, 2016), and that the degree of imprecision (or uncertainty) is not the same for all choice options. Indeed, it has been shown that decision makers hold highly variable beliefs about the certainty of their value estimates for different options, and that those beliefs are relatively stable within individuals (Lee & Coricelli, 2020; Lee & Daunizeau, 2020, 2021; Gwinn

& Krajbich, 2020; Polania, Woodford, & Ruff, 2019). It has further been shown that the variability

in value (un)certainty has a clear and systematic impact on both choice and RT (Lee & Coricelli,

2020; Lee & Daunizeau, 2020, 2021). Specifically, the judged value certainty, C, correlates

positively with choice consistency (the equivalent of choice accuracy for preferential choice) and

negatively with RT[2]. Furthermore, the difference in certainty (in favor of the higher-valued

alternative), dC, is associated with higher choice consistency and faster RT. This provides a

qualitative benchmark that any DDM variant that includes option-specific uncertainty should be

able to account for. To summarize these previous findings, we pooled the data from all four studies

(n=191) and conducted mixed model regression analyses: logistic for consistency, linear for

log(RT). The random effects variables were the different studies and individual participants, and

the fixed effects variables were dV, C, and dC (see Figure 2). We focused here on the effects of

dV, C, and dC — the former is expected in any DDM, while the latter two are challenges that we

aim to account for. The actual data also shows an effect of the average value (V = V1 + V2) on

RT (see Figure S4 in the Supplementary Material). Such dependence is usually not obtained in

DDM type models, and therefore we chose not to focus on it here. However, as shown in the

Supplementary Material, the DDM extension that we propose does account for this effect.

---

[2] As shown in Figure 2, we regressed the various measures of certainty on log(RT) rather than on RT. This is because whereas the RT variable is not normally distributed (it has a relatively long right tail), the transformed variable of log(RT) is normally distributed and is more adequate for statistical testing (see Glockner & Betch, 2008). Our conclusions hold if we use RT (without the log-transformation).

Figure 2: *Previous results (pooled across four studies, n=191) demonstrate the relationships between value difference (dV = V1 - V2), value certainty (C = C1 + C2), and certainty difference (dC = C1 - C2) with choice consistency and log(RT). We define V1 (V2) as the higher (lower) value rating, and C1 (C2) as the certainty report for the option with the higher (lower) value rating. The **upper** panels show the fixed effects coefficients from mixed effect logistic regression on choice consistency (upper left) and linear regression on log(RT) (upper right). Error bars represent standard errors). Significance stars represent *** p < .001, ** p < .01, * p < .05 (one-sided t-tests). The **lower** panels show the isolated relationships between log(RT) and C (left) and between log(RT) and dC (right). All variables were z-scored within participant prior to binning. Note: the distribution of C was highly skewed, so we show the data binned by quantiles. (Lee & Daunizeau, 2020, 2021; Lee & Coricelli, 2020; Gwinn & Krajbich, 2020)*

A natural way in which one could introduce option-specific noise in the preferential DDM, consistent with the assumptions of the optimal policy model (Tajima et al, 2016; Fudenberg et al, 2018), would be to assume that the diffusion noise parameter increases with value uncertainty. This is because alternatives whose values are less certain will have greater variance in their representations, from which the evidence samples are drawn. However, on its own, this would lead to the wrong qualitative predictions, with regards to the RT x certainty benchmark. Specifically, under such a model, diffusion noise would decrease as certainty increases, resulting in longer RT. Thus, unlike what we see in experimental data, such a model would predict that people would speed up when they are less certain of the options' values (all else equal). Moreover, this prediction is not specific to the classic DDM, but applies to the broader class of evidence accumulation-to-bound models (e.g., independent accumulators (Vickers, 1970; Brown & Heathcote, 2006); *leaky competing accumulator* (LCA; Usher & McClelland, 2001)), which also predict that higher noise in the system will result in *faster* responses, in direct contrast to the empirical data[3]. An alternative way to include option-specific noise in the DDM could be to assume that (only) the response boundary increases with value uncertainty (keeping the diffusion noise independent). Such a model could account for the observed negative correlation between RT and certainty sum, but not certainty difference (between options; see Figure 2, bottom right panel). Furthermore, such a model would also predict that choice consistency (i.e., in favor of the higher rated option) decreases with certainty (resulting from a decrease in the response boundary, all else equal), which contradicts the empirical data. One possibility, which we will consider here (inspired by the DFT model) is that the magnitude of value uncertainty has a simultaneous effect on both the diffusion noise and the response boundary (both increasing with uncertainty; Busemeyer & Townsend, 1993; see Table 4 in particular). In principle, in such a case, the impact of value certainty on choice accuracy would thus involve two opposing factors (noise and boundary), which may allow a way to account for some aspects of the data (though note that this is not likely to provide an account for the dependency of accuracy and RT on dC). Another hypothesis is that the option-specific

---

[3] In the *linear ballistic accumulator* (LBA) model (Brown & Heathcote, 2006), the within-trial noise is replaced by between-trial noise in the starting point and drift rate (in the DDM, such between-trial variability is sometimes introduced in addition to the within-trial noise (Ractcliff & McKoon, 2008). Increasing those components with option-uncertainty, however, would have a similar effect as increasing the within-trial diffusion noise in the DDM (i.e., a speedup effect and reduced accuracy; see the Supplementary Material).

value certainty affects the DDM process through the drift rate. We provide preliminary support for
this conclusion in the Supplementary Material, using median splits on certainty and comparing the
fitted model drift rate parameters.

The aim of this paper is to examine a number of DDM variants for preferential choice,
which could account for the impact of value certainty on choice and RT in behavioral data. In
particular, we will first present a variety of extensions of the classic DDM, each of which
incorporates the concept of option-specific value certainty in a unique and realistic way. We start
with the classic DDM (Model 1), which has no option-specific noise, and progress to models that
include option-specific certainty in the sampling noise or in the response boundary (Models 2-3;
although we know that these models are less likely to succeed, we believe that it is instructive to
formally test them and to examine their potential failures). We then examine a model (2+3) in
which value certainty modulates both the sampling noise and response boundary, as well as models
(4-5) based on signal-to-noise ratio concepts in the formulation of the drift rate. We fit each of
these models to experimental data from four empirical datasets. Finally, we quantitatively compare
the performance of each model across the different datasets and use the results to suggest the best
recommended approach for future studies to incorporate option-specific value certainty in models
derived from the DDM. To anticipate our results, we find that a signal-to-noise DDM variant
provides the best fit to the data and accounts well for all qualitative benchmarks.

METHODS

Computational models

In each of the models described below, we consider decisions between two alternatives,
with values, V1 and V2, (with V1>V2), and with uncertainties, $\sigma 1$, $\sigma 2$, respectively. Evidence of
these values is integrated over deliberation time, subject to noise. The evidence accumulator for
each decision is initialized at a neutral starting point (i.e., zero evidence or default bias in favor of
either option), and evolves until reaching one of two symmetric boundaries (see Figure 1 above).
For each decision, the output of the model is: the choice (ch = {0, 1}), determined by which
boundary was reached; RT, determined by the number of integration time steps elapsed before that
boundary was reached.

Model 1: Classic DDM

As a baseline default model for comparison (without any option-specific certainty term),
we first consider the classic DDM. In this model, the equations that govern the evidence
accumulation process are:

$$E_{t+1} = E_t + d\Delta$$

$$\Delta \sim N(\mu_1 - \mu_2, \sigma^2)$$

where E represents the cumulative balance of evidence that one option is better than the other, t
represents deliberation time, Δ represents the incremental evidence in favor of one option over the
other, d is a sensitivity parameter, μ_i is the value estimate of option i, and σ^2 represents processing
noise in the evidence accumulation and/or comparator systems. In the classic DDM, choice
probability and expected response time can be analytically calculated using the following
equations (Alos-Ferrer, 2018):

$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{-2\theta d}{\sigma^2}(\mu_1 - \mu_2)\right)}}$$

$$E[RT] = \frac{\theta(2 * p(ch = 1) - 1)}{d(\mu_1 - \mu_2)}$$

where θ is the height of the evidence accumulation boundary where a decision is triggered (shown
here as the upper boundary, for a choice of option 1), p(ch=1) is the probability that the upper
boundary will be reached (rather than the lower boundary), and RT is the expected time at which
the accumulation process will end (by reaching one of the boundaries)[4]. Because this system of
equations is over-parameterized, one of the free parameters must be fixed for a practical
application of the equations. In this work, for simplicity, we will fix to a value of 1 (when fitting
the models) the boundary parameter θ. Choice probability and RT will thus be functions of the
drift rate coefficient *d*, the diffusion (noise) coefficient σ^2, and the height of the response boundary

---

[4] In the standard version of the DDM, the RT distribution for correct and incorrect responses is identical. In a more complex version, additional variability parameters are introduced that allow to account for asymmetries between the RT distributions of correct and incorrect responses (see Ratcliff & McKoon, 2008 for a review). We only consider the standard DDM without the variability parameters, as those cannot change the impact of value certainty on accuracy and RT illustrated in Figures 2 and 3.

θ (see Figure 3). As expected, accuracy increases and RT decreases with the drift rate. The noise,
however, reduces both accuracy and RT, and the boundary increases both accuracy and RT.

Figure 3: *Dependency of choice consistency (upper row) and of RT (bottom row) on DDM*
parameters: drift rate (left/blue), diffusion noise (middle/red) and response boundary
(right/green). With all other parameters fixed: an increase in drift rate leads to an increase in the
probability of choosing the best option from 0.5 (random guess) to 1, and a decrease in RT; an
increase in processing noise leads to a decrease in the probability of choosing the best option, and
a parallel decrease in RT; an increase in response boundary leads to an increase in the probability
of choosing the best option, and an increase in RT.

Model 2: Certainty-Adjusted Diffusion Noise

One simple potential solution to incorporate option-specific uncertainty into the DDM
would be to model the process as:

$$E_{t+1} = E_t + d$$

$$\Delta \sim N\left(\mu_1 - \mu_2, \sigma^2 + \gamma(\sigma_1^2 + \sigma_2^2)\right)$$

This corresponds to the classic DDM equation capturing the evolution of the accumulated
evidence across time, but with σ_i^2 representing the uncertainty about the value estimate of option

i, and γ serving as a sensitivity parameter that controls the impact of value certainty on the level
of diffusion noise. The only difference between this formulation and the classic one is that here
the variance of Δ is specific to the options in the current choice set, whereas in the classic DDM it
is fixed across choices (for an individual decision maker). A direct result of this reformulation is
that choices between options with greater value uncertainty (lower C) will be more stochastic and
take less time (on average), as can be seen by examining the (revised) DDM equations for choice
probability and expected response time (see also Figure 3, center panels):

233
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{-2\theta d(\mu_1 - \mu_2)}{\sigma^2 + \gamma\left(\sigma_1^2 + \sigma_2^2\right)}\right)}}$$

234
$$E[RT] = \frac{\theta(2 * p(ch = 1) - 1)}{d(\mu_1 - \mu_2)}$$

While the predicted relationship between certainty and choice consistency would match empirical
observations, the relationship between certainty and RT would be inconsistent with the empirical
data.

Model 3: Certainty-Adjusted Response Boundary

Another potential solution could be to allow the magnitude of the response boundary to
vary as a function of option-specific (or trial-specific) value certainty. The evidence sampling
process would be identical to that of the classic DDM:

242
$$E_{t+1} = E_t + d\Delta$$

243
$$\Delta \sim N(\mu_1 - \mu_2, \sigma^2)$$

244 Under this model, the height of the boundary would increase as the value certainty of the
245 pair of options decreases, on a trial-by-trial basis. Choice probability and mean RT would thus be
246 calculated using the following equations:

247
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{-2\theta\left(1 + \gamma(\sigma_1^2 + \sigma_2^2)\right)d}{\sigma^2}(\mu_1 - \mu_2)\right)}}$$

248
$$E[RT] = \frac{\theta\left(1 + \gamma(\sigma_1^2 + \sigma_2^2)\right)(2 * p(ch = 1) - 1)}{d(\mu_1 - \mu_2)}$$

where γ is a sensitivity parameter that controls the impact of certainty on the magnitude of the

response boundary. As can be seen in the equations, increasing value uncertainty (all else equal)

will result in higher choice consistency and higher RT (see Figure 3, right panels). While the

predicted relationship between certainty and RT would match empirical observations, the

relationship between certainty and choice consistency would be inconsistent with the empirical

data.

Model 2+3: Certainty-Adjusted Diffusion Noise and Response Boundary

Decision Field Theory (DFT; Townsend & Busemeyer, 1993; see Table 4) suggests that

value certainty simultaneously modulates the diffusion noise and the response boundary. We

therefore considered such a formulation in our set of models. The idea is that while uncertainty is

expected to increase the sampling noise (see also Tajima et al, 2016), participants are likely to

compensate for the reduced certainty by increasing the *caution parameter* – the response boundary.

In this hybrid model, choice probability and mean RT would thus be calculated using the following

equations:

$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{-2\theta\left(1+\lambda(\sigma_1^2+\sigma_2^2)\right)d(\mu_1-\mu_2)}{\sigma^2+\gamma(\sigma_1^2+\sigma_2^2)}\right)}}$$

$$E[RT] = \frac{\theta\left(1 + \lambda(\sigma_1^2 + \sigma_2^2)\right)(2 * p(ch = 1) - 1)}{d(\mu_1 - \mu_2)}$$

265 Here, the boundary varies as a function of value certainty independently of the diffusion noise (as

266 thus making Model 2+3 a direct hybrid of Models 2 and 3) [5].

---

[5] In the DFT model, the diffusion noise has an additive component that that originates from the correlation of the value estimates of the options. This was relevant for the choice task the model was applied to, in which values estimates could be correlated since they depended on a common event. As this is not likely to be the case for choices between arbitrary pairs of food items, we assume that there is no correlation between the value fluctuations and thus that the diffusion noise simply sums with the variance of the two options ($\sigma_1^2 + \sigma_2^2$). Note that in this model we allowed different sensitivity parameters, γ and λ, for the dependency of the diffusion noise and of the boundary on value uncertainty; we also considered a less flexible alternative, consistent with the DFT formulation. Here, the response boundary θ is an increasing function of the square root of the diffusion noise, which itself is an increasing function of value uncertainty $\sigma_1^2 + \sigma_2^2$. However, this model was dominated by the model we present above during model comparison, so we chose not to include it in our analyses.

Model 4: Certainty-Adjusted Drift Rate (value difference)

One possible way in which the concept of option-specific value (un)certainty could be

incorporated into the DDM would be through a signal-to-noise dependency in the drift rate. The

drift rate in the DDM symbolizes the accumulation of evidence for one option over the other, equal

to the value estimate of one option minus the value estimate of the other option (scaled by a fixed

term). The accumulator variable is referred to as "evidence" because the probability distributions

controlling it (or the neural activity driving it) are thought to provide reliable signal that will

accurately inform the decision. If the value representations of different options can have different

levels of uncertainty, it stands to reason that the reliability of the "evidence" that these signals

provide about the correct decision will also be different. As such, evidence drawn from a more

reliable source (i.e., one with higher certainty) should be weighted more heavily. Under this

framework, the equation governing the DDM process would be:

279
$$E_{t+1} = E_t + d\Delta$$

280
$$\Delta \sim N\left(\frac{\mu_1 - \mu_2}{\sqrt{\sigma_1^2 + \sigma_2^2}}, \sigma^2\right)$$

where σ^2 (without a subscript) is the noise in the system unrelated to specific choice options. The

only difference between this formulation and the classic one is that here the mean of the option

value difference is divided by its standard deviation ($\sqrt{\sigma_1^2 + \sigma_2^2}$). A direct result of this

reformulation is that choices between options with greater value uncertainty will be more

stochastic and also take *more* time (on average), as can be seen by examining the (revised) DDM

equations for choice probability and expected response time:

287
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{-2\theta d(\mu_1 - \mu_2)}{\sigma^2 \sqrt{\sigma_1^2 + \sigma_2^2}}\right)}}$$

288
$$E[RT] = \frac{\theta(2 * p(ch = 1) - 1)}{\frac{d(\mu_1 - \mu_2)}{\sqrt{\sigma_1^2 + \sigma_2^2}}}$$

289   Here the impact of option-specific uncertainty on RT is more complex. First, greater

290 uncertainty decreases RT through its effect on choice stochasticity (as before). Second, greater

291 uncertainty directly increases RT by diminishing the slope of the drift rate. The second effect
292 dominates.

Model 5: Certainty-Adjusted Drift Rate (option values)

An alternative variant of the DDM in which the drift rate is altered by the option-specific
value certainty could be one in which the evidence in favor of each option i is scaled by its own
precision term σ_i^2, as is the case, for example, in multi-sensory integration (Drugowitsch et al,
2014; Fetsch et al, 2012). The drift rate would thus become the accumulation of adjusted evidence
for one option over the other, equal to the precision-weighted value estimate of one option minus
the precision-weighted value estimate of the other option (scaled by a fixed term). Here, the
evidence drawn from a more reliable source (i.e., one with higher certainty) will be weighted more
heavily (as in Model 4), but prior to comparison between the alternative options. Note that here
the certainty weighting is truly specific to each option, whereas in Model 4 the certainty weighting
is specific to the pair of options. Under this framework, the equation governing the DDM process
would be:

305
$$E_{t+1} = E_t + d\Delta$$

306
$$\Delta \sim N\left(\frac{\mu_1}{\sigma_1} - \frac{\mu_2}{\sigma_2}, \sigma^2\right)$$

The only difference between this formulation and the classic one is that here the mean of
the option value difference is adjusted by the standard deviations of the individual choice options.
Because the evidence in favor of each option (prior to comparison) will be scaled by its own
specific (and importantly, potentially different) precision term, the impact on both choice
stochasticity and response time could go in either direction. This can be seen by examining the
(revised) DDM equations for choice probability and expected response time:

313
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{-2\theta d}{\sigma^2}\left(\frac{\mu_1}{\sigma_1} - \frac{\mu_2}{\sigma_2}\right)\right)}}$$

314
$$E[RT] = \frac{\theta(2 * p(ch = 1) - 1)}{d\left(\frac{\mu_1}{\sigma_1} - \frac{\mu_2}{\sigma_2}\right)}$$

Here the impact of option-specific uncertainty on both choice and RT is more complex than in the previous models. If the value signal for one option has both a larger mean and a smaller variance, relative to the other option, the effective drift rate will be higher than in the classic DDM (e.g., choices will be less stochastic and faster). On the other hand, if the option with the larger mean value also has a higher variance (due to its value uncertainty), the effective drift rate will be lower than in the classic DDM (e.g., choices will be more stochastic and slower).

Materials and Design

The data for this study was taken from 4 experiments reported in the literature that examined choices between pairs of food items, whose ratings (value and certainty) were measured for each participant before the pairs were presented for binary choice.

Dataset 1

The first dataset we examined was from Lee & Daunizeau (2020). In this study, participants made choices between various snack food options based on their personal subjective preferences. Value estimates for each option were provided in a separate rating task prior to the choice task. Participants used a slider scale to respond to the question, "Does this please you?" After each rating, participants used a separate slider scale to respond to the question, "Are you sure?" This provided a measure of value estimate certainty for each item. During the choice task, participants were presented with pairs of snack food images and asked, "What do you prefer?" After each choice, participants used a slider scale to respond to the question, "Are you sure about your choice?" to provide a subjective report of choice confidence. This dataset contained 51 subjects, each of whom were faced with 54 choice trials.

Dataset 2

The second dataset we examined was from Lee & Daunizeau (2021). In this study, participants made choices between various snack food options based on their personal subjective preferences. Value estimates for each option were provided in a separate rating task prior to the choice task. Participants used a slider scale to respond to the question, "How much do you like this item?" After each rating, participants used the same slider scale to respond to the question, "How certain are you about the item's value?" by indicating a zone in which they believed the value of the item surely fell. This provided a measure of value estimate certainty for each item. During the

choice task, participants were presented with pairs of snack food images and asked, "Which do you prefer?" After each choice, participants used a slider scale to respond to the question, "Are you sure about your choice?" to provide a subjective report of choice confidence. This dataset contained 32 subjects, each of whom were faced with 74 choice trials.

Dataset 3

The third dataset we examined was from Lee & Coricelli (2020). In this study, participants made choices between various snack food options based on their personal subjective preferences. Value estimates for each option were provided in a separate rating task prior to the choice task. Participants used a slider scale to respond to the question, "How pleased would you be to eat this?" After each rating, participants used a six-point descriptive scale to respond to the question, "How sure are you about that?" This provided a measure of value estimate certainty for each item. During the choice task, participants were presented with pairs of snack food images and asked, "Which would you prefer to eat?" After each choice, participants used a slider scale to respond to the question, "How sure are you about your choice?" to provide a subjective report of choice confidence. This dataset contained 47 subjects, each of whom were faced with 55 choice trials.

Dataset 4

The fourth dataset we examined was from Gwinn & Krajbich (2020). In this study, participants made choices between various snack food options based on their personal subjective preferences. Value estimates for each option were provided in a separate rating task prior to the choice task. Participants used a 10-point numerical scale to respond to the prompt, "Please indicate how much you want to eat this item." After each rating, participants used a seven-point numerical scale to respond to the prompt, "Please indicate how confident you are in your rating of this item." This provided a measure of value estimate certainty for each item. During the choice task, participants were presented with pairs of snack food images and instructed to choose the one that they preferred to eat. Choice confidence was not measured in this study. This dataset contained 36 subjects, each of whom were faced with 200 choice trials.

Computational Model-Fitting Procedure

We fit the experimental data to each of the models that we described above. We then performed Bayesian model comparison to determine which of the models (if any) performed better

than the others across the population of participants. For this model fitting and comparison exercise, we relied on the Variational Bayesian Analysis toolbox (VBA, available freely at https://mbb-team.github.io/VBA-toolbox/; Daunizeau, Adam, & Rigoux, 2014) with Matlab R2020a. Within participant and across trials, we entered the experimental variables {left option value, right option value, left option certainty, right option certainty} as input and {choice = 1 for left option, 0 for right option; RT} as output. We also provided the model-specific mappings from input to output as outlined in the analytical formulas above. The parameters to be fitted included all of the d, σ^2, γ, λ, and α terms described above in the model formulations. VBA requires prior estimates for the free parameters, for which we set the mean equal to one and the variance equal to e (to allow an arbitrarily-large step size during the gradient descent search algorithm, yet constrain the algorithm to a reasonable search space) for each parameter. VBA then recovers an approximation to both the posterior density on unknown variables and the model evidence (which is used for model comparison). We used the VBA_NLStateSpaceModel function to fit the data for each participant individually, followed by the VBA_groupBMC function to compare the results of the model fitting across models for the full group of participants.

The benefit of using VBA to fit the data to our models spawns from the nature of the preferential choice data itself, where the value estimate and certainty of each option necessarily varies from one trial to the next (i.e., no two choice trials are ever alike). Other DDM parameter estimation approaches that are popular in the literature do not optimally handle this kind of experimental design, because they lack the trial repetitions per condition (in this case, value and certainty ratings) necessary to provide empirical estimates of RT distributions (or moments) in each condition. The VBA toolbox is computationally efficient, as it relies on Variational Bayesian analysis under the Laplace approximation. This iterative algorithm provides a free-energy approximation for the model evidence, which represents a natural trade-off between model accuracy (goodness of fit, or log likelihood) and complexity (degrees of freedom, or KL divergence between flat priors and fitted parameter estimates; see Friston et al., 2007; Penny, 2012). Additionally, the algorithm provides an estimate of the posterior density over the model free parameters, starting with Gaussian priors. Individual log model evidence scores are then provided as input to the group-level random-effect Bayesian model selection (BMS) procedure. BMS provide an exceedance probability that measures how likely it is that a given model is more frequently implemented, relative to all other models under consideration, in the population from

which participants were drawn (Rigoux et al., 2014; Stephan et al., 2009). This approach to fitting

and comparing variants of DDM has already been successfully demonstrated in previous studies

(Lopez-Persem, Domenech, & Pessiglione, 2016; Feltgen & Daunizeau, 2021).

As an initial check to verify that our model-fitting procedure is suitable for this specific

analysis, we performed a test of model recovery. Specifically, we simulated synthetic data for each

participant in our experiments based on each of the models. We took as the model input the actual

rating data for each participant in Studies 1-4 (value estimate and certainty for each option) and

we simulated the set of choices and their RT for each participant, separately according to each of

our models and using the best-fit parameters for each participant for each model. Finally, we fit all

simulated data (per participant) to each of our models and performed the same formal model

comparison as with our real experimental data. The results of this procedure can be seen in Figure

4 as a model confusion matrix. This matrix shows, for each true generative model, the percentage

of simulated participants (under that model) that were attributed to each of the best fit models by

our model-fitting procedure. As shown in the matrix, overall confusion was low, as the procedure

attributed the true model as the best fit model for the majority of participants. Model 2 appears to

be the only model in our set of models that our model-fitting procedure was unable to accurately

recognize, wrongly classifying most participants simulated under Model 2 as having been

simulated under Model 1.

Figure 4: *Model recovery analysis. Each cell in the "confusion matrix" summarizes the percentage of simulated participants (under each true model) for which our model-fitting procedure attributed each of the (best fit) models. Note that for every model other than Model 2, our fitting procedure attributed the true model as the best fit model for the majority of simulated participants.*

RESULTS

Before we present the quantitative model comparison results, we show the qualitative predictions that each model (simulated under its optimal parameters) makes with respect to the effects of value difference, value certainty, and certainty difference on choice consistency and RT, and how this compares to the experimental data (see Figure 5). For the synthetic data, we first simulated 10^4 trials for each model, based on random input (option values and certainties) with the same range as in our experimental data (0 to 1; simulated data uniformly distributed). The model parameters we used were the best fit parameters to the experimental data, with mean and variance equal to the mean and variance across all participants in all studies (simulated data normally distributed). We then performed GLM regressions: dV, C, and dC on choice (binomial) and on RT (linear). (Note: we coded the data such that option 1 always had the higher value.) A preliminary inspection of the results suggests that Model 5 is the only model that accounts for all of the qualitative benchmarks of the certainty-RT correlations, in particular the decrease in RT with both certainty sum and certainty difference.

Figure 5: *Qualitative predictions of the effects of value difference (dV = V1 - V2), value certainty (C = C1 + C2), and certainty difference (dC = C1 - C2) on choice consistency and log(RT) (shown for the best fit parameters; bar heights represent mean GLM beta weights based on 10^4 simulated trials for each model; top row shows experimental data, n=191). All log(RT) data were z-scored prior to entering the GLM regressions and prior to binning for the quantile curve plots.*

448  We note that Models 2 and 3 cannot possibly account for the experimental data due to their

449 qualitative predictions (see Figure 5). Model 2 necessitates a negative relationship between value

450 certainty and choice consistency and a positive relationship between value certainty and RT, which

451 is the opposite of what we observe in all of the empirical datasets. Model 3 necessitates a negative

452 relationship between value certainty and choice consistency. Furthermore, as expected, neither the

453 "noise model" nor the "boundary model" received any support when quantitatively compared with

454 the classic DDM. However, it was not clear (from basic DDM principles) how a hybrid model in

455 which value certainty modulates both the diffusion noise (as in Model 2) and the response

456 boundary (as in Model 3) would perform. Previous work based on *Decision Field Theory* has

457 shown that such a hybrid model accounts for a variety of puzzling experimental phenomena (such

458 as the violation of independence axioms in risky choice; Townsend & Busemeyer, 1993). The

459 results of our preliminary simulations suggest that the hybrid Model 2+3 should perform better

460 than either Model 2 or Model 3 alone. However, it is important to note that this model cannot

461 account for the impact of certainty difference (dC) on either choice or RT.

462 *Quantitative Model Comparison*

463  The classic DDM, our Model 1, has been validated countless times for its ability to account

464 for two-alternative forced choice responses and mean response times. The other models described

465 above are new and have therefore never been tested with empirical data. We thus performed a

466 formal model comparison of our full set of models, with Model 1 serving as a benchmark. The

467 results are displayed in Figure 6, which shows the support that each model received within

468 participants (left panel) and across participants (right panel).

Figure 6: *Model comparison results. We show here the probability that each model best accounted for the data at the participant level (left panel), across the four studies we examined; each cell represents the probability that the model (column) best represents the behavior of the participant (row). The dashed lines serve to indicate which participants belonged to each of the four datasets. We also show the probability that each model best explains the data across the participant population (right panel), across all studies. The dashed line indicates chance level if all models were equally probable a priori.*

Across all four studies, Models 2 and 3 were each dominated by Model 1 (the classic DDM) as anticipated, as were Models 2+3 and Model 4. Model 5, in contrast, dominated Model 1. In recognition of the possibility that the impact of certainty on the drift rate might be non-linear, we considered a variation of our best-fitting model (Model 5), which we label Model 5*. Here, the only change was the inclusion of an exponent α in computation of the signal to noise ratio:

$$\Delta \sim N\left(\frac{\mu_1}{\sigma_1{}^\alpha} - \frac{\mu_2}{\sigma_2{}^\alpha}, \sigma^2\right)$$

In a formal comparison of Model 5 against Model 5*, Model 5* was the clear winner with an estimated model frequency (across participants) of 0.994. Most participants showed a concave impact of value certainty on evidence accumulation, with 70% of participants having $\alpha < 1$ (28% had $\alpha > 1$). Across participants, the median recovered α was 0.68 with a median average deviation of 0.46. This suggests that the marginal impact of value certainty is greater for options with relatively low certainty compared to those with relatively high certainty, for most participants. However, we note that these results should be interpreted with caution, because the parameter estimates of the α parameter were not very reliable. The parameters for each participant were estimated using very few trials, which presents a challenge to accurately fitting exponential parameters in addition to the linear ones. In particular, our parameter recovery exercise showed that the exponent parameters were often overestimated.

DISCUSSION

The aim of this study was to examine a number of extensions of the drift-diffusion model for preferential choice that include option-specific value certainty, and to probe them in their ability to account for benchmark data on the dependency of choice and RT on value certainty. As illustrated in Figure 2, the experimental data that we examined show that value certainty has a

clear and systematic impact on both choice and RT, thereby motivating an extension of the classic

DDM without option-specific noise and providing constraints on the way one can introduce option-

specific noise into the models. As we have shown, perhaps the simplest DDM extension, in which

(only) the evidence accumulation noise decreases with value certainty, produces the wrong

qualitative prediction: RT increases with certainty (as certainty reduces the stochasticity of the

system, which slows down RT; see Figure 3, center panels). Moreover, the problem with this

method of introducing option-specific value certainty in modeling value-based decisions is not

particular to the DDM, but also applies to the broader class of evidence accumulation-to-bound

models (e.g., LCA (Usher & McClelland, 2001) or independent accumulators (Vickers, 1970)), in

which noise decreases RT.

An alternative way to introduce option-specific value uncertainty in the DDM could be to

assume that the uncertainty affects the response boundary (only). Accordingly, decision makers

would compensate for their uncertainty by increasing the height of the boundary. While such a

model could account for the negative correlation between RT and certainty (C) shown in Figure 2,

it would not be able to account for the negative correlation between RT and certainty difference

(dC). Moreover, such a model would predict that choices become more stochastic as value

certainty increases (due to the lower boundary; see Figure 3, right panels), which is both

counterintuitive and in contrast to the experimental data. Those theoretical predictions are

supported by our model comparison (see Figures 5 and 6). We also considered a model in which

both diffusion noise and response boundary were simultaneously modulated by value certainty,

which can make the correct predictions in terms of C. However, this model does not predict any

impact of dC on either choice consistency or RT (see Figure 5). Thus, we believe that the way in

which value uncertainty affects the decision process is via its impact on the drift rate. (We provided

additional support for this conclusion in the Supplementary Material, using median splits on

certainty and comparing the fitted model parameters.)

In order to understand the way in which value certainty affects the drift rate, we examined

and tested two core DDM variants built on certainty-modulated drift rates. These models (4 and 5)

were based on signal-to-noise principles. While Model 4 was able to account for some of the

qualitative relationships in the data, only Model 5/5* accounted for all of them.[6] In this model, the

drift rate of the diffusion process is not simply the fluctuating difference in the values of the options

(Tajima et al, 2016), but rather a difference between the ratios of the mean values and their

corresponding value uncertainties. This mechanism has a normative flavor, as it penalizes values

that are associated with uncertain alternatives. Some similar types of signal-to-noise models have

also been supported by data in perceptual choice tasks. For example, de Gardelle and Summerfield

(2011) examined choices in which an array of eight visual patches of variable color or shape are

compared (in binary choice) to a reference (color or shape). By independently varying the set mean

distance (in the relevant dimension) from the reference as well as the set variance, they found that

both independently affect choice accuracy and RT. In particular, the set variance (which is the

analog of our value uncertainty) reduces choice accuracy and increases RT. As shown by de

Gardelle and Summerfield (2011), a signal-to-noise model can account for this dependency.

Furthermore, the random dot motion task that is widely used alongside the DDM in perceptual

decision-making studies provides a signal-to-noise ratio as input for the drift rate (e.g., Gold &

Shadlen, 2007); for this task, drift rate is typically determined by the motion coherence, which is

composed of the number of dots moving in the same direction (signal) as well as the number of

dots moving randomly (noise).

Future work is also needed to examine the neural mechanism that extracts the drift rate

from fluctuating values (sampled from memory or prospective imagination; Bakkour et al, 2019;

Poldrack et al, 2001; Schacter, Addis, & Buckner, 2007) and that reduces the drift rate of more

strongly fluctuating items. One interesting possibility is that the effective drift rate might be

modulated by the temporal congruency of the evidence in successive samples (Glickman, Moran,

& Usher, 2020); the higher the value certainty, the lower the variance of the value signals, thus

leading to a higher probability that successive samples will provide consistent choice evidence.

Future research is also needed to understand the nonlinearity indicated in this mechanism

(deviation of α exponent from 1), to examine if the effects of value uncertainty on choice correlate

with risk- or ambiguity-aversion at the level of individual participants, and to integrate this type

of model with dynamical attentional affects as in the attentional drift-diffusion model (aDDM;

---

[6] From this point forward, we will refer to the best fit model, Model 5*, as the signal-to-noise DDM, or *snDDM*. Note that Model 5 is a special case of Model 5*, where the exponent parameter α is equal to 1; Model 4 is a special case when dC = 0.

Krajbich et al, 2010; Sepulveda et al, 2020). Recent work has suggested that attention allocated
toward a particular choice option will serve to reduce the uncertainty about the value of that option
(Callaway, Rangel, & Griffiths, 2020; Jang, Sharma, & Drugowitsch, 2020). If true, this would
provide a link between this current work and the so-called gaze bias that has already been well
established (Krajbich, Armel, & Rangel, 2010; Sepulveda et al, 2020).

Response Time Distributions

One of the strengths of DDM-type models is their ability to account not only for accuracy
and mean RT, but also for how the full distribution of RT varies with task conditions (see Ratcliff
& McKoon, 2008 for a review). In this study, we did not fit RT distributions for a pragmatic reason:
the preference data (Studies 1-4) do not have enough repeated trials per condition (i.e., specific
combinations of value and certainty rating) needed to estimate RT distributions. Nevertheless, we
thought to examine predictions for how RT distributions vary with the predicted drift rate (which
is a function of the value estimates, V1 and V2, and their certainties, C1 and C2) in the signal-to-
noise model. As we report in the Supplementary Material (see Figure S2), we find the typical
rightward-skewed RT distributions in data simulated under the snDDM. Moreover, we find the
expected patterns with respect to trials with low versus high predicted drift rate. The experimental
data appear to confirm the model predictions (see Figure S2). This suggests that the snDDM can
account for patterns in the full RT distributions, in addition to all of the qualitative and quantitative
support that we reported above. However, future studies that include more trials per condition (i.e.,
each specific level of dV and C) will be needed to fully validate this.

Impact of Overall Value on Response Time

In this study, we focused on the impact of value difference, certainty sum, and certainty
difference on choice behavior. Yet there remains an additional element that we left out: value sum
(V = V1 + V2). A number of previous studies have exposed a clear and robust effect of overall
value (of the options being considered during a particular decision) on RT, where responses are
typically faster for higher-valued options (Smith & Krajbich, 2019; Polania et al, 2014; Hunt et al,
2012; Teodorescu, Moran & Usher, 2016; Sepulevada et al, 2020). We also found this effect in the
datasets we examined (see the Supplementary Material). As shown in Figure S4, the value sum
(V) has a clear effect on RT (though somewhat smaller than the effect of dV). People decide faster
(all else equal) on a choice between a pair of alternatives with ratings 0.9 and 0.8, respectively,

than on a choice between alternatives with ratings of 0.2 and 0.1, respectively. This absolute value

effect is naturally explained in race or competing accumulator models (Vickers, 1970) and even in

the LCA (Teodorescu, Moran & Usher, 2016), but is more challenging to explain within a diffusion

model, in which the preference formation is based on relative evidence and unrelated to absolute

overall evidence. Interestingly, although not predicted and therefore not discussed so far, it turns

out that the snDDM can account for the negative relationship between value sum and RT (see

Figure S4). The snDDM account of this effect is an emergent one (i.e., there was nothing in the

model that was explicitly dependent on V). We briefly comment on this in the Supplementary

Material, and leave it to future studies to explore the effect further.

Relation to Other Models and Additional DDM Variants

The most influential model for preferential choice is the *decision field theory* (DFT;

Busemeyer & Townsend, 1993; Busemeyer & Diederich, 2002; Busemeyer et al, 2019; Roe et al.,

2001). The DFT model was the first model that adopted the sequential sampling (integration to

bound) concept in its account of preferential choice. The model was applied to risky choice

(Busemeyer & Townsend, 1993), where it was able to account for violations of the independence

axiom (the Meyer effect; Busemeyer & Townsend, 1993, Fig. 2), and to multi-attribute decisions

(Roe et al., 2001), where it accounted for contextual reversal effects (the attraction, similarity, and

compromise effects; see also Johnson & Busemeyer, 2005, for an account of preference reversals

between choice and price elicitation in risky choice). Moreover, the DFT proposed a mechanism

for the stochastic nature of the evidence accumulation process, which it related to fluctuations in

attention between event outcomes or choice option attributes. The DFT model predicts that the

magnitude of this stochastic component (the diffusion noise term) increases with the outcome

variance associated with the options (which in turn reflects uncertainty). While in this study we do

not focus on risky choice nor on explicit multi-attribute choices (we focus here on holistic and

potential memory-based choices between food items), and we only consider relatively simple

DDM variants, it is important to discuss how our results compare with DFT-inspired uncertainty

modulations.

As we described above, the DFT suggests that uncertainty has a dual effect on the

preference formation — it increases both the diffusion noise and the response boundary (as a type

of compensation to the former). Our Model 2+3 version of the DDM was based on this idea, and

it successfully accounted for the qualitative effect of average certainty (C) on choice consistency

and on RT. This model version could not, however, account for the effect of certainty difference

(dC; the certainty of the high-value option minus the certainty of the low-value option) on choice

or on RT. This DDM extension was also outperformed by the snDDM in its quantitative account

of the data. Obviously, this is not a test of the DFT model (which is more complex, including other

factors such as an information leak, and which has not yet been applied to the type of tasks we

examined here), but merely suggests that value certainty is more likely to operate (in our task) via

a direct effect on the drift rate (rather than via modulation of the diffusion noise and of the response

boundary). Future work will be needed to examine in detail how the DFT accounts for choice and

RT for this type of preferential choice data.

We have focused here on simple DDM variants that include fixed (i.e., static across

decision time) response boundaries. While this is the type of DDM that has been explored most in

experimental psychology (Ratcliff, 1978; Ratcliff & Rouder, 1998), other contemporary versions

of the DDM sometimes include response boundaries that collapse over time (Evans, Trueblood, &

Holmes, 2020; Hawkins et al, 2015; Palestro et al, 2018; Glickman & Usher, 2019; Glickman et

al, 2019). Such collapsing bounds were first introduced to truncate the long tails of the RT

distributions that are predicted by a fixed-boundary DDM, especially for error trials (Milosavljevic

et al, 2010). Some researchers have since included some form of collapsing bound parameter in

their usage of the DDM, because it tends to provide better fits to the data (for preferential choice

see Glickman et al, 2019). Furthermore, Tajima et al (2016) and Fudenberg (2018) proved that the

DDM was an optimal model only in the case that the response boundary decreased across

deliberation time (although their work related to series of sequential decisions that created an

explicit opportunity cost of time, thereby mandating the collapsing bounds). In the current study,

we remained agnostic as to the shape of the DDM response boundary (fixed versus collapsing).

Whereas we relied on the fixed-boundary DDM in our model comparison, we verified that the

inclusion of a collapsing boundary does not change any of our conclusions (see Figure S1A in the

Supplementary Material), showing that, like for the fixed-boundary DDM (see Figure 3, center

panel), RT in the collapsing-boundary model also decreases with sampling noise that could be

associated with value-uncertainty (contrary to the data; see Figure 2). This can easily be understood

by examining the basic relationships between sampling noise and both response probability and

RT. If there were no noise in the sampling/accumulation process, the evidence would simply

accumulate along a straight trajectory until reaching a bound. Adding noise perturbs the trajectory

(at any point in time) and thus increases the probability that the response boundary will be reached

sooner. This is because the probability of reaching a boundary on the next time step is a decreasing

function of the current distance from the accumulator to the boundary, and an increasing function

of both the mean and the *variance* of the trajectory (which reflects the accumulation noise). Note

that this will hold for any response boundary, regardless of the shape. Thus, models such as our

Model 2 will also fail even when including collapsing boundaries (see Figure S1A in the

Supplementary Material for a confirmation of this via simulations).

Some versions of the DDM (Ratcliff & Rouder, 1998) include additional parameters, which

we did not examine in this study. Such parameters include non-decision time (time for encoding

and response, or T_{er}), starting point bias (and starting point variability), and drift rate variability.

While it might be mathematically possible to account for the effects of option-specific certainty

by allowing T_{er} to vary as a function of certainty, we believe that this would not be a principled

approach. Because T_{er} corresponds to time spent on perceptual (encoding) or motor (response)

processes, we see no reason why this should vary with value certainty (determined by valuation

processes). Similarly, we did not consider a starting point bias in the current study, as such a bias

is typically meant to capture prior expectations about which option is more likely to be better. In

the experimental paradigms that we considered, there are no such prior expectations – the choice

pairs were randomly created, each option was randomly assigned as either option 1 or option 2,

and the sequence of trials was randomized. Therefore, as participants proceeded to make choices

across trials, any potential biasing effects from one trial should be obliterated by the effects from

all other trials, leaving the prior expectations about which option will be better (on the next trial)

centered on zero. It remains possible that participants might have an inherent preference to pick

the option displayed on the left (or right) side of the screen, regardless of content. However, we

checked the data and found no evidence for a response side bias (see the Supplementary Material

for details). It would also be theoretically reasonable to consider starting point variability (around

a mean of zero). This could be due to a carry-over effect of the previous trial, where residual neural

activation might lead the decision maker to be inclined to repeat the same choice on the next trial.

However, this implies that starting point variability is caused by information from previous trials,

and not from the current trial (for which deliberation will have not yet begun). Hence, the starting

point variability is unlikely to be related to the option-specific value certainty on the current trial

and therefore cannot mediate the impact of certainty on either choice or RT. However, see the

Supplementary Material for a confirmation (via simulations) that even if one were to imagine that

starting point variability increased as a function of value uncertainty (a type of noisier responding),

it would have no impact on the mean RT (see Figure S1C), and it could definitely not account for

the slowdown with value uncertainty shown in the data (see Figure 2). Finally, we have not

considered drift rate variability in the current study. In some versions of the DDM (see Ratcliff &

McKoon, 2008), the drift rate on a given trial is a random variable selected from a normal

distribution whose mean equals the value difference of the options on that trial and with a variance

that corresponds to either attentional or memory biases that are specific to that trial (so if the same

choice is presented again, that drift may vary). In principle, it could be possible (but less likely,

we think) that value uncertainty would affect the drift rate variability (across trials) rather than

within-trial sampling noise. As we show in the Supplementary Material (Figure S1C), however,

such an assumption would not help, because increasing the variability in the drift rate parameter d

across trials with uncertainty, would have a similar effect on choice and RT as increasing the

within-trial diffusion noise (sampling variability) – higher variability will yield lower choice

consistency and faster RT, contrary to the data (see Figure S1C in the Supplementary Material for

a confirmation of this via simulations).

Another approach to the role of uncertainty on choice has been suggested in a recent study

by Li and Ma (2020), who proposed a model, in which the decision maker chooses based on a

comparison of the subjective utility of each option, where utility is a linear combination of the

mean of the value estimate and the standard deviation of the value estimate (derived via Bayesian

updating with the sequential evidence samples), suggesting that people explicitly prefer options

that are both more valuable *and* those whose value estimates are more certain. Note that such a

model does not modulate any of the standard DDM parameters as a function of value certainty,

but rather includes a separate evidence accumulator variable that tracks the difference in certainty

between choice options. We tested a model of this variety against our winning model and found

that although it did not win the competition, it did garner a non-trivial amount of support (see the

Supplementary Material for the formulation of Model 6 and the model comparison results). It

would be interesting for future work to explore the similarities and differences between this type

of model and the snDDM.

Choice Confidence

While we have focused here on how value certainty affects choice and RT, the experimental data also importantly show a marked and systematic effect of value certainty on choice confidence. In particular, higher value certainty sum (C) and certainty difference (dC) both lead to higher choice confidence. This pattern raises a further challenge for most accumulation-to-bound style choice models that aim to account for both RT and choice confidence. For example, in the balance of evidence (BOE) type models (Vickers & Packer, 1982; De Martino et al, 2013), confidence corresponds to the difference in the activation of two accumulators that independently race to a response boundary. If we were to naively introduce option-specific noise in such models, they would predict, contrary to the data, that the confidence becomes larger for options with more value uncertainty (as the noise increases the BOE). Similarly, if we were to model confidence using a DDM with collapsing boundaries (e.g., Tajima et al, 2016; Fudenberg et al, 2018), with confidence corresponding to the height of the boundary at the time the choice is made, naively introducing option-specific noise would once again provide us with a prediction opposite from what we see in the data. For uncertain alternatives, there would be more noise in the evidence accumulation process, resulting in faster choices and therefore higher boundaries, and thus higher confidence (in fact, this would be true for any model that assumes that confidence decreases with RT; Kiani & Shadlen, 2009).

There are very few value-based choice studies that simultaneously examined value certainty and choice confidence (but see Lee & Daunizeau, 2020, 2021; Lee & Coricelli, 2020; De Martino et al, 2013). We have not modeled choice confidence here, as there are many potential ways to do this, with substantial divergence among them (Vickers & Packer, 1982; Kiani & Shadlen, 2009; Pleskac & Busemeyer, 2010; De Martino et al, 2013; Moran, Teodorescu, & Usher, 2015; Calder-Travis, Bogacz, & Yeung, 2020; see Calder-Travis et al, 2020). Nevertheless, all of these models strive to predict a strong negative correlation between RT and choice confidence, as has been demonstrated in a plethora of experimental data. We note that in the data we examined, the impact of value certainty on choice confidence was essentially the reverse of its effect on RT (see Supplementary Material). While we did not explore this further, it suggests that a signal-to-noise DDM could also capture the dependency of choice confidence on value certainty. Future

work is needed to determine how signal detection style DDM variants might be extended towards

an optimal unified account of choice, RT, and choice confidence.

Funding

This research was supported by a grant from the United States-Israel Binational Science

Foundation/CNCRS (grant no 2014612 to MU).

Acknowledgments

We wish to thank Konstantinos Tsestsos and Giovanni Pezzulo for a critical reading of an early

version of the manuscript and for helpful discussions. We thank Ian Krajbich for generously

sharing his data with us. DL also wishes to thank Antonio Rangel for his supporting

encouragement at the initiation of this project.

BIBLIOGRAPHY

Alós-Ferrer, C. (2018). A Dual-Process Diffusion Model. Journal of Behavioral Decision Making, 31(2), 203-218.

Bakkour, A., Palombo, D. J., Zylberberg, A., Kang, Y. H., Reid, A., Verfaellie, M., ... & Shohamy, D. (2019). The hippocampus supports deliberation during value-based decisions. elife, 8, e46080.

Basten, U., Biele, G., Heekeren, H. R., & Fiebach, C. J. (2010). How the brain integrates costs and benefits during decision making. Proceedings of the National Academy of Sciences, 107(50), 21767-21772.

Bogacz, R., Brown, E., Moehlis, J., Holmes, P., & Cohen, J. D. (2006). The physics of optimal decision making: a formal analysis of models of performance in two-alternative forced-choice tasks. Psychological review, 113(4), 700.

Boldt, A., & Yeung, N. (2015). Shared neural markers of decision confidence and error detection. *Journal of Neuroscience*, *35*(8), 3478-3484.

Brown, S. D., & Heathcote, A. (2008). The simplest complete model of choice response time: Linear ballistic accumulation. Cognitive psychology, 57(3), 153-178.

Busemeyer, J. R., & Diederich, A. (2002). Survey of decision field theory. Mathematical Social Sciences, 43(3), 345-370.

Busemeyer, J. R., Gluth, S., Rieskamp, J., & Turner, B. M. (2019). Cognitive and neural bases of multi-attribute, multi-alternative, value-based decisions. Trends in cognitive sciences, 23(3), 251-263.

Busemeyer, J. R., & Townsend, J. T. (1993). Decision field theory: a dynamic-cognitive approach to decision making in an uncertain environment. Psychological review, 100(3), 432.

Calder-Travis, J., Bogacz, R., & Yeung, N. (2020). Bayesian confidence for drift diffusion observers in dynamic stimuli tasks. bioRxiv.

Calder-Travis, J., Charles, L., Bogacz, R., & Yeung, N. (2020). Bayesian confidence in optimal decisions. psyArXiv.

Callaway F, Rangel A, Griffiths TL (2021) Fixation patterns in simple choice reflect optimal information sampling. PLoS Comput Biol 17(3): e1008863.

Daunizeau, J., Adam, V., & Rigoux, L. (2014). VBA: a probabilistic treatment of nonlinear models for neurobiological and behavioural data. *PLoS Comput Biol*, *10*(1), e1003441.

De Gardelle, V., & Summerfield, C. (2011). Robust averaging during perceptual judgment. Proceedings of the National Academy of Sciences, 108(32), 13341-13346.

De Martino, B., Fleming, S. M., Garrett, N., & Dolan, R. J. (2013). Confidence in value-based choice. *Nature neuroscience*, *16*(1), 105-110.

Drugowitsch, J., DeAngelis, G. C., Klier, E. M., Angelaki, D. E., & Pouget, A. (2014). Optimal multisensory decision-making in a reaction-time task. eLife, 3, e03005.

Evans, N. J., Trueblood, J. S., & Holmes, W. R. (2020). A parameter recovery assessment of time-variant models of decision-making. Behavior research methods, 52(1), 193-206.

Feltgen, Q., & Daunizeau, J. (2021). Fitting drift-diffusion decision models to trial-by-trial data. bioRxiv.

Fetsch, C. R., Pouget, A., DeAngelis, G. C., & Angelaki, D. E. (2012). Neural correlates of reliability-based cue weighting during multisensory integration. Nature neuroscience, 15(1), 146-154.

Friston, K., Mattout, J., Trujillo-Barreto, N., Ashburner, J., & Penny, W. (2007). Variational free energy and the Laplace approximation. Neuroimage, 34(1), 220-234.

Fudenberg, D., Strack, P., & Strzalecki, T. (2018). Speed, accuracy, and the optimal timing of choices. American Economic Review, 108(12), 3651-84.

Glickman, M., & Usher, M. (2019). Integration to boundary in decisions between numerical sequences. Cognition, 193, 104022.

Glickman, M., Moran, R., & Usher, M. (2020). Evidence integration and decision-confidence are modulated by stimulus consistency. bioRxiv.

Glickman M, Sharoni O, Levy DJ, Niebur E, Stuphorn V, & Usher M (2019). The formation of preference in risky choice. PLOS Computational Biology.

Glöckner, A., & Betsch, T. (2008). Multiple-reason decision making based on automatic processing. Journal of experimental psychology: Learning, memory, and cognition, 34(5), 1055.

Gold, J. I., & Shadlen, M. N. (2007). The neural basis of decision making. *Annual review of neuroscience*, *30*.

Gwinn, R., & Krajbich, I. (2020). Attitudes and attention. *Journal of Experimental Social Psychology*, *86*, 103892.

Hawkins, G. E., Forstmann, B. U., Wagenmakers, E. J., Ratcliff, R., & Brown, S. D. (2015). Revisiting the evidence for collapsing boundaries and urgency signals in perceptual decision-making. Journal of Neuroscience, 35(6), 2476-2484.

Heng, J. A., Woodford, M., & Polania, R. (2020). Efficient sampling and noisy decisions. eLife, 9, e54962.

Hunt, L. T., Kolling, N., Soltani, A., Woolrich, M. W., Rushworth, M. F., & Behrens, T. E. (2012). Mechanisms underlying cortical activity during value-guided choice. Nature neuroscience, 15(3), 470-476.

Jang, A. I., Sharma, R., & Drugowitsch, J. (2021). Optimal policy for attention-modulated decisions explains human fixation behavior. eLife, 10, e63436.

Kiani, R., & Shadlen, M. N. (2009). Representation of confidence associated with a decision by neurons in the parietal cortex. Science, 324(5928), 759-764.

Krajbich, I., Armel, C., & Rangel, A. (2010). Visual fixations and the computation and comparison of value in simple choice. *Nature neuroscience*, *13*(10), 1292-1298.

Lebreton, M., Abitbol, R., Daunizeau, J., & Pessiglione, M. (2015). Automatic integration of confidence in the brain valuation signal. *Nature neuroscience*, *18*(8), 1159-1167.

Lee, D., & Coricelli, G. (2020). An Empirical Test of the Role of Value Certainty in Decision Making. Frontiers in psychology, 11, 574473.

Lee, D., & Daunizeau, J. (2020). Choosing what we like vs liking what we choose: How choice-induced preference change might actually be instrumental to decision-making. *PLoS One*, *15*(5), e0231081.

Lee, D., & Daunizeau, J. (2021). Trading Mental Effort for Confidence in the Metacognitive Control of Value-Based Decision-Making. *eLife*, in press.

Li, Z., & Ma, W. (2020). An uncertainty-based model of the effects of fixation on choice. PsyArXiv. doi.org/10.31234/osf.io/ajmwx

Lopez-Persem, A., Domenech, P., & Pessiglione, M. (2016). How prior preferences determine decision-making frames and biases in the human brain. eLife, 5, e20317.

Milosavljevic, M., Malmaud, J., Huth, A., Koch, C., & Rangel, A. (2010). The drift diffusion model can account for the accuracy and reaction time of value-based choices under high and low time pressure. Judgment and Decision Making, 5(6), 437.

Moran, R., Teodorescu, A. R., & Usher, M. (2015). Post choice information integration as a causal determinant of confidence: Novel data and a computational account. Cognitive psychology, 78, 99-147.

Palestro, J. J., Weichart, E., Sederberg, P. B., & Turner, B. M. (2018). Some task demands induce collapsing bounds: Evidence from a behavioral analysis. Psychonomic bulletin & review, 25(4), 1225-1248.

Penny, W. D. (2012). Comparing dynamic causal models using AIC, BIC and free energy. Neuroimage, 59(1), 319-330.

Philiastides, M. G., & Ratcliff, R. (2013). Influence of branding on preference-based decision making. Psychological science, 24(7), 1208-1215.

Pleskac, T. J., & Busemeyer, J. R. (2010). Two-stage dynamic signal detection: a theory of choice, decision time, and confidence. Psychological review, 117(3), 864.

Polanía, R., Krajbich, I., Grueschow, M., & Ruff, C. C. (2014). Neural oscillations and synchronization differentially support evidence accumulation in perceptual and value-based decision making. Neuron, 82(3), 709-720.

Polania, R., Woodford, M., & Ruff, C. C. (2019). Efficient coding of subjective value. *Nature neuroscience*, *22*(1), 134-142.

Polanía, R., Moisa, M., Opitz, A., Grueschow, M., & Ruff, C. C. (2015). The precision of value-based choices depends causally on fronto-parietal phase coupling. Nature communications, 6(1), 1-10.

Poldrack, R. A., Clark, J., Paré-Blagoev, E. A., Shohamy, D., Moyano, J. C., Myers, C., & Gluck, M. A. (2001). Interactive memory systems in the human brain. Nature, 414(6863), 546-550.

Ratcliff, R. (1978). A theory of memory retrieval. Psychological review, 85(2), 59.

Ratcliff, R. (1979). Group reaction time distributions and an analysis of distribution statistics. Psychological bulletin, 86(3), 446.

Ratcliff, R., & McKoon, G. (2018). Modeling Numerosity Representation with an Integrated Diffusion Model. Psychological Review, 125, 2, 183–217.

Ratcliff, R., & McKoon, G. (2008). The diffusion decision model: theory and data for two-choice decision tasks. Neural computation, 20(4), 873-922.

Ratcliff, R., & Rouder, J. N. (1998). Modeling response times for two-choice decisions. Psychological science, 9(5), 347-356.

Ratcliff, R., Smith, P. L., Brown, S. D., & McKoon, G. (2016). Diffusion decision model: Current issues and history. *Trends in cognitive sciences*, *20*(4), 260-281.

Ratcliff, R., & Tuerlinckx, F. (2002). Estimating parameters of the diffusion model: Approaches to dealing with contaminant reaction times and parameter variability. Psychonomic bulletin & review, 9(3), 438-481.

Ratcliff, R., Van Zandt, T., & McKoon, G. (1999). Connectionist and diffusion models of reaction time. Psychological review, 106(2), 261.

Ratcliff, R., Voskuilen, C., & Teodorescu, A. (2018). Modeling 2-alternative forced-choice tasks: Accounting for both magnitude and difference effects. Cognitive psychology, 103, 1-22.

Rigoux, L., Stephan, K. E., Friston, K. J., & Daunizeau, J. (2014). Bayesian model selection for group studies—revisited. Neuroimage, 84, 971-985.

Roe, R. M., Busemeyer, J. R., & Townsend, J. T. (2001). Multialternative decision field theory: A dynamic connectionist model of decision-making. Psychological Review, 108,370–392.

Schacter, D. L., Addis, D. R., & Buckner, R. L. (2007). Remembering the past to imagine the future: the prospective brain. Nature reviews neuroscience, 8(9), 657-661.

Sepulveda, P., Usher, M., Davies, N., Benson, A. A., Ortoleva, P., & De Martino, B. (2020). Visual attention modulates the integration of goal-relevant evidence and not value. eLife, 9, e60705.

Smith, S. M., & Krajbich, I. (2019). Gaze amplifies value in decision making. Psychological science, 30(1), 116-128.

Stephan, K. E., Penny, W. D., Daunizeau, J., Moran, R. J., & Friston, K. J. (2009). Bayesian model selection for group studies. Neuroimage, 46(4), 1004-1017.

Tajima, S., Drugowitsch, J., & Pouget, A. (2016). Optimal policy for value-based decision-making. *Nature communications*, *7*(1), 1-12.

Teodorescu, A. R., Moran, R., & Usher, M. (2016). Absolutely relative or relatively absolute: violations of value invariance in human decision making. Psychonomic bulletin & review, 23(1), 22-38.

Turner, B. M., Schley, D. R., Muller, C., & Tsetsos, K. (2018). Competing theories of multialternative, multiattribute preferential choice. Psychological Review, 125(3), 329–362.

Usher, M. and McClelland, J. L. (2004). Loss aversion and inhibition in dynamical models of multialternative choice (proofs version). Psychological Review, 111 (3), 757-769

Usher, M., & McClelland, J. L. (2001). The time course of perceptual choice: the leaky, competing accumulator model. Psychological review, 108(3), 550.

Vickers, D. (1970). Evidence for an accumulator model of psychophysical discrimination. Ergonomics, 13(1), 37-58.

Vickers, D., & Packer, J. (1982). Effects of alternating set for speed or accuracy on response time, accuracy and confidence in a unidimensional discrimination task. Acta psychologica, 50(2), 179-197.

Wickelgren, W. A. (1977). Speed-accuracy tradeoff and information processing dynamics. Acta psychologica, 41(1), 67-85.

Supplementary Material

Simulation Results for the Impact of Value Certainty on Other DDM Parameters

In the Discussion section of the main manuscript, we explained why additional DDM
variants (collapsing boundaries) or other assumptions about the impact of value certainty on the
additional parameters in the standard DDM (drift rate variability, starting point variability) would
not change the conclusions we presented here. Here we present simulation results that provide
additional support to our conceptual explanations (Figure S1).

We start with the collapsing-boundary DDM variant. We show via simulations that if value
uncertainty increased the sampling noise, this would decrease RT, just like in the fixed-boundary
DDM (Figure S1A). We then consider the classic DDM with fixed boundaries and we examine
whether value certainty could be introduced via an effect on either starting point variability or on
drift rate variability (we consider the possibility that uncertainty might increase those variability
parameters, so as to make to make the choice more random). We show via simulations that if value
uncertainty increased the drift rate variability, this would decrease RT (Figure S1B). We also show
via simulations that if value uncertainty increased the starting point variability, this would have no
impact on mean RT (Figure S1C).

Figure S1: *Simulated results demonstrating the relationship between additional DDM*
parameters and RT. For each plot, we simulated 10^4 trials. Panel A show the relationships between
diffusion noise and RT (see Figure 3, center panel), when the response boundaries collapse over
time. Panel B shows the relationships between drift rate variability and RT. Panel C shows the
relationship between starting point variability and RT. Each data point represents one simulated
trial. The solid black lines indicate the mean RT as a function of the independent variable of
interest.

 Response Time Distributions as a Function of snDDM Predicted Drift Rate

To verify that the snDDM can account for patterns in the full RT distributions (rather than
only in mean RT), we performed an RT quantile analysis as advocated by previous studies
(Ratcliff, 1979; Ratcliff, Van Zandt, & McKoon, 1999; Ratcliff & Tuerlinckx, 2002). Because we
had many participants but only few trials per participant, we utilized a technique recommended by
Ratcliff (1979), in which the independent variable of interest is summarized by its 10^{th}, 30^{th}, 50^{th},
70^{th}, and 90^{th} percentiles for each participant and then averaged across participants. We present
this summary for all subjects pooled together across studies in Figure S2, for the key snDDM
variable of interest: (signal-to-noise difference) SN = V1*C1 – V2*C2. To obtain this data, we
first separated the trials by a median split (within participant) and then divided the trials within
each half of the median split by choice consistency (i.e., whether the higher-rated option was
chosen or not). This provided us with four bins (high SN / inconsistent, low SN / inconsistent, low
SN / consistent, high SN / consistent), within which we found the RT percentiles for each
participant and then took the average across participants (Figure S2, left panel). The location on
the x-axis represents the mean consistency level (across participants) for trials within each of the
categories. To compare the empirical results with snDDM predictions, we simulated choice and
RT using the real experimental input variables (V1, V2, C1, C2) and the best fit parameters for
each participant. We then performed the identical analysis as with the experimental data (Figure
S2, right panel). We found that the patterns across the full range of RT were similar between the
simulated and the experimental data. In particular, in both, we can see the skew in the RT
distribution (a larger separation for the higher RT quantiles compared the lower ones), and the fact
that while the higher quantiles show a characteristic inverse U-shape, the lower quantiles appear
to decrease monotonically with choice consistency.

Figure S2: Quantile probability plots: *RT percentiles as a function of response consistency (equivalent to choice accuracy), for different levels (median split) of the predicted signal-to-noise drift rate (SN). For each point, the x-coordinate represents the categorical bin {high SN / inconsistent choice, low SN / inconsistent choice, low SN / consistent choice, high SN consistent choice} and the y-coordinate represents the cross-participant mean of the RT quantile {0.1, 0.3, 0.5, 0.7, 0.9}. The left panel summarizes the experimental data (pooled across Studies 1-4) and the right panel summarizes the snDDM simulated output using real experimental input.*

Below we illustrate the snDDM predicted relationship between C / dC and drift rate, which is inversely related to RT (Figure S3, left panel). The figure shows predictions across the entire (normalized) range of possible values of C1 and C2, of which C and dC are composed. We compare this to the experimental data (Figure S3, right panel), which summarizes RT (across participants) as a function of C1 and C2 (and thus C and dC). Note that while the match is not exact, the expected qualitative patterns hold between the theoretical and empirical data (i.e., RT is a decreasing function of both C and dC).

Figure S3: *Drift rate under the snDDM is an increasing function of both dC and C (left panel). Averaged across the full normalized range of value (both V1 and V2), drift rate increased as either option increased in certainty in isolation (i.e., dC increased; moving upwards or rightwards in the plot), but also as both options increased simultaneously (i.e., C increased while dC remained constant; moving at a diagonal towards the upper right of the plot). Average RT (across participants) in Studies 1-4 shows the opposite pattern as the theoretical drift rate, as expected.*

Impact of Overall Value on Response Time

Beyond the effects of value difference (dV), certainty sum (C), and certainty difference (dC) on RT, we also observed a strong effect of value sum (V). Specifically, higher V corresponds to lower RT (see Figure S4). As it turns out, the snDDM is able to predict this relationship (none of the other models can account for this; see Figure S4). This is especially interesting, given the fact that V itself never directly enters the model. Instead, the impact of V likely arises from the nature of the signal-to-noise structure of the model. The basic idea would be that the value signals for lower-valued options would be washed away by the detrimental impact of uncertainty, whereas the value signals for higher-valued options would be more resilient. For example, imagine two separate choice pairs: a) V1 = 0.2, V2 = 0.1, C1 = 0.8, C2 = 0.2; b) V1 = 0.9, V2 = 0.8, C1 = 0.8, C2 = 0.2. Under the standard DDM, both choice pairs would have the same mean evidence (e) dV = V1 - V2 = 0.1. Under the snDDM, the mean evidence for each pair is distinct: a) V1*C1 = 0.16, V2*C2 = 0.02 → e = 0.14; b) V1*C1 = 0.72, V2*C2 = 0.16 → e = 0.56. The larger mean evidence with larger overall value effectively makes the choice easier (i.e., the drift rate is higher), thus leading to lower RT. The same phenomena holds if the higher-valued option within each choice

pair is the one with higher certainty, for example: a) V1 = 0.2, V2 = 0.1, C1 = 0.2, C2 = 0.8; b)

V1 = 0.9, V2 = 0.8, C1 = 0.2, C2 = 0.8. Now, the mean evidence for each pair is: a) V1*C1 = 0.04,

V2*C2 = 0.08 → e = -0.04; b) V1*C1 = 0.18, V2*C2 = 0.64 → e = -0.46. Again, the higher-valued

choice is effectively easier (greater absolute evidence magnitude), and the choice can be made

more quickly. Note that if both options within a choice pair had the same level of certainty, there

would be no impact of V on RT, which explains why none of the other models we examined were

able to account for that effect.

1005

1006

Figure S4: *Impact of overall value on response time. The upper panel shows the fixed effects*

coefficients from mixed effect linear regression of value difference (dV) and value sum (V) on

log(RT). Participants pooled across four studies, n=191; error bars represent standard errors;

significance stars represent *** p < .001, one-sided t-tests. The bottom panel shows the same

analysis for data simulated under each of the models we compared. Note that only Model 5

qualitatively matches the data.

1013   Beyond the illustrative examples above, we also explored the relationship between V and

RT in a more systematic manner. Specifically, we examined a 100x100x100x100 grid of V1, V2,

C1, and C2 across a standardized range of 0:0.01:1. Within each cell in the grid, we calculated the

drift rate as |V1*C1 – V2*C2| (we took the absolute value because it is the magnitude of the drift

that impacts RT; the sign is irrelevant). We then averaged across all values of C1 and C2, leaving

us with a two-dimensional array (V1 x V2) of calculated drift rates (see Figure S5). As expected,

the drift rates were highest when |dV| was largest (so when V1 and V2 had much different values).

Beyond that, there was a clear positive relationship between drift rate and V, explaining the

negative relationship between V and RT (which is inversely related to drift rate).

1022

Figure S5: *Drift rate under the snDDM is an increasing function of both dV and V (left panel).*

Averaged across the full normalized range of certainty (both C1 and C2), drift rate increased as

either option increased in value in isolation (i.e., dV increased; moving upwards or rightwards in

the plot), but also as both options increased simultaneously (i.e., V increased while dV remained

constant; moving at a diagonal towards the upper right of the plot). Average RT (across

participants) in Studies 1-4 shows the opposite pattern as the theoretical drift rate, as expected.

Choice Confidence

In this study, we chose not to include choice confidence in our model predictions, as there

is not currently an agreed-upon standard for doing so. Nevertheless, we did briefly examine this

variable in those datasets that contained it (Studies 1-3; Lee & Daunizeau, 2020 2021; Lee &

Coricelli, 2020). In general, choice confidence exhibited patterns qualitatively opposite to those

exhibited by RT. Specifically, the mixed model fixed effects coefficients for dV, C, and dC were

of similar magnitude as those for RT, but were all positive (whereas for RT, they were all negative;

see Figure S6).

1037

Figure S6: *The fixed effects coefficients from mixed effect linear regression of dV, C, and dC on choice confidence (left panel) and log(RT) (right panel). Participants pooled across three studies, n=155; error bars represent standard errors; significance stars represent *** p < .001, one-sided t-tests.*

DDM Median Split Parameter Comparison

To provide additional corroborating evidence that value certainty should relate to the drift rate and not the diffusion noise parameter, we performed classic median-split analyses on both the sum and the difference of the value certainties of the choice pair options. For each participant across all studies, we divided the trials based on a participant-specific median split of either certainty sum or certainty difference. We then fit the classic DDM separately for each of the subsets of data and compared the fitted parameters across subsets. As shown in Figure S7, in both the certainty sum median split and the certainty difference median split, the drift rate parameter was larger in the high versus low subsets; the noise parameters did not differ across subsets. This shows that our conclusion that Model 5 (with drift rate adjusted by certainty) is valid, while Model 2 (with diffusion noise adjusted by certainty) is not.

1053

Figure S7: *DDM parameter comparison across median splits of certainty sum (top panels) and certainty difference (bottom panels). Each data point represents one participant. Across the population, the fitted drift rate parameters were larger for the high certainty trials as well as for the high certainty difference trials (left panels); the fitted diffusion noise parameters were not different across high and low certainty or certainty difference (right panels).*

Starting Point Bias towards Left or Right Option

1059

As a simple check to see if the participants in the studies we examined had a tendency to favor the options presented on the left or right side of the screen (independent of option values), we calculated a simple percentage of the total choice trials for which the participants chose the left option (since the value of each option was orthogonal to its physical location on the screen). Overall, the participants chose the left option on 50% of the trials. At the individual level, 91% of

participants chose the left option on 40-60% of trials. We tested that these individual deviations

from 50% were due to chance rather than bias. We thus calculated, for each participant, the

standard deviation of a binomial distribution ($\sqrt{p * q * n}$) with a 50% probability (p = q = 0.5) and

n = number of trials for that participant (which varied across datasets). We then computed a t-

statistic for each participant by subtracting n/2 from the number of left responses and dividing by

the standard deviation. Using t = +/- 1.96 as a cutoff for a 95% confidence interval that the leftward

response rate for each participant was not statistically different than 0.5, we found that only 8% of

participants fell outside of this confidence interval. We thus concluded that there was no

meaningful side response bias in our data.

1074   *Model 6: Separate Accumulator for Certainty*

In this study, we examined a number of alternative ways in which the traditional DDM

parameters (drift rate, diffusion noise, and response boundary) could be modulated by option-

specific value certainty. Yet a fundamentally different way in which the DDM could include value

certainty is in the form of an independent evidence accumulator—a secondary drift, with a rate

proportional to the difference in certainty between the choice options. In this way, the decision

about which option to choose would be influenced both by which value estimate was higher (via

the primary, standard drift) and by which value estimate was more certain (via the secondary, novel

drift). The secondary drift might represent an aversion to uncertainty, where the deliberation

process would be both slowed by the uncertainty and the decision pulled towards the more certain

option. This would imply that the decision maker prefers options that are more valuable, but also

options for which the value is more certain. Other ongoing research has proposed a model based

on a similar principle (Li & Ma, 2020). Under our version of this framework, the equation

governing the DDM process would be:

1088
$$E_{t+1} = E_t + d_v \Delta_v + d_c \Delta_c$$

1089
$$\Delta_v \sim N(\mu_1 - \mu_2, \sigma_v^2)$$

1090
$$\Delta_c \sim N\left(\frac{1}{\sigma_1} - \frac{1}{\sigma_2}, \sigma_c^2\right)$$

where Δ_v and Δ_c are the incremental evidence for value and certainty, respectively, and d_v and d_c

are scalars (it is assumed that d_v will always be positive, but d_c could take either sign). The

inclusion of a secondary drift rate (in essence, a parallel evidence stream that monitors value

certainty rather than value itself) can have either a positive or a negative impact on both choice

and RT, depending on whether the option with the higher mean evidence stream is the same as or

different than the option with the higher evidence reliability (as well as on the sign of d_c). This can

be seen by examining the (revised) DDM equations for choice probability and expected response

time:

1099
$$p(ch = 1) = \frac{1}{1 + e^{\left(\frac{-2\theta}{\sigma_v^2 + \sigma_c^2}\left[d_v(\mu_1 - \mu_2) + d_c\left(\frac{1}{\sigma_1} - \frac{1}{\sigma_2}\right)\right]\right)}}$$

1100
$$E[RT] = \frac{\theta(2 * p(ch = 1) - 1)}{d_v(\mu_1 - \mu_2) + d_c\left(\frac{1}{\sigma_1} - \frac{1}{\sigma_2}\right)}$$

Here, the secondary drift rate will result in more consistent and faster choices if the sign of $\mu_1 - \mu_2$

is the same as that of $\frac{1}{\sigma_1} - \frac{1}{\sigma_2}$, but it will result in less consistent and slower choices otherwise. This,

of course, is under the assumption that the decision maker is uncertainty averse (i.e., $d_c > 0$). If the

decision maker were uncertainty seeking (i.e., $d_c < 0$), the opposite predictions would hold.

Model 6 could account for the empirical relationship between certainty difference (dC) and

both choice consistency and RT. However, Model 6 would fail to account for the dependency of

consistency and RT on certainty sum (C) (see Figure S8).

1108

Figure S8: *Qualitative predictions of the effects of value difference (dV = V1 - V2), value certainty*

(C = C1 + C2), and certainty difference (dC = C1 - C2) on choice consistency and log(RT) (shown

for the best fit parameters; bar heights represent mean GLM beta weights based on 10^4 simulated

trials for Model 6). The log(RT) data were z-scored prior to binning for the quantile curve plots.

To test the ability of this alternative formulation to explain the data, we ran a formal model

comparison between our winning model (Model 5*) and Model 6. It is worth noting that while

Model 5* dominated the model comparison overall (exceedance probability = 1), Model 6

nevertheless received a non-trivial amount of support. Approximately 33% of participants (pooled

across all datasets) had data that were best fit by Model 6 (see Figure S9). This suggests that

whereas Model 5* might be more prevalent across the general population, there may exist

individual differences in decision-making apparatus such that Model 6 represents a sizeable

minority of the population.

1121

Figure S9: Model comparison results for Model 5 vs Model 6. We show here the probability that

each model best accounted for the data at the participant level (left panel), across the studies we

examined; each cell represents the probability that the model (column) best represents the

behavior of the participant (row). The dashed lines separate participants from each of the different

studies, for visualization purposes only. We also show the probability that each model best

explains the data across the participant population (right panel), across all studies. The dashed

line indicates chance level if all models were equally probable a priori.

The final analysis we performed with respect to Model 6 was to include it in the model

recovery exercise discussed in the main text. It turns out that Models 5* and 6 had a substantial

level of confusion between them. Specifically, data generated by a true Model 5* was most often

attributed to a best fit Model 6 (see Figure S10). This suggests that the results of the formal

comparison between Models 5* and 6 (see above) should be interpreted with caution. Specifically,

participants for whom Model 6 appeared to be the best model might actually behave according to

Model 5* and our model-fitting procedure was simply unable to make that distinction.

1136

1137 ***Figure S10:*** *Model recovery analysis. Each cell in the "confusion matrix" summarizes the*

1138 *percentage of simulated participants (under each true model) for which our model-fitting*

1139 *procedure attributed each of the (best fit) models. Note that with Model 6 included in the analysis,*

1140 *Model 5* is no longer accurately recovered. Instead, most participants simulated under true*

1141 *Model 5* are now attributed to best fit model 6.*