Title: Intra and inter-annual climatic conditions have stronger effect than grazing intensity on

root growth of permanent grasslands

Running head: Root production in grazed grasslands

List of authors: Catherine Picon-Cochard[1*], Nathalie Vassal[1,2], Raphaël Martin[1], Damien

Herfurth[1,2], Priscilla Note[1,3], Frédérique Louault[1]

Institute of origin:

1. Université Clermont Auvergne, INRAE, VetAgro Sup, UREP, 63000 Clermont-

Ferrand, France

2. Université Clermont Auvergne, VetAgro Sup, 63370 Lempdes, France

3. INRAE, UE1414, Herbipôle, 63122 Saint-Genès-Champanelle, France

*: Corresponding author: Catherine Picon-Cochard, Tel: (+)33 443 761 615; email:

catherine.picon-cochard@inrae.fr

Keywords: annual root and above-ground production; ingrowth core; leaf and root traits; root

dynamics; rotational grazing; soil moisture; soil temperature;

Type of paper: Regular article

Abstract

- **Background and Aims**. Understanding how direct and indirect changes in climatic conditions, management, and species composition affect root production and root traits is of prime importance for grassland C sequestration service delivery. We aim studying the effects of climatic variability and gradient of herbage utilization by grazing on root production during two years. Root and leaf traits of the communities were determined to detect their capacity to predict above and below-ground production (ANPP, BNPP).

- **Methods**. A long-term field experiment was used to compare after 10 years of treatment application, the effects of abandonment, low (Ca-) and high (Ca+) grazing intensities, respectively 15.2 cm and 7.7cm mean residual plant height, induced by grazing rotation on upland fertile grasslands. Ingrowth-cores and exclusion cages were used to measure, respectively, root and shoot mass production several times each year and at annual scale. Root and leaf traits of the communities were measured close to the vegetation peak growing season.

- **Results**. We observed strong seasonal root production across treatments in both a wet and a dry year but response to grazing intensity was hardly observed within growing seasons. In abandonment, spring and autumn peaks of root growth were delayed by about one month compared to the two cattle treatments, possibly due to later canopy green-up induced by lower soil temperature and accumulation of litter. BNPP was slightly lower in abandonment compared to cattle treatments only during the dry year, whereas decline of ANPP in abandonment compared to Ca+ was observed the wet year. In response to drought (case of the second year), the root-to-shoot biomass ratio was stable in cattle treatments but declined in abandonment. The higher allocation to root mass can be beneficial for plant communities under drier conditions.

- **Conclusions**. Rotational grazing pressure and climatic conditions variability had very limited effects on root growth seasonality although drought had stronger effects on BNPP than

on ANPP. The stability of the root-to-shoot biomass ratio during the dry year put into evidence

higher resistance to drought of grazed grassland communities compared to abandonment.

Introduction

Permanent grasslands provide many services that tie in to human activities through livestock

products, but also contribute to regulate greenhouse gas emission, because their soils

accumulate large amounts of carbon in organic matter fractions. Root activity (growth,

exudation, turnover) contributes to C and N inputs but is also a main determinant of both

nutrients and water uptake, essential to fix atmospheric CO_2 and to produce biomass.

Intensification of management practices may affect these services as well as climate variability

(Conant et al. 2001; Jones and Donnelly 2004; Soussana and Duru 2007). Thus, improving our

understanding of grassland roots dynamics under different management and climatic conditions

may help to identify management options to maintain forage production and C sequestration

abilities of this ecosystem and thus its sustainability.

Different practices of management modify forage production and the amount of soil C and

N through direct effects of defoliation, fertilization or returns of excreta to soil on root growth

and soil abiotic factors and indirect effects through species composition changes (Bardgett and

Wardle 2003; Dawson et al. 2000; Soussana et al. 2004). In mown grasslands, it has been shown

that root mass production is generally lower when grass is frequently mown and fertilised

(Leuschner et al. 2013; Picon-Cochard et al. 2009). This may be explained by changes in root-

to-shoot allocation, with increase of above-ground growth in order to maximize light capture.

The complexity of these phenomena in grazed grassland is greater than in mown systems owing

to animals' selective defoliation of plant species, and also because returns to soil are spatially

heterogeneous (Rossignol et al. 2011). In addition, level of soil fertility may buffer the degree

of root response to defoliation in grazed grasslands as plants exhibit specific responses to

defoliation in fertile and unfertile grasslands (Duru et al. 1998). Overall, this can explain why

no clear trend is found for the effects of grazing on above- and below-ground production (e.g.

see syntheses of Milchunas and Lauenroth (1993) and McSherry and Ritchie (2013)), although

two meta-analyses emphasized negative effect of grazing intensity on above- and below-ground

carbon stocks compared to ungrazed systems (Zhou et al. 2017; Li et al. 2018). In addition,

repeated defoliations induced by grazing or mowing of grassland can simultaneously increase

soil temperature and soil moisture (Moretto et al. 2001; Pineiro et al. 2010; Smith et al. 2014).

Soil moisture can also be modified by high stocking rate through changes of soil bulk density

due to soil compaction and by changes of leaf area index after defoliation (Pineiro et al. 2010).

These direct effects of grazing on soil abiotic factors should affect root growth of grazed

grassland, although all these phenomena are not very well documented in field conditions.

Species composition change induced by management is also an important determinant of

above- and below-ground response in grazed grassland. Intensive practices (high grazing

intensity, fertilization) generally favour the development of fast growing species (exploitative

strategy) whereas at the opposite extensive practices (low grazing intensity, absence of

fertilization) favour slow growing species (conservative strategy) (Klumpp et al. 2009; Louault

et al. 2005; Soussana and Lemaire 2014; Wardle et al. 2004). Root-to-shoot biomass allocation,

but also functional traits (used as proxies of ecosystems properties like ANPP or BNPP, e.g.

Laliberté and Tylianakis 2012), are thus likely to change in response to intensification of

practices, e.g. from ungrazed to intensive grazed temperate grassland (Klumpp and Soussana

2009) or in alpine meadows, steppes and desert-steppes (Zeng et al. 2015). Overall, according

to Ziter and MacDougall (2013), the uncertainty surrounding nutrient-defoliation responses

makes it difficult to predict whether C storage will be higher in managed compared to

unmanaged grasslands. Thus, soil fertility should be considered when comparing different

grazing intensities in grassland (Louault et al. 2005).

Increased climate variability is another source of response uncertainty in managed

ecosystems. As more frequent and longer period of drought associated with heat waves may

threaten and shape the long-term dynamics of perennial ecosystems such as grasslands

(Brookshire and Weaver 2015), it is important to understand how above- and below-ground

compartments respond to climate variability. However, there are few data on above- and below-

ground biomass responses to drought for grassland (Byrne et al. 2013; Wilcox et al. 2015; Li et

al. 2018), although some evidence shows that the 'slow' trait strategy (resource conservation)

is associated with drought tolerance (Pérez-Ramos et al. 2012; Reich 2014). Changes in root

morphology and functioning may thus be important determinants in plant adaptive strategies to

drought, and have been less studied than above-ground plant responses (Biswell and Weaver

1933; Dawson et al. 2000; McInenly et al. 2010). However, there are not enough data to make

generalizations about combined impacts of management and climatic conditions variability

such as precipitation reduction on root and shoot biomass production and plant traits defining

plant strategies related to resource use and grazing intensity.

The study was carried out in a long-term field experiment for which controlled grazing

intensity had been applied for 10 years. We compared abandonment of grazing and two levels

of herbage utilization by grazing based on five rotations per year. In two consecutive years, the

ingrowth core method was used to measure monthly root biomass production and calculate

annual root production (BNPP); ANPP was measured by grazing exclusion cages and

community-weighted mean leaf and root traits were assessed the first year. We tested the

following hypotheses: (i) high grazing intensity increases above-ground mass at the expense of

root production as a result of the direct negative effect of defoliation on root growth, whatever

the climatic conditions, (ii) inter-annual climatic conditions modulate above and below-ground

biomass production response to grazing intensity as a consequence of higher presence of

defoliation tolerant and drought-sensitive species (*Lolium perenne* or *Trifolium repens*) in the

high grazing intensity treatment; (iii) root traits respond to treatment and is a determinant of

BNPP, as observed for leaf traits for ANPP.

Materials and methods

<u>Site characteristics</u>

The experiment took place in the long-term observatory network (ACBB-SOERE) located at

St-Genès-Champanelle, France (45°43′N, 03°01′E, 880 m a.s.l.). The local climate is semi-

continental with oceanic influences (mean annual temperature 8.5 °C, mean annual

precipitation 784 mm, Table 1). The site supports mesotrophic multi-specific permanent

grassland, dominated by species with high Ellenberg indicator values for N (Schaffers and

Sykora 2000), indicating a high level of fertility for the site (Table S1; Louault et al. 2017). The

soil is a cambisol with a sandy loam texture, developed on granitic bedrock. Differences in local

soil composition and profile led us to consider two blocks characterized respectively by a eutric

cambisol (54% sand; 26% silt; 20% clay; 7.0% organic matter; pH: 5.9) and a colluvic cambisol

(50% sand; 26% silt; 24% clay; 7.4% organic matter; pH: 6.0) including some volcanic

materials.

<u>Management</u>

Prior to the installation of this experiment in 2005, the study area had been used for intensive

hay and silage production (combining grazing, mowing and fertilization), with mineral

fertilization, and two years preceding the start of the experiment (2003 and 2004), the grassland

site was mown three times per year without fertilization. Then, from 2005, the grassland had

been managed for 10 years with a gradient of grazing intensity resulting from three treatments:

abandonment (Ab), low (Cattle-) and high (Cattle+) level of herbage utilization obtained by

modification of stocking density (6.9 and 13.8 LSU ha^{-1}, livestock unit, respectively) with five

grazing rotations each year: mid-April, late May, early July, September and November, lasting

on average 9.6, 9.0, 10.7, 8.6, and 2.1 days, respectively. The two cattle treatments

corresponded to two levels of herbage utilization by grazing, and had on average 15.2 ± 0.5 cm

(mean $\pm$ se, Cattle-) and 7.7 ± 0.2 cm (Cattle+) residual plant height at the end of each grazing

rotation, respectively. For each treatment, two replicate plots were set up per block, resulting in

four replicates per treatment, and a total of 12 plots (2 blocks x 2 plots x 3 treatments). The

average distance between the two blocks is about 230 m and all treatments are randomized

within each block. The size of the plots differs according to treatments: 2200 m^2 for the two

cattle treatments and 400 m^2 for the abandonment.

<u>Climatic and edaphic conditions</u>

Daily precipitation (mm) and air temperature (°C) were measured for the two years, and

recorded with a meteorological station located at the site. An aridity index was calculated as

precipitation minus potential evapotranspiration (P - PET, mm) with the Penman-Monteith

equation. Daily soil temperature (°C) was measured with thermocouple sensors (home-made

copper-constantan sensors) inserted at 20 cm depth in each plot and recorded with a HOBO

data logger (U12-014, Onset Instruments, MA, USA). Daily soil volumetric water content

(SWC, m^3 m^{-3}) of each plot was measured with two probes (ECHO-10, Decagon, USA),

inserted horizontally at 20 cm depth, and connected to dataloggers (EM5 and EM50, Decagon,

USA). From January 2014 to November 2015 (DOY 132–326), SWC was measured every 30

min and averaged at daily scale. For each plot, average values of the two probes were used.

Daily relative soil water content data are shown and calculated as the ratio:

$RSWC = \frac{SWC - SWCmin}{SWCmax - SWCmin}$, where SWC is the soil moisture at a given day, SWCmin is the

minimum value of soil moisture and SWCmax is the maximum value of soil moisture, both

observed during the two years. For soil temperature and RSWC, values were averaged

according to root growth time scale.

<u>Root growth and root mass</u>

Six months beforehand, shallow (0-20 cm) soil was collected on each of the two blocks of the

site and sieved (5 mm mesh size) to remove stones and coarse organic matter, and then left

unused outside covered under a shelter and protected from direct sunlight. Thereafter, this air-

dried soil was used to fill the ingrowth-core each month.

In December 2013 and for each of the 12 plots, soil cores were collected with an auger (8 cm

diameter, 0-20 cm depth) at four locations representative of the plant community in the

treatment. On average mean distance between locations are 19.8 m ± 0.2, 21.7 m ± 0.1 and 17.2

m ± 0.2 for Ca+, Ca- and Ab (mean ± SD, see Fig S1), respectively. After core harvest, each

hole was filled with a plastic net (8 mm mesh size) containing a fixed volume of air-dried sieved

soil (ingrowth core), collected six months beforehand. Then, about each month and for two

years (2 x 10 times), ingrowth cores, containing soil and the root and rhizome material that had

grown therein, were extracted, and then replenished with another fixed volume of dry sieved

soil. Thus monthly and annual root production (BNPP, $g\ m^{-2}\ y^{-1}$) were measured from February

2014 to December 2015. Root production period ranged on average 36.5 days, but with longer

and shorter periods in winter and spring-summer, respectively (Table 1). In periods with

absence of precipitation, a fixed volume of water was added to adjust soil humidity to field

conditions. After collection, the ingrowth cores were transported to the laboratory and

immediately stored at 4 °C before processing in the next five days. The roots were washed

under tap water and with a 200 μm sieve, and then oven-dried (48 h, 60 °C).

In order to measure root mass stock, soil cores were collected three times (December 2013,

March and June 2014) with the same auger and near the ingrowth cores locations. These

samples were stored in the freezer (-18°C), and after defrosting, the roots were washed with the

same procedure as that used for the ingrowth cores, and then oven-dried (48 h, 60 °C).

Root traits

Subsamples of washed roots collected with the ingrowth cores collected in June 2014, were

fresh weighed, and then frozen (-18 °C) before morphology analysis. After defrosting, roots

were stained with methylene blue (5 g L^{-1}) for about 5-10 minutes, rinsed in water, spread in a

transparent glass box containing a thin layer of water, and covered with a transparent plastic

sheet. High resolution images were recorded with a double light scanner (800 dpi, perfection

V700, Epson, JA) and analyzed with WinRhizo software (PRO 2012b, Regent Instruments,

CA) with the automatic procedure. Two scans per location were recorded and separately

analyzed to measure root length (m), root volume (cm^3), root surface area (m^2), average root

diameter (mm) and length by class diameter (13 classes: 11 with 0.1 mm interval and 2 with

0.5 mm interval). Specific root length (m g^{-1}), root tissue density (g cm^{-3}) and specific root area

(m^2 g^{-1}) were calculated for fine roots as in Picon-Cochard et al. (2012).

Botanical composition

Species contribution (%) was visually observed on a circle (20 cm diameter) around each

ingrowth core location in April (cattle treatments) and May (abandonment) 2014. For each

zone, a score on a ten-point scale was allocated to species present according to their volume

occupancy, and the percentage of each species was calculated at the plot scale by averaging

values of the four zones. The list of species and their relative contributions is given in Table

S3.

Above-ground biomass production

On each plot and on each sampling date, four fenced sampling areas (0.6 × 0.6 m) were used to measure accumulation of above-ground biomass after above-ground standing biomass was clipped at 5.5 cm. At each sampling date, biomass was sampled at a height of 5.5 cm, oven-dried and weighed. Measurements were made five times in the course of the year, before each grazing event in Cattle+ and Cattle- plots, and three times (spring, summer, autumn) in abandonment plots. Sampling areas were moved within the plot at each measurement date during the year. Annual above-ground net primary production (ANPP, $g\,m^{-2}\,y^{-1}$) was calculated as the sum of the successive biomass accumulations along the year.

Leaf traits

Community-weighted mean (CWM) trait values of leaf dry matter content (LDMC), specific leaf area (SLA) and reproductive plant height (H) were calculated for each ingrowth core zone using (i) the relative contribution of the dominant species to the community (i.e. species that account for at least 85% of the cumulated species contribution of the community) measured in 2014, and (ii) leaf trait measurements made at plot scale in 2006 and 2007. Traits were measured on ten vegetative plants using standard protocols (see methods in Louault et al. 2005). Reproductive plant height was measured on mature plants located in fenced zones to allow full plant development. CWM is expressed with the following equation: $CWM = \sum p_i \times trait_i$, where p_i is the relative contribution of species i to the community and $trait_i$ the trait of species i.

Statistical analyses

For a given date, root mass and root traits collected at each location (four ingrowth-cores in each plot), averages of data coming from the four locations were used to have a single value for

each of the 12 plots and test for the effect of treatment and dates. Before ANOVA, normality of residuals was inspected with quantile-quantile plots of model residuals, and variance homogeneity was confirmed by checking the plots of model residuals *vs.* fitted values. Data were transformed if they deviated from ANOVA assumptions (square root, ln, reciprocal). Linear mixed effects models as available in the R 'nlme' package (Pinheiro et al. 2015) were used to perform repeated measure ANOVAs to test the effects of treatments, dates and their interactions on values of root growth, soil temperature, RSWC, and root mass stock, with plots nested in block as a random factor accounting for temporal pseudo-replication. For root growth dynamics, soil temperature and RSWC (Fig 1, Table S2), dates correspond to 20 dates and for root mass stock, dates correspond to three harvest dates (Table 2). For BNPP, ANPP and root to shoot ratio (BNPP/ANPP), data were analyzed using a nested mixed model procedure, with treatments and year used as fixed factors with plot nested in block as random factors. For leaf and root traits data, treatments were used as fixed factors with plots nested in block as a random factor. *Post hoc* tests were performed to compare significance levels across fixed factors with a Tukey test ('lsmeans' package). Principal component analyses (PCA) were performed for each year to analyze relationships between leaf and root traits, soil temperature, RSWC, root mass stock, ANPP and BNPP measured at plot level; treatments were considered as supplementary categories ('FactoMineR' package). This statistic approach allows comparing sets of traits and properties relationships in order to detect response and effect traits, but also to analyse multiple dimensions of traits relationships, not possible with pairs of correlation.

All statistical analyses were performed in the R environment (version 3.5.2, R Core team 2012) using RStudio (Version 1.1.463). Scripts are shown in Table S4.

Results

Climatic conditions during the experiment

Compared with average long-term climatic data for the site, the first and second years of the experiment had higher (+92 mm) and lower (-199 mm) precipitation, respectively (Table 1). Potential evapotranspiration (PET) in the second year was also higher than the long-term average (difference of 73 mm), leading to a negative annual climatic water balance (P - PET = -181 mm and a deficit of 271 mm compared to the long-term average). Annual temperature in the two experimental years was similar and about 0.8°C higher than the long-term average for the site (Table 1). At monthly time scale and during part of the growing season (March to September), in comparison with the first year, the second year had a cumulated water deficit difference of -266 mm and a temperature warmer by +1.9 °C than the first year. Larger differences between the two years occurred in June-July with higher temperature (+6 °C), higher water deficit (P-PET = -152.6 mm) and less precipitation (-81%) in the second year.

Dynamics of soil temperature and relative soil water content

Soil temperature was significantly affected by treatment, dates and treatment × dates (Figure 1; Table S2). For most of the dates (February to October), abandonment treatment had lower soil temperature (1.76 °C, on average) than the grazing treatments, whereas the Cattle- treatment showed significant lower soil temperature (-0.64 °C) than the Cattle+ treatment. However, this was significantly observed for a limited number of dates in early summer of both years. Relative soil water content (RSWC) fluctuated from 0.6-0.7 at the beginning of spring to 0.38 in June in the wet year and to 0.2 during the dry year, which is in accordance to variation of the atmospheric aridity index (P-PET). In the case of the dry year, from summer until autumn, RSWC remained lower than 0.4 and the aridity index was negative.

<u>Root growth dynamics</u>

Root growth was affected by date and treatment × date interaction (Figure 1). Each year, peak of root growth occurred twice, in spring and autumn, and growth was markedly reduced in summer and winter. Only in the second year did growth stop in summer, and it was significantly lower than the first year. Regarding treatment effect, abandonment showed significant lower root growth than the two grazing treatments for the spring period in both years, and for the autumn of the second year. While in autumn 2014, a delay of growth peaks was always observed, which led to a two-fold higher root growth for abandonment *vs.* the two cattle treatments (end of September: date 8). The two grazing treatments had similar root growth across years and seasons.

<u>Seasonal root mass stock, BNPP, ANPP and root-to-shoot biomass ratio</u>

Stock of root mass did not change through season and across treatment (Table 2). BNPP, ANPP and root-to-shoot biomass ratio (R/S) were significantly lower during the second year, with a stronger effect on BNPP (-44% on average) than ANPP (-24%) (Figure 2, Table 3). Only the abandonment treatment maintained their value of ANPP in the second year, which led to a 48% decline in R/S (significant treatment × year, $P < 0.01$, Table 3). Accordingly, treatment effect was only observed for BNPP the second year, with a decline of 24% for abandonment compared to cattle grazing treatments and for ANPP the first year: Cattle+ having 22% and 68% higher values than Cattle- and abandonment, respectively, while Cattle- had 38% higher ANPP than abandonment.

<u>Species composition, leaf and root traits</u>

Abandonment treatment was characterized by the dominance of tall grass species: 76% in all with 27.2% of *Alopecurus pratensis*, 18.8% of *Elytrigia repens*, 11.3% of *Poa pratensis* and

10.3% of *Arrhenatherum elatius*, the presence of some forbs (19%) and the absence of legumes

(Table S3 and Table 4). The two cattle treatments differed from abandonment treatment by

equal presence of *Taraxacum officinale* (18% on average) and *Trifolium repens* (17% on

average). Difference also concerns grass species (56% in total) with the dominance of *Dactylis*

glomerata (22.2%), *A. pratensis* (7.6%) and *Festuca arundinacea* (5.6%) for Cattle- and *Lolium*

perenne (13.6%), *D. glomerata* (9.1%) and *Poa trivialis* (7.2%) for Cattle+. Thus, the Cattle+

treatment had a higher percentage of *L. perenne* than Cattle- (Table S3).

Community-weighted mean leaf traits (CWM) were significantly modified by the

treatments. Plant height and LDMC were significantly higher (P < 0.05 and P < 0.0001,

respectively; Table 4) in abandonment than in the two cattle grazed treatments, whereas SLA

was lower (P < 0.05). Unlike leaf traits, root traits were only slightly affected by the treatments.

Specific root length (SRL, P < 0.1) and specific root area (SRA, P < 0.05) were lower in

abandonment treatment than in Cattle-, but not Cattle+. For other root traits (diameter, RTD

and root length % by class diameter) no between-treatment differences were observed (Table

4).

<u>Co-variation of traits and production</u>

The two main axes of the standardized PCA explained 60.1% and 56.8% of the community trait

and production variation in 2014 and 2015, respectively (Figure 3). For the first year, the first

PCA axis (PC1), accounting for 43.4% of the total variation, was significantly related to leaf

and root traits, ANPP and soil temperature. Soil temperature, SRA and ANPP had positive

loadings, and diameter, plant height and LDMC had negative loadings (Table 5). The second

PCA axis (PC2), accounting for 16.7% of the total variation, was significantly and positively

related to root diameter and negatively to SRA. For the second year, the first PCA axis (PC1),

accounted for 37.4% of the total variation, and was significantly related to leaf and root traits,

ANPP and BNPP. BNPP and SRA had negative loadings, and root diameter, plant height and ANPP had positive loadings (Table 5). The second PCA axis (PC2), accounting for 19.4% of the total variation, was significantly and positively related to RSWC and stock of root mass averaged across three dates. Finally, abandonment treatment was significantly related to PC1s with negative and positive loadings for the first and the second year, respectively.

Discussion

Ten years of contrasted management had strongly modified the functional diversity and above-ground production of this fertile upland grassland (Herfurth et al. 2015; Louault et al. 2017). Accordingly, we expected that above-ground biomass patterns would be mirrored below-ground, especially during the periods of grazing. Here we first discuss within-year differences of root growth, followed by inter-annual variation responses to grazing intensity and climatic conditions variability between the two contrasting years, and last we analyze relationships between traits and above- and below-ground production.

Seasonality of root growth was independent of grazing intensity and climatic conditions

As expected, root growth of permanent grassland is affected by seasons and peaks in spring and autumn (Garcia-Pausas et al. 2011; Pilon et al. 2013; Steinaker and Wilson 2008), but unexpectedly, grazing pressure applied by rotations and climatic conditions variability had very limited effects on this seasonality. This means that at below-ground level, plant community behavior was not affected by rotational grazing management nor by climatic conditions variability, although a severe drought occurred in summer of the second year. Only the abandonment treatment showed a delayed root growth peak in spring. This delay is probably the result of slower shoot budburst and reduced capacity to produce new green leaves in dense litter canopy, especially at the beginning of the growing season in spring (data not shown).

Moreover, the tall and dense canopy of the abandonment treatment strongly modified soil temperature, with cooler soil conditions as expected in such abandoned vegetation (Picon-Cochard et al. 2006; Zhou et al. 2017; Zhu et al. 2016). As shown in some studies, light or soil water and nutrient availabilities (Edwards et al. 2004; Garcia-Pausas et al. 2011; Steinaker and Wilson 2008) are other abiotic factors determining dynamics of root growth in grasslands, as root peaks were observed before the peak of soil temperature in summer when negative climatic water balance occurred, especially in the second year. Nevertheless, plants growing in abandonment offset their slower root growth by producing similar root biomass at annual scale, especially during the wet year. The presence of tall grass species such as *A. pratensis*, *A. elatius* and *E. repens* with plant trait syndromes related to both disturbance and resource conservation strategies (lower SLA and SRL and higher plant height and root depth; Pagès and Picon-Cochard 2014) might explain their capacity to produce higher root biomass on a shorter-term period before canopy senescence onset. Also pre-existing soil fertility can be maintained in conditions of very low levels of herbage utilization (near-abandonment), because of the absence of biomass exportation and increased internal recycling of N within senescent plants, both contributing to an increase in total N available for plant growth (Loiseau et al. 2005).

The similar root growth dynamics of the two cattle treatments was unexpected, considering that infrequent defoliation and moderate excreta returns to the soil might increase root biomass production at the expense of shoot biomass (Klumpp et al. 2009). The absence of effect on root growth and BNPP means that grazing applied on plant communities by rotations was too short but enough to observe effect on ANPP, in wet conditions. Worldwide there are different ways to manage grassland by grazing (Huyghe et al. 2014), rotational or permanent grazing options with different stocking densities, durations, types of herbivores. In general, this management creates high spatial heterogeneity within the plots due to animals' selective defoliation of plant species, and also because returns to soil are spatially heterogeneous. Thus in grazed grassland,

disturbance creates patches of vegetation, which should affect locally root growth and below-

ground biomass of plant communities if intensity of grazing is sufficient. The complexity of

these phenomena in grazed grassland is greater than in mown systems owing (Rossignol et al.

2011).

Then, again, the confounding effect of soil fertility and defoliation may mask a clear

response of the below-ground compartment in grazed grasslands. In view of that, we postulate

that root growth in Cattle+ treatment was favored by the higher soil temperature compensating

for the negative effects of frequent defoliation on root growth while the cooler soil conditions

encountered in Cattle- might have slowed root growth. Soil moisture is a main determinant of

plant growth and can be affected by cattle treatments. Some studies showed an increase in

grazed compared ungrazed treatment due to lower leaf area index in the grazed conditions

(Moretto et al. 2001; Pineiro et al. 2010), or an absence of effects in others (LeCain et al. 2002;

Smith et al. 2014). The presence of herbivores can increase soil bulk density and consequently

modify soil moisture. In our field conditions and after 10 years of treatments application, the

absence of effect on soil moisture can be due to several reasons. A first determinant to consider

is the soil density, which is expected to be higher in grazed plots, but in case of rotational

grazing this effect is less clear as cattle spend less time than in continuous grazing systems. A

second determinant is the functional composition of community regarding both the response to

defoliation and their water use strategies. A last determinant of this response can be linked to

the temporal scale used (monthly-based) which could buffer shorter-term response.

We should also consider the level of soil fertility and species composition as drivers of root

growth and trait plasticity (Dawson et al. 2000). The soil fertility of our site, reflected by the

nitrogen nutrition index (NNI, Lemaire and Gastal 1997), was very similar along our grazing

intensity gradient (Table S1), at least in 2014. Thus in our site we had the opportunity to

compare grazing intensity effect at equivalent soil fertility. Knowing that root trait plasticity

generally shows larger differences with respect to soil fertility than by cutting or defoliation

(Leuschner et al. 2013; Picon-Cochard et al. 2009), we can expect that under similar soil fertility

grazing intensity had a less pronounced effect on root growth. Indeed, the higher presence of

species tolerating defoliation, with shorter stature and root system (*L. perenne*, *P. trivialis*), but

having higher shoot and root growth capacity after defoliation and also higher rhizosphere

activity (Dawson et al. 2000), probably compensated for the negative effect of defoliation in

the Cattle+ treatment. Also the sampling depth might have had an effect, as we expect that

harvesting root systems deeper than 20 cm should give more contrasting root growth response

across the two cattle treatments according to the grass species composition due to species-

specific differential root depth distribution (Xu et al. 2014). Taken together, we provide

evidence that higher soil temperature, high soil fertility and species composition have

moderated root growth response along our grazing intensity gradient. The difficulty to assign

species composition in root mixtures, however, makes it difficult to draw firm conclusions.

<u>Climatic conditions variability shaped responses of ANPP, BNPP and root-to-shoot biomass</u>

<u>production ratio along the grazing intensity gradient</u>

According to meta-analyses and recent results (McSherry and Ritchie 2013; Zeng et al. 2015;

Zhou et al. 2017; Li et al. 2018), grazing intensity generally has negative effects on above- and

below-ground biomass of grasslands whatever the climatic conditions or vegetation type,

although these effects can be modulated by levels of grazing intensity. Our results do not

confirm these findings, because ANPP and BNPP increased in response to grazing intensity

compared to abandonment, in the wet and the dry year, respectively. Methodology issues for

estimating ANPP and BNPP in grazed grasslands should thus be taken into account, as some

papers report either biomass stock or fluxes measured once at peak of growth or at several

periods (Scurlock et al. 2002), but also estimation of BNPP from indirect measurements (e.g.

Zeng et al. 2015). Mass based on stock gives a snapshot of plant functioning, generally

including mixtures of living and senescent tissues, thus depending on abiotic factors and plant

growth, whereas measurements based on new shoot and root biomass reflect the growth

potential of grasslands. We are aware that these methods are very different, but in response to

grazing intensity, BNPP measured with ingrowth cores gave similar results as root mass stock

assessed at three seasons. Another point to consider is the number of samples used to compare

treatments and detect significant differences. In grasslands, the coefficient of variation of root

dry weight in auger samples from grassland are generally between 30 and 50% (Bengough et

al., 2000). According to these authors, our sampling protocol (with 16 samples) is adapted to

detect at least 35% differences between treatments, whereas to detect less than 10% differences,

more than 100 replicates should be collected. Thus, we cannot rule out that collecting more

samples should have highlighted significant differences across treatments. Nevertheless, we

had to find compromise between more frequent samplings (20 dates) to study seasonal

dynamics of root growth compared with more samples collected at plot level but less frequently.

Climatic aridity (P-PET) had stronger effects on ANPP and BNPP than grazing intensity,

because severe drought had a direct negative effect on plant growth. In comparison with another

experiments located alongside ours, 80% of canopy senescence was reached for a cumulated

aridity of -156 mm (Zwicke et al. 2013). As this index reached -303 mm from March to August,

this confirmed that a severe drought occurred in the second year of our experiment, and

explained root growth cessation in summer. At annual scale, ANPP of the two cattle treatments

showed lower resistance to increased aridity (resistance defined as $ANPP_{year2} / ANPP_{year1}$, being

equal to 0.63) than abandonment treatment (ratio=1). For BNPP, results were inversed, leading

to a lower resistance of root-to-shoot biomass ratio in abandonment than in the two cattle

treatments. The absence of root growth modification by grazing at annual scale the wet year

reflects well the change in root-to-shoot biomass allocation, albeit not significant. Other

processes such as root turnover (mortality, rhizodeposition) are expected to change in grazed

vs. ungrazed grassland. For our site Herfurth et al. (2015) observed similar root mass stock

along a grazing disturbance gradient as in the present study, but by using a simplified C flux

model, these authors showed that the Cattle+ treatment tended to accelerate C cycling in plant

communities, resulting in a higher quantity of C allocated to the soil organic matter continuum.

Taken together, these results suggest that the slight BNPP increase under grazing may occur

with an increase in rhizodeposition, because root turnover calculated as BNPP to root mass

stock ratio (data not shown, Lauenroth and Gill 2003) was not different across treatments.

Furthermore, our results suggest that grazing treatments slow down the negative effect of

aridity on the root-to-shoot biomass ratio, underling that these treatments seem to be better

adapted to buffering the negative effect of drought on grassland root production than for

abandoned grasslands. This is consistent with previous work showing that moderate grazing

could be more beneficial than no grazing for drought resistance and recovery of ANPP and

BNPP (Frank 2007; Xu et al. 2012), and that BNPP was more resistant than ANPP to change

in precipitation (Yan et al. 2013). Other studies showed no prevalence effects of grazing,

drought or fire observed on grassland production in North America and South Africa (Koerner

and Collins 2014). Nevertheless, this points to a need for further research to determine whether

grazing pressure has additive or combined effects on drought response of grasslands (Ruppert

et al. 2015).

<u>Community-weighted mean leaf and root traits as predictors of ANPP and BNPP</u>

As shown by other studies (e.g. Diaz et al. 2007; Laliberté and Tylianakis 2012; Louault et al.

2017; Zheng et al. 2015), disturbance induced by grazing pressure has profound effects on plant

community and functional traits by selecting tolerant species to defoliation such as *L. perenne*,

P. trivialis or *T. repens*, with possible cascading effects on multiple ecosystem functions. With

the capacity to regrow quickly after defoliation, these species generally exhibited high values

of SLA and low values of LDMC and plant height. They contrast with species adapted to fertile

soil, but with a slower regrowth capacity after defoliation such as *D. glomerata* or *F.*

arundinacea, with opposite leaf trait values. In abandonment, competition for light tends to

select plants with trait syndromes related to disturbance and conservative strategies (tall plants,

low SLA and high LDMC values). Thus, the CWM traits of the community will depend on the

balance between these species groups, which are expected to affect ANPP and BNPP (Klumpp

et al. 2009; Milchunas and Lauenroth 1993). Although the presence of tolerant and intolerant

species to defoliation in both cattle treatments, leaf trait values were similarly and positively

related to ANPP, and only differed from traits of species present in the abandonment treatment.

This means that cessation of grazing strongly differentiated plant communities, whereas within

the two cattle treatments differences were slighter.

For the below-ground compartment, we expected that above-ground differences were

mirrored by the root growth and traits, assuming that higher root diameter values, and lower

SRL and SRA values are associated with lower BNPP in abandonment compared with the two

cattle treatments. Although root response to grazing (mainly through defoliation) generally

reported reduction of root mass or root length (Dawson et al. 2000) our study did not confirm

these assumptions. The contrasting results are possibly due to variable abundance of tolerant

species to defoliation or with confounding effects of both defoliation and level of soil fertility

on roots of grazed grasslands (Leuschner et al. 2013; Picon-Cochard et al. 2009; Yan et al.

2013; Ziter and McDougall 2013). Thus, root growth reductions associated with grazing may

have a greater impact in locations where grazer-mediated nitrogen return is spatially decoupled

from defoliation (McInenly et al. 2010). Further, higher specific root area (SRA) observed in

Cattle- than in abandonment and Cattle+ treatments should reflect higher presence of species

with fine roots such as *D. glomerata* or *H. lanatus* (Picon-Cochard et al. 2012), because soil

fertility approximated by NNI was near comparable across treatments.

Conclusions

Near similar functional traits of the plant communities and similar soil fertility across the two

cattle treatments explained the absence of changes in root mass production for these treatments.

Our site disentangled confounding effects of fertility and defoliation on root production, which

is not generally the case for other studies. Thus, our results suggest the prevalence of a soil

fertility effect on root production response rather than a defoliation effect. Besides, the strong

effect of climatic conditions variability on ANPP and BNPP observed at short term could

increase in the future as more frequent climatic extremes are expected. It is thus necessary to

improve our knowledge at larger time scale on the grazing practices allowing higher resilience

of grasslands to more frequent and intense climatic events such as drought and heat waves.

Data accessibility

Data are available online: https://zenodo.org/record/4034903#.YA129-fjJPZ
538

Acknowledgments

We thank staff from INRAE UREP: V. Guillot and E. Viallard for their technical expertise in

field measurements, D. Colosse and S. Toillon for the soil temperature database, and S.

Revaillot, A. Bartout, L. Bulon and S. Sauvat and M Mattei (VetAgro Sup) for their help in root

sample measurements, and the staff of INRAE UE1414 Herbipôle. The experiment is part of

the SOERE-ACBB project (http://www.soere-acbb.com/) funded by Allenvi and the French

National Infrastructure AnaEE-F through ANR-11-INBS-0001. Data of the weather station are

coming from the platform INRAE CLIMATIK (https://intranet.inrae.fr/climatik/, in French)

managed by the AgroClim laboratory of Avignon, France. DH received a doctoral fellowship

from VetAgro Sup and DGER pole "ESTIVE". The present work falls within the thematic area

of the French government IDEX-ISITE initiative 16-IDEX-0001 (CAP 20-25).

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the

content of this article

References

Bardgett RD, Wardle DA (2003) Herbivore-mediated linkages between aboveground and

belowground communities. Ecology 84:2258-2268

Bengough AG, Castrignano A, Pagès L, van Noordwijk M. 2000. Sampling strategies, scaling,

and statistics. In: Smit AL, Bengough AG, Engels C, van Noordwijk M, Pellerin S, van

de Geijn SC, eds. *Root Methods.* Berlin, Germany: Springer, 147-173.

Biswell H, Weaver JE (1933) Effect of frequent clipping on the development of roots and tops

of grasses in prairie sod. Ecology 14:368-390

Brookshire ENJ, Weaver T (2015) Long-term decline in grassland productivity driven by

increasing dryness. Nature Commun 6:7148

Byrne KM, Lauenroth WK, Adler PB (2013) Contrasting Effects of Precipitation Manipulations

on Production in Two Sites within the Central Grassland Region, USA. Ecosystems

16:1039-1051

Conant RT, Paustian K, Elliott ET (2001) Grassland management and conversion into

grassland: effects on soil carbon. Ecol App 11:343-355

Dawson LA, Grayston SJ, Paterson E (2000) Effects of grazing on the roots and rhizosphere of

grasses. In: Lemaire G, Hodgson J, de Moraes A, Nabinger C, De F. Carvalho PC (eds)

Grassland ecophysiology and grazing ecology. CABI Publishing, Wallingford, pp 61–84

Diaz S, Lavorel S, McIntyre S, Falczuk V, Casanoves F, et al. (2007) Plant trait responses to

grazing-a global synthesis. Glob Chang Biol 13:313-341

Duru M, Balent G, Gibon A, Magda D, Theau JP, Cruz P, Jouany C (1998) Fonctionnement et

dynamique des prairies permanentes. Exemple des Pyrénées centrales. Fourrages 153:97-

113

Edwards EJ, Benham DG, Marland LA, Fitter AH (2004) Root production is determined by

radiation flux in a temperate grassland community. Glob Chang Biol 10**:**209-227

Frank DA (2007) Drought effects on above-and belowground production of a grazed temperate

grassland ecosystem. Oecologia 152:131–139

Garcia-Pausas J, Casals P, Romanyà J, Vallecillo S, Sebastià M-T (2011) Seasonal patterns of

belowground biomass and productivity in mountain grasslands in the Pyrenees. Plant Soil

340:315–326

Herfurth D, Vassal N, Louault F, Alvarez G, Pottier J, Picon-Cochard C, Bosio I, Carrère P

(2015) How does soil particulate organic carbon respond to grazing intensity in

permanent grasslands? Plant Soil 394:239-255

Huyghe C, De Vliegher A, van Gils B, Petters A (2014) Grasslands and herbivore production

in Europe and effects of common policies. In, Huyghe C, De Vliegher A, van Gils B,

Petters A eds, Quæ editions, Versailles, France, 287pp.

Jones MB, Donnelly A (2004) Carbon sequestration in temperate grassland ecosystems and the

influence of management, climate and elevated CO_2. New Phytol 164:423-439

Klumpp K, Fontaine S, Attard E, Le Roux X, Gleixner G, Soussana JF (2009) Grazing triggers

soil carbon loss by altering plant roots and their control on soil microbial community. J

Ecol 97:876–885.

Klumpp K, Soussana J-F (2009) Using functional traits to predict grassland ecosystem change:

a mathematical test of the response-and-effect trait approach. Glob Chang Biol 15:2921-

2934

Koerner SE, Collins SL (2014) Interactive effects of grazing, drought, and fire on grassland

plant communities in North America and South Africa. Ecology 95:98-109

Laliberté E, Tylianakis JM (2012) Cascading effects of long-term land-use changes on plant

traits and ecosystem functioning. Ecology 93:145–155

Lauenroth WK, Gill R (2003) Turnover of root systems. In: de Kroon H, Visser EJW (eds) Root

Ecology, Springer-Verlag, Berlin, pp 61-83

LeCain DR, Morgan JA, Schuman GE, Reeder JD, Hart RH (2002) Carbon exchange and

species composition of grazed pastures and exclosures in the shortgrass steppe of

Colorado. Agriculture, Ecosystems & Environment 93(1): 421-435.

Lemaire G, Gastal F (1997) N uptake and distribution on plant canopy. In: Lemaire, G (ed)

Diagnosis of the nitrogen status in crops, Springer-Verlag, Berlin, pp 3-43

Leuschner C, Gebel S, Rose L (2013) Root trait responses of six temperate grassland species to

intensive mowing and NPK fertilisation: a field study in a temperate grassland. Plant Soil

373:687-698

Li, W., X. Li, Y. Zhao, S. Zheng, and Y. Bai. 2018. Ecosystem structure, functioning and

stability under climate change and grazing in grasslands: current status and future

prospects. Curr Opin Environ Sustain 33:124-135.

Loiseau P, Louault F, Le Roux X, Bardy M (2005) Does extensification of rich grasslands alter

the C and N cycles, directly or via species composition? Basic App Ecol 6:275-287

Louault F, Pillar VD, Aufrere J, Garnier E, Soussana JF (2005) Plant traits and function types

in response to reduced disturbance in a semi-natural grassland. J Veg Sci 16:151–160

Louault F, Pottier J, Note P, Vile D, Soussana JF, Carrère P (2017) Complex plant community

responses to modifications of disturbance and nutrients availability in productive

grasslands. J Veg Sci, in press

McInenly LE, Merrill EH, Cahill JF, Juma NG (2010) Festuca campestris alters root

morphology and growth in response to simulated grazing and nitrogen form. Funct Ecol

24:283-292

McSherry ME, Ritchie ME (2013) Effects of grazing on grassland soil carbon: a global review.

Glob Chang Biol 19:1347–1357

Milchunas DG, Lauenroth WK (1993) Quantitative effects of grazing on vegetation and soils

over a global range of environments. Ecol Mono 63:327-366

Moretto AS, Distel RA, Didoné NG (2001) Decomposition and nutrient dynamic of leaf litter

and roots from palatable and unpalatable grasses in a semi-arid grassland. App Soil Ecol

18: 31–37

Pagès L, Picon-Cochard C (2014) Modelling the root system architecture of Poaceae. Can we

simulate integrated traits from morphological parameters of growth and branching? New

Phytol 204:149-158

Pérez-Ramos IM, Roumet C, Cruz P, Blanchard A, Autran P, Garnier E (2012) Evidence for a

'plant community economics spectrum' driven by nutrient and water limitations in a

Mediterranean rangeland of southern France. J Ecol 100:1315-1327

Picon-Cochard C, Coll L, Balandier P (2006) The role of below-ground competition during

early stages of secondary succession: the case of three-year-old Scots pine (*Pinus*

sylvestris L.) seedlings in an abandoned grassland. Oecologia 148:373-383

Picon-Cochard C, Pilon R, Revaillot S (2009) Plasticity of grass root functional traits and root

mass in response to cutting frequency and N fertilisation. Proceedings of the 7th ISRR

Symposium, Root Research and Applications (RootRAP), Vienne September 2–4, 2009. 4pp

Picon-Cochard C, Pilon R, Tarroux E, Pagès L, Robertson J, Dawson L (2012) Effects of species, root branching order and season on root traits of 13 perennial grass species. Plant Soil 353:47-57

Pilon R, Picon-Cochard C, Bloor JMG, Revaillot S, Kuhn E, Falcimagne R, Balandier P, Soussana J-F (2013) Grassland root demography responses to multiple climate change drivers depend on root morphology. Plant Soil 364:395-408

Pineiro G, Paruelo JM, Oesterheld M, Jobbagy EG (2010) Pathways of grazing effects on soil organic carbon and nitrogen. Rangeland Ecol Manage 63:109–119

Pinheiro J, Bates D, Debroy S, Sarkar D and R Core Team (2015) nlme: linear and non linear mixed effect models. R Package Version 3.1-119

R Core Team (2012) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Reich PB (2014) The world-wide 'fast–slow' plant economics spectrum: a traits manifesto. J Ecol 102:275-301

Rossignol N, Chadoeuf J, Carrère P, Dumont B (2011) A hierarchical model for analysing the stability of vegetation patterns created by grazing in temperate pastures. App Veg Sci 14:189-199.

Ruppert JC, Harmoney K, Henkin Z, Snyman HA, Sternberg M, Willms W, Linstädter A (2015) Quantifying drylands' drought resistance and recovery: the importance of drought intensity, dominant life history and grazing regime. Glob Chang Biol 21:1258-1270

Schaffers A, Sykora K (2000) Reliability of Ellenberg indicator value for moisture, nitrogen and soil reaction: a comparison with field measurements. J Veg Sci 11:225-244

Scurlock JMO, Johnson K, Olson RJ (2002) Estimating net primary productivity from grassland

biomass dynamics measurements. Glob Chang Biol 8:736-753

Smith SW, Woodin SJ, Pakeman RJ, Johnson D, van der Wal R (2014) Root traits predict

decomposition across a landscape-scale grazing experiment. New Phytol 203:851-862.

Soussana JF, Loiseau P, Vuichard N, Ceschia E, Balesdent J, Chevallier T, Arrouays D (2004)

Carbon cycling and sequestration opportunities in temperate grasslands. Soil Use Manag

20:219-230

Soussana JF, Duru M (2007) Grassland science in Europe facing new challenges: biodiversity

and global environmental change. CAB Reviews: Perspectives in Agriculture, Veterinary

Science, Nutrition and Natural Resources 272:1-12

Soussana JF, Lemaire G (2014) Coupling carbon and nitrogen cycles for environmentally

sustainable intensification of grasslands and crop-livestock systems. Agr Ecosyst Environ

190:9-17

Steinaker DF, Wilson SD (2008) Phenology of fine roots and leaves in forest and grassland. J

Ecol 96:1222-1229

Wardle DA, Bardgett RD, Klironomos JN, Setala H, van der Putten WH, Wall DH (2004)

Ecological linkages between aboveground and belowground biota. Science 304:1629-

1633

Wilcox KR, von Fischer JC, Muscha JM, Petersen MK, Knapp AK (2015) Contrasting above-

and belowground sensitivity of three Great Plains grasslands to altered rainfall regimes.

Glob Chang Biol 21:335-344

Xu X, Sherry RA, Niu S, Li D, Luo Y (2012) Net primary productivity and rain-use efficiency

as affected by warming, altered precipitation, and clipping in a mixed-grass prairie. Glob

Chang Biol 19:2753-2764

Xu X, Luo Y, Shi Z, Zhou X, and Li D (2014) Consistent proportional increments in responses
of belowground net primary productivity to long-term warming and clipping at various
soil depths in a tallgrass prairie. Oecologia 174, 1045-1054

Yan L, Zhou G, Zhang F (2013) Effects of different grazing intensities on grassland production
in China: A meta-analysis. PLoS ONE 8:e81466

Zeng C, Wu J, Zhang X (2015) Effects of grazing on above- vs. below-ground biomass
allocation of alpine grasslands on the northern tibetan plateau. PLoS ONE 10:e0135173

Zheng S, Li W, Lan Z, Ren H, Wang K (2015) Functional trait responses to grazing are
mediated by soil moisture and plant functional group identity. Sci Rep 5:18163

Zhou G, Zhou X, He Y, Shao J, Hu Z, Liu R, Zhou H, Hosseinibai S (2017) Grazing intensity
significantly affects belowground carbon and nitrogen cycling in grassland ecosystems:
a meta-analysis. Glob Chang Biol 23:1167-1179

Zhu J, Zhang Y, Liu Y (2016) Effects of short-term grazing exclusion on plant phenology and
reproductive succession in a Tibetan alpine meadow. Sci Rep 6:27781

Ziter C, MacDougall AS (2013) Nutrients and defoliation increase soil carbon inputs in
grassland. Ecology 94:106-116

Zwicke M, Alessio GA, Thiery L, Falcimagne R, Baumont R, Rossignol N, Soussana J-F,
Picon-Cochard C (2013) Lasting effects of climate disturbance on perennial grassland
above-ground biomass production under two cutting frequencies. Glob Chang Biol
19:3435–3448.

Table 1. Air temperature (°C), precipitation (P, mm), potential evapotranspiration (PET) and

climatic water balance: cumulated (P - PET, mm) and calculated for the 28 y period 1986-2013,

mean values ± SD) and measured for the 10 dates in 2014 and 2015 corresponding to

measurements of root growth and averaged (temperature) or summed (P, PET, P-PET) at annual

scale.

Year	Dates	Air temperature	Precipitation	PET	P-PET
	Annual long-term average	**8.5 ± 0.6**	**784 ± 1376**	**693 ± 96**	**91 ± 195**
2014	December 12 – February 23	3.7	98	37.5	60.5
	February 24 – March 23	5.3	27	46.3	-19.3
	March 24 – April 21	7.2	23.5	68.7	-45.2
	April 22 – May 25	9.2	79.5	103.1	-23.6
	May 26 – June 22	14.2	58	110.2	-52.2
	June 23 –July 20	15.1	136.5	93.9	42.6
	July 21 – August 24	14.4	90.5	100.5	-10
	August 25 – September 29	13.7	141.8	79.5	62.3
	September 30 – October 29	11.7	69	36.3	32.7
	October 30 – December 14	5.3	111	10.9	72.1
	Annual	**9.2**	**876**	**691**	**157.7**
2015	December 15 – March 1	1.3	132.5	31	101.5
	March 2 – March 29	4.5	36.5	36.8	-0.3
	March 30 – April 23	8.5	17.5	66.4	-48.9
	April 24 – May 28	11.0	66	113.6	-47.6
	May 29 – June 28	15.5	62.5	129.1	-66.6
	June 29 –July 23	21.1	26	136	-110
	July 24 – August 27	16.4	94.5	124.6	-30.1
	August 28 – September 24	12.8	77	66.3	10.7
	September 25 – October 29	7.8	55	36.1	18.9
	October 30 – December 11	7.0	54.5	25.1	29.4
	Annual	**9.4**	**585**	**766**	**-180.9**

Table 2. a) Repeated measure ANOVA is shown for treatment, date (December 2013, March 2014, June 2014) and interaction effects on root mass (g m^{-2}). Numerator (num), denominator (den) of degree of freedom (DF) and F values are shown. b) Root mass (g m^{-2}) of abandonment, low (Cattle-) and high (Cattle+) stocking density treatments measured in winter (December 12 2013), spring (March 20 2014), summer (June 20 2014) and averaged across the three dates. Means ± se are shown, n = 4. Superscripts [ns] correspond to P > 0.05.

a)	num/den DF	*F-value*	
Treatment	2/8	1.151[ns]	
Date	2/18	2.027[ns]	
Treatment × date	4/18	1.340[ns]	
b) Date	Abandonment	Cattle-	Cattle+
December 2013	636.4 ± 133.1	403.3 ± 66.4	496.5 ± 20.6
March 2014	559.1 ± 166.2	609.2 ± 45.3	719.8 ± 47.5
June 2014	574.2 ± 84.8	482.2 ± 38.6	591.2 ± 101.7
3 dates average	589.9 ± 99.9	498.2 ± 43.6	602.5 ± 44.4

744 Table 3. Repeated measure ANOVA is shown for treatment, year and interaction effects on

745 annual root production (BNPP, g m^{-2} y^{-1}), annual above-ground production (ANPP, g m^{-2} y^{-1})

746 and root to shoot ratio (R/S). Numerator (num), denominator (den) of degree of freedom (DF),

747 *F* values are shown. Superscripts [ns, *, **, ***] correspond to P > 0.05, P < 0.05, P < 0.01, P < 0.001,

748 respectively.

		BNPP	ANPP	R/S
	num/den DF	*F-value*	*F-value*	*F-value*
Treatment	2/8	2.51^{ns}	8.10*	0.46^{ns}
Year	1/9	70.72***	83.77***	13.09**
Treatment × Year	2/9	3.83^{ns}	22.21**	9.52**

Table 4. Root traits measured from ingrowth core collected in June 2014 and leaf traits measured from botanical observation in abandonment (May 2014), Cattle- and Cattle+ (April 2014) treatments. Diameter: root diameter (mm); SRL: specific root length (m g^{-1}); RTD: root tissue density (g cm^{-3}); SRA: specific root area (m^2 g^{-1}); % 0-0.1 mm: percentage of length in the class diameter 0-0.1 mm; % 0.1-0.2 mm: percentage of length in the class diameter 0.1-0.2 mm; % 0.2-0.3 mm: percentage of length in the class diameter 0.2-0.3 mm; % > 0.3 mm: percentage of length in the class diameter > 0.3 mm; Community-weighted mean (CWM) Height: plant height (cm); SLA: specific leaf area (cm^2 g^{-1}); LDMC: leaf dry matter content (g g^{-1}). Means ± se are shown (n = 4). num/den DF: numerator and denominator of degree of freedom. Superscripts $^{ns, +, *, **, ***}$ correspond to $P > 0.1$, $P \leq 0.1$, $P < 0.05$, $P < 0.01$, $P < 0.001$, respectively. For SRL and SRA, different letters correspond to significant differences between treatments.

	num/den DF	F-value	Abandonment	Cattle-	Cattle+
Root traits					
Diameter	2/8	1.61^{ns}	0.240 ± 0.015	0.210 ± 0.006	0.222 ± 0.015
SRL	2/8	3.71^{+}	237.2 ± 26.3 b	332.7 ± 30.4 a	277.8 ± 23.8 ab
RTD	2/8	0.55^{ns}	0.099 ± 0.007	0.095 ± 0.003	0.102 ± 0.007
SRA	2/8	4.96*	0.137 ± 0.011 b	0.182 ± 0.008 a	0.155 ± 0.01 ab
% 0-0.1 mm	2/8	1.28^{ns}	28.5 ± 1.1	32.9 ± 5.5	28.8 ± 2.6
% 0.1-0.2 mm	2/8	0.46^{ns}	37.7 ± 4.4	37.7 ± 2.2	39.1 ± 1.8
% 0.2-0.3 mm	2/8	0.30^{ns}	16.6 ± 1.2	16.2 ± 2.4	17.1 ± 1.9
% > 0.3 mm	2/8	1.22^{ns}	17.2 ± 5.0	13.2 ± 1.3	15.1 ± 2.1
Leaf traits					
CWM_Height	2/8	8.45*	93.0 ± 3.5 a	72.8 ± 7.0 b	68.6 ± 3.8 b
CWM_SLA	2/8	5.30*	205.1 ± 5.7 b	231.8 ± 7.3 a	225.5 ± 7.1 ab
CWM_LDMC	2/8	11.22*	0.261 ± 0.008 a	0.227 ± 0.007 b	0.213 ± 0.010 b

Table 5. Contribution of the different variables to the first two axes of the principal component analysis (PCA) calculated for 2014 and 2015. Variables used in the PCA were annual relative soil water content (RSWC), annual soil temperature (Tsoil, °C), root diameter (Diam, mm), specific root area (SRA, $m^2\,g^{-1}$), root mass averaged over three dates (RootMass, $g\,m^{-2}$), annual root production (BNPP, $g\,m^{-2}\,y^{-1}$), plant height (Height, cm), leaf dry matter content (LDMC, $g\,g^{-1}$), annual above-ground production (ANPP, $g\,m^{-2}\,y^{-1}$). Treatments were added as supplementary categories.

Variable	2014		2015	
	Axis 1 (43.4 %)	Axis 2 (16.7 %)	Axis 1 (37.4 %)	Axis 2 (19.4 %)
RSWC	0.62	0.44	-0.21	**0.64**
Tsoil	**0.91**	0.09	-0.58	0.52
Diam	**-0.64**	**0.75**	**0.78**	0.53
SRA	**0.62**	**-0.58**	**-0.69**	-0.48
RootMass	-0.06	0.22	-0.07	**0.60**
BNPP	0.21	-0.23	**-0.71**	0.35
Height	**-0.82**	-0.07	**0.83**	-0.19
LDMC	**-0.83**	-0.12	**0.61**	0.03
ANPP	**0.71**	0.54	**0.57**	0.20
Suppl. Categories				
Abandonment	**-2.62**	-0.24	**2.04**	-0.27
Cattle-	1.07	-0.55	-1.21	-0.62
Cattle+	0.70	0.18	-0.83	0.90

Contribution in bold indicates significant correlation of the variables on the PCA axis ($P < 0.05$).

Figure captions

Figure 1. Dynamics of root growth (g m^{-2} day^{-1}), soil temperature (°C), relative soil water

content and an aridity index (P-PET, mm) (hashed bars), measured over two years for

abandonment, low (Cattle-) and high (Cattle+) stocking density treatments. Vertical bars

correspond to 1 se (n = 4). Insets indicate P values from repeated measure two-tailed ANOVA

(Treat: treatment, dates and interaction for main treatments). *: $P < 0.05$; x: $P \leq 0.1$. For soil

temperature, *# corresponds to significant differences between all treatments (Abandonment <

Cattle- < Cattle+).

Figure 2. Annual root biomass production (BNPP, g m^{-2} y^{-1}), annual above-ground biomass

production (ANPP, g m^{-2} y^{-1}) and root-to-shoot biomass ratio measured in 2014 and 2015 for

abandonment, low (Cattle-) and high (Cattle+) grazing intensiy treatments. Vertical bars

correspond to 1 se (n = 4). Within a year, different letters correspond to significant differences

at $P < 0.05$.

Figure 3. Principal component analysis (PCA) combining leaf and root traits, above- and below-

ground net primary production, root mass stock, relative soil water content and soil temperature

measured in 2014 (a) and 2015 (b) for abandonment, low (Cattle-) and high (Cattle+) stocking

density treatments. Data of each plot were used in each PCA. The first two axes are shown.

Arrows show projections of the variables within the PCA. RSWC: relative soil water content;

Tsoil: soil temperature (°C), Diam: root diameter (mm), SRA: specific root area (m^2 g^{-1}),

RootMass: root mass averaged over 3 dates (g m^{-2}), BNPP: annual root production (g m^{-2} y^{-1}),

Height: plant height (cm), LDMC: leaf dry matter content (g g^{-1}) and ANPP: annual above-

ground production (g m^{-2} y^{-1}).

Figure 1

Figure 2

Figure 3

Table S1. Nitrogen nutrition index (NNI %, Lemaire and Gastal 1997, Cruz et al. 2006)

measured on forage regrowth of May in 2014 and 2015 on the non-leguminous part to assess

the effect of treatments on N availability according to grazing intensity. When legumes were

below 4.5% in the herbage mass, NNI was assessed using the procedure defined by Cruz et al

(2006) based on the total forage and the legume contribution. The P-values are associated with

a nested mixed model: treatment used as fixed factor with plots nested in blocks as random

factors. Mean ± se is shown (n = 4). For each year, different letters correspond to significant

differences at $P < 0.05$.

Year	P-value	Abandonment	Cattle-	Cattle+
2014	0.146	65.64 ± 3.10 a	59.54 ± 1.78 a	63.72 ± 2.86 a
2015	0.018	69.72 ± 1.19 a	61.71 ± 1.53 b	69.25 ± 2.09 a

For each year, different letters correspond to significant differences at *: $P < 0.05$; **: $P < 0.01$;

***: $P < 0.001$; ns: $P > 0.05$.

References

Lemaire G, Gastal F (1997) N uptake and distribution on plant canopy. In: Lemaire, G (ed.)

Diagnosis of the nitrogen status in crops, pp. 3-43. Springer-Verlag, Berlin, DE.

Cruz P, Jouany C, Theau J-P, Petibon P, Lecloux E, Duru M (2006) L'utilisation de l'indice de

nutrition azotée en prairies naturelles avec présence de légumineuses. Fourrages 187:369-376.

Table S2. Repeated measure ANOVA is shown for root growth (g m^{-2} day^{-1}), soil temperature

(Tsoil, °C) and relative soil water content (RSWC) responses to treatment, dates (d1 to d20)

and interaction effects. Numerator (num), denominator (den) of degree of freedom (DF) and F

values are shown. Superscripts $^{ns,\ **,\ ***}$ correspond to P > 0.05, P < 0.001, P < 0.0001,

respectively.

Variables	Treatment		Dates		Treat. x Dates	
	num/den DF	F-value	num/den DF	F-value	num/den DF	F-value
Root growth	2/8	1.80^{ns}	19/171	50.40^{***}	38/171	2.096^{**}
Tsoil	2/8	33.93^{***}	19/166	944.83^{***}	38/166	9.75^{***}
RSWC	2/8	1.914^{ns}	19/163	25.287^{***}	38/163	1.097^{ns}

Table S3. Species contribution (%) in the community present around the ingrowth core measured in April and May 2014 for Cattle-, Cattle+ and Abandonment, respectively. Mean ± se is shown (n = 4). For each species, different letters correspond to significant differences at *: P < 0.05; **: P < 0.01; ***: P < 0.001; ns: P > 0.05.

Group	Species	*P-value*	Abandonment	Cattle-	Cattle+
Grasses	*Agrostis capillaris*	ns	0.0 ± 0.0	0.6 ± 0.6	1.7 ± 1.2
	Arrhenatherum elatius	ns	10.3 ± 6.8	2.2 ± 2.2	2.5 ± 2.5
	Alopecurus pratensis	**	27.2 ± 7.9 a	7.8 ± 3.3 b	3.3 ± 1.7 b
	Dactylis glomerata	*	3.1 ± 2.7 b	22.2 ± 9.8 a	9.1 ± 3.8 ab
	Elytrigia repens	*	18.8 ± 9.9 a	2.8 ± 1.8 b	3.8 ± 2.7 b
	Festuca arundinacea[x]	ns	5.0 ± 2.3	5.6 ± 2.1	6.3 ± 2.2
	Holcus lanatus	*	0.0 ± 0.0 b	4.7 ± 1.6 a	3.4 ± 1.9 a
	Lolium perenne	***	0.0 ± 0.0 b	0.9 ± 0.9 b	13.6 ± 3.8 a
	Poa pratensis	ns	11.3 ± 2.2	3.1 ± 1.5	3.4 ± 2.5
	Poa trivialis	*	0.0 ± 0.0 b	5.0 ± 2.5 a	7.2 ± 2.4 a
	Trisetum flavescens	ns	0.0 ± 0.0	2.2 ± 1.3	0.6 ± 0.4
Forbs	*Achillea millefolium*	ns	1.3 ± 0.9	3.8 ± 2.4	3.1 ± 2.3
	Anthriscus sylvestris	ns	2.5 ± 2.1	0.0 ± 0.0	0.0 ± 0.0
	Cerastium fontanum	ns	0.0 ± 0.0	1.3 ± 0.9	0.0 ± 0.0
	Cerastium glomeratum	ns	0.0 ± 0.0	0.0 ± 0.0	0.3 ± 0.3
	Cirsium arvense	ns	5.0 ± 3.5	0.0 ± 0.0	0.0 ± 0.0
	Hypocheris radicata	ns	0.0 ± 0.0	0.9 ± 0.9	0.0 ± 0.0
	Ranunculus acris	ns	0.0 ± 0.0	0.0 ± 0.0	3.8 ± 3.8
	Stellaria graminea	ns	0.6 ± 0.6	0.6 ± 0.4	0.0 ± 0.0
	Taraxacum officinale agg.	**	0.0 ± 0.0 b	17.5 ± 1.8 a	19.1 ± 6.0 a
	Urtica dioïca	*	9.7 ± 4.9 a	0.0 ± 0.0 b	0.0 ± 0.0 b
	Veronica serpyllifolia	ns	0.0 ± 0.0	0.3 ± 0.3	0.0 ± 0.0
Legumes	*Lathyrus pratensis*	ns	0.0 ± 0.0	0.3 ± 0.3	0.3 ± 0.3
	Trifolium pratense	ns	0.0 ± 0.0	0.0 ± 0.0	0.3 ± 0.3
	Trifolium repens	***	0.0 ± 0.0 b	16.3 ± 4.0 a	17.7 ± 2.5 a

[x]: new species name: *Schedonorus arundinaceus*

879 S4: R scripts used in the paper
880

881 1) Root growth, soil temperature and relative soil water content
882
883
884

```
> lme1 <- lme (variable ~ trait*date, random = ~ 1 | bloc/ID,
+                    data=IGC_ID, method="ML")
> anova(lme1)
```

```
plot(lme1)
> par(mfrow=c(2,2))
> plot(residuals(lme1))
> qqPlot(residuals(lme1))
> hist(residuals(lme1))
```

891
892
893

```
> lme1.lsmeans <-lsmeans(lme1, pairwise ~  trait | date)
> print(lme1.lsmeans[[2]])
```

895 2) Root mass stock measured at three dates
896
897
898

```
> lme1 <- lme (StockRac ~ trait*date, random = ~ 1 | bloc/ID,
+                 data=Stock_ID, method="ML")
> anova(lme1)
```

```
plot(lme1)
> par(mfrow=c(2,2))
> plot(residuals(lme1))
> qqPlot(residuals(lme1))
> hist(residuals(lme1))
```

905
906
907

```
> lme1.lsmeans <-lsmeans(lme1, pairwise ~  trait | date)
> print(lme1.lsmeans[[2]])
```

909 3) Root mass stock: averaged of the three dates, root and leaf traits
910
911
912

```
> lme1 <- lme (variable ~ trait, random = ~ 1 | bloc/ID,
+                    data=Stock3, method="ML")
> anova(lme1)
```

```
plot(lme1)
> par(mfrow=c(2,2))
> plot(residuals(lme1))
> qqPlot(residuals(lme1))
> hist(residuals(lme1))
```

919
920
921

```
> lme1.lsmeans <-lsmeans(lme1, pairwise ~  trait)
> print(lme1.lsmeans[[2]])
```

923 4) BNPP, ANPP, Root shoot mass ratio, NNI
924
925
926
927

```
> lme1 <- lme (variable ~ trait*year, random = ~ 1 | bloc/ID,
+                    data=BNPP, method="ML")
> anova(lme1)
```

```
plot(lme1)
> par(mfrow=c(2,2))
> plot(residuals(lme1))
> qqPlot(residuals(lme1))
> hist(residuals(lme1))
```

934
935
936

```
> lme1.lsmeans <-lsmeans(lme1, pairwise ~  trait | year)
> print(lme1.lsmeans[[2]])
```

937
938

941 Fig S1 : Scheme of the plots and blocks on the experimental site