In-situ observation of mitochondrial biogenesis as the early event of apoptosis

Chang-Sheng Shao [1,2], Xiu-Hong Zhou [3], Yu-Hui Miao [4], Peng Wang[1], Qian-Qian Zhang [1,2] and Qing Huang [1,2] *

[1] CAS Key Laboratory of High Magnetic Field and Ion Beam Physical Biology, Hefei Institutes of Physical Science, Chinese Academy of Sciences (CAS), Hefei 230031, China;

[2] Science Island Branch of Graduate School, University of Science and Technology of China, Hefei 230026, China;

[3] Center of Biology, Anhui Agricultural University, Hefei, China;

[4] Department of Medical Oncology, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China;

*Corresponding author:

Prof. Qing Huang

Hefei Institutes of Physical Science, Chinese Academy of Sciences, P. O. Box 1138, Hefei 230031, P.R. China

Tel: 86-551-65595261; Fax: 86-551-65595261

E-mail: huangq@ipp.ac.cn

Abstract

Mitochondrial biogenesis is a cell response to external stimuli which is generally believed to suppress apoptosis. However, during the process of apoptosis, whether mitochondrial biogenesis occurs in the early stage of the apoptotic cells remains unclear. To address this question, we constructed the COX8-EGFP-ACTIN-mCherry HeLa cells with recombinant fluorescent proteins respectively tagged on the nucleus and mitochondria, and monitored the mitochondrial changes in living cells exposed to gamma-ray radiation. Besides in situ detection of mitochondrial fluorescence changes, we also examined the cell viability, nuclear DNA damage, reactive oxygen species (ROS), mitochondrial DNA copy number and expression of transcription genes related to mitochondrial biogenesis as well as the apoptosis biomarkers. As a result, we confirmed that significant mitochondrial biogenesis took place preceding the radiation-induced apoptosis, and the change of mitochondrial biogenesis at early time was closely correlated with the apoptotic cells at late stage. The involved mechanism was also discussed.

Keywords: mitochondrial biogenesis; apoptosis; fluorescent proteins; ionizing radiation; living cells; DNA damage

Abbreviations: AICAR, 5-Aminoimidazole-4-carboxamide ribotide; ANOVA, Analysis of Variance; CCK-8, Cell Counting Kit-8; CytoC, Cytochrome C; DMEM, Dulbecco's modified Eagle's medium; DSB, Double strand breakage; *E. coli, Escherichia coli*; HCS, High-content screening; H2DCFDA, 2',7'-

dichlorodihydrofluorescein diacetate; HR, Homologous recombination mediated repair;

LDH, Lactate Dehydrogenase; MMP, Mitochondrial membrane potential; MFI, mean

fluorescence intensity; Mfn1, Mitofusin 1; mtDNA, Mitochondrial DNA; nDNA,

nuclear DNA; NHEJ, Non homologous end joining; PI, Propidium iodide; PARP, Poly

ADP-Ribose Polymerase; PGC-1α, Peroxisome proliferator-activated receptor-gamma

coactivator (PGC)-1alpha; RT-qPCR, Real-Time Quantitative Reverse Transcription

PCR; ROS, reactive oxygen species; SD, standard derivation; SIRT, Silent information

regulator; TMRE, Tetramethylrhodamine ethyl ester; TFAM, transcription factor A,

mitochondrial.

Synopsis

This study revealed that significant mitochondrial biogenesis could take place even when the cells were destined to the fate of apoptosis, and therefore for the first time explicitly demonstrated that mitochondrial biogenesis could be positively correlated with apoptosis given that these events were evaluated at right separate time points with consideration of their logical time-sequence of occurrence.

Highlights

- A dual fluorescence reporter system was constructed for in-situ observation of mitochondrial biogenesis in living cells.
- Significant mitochondrial biogenesis in the radiation-induced apoptosis was observed.
- Positive correlation of early mitochondrial biogenesis with late apoptosis was

68   explicitly revealed.

69 ●   Assessment of the early event of mitochondrial biogenesis is critical for prediction

70   of the late fate of cells.

Introduction

Mitochondrial biogenesis is a physiological response of cells to external stress and

development signals that may cause increase of energy demand, and it plays an

important role in cell metabolism regulation, signal transduction and mitochondrial

ROS regulation (Luo *et al*, 2016; Vyas *et al*, 2016). Mitochondrial biogenesis maintains cell

homeostasis by ensuring the quality of mtDNA and regulating the renewal of organelles

(Yambire *et al*, 2019), and it has been extensively explored in recent years due to its related

interests in human aging, neurodegenerative diseases, cell metabolic diseases and

tumors(Fanibunda *et al*, 2019).

Since mitochondrial regulation plays a very critical role in controlling the cell fate,

it is also intriguing for researchers to explore the relationship between mitochondrial

biogenesis and apoptosis. Generally, apoptosis is mediated by the activation of the

caspase pathways which are accompanied with mitochondrial damage accompanied by

destruction of electron transfer, oxidative phosphorylation, ATP production and change

of cell redox potential(Burke, 2017). In fact, some studies have claimed that there is a

negative correlation between mitochondrial biogenesis and apoptosis, i.e., apoptosis

may be substantially suppressed by mitochondrial biogenesis. For example, it has been

reported that stress-induced apoptosis can be inhibited by enhanced mitochondrial

biogenesis (Dam *et al*, 2013), or, on the other hand, apoptosis and cell cycle arrest may

occur by inhibiting mitochondrial biogenesis(Cao *et al*, 2017). Also, in the study of

radiation-induced apoptosis, mitochondrial biogenesis was observed and analyzed (Rai

et al, 2018), and it was claimed that enhancement of mitochondrial biogenesis could

reduce the proportion of apoptosis caused by ionizing radiation(Yu *et al*, 2013). However,

regarding the whole process of apoptosis, whether and how mitochondrial biogenesis

takes place in the early period in the apoptotic cells exposed to the external stimuli such

as ionizing radiation remains elusive.

To study the process of mitochondrial biogenesis related to apoptosis, it is thus

crucial to monitor the mitochondrial changes in the cells. At present, there are not many

methods available to study the influence of external factors on mitochondrial changes.

One of the conventional methods is to probe the change of mitochondrial gene copy

numbers(Yu, 2011). Another approach is to use fluorescent proteins. Fluorescent proteins

are beneficial for studying the living cells with reporting gene expression, which can

provide information on protein locations and expression levels as well as the involved

biochemical activities(Newman *et al*, 2011). Actually, study of mitochondrial process

using fluorescent proteins is now attracting increasing attention (Melentijevic *et al*, 2017;

Katajisto *et al*, 2015; Ruan *et al*, 2017). Recent development is to make use of multiple

different fluorescent proteins with different excitation/emission spectra, so that the

precision of transcriptional measurements can be ensured by using the reference color

for normalization (Miyashiro & Goulian, 2007). In such a circumstance, the ratio of dual

fluorescence can be used to eliminate the possible fluorescence measurement

errors(Miyashiro & Goulian, 2007). So, with application of this dual fluorescence approach,

nowadays researchers are able to observe and analyze the temporal and spatial life

trajectories of living cells (Jadhav & Shivdasani, 2019), and maybe more importantly, to

scrutinize the early events leading to various bio-effects such as apoptosis as end-point

fate in the living cells (Yu *et al*, 2020).

Therefore, in the present work, we attempted to apply the dual recombinant

fluorescent proteins to monitor the mitochondrial biogenesis induced by radiation, and

in this way, to examine whether and how the early event of mitochondrial biogenesis

would be associated with the prominent bio-effect such as apoptosis. For the effect of

radiation as the external stimuli to the cells, it is known that ionizing radiation (such as

α-, γ-, X-rays, protons, heavy ions) can readily cause various detrimental damages to

living cells, which can lead to the change of genetic information, cell mutations,

genomic instability, and apoptosis (O'Driscoll & Jeggo, 2006). Mitochondria, as important

subcellular organelles, are one of the major targets of ionizing radiation as they occupy

a considerable proportion of the cell volume (4-25%)(Leach *et al*, 2001). On the other

hand, radiation can also effectively induce mitochondrial biogenesis (Das *et al*, 2020). For

the establishment of the dual fluorescent system for in-situ observation of

mitochondrial biogenesis in living cells, we constructed the stable fluorescent reporter

cell lines by lentiviral transfection to achieve mitoGFP and nuclear RFP. As a result, we

fulfilled the task of in-situ and real-time observation of mitochondrial changes in the

living cells, and observed the effect of mitochondrial biogenesis occurring in the early

process of the apoptotic cells. By analyzing the fluorescence ratio of mitoGFP and

nuclear mCherry fluorescence, we could therefore examine the relationship between

the early mitochondrial biogenesis and late apoptosis, and testified that the increased

mitochondria number in the early time could be positively correlated with the apoptotic

cell population. The detailed mitochondrial/apoptotic processes were analyzed, and the

involved mechanism was discussed.

Results

Construction of dual-fluorescent stably transfected cell lines

In order to construct a universal dual-fluorescence gene reporter vector, we

considered to make use of the simplicity of vector construction and referred to the

previously reported dual-fluorescence reporter system, and so we chose the CMV

promoter to express the dual-fluorescence target proteins. The mitochondrial reporter

expression system is generally localized in the stroma, targeting the leader peptide

(MSVLTPLLLRGLTGSARRLPVPRAKIHSLGDP) on the VIIIa subunit of

cytochrome C (COX8) (Haggie & Verkman, 2002) (Fig 1A). As such, we constructed the

pLVX-mCherry-actin lentiviral vector (Fig 1B).

Figure 1.

A. Schematic illustration of the pEGFP-cox8 reporter system.

B. Schematic illustration of the pLVX-actin-mCherry reporter system.

C. Fluorescence images of HeLa cells transfected with EGFP-tagged COX8.

D. Fluorescence images of HeLa cells transfected with mCherry-tagged Actin.

To establish the cell line stably co-expressing COX8-EGFP and ACTIN-mCherry,

we transfected the HeLa cells with the plasmid containing the expression sequence of

COX8-EGFP, and obtained the COX8-EGFP expression cells. After screening by G418

for 2 weeks, the stably transfected cell line was obtained. The COX8-EGFP-mCherry-

ACTIN co-expressing HeLa cell line was further established based on the COX8-EGFP

expressing cells with lentiviral transfection system. Fig 1 shows the fluorescence

microscope examination of the cells, confirming the co-expressions of COX8-EGFP

and mCherry-ACTIN.

In order to further confirm the GFP expression right on the targeted mitochondria,

we then labeled the mitochondria with TMRE dye in the GFP-tagged cells and then

examined the dye-stained cells with a confocal microscope. As shown in Figure 2, the

COX8 protein fluorescence (Green) proteins and mitochondrial fluorescence (Red) were indeed co-localized in the cells.

Figure 2.

Confocal microscopy observation of co-localization of GFP-COX8 on the mitochondria. The GFP-COX8-HeLa cells were stained with TMRE (200 nM) for 20 minutes and before the examination by confocal microscopy. Scale bar: 10 μm.

Furthermore, in order to verify the universal validity of our as-constructed dual fluorescence reporting system for observing mitochondrial biogenesis, we also used mitochondrial biogenesis's initiator 5-Aminoimidazole-4-carboxamide ribotide (AICAR) as for the in-depth testing, or, as the positive control(Komen & Thorburn, 2014). The applied drug concentration was 0.5 mM. As shown in the Fig 3A-D, compared with the non-AICAR treatment group, the AICAR treated group showed increased fluorescence ratio with time. This experiment therefore confirmed that the dual fluorescence system we constructed is feasible and effective for monitoring mitochondrial biogenesis.

Figure 3.

A-D. The cells were treated with mitochondrial biogenesis inducer drug (AICAR, 0.5 mM) and

observed with high-content imaging and screening (HCS) system after different treatment hours.

In the figures, green color indicates the mitochondria encoding COX8-EGFP, red color shows

the actin encoded by actin-mCherry, and blue color represents the DAPI stained nuclei. Scale

bar: 25 μm.

E. The mean fluorescence intensity (MFI) ratios of HeLa cells treated by AICAR after different

hours, in comparison with the non-AICAR treated cells as the control group. Bar graphs are

presented as mean ± sd, n = 3.

Moreover, in order to further validate the effectiveness of the as-constructed

fluorescent reporter system, we also testified the co-expression levels of GFP protein

together with the mitochondrial target protein (e.g. COX8) for the positive control with

the cells treated by AICAR. Technically, CytoC was measured in the experiment instead

of COX8 as COX8 is the subunit of CytoC. As shown in Fig 4, both GFP and CytoC

were in the same trend of expression of levels for the mitochondrial biogenesis proteins

including PGC1-α, NRF1 and TFAM. Besides, we also examined the AICAR-induced

bio-marker proteins expression levels related to mitochondrial biogenesis. Fig. 4 shows

that for the cells treated with AICAR, the expression levels of GFP, CytoC, PGC1-α,

NRF1 and TFAM increased significantly in comparison with the untreated group. Also,

with the increase of AICAR treatment time, the protein expression levels of

mitochondrial biogenesis marker proteins (PGC1, NRF1, TFAM) also increased

concomitantly (see Fig EV1). All these results therefore confirmed the validity of our

fluorescence report system suitable for mitochondrial biogenesis monitoring.

Figure 4.

A. HeLa cells were treated with mitochondrial biogenesis inducer drug (AICAR, 0.5 mM).

Total proteins were extracted after 12 hours.

B.   The graph shows the comparison of the relative intensity for expressions of GFP and

mitochondrial biogenesis marker proteins between the non-AICAR treated group and the

AICAR treated group.

In-situ observation of the dual fluorescence reporter system for the evaluation of

the radiation-induced mitochondrial changes

With the establishment of this dual-fluorescence reporter system, we then employ

the laser fluorescence confocal microscope to observe the biological effect in the living

cells with the treatment of gamma-ray irradiation. The COX8-EGFP encoding

mitochondria (Green), the ACTIN-mCherry encoding actin (Red), and the DAPI-

stained nuclei were observed. The representative images together with the

corresponding quantitative analysis of fluorescence intensity are shown in Fig 5.

Figure 5.

A-D.   HeLa-cox8-EGFP-actin-mCherry cells were exposed to 1 Gy, 2 Gy, 4 Gy gamma-ray

and incubated for 24 h and observed with confocal laser scanning microscope. Green color

indicates the mitochondria encoding COX8-EGFP, red color shows the actin encoded by

actin-mCherry, and blue color represents the DAPI stained nuclei. Scale bar: 25 μm.

E. The ratio comparison for the mean fluorescence intensity (MFI) of GFP over mCherry

fluorescence intensities quantified in the figure 2A-D. The MFI was determined by ImageJ

software. Bar graphs are presented as mean ± sd, n = 3.

Fig 5 presents the fluorescent images of the cells by confocal microscopy,

indicating that the intracellular mitochondrial content increased with radiation dose.

Compared with the control group (0 Gy), the irradiated group showed mitochondrial

fluorescence enhancement, and it was significantly enhanced especially for the 1 and 2 Gy cases (Fig 5A-D). The fluorescent images intuitively show that the number of mitochondria increased with increasing radiation dose, confirming that the dual fluorescence reporter system can directly reflect the relevant mitochondrial responses and changes in the living cells.

To investigate the mitochondrial fluorescence enhancement of the cells exposed to ionizing radiation, the mean fluorescence intensity (MFI) from the dual fluorescence reporter system was utilized. This MFI ratio change was also obtained by employing the High Content Imaging and Screening (HCS) system, with the results shown in Fig 6. The HCS living cell workstation recorded the changes of cell fluorescence within 24 hours after cell irradiation (the videos were also recorded, see Videos S1, S2, S3 in the Supporting Information). The results indicate that for 1 Gy, the fluorescence ratio GFP/mCherry increased with time, while for 2 Gy and 4 Gy cases, the fluorescence ratio increased significantly with the irradiation dose during the first interval of 1-12 hours, but afterwards, it decreased to some extent, though still higher than that of the control (non-irradiated cells). The reduction in MFI suggested that the mitochondria of the irradiated cells with higher radiation doses were actually more detrimentally damaged by the radiation than that received lower radiation doses. Fig EV2 shows the numerical changes of two fluorescence channels detected by flow cytometry. The MFI ratio of EGFP$^+$: mCherry$^+$ was calculated, and it was found that the fluorescence ratio was positively correlated with the radiation dose. So the result by the flow cytometry

(Fig EV2) also confirms there were significant differences between the irradiated and

non-irradiated groups.

Figure 6.

The cellular mean fluorescence intensity (MFI) measured by High Content Imaging and

Screening (HCS) system. HeLa-Cox8-EGFP-actin-mCherry cells were exposed to 1 Gy, 2 Gy,

and 4 Gy gamma-ray respectively. MFI changed with time after the γ-ray irradiation.

Radiation effect on cell viability, survival rate, cell cycle, apoptosis, DNA damage

and mitochondrial membrane damage

To investigate the radiation effect, the cell viability was measured by CCK-8

method, showing that the cell viability decreased with the increase of radiation dose

(Fig 7A). The toxic effect of radiation on cells was also examined using LDH kit, which

revealed that the radiation-induced toxicity was indeed increased significantly with

radiation dose (Fig 7B). Also, the colony-forming method was used to analyze the

inhibitory effect of γ-ray on HeLa cell survival. With increase of γ-ray irradiation dose,

the cell survival rate was only 30% after 4 Gy irradiation (Fig 7C), confirming that the

cell survival rate decreased with the rise of radiation dose.

Figure 7.

A.   CCK-8 assay for cell viability.

B.   LDH assay for cell cytotoxicity.

C.   Colony formation assay of HeLa cells after 14 days under different radiation doses.

D.   Flow cytometry results of Annexin V-PI assay.

E.   Apoptotic cells at different radiation doses.

F.   Cell cycle examination. The experiment was performed 3 times. Data are presented as

Mean±SD; $P < 0.01$(**); $P < 0.001$(***); ns: not significant compared with untreated cells.

In addition, the cell cycle and apoptosis were assessed for the cells irradiated with

γ-ray. Apoptotic cells were detected by flow cytometry after irradiation using Annexin

V-PI methods (Fig 7D). Through statistics and comparison of the number of apoptotic

cells, it was found that with the increase of radiation dose, the number of apoptotic cells

increased significantly (Fig 7E). Cell cycle was analyzed by flow cytometry, and the

results showed that all groups of cells had G2 arrest after 24 hours of irradiation,

especially at 2 Gy and 4 Gy irradiation doses showed significant G2 arrest (Fig 7F).

Furthermore, in order to understand the radiation effects, the radiation induced

damages on DNA and mitochondrial membrane, together with the accompanied ROS

levels were also examined. For such a purpose, the immunofluorescence method was

used to detect DNA damage, H_2DCFDA probe was used to label ROS, and TMRE was

used to label mitochondrial membrane potential. Fig 8A shows that with increase of the

radiation dose, the foci of the γ-H2AX increased significantly, and the statistical results

showed that the ROS level in the irradiated cells increased with the increase of the

radiation dose (Fig 8B- D). At the dose of 2 Gy, the ROS level was the highest, and for

the 4 Gy group, there was a significant difference compared with the non-irradiated

group. In the meantime, the mitochondrial membrane potential decreased significantly

with the increase of radiation dose (Fig 8E, F).

Figure 8.

A. Representative images for DNA damage measured instantly (within 30 minutes) after the

    Hela cells were exposed to radiation with different radiation doses. Green: γ-H2AX; red:

    DNA stained with PI; Scale bars: 20 μm.

B. The graph shows the percentage of population with the noted number of γ-H2AX foci with

    different irradiation doses.

C-D. ROS assessment using H_2DCFDA assay at 6 h.

E- F. MMP assay using TMRE at 24 h.

As the ionizing radiation induced DSB of DNA, we expected that there would also

be enhanced DNA repair process in the irradiated cells. Indeed, we observed the

fluorescence from γ-H2AX and the co-localization of 53BP1 foci after the radiation

exposure. Fig 9 shows the number of foci increased significantly with the increase of

radiation dose, while the number of foci of 53BP1 was the largest at 1 Gy but then

decreased with radiation dose, implying that at higher radiation dose the cell received

severer damage son the DNA repair ability became weaker.

Figure 9.

A. Representative images for DNA damage detection when the cells were exposed to

radiation with different radiation doses. Green: γ-H2AX; red: 53BP1; Blue: DNA nucleus

stained with DAPI; Scale bars: 10 μm.

321 B. The percentage of population with the noted number of γ-H2AX foci corresponding to the

322   respective fluorescence images.

323 C. The percentage of population with the noted number of 53BP1 foci corresponding to the

324   respective fluorescence images. At least 100 nuclei per genotype were used for

325   quantification. Image auto-quantification was performed with CellProfiler version 4

326   (www.cellprofiler.org).

Verification of mitochondrial biogenesis and apoptosis

To further verify the mitochondrial biogenesis induced by radiation, the copy number of mitochondrial DNA was evaluated. Total DNA was extracted from HeLa cells 6-24 hours after irradiation. As shown in Fig 10, the copy number of mitochondrial DNA changed significantly after cell irradiation. Indeed, under the same radiation dose, mitochondrial DNA number increased significantly compared with that of the non-irradiated cells.

Figure 10.

Effect of gamma-irradiation on mitochondrial DNA copy number. HeLa cells were exposed to gamma rays at doses of 1 Gy, 2 Gy and 4 Gy. After 6, 12, 24 hours, total genomic DNA was extracted and used as a template to detect the mtDNA/nDNA ratio by RT-qPCR. Data are presented as Mean±SD; #: significant compared with unirradiated group; $**P < 0.01$; $***P < 0.001$.

Furthermore, to explore the correlation between mitochondrial biogenesis and

apoptosis under the condition of radiation, the related marker proteins were also

examined. For the mitochondrial change, the related protein expression levels of PGC-

1α, NRF1, TFAM and Mfn1 at different times (6 h, 12 h, 24 h) with different radiation

doses (0 Gy, 1 Gy, 2 Gy, 4 Gy) were assessed by Western blots (Figure 11). At 6 hours,

the expression level of TFAM protein was increased with radiation dose. On the other

hand, for the apoptosis, the related protein expression levels of apoptosis-related marker

proteins (BAX, Bcl-2, CytoC, Caspase-3, Cleaved Caspase-9, PARP) were examined.

As shown in Figure 8, the expression levels of crucial apoptosis proteins did not differ

significantly at early time (6 h), but their expression levels increased significantly

afterwards (12 h, 24 h). So these results unambiguously demonstrate that the

mitochondrial biogenesis at early time (6 h) was indeed positively correlated to the

apoptosis in the late stage (24 h).

Figure 11.

Western blot (WB) assay of the related proteins after radiation treatment. HeLa cells were treated with different doses of irradiation (0, 1, 2, 4 Gy), and total proteins were extracted at different time points. The WB assay shows the changes of expression of mitochondrial biogenesis marker proteins (PGC-1α, NRF1, TFAM and Mfn1) and apoptotic marker proteins (BAX, Bcl-2, CytoC, Caspase-3, Cleaved Caspase-9, PARP) with time.

Discussion

Mitochondria as the cell energy factory play a vital role in cell homeostasis(Lill & Freibert, 2020). In response to external stimuli, mitochondrial biogenesis can be elicited in order to provide the cells with extra energy(Barros & McStay, 2019). Previously, many studies had reported that mitochondrial biogenesis could suppress apoptosis(Yu *et al*, 2013), i.e., there would be negative correlation between mitochondrial biogenesis and apoptosis. Generally, these studies mainly concerned the cell response to relatively small stimulus or stress. But what would happen if the external stimulus or stress is

large enough so that the final fate of cell death or apoptosis cannot be avoided? Would

mitochondrial biogenesis still go along with apoptosis in this case? And if there would

be mitochondrial biogenesis occurring as the early event of the apoptosis, how would

it change with time? With these questions, we thus initiated this research, and applied

radiation approach to induce both mitochondrial biogenesis and apoptosis in the cells,

so as to explore the time-dependent relationship between mitochondrial biogenesis and

apoptosis. As a result, we have achieved the following results with some new findings

and elaborated understandings.

Firstly, we have constructed the dual fluorescence report system that can be

effectively used for mitochondrial biogenesis study. To probe the changes taking place

in mitochondria, we especially developed the approach using mitochondrial-tagged

recombinant fluorescent proteins. The fluorescent protein reporting system has the

advantages of real-time, convenient, fast, and non- destructive(Betzig *et al*, 2006). Besides,

it has also an advantage in the study of subcellular organ localization and protein stress

regulation(Llopis *et al*, 1998). Previously, people often used fluorescent dye probes for

studying effect involved with mitochondria. For example, Mukhopadhyay et al. used

Annexin V and Sytox Green and MitoSOX Red fluorescent probes to study the

detection of apoptosis and mitochondrial superoxide so as to establish the method for

detecting living cells by laser confocal scanning microscopy and flow cytometry

(Mukhopadhyay *et al*, 2007). But the disadvantages of fluorescent dyes are the toxicity to

cells, as well as the light quenching under long-period real-time observation. Compared

to fluorescent dyes, fluorescent proteins are more stable and have minimal toxicity and

can generate visible fluorescence in vivo (Westermann & Neupert, 2000). Actually, GFP

reporter protein is now commonly used to detect mitochondrial changes(Mahajan *et al*,

1998; Hanson *et al*, 2004). In this work, in particular, we constructed the dual

fluorescence reporting system based on fluorescence proteins for monitoring the

mitochondrial biogenesis process. In order to verify the effectiveness of this system

designed for monitoring mitochondrial biogenesis, in this study, we applied AICAR to

activate mitochondrial biogenesis as the positive control (Komen & Thorburn, 2014).

AICAR is an activator of AMP-activated protein kinase (AMPK) that can permeate cell

membranes(Rai *et al*, 2015). Among them, AMPK (AMP activated protein kinase) is

an activator of mitochondrial biogenesis and key regulator of energy homeostasis

caused by external factors(Kim *et al*, 2016). The main function is to phosphorylate

PGC-1α protein, or affect the transduction of SIRT1 signaling pathway(Mihaylova &

Shaw, 2011). Our result clearly showed that AICAR indeed induced the up-regulation

of key protein levels of mitochondrial biogenesis, confirming that the as-constructed

dual fluorescent protein reporter system could track the event of mitochondrial

biogenesis effectively. Also, this dual fluorescent protein reporter system is not only

suitable for the real-time in situ observation for a long time, but also facilitates the

precise fluorescence evaluation with the reference fluorescent protein for intensity

normalization. Therefore, by analyzing the ratio of the fluorescence intensity of

mitochondrial DNA to that of nuclear DNA expression, the early event of mitochondrial

biogenesis under irradiation condition could be readily observed and evaluated, and

with this approach we could thus sensitively track the early process that which

eventually led to the late event of apoptosis.

Secondly, with the as-constructed dual fluorescence report system, we have

observed the radiation-induced mitochondrial biogenesis together with the apoptosis.

Ionizing radiation has been used for cancer therapy for they can directly or indirectly

cause DNA damage and kill the cancer cells(Spitz et al, 2004).　With DNA damage in

the nucleus, cells may undergo cell cycle arrest required for repairing the damage or

cell death including apoptosis(Bernstein et al, 2002). In parallel, mitochondria is also

considered to be involved in the radiation induced effects, where more reactive oxygen

species (ROS) would be produced which may affect the cells profoundly. Actually, it

has been well documented that exposure of cells to ionizing radiation could activate

ROS to produce oxidases, regulate antioxidants, and alter metabolic activity in response

to oxidative damage(Azzam et al, 2012). In both the nucleic or mitochondrial processes,

more ATP is required to elicit mitochondria biogenesis either directly or indirectly(Kulms

et al, 2002; Srinivas et al, 2019). So it is not surprising that mitochondria biogenesis could

take place as one of the consequences of radiation-induced bio-effects (Galluzzi & Kroemer,

2008). But interestingly, in our fluorescence measurements, we found that the

fluorescence ratio increased and reached the maximum at 6 h after irradiation, but

afterwards the fluorescence ratio decreased gradually. According to our established

mitochondrial fluorescence report system, this suggests that in the early stage,

mitochondrial biogenesis was indeed initiated and enhanced. In this process of

mitochondrial biogenesis, mitochondrial DNA and mitochondrial mass were

maintained to keep the cell homeostasis and metabolism(Huangyang et al, 2020). But in

the late stage of the radiation effect, the fluorescence ratio decreased, showing that the

function of mitochondrial biogenesis gradually diminished. We can also understand this

change by relating it to the ROS elevation. There was increase of ROS caused by

radiation, which continuously increased the oxidative stress and damaged the mtDNA

and the mitochondrial integrity. For the role of ROS, it is generally accepted that normal

range of ROS may induce stress response by changing the expression of related nuclear

genes, and certain level of ROS can save cells by maintaining energy metabolism

(Srinivas et al., 2019); but when the ROS level exceeds the tolerant threshold, ROS can

then induce the transition of mitochondrial membrane permeability, leading to the

activation of the caspase pathway and promoting cell apoptosis. In our experiment, we

indeed detected the increase of ROS level and the decrease of mitochondrial membrane

potential. Also, given the increase of radiation dose, we detected the related caspase

related proteins and found that the increase of pro-apoptotic proteins such as

cytochrome c. Release of cytochrome c is an indicator for the apoptosis, in which the

oxidative stress would damage mtDNA and other components of the cells(Lee & Wei,

2005). Here, we especially notified that the mitochondrial fluorescence intensity change

rates were quite different after the cell exposure to the radiation. The larger the

irradiation dose, the higher the fluorescence ratio reduction rate. All these observations

are actually the typical features for the radiation-induced apoptotic process.

Thirdly, our new observation have revealed the more subtle but very important

relationship which had largely been ignored by previous studies. We noticed that in the

beginning of the process, the mitochondrial biogenesis was actually increasingly

associated with the apoptotic stress, and the trend of fluorescence change was consistent

with the number of apoptotic cells. This is actually contrary to the conventional view

on the relationship between mitochondrial biogenesis and apoptosis. In the past, there

were few reports concerning the dynamic process of mitochondrial biogenesis, and the

mainstream opinion or understanding on the relationship between mitochondrial

biogenesis and apoptosis was that these two processes normally appear to be

antagonistic. For example, many studies showed that the apoptosis occurred

simultaneously with the suppression, or, that apoptosis took place with even missing

the observation of mitochondrial biogenesis (Martins *et al*, 2015; Vayssiere *et al*, 1994;

Zhang *et al*, 2017; Lin *et al*, 2020). This is actually not true or accurate, because

mitochondrial biogenesis as an earlier event did occur preceding the late state of

apoptosis, as we clearly demonstrated in this study.

The reason for this misleading of understanding is largely due to the ignorance of

the investigation of the dynamic process of the cells. In the past, people tended to

choose just several time check points for inspecting the apoptosis and mitochondrial

biogenesis. But which time check points are most suitable for the correct evaluation?

What cautions should be taken to avoid the wrong evaluation when especially

concerning the relationship between the processes in the involvement? To solve this,

we actually need to establish more advanced method to studying the cellular effect in

its dynamic process, and so to scrutinize the processes of both mitochondrial biogenesis

and apoptosis in their logical time sequence. This is simply just one of the main

purposes of this study. It is understandable why people had not focused on the study of

processes, as we realized that there had been lack of tool/method for the monitoring the

whole process regarding the mitochondrial changes in situ and in a nondestructive way.

For example, the traditional methods include using real-time quantitative PCR to detect

the copy number of mitochondrial DNA, or using Western blot assay to identify the

changes of marker protein level of mitochondrial biogenesis. These methods cannot

achieve in-situ and real-time observation in living cells. On the contrary, the

fluorescence protein method can be used to observe the mitochondrial biogenesis in

real time as especially useful for studying the early events prior to apoptosis. Therefore,

in this work, we at first attempted to construct the fluorescent reporting system which

could let us to track the change of mitochondrial biogenesis. Based on the fluorescence

signal, the dual fluorescence ratio can help us to predict the radiation effects or the cell

fate according to the fluorescence change of these irradiated cells. In particular, from

the fluorescence data obtained from Fig 6, we can calculate the fluorescence ratios, and

so we can retrieve the MFI ratios in each irradiation dose group (0 Gy, 1 Gy, 2 Gy, 4

Gy) at an early time, as shown in Fig 12 A&B. Strikingly, this result is consistent with

the cells in the apoptotic state at 24 h (Fig 7E). In the late stage (24 hours), the

fluorescence ratio becomes smaller as the irradiation dose increases. It is

understandable because the apoptosis is actually dependent on the early biochemical

events in the cells(Holler *et al*, 2000). In this way, we may therefore infer or predict the

late apoptotic state of cells by observing the early-staged changes in terms of

mitochondrial fluorescence ratio. On the other hand, it is also seen that this fluorescence

ratio was significantly reduced after 24 h (Fig 12C), which is not concomitant with the

apoptosis at 24 h (Fig 7E). This suggest that it is very critical to evaluate both the

processes by choosing the right time check points.

Figure 12.

Analysis of mitochondrial biogenesis and apoptosis.

A.   Analysis of EGFP/mCherry ratio in High Content Screening (HCS) at 6h.

B.   Analysis of EGFP/mCherry ratio in High Content Screening (HCS) at 12 h.

C.   Analysis of EGFP/mCherry ratio in High Content Screening (HCS) at 24 h.

Lastly, we have also tried to understand the mechanism for the mitochondrial

biogenesis related to the radiation-induced apoptosis. Generally, it is believed that the

biogenesis of mitochondria is a very complex biological process, which can maintain

the cell homeostasis by controlling the self-renewal of subcellular organelles and

maintaining mitochondrial DNA (Scarpulla, 2008). The initiation of mitochondrial

biogenesis is normally triggered by the reception of energy demand signals caused by

external stress (Popov, 2020), while at the same time, mitochondrial biogenesis itself is

a protection way for cell's self-renewal. Since it is well-known that ionizing radiation

is an effective way to stimulate the cell mitochondrial biogenesis, we thus employed

the tool of ionizing radiation to treat the cells. As a result, we have successfully proved

that some marker proteins of mitochondrial biogenesis are involved in the biological

effects of radiation. We have found that the protein expression levels of nuclear

respiratory factor1 (NRF1), peroxisome proliferative activated receptor, gamma,

coactivator 1 alpha (PGC-1α), transcription factor A, mitochondrial (TFAM) increased

with radiation dose. NRF1 is a transcription factor with a vital role in mitochondrial

functional genes(Virbasius *et al*, 1993). PGC-1α is a transcriptional coactivator which is a

key inducer of mitochondrial biogenesis (Wu *et al*, 1999). TFAM is a key activator

of mitochondrial (mt) DNA transcription. mtDNA is reported to be very susceptible to

oxidative stress(May-Panloup *et al*, 2005). With the confirmation of these factors, we have

thus also again unambiguously confirmed that this mitochondrial biogenesis is indeed

related to radiation-induced cell adaptation and apoptosis.

But how to explain the suppression or disappearance of mitochondrial biogenesis

for the higher dose of radiation observed in the late stage of the cell process which

corresponds to the apoptosis? In our experiments, we observed that mitochondrial

biogenesis increased in the early stage of the cell process, but with the increase of

radiation dose, the mitochondrial biogenesis decreased substantially in the late stage.

We speculate that this change of mitochondrial biogenesis is related to DNA repair

ability. It was previously reported that mitochondrial biogenesis can enhance DNA

repair ability, and mitochondrial biogenesis is closely related to DNA repair ability(Li

et al, 2020) (Fu *et al*, 2008). For example, it was reported that doxorubicin, a drug

which can induce DSBs of DNA as accompanied by mitochondrial biogenesis (Kluza

et al, 2004). Normally, there are two main repair pathways of DSBs in mammalian cells:

non homologous end joining (NHEJ) and homologous recombination mediated repair

(HR) (Löbrich & Jeggo, 2005). Mitochondrial biogenesis can promote DNA repair by

inducing chromatin changes in DSB and regulating the activity of DNA repair factors,

and this upstream factor has be found to be the silent information regulator (SIRT)

(Paredes & Chua, 2016). SIRT is a family of sirtuins that depend on NAD^+ lysine

deacetylases, which may play an important role in mitochondrial biogenesis and cell

metabolism(Chalkiadaki & Guarente, 2015). On the other hand, DSB repair can be

achieved by NHEJ which can activate DNA-PK (Davis *et al*, 2014), while DNA-PK

can activate AMPK signal pathway, which is just closely related to mitochondrial

biogenesis (Park *et al*, 2017). In our experiment, we have indeed observed the change

of DNA repair as reflected by the change of 53BP1, and this change is just concomitant

with the mitochondrial biogenesis.

So, with the above findings and discussion of mechanisms, we can therefore

concisely describe the process of radiation affecting cells as schematically depicted in

Fig. 13. After a certain dose of radiation, DNA damage occurs, and the cell cycle is

blocked in order to initiate the repair process. At the same time, radiation produces a

large amount of ROS in the cell and affects the homeostasis of the cell. To resist the

radiation stress or damage, mitochondria then start to provide more energy for DNA

repair and maintenance of cell metabolism and homeostasis, leading to mitochondrial

biogenesis. Next, as ROS continues to increase, mitochondrial damage occurs, leading

to the decrease in membrane potential, which finally leads to the release of the pro-

apoptotic protein cytochrome c, activation of the apoptosis caspase pathway, and so the

programmed cell death.

Figure 13.

Schematic diagram of biochemical events after ionizing radiation.

Conclusions

In this work, we have constructed a dual fluorescence reporter system for achieving

the real-time observation of mitochondrial changes, which are related to

mitochondrial biogenesis and apoptosis induced by ionizing radiation. Particularly, we

have scrutinized the relationship between this mitochondrial biogenesis and apoptosis

by considering the time sequence of occurrence, showing that these two processes are

not antagonistic if they are evaluated at right time points. In other words, they may be

positively correlated if we relate the mitochondrial biogenesis taking place at the early

time to the apoptosis occurring at the late stage. Therefore, this work not only provides

an effective method for directly observing the mitochondrial biogenesis process in

580　living cells, but also demonstrates how to evaluate the biological effects correctly with

581　right timing, with which we may thus be able to predict the cell fate such as apoptosis

582　through the cell's early event observation.

Materials and methods

Reagents

The *Escherichia coli* DH5a and DNA Ladder were purchased from Transgen

(Beijing, China). *Nhe*I, *Hind*III, *EcoR*I, *Not*I restriction enzymes, T4 DNA Ligase, and

Q5 High-Fidelity DNA Polymerase were purchased from NEB (MA, USA). Plasmid

pEGFP-N1 was purchased from Clonetech Biotechnology (CA, USA). Plasmid pLVX-

mCherry, PUC57 were stored in our laboratory. Plasmids of psPAX2 (# 12260) and

pMD.2G (#12259) were purchased from Addgene. PCR purification and gel extraction

kits were obtained from Axygen (CA, USA). TRIzol™ reagent, Lipofectamine 2000

were purchased from Invitrogen (CA, USA). Primescript II 1st strand cDNA Synthesis

Kit, RT-qPCR kit and Total genomic DNA extraction kit were purchased from

TAKARA (Dalian, China). G418, DAPI was purchased from Beyotime Institute of

Biotechnology (Shanghai, China). Reactive oxygen species (ROS) probe 2',7'-

dichlorodihydrofluorescein diacetate (H2DCFDA, D6883) was purchased from Sigma-

Aldrich. Tetramethylrhodamine ethyl ester (TMRE, T4057) was purchased from

Useverbright. 5-Aminoimidazole-4-carboxamide ribotide (AICAR) was purchase from

Selleck (MA, USA). The antibodies of PGC-1α, NRF1, TFAM, Mfn1, BAX, Bcl-2,

CytoC, Caspase-3, Cleaved Caspase-9, PARP, γH2AX(S139), GFP were purchased

from Bimake (MA, USA), 53BP1, anti-β-Actin antibody was purchased from CST (MA,

USA). Alexa Fluor® 488 Goat Anti-Mouse (IgG) secondary antibody, Alexa Fluor®

647 Goat Anti-Rat (IgG) secondary antibody were purchased from Invitrogen (CA,

USA).

Creation of stably transfected cell lines

Generation of plasmids. Firstly, the pEGFP-N1-COX8 expression plasmid was

constructed. Briefly, the DNA sequence of cytochrome c oxidase 8A subunit was found

from Genbank (NM_004074.2) (Hendrickson et al., 2010). Then two complementary

oligonucleotide strands were obtained by amplification using PCR, and the restriction

sites of the *Nhe*I and *Hind*III were added at both ends. The sequences of oligonucleotide

strands were listed in S1 Table. The DNA sequence (COX8A) was ligated on the PUC57

plasmid. Then, the appropriate oligonucleotides were annealed and ligated into pEGFP-

N1 cloning cut with *Nhe*I and *Hind*III. After incubation overnight at 37°C, the target

plasmids were extracted from the bacterial fluid using the EndoFree Maxi Plasmid kit.

Secondly, the PLVX-mCherry-Actin expression plasmid was constructed. Total

RNA was extracted from HeLa cells by Trizol reagent and served as the template for

cDNA synthesis, and then it was reversely transcribed using Primescript II 1st strand

cDNA synthesis kit. The primers were designed according to the sequence of actin

(Genbank no. NM_10277): *EcoR*I tailed forward (5′-

gtcGAATTCatggatgatgatatcgccgcgc-3′) and *Not*I tailed reverse (5′-

TATGCGGCCGCctagaagcatttgcggtggac-3′) primers which are named as Fw-actin-

EcoRI and Rv-actin-NotI, respectively. The length of the amplified product was 1128

bp. The PCR product was separated by electrophoresis, and the target fragment was

recovered after gel cutting and purification. The target gene fragment (actin) and the

expression vector (LVX-mCherry) with the same restriction site (*EcoR*I and *Not*I) were

digested by restriction enzymes respectively, and then ligated with T4 ligase and

transformed into *E. coli* DH5a for plasmid amplification.

Production of lentiviral supernatant and transduction of COX8-EGFP-mCherry-

ACTIN-HeLa. Briefly, positive monoclonal cells (COX8-EGFP-HeLa cells) were

constructed. When HeLa cells grew to the concentration of 80% in complete medium,

they were then transfected with pEGFP-COX8 plasmid with lipofectamine 2000. After

24-48 h of transfection, G418 was added to screen positive monoclonal cells.

The lentiviral supernatant was prepared according to the protocol provided by

addgene (http://www.addgene.org/tools/protocols/plko/#A). When the confluence of

HEK293T cells cells in the 6-well plate reached 80%, Lipofectamine 2000 was used to

transfect the pLVX-mCherry-ACTIN plasmid and the viral packaging helper plasmid

psPAX2, pMD.2G into the cells. After incubation for 8h, the medium was replaced with

the fresh medium containing serum, and the supernatant was collected 24 hours and 48

hours later. The virus supernatant was filtered through a 0.45 μm pore so as to infect

COX8-EGFP-HeLa cells. Next, the COX8-EGFP-HeLa cells were cultured in a 6-well

plate to 70% confluence, then the medium was removed, and the virus supernatant was

mixed with fresh medium containing 8 μg/ml polybrene in equal proportions and added

to the culture plate. The fresh medium was replaced after 24 hours and the medium

containing 1 μg/ml puromycin was changed at 48 hours to start screening to obtain

stable cell lines.

Cell Culture, Synchronization and Gamma irradiation

HeLa, HEK-293T cell lines were purchased from American Type Cultures

Collection (ATCC, Manassas, VA). The HeLa and HEK-293T cells were cultured in

Dulbecco's modified Eagle's medium (DMEM, HyClone) containing 10% fetal bovine

serum (FBS, Gibco BRL). All cells were maintained at 37 °C in 5% CO2.

The cell cycle synchronization method was detailed in the previous

works(Rodríguez-Martínez et al., 2020). The cells were incubated with thymidine (2.5 mM)

for 18 hours, washed once with PBS buffer and replaced with fresh medium for 14

hours, and then 2.5 mM thymidine was added again to the cells which were then

incubated for 18 hours. The efficiency of synchronization was tested by flow cytometry

using cell cycle analysis kit (Multisciences).

HeLa cells were gamma-ray irradiated with a Biobeam Cs137 irradiator (cat no.

GM 2000; Gamma-Service Medical, Leipzig, Germany) at 1, 2, and 4 Gy witho the

dose rate of 3.37 Gy/min. In the control group, the cells were taken out of the incubator

without irradiation.

Cell viablity, LDH measurements and colony-formation assays

Cell viability was measured with CCK-8 assay kit (Dojindo, Japan). Briefly, 24 h prior to irradiation treatment, ca. 6000 cells were plated in 96-well plates in a final volume of 100 μl, and then cultured for 24h. And after adding the CCK-8 solution, the plate was incubated for another 1 h. The optical density was measured at 450 nm using a microplate reader (SpectraMax M5, Molecular Devices, USA). The amount of LDH released into the medium was assayed using the LDH-Cytotoxicity Assay Kit II (Dojindo, Japan) according to the manufacturer's instructions. For the colony formation assay, 500 cells were seeded in 6-well plate. After being irradiated, the cells were cultured at 37℃ for about 2 weeks. After carefully washing twice with PBS, 4% paraformaldehyde was added to fix the cells. The the cells were then stained with crystal violet solution and evaluated.

Analysis of mitochondrial membrane potential and ROS

TMRE-Mitochondrial membrane potential (MMP) assay kit (Useverbright, USA) was used to measure MMP. The cells were incubated with culture medium containing 200 nM TMRE for 45 minutes at 37 °C, then rinsed with PBS 3 times, followed by incubation with medium containing 50 nM TMRE. For the quantitative assessment, the TMRE staining was evaluated using SpectraMax M5 microplate reader.

Reactive oxygen species (ROS) measurement was performed using H2DCFDA (Sigma-Aldrich, USA) probe. 4×10^3 cells were seeded into 96-well plates and incubated overnight. Before irradiation, the cells were incubated with HBSS solution

containing 5 μM H₂DCFDA for 15 minutes and then irradiated. After 30 minutes, the

cells were washed 3 times, followed by the fluorescence measurement using a

microtiter plate reader.

Annexin V-FITC/PI staining and cell cycle analysis

The HeLa cells were seeded at 2×10^5 cell /well in 6-well plates and irradiation

with gamma-ray. After 24 h, the cells were fixed with cold ethanol and then stained with

propidium iodide. The cells (about 20,000) were analyzed by flow cytometry (Beckman

CytoFLEX, USA). The total cells were washed once and resuspended in Annexin V

binding buffer, and then stained with 5 μl Annexin V and 5μl PI (Roche, Germany) for

20 min and analyzed with Beckman CytoFLEX flow cytometry. Ten thousand events

were collected for each sample.

Measurement of mtDNA copy number

Total genomic DNA of HeLa cells was extracted after irradiation using Total

genomic DNA extract kit (TAKARA, China). Human nuclear GAPDH gene (forward:

5′-   CAGAACATCATCCCTGCCTCTAC-3′;   reverse: 5′-

AAGGGTTGTAGTAGCCCGTAG-3′) and the mtDNA ND1 gene (forward: 5′-

TTCTAATCGCAATGGCATTCCT-3′;   reverse: 5′-

AAGGGTTGTAGTAGCCCGTAG-3′), as well as a primer were used to detect the

mtDNA copy number. The real-time quantitative PCR analysis was performed using

SYBR Premix Ex Taq II (TAKARA, China) on LightCycler 480 PCR System (Roche,

Germany). The calculations were analyzed with $2^{-\Delta\Delta Ct}$ method.

Analysis of mean fluorescence intensity（MFI）of dual fluorescent protein

First, MFI of the dual fluorescent protein was analyzed using a flow cytometer. The

cells were treated with different radiation doses and collected at different time points,

washed once with PBS buffer, then assessed by flow cytometry (Beckman CytoFLEX,

USA). The data were analyzed using Flow Jo software. The ratio of the MFI of EGFP

to MFI of mecherry was used for calculation and analysis. Each irradiated dose sample

group was analyzed for 10,000 cells, and three dishes were repeated.

Furthermore, the High-Content Screening system Cell Insight CX5 HCS (Thermo

Fisher, Waltham, MA) was used for automatic photo quantitative analysis(Sun *et al*, 2016).

The dual-fluorescent protein HeLa cells were seeded in 96-well plates by irradiated 1

Gy, 2 Gy, 4 Gy gamma ray. The nuclei were stained with Hoechst 33342 (Invitrogen,

H3570). Fluorescence ratios of protein expression of mitochondrial expressed COX8-

EGFP and actin-mCherry were statistically analyzed by cellomics software (Thermo

Fisher, Waltham, MA).

Laser scanning confocal fluorescence microscopy

All the images were obtained using 488 nm and 561 nm lasers to excite cox8-

EGFP-actin-mCherry, and 405 nm laser lines were used to excite DAPI. The imaging

was performed in line mode, the z-stack was over-sampled and photographed at the

height of 0.7 microns, and the image was acquired on the LEICA SP8 confocal

microscope system. The contrast and brightness of the image were adjusted by

Application Suite X (LEICA).

Western blot analysis and immunofluorescence staining

Total proteins were extracted from cells using RIPA lysis buffer with a protease

inhibitor cocktail (Roche, Germany). The proteins were quantified using Thermo

Scientific Pierce BCA Protein Assay. The protein level of Bax, Bcl-2, caspase-3,

cleaved caspase-9, Cytoc, PARP, TFAM, Mfn1, PGC-1α, GFP and loading control β-

Actin were determined by Western blot assay.

For immunofluorescence staining, HeLa cells were grown on chamber slides. After

γ-ray irradiation or mock irradiation, they were fixed with cold methanol for 10 minutes

and permeated with 0.5% Triton X-100 in PBS for 20 minutes. After blocking for 1

hour, the cells were incubated with the primary antibody overnight and then incubated

with the fluorescently labeled secondary antibody for 1 hour at 37°C. Samples

investigated by laser confocal microscopy (LEICA). Image auto-quantification and co-

localized foci were recognized by CellProfiler version 4

(www.cellprofiler.org)(McQuin *et al*, 2018).

Statistical analysis

739 Data were presented as mean ± standard deviation (SD). Statistical significance was

740 assessed by one-way or two-way ANOVA via GraphPad Prism software (Prism version

741 8, San Diego, CA, USA). *P*-Value < 0.05 was considered as statistically significant.

Data availability

This study includes no data deposited in external repositories.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grants No. 11635013 and No. 11775272) and Special Repair and Purchase Fund for Central-level scientific institutions (No. Y79XG13361).

Author contributions

QH conceived the idea and supervised the research. CS and QH designed and performed the experiments, analyzed and interpreted data, performed statistical analysis, and wrote the manuscript. XZ, YM, PW and QZ provided technical assistance in the experiments. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

757 **Supplementary Information**

759 **Table S1**

760 Oligonucleotides for amplify PCR product of COX8

Oligonucleotides	Sequence 5′ → 3′	size (bp)
Oligo1	CTAGAGATCTGGTACCGTCGACTGCAG AGATCCGCTAGGGGTACTCCGTG CCATCATGTCCGTCCTGACGCCGC TGCTGCTGCGGGGCTTGACAGGC TCGGCCCGGCGGCTCCCAGTGCC GCGCGCCAAGATCCATTCGTTGC CGCCGGAGGGG	154
Oligo2	AGCTTCCCCTCCGGCGGCAACGAATGG ATCTTGGCGCGCGGCACTGGGAG CCGCCGGGCCGAGCCTGTCAAGC CCCGCAGCAGCAGCGGCGTCAG GACGGACATGATGGCACGGAGTA CCCCTAGCGGATCTCTGCAGTCGA CGGTACCAGATCT	155

Figure EV1. Western blot assay for verifying the effect of AICAR on mitochondrial biogenesis.

HeLa cells were treated with mitochondrial biogenesis inducer drug (AICAR, 0.5 mM), and the whole cell extracts were collected at different time points. The mitochondrial biogenesis marker proteins (PGC-1α, NRF1 and TFAM) are indicated in the plot.

Figure EV2. The mean fluorescence intensity (MFI) of HeLa cells was determined by flow cytometry.

HeLa-cox8-EGFP-actin-mCherry cells were exposed to 1 Gy，2 Gy，4 Gy gamma-ray and incubated for the indicated periods. The analysis and comparison MFI obtained from the dual-fluorescent measurements by flow cytometry.

S1 Video

S2 Video

S3 Video

In-situ real-time video of dual fluorescence reporter system of irradiated cells (Video S1, Video

S2, Video S3)

References:

Azzam EI, Jay-Gerin J-P & Pain D (2012) Ionizing radiation-induced metabolic oxidative stress
and prolonged cell injury. *Cancer Lett* 327: 48–60

Barros MH & McStay GP (2019) Modular Biogenesis of Mitochondrial Respiratory Complexes.
Mitochondrion 50: 94–114

Bernstein C, Bernstein H, Payne CM & Garewal H (2002) DNA repair/pro-apoptotic dual-role
proteins in five major DNA repair pathways: fail-safe protection against carcinogenesis. *Mutat*
Res Rev Mutat Res 511: 145–178

Betzig E, Patterson GH, Sougrat R, Lindwasser OW, Olenych S, Bonifacino JS, Davidson MW,
Lippincott-Schwartz J & Hess HF (2006) Imaging Intracellular Fluorescent Proteins at
Nanometer Resolution. *Science* 313: 1642–1645

Burke PJ (2017) Mitochondria, Bioenergetics and Apoptosis in Cancer. *Trends Cancer* 3: 857–
870

Cao C, Yu H, Wu F, Qi H & He J (2017) Antibiotic anisomycin induces cell cycle arrest and
apoptosis through inhibiting mitochondrial biogenesis in osteosarcoma. *J Bioenerg Biomembr*
49: 437–443

Chalkiadaki A & Guarente L (2015) The multifaceted functions of sirtuins in cancer. *Nat Rev*
Cancer 15: 608–624

Dam AD, Mitchell AS & Quadrilatero J (2013) Induction of mitochondrial biogenesis protects
against caspase-dependent and caspase-independent apoptosis in L6 myoblasts. *Biochimica Et*
Biophysica Acta Bba - Mol Cell Res 1833: 3426–3435

Das S, Joshi MB, Parashiva GK & Rao SBS (2020) Stimulation of cytoprotective autophagy and
components of mitochondrial biogenesis / proteostasis in response to ionizing radiation as a
credible pro-survival strategy. *Free Radical Bio Med*

Davis AJ, Chen BPC & Chen DJ (2014) DNA-PK: A dynamic enzyme in a versatile DSB repair
pathway. *Dna Repair* 17: 21–29

Fanibunda SE, Deb S, Maniyadath B, Tiwari P, Ghai U, Gupta S, Figueiredo D, Weisstaub N,
Gingrich JA, Vaidya ADB, *et al* (2019) Serotonin regulates mitochondrial biogenesis and
function in rodent cortical neurons via the 5-HT2A receptor and SIRT1–PGC-1α axis. *Proc*
National Acad Sci 116: 201821332

Fu X, Wan S, Lyu YL, Liu LF & Qi H (2008) Etoposide Induces ATM-Dependent Mitochondrial
Biogenesis through AMPK Activation. *Plos One* 3: e2009

Galluzzi L & Kroemer G (2008) Necroptosis: a specialized pathway of programmed necrosis. *Cell*
135: 1161–3

Haggie PM & Verkman AS (2002) Diffusion of Tricarboxylic Acid Cycle Enzymes in the
Mitochondrial Matrixin Vivo: EVIDENCE FOR RESTRICTED MOBILITY OF A
MULTIENZYME COMPLEX. *J Biol Chem* 277: 40782–40788

Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY & Remington SJ (2004)
Investigating Mitochondrial Redox Potential with Redox-sensitive Green Fluorescent Protein
Indicators. *J Biol Chem* 279: 13044–13053

Holler N, Zaru R, Micheau O, Thome M, Attinger A, Valitutti S, Bodmer J-L, Schneider P, Seed
B & Tschopp J (2000) Fas triggers an alternative, caspase-8–independent cell death pathway
using the kinase RIP as effector molecule. *Nat Immunol* 1: 489–495

Huangyang P, Li F, Lee P, Nissim I, Weljie AM, Mancuso A, Li B, Keith B, Yoon SS & Simon
MC (2020) Fructose-1,6-Bisphosphatase 2 Inhibits Sarcoma Progression by Restraining
Mitochondrial Biogenesis. *Cell Metab* 31: 1032

Jadhav U & Shivdasani RA (2019) Dissecting Cell Lineages: From Microscope to Kaleidoscope.
Cell 176: 949–951

Katajisto P, Döhla J, Chaffer CL, Pentinmikko N, Marjanovic N, Iqbal S, Zoncu R, Chen W,
Weinberg RA & Sabatini DM (2015) Stem cells. Asymmetric apportioning of aged
mitochondria between daughter cells is required for stemness. *Sci New York N Y* 348: 340–3

Kim J, Yang G, Kim Y, Kim J & Ha J (2016) AMPK activators: mechanisms of action and
physiological activities. *Exp Mol Medicine* 48: e224–e224

Kluza J, Marchetti P, Gallego M-A, Lancel S, Fournier C, Loyens A, Beauvillain J-C & Bailly C
(2004) Mitochondrial proliferation during apoptosis induced by anticancer agents: effects of
doxorubicin and mitoxantrone on cancer and cardiac cells. *Oncogene* 23: 7018–7030

Komen JC & Thorburn DR (2014) Turn up the power – pharmacological activation of
mitochondrial biogenesis in mouse models. *Brit J Pharmacol* 171: 1818–1836

Kulms D, Zeise E, Pöppelmann B & Schwarz T (2002) DNA damage, death receptor activation
and reactive oxygen species contribute to ultraviolet radiation-induced apoptosis in an
essential and independent way. *Oncogene* 21: 5844–5851

Leach JK, Tuyle GV, Lin PS, Schmidt-Ullrich R & Mikkelsen RB (2001) Ionizing radiation-
induced, mitochondria-dependent generation of reactive oxygen/nitrogen. *Cancer Res* 61:
3894–901

Lee H-C & Wei Y-H (2005) Mitochondrial biogenesis and mitochondrial DNA maintenance of
mammalian cells under oxidative stress. *Int J Biochem Cell Biology* 37: 822–834

Li S, Shi B, Liu X & An H-X (2020) Acetylation and Deacetylation of DNA Repair Proteins in
Cancers. *Frontiers Oncol* 10: 573502

Lill R & Freibert S-A (2020) Mechanisms of Mitochondrial Iron-Sulfur Protein Biogenesis. *Annu*
Rev Biochem 89: 471–499

Lin WT, Nithiyanantham S, Hsieh DJ, Chen R, Day C, Liao JY, Kuo C, Mahalakshmi B, Kuo W
& Huang C (2020) Bioactive peptides attenuate cardiac apoptosis in spontaneously
hypertensive rat hearts through activation of autophagy and mitochondrial biogenesis pathway.
Environ Toxicol 35: 804–810

Llopis J, McCaffery JM, Miyawaki A, Farquhar MG & Tsien RY (1998) Measurement of
cytosolic, mitochondrial, and Golgi pH in single living cells with green fluorescent proteins.
Proc National Acad Sci 95: 6803–6808

Löbrich M & Jeggo PA (2005) Harmonising the response to DSBs: a new string in the ATM bow.
Dna Repair 4: 749–759

Luo C, Widlund HR & Puigserver P (2016) PGC-1 Coactivators: Shepherding the Mitochondrial
Biogenesis of Tumors. *Trends Cancer* 2: 619–631

Mahajan NP, Linder K, Berry G, Gordon GW, Heim R & Herman B (1998) Bcl-2 and Bax
interactions in mitochondria probed with green fluorescent protein and fluorescence resonance
energy transfer. *Nat Biotechnol* 16: 547–552

Martins LAM, Vieira MQ, Ilha M, Vasconcelos M de, Biehl HB, Lima DB, Schein V, Barbé-
Tuana F, Borojevic R & Guma FCR (2015) The Interplay Between Apoptosis, Mitophagy and
Mitochondrial Biogenesis Induced by Resveratrol Can Determine Activated Hepatic Stellate
Cells Death or Survival. *Cell Biochem Biophys* 71: 657–672

May-Panloup P, Vignon X, Chrétien M-F, Heyman Y, Tamassia M, Malthièry Y & Reynier P
(2005) Increase of mitochondrial DNA content and transcripts in early bovine embryogenesis
associated with upregulation of mtTFA and NRF1 transcription factors. *Reprod Biol Endocrin*
3: 65

McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, Doan M, Ding
L, Rafelski SM, Thirstrup D, *et al* (2018) CellProfiler 3.0: Next-generation image processing
for biology. *Plos Biol* 16: e2005970

Melentijevic I, Toth ML, Arnold ML, Guasp RJ, Harinath G, Nguyen KC, Taub D, Parker JA,
Neri C, Gabel CV, *et al* (2017) C. elegans neurons jettison protein aggregates and
mitochondria under neurotoxic stress. *Nature* 542: 367–371

Mihaylova MM & Shaw RJ (2011) The AMPK signalling pathway coordinates cell growth,
autophagy and metabolism. *Nat Cell Biol* 13: 1016–1023

Miyashiro T & Goulian M (2007) Methods in Enzymology. *Methods Enzymol* 423: 458–475

Mukhopadhyay P, Rajesh M, Haskó G, Hawkins BJ, Madesh M & Pacher P (2007) Simultaneous
detection of apoptosis and mitochondrial superoxide production in live cells by flow cytometry
and confocal microscopy. *Nat Protoc* 2: 2295–2301

Newman RH, Fosbrink MD & Zhang J (2011) Genetically encodable fluorescent biosensors for
tracking signaling dynamics in living cells. *Chem Rev* 111: 3614–66

O'Driscoll M & Jeggo PA (2006) The role of double-strand break repair — insights from human
genetics. *Nat Rev Genet* 7: 45–54

Paredes S & Chua KF (2016) SIRT7 clears the way for DNA repair. *Embo J* 35: 1483–1485

Park S-J, Gavrilova O, Brown AL, Soto JE, Bremner S, Kim J, Xu X, Yang S, Um J-H, Koch LG,
et al (2017) DNA-PK Promotes the Mitochondrial, Metabolic, and Physical Decline that
Occurs During Aging. *Cell Metab* 25: 1135-1146.e7

Popov L (2020) Mitochondrial biogenesis: An update. *J Cell Mol Med* 24: 4892–4899

Rai PK, Russell OM, Lightowlers RN & Turnbull DM (2015) Potential compounds for the
treatment of mitochondrial disease. *Brit Med Bull* 116: 5–18

Rai Y, Pathak R, Kumari N, Sah DK, Pandey S, Kalra N, Soni R, Dwarakanath BS & Bhatt AN
(2018) Mitochondrial biogenesis and metabolic hyperactivation limits the application of MTT
assay in the estimation of radiation induced growth inhibition. *Sci Rep-uk* 8: 1531

Ruan L, Zhou C, Jin E, Kucharavy A, Zhang Y, Wen Z, Florens L & Li R (2017) Cytosolic
proteostasis through importing of misfolded proteins into mitochondria. *Nature* 543: 443–446

Scarpulla RC (2008) Transcriptional Paradigms in Mammalian Mitochondrial Biogenesis and
Function. *Physiol Rev* 88: 611–638

Spitz DR, Azzam EI, Li JJ & Gius D (2004) Metabolic oxidation/reduction reactions and cellular
responses to ionizing radiation: A unifying concept in stress response biology. *Cancer Metast
Rev* 23: 311–322

Srinivas US, Tan BWQ, Vellayappan BA & Jeyasekharan AD (2019) ROS and the DNA damage
response in cancer. *Redox Biol* 25: 101084

Sun J, Li N, Oh K-S, Dutta B, Vayttaden SJ, Lin B, Ebert TS, Nardo DD, Davis J, Bagirzadeh R,
et al (2016) Comprehensive RNAi-based screening of human and mouse TLR pathways
identifies species-specific preferences in signaling protein use. *Sci Signal* 9: ra3–ra3

Vayssiere JL, Petit PX, Risler Y & Mignotte B (1994) Commitment to apoptosis is associated
with changes in mitochondrial biogenesis and activity in cell lines conditionally immortalized
with simian virus 40. *Proc National Acad Sci* 91: 11752–11756

Virbasius CA, Virbasius JV & Scarpulla RC (1993) NRF-1, an activator involved in nuclear-
mitochondrial interactions, utilizes a new DNA-binding domain conserved in a family of
developmental regulators. *Gene Dev* 7: 2431–2445

Vyas S, Zaganjor E & Haigis MC (2016) Mitochondria and Cancer. *Cell* 166: 555–566

Westermann B & Neupert W (2000) Mitochondria-targeted green fluorescent proteins: convenient
tools for the study of organelle biogenesis inSaccharomyces cerevisiae. *Yeast* 16: 1421–1427

Wu Z, Puigserver P, Andersson U, Zhang C, Adelmant G, Mootha V, Troy A, Cinti S, Lowell B,
Scarpulla RC, *et al* (1999) Mechanisms Controlling Mitochondrial Biogenesis and Respiration
through the Thermogenic Coactivator PGC-1. *Cell* 98: 115–124

Yambire KF, Fernandez-Mosquera L, Steinfeld R, Mühle C, Ikonen E, Milosevic I & Raimundo
N (2019) Mitochondrial biogenesis is transcriptionally repressed in lysosomal lipid storage
diseases. *Elife* 8: e39598

Yu H, Zhao W, Xie M, Li X, Sun M, He J, Wang L & Yu L (2020) Real-Time Monitoring of Self-
Aggregation of β-Amyloid by a Fluorescent Probe Based on Ruthenium Complex. *Anal Chem*
92: 2953–2960

Yu J, Wang Q, Chen N, Sun Y, Wang X, Wu L, Chen S, Yuan H, Xu A & Wang J (2013)
Mitochondrial transcription factor A regulated ionizing radiation-induced mitochondrial
biogenesis in human lung adenocarcinoma A549 cells. *J Radiat Res* 54: 998–1004

Yu M (2011) Generation, function and diagnostic value of mitochondrial DNA copy number
alterations in human cancers. *Life Sci* 89: 65–71

Zhang T, Ikejima T, Li L, Wu R, Yuan X, Zhao J, Wang Y & Peng S (2017) Impairment of
Mitochondrial Biogenesis and Dynamics Involved in Isoniazid-Induced Apoptosis of HepG2
Cells Was Alleviated by p38 MAPK Pathway. *Front Pharmacol* 8: 753

948