Monitoring mitochondrial biogenesis as the early event for prediction of radiation-induced apoptosis

Chang-Sheng Shao [1,2], Xiu-Hong Zhou [3], Yu-Hui Miao [4], Qian-Qian Zhang [1,2] and Qing Huang [1,2] *

[1] CAS Key Laboratory of High Magnetic Field and Ion Beam Physical Biology, Hefei Institutes of Physical Science, Chinese Academy of Sciences (CAS), Hefei 230031, China;

[2] Science Island Branch of Graduate School, University of Science and Technology of China, Hefei 230026, China;

[3] Center of Biology, Anhui Agricultural University, Hefei, China;

[4] Department of Medical Oncology, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China;

*Corresponding author:

Prof. Qing Huang

Hefei Institutes of Physical Science, Chinese Academy of Sciences, P. O. Box 1138, Hefei 230031, P.R. China

Tel: 86-551-65595261; Fax: 86-551-65595261

E-mail: huangq@ipp.ac.cn

ABSTRACT

Mitochondrial biogenesis is a cell response to external stimuli which is generally believed to be negatively correlated with apoptosis. However, in the process of apoptosis, whether mitochondrial biogenesis occurs in the early period of the apoptotic cells exposed to the external stimuli such as radiation remains unclear. To address this question, we constructed the COX8-EGFP-ACTIN-mCherry HeLa cells with recombinant fluorescent proteins respectively tagged on the nucleus and mitochondria, and monitored the mitochondrial changes in living cells exposed to the gamma-ray radiation. Besides the in situ and real-time detection of mitochondrial fluorescence changes, we also examined the cell viability, nuclear DNA damage, mitochondrial DNA copy number and the expression of transcription genes of mitochondrial biogenesis and apoptosis marker proteins. As such, we confirmed that mitochondrial biogenesis is taking place preceding apoptosis, and enhancement of mitochondrial biogenesis at at early time is actually positively correlated with the apoptotic cells at late stage.

Keywords: mitochondrial biogenesis; apoptosis; fluorescent proteins; ionizing radiation; living cells; early events

Abbreviations: ANOVA, Analysis of Variance; CCK-8, Cell Counting Kit-8; CytoC, Cytochrome C; DMEM, Dulbecco's modified Eagle's medium;; *E. coli, Escherichia coli*; HCS, High-content screening; H2DCFDA, 2',7'-dichlorodihydrofluorescein diacetate; LDH, Lactate Dehydrogenase; MMP, Mitochondrial membrane potential; MFI, mean fluorescence intensity; Mfn1, Mitofusin 1; mtDNA, Mitochondrial DNA;

44   nDNA, nuclear DNA; PI, Propidium iodide; PARP, Poly ADP-Ribose Polymerase;

45   PGC-1α, Peroxisome proliferator-activated receptor-gamma coactivator (PGC)-1alpha;

46   RT-qPCR, Real-Time Quantitative Reverse Transcription PCR; ROS, reactive oxygen

47   species; SD, standard derivation; TMRE, Tetramethylrhodamine ethyl ester; TFAM,

48   transcription factor A, mitochondrial.

Introduction

Mitochondrial biogenesis is a physiological response of cells to external stress and development signals which cause increased energy demand, and it plays an important role in cell metabolism regulation, signal transduction and mitochondrial ROS regulation (Luo et al., 2016; Vyas et al., 2016). Mitochondrial biogenesis maintains cell homeostasis by ensuring the quality of mtDNA and regulating the renewal of organelles(Yambire et al., 2019), and it has been extensively explored in recent years due to its related interests in human aging, neurodegenerative diseases, cell metabolic diseases and tumors(Fanibunda et al., 2019).

Since mitochondrial regulation plays a very critical role in controlling the cell fate, it is intriguing for researchers to explore the relationship between mitochondrial biogenesis and apoptosis. Generally, apoptosis is mediated by the activation of the caspase pathways which are accompanied with mitochondrial damage with destruction of electron transfer, oxidative phosphorylation and ATP production, change of cell redox potential (Burke, 2017). In fact, some studies have claimed that there is a negative correlation between mitochondrial biogenesis and apoptosis, i.e., apoptosis can be suppressed by preceding mitochondrial biogenesis. For example, it has been reported that stress-induced apoptosis can be inhibited by increasing mitochondrial biogenesis (Dam et al., 2013), or, on the other hand, apoptosis and cell cycle arrest can be induced by inhibiting mitochondrial biogenesis (Cao et al., 2017). Also, in the study of radiation-induced apoptosis, mitochondrial biogenesis was observed and analyzed (Rai et al., 2018), and it was claimed that enhancement of mitochondrial biogenesis could reduce the proportion of apoptosis cuased by ionizing

radiation (Yu et al., 2013) . But considering the whole process of apoptosis, whether

mitochondrial biogenesis takes place in the early period in the apoptotic cells exposed

to the external stimuli such as radiation remains unclear.

To study the process of mitochondrial biogenesis prior to apoptosis, it is crucial to

monitor the mitochondrial changes in the cells. At present, there are not many

methods available to study the influence of external factors on mitochondrial changes.

One of the conventional methods is to detect the change of mitochondrial gene copy

numbers (Yu, 2011) . Another approach is to use fluorescent proteins. Fluorescent

proteins are beneficial for studying the living cells with reporting gene expression,

which can provide information on protein locations and expression levels as well as

the involved biochemical activities (Newman et al., 2011) . Actually, study of

mitochondria using fluorescent proteins is now emerging (Katajisto et al., 2015;

Melentijevic et al., 2017; Ruan et al., 2017). The more recent development is to make use

of multiple different fluorescent proteins with different excitation/emission spectra, so

that the precision of transcriptional measurements can be ensured by using the

reference color for normalization (Miyashiro and Goulian, 2007). In such a circumstance,

the ratio of dual fluorescence can be used to eliminate the possible fluorescence

measurement errors (Miyashiro and Goulian, 2007) . So with application of this dual

fluorescence approach researchers can observe and analyze the temporal and spatial

life trajectories of living cells (Jadhav and Shivdasani, 2019) , and maybe more

importantly, to scrutinize the early events leading to various bio-effects in the living

cells (Yu et al., 2020).

Therefore, in the present work, we attempted to apply the dual recombinant

fluorescent protein approach to monitor the mitochondrial biogenesis induced by

radiation, and in this way, to examine whether and how the early event of

mitochondrial biogenesis would be associated with apoptosis. For the effect of

radiation as the external stimuli to the cells, it is known that ionizing radiation (such

as α-, γ-, X-rays, protons, heavy ions) can easily cause various damages to living cells,

which can lead to the change of genetic information, cell mutations, genomic

instability, and apoptosis (O'Driscoll and Jeggo, 2006) . Mitochondria, as subcellular

organelles, are one of the main targets of ionizing radiation as they occupy a

considerable proportion of the cell volume (4-25%) (Leach et al., 2001) . On the other

hand, radiation can effectively induce mitochondrial biogenesis (Das et al., 2020) . For

the establishment of the dual fluorescent system for in-situ observation of

mitochondrial biogenesis in living cells, we constructed the stable fluorescent reporter

cell lines by lentiviral transfection to achieve mitoGFP and nuclear RFP, and so we

fulfilled the task of real-time in-situ observation of mitochondrial changes in the

living cells. As a result, we observed the effect of mitochondrial biogenesis occurring

in the early process of the apoptotic cells, and by analyzing fluorescence ratio of

mitoGFP and nuclear mCherry fluorescence, we could thus examine the relationship

between the early mitochondrial biogenesis and late apoptosis, and so discovered that

the increased mitochondria number in the early time was actually positively correlated

with the apoptotic cell population. The mitochondrial and apoptotic processes were

analyzed and the involved mechanism was discussed.

Results

Construction of dual-fluorescent stably transfected cell lines

In order to construct a universal dual-fluorescence gene reporter vector, we considered to make use of the simplicity of vector construction and referred to the previously reported dual-fluorescence reporter system, and so we chose the CMV promoter to express the dual-fluorescence target proteins. The mitochondrial reporter expression system is generally localized in the stroma, targeting the leader peptide (MSVLTPLLLRGLTGSARRLPVPRAKIHSLGDP) on the VIIIa subunit of cytochrome C (COX8) (Haggie and Verkman, 2002)(Figure 1A). As such, we constructed the pLVX-mCherry-actin lentiviral vector (Figure 1B).

Figure 1. Generation of the dual-fluorescent protein reporter system. (**A**) Schematic illustration of the pEGFP-cox8 reporter system. (**B**) Schematic illustration of the

pLVX-actin-mCherry reporter system. (**C**) Fluorescence images of HeLa cells transfected

with EGFP-tagged COX8. (**D**) Fluorescence images of HeLa cells transfected with

mCherry-tagged Actin.

To establish the cell line stably co-expressing COX8-EGFP and ACTIN-mCherry,

we transfected the HeLa cells with the plasmid containing the expression sequence of

COX8-EGFP, and obtained the COX8-EGFP expression cells. After screening by

G418 for 2 weeks, the stably transfected cell line was obtained. The

COX8-EGFP-mCherry-ACTIN co-expressing HeLa cell line was further established

based on the COX8-EGFP expressing cells with lentiviral transfection system. The

co-expressions of COX8-EGFP and mCherry-ACTIN were confirmed as shown by

the fluorescence microscope examination (Figure 1C, D).

In-situ observation with dual fluorescence reporter system

To corroborate our dual-fluorescence reporter system to detect the biological

effect in the irradiated living cells, the laser fluorescence confocal microscope was

employed. The COX8-EGFP encoding mitochondria (Green), the ACTIN-mCherry

encoding actin (Red), and the DAPI-stained nuclei were observed as shown in Figure

2.

Figure 2. HeLa-COX8-EGFP-ACTIN-mCherry cells exposed to gamma-radiation. The

cells were treated with different doses of γ-ray and observed with confocal laser scanning

microscope. In the figure, green color indicates the mitochondria encoding COX8-EGFP, red

color shows the actin encoded by actin-mCherry, and blue color represents the DAPI stained

nuclei. Scale bar: 25 μm.

The intracellular mitochondrial content gradually increased as confirmed by laser

confocal microscopy visually. Compared with the control group (0 Gy), the irradiated

group showed mitochondrial fluorescence enhancement, and it was significantly

increased for 2 and 4 Gy irradiation (Figure 2). The fluorescent images intuitively

showed that the number of mitochondria increased with increasing radiation dose,

confirming that the dual fluorescence reporter system could directly reflect the

relevant mitochondrial responses and changes in living cells.

Evaluation of the radiation-induced mitochondrial change using dual-fluorescent reporter system

To investigate the mitochondrial fluorescence enhancement of the cells receiving ionizing radiation, the mean fluorescence intensity (MFI) from dual fluorescence reporter system was utilized. Figure 3A shows the numerical changes of two fluorescence channels detected by flow cytometry. The MFI ratio of EGFP$^+$: mCherry$^+$ was calculated and it was found that the fluorescence ratio was positively correlated with the radiation dose. As shown in Figure 3A, there were significant differences between the irradiated and non-irradiated groups. Besides, this MFI ratio change was observed and recorded in real time by employing High-content Imaging (HCS) system, as shown by Figure 3B. This living cell workstation was employed to record the changes of cell fluorescence within 30 hours after cell irradiation, and the related videos were provided in the Supporting Information (Video 1, 2, 3). The analysis of the fluorescence data indicated that the fluorescence ratio increased significantly with the irradiation dose during the first interval of 1-6 hours, but afterwards, the fluorescence ratio decreased, and the 4 Gy group had a larger declining rate. Lower radiation dose corresponded to the lower rate of reduction of the fluorescence ratio (such as sham group and 1 Gy group). Therefore, the reduction in MFI suggested that the mitochondria of the irradiated cells with higher radiation doses were actually more damaged by the radiation.

Figure 3. The mean fluorescence intensity (MFI) of HeLa cells was determined by flow cytometry and High-content Imaging (HCS) Analysis system. HeLa-cox8-EGFP-actin-mCherry cells were exposed to 1 Gy，2 Gy，4 Gy gamma-ray and incubated for the indicated times. The MFI of HeLa cells by irradiation was determined by flow cytometry and HCS system. Normalized to the average MFI of the positive EGFP cells ratio on the average MFI of the positive mCherry cells. (**A**) The MFI of dual-fluorescent was detected by flow cytometry. (**B**) The MFI of HeLa cells by irradiation was recorded by HCS.

Radiation effects on cell viability, survival rate, cell cycle, apoptosis, DNA

damage and mitochondrial membrane damage

To investigate the radiation effect, the cell viability was measured by CCK-8

method. The results showed that cell viability decreased gradually with the increase of

radiation dose (Figure 4A). The toxic effect of radiation on cells was also examined

using LDH kit, which revealed that the radiation-induced toxicity was indeed

increased significantly with radiation dose (Figure 4B). Also, the colony-forming

method was used to analyze the inhibitory effect of γ-ray on HeLa cell survival. With

increase of γ-ray irradiation dose, the cell survival rate was only 30% after 4 Gy

irradiation (Figure 4C), confirming that the cell survival rate decreased with the rise

of radiation dose.

Figure 4. Detection of the radiation effect on HeLa cells with different radiation doses.
(**A**) CCK-8 assay for cell viability. (**B**) LDH assay for cell cytotoxicity. (**C**) Colony formation assay of HeLa cells after 14 days under different radiation doses. (**D**) Flow cytometry results of Annexin V-PI assay. (**E**) Apoptotic cells rate. (**F**) Cell cycle examination. The experiment was performed 3 times. Data are presented as Mean±SD; $P < 0.01$(**); $P < 0.001$(***); ns: not significant compared with untreated cells.

In addition, the cell cycle and apoptosis were assessed for the cells irradiated with γ-ray. Apoptotic cells were detected by flow cytometry after irradiation using Annexin V-PI methods (Figure 4D). Through statistics and comparison of the number of apoptotic cells, the results showed that with the increase of radiation dose, the number

of apoptotic cells increased significantly (Figure 4E). Cell cycle was analyzed by flow

cytometry. The results showed that all groups of cells had G2 arrest after 24 hours of

irradiation, especially at 2 Gy and 4 Gy irradiation doses showed significant G2 arrest

(Figure 4F).

Furthermore, in order to understand the radiation effects, the radiation induced

DNA and mitochondrial membrane damage as well the accompanied ROS change

were also examined. For such a purpose, the immunofluorescence method was used to

detect DNA damage, H_2DCFDA probe was used to label ROS, and TMRE was used

to label mitochondrial membrane potential. Figure 5A showed that with increase of

the radiation dose, the foci of the gamma-H2AX increased significantly, and the

statistical results showed that the ROS level in the irradiated cells increased with the

increase of the radiation dose (Figure 5B- D). At the dose of 2 Gy, the ROS level was

the highest, and for the 4 Gy group, there was a significant difference compared with

the non-irradiated group. In the meantime, the mitochondrial membrane potential

decreased significantly with the increase of radiation dose (Figure 5E, F).

Figure 5. Quantitative analysis of DNA damage level, ROS change level and mitochondrial membrane potential change level after irradiation. (**A**) Representative images of DNA damage detection at different radiation doses when Hela cells were exposed to radiation and measured instantly (within 30 minutes). Green: γ-H2AX; red: DNA stained with PI; Scale bars: 20 μm. (**B**) The graphs show the percentage of populations with the noted number of γ-H2AX foci with different irradiation doses. (**C-D**) ROS assessment using H2DCFDA assay at 6 h. (**E-F**) MMP assay using TMRE at 24 h.

Verification of mitochondrial biogenesis and apoptosis

Detection of copy number changes of mitochondrial DNA. To furhter verify the

mitochondrial biogenesis induced by radiation, the copy number of mitochondrial

DNA was examined. Total DNA was extracted from HeLa cells 6-24 hours after

irradiation. As shown in Figure 6, the copy number of mitochondrial DNA also

changed after irradiation of the cells. Indeed, under the same radiation dose,

mitochondrial DNA number increased with time and compared with the

non-irradiated group, and there showed the significant difference. Also, the analysis

and comparison of different irradiation groups at the same time point showed that the

number of mitochondrial DNA copy increased with radiation dose.

Figure 6. Effect of gamma-irradiation on mitochondrial DNA copy number. HeLa cells

were exposed to gamma rays at doses of 1 Gy, 2 Gy and 4 Gy. After 6, 12, 24 hours, total

genomic DNA was extracted and used as a template to detect the mtDNA/nDNA ratio by

RT-qPCR. Data are presented as Mean±SD; #: significant compared with unirradiated group;

**$P < 0.01$; ***$P < 0.001$.

Detection of the related protein expression levels. To explore the correlation

between mitochondrial biogenesis and apoptosis under the condition of radiation, the

related marker proteins were examined by Western analysis. For the mitochondrial

change, the related protein expression levels of PGC-1α, NRF1, TFAM and Mfn1 at

different times (6h, 12h, 24h) with different radiation doses (0 Gy, 1 Gy, 2 Gy, 4 Gy)

were assessed by Western blot (Figure 7). At 6 hours, the expression level of TFAM

protein was increased with radiation dose. On the other hand, for the apoptosis, the

related protein expression levels of apoptosis-related marker proteins (BAX, Bcl-2,

CytoC, Caspase-3, Cleaved Caspase-9, PARP) were examined. As shown in Figure 7,

the expression levels of crucial apoptosis proteins did not differ significantly at early

time (6 h), but their expression levels increased significantly afterwards (12 h, 24 h).

So these results suggested that there was positive correlation between mitochondrial

biogenesis in the early time (6h) and the apoptosis in the late time (24 h).

Figure 7. WB assay of the related proteins after radiation treatment. HeLa cells were treated with different doses of irradiation (0, 1, 2, 4 Gy), and total protein was extracted at different time points. Expression level of mitochondrial biogenesis marker proteins (PGC-1α, NRF1, TFAM and Mfn1) and apoptotic marker proteins (BAX, Bcl-2, CytoC, Caspase-3, Cleaved Caspase-9, PARP) by Western blotting.

Discussion

Mitochondria as the cell energy factory play a vital role in cell homeostasis(Lill and Freibert, 2020) In response to external stimuli, mitochondrial biogenesis can be elicited in order to provide the cells with extra energy (Barros and McStay, 2019) . Previously, many studies reported that mitochondrial biogenesis could suppress apoptosis(Yu et al., 2013) i.e., there would be negative correlation between mitochondrial biogenesis and apoptosis. Generally, these studies mainly focused on the cell response to the relatively small stimulus or stress from outside. What would happen if the external

stimulus or stress is large enough so that the final fate of apoptosis cannot be avoided?

Would mitochondrial biogenesis still go along with the apoptosis in this case? And if

it would occur, could we predict the fate of the cells by evaluating the mitochondrial

biogenesis as the early event of the cell fate such as apoptosis? With these questions,

we thus initiated this research, and applied radiation to induce both mitochondrial

biogenesis and apoptosis in the cells, to explore how the mitochondrial biogenesis

would be correlated with the apoptosis. As the consequences, we have the following

achievements, findings and elaborated understandings.

Firstly, we have constructed the dual fluorescence report system that can be

effectively used for mitochondrial biogenesis study. To probe the changes taking place

in mitochondria, we especially developed the approach using mitochondrial-tagged

recombinant fluorescent proteins. The fluorescent protein reporting system has the

advantages of real-time, convenient, fast, and non-destructive (Betzig et al., 2006) .

Besides, it has also an advantage in the study of subcellular organ localization and

protein stress regulation (Llopis et al., 1998) . Previously, people often used fluorescent

dye probes for studying effect involved with mitochondria. For example,

Mukhopadhyay et al. used Annexin V and Sytox Green and MitoSOX Red fluorescent

probes to study the detection of apoptosis and mitochondrial superoxide so as to

establish the method for detecting living cells by laser confocal scanning microscopy

and flow cytometry (Mukhopadhyay et al., 2007) . But the disadvantages of fluorescent

dyes are their toxicity to cells, as well as the light quenching under long-period

real-time observation. Compared to fluorescent dyes, fluorescent proteins are stable

and have minimal toxicity and can generate visible fluorescence in vivo without the

need for external cofactors (Westermann and Neupert, 2000). Actually, GFP reporter

protein is now commonly used to detect mitochondrial changes (Mahajan et al.,

1998) (Hanson et al., 2004). In this work, in particular, we constructed the dual

fluorescence reporting system based on fluorescence proteins for the monitoring of

mitochondrial changes. This dual fluorescent protein reporter system can not only for

the real-time in situ observation for a long time, but also help the precise fluorescence

evaluation with the reference fluorescent protein for intensity normalization.

Therefore, by analyzing the ratio of the fluorescence intensity of mitochondrial DNA

to nuclear DNA expression, the early event of mitochondrial biogenesis under

irradiation condition have been readily observed and evaluated, and with this

approach we can actually sensitively track the early process which eventually leads to

the late event of apoptosis.

Secondly, with the as-constructed dual fluorescence report system, we have

observed radiation-induced mitochondrial biogenesis preceding the cell apoptosis.

Ionizing radiation has been used for cancer therapy for they can directly or indirectly

cause DNA damage and kill the cancer cells (Spitz et al., 2004). With DNA damage in

the nucleus, cells may undergo cell cycle arrest required for repairing the damage or

cell death including apoptosis (Bernstein et al., 2002). In parallel, mitochondria is also

considered to be the key target for the radiation induced effects, where more reactive

oxygen species (ROS) could be produced to cause secondary damage to the cells.

Actually, it has been well documented that exposure of cells to ionizing radiation

could activate ROS to produce oxidases, regulate antioxidants, and alter metabolic

activity in response to oxidative damage (Azzam et al., 2012) . In both the nucleic or

mitochondrial processes, more ATP is required to elicit mitochondria biogenesis either

directly or indirectly (Kulms et al., 2002; Srinivas et al., 2019) Therefore, mitochondria

biogenesis could take place as one of the consequences of radiation-induced

bio-effects (Galluzzi and Kroemer, 2008). Interestingly, in our fluorescence measurements,

we found that the fluorescence ratio increased and reached the maximum at 6 h after

irradiation, but the fluorescence ratio gradually decreased afterwards. So, in the early

stage, the fluorescence ratio gradually increased, indicating that mitochondrial

biogenesis was enhanced. In this process of mitochondrial biogenesis, mitochondrial

DNA and mitochondrial mass are maintained to keep the cell homeostasis and

metabolism (Huangyang et al., 2020) . But in the late stage of the radiation effect, the

mitochondrial biogenesis gradually decreased. This may be due to the increase of

ROS caused by radiation, which continuously increased the oxidative stress that

damaged the mtDNA and the mitochondrial integrity. Generally, normal range of ROS

may induce stress response by changing the expression of related nuclear genes, and

certain level of ROS can save cells by maintaining energy metabolism (Srinivas et al.,

2019) . But when the ROS level exceeds the tolerant threshold, ROS can then induce

the transition of mitochondrial membrane permeability, leading to the activation of the

caspase pathway and promoting cell apoptosis. In our experiment, we indeed detected

the increase of ROS level and the decrease of mitochondrial membrane potential. In

the meantime, we detected the related caspase related proteins and found that the

increase of pro-apoptotic proteins such as cytochrome c. Release of cytochrome c is

an indicator for the apoptosis, in which the oxidative stress would damage mtDNA

and other components of the cells(Lee and Wei, 2005). Here, we especially noted that the

mitochondrial fluorescence intensity decrease rates were quite different after the cell

exposure to the radiation. The larger the irradiation dose, the higher the fluorescence

ratio reduction rate. This actually confirms the radiation-induced apoptotic process.

The mitochondrial damage can be regarded as an early event or signal for

apoptosis(Goldstein et al., 2000) As such, the use of dual fluorescence reporter system

applied in this work allowed us to observe in situ the cell dynamic process leading to

apoptosis, and we could localize fluorescent proteins on the mitochondria in the dual

fluorescence reporter system and distinguished the apoptosis process induced by

gamma-ray irradiation.

Thirdly, our results have revealed the subtle but critical relationship which had

not been reported previously. The mitochondrial biogenesis was increasingly

associated with the apoptotic stress, and the trend of fluorescence change was

consistent with the number of apoptotic cells, and this is actually a new phenomenon

contrary to the traditional view on the relationship between mitochondrial biogenesis

and apoptosis. In the past, there were few reports regarding the dynamic observation

of mitochondrial biogenesis in the early stage. The main reason may be due to lack of

in-situ observation method that can deliver the instant and reliable analysis of

mitochondrial biogenesis. For example, the traditional methods include using

real-time quantitative PCR to detect the copy number of mitochondrial DNA, or using

WB to identify the changes of marker protein level of mitochondrial biogenesis.

These methods cannot achieve in-situ and real-time observation in living cells. On the

contrary, the fluorescence protein method can be used to observe the mitochondrial

biogenesis in real time as especially useful for studying the early events prior to

apoptosis. Based on the fluorescence signal, the dual fluorescence ratio can help us to

predict the radiation effects or the cell fate according to the fluorescence change of

these irradiated cells. For example, from the data obtained from Fig. 3B, the

fluorescence ratios can be calculated, and we can retrieve the MFI ratios in each

irradiation dose group (0 Gy, 1 Gy, 2 Gy, 4 Gy) at an earlier time (e.g. 6 h), as shown

in Figure 8A. Strikingly, this result is consistent with the cells in the apoptotic state at

24 h (Figure 8B). It is understandable because the apoptosis is actually dependent on

the early biochemical events in the cells(Holler et al., 2000). In this way, we can therefor

infer or predict the late apoptotic state of cells by observing the changes in

mitochondrial fluorescence ratio in the early stage.

Figure 8. Analysis of mean fluorescence intensity and apoptosis percentage. (**A**) Analysis

of EGFP/mCherry ratio in High Content Screening (HCS) at 6h. (**B**) Analysis of apoptotic

cells percentage at 24h.

Lastly, we have also tried to understand the mechanism for the mitochondrial

biogenesis related to radiation induced apoptosis. For this purpose, we proved that

some marker proteins of mitochondrial biogenesis were involved in the biological

effects of radiation. We found that the protein and mRNA expression levels of nuclear

respiratory factor1 (NRF1), peroxisome proliferative activated receptor, gamma,

coactivator 1 alpha (PGC-1α), transcription factor A, mitochondrial (TFAM)

increased with increasing irradiation dose. NRF1 is a transcription factor with an vital

role in mitochondrial functional genes (Virbasius et al., 1993)　PGC-1α is a

transcriptional coactivator which is a key inducer of mitochondrial biogenesis (Wu et

al., 1999). TFAM is a key activator of mitochondrial (mt) DNA transcription. mtDNA

is reported to be very susceptible to oxidative stress(May-Panloup et al., 2005). With the

confirmation of these factors, we thus confirmed that this mitochondrial biogenesis is

indeed related to radiation induced cell adaptation and apoptosis.

So, with the above findings, we can therefore describe the process of radiation

affecting cells as depicted in Figure 9. First, after a certain dose of radiation, DNA

damage occurs, and the cell cycle is blocked in order to initiate the repair process. At

the same time, radiation produces a large amount of ROS in the cell and affects the

homeostasis of the cell. To resist the radiation stress or damage, mitochondria then

start to provide more ATP for DNA repair and maintenance of cell metabolism and

homeostasis, leading to mitochondrial biogenesis. Next, as ROS continues to increase,

mitochondrial damage occurs, leading to the decrease in membrane potential, which

finally leads to the release of the pro-apoptotic protein cytochrome c, activation of the

apoptosis caspase pathway, and so the programmed cell death.

Figure 9. Schematic diagram of biochemical events after ionizing radiation.

SUMMARY

We have constructed a dual fluorescence reporter system, and with the real-time

observation and the analysis of the cell fluorescence ratio, we have confirmed the

occurrence of mitochondrial biogenesis and apoptosis induced by radiation, and in

particular, we have investigated the relationship between this mitochondrial

biogenesis and apoptosis, and found there is positive correlation between them with

the checkpoint at early stage of cell process which leads to apoptosis. Compared with

previous work, this work not only demonstrates how to monitor the mitochondrial

biogenesis process but also suggest the way how to predict the radiation-induced

biological effects such as apoptosis using the early event observation.

Materials and methods

Reagents

The *Escherichia coli* DH5a and DNA Ladder were purchased from Transgen (Beijing, China). *Nhe*I, *Hind*III, *EcoR*I, *Not*I restriction enzymes, T4 DNA Ligase, and Q5 High-Fidelity DNA Polymerase were purchased from NEB (MA, USA). Plasmid pEGFP-N1 was purchased from Clonetech Biotechnology (CA, USA). Plasmid pLVX-mCherry, PUC57 were stored in our laboratory. Plasmids of psPAX2 (# 12260) and pMD.2G (#12259) were purchased from Addgene. PCR purification and gel extraction kits were obtained from Axygen (CA, USA). TRIzol™ reagent, Lipofectamine 2000 were purchased from Invitrogen (CA, USA). Primescript II 1st strand cDNA Synthesis Kit, RT-qPCR kit and Total genomic DNA extraction kit were purchased from TAKARA (Dalian, China). G418, DAPI was purchased from Beyotime Institute of Biotechnology (Shanghai, China). Reactive oxygen species (ROS) probe 2',7'-dichlorodihydrofluorescein diacetate (H2DCFDA, D6883) was purchased from Sigma-Aldrich. Tetramethylrhodamine ethyl ester (TMRE, T4057) was purchased from Useverbright. The antibodies of PGC-1α, NRF1, TFAM, Mfn1, BAX, Bcl-2, CytoC, Caspase-3, Cleaved Caspase-9, PARP, γH2AX(S139) were purchased from Bimake, anti-β-Actin antibody was purchased from CST(MA, USA).

Creation of stably transfected cell lines

Generation of plasmids. Firstly, the pEGFP-N1-COX8 expression plasmid was constructed. Briefly, the DNA sequence of cytochrome c oxidase 8A subunit was found from Genbank (NM_004074.2) (Hendrickson et al., 2010) . Then two complementary oligonucleotide strands were obtained by amplification using PCR, and the restriction sites of the *Nhe*I and *Hind*III were added at both ends. The sequences of oligonucleotide strands were listed in S1 Table. The DNA sequence (COX8A) was ligated on the PUC57 plasmid. Then, the appropriate oligonucleotides were annealed and ligated into pEGFP-N1 cloning cut with *Nhe*I and *Hind*III. After incubation overnight at 37°C, the target plasmids were extracted from the bacterial fluid using the EndoFree Maxi Plasmid kit.

Secondly, the PLVX-mCherry-Actin expression plasmid was constructed. Total RNA was extracted from HeLa cells by Trizol reagent and served as the template for cDNA synthesis, and then it was reversely transcribed using Primescript II 1st strand cDNA synthesis kit. The primers were designed according to the sequence of actin (Genbank no. NM_10277): *EcoR*I tailed forward (5′-gtcGAATTCatggatgatgatatcgccgcgc-3′) and *Not*I tailed reverse (5′-TATGCGGCCGCctagaagcatttgcggtggac-3′) primers which are named as Fw-actin-EcoRI and Rv-actin-NotI, respectively. The length of the amplified product was 1128 bp. The PCR product was separated by electrophoresis, and the target fragment was recovered after gel cutting and purification. The target gene fragment

(actin) and the expression vector (LVX-mCherry) with the same restriction site

(*EcoR*I and *Not*I) were digested by restriction enzymes respectively, and then ligated

with T4 ligase and transformed into *E. coli* DH5a for plasmid amplification.

Production   of   lentiviral   supernatant   and   transduction   of

COX8-EGFP-mCherry-ACTIN-HeLa.   Briefly,   positive   monoclonal   cells

(COX8-EGFP-HeLa cells) were constructed. When HeLa cells grew to the

concentration of 80% in complete medium, they were then transfected with

pEGFP-COX8 plasmid with lipofectamine 2000. After 24-48 h of transfection, G418

was added to screen positive monoclonal cells.

The lentiviral supernatant was prepared according to the protocol provided by

addgene (http://www.addgene.org/tools/protocols/plko/#A). When the confluence of

HEK293T cells cells in the 6-well plate reached 80%, Lipofectamine 2000 was used

to transfect the pLVX-mCherry-ACTIN plasmid and the viral packaging helper

plasmid psPAX2, pMD.2G into the cells. After incubation for 8h, the medium was

replaced with the fresh medium containing serum, and the supernatant was collected

24 hours and 48 hours later. The virus supernatant was filtered through a 0.45 μm

pore so as to infect COX8-EGFP-HeLa cells. Next, the COX8-EGFP-HeLa cells were

cultured in a 6-well plate to 70% confluence, then the medium was removed, and the

virus supernatant was mixed with fresh medium containing 8 μg/ml polybrene in

equal proportions and added to the culture plate. The fresh medium was replaced after

24 hours and the medium containing 1 μg/ml puromycin was changed at 48 hours to

start screening to obtain stable cell lines.

Cell Culture, Synchronization and Gamma irradiation

HeLa, HEK-293T cell line was purchased from American Type Cultures

Collection (ATCC, Manassas, VA). The HeLa and HEK-293T cells were cultured in

Dulbecco's modified Eagle's medium (DMEM, HyClone) containing 10% fetal bovine

serum (FBS, Gibco BRL). All cells were maintained at 37 °C in 5% CO_2.

The cell cycle synchronization method was detailed in the previous

works (Rodríguez-Martínez et al., 2020) . The cells were incubated with thymidine (2.5

mM) for 18 hours, washed once with PBS buffer and replaced with fresh medium for

14 hours, and then 2.5 mM thymidine was added again to the cells which were then

incubated for 18 hours. The efficiency of synchronization was tested by flow

cytometry using cell cycle analysis kit (Multisciences).

HeLa cells were gamma-ray irradiated with a Biobeam Cs137 irradiator (cat no.

GM 2000; Gamma-Service Medical, Leipzig, Germany) at 1, 2, and 4 Gy witho the

dose rate of 3.37 Gy/min. In the control group, the cells were taken out of the

incubator without irradiation.

 Cell viablity, LDH measurements and colony-formation assays

Cell viability was measured with CCK-8 assay kit (Dojindo, Japan). Briefly, 24 h

prior to irradiation treatment, ca. 6000 cells were plated in 96-well plates in a final

volume of 100 μl, and then cultured for 24h. And after adding the CCK-8 solution, the

plate was incubated for another 1 h. The optical density was measured at 450 nm

using a microplate reader (SpectraMax M5, Molecular Devices, USA). The amount of

LDH released into the medium was assayed using the LDH-Cytotoxicity Assay Kit II

(Dojindo, Japan) according to the manufacturer's instructions. For the colony

formation assay, 500 cells were seeded in 6-well plate. After being irradiated, the cells

were cultured at 37℃ for about 2 weeks. After carefully washing twice with PBS, 4%

paraformaldehyde was added to fix the cells. The the cells were then stained with

crystal violet solution and evaluated.

Analysis of mitochondrial membrane potential and ROS

TMRE-Mitochondrial membrane potential (MMP) assay kit (Useverbright, USA)

was used to measure MMP. The cells were incubated with culture medium containing

200 nM TMRE for 45 minutes at 37 °C, then rinsed with PBS 3 times, followed by

incubation with medium containing 50 nM TMRE. For the quantitative assessment,

the TMRE staining was evaluated using SpectraMax M5 microplate reader.

Reactive oxygen species (ROS) measurement was performed using H2DCFDA

(Sigma-Aldrich, USA) probe. 4×10^3 cells were seeded into 96-well plates and

incubated overnight. Before irradiation, the cells were incubated with HBSS solution

containing 5 μM H2DCFDA for 15 minutes and then irradiated. After 30 minutes, the

cells were washed 3 times, followed by the fluorescence measurement using a

microtiter plate reader.

523   Annexin V-FITC/PI staining and cell cycle analysis

The HeLa cells were seeded at 2×10^5 cell /well in 6-well plates and irradiation

with gamma-ray. After 24 h, the cells were fixed with cold ethanol and then stained

with propidium iodide. The cells (about 20,000) were analyzed by flow cytometry

(Beckman CytoFLEX, USA). The total cells were washed once and resuspended in

Annexin V binding buffer, and then stained with 5 μl Annexin V and 5μl PI (Roche,

Germany) for 20 min and analyzed with Beckman CytoFLEX flow cytometry. Ten

thousand events were collected for each sample.

531   Measurement of mtDNA copy number

Total genomic DNA of HeLa cells was extracted after irradiation using Total

genomic DNA extract kit (TAKARA, China). Human nuclear GAPDH gene (forward:

5′-   CAGAACATCATCCCTGCCTCTAC-3′;   reverse: 5′-

AAGGGTTGTAGTAGCCCGTAG-3′) and the mtDNA ND1 gene (forward:

5′-TTCTAATCGCAATGGCATTCCT-3′;   reverse: 5′-

AAGGGTTGTAGTAGCCCGTAG-3′), as well as a primer were used to detect the

mtDNA copy number. The real-time quantitative PCR analysis was performed using

SYBR Premix Ex Taq II (TAKARA, China) on LightCycler 480 PCR System (Roche,

Germany). The calculations were analyzed with $2^{-\Delta\Delta Ct}$ method.

Analysis of mean fluorescence intensity（MFI）of dual fluorescent protein

First, MFI of the dual fluorescent protein was analyzed using a flow cytometer.

The cells were treated with different radiation doses and collected at different time

points, washed once with PBS buffer, then assessed by flow cytometry (Beckman

CytoFLEX, USA). The data were analyzed using Flow Jo software. The ratio of the

MFI of EGFP to MFI of mecherry was used for calculation and analysis. Each

irradiated dose sample group was analyzed for 10,000 cells, and three dishes were

repeated.

Furthermore, the High-Content Screening system CellInsight CX5 HCS (Thermo

Fisher, Waltham, MA) was used for automatic photo quantitative analysis (Sun et al.,

2016) . The dual-fluorescent protein HeLa cells were seeded in 96-well plates by

irradiated 1 Gy, 2 Gy, 4 Gy gamma ray. The nuclei were stained with Hoechst 33342

(Invitrogen, H3570). Fluorescence ratios of protein expression of mitochondrial

expressed COX8-EGFP and actin-mCherry were statistically analyzed by cellomics

software (Thermo Fisher, Waltham, MA).

Laser scanning confocal fluorescence microscopy

All the images were obtained using 488 nm and 561 nm lasers to excite

cox8-EGFP-actin-mCherry, and 405 nm laser lines were used to excite DAPI. The

imaging was performed in line mode, the z-stack was over-sampled and photographed

at the height of 0.7 microns, and the image was acquired on the LEICA SP8 confocal

microscope system. The contrast and brightness of the image were adjusted by

Application Suite X (LEICA).

Western blot analysis and immunofluorescence staining

Total proteins were extracted from cells using RIPA lysis buffer with a protease

inhibitor cocktail (Roche, Germany). The proteins were quantified using Thermo

Scientific Pierce BCA Protein Assay. The protein level of Bax, Bcl-2, caspase-3,

cleaved caspase-9, Cytoc, PARP, TFAM, Mfn1, PGC-1α and loading control β-Actin

were determined by Western blot assay.

For immunofluorescence staining, cells were grown on chamber slides. After

gamma-ray irradiation or mock irradiation, they were fixed with cold methanol for 10

minutes and permeated with 0.5% Triton X-100 in PBS for 20 minutes. After blocking

for 1 hour, the cells were incubated with the primary antibody overnight and then

incubated with the fluorescently labeled secondary antibody for 1 hour at 37°C.

Statistical analysis

575 Data were presented as mean ± standard deviation (SD). Statistical significance

576 was assessed by one-way or two-way ANOVA via GraphPad Prism software (Prism

577 version 8, San Diego, CA, USA). P-Value < 0.05 was considered as statistically

578 significant.

Acknowledgments

The authors would like to thank Dr. J. Yan for some preliminary research related to this work. We would also like to thank Drs. Zhiming Zheng and Peng Wang at the Hefei Institutes of Physical Science, Chinese Academy of Sciences for extending their High-content imaging Analysis System facility and providing some consumables for this study. This work was supported by the National Natural Science Foundation of China (grants No. 11635013 and No. 11775272) and Special Repair and Purchase Fund for Central-level scientific institutions (No. Y79XG13361).

Supplementary Information

S1 Table

Oligonucleotides for amplify PCR product of COX8

Oligonucleotides	Sequence 5' → 3'	size (bp)
Oligo1	CTAGAGATCTGGTACCGTCGACTGCAG AGATCCGCTAGGGGTACTCCGTG CCATCATGTCCGTCCTGACGCCGC TGCTGCTGCGGGGCTTGACAGGC TCGGCCCGGCGGCTCCCAGTGCC GCGCGCCAAGATCCATTCGTTGCC GCCGGAGGGG	154
Oligo2	AGCTTCCCCTCCGGCGGCAACGAATGG ATCTTGGCGCGCGGCACTGGGAG CCGCCGGGCCGAGCCTGTCAAGC CCCGCAGCAGCAGCGGCGTCAG GACGGACATGATGGCACGGAGTA CCCCTAGCGGATCTCTGCAGTCGA CGGTACCAGATCT	155

609   S1 Video

610   S2 Video

611   S3 Video

613   In-situ real-time video of dual fluorescence reporter system of irradiated cells

References:

Azzam EI, Jay-Gerin J-P, Pain D. 2012. Ionizing radiation-induced metabolic oxidative stress and prolonged cell injury. *Cancer Lett* **327**:48–60. doi:10.1016/j.canlet.2011.12.012

Barros MH, McStay GP. 2019. Modular Biogenesis of Mitochondrial Respiratory Complexes. *Mitochondrion* **50**:94–114. doi:10.1016/j.mito.2019.10.008

Bernstein C, Bernstein H, Payne CM, Garewal H. 2002. DNA repair/pro-apoptotic dual-role proteins in five major DNA repair pathways: fail-safe protection against carcinogenesis. *Mutat Res Rev Mutat Res* **511**:145–178. doi:10.1016/s1383-5742(02)00009-1

Betzig E, Patterson GH, Sougrat R, Lindwasser OW, Olenych S, Bonifacino JS, Davidson MW, Lippincott-Schwartz J, Hess HF. 2006. Imaging Intracellular Fluorescent Proteins at Nanometer Resolution. *Science* **313**:1642–1645. doi:10.1126/science.1127344

Burke PJ. 2017. Mitochondria, Bioenergetics and Apoptosis in Cancer. *Trends Cancer* **3**:857–870. doi:10.1016/j.trecan.2017.10.006

Cao C, Yu H, Wu F, Qi H, He J. 2017. Antibiotic anisomycin induces cell cycle arrest and apoptosis through inhibiting mitochondrial biogenesis in osteosarcoma. *J Bioenerg Biomembr* **49**:437–443. doi:10.1007/s10863-017-9734-8

Dam AD, Mitchell AS, Quadrilatero J. 2013. Induction of mitochondrial biogenesis protects against caspase-dependent and caspase-independent apoptosis in L6 myoblasts. *Biochimica Et Biophysica Acta Bba – Mol Cell Res* **1833**:3426–3435. doi:10.1016/j.bbamcr.2013.04.014

Das S, Joshi MB, Parashiva GK, Rao SBS. 2020. Stimulation of cytoprotective autophagy and components of mitochondrial biogenesis / proteostasis in response to ionizing radiation as a credible pro-survival strategy. *Free Radical Bio Med*. doi:10.1016/j.freeradbiomed.2020.01.015

Fanibunda SE, Deb S, Maniyadath B, Tiwari P, Ghai U, Gupta S, Figueiredo D, Weisstaub N, Gingrich JA, Vaidya ADB, Kolthur-Seetharam U, Vaidya VA. 2019. Serotonin regulates mitochondrial biogenesis and function in rodent cortical neurons via the 5-HT2A receptor and SIRT1–PGC-1α axis. *Proc National Acad Sci* **116**:201821332. doi:10.1073/pnas.1821332116

Galluzzi L, Kroemer G. 2008. Necroptosis: a specialized pathway of
programmed necrosis. *Cell* **135**:1161–3.
doi:10.1016/j.cell.2008.12.004

Goldstein JC, Waterhouse NJ, Juin P, Evan GI, Green DR. 2000. The
coordinate release of cytochrome c during apoptosis is rapid, complete
and kinetically invariant. *Nat Cell Biol* **2**:156–162.
doi:10.1038/35004029

Haggie PM, Verkman AS. 2002. Diffusion of Tricarboxylic Acid Cycle Enzymes
in the Mitochondrial Matrixin Vivo: EVIDENCE FOR RESTRICTED MOBILITY
OF A MULTIENZYME COMPLEX. *J Biol Chem* **277**:40782–40788.
doi:10.1074/jbc.m207456200

Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, Remington
SJ. 2004. Investigating Mitochondrial Redox Potential with
Redox-sensitive Green Fluorescent Protein Indicators. *J Biol Chem*
279:13044–13053. doi:10.1074/jbc.m312846200

Hendrickson SL, Lautenberger JA, Chinn LW, Malasky M, Sezgin E, Kingsley
LA, Goedert JJ, Kirk GD, Gomperts ED, Buchbinder SP, Troyer JL, O'Brien
SJ. 2010. Genetic variants in nuclear-encoded mitochondrial genes
influence AIDS progression. *Plos One* **5**:e12862.
doi:10.1371/journal.pone.0012862

Holler N, Zaru R, Micheau O, Thome M, Attinger A, Valitutti S, Bodmer J-L,
Schneider P, Seed B, Tschopp J. 2000. Fas triggers an alternative,
caspase-8–independent cell death pathway using the kinase RIP as
effector molecule. *Nat Immunol* **1**:489–495. doi:10.1038/82732

Huangyang P, Li F, Lee P, Nissim I, Weljie AM, Mancuso A, Li B, Keith B,
Yoon SS, Simon MC. 2020. Fructose-1,6-Bisphosphatase 2 Inhibits
Sarcoma Progression by Restraining Mitochondrial Biogenesis. *Cell*
Metab **31**:1032. doi:10.1016/j.cmet.2020.04.009

Jadhav U, Shivdasani RA. 2019. Dissecting Cell Lineages: From Microscope
to Kaleidoscope. *Cell* **176**:949–951. doi:10.1016/j.cell.2019.01.054

Katajisto P, Döhla J, Chaffer CL, Pentinmikko N, Marjanovic N, Iqbal S,
Zoncu R, Chen W, Weinberg RA, Sabatini DM. 2015. Stem cells. Asymmetric
apportioning of aged mitochondria between daughter cells is required
for stemness. *Sci New York N Y* **348**:340–3. doi:10.1126/science.1260384

Kulms D, Zeise E, Pöppelmann B, Schwarz T. 2002. DNA damage, death receptor
activation and reactive oxygen species contribute to ultraviolet

radiation-induced apoptosis in an essential and independent way.
Oncogene **21**:5844–5851. doi:10.1038/sj.onc.1205743

Leach JK, Tuyle GV, Lin PS, Schmidt-Ullrich R, Mikkelsen RB. 2001.
Ionizing radiation-induced, mitochondria-dependent generation of
reactive oxygen/nitrogen. *Cancer Res* **61**:3894–901.

Lee H-C, Wei Y-H. 2005. Mitochondrial biogenesis and mitochondrial DNA
maintenance of mammalian cells under oxidative stress. *Int J Biochem
Cell Biology* **37**:822–834. doi:10.1016/j.biocel.2004.09.010

Lill R, Freibert S-A. 2020. Mechanisms of Mitochondrial Iron–Sulfur
Protein Biogenesis. *Annu Rev Biochem* **89**:471–499.
doi:10.1146/annurev-biochem-013118-111540

Llopis J, McCaffery JM, Miyawaki A, Farquhar MG, Tsien RY. 1998.
Measurement of cytosolic, mitochondrial, and Golgi pH in single living
cells with green fluorescent proteins. *Proc National Acad Sci*
95:6803–6808. doi:10.1073/pnas.95.12.6803

Luo C, Widlund HR, Puigserver P. 2016. PGC-1 Coactivators: Shepherding
the Mitochondrial Biogenesis of Tumors. *Trends Cancer* **2**:619–631.
doi:10.1016/j.trecan.2016.09.006

Mahajan NP, Linder K, Berry G, Gordon GW, Heim R, Herman B. 1998. Bcl-2
and Bax interactions in mitochondria probed with green fluorescent
protein and fluorescence resonance energy transfer. *Nat Biotechnol*
16:547–552. doi:10.1038/nbt0698-547

May-Panloup P, Vignon X, Chrétien M-F, Heyman Y, Tamassia M, Malthièry
Y, Reynier P. 2005. Increase of mitochondrial DNA content and
transcripts in early bovine embryogenesis associated with upregulation
of mtTFA and NRF1 transcription factors. *Reprod Biol Endocrin* **3**:65.
doi:10.1186/1477-7827-3-65

Melentijevic I, Toth ML, Arnold ML, Guasp RJ, Harinath G, Nguyen KC, Taub
D, Parker JA, Neri C, Gabel CV, Hall DH, Driscoll M. 2017. C. elegans
neurons jettison protein aggregates and mitochondria under neurotoxic
stress. *Nature* **542**:367–371. doi:10.1038/nature21362

Miyashiro T, Goulian M. 2007. Methods in Enzymology. *Methods Enzymol*
423:458–475. doi:10.1016/s0076-6879(07)23022-8

Mukhopadhyay P, Rajesh M, Haskó G, Hawkins BJ, Madesh M, Pacher P. 2007.
Simultaneous detection of apoptosis and mitochondrial superoxide

production in live cells by flow cytometry and confocal microscopy.
Nat Protoc **2**:2295–2301. doi:10.1038/nprot.2007.327

Newman RH, Fosbrink MD, Zhang J. 2011. Genetically encodable fluorescent
biosensors for tracking signaling dynamics in living cells. *Chem Rev*
111:3614–66. doi:10.1021/cr100002u

O'Driscoll M, Jeggo PA. 2006. The role of double-strand break repair —
insights from human genetics. *Nat Rev Genet* **7**:45–54.
doi:10.1038/nrg1746

Rai Y, Pathak R, Kumari N, Sah DK, Pandey S, Kalra N, Soni R, Dwarakanath
BS, Bhatt AN. 2018. Mitochondrial biogenesis and metabolic
hyperactivation limits the application of MTT assay in the estimation
of radiation induced growth inhibition. *Sci Rep-uk* **8**:1531.
doi:10.1038/s41598-018-19930-w

Rodríguez-Martínez M, Hills SA, Diffley JFX, Svejstrup JQ. 2020.
Multiplex Cell Fate Tracking by Flow Cytometry. *Methods Protoc* **3**:50.
doi:10.3390/mps3030050

Ruan L, Zhou C, Jin E, Kucharavy A, Zhang Y, Wen Z, Florens L, Li R. 2017.
Cytosolic proteostasis through importing of misfolded proteins into
mitochondria. *Nature* **543**:443–446. doi:10.1038/nature21695

Spitz DR, Azzam EI, Li JJ, Gius D. 2004. Metabolic oxidation/reduction
reactions and cellular responses to ionizing radiation: A unifying
concept in stress response biology. *Cancer Metast Rev* **23**:311–322.
doi:10.1023/b:canc.0000031769.14728.bc

Srinivas US, Tan BWQ, Vellayappan BA, Jeyasekharan AD. 2019. ROS and the
DNA damage response in cancer. *Redox Biol* **25**:101084.
doi:10.1016/j.redox.2018.101084

Sun J, Li N, Oh K-S, Dutta B, Vayttaden SJ, Lin B, Ebert TS, Nardo DD,
Davis J, Bagirzadeh R, Lounsbury NW, Pasare C, Latz E, Hornung V, Fraser
IDC. 2016. Comprehensive RNAi-based screening of human and mouse TLR
pathways identifies species-specific preferences in signaling protein
use. *Sci Signal* **9**:ra3–ra3. doi:10.1126/scisignal.aab2191

Virbasius CA, Virbasius JV, Scarpulla RC. 1993. NRF-1, an activator
involved in nuclear-mitochondrial interactions, utilizes a new
DNA-binding domain conserved in a family of developmental regulators.
Gene Dev **7**:2431–2445. doi:10.1101/gad.7.12a.2431

Vyas S, Zaganjor E, Haigis MC. 2016. Mitochondria and Cancer. *Cell*
166:555–566. doi:10.1016/j.cell.2016.07.002

Westermann B, Neupert W. 2000. Mitochondria-targeted green fluorescent
proteins: convenient tools for the study of organelle biogenesis
inSaccharomyces cerevisiae. *Yeast* **16**:1421–1427.
doi:10.1002/1097-0061(200011)16:15<1421::aid-yea624>3.0.co;2-u

Wu Z, Puigserver P, Andersson U, Zhang C, Adelmant G, Mootha V, Troy A,
Cinti S, Lowell B, Scarpulla RC, Spiegelman BM. 1999. Mechanisms
Controlling Mitochondrial Biogenesis and Respiration through the
Thermogenic Coactivator PGC-1. *Cell* **98**:115–124.
doi:10.1016/s0092-8674(00)80611-x

Yambire KF, Fernandez-Mosquera L, Steinfeld R, Mühle C, Ikonen E,
Milosevic I, Raimundo N. 2019. Mitochondrial biogenesis is
transcriptionally repressed in lysosomal lipid storage diseases. *Elife*
8:e39598. doi:10.7554/elife.39598

Yu H, Zhao W, Xie M, Li X, Sun M, He J, Wang L, Yu L. 2020. Real-Time
Monitoring of Self-Aggregation of β-Amyloid by a Fluorescent Probe
Based on Ruthenium Complex. *Anal Chem* **92**:2953–2960.
doi:10.1021/acs.analchem.9b03566

Yu J, Wang Q, Chen N, Sun Y, Wang X, Wu L, Chen S, Yuan H, Xu A, Wang J.
2013. Mitochondrial transcription factor A regulated ionizing
radiation-induced mitochondrial biogenesis in human lung
adenocarcinoma A549 cells. *J Radiat Res* **54**:998–1004.
doi:10.1093/jrr/rrt046

Yu M. 2011. Generation, function and diagnostic value of mitochondrial
DNA copy number alterations in human cancers. *Life Sci* **89**:65–71.
doi:10.1016/j.lfs.2011.05.010

785