A negative feedback model to explain regulation of SARS-CoV-2 replication and transcription

Shan Gao[1$], Zhi Cheng[2$], Xiufeng Jin[1], Fang Wang[3], Yibo Xuan[4]

Hao Zhou[1], Chang Liu[1], Jishou Ruan[5], Guangyou Duan[6]*, Xin Li[1]*

[1] College of Life Sciences, Nankai University, Tianjin, Tianjin 300071, P.R.China;
[2] State Key Laboratory of Medicinal Chemical Biology, Nankai University, Tianjin, Tianjin 300071, P.R.China;
[3] The Second Hospital of Tianjin Medical University, Tianjin 300211, P. R. China;
[4] College of Life Sciences, Hebei Normal University, Shijiazhuang, Hebei 050024, P.R.China;
[5] School of Mathematical Sciences, Nankai University, Tianjin, Tianjin 300071, P.R.China;
[6] School of Life Sciences, Qilu Normal University, Jinan, Shandong 250200, P.R.China

[$] These authors contributed equally to this paper.

* Corresponding authors.

XL：lix1980@nankai.edu.cn

GD：guangyou.duan@qlnu.edu.cn

Abstract

Background: Coronavirus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Although a preliminary understanding of the replication and transcription mechanisms of SARS-CoV-2 has recently emerged, their regulation remains unclear.

Results: Based on reanalysis of public data, we propose a negative feedback model to explain the regulation of replication and transcription in—but not limited to—SARS-CoV-2. The key step leading to new discoveries was the identification of the cleavage sites of nsp15—an RNA uridylate-specific endoribonuclease, encoded by CoVs. According to this model, nsp15 regulates the synthesis of subgenomic RNAs (sgRNAs) and genomic RNAs (gRNAs) by cleaving transcription regulatory sequences in the body. The expression level of nsp15 determines the relative proportions of sgRNAs and gRNAs, which in turn change the expression level of nps15 to reach equilibrium between the replication and transcription of CoVs.

Conclusions: The replication and transcription of CoVs are regulated by a negative feedback mechanism that influences the persistence of CoVs in hosts. Our findings enrich fundamental knowledge in the field of gene expression and its regulation, and provide new clues for future studies. One important clue is that nsp15 may be an important and ideal target for the development of drugs (e.g. uridine derivatives) against CoVs.

Keyword: coronavirus; transcription; replication; regulation model; nsp15

Introduction

Coronavirus disease 2019 (COVID-19) is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). As enveloped viruses composed of a positive-sense, single-stranded RNAs, CoVs have the largest genomes (26–32 kb) among all RNA virus families. SARS-CoV-2 has a genome of ∼30 kb [1]. In addition to ORF1a and 1b (**Figure 1A**), the SARS-CoV-2 genome has sequences encoding four conserved structural proteins [spike protein (S), envelope protein (E), membrane protein (M), and nucleocapsid protein (N)] and six accessory proteins (ORF3a, 6, 7a, 7b, 8, and 10) that are yet to be experimentally verified.

Figure1 Replication and transcription of SARS-CoV-2

The elements used to represent the SARS-CoV-2 genome (GenBank: MN908947.3) were originally used in the previous study [2]. **A**. TRS-L: transcription regulatory sequence in the body; TRS-B: transcription regulatory sequence in the leader. **B**. gRNA(+):genomic RNAs; gRNAs(-): antisense genomic RNAs.

The genomic RNA, as a positive-sense RNA [gRNA(+)], is used as a template for the translation of ORF1a and ORF1b, and the replication and transcription of the SARS-CoV-2 genome. Recently, the translation of 10 other proteins (S, E, M, N, ORF3a, 6, 7a, 7b, 8 and 10) of SARS-CoV-2 has been reported in a study [2] that

directly validates the prevailing "leader-to-body fusion" model using Nanopore

RNA-seq—a direct RNA sequencing method [3]. In the model validated in that study

(**Figure 1A**), replication and transcription require gRNA(+) as the template for the

synthesis of antisense genomic RNAs [gRNAs(-)] and antisense subgenomic RNAs

[sgRNAs(-)] by RNA-dependent RNA polymerase (RdRP). When RdRP pauses, as it

crosses a body transcription regulatory sequence (TRS-B) and switches the template

to the leader TRS (TRS-L), sgRNAs(-) are formed. Otherwise, RdRP reads gRNAs(+)

continuously, without interruption, resulting in gRNAs(-). Thereafter, gRNAs(-) and

sgRNAs(-) are used as templates to synthesize gRNAs(+) and sgRNAs(+),

respectively; sgRNAs(+) are used as templates for the translation of the 10 proteins

mentioned above.

TRS-L usually forms the first 60–70 nts of the 5' UTR in a CoV genome. TRS-B

with varied length is located immediately upstream of ORFs except ORF1a and 1b

(**Figure 1A**). During antisense-strand synthesis, the discontinuous transcription

(referred to as polymerase jumping or template switching) by RdRP is described by

the "leader-to-body fusion" model that is conserved in the entire nidovirus order,

including arteriviruses, toroviruses, roniviruses and animal coronaviruses. Although

investigation was conducted to determine the underlying mechanisms in previous

studies [4], the molecular basis remain unknown. In the present study, we aimed to

determine the molecular basis of the "leader-body fusion" model and construct a

model to explain the regulation of SARS-CoV-2 replication and transcription.

Results

Reanalysis of leader-body junction regions

Using public Nanopore RNA-seq data (**Materials and Methods**), 575,106 out of

879,679 reads from a SARS-CoV-2-infected Vero cell sample were aligned to the

SARS-CoV-2 reference genome. Among all aligned reads, 575,106 sense reads

represented gRNAs(+) or sgRNAs(+), while 30 antisense reads represented gRNAs(-)

or sgRNAs(-). This exceedingly high ratio (575,106 vs. 30) between sense and

antisense reads may be the result of significant differences between the

gRNAs(+)/sgRNAs(+) and gRNAs(-)/sgRNAs(-) degradation efficiencies. This

phenomenon, however, was not reported in the previous study [2]. Another

explanation for the high ratio is that gRNAs(+)/sgRNAs(+) are protected by binding

to the N proteins. Another super high ratio (198,198,542 vs. 11,820,438) between

contiguous and junction-spanning reads was reported in that previous study [2]. This

suggested that there were significant differences between gRNAs(+)/gRNAs(-) and

sgRNAs(+)/sgRNAs(-).

By reanalysis of junction-spanning reads, we found that TRS-Bs share ~12 nt

junction regions with TRS-Ls in SARS-CoV-2 (**Figure 1A**). Most junction regions in

eight protein-coding genes (*S, E, M, N, ORF3a, 6, 7a and 8*) exhibit high similarity to

those of TRS-Ls and are defined as canonical junction regions [2]. Accordingly, the

sgRNAs(+) and sgRNAs(-) containing these canonical junction regions are defined as

canonical sgRNAs(+) and sgRNAs(-), respectively. The junction regions of *ORF7b*

exhibit high diversity in their sequences. Whether junction regions exist in *ORF10*

remains unclear [2]. Our analysis of canonical junction regions determined their core

sequence (CS) of SARS-CoV-2 to be ACGAAC (**Figure 1A**).

Further investigation of "leader-to-body fusion" model

Further analysis of betacoronavirus subgroup B (**Materials and Methods**)

showed that ACGAAC is a highly conserved sequence among their genomes,

particularly in their putative TRS-Bs. This suggests that canonical junction regions

contain a specific motif for enzyme reorganization. Since all CoVs and even viruses

in the entire nidovirus order adhere to the "leader-to-body fusion" model [2], this or

these enzymes should be encoded by the *ORF1a* or *1b* gene, given their likelihood to

be translated. After analysing 16 non-structural proteins (nsp1–16) encoded by the

ORF1a or *1b* genes, we determined that nsp15 (a nidoviral RNA uridylate-specific

endoribonuclease, NendoU [5]) is most likely to function in these junction regions,

given that a homolog of nsp15 has cleavage sites containing "GU" [6]. Thus, the

cleavage site of nsp15 was identified to follow the motif "GTTCGT|N" (the vertical

line indicates the breakpoint and N indicates any nucleotide base), read in the

antisense strands of CoV genomes. Furthermore, we found that almost all the genomic

sites containing the motif "GTTCGT" have polyT (not less than three T) at the tail,

which ensures the presence of at least one U for nsp5 cleavage.

128   Table1   The genomic sites of motif GTTCGT

Motif	Position	Region(Start-end)	Type
GTTCGT	16014	nsp12(13483-16236)*	ICS
	28198	ORF8(27894-28259)	ICS
	28233	ORF8(27894-28259)	ICS
ACGAAC	70	5' UTR(1-265)	TRS-L
	21556	S(21563-25384)	TRS-B
	25385	ORF3a(25393-26220)	TRS-B
	26237	E(26245-26472)	TRS-B
	26473	M(26523-27191)	TRS-B
	27041	ORF6(27202-27387)	TRS-B
	27388	ORF7a(27394-27759)	TRS-B
	27888	ORF8(27894-28259)	TRS-B
	28260	N(28274-29533)	TRS-B

The motifs (in the first column) were mapped to the SARS-CoV-2 genome (GenBank: MN908947.3)
using positions (in the second column) of the first nucleotides. ACGAAC indicated the motif
GTTCGT, read on the antisense strand of the SARS-CoV-2 genome. The third column shows the
regions influenced by the motifs (in the first column). * this site is located in the CDS of RdRP
(nsp12). TRS-L: transcription regulatory sequence in the body; TRS-B: transcription regulatory
sequence in the leader; ICS: internal cleavage site; RdRP: RNA-dependent RNA polymerase.

Upon searching for "GTTCGT" in the genomes of betacoronavirus subgroup B,

the occurrence of "GTTCGT" on the antisense strand was found to be more than 1.6

times that on the sense strand. In particular, "GTTCGT" occurred 3 and 9 times

(**Table 1**) on the sense and antisense strands of the SARS-CoV-2 genome,

respectively. These findings suggest that the basic function of nsp15 involves the

degradation of gRNAs and gsRNAs and that the high ratio between sense and

antisense reads (**see above**) results from substantially more cleavage of

gRNAs(-)/gsRNAs(-) than that of gRNAs(+)/gsRNAs(+). Among the three sites

containing "GTTCGT" on the sense strand of the SARS-CoV-2 genome (referred to

as internal cleavage sites—ICSs), one is located in the coding sequence (CDS) of

RdRP (nsp12), while the other two are located in the *ORF8* gene (**see below**).

These two ICSs are also located in *ORF8* of most SARS-CoV-2, SARS2-like

CoV and SARS-like CoV genomes; however, they are absent in the genomes of

SARS-CoVs obtained from humans (GenBank: AY274119 and AY278489) and

SARS-like CoV genomes from civets (GenBank: AY304486, AY515512 and

AY572034). One of the two ICSs is present in the genome of the SARS-like CoV

strain WIV1 from bats (GenBank: KF367457), which is the ancestor of SARS-CoV.

Deletions of *ORF8* were reported to be associated with attenuation of SARS-CoV

(GenBank: AY274119) [8] and SARS-CoV-2 [10]. The *ORF8* gene of SARS-CoV is

considered to have played a significant role in adaptation to human hosts following

interspecies transmission [7] via the modification of viral replication [8]. The loss of

two nsp15 ICSs in ORF8 of SARS-CoV is the key clue revealing the functions of

ORF8 and the pandemic of SARS-CoV.

Based on the above analyses, we modified the "leader-body fusion" model and

proposed the molecular basis of it. In our model, nsp15 cleaves synthesized gRNAs(-)

and sgRNAs(-). The cleavage occurs at TRS-Bs(-) synthesized using TRS-Bs as

templates. Next, the free 3' ends (~6 nts) of TRS-Bs(-) hybridize the junction regions

of TRS-Ls for template switching. Alternatively, the cleaved TRS-Ls(-) are used as

templates to synthesize TRS-Ls that then hybridize the junction regions of the cleaved

sgRNAs(-) without TRS-Ls(-) for template switching and gRNAs(+)/sgRNAs(+)

synthesis. These findings suggests that RdRP (nsp12) is so active that it must have a

very high enzyme activity.

Some non-canonical sgRNAs(+) and sgRNAs(-) are synthesized as a result of

occasional hybridization between the free 3' ends of TRS-Bs(-) and highly similar

sequences of TRS-L junction regions. The sgRNAs(-) without TRS-Ls are also

synthesized due to missing hybrids. In the previous study [2], sgRNAs(-) without

TRS-Ls, non-canonical sgRNAs(+) and sgRNAs(-) were reported, supporting our

model. Furthermore, cleavage also occurs at ICSs on the sense strands, after which the

free 3' ends of the cleaved 5' ends hybridize the antisense strands to synthesize

recombinant sgRNAs(+) or even gRNAs(+). This may also contributes to the multiple

recombination events in betacoronavirus genomes.

Proposal of a negative feedback model

Based on the above analyses, we proposed a negative feedback model (**Figure**

1B) to explain the regulation of replication and transcription in—but not limited

to—CoVs. In this model (**Figure 1**), nsp15 regulates the synthesis of subgenomic

RNAs (sgRNAs) or genomic RNAs (gRNAs) by the cleavage of TRS-Bs(-). The

expression level of nsp15 determines the relative proportions of sgRNAs and gRNAs.

An increase in nsp15 expression results in less gRNAs(-) and more gsRNAs(-), after

which fewer gRNAs(+) and more gsRNAs(+) are synthesized, respectively. A

decrease in gRNAs(+) results in a decrease of nsp15 expression, given that nsp15 is

synthesized using gRNA(+) as the template. Furthermore, the nsp15 ICS in the CDS

of SARS-CoV-2 nsp12 (**Table 1**) enhances the negative feedback. Via this negative

feedback mechanism, CoVs reach equilibrium between the replication and

transcription (**Figure 1B**); thus, this mechanism is important for the persistence of

CoVs in hosts.

Our negative feedback model is based on the determination of the molecular

basis of the "leader-body fusion" model. This molecular basis is associated to the

cleavage function of nsp15, which is different from other models proposed in the

previous studies [4], mainly due to the integration of information from many aspects,

particularly considering: (1) The direct RNA sequencing data [2]; (2) The nsp15

structure in complex with GpU (pdb code: 6X1B); (3) the polyT at the tail of

"GTTCGT"; and (4) the nsp15 ICSs in *ORF8* (**see above**). These discoveries

confirmed the identification of the cleavage sites of nsp15. However, they were not

used to construct the models in the previous studies [4].

The necessity of negative feedback regulation

In our previous study [7], we proposed that first hairpins (immediately upstream

of ORF1a) have an important role in the functions (e.g. regulation of translational

initiation) of ribosome binding sites (RBSs) in 5' UTRs of the SARS-CoV-2 genomes.

SARS-CoV and SARS-CoV-2 have an identical first hairpin, which may enhance the

translation of downstream genes. To indirectly prove that negative feedback is a basic

mechanism, we designed preliminary experiments to show that first hairpins from

SARS-CoV-2 enhance the translation of the downstream enhanced green fluorescent

proteins (EGFPs); however, over-expression of EGFP without negative feedback

regulation will cause cell death.

212   Figure2 Regulation of EGFP expression by three RBSs

A. The plasmid pEGFP-C1 was used as control, since it contains 17-nt sequences, encoding the first

hairpin from Cytomegalovirus (CMV). Two other types of plasmids pSARS and pCoV-ba were

proceeded by replacement of first hairpins from SARS-CoV-2 and betacoronavirus subgroup A,

respectively (**Materials and Methods**). **B**. HEK293 cells were transfected with three types of plasmids.

At each time after transfection, cell viability and cell death were meausured using MTT and LDH,

respectively.

In total, three types of plasmids containing EGFP reporter genes—named

pEGFP-C1, pSARS, and pCoV-ba (betacoronavirus subgroup A)—were used in the

experiments (**Figure 2A**). The plasmid pEGFP-C1 was used as a control, given that it

contains 17-nt sequences, encoding the first hairpin from Cytomegalovirus (CMV).

Two types of plasmids proceeded by 30- and 29-nt inserts were used to evaluate their

influence on translation (**Materials and Methods**). These two inserts encoded the two

first hairpins from SARS-CoV-2 and the subgroup A of betacoronaviruses,

respectively. Comparing the fluorescent brightness of cells transfected with three

types of plasmids, the expression of EGFP in pSARS was markedly higher than that

in pEGFP-C1 and pCoV-ba (**Supplementary 1**). Moreover, pSARS caused cell death

at   48   hours   after   plasmid   transfection.   We   then   performed

3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2-H-tetrazolium   bromide   (MTT)   and

lactate dehydrogenase (LDH) experiments to further evaluate the influence of plasmid

transfection (**Materials and Methods**). Both MTT and LDH experiments consistently

suggested that pSARS caused significantly more HEK293T and Hela cell death at 48

hours after plasmid transfection, than pEGFP-C1 and pCoV-ba. However, this

phenomenon appeared in HEK293 cells at 56 hours after plasmid transfection (**Figure**

2B). Given that the only difference among the three types of plasmids is their 17-, 30-,

and 29-nt sequences encoding different hairpins, we concluded that these hairpins

determined the translation efficiency of the downstream EGFPs. The hairpin in

pSARS resulted in the over-expression of EGFP, which caused more cell death. To

determine whether the factor acts at the translation level and to rule out other possible

factors that may exert influence at the replication or transcription level, we performed

the following experiments: (1) using HEK293 cells to rule out the possible influence

by the differences of plasmid copy numbers, since all three types of plasmids

containing the SV40 origins can be replicated to a copy number of between 400~1000

plasmids per cell within HEK293T; and (2) using qPCR to rule out the possible

influence by differential transcription (**Supplementary 1**).

Conclusion and Discussion

In the present study, we determined the molecular basis of the "leader-body fusion" model and constructed a negative feedback model to explain the regulation of SARS-CoV-2 replication and transcription. The key step that led us to new discoveries was our identification of the cleavage sites of nsp15, an RNA uridylate-specific endoribonuclease (NendoU). Our model does not rule out other RNAs or proteins involved in these two models. However, current studies involving NendoU remain contradictory in terms of their findings regarding fundamental questions [5]: (1) NendoU is conserved among coronaviruses, arteriviruses and toroviruses, is it present in nonvertebrate-infecting representatives of the nidoviruses order? (2) is nsp15 indispensable in viral replication and living? (3) is nsp15 responsible for protein interference with the innate immune response? and (4) does nsp15 degrade viral RNA to hide it from host defences? Nsp15 cleavage sites in relevant genes provide new clues to answer these questions.

The Kozak consensus sequence GCCACCAUGG (**Figure 1B**) is not necessary for protein translation in eukaryotes. The hairpins immediately upstream of ORF1a (first hairpins) play an important role in the functions of CoV RBSs. Hairpin with proper structures may enhance the translation of their downstream proteins. These findings can be used to design vaccine, drug and expression vectors. Based on our preliminary experiments, we concluded that the hairpin in pSARS resulted in the over-expression of EGFP, which caused cell death. These facts supported that the negative feedback is necessary to prevent the over-expression of viral genes.

Our findings enrich fundamental knowledge in the field of gene expression and its regulation, and provide new clues for future studies. The template switching ability and the high ratio between contiguous and junction-spanning reads indicated that RdRP (nsp12) has a very high enzyme activity. Relatively few junction-spanning reads indicated that nsp15 does not have a high enzyme activity. This suggested that

nsp15 may be an important and ideal target for the development of drugs against

CoVs. The most recent nsp15 structure in complex with GpU (PDB: 6X1B) shows

that the uridine in GpU binds to the active site of nsp15. Thus, uridine derivatives,

such as Trifluridine, Tipiracil, Tegafur, Carmofur, Furtulon, etc, are potential

inhibitors of this enzyme.

Materials and Methods

1,265 genome sequences of betacoronaviruses were downloaded from the NCBI

GenBank database in our previous study [8]. Nanopore RNA-seq data was

downloaded from the website (https://osf.io/8f6n9/files/) for the reanalysis of

leader-body junction regions. The results were confirmed using Illumina RNA-seq

data from the NCBI SRA database under the accession number SRP251618. Data

cleaning and quality control were performed using Fastq_clean [9]. Statistics and

plotting were conducted using the software R v2.15.3 with the package ggplot2 [10].

The 5' and 3' ends of gRNAs(+) and sgRNAs(+) were observed and double-checked

using the software Tablet v1.15.09.01 [11].

In the present study, three types of plasmids pEGFP-C1 (maintained in the lab of

Hao Zhou), pSARS and pCoV-ba, and three types (Hela, HEK293 and HEK293T) of

cell maintained in our lab were used for transfection. To construct pSARS, pEGFP-C1

was PCR amplified, using primers fVR and rRBS2 (**Supplementary 1**). Then the

linear PCR product was self-ligated into a plasmid by homologous recombination

technology using ClonExpress II One Step Cloning Kit (Vazyme Biotech, China).

Following the same procedure, primers fVR and rRBS3 (**Supplementary 1**) were

used to construct pCoV-ba. The cells were cultured in Dulbecco's Modified Eagle

Medium (DMEM) media supplemented with 10% fetal bovine serum. For each

experiment, 100,000 cells were seeded into one well of a 6-well plate for plasmid

transfection. After 12 hours (0 hour in **figure 2B**), transfection of 1 μg of plasmid into

one well was performed using 3 μL PolyJet (SignaGen Laboratories, USA), following

the manufacturer's instructions. The medium was changed at 12 hours after plasmid

transfection. MTT (200 µl, 5 mg/ml) , and 1.8 ml medium was added to each well and

cultured at 37 °C for 4 h. Then, the cells in each well were removed of medium,

washed with PBS, then mixed with 1000 µl DMSO to dissolve the formazan product.

Finally, formazan absorbance was measured by a microplate reader with a wavelength

of 492 nm (Thermo Labsystems Helsinki, Finland). LDH experiments were

performed using LDH cytotoxicity assay detection kit (Beyotime, China), following

the manufacturer's instructions.

Supplementary information

313   Additional file 1: Figure S1. Comparison of fluorescent brightness. Table S1.

314 Primers for plasmid construction.

Abbreviations

COVID-19: Coronavirus disease 19; CoVs: coronavirus; SARS-CoV: Severe

Acute Respiratory Syndrome Coronavirus; SARS-CoV-2: SARS Coronavirus 2;

SARS-like CoV: SARS-CoV like Coronavirus; SARS-2 like CoV: SARS-CoV-2 like

Coronavirus; gRNA: genomic RNA; sgRNA: subgenomic RNA; RBS: ribosome

binding site; ORF: open reading frame; CDS: coding sequence; TRS-L: transcription

regulatory sequence in the leader; TRS-B: transcription regulatory sequence in the

body; ICS: internal cleavage site; RdRP: RNA-dependent RNA polymerase; nsp:

non-structural protein; NendoU: nidoviral RNA uridylate-specific endoribonuclease;

CMV: Cytomegalovirus; MTT:

3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2-H-tetrazolium bromide; LDH: lactate

dehydrogenase.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent to publish

Not applicable.

Availability of data and materials

All data used in the present study was download from the public data sources.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Natural Science Foundation of Tianjin (18JCYBJC28000) to Fang Wang and Tianjin Key Research and Development Program of China (19YFZCSY00500) to Shan Gao. The funding bodies played no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Authors' contributions

Shan Gao conceived the project. Xin Li and Guangyou Duan supervised this study. Guangyou Duan and Xiufeng Jin conducted programming. Zhi Cheng, Hao Zhou and Fang Wang conduced the experiments. Shan Gao and Yibo Xuan downloaded, managed and processed the data. Shan Gao drafted the main manuscript text. Shan Gao, Chang Liu and Jishou Ruan revised the manuscript.

Acknowledgments

We are grateful for the help from the following faculty members of College of

Life Sciences at Nankai University: Xuetao Cao, Deling Kong, Quan Chen, Wenjun

Bu, Ting Ma, Tao Zhang, Dawei Huang, Mingqiang Qiao, Yanqiang Liu, Qiang

Zhao, Bingjun He and Zhen Ye. This manuscript was online as a preprint on Aug

24nd, 2020 at https://www.biorxiv.org/content/10.1101/2020.08.23.263327v1.

REFERENCES

1.   R.L. Graham and R.S. Baric, *Recombination, Reservoirs, and the Modular*
Spike: Mechanisms of Coronavirus Cross-Species Transmission. Journal of
Virology, 2010. **84**(7): p. 3134-3146.
2.   D. Kim, J.-Y. Lee, J.-S. Yang, J.W. Kim, V.N. Kim and H. Chang, *The*
Architecture of SARS-CoV-2 Transcriptome. Cell, 2020. **181**(4): p. 914-921.
3.   X. Xu, H. Ji, X. Jin, Z. Cheng, X. Yao, Y. Liu, Q. Zhao, T. Zhang, J. Ruan, W.
Bu, Z. Chen and S. Gao, *Using pan RNA-seq analysis to reveal the ubiquitous*
existence of 5' and 3' end small RNAs. Frontiers in Genetics, 2019. **10**: p. 1-11.
4.   S. Zuniga, I. Sola, S. Alonso and L. Enjuanes, *Sequence Motifs Involved in the*
Regulation of Discontinuous Coronavirus Subgenomic RNA Synthesis. Journal
of Virology, 2004. **78**(2): p. 980-994.
5.   Y. Kim, R. Jedrzejczak, N.I. Maltseva, M. Wilamowski, M. Endres, A.
Godzik, K. Michalska and A. Joachimiak, *Crystal structure of Nsp15*
endoribonuclease NendoU from SARS-CoV-2. Protein Science, 2020. **29**(7): p.
1596-1605.
6.   D.D. Nedialkova, R. Ulferts, E. van den Born, C. Lauber, A.E. Gorbalenya, J.
Ziebuhr and E.J. Snijder, *Biochemical Characterization of Arterivirus*
Nonstructural Protein 11 Reveals the Nidovirus-Wide Conservation of a
Replicative Endoribonuclease. Journal of Virology, 2009. **83**(11): p.
5671-5682.
7.   G. Duan, J. Shi, Y. Xuan, J. Chen, C. Liu, J. Ruan, S. Gao and X. Li, *5' UTR*
Barcode of the 2019 Novel Coronavirus Leads to Insights into Its Virulence.
Chinese Journal of Virology (In Chinese), 2020. **36**(3): p. 365-369.
8.   X. Li, G. Duan, W. Zhang, J. Shi, J. Chen, S. Chen, S. Gao and J. Ruan, *A*
Furin Cleavage Site Was Discovered in the S Protein of the 2019 Novel
Coronavirus. Chinese Journal of Bioinformatics (In Chinese), 2020. **18**(2): p.
103-108.
9.   M. Zhang, F. Zhan, H. Sun, X. Gong, Z. Fei and S. Gao. *Fastq_clean: An*
optimized pipeline to clean the Illumina sequencing data with quality control.

394   in *Bioinformatics and Biomedicine (BIBM), 2014 IEEE International*
395   *Conference on*. 2014. IEEE.
396 10. H. Wickham, *ggplot2: elegant graphics for data analysis*. 2009: Springer
397   Science & Business Media.
398 11. I. Milne, G. Stephen, M. Bayer, P.J. Cock, L. Pritchard, L. Cardle, P.D. Shaw
399   and D. Marshall, *Using Tablet for visual exploration of second-generation*
400   *sequencing data*. Briefings in bioinformatics, 2012: p. bbs012.
401

A.

pCoV-ba   ···GCT GGTTTAGTGAACCGTCCCA-TCACA ATG ···

```
          A T
         C   A
         C-G
         C-G
```

pSARS   ···GCT GGTTTAGTGAACCGTAGGT-AGGTAAG ATG ···

```
          G G
         G   T
         C-G  T
         C-G
         T-A
         G-C C
         C-G
         T-A
         T-A
```

pEGFP-C1   ···GCT GGTTTAGTGAACCGTCAG-CGCCACC ATG ···

```
          C G
         G   C
         A-T
         T-A
         C   C
         G-C
         C-G
        T C-G
         A-T   Kozak sequence
```

RBS

CMV-promoter   EGFP gene

ori

B.

MTT

LDH

pCoV-ba
pSARS
pEGFP-C1