Title

Whole-organism eQTL mapping at cellular resolution with single-cell sequencing

Authors

Eyal Ben-David[1,2*], James Boocock[1], Longhua Guo[1], Stefan Zdraljevic[1], Joshua S. Bloom[1*],

Leonid Kruglyak[1*]

(*) To whom correspondence should be addressed: eyal.bendavid@mail.huji.ac.il,

jbloom@mednet.ucla.edu, lkruglyak@mednet.ucla.edu

Affiliations

1. Department of Human Genetics, Department of Biological Chemistry, and Howard Hughes

Medical Institute, University of California, Los Angeles, CA 90095, USA.

2. Department of Biochemistry and Molecular Biology, Institute for Medical Research Israel-

Canada, The Hebrew University School of Medicine, Jerusalem, Israel

Abstract

Genetic regulation of gene expression underlies variation in disease risk and other complex traits. The effect of expression quantitative trait loci (eQTLs) varies across cell types; however, the complexity of mammalian tissues makes studying cell-type eQTLs highly challenging. We developed a novel approach in the model nematode *Caenorhabditis elegans* that uses single cell RNA sequencing to map eQTLs at cellular resolution in a single one-pot experiment. We mapped eQTLs across cell types in an extremely large population of genetically distinct *C. elegnas* individuals. We found cell-type-specific *trans*-eQTL hotspots that affect the expression of core pathways in the relevant cell types. Finally, we found single-cell-specific eQTL effects in the nervous system, including an eQTL with opposite effects in two individual neurons. Our results show that eQTL effects can be specific down to the level of single cells.

Introduction

Gene expression differences have a strong genetic basis, and genome-wide studies have identified thousands of regions affecting gene expression, termed expression quantitative trait loci (eQTLs) [1]. eQTLs have been proposed to underlie genetic associations with complex traits and diseases [2,3], and genome-wide eQTL mapping holds great promise for uncovering the molecular underpinnings of phenotypic variation.

Studies in purified blood cell populations [4–6] and computational analyses in human tissues [7,8] indicate that many eQTLs are cell-type specific. Studying eQTLs in relevant tissues and cell types is thus important for understanding their role in trait variation [9]. However, major challenges remain for cell-type-specific eQTL mapping. First, studying multiple tissues or cell types is laborious and expensive, and has mostly been limited to large consortia [10]. Additionally, diseases can be associated with specific subsets of cells that may not be readily separable from the rest of the tissue, limiting the insight that can be gleaned from studying bulk tissue samples [11].

The recently developed technology of single-cell RNA sequencing (scRNA-seq) has the potential to significantly mitigate both of the above issues. In scRNA-seq, gene expression is profiled at the level of individual cells, and different cell populations are identified based on their gene expression profiles. This approach allows simultaneous measurement of expression in multiple tissues and cell types in a "one-pot" experiment, reducing the number of samples that need to be processed and profiled separately. Furthermore, distinct cell types can be interrogated directly, facilitating eQTL mapping in disease-relevant but rare cell types.

The size and complexity of mammalian tissues has so far limited the ability to use scRNA-seq for eQTL mapping, and studies have focused on purified cell populations and cell lines [12,13].

To pilot a whole-organism approach, we therefore turned to the model nematode *Caenorhabditis*

elegans. One of the mainstays of modern genetic research, *C. elegans* has an invariant cell

lineage that leads to each individual having the same number and identity of cells [14]. Decades

of research have uncovered the functions of many of those cells, as well as expression markers

that uniquely identify them [15,16]. These features and resources make *C. elegans* exceptionally

well-suited for using scRNA-seq to map eQTLs across cell types in the natural physiological

context of a whole animal, down to cellular resolution.

Results

Cell-type-specific eQTL mapping in a single one-pot experiment

Genome-wide eQTL mapping involves acquiring genotypes and gene expression profiles for a

genetically diverse cohort. We recently developed a method, *C. elegans* extreme quantitative

trait locus (ceX-QTL) mapping, for genetic analysis of complex traits in extremely large

populations of segregants [17]. The method takes advantage of a mutation in the gene *fog-2* that

forces the normally hermaphroditic *C. elegans* to reproduce via obligate outcrossing, allowing

us to propagate a large crossing experiment for multiple generations.

Here we build on ceX-QTL by combining it with single-cell RNA sequencing (scRNA-seq)

to carry out eQTL mapping at cellular resolution in a single one-pot experiment (Fig. 1A). In this

approach, a large heterogeneous pool of cells from thousands of genetically distinct individuals

is profiled using scRNA-seq, cell types are inferred by clustering scRNA-seq profiles and

studying known cell-type markers, and genotype information is reconstructed using expressed

genetic variants, enabling eQTL mapping in multiple cell types simultaneously.

We propagated a cross between the laboratory strain N2 and a highly divergent isolate from Hawaii, CB4856, for four generations, generating a pool of 200,000 genetically distinct F4 segregants. We dissociated the segregant pool to single cells at the L2 larval stage and profiled the cells with scRNA-seq. We identified clusters in a Uniform Manifold Approximation and Projection (UMAP) of the dataset [18–20], and determined their cell-type identities using known markers [21,22]. Our final dataset comprises 55,508 cells classified into 19 different cell types (Fig. 1B) (Table S1). The observed number of cells of each type was strongly correlated with the known cell-type abundance in L2 larvae (Spearman's ρ = 0.87, p = 2.2 x 10^{-6} , Supplementary Fig. 1).

Most of the cells in our sample were expected to carry unique genotypes (Methods). This design is advantageous for eQTL mapping because it maximizes the sample size [23], but requires *de novo* genotype calling, since the genotype of each cell is unknown beforehand. Rather than assign deterministic genotype calls based on sparse scRNA-seq data, we derived genotype probabilities for each cell using a Hidden Markov Model (HMM) (Supplementary Fig. 2). We then performed eQTL mapping with these genotype probabilities in a negative binomial modeling framework (Methods).

eQTL mapping in multiple cell types

We mapped 1,718 *cis* eQTLs in 1,294 genes, and 451 *trans* eQTLs in 390 genes, at a false discovery rate (FDR) of 10% across the different cell types (Fig. 2A-B, Table S2). The number of eQTLs detected in each cell type was strongly correlated with the number of cells of that type (Spearman's ρ = 0.91, p < 2.2 x 10^{-16}). In cell types with >1000 cells, we mapped between 52 and 415 eQTLs (Table S1). For 1,071 of the 1,294 genes with a *cis* eQTL (83%), the eQTL was

detected in only one cell type. For 208 of the remaining 223 genes (93%), the direction of the

eQTL effect was the same in all cell types in which it was detected.

We studied to what degree our *cis* eQTL results were concordant with gene expression

differences between the parents. We generated a scRNA-seq dataset from 6,721 N2 and 3,104

CB4856 cells, and used a classifier trained on the segregant dataset to identify cell types in the

parental scRNA-seq dataset. We then carried out a differential expression analysis in each cell

type. We found 870 differentially expressed genes (at a >2 fold change and FDR of 10%), of

which 201 (23%) had a *cis* eQTL in the same tissue (OR = 18.8, *p* < 2.2 x 10^{-16}, Fisher's Exact

Test). 191 of these *cis* eQTL (95%) showed the same direction of effect as the parental

difference. Further, the effect sizes of the significant *cis* eQTLs were strongly correlated with the

sizes of the parental differences (Spearman's ρ = 0.66, *p* < 2.2 x 10^{-16}) (Supplementary Fig. 3-

4). These results provide independent support for our *cis*-eQTL mapping, and show that for a

sizable fraction of the genes, those *cis* eQTLs are a major cause of differential gene expression

between the strains.

Comparison between bulk and single-cell *cis*-eQTL mapping

To investigate the relationship between single-cell and bulk eQTL mapping, we compared our

single-cell eQTLs to those previously identified in a panel of 200 recombinant inbred lines (RILs)

generated from crossing N2 and CB4856 [24]. In the bulk study, a large population of whole worms

from each RIL was recovered at a late larval stage, L4, and profiled on expression microarrays.

We reanalyzed data for 11,535 genes expressed in both datasets and identified 981 *cis* eQTLs

in the bulk dataset (at an FDR cutoff of 10%). Despite major differences in experimental design,

including the developmental stage of the worms, the overlap with the single-cell *cis* eQTLs was

highly significant, with 335 *cis* eQTLs shared between the studies (Odds Ratio = 7.2, $p < 2.2 \times$

10^{-16}, Fisher's exact test) (Fig. 2C). These shared loci represented 34% of the bulk *cis* eQTLs

and 32% of the single-cell *cis* eQTLs. Furthermore, the bulk and single-cell eQTL effect sizes

were highly correlated (Spearman's ρ = 0.64, $p < 2.2 \times 10^{-16}$) (Fig. 2D). Lastly, single-cell eQTLs

detected in multiple cell types were more likely to also be seen in the bulk study: 50% of the

genes with *cis* eQTLs detected in multiple cell types were also identified in bulk, compared to

28% of the eQTLs detected in only one cell type (OR = 2.58, p = 2.1 $\times 10^{-8}$) (Fig. 2C). This

observation suggests that the single-cell eQTL mapping approach improves the power to detect

cell-type specific effects.

Shared and cell-type specific *trans*-eQTL hotspots

We observed that 90 of the 451 *trans* eQTLs clustered at 5 hotspots, each containing 12-31

eQTLs (Supplementary Fig. 5, Table S3). A hotspot on Chr I was identified independently in

both neurons and seam cells; the top associated variant (Chr. I:10890182) was the same for

both cell types (Supplementary Fig. 5A-B). The other hotspots were identified in the body wall

muscle (on Chr. I) (Supplementary Fig. 5C), the intestine (on Chr. V) (Supplementary Fig 5D),

and neurons (two distinct hotspots on Chr. III) (Supplementary Fig. 5B).

To test whether the target genes of these 5 hotspots are involved in coherent biological

processes, we relaxed the FDR threshold to 20%, which increased the number of genes linked

to each hotspot to 21-42, and performed Gene Ontology (GO) enrichment analysis (Table S4).

For three of the hotspots, we found significant enrichments that were consistent with the cell-

type specificity of the hotspot. The targets of the hotspot detected in intestinal cells were weakly

enriched for genes involved in the innate immune response (FDR-corrected p = 0.042), a major

role of that tissue [25]. The targets of the hotspot detected in the body wall muscle were enriched

for genes associated with the term *myofilament* (FDR corrected p = 6.4 x 10^{-8}), *actin cytoskeleton*

(FDR-corrected p = 4.2 x 10^{-6}), and related terms. The enrichment was driven by the genes *mup-*

2, tni-1, tnt-2, mlc-2, mlc-3, lev-11 and *act-4*. *mlc-2* and *mlc-3* encode a myosin light chain, and

act-4 encodes an actin protein. *lev-11* encodes a tropomyosin, and *mup-2*, *tni-1*, and *tnt-2*

encode 3 of the 4 proteins in *C. elegans* that are expressed in the body-wall muscle and form

troponin complexes, highly conserved regulators of muscle contraction [26] (Supplementary Fig.

6).

The targets of the neuronal hotspot on the right arm of Chr III were enriched for genes

involved in *vesicle localization* (FDR-corrected p = 7.5 x 10^{-3}), as well as for BMP receptor

binding genes (FDR-corrected p = 2.8 x 10^{-3}). The latter enrichment was driven by *dbl-1* and *tig-*

2, orthologs of human bone morphogenetic protein (BMP) genes BMP5 and BMP8 and ligands

of the transforming growth factor beta (TGF-β) pathway [27]. Notably, *dbl-1* was discovered as a

gene that regulates body size in *C. elegans* [28], the hotspot peak marker is located <300 kb from

the peak of a QTL we previously identified for body size [29], and the corresponding confidence

intervals overlap (Table S3), suggesting that differential regulation of the TGF-β pathway is

involved in variation in body size between N2 and CB4856.

Cell-specific eQTL effects in the *C. elegans* nervous system

C. elegans is a premier model for studying neurobiology at the cellular level, which is aided by

its invariant cell lineage and the diverse functions associated with specific individual neurons.

Importantly, many of the neurons are highly variable in their gene expression, and express

specific gene markers [16]. To identify specific subtypes of neuronal cells, we separately clustered

the 12,467 cells identified as neurons and compared the clusters to previous *C. elegans* scRNA-

seq datasets, including the recently published *C. elegans* Neuronal Gene Expression Map &

Network (CeNGEN) [21,22,30,31] (Table S5). The neurons fell into 81 distinct clusters, ranging from

17 to 872 cells. We mapped these clusters onto 100 (83%) of the 120 neuronal clusters identified

in CeNGEN (Supplementary Fig. 7). We also identified CEM neurons, which are male specific

and absent from CeNGEN, based on the expression of the marker *cwp-1* [32].

We mapped *cis* eQTLs in each of the single neuronal subtypes (sn-eQTLs) and identified

a total of 163 sn-eQTLs in 132 genes at an FDR of 10% (Fig. 3A, Table S6). Of these, 117 (88%)

were identified in only a single neuronal subtype. Functional annotation of sn-eQTLs identified

25 genes involved in signaling (FDR-corrected p = 0.047), including 12 genes involved in G-

protein Coupled Receptor (GPCR) signaling (FDR-corrected p = 0.047) and 8 genes involved in

neuropeptide signaling (FDR-corrected p = 9.9 x 10^{-3}).

We compared the sn-eQTLs to those identified when all neurons were analyzed jointly

("pan-neuronal mapping"), and found that a sizable fraction of the sn-eQTLs did not have

evidence for a pan-neuronal signal: 92 were not identified pan-neuronally at an FDR of 10%,

and 69 were not identified even at a highly permissive FDR of 50%, suggesting that they exert

their effects only in specific neuronal subtypes (Fig. 3A). Regardless of statistical significance,

pan-neuronal eQTLs should have consistent effect directions across neuronal subtypes, while

subtype-specific eQTLs should not. We therefore compared the direction of effect of each sn-

eQTL in the subtype in which it was detected with its direction of effect in the set of all neurons

excluding that subtype. Among the 69 sn-eQTLs with no signal in the pan-neuronal mapping

even at the permissive FDR, the direction of the effect was concordant for 33 and discordant for

36, not significantly different from chance (p > 0.5; binomial test), as would be expected if these

effects are truly subtype-specific (Fig. 3B). In contrast, among the 94 that had a pan-neuronal

signal at an FDR of 50%, the direction of the effect was concordant for 88 and discordant for

only 6 ($p < 0.000001$; binomial test), consistent with differences in detection arising from limited

statistical power.

In a striking case, we observed an sn-eQTL in the neuropeptide gene *nlp-21* that showed

significant and opposing effects in two neurons (Fig. 3C-D). In the RIC neuron, higher *nlp-21*

expression was associated with the CB4856 allele (β = 4.4, FDR-corrected *p* = 0.03), while in

the RIM neuron, higher *nlp-21* expression was associated with the N2 allele (β = -5.4, FDR-

corrected *p* = 9.8×10^{-7}). In the pan-neuronal mapping, no significant effect is observed for this

gene. We identified the RIC and RIM neurons in the parental dataset, and although the small

number of cells in each group (35 and 27, respectively, with only 9 and 5 of them from CB4856)

was insufficient for statistical testing, the directions of the differences agreed with the eQTL

effects (Fig. 3E). These results provide direct evidence that eQTLs can be specific down to the

cellular level.

Discussion

We used scRNA-seq to map eQTLs in *C. elegans* across cell types in a single one-pot

experiment. Earlier scRNA-seq eQTL mapping studies were limited in sample size to at most

~100 individuals, but nevertheless highlighted the potential of this approach to identify cell-type

[13] and developmental [12] eQTLs, as well as loci affecting expression variance [33]. Our novel

approach allowed us to map eQTLs in tens of thousands of genotypically distinct individuals and

enabled detection of both *cis* and *trans* eQTLs, as well as resolution of their effects down to the

level of specific cells.

One of the major factors affecting gene expression studies is variation resulting from uncontrolled environmental differences between individuals that are grown or processed separately. By using scRNA-seq, we were able to process all individuals jointly. After the initial parental cross, all subsequent steps carried out over the course of five *C. elegans* life-cycles (three weeks) were performed in bulk, limiting any confounding environmental factors. To minimize the influence of genotype on development, we synchronized the worms at the first larval stage, L1, and collected samples at the L2 stage, limiting the time for differences to accumulate post-synchronization. Even careful synchronization is not expected to completely remove the effects of genetic variation on developmental timing, and such variation can be combined with gene expression time course data collected during development to increase the power of eQTL mapping and to study the developmental dynamics of eQTLs [34]. This raises the possibility that future scRNA-seq studies of *C. elegans* across developmental stages would open the door to a similar analysis of our single-cell eQTL dataset.

Previous work suggested the existence of cell-type specific eQTL hotspots in *C. elegans* based on the expression patterns of hotspot targets [34]. We discovered three hotspots that are cell-type specific, with targets that are involved in core functions performed by these cell types. Recently, eQTL hotspots have been identified in human blood cells [35,36], as well as in cell lines [37]. These results suggest hotspot and *trans* eQTL discovery is facilitated by expression studies that can distinguish cell types and point to a larger role of hotspots in the genetics of gene expression in animals.

A comparison of the single-cell *cis* eQTLs to those mapped in a previous whole-worm eQTL study from our laboratory showed a highly significant overlap despite major differences in experimental design. These results join accumulating evidence that a sizable fraction of *cis*

eQTLs have robust, consistent effects [10,38], and show that many of the effects are conserved

across worm development. The strong overlap of *cis* eQTLs mapped by scRNA-seq and by

whole-worm analysis also suggests that the effect of many *cis* eQTLs is conserved across cell

types. By generating an extremely large number of unique segregants, our method enables

scaling up the number of studied individuals simply by sequencing a larger number of cells.

Thus, the increasing throughput of single-cell technologies and sequencing platforms will enable

future work to study cell-type specificity of *cis* and *trans* eQTLs in greater detail.

Lastly, we discovered *cis* eQTLs that act in single subtypes of *C. elegans* neurons,

including many that were not found when all neurons were analyzed jointly. Importantly, we also

discovered an eQTL that influences expression of the gene *nlp-21* in opposing directions in two

different neurons. Such antagonism could result from context-dependent effects of the same

variant, or from different variants acting in each cell. Our study joins a growing body of work that

identified eQTLs that are specific to cell types [5–8,13,39,40], as well as developmental [12,34] and

environmental [41–43] contexts. Distinct genetic effects are found across cell types and conditions,

and large-scale study of such effects is therefore crucial for gaining a comprehensive

understanding of regulatory variation.

Figures

Figure 1. Whole-organism eQTL mapping with single-cell RNA-sequencing (scRNA-seq). (A) A large population of segregants is dissociated to single cells. Each cell in the suspension has an unknown genotype and cell-type identity. The suspension is profiled using scRNA-seq. Cell-type identity is inferred by clustering cells and comparing the expression of known marker genes. Genotypes are reconstructed from expressed single-nucleotide variants (SNVs). (B) The UMAP projection of 55,508 scRNA-seq expression profiles from approximately 200,000 *C. elegans* F4 segregants collected at the L2 larval stage is shown. Cells are colored based on the inferred cell type.

Figure 2. eQTL mapping in cell types. (A) A genome-wide map of eQTLs across all cell types is shown. The position of the eQTLs is shown on the x-axis, while the y-axis shows the position of the associated transcripts. Points along the diagonal are *cis* eQTLs (those mapping to nearby genes). **(B)** The number of *cis* and *trans* eQTLs mapped in each cell type. **(C)** The overlap between a previous study that mapped eQTLs in whole worms in a panel of recombinant inbred lines (RIL) and our dataset.

274 (Left) The proportion of genes with a *cis* eQTL in at least one dataset, out of all genes tested. (Middle)

275 Of the 1,688 significant *cis*-eQTL genes, 355 had a *cis* eQTL in both datasets, representing a highly

276 significant enrichment. (Right) Hits mapped in more than one cell type were more likely to also be

277 found in the whole-worm ("bulk") dataset. (D) Quantitative comparison between normalized effect

278 sizes in our dataset and in the whole-worm dataset.

Figure 3. Neuron-specific eQTL mapping. (A) *cis*-eQTLs mapped in single neuronal subtypes (sn-eQTLs) are shown. The top three rows indicate whether the eQTL was mapped pan-neuronally at a 10% FDR threshold (row 1), at a 50% FDR threshold (row 2), and whether the sign of the effect estimate ("effect direction") was the same in the pan-neuronal and single cell mapping (row 3). (B) Comparing the effect direction between the sn-eQTL mapping and mapping in a set of neurons excluding the sn-eQTL neuron shows evidence for subtype-specific effects. The number of genes showing the same ("purple") or opposite ("turquoise") effect directions is shown for genes with pan-neuronal FDR > 50% (top) and < 50% (bottom). (C-D) An eQTL with antagonistic effects in two

neurons. Higher expression of the gene *nlp-21* in the RIM neuron is associated with the N2 allele (C),

while higher expression in the RIC neuron is associated with the CB4856 allele (D). In C and D, a

linear fit is shown for illustration. All p values are FDR-corrected. Read counts were normalized to the

number of UMIs in each cell and log-transformed. (E) Expression of *nlp-21* in the parental dataset.

The direction of effect is concordant between the left panel and (C) (RIM neuron) and between the

middle panel and (D) (RIC neuron). Horizontal lines are averages.

Methods

C. elegans culturing

C. elegans strains were cultured at 20°C using standard conditions with the exception that the agar in the nematode growth media (NGM) was replaced with a 4:6 mixture of agarose and agar (NGM+agarose), to prevent burrowing of the CB4856 strain. Parental strains used were QX2314 (N2 *fog-2*(q71) V; *hsp-90p*::GFP II) and PTM299 (CB4856 *fog-2*(kah89)). Large segregant panels were generated as before [17]. Briefly, 500 L4 males from PTM299 and 500 L4 hermaphrodites from QX2314 were seeded on a plate for 30 hours, and gravid worms and eggs were collected and bleached. Eggs were synchronized to L1 larvae for 24h, and seeded on 10cm NGM+agarose plates. In each generation, gravid worms were bleached, their progeny synchronized for 24h and seeded. The entire process was repeated up to F4, with 3-4 days per generation.

Cell extraction and sequencing

192,000 F4 were seeded on four 10cm NGM+agarose plates seeded with OP50. L2 were recovered after 24 hours, and staging was validated under a stereomicroscope. L2 cell dissociation was carried out as previously described [44], implementing modifications from a later study [45], as well as our own. Worms were recovered off the plates and washed three times in M9. Lysis was then done with an SDS-DTT solution (200 mM DTT, 0.25% SDS, 20 mM HEPES, pH 8.0, 3% sucrose) in a hula mixer set on low speed to prevent worms from settling. The lysate was observed under the stereoscope every two minutes, and lysis was stopped when a blunted head shape appeared in the majority of worms [45], after ~4 minutes. Worms were then washed quickly three times in 1ml of M9, and two additional times in 1ml of egg buffer (118 mM NaCl,

48 mM KCl, 2 mM CaCl2, 2 mM MgCl2, 25 mM HEPES, pH 7.3, osmolarity adjusted to 340

mOsm with sucrose). Worms were then resuspended in 0.5ml of 20mg/ml Pronase E that was

freshly prepared in L15 media supplanted with 2% fetal bovine serum (L15-FBS) and adjusted

to 340 mOsm with sucrose. Worm dissociation was done by continuous pipetting on the side of

the tube, and monitored every 2-3 minutes on a microscope equipped with a 40x phase contrast

objective lens. Dissociation was stopped when few intact worms remained and a high density of

cells was visible. 0.5ml of L15-FBS was added to stop the reaction, and the lysate was spun for

6 minutes at 500g at 4°C. The cell pellet was resuspended in PBS (PBS was adjusted to 340

mOsm with sucrose). Cell suspension was spun for 1 minute in 100g at 4°C to remove remaining

undigested worms, counted and diluted to 1M cells/ml in osmolarity adjusted PBS, and loaded

directly onto 5 lanes of 3' Chromium single-cell RNA-sequencing flow cells (10x Genomics),

targeting 10,000 cells on each lane. Library prep was carried out according to manufacturer's

protocol. Prepared libraries were sequenced together on an S4 lane of Novaseq 6000. A paired-

end 2x150 run was done, to maximize the recovery of single-nucleotide variants. In all

downstream processing, each of the five 3' Chromium lanes processed concurrently was treated

as a separate "batch", and lane identity corresponds to the "batch" identity for the rest of the

methods.

Single-cell RNA-sequencing data processing

Raw sequencing reads were analyzed using *CellRanger* (Ver 3.0.2). We used a gene transfer

format (GTF) file that was corrected for misannotation of 3' untranslated regions (3'UTR) that

was generated in a previous study [22]. *C. elegans* cell-types differ widely in the number of UMIs

that are recovered using scRNA-seq. Therefore, a simple UMI cutoff, as is commonly used, may

be biased for cell-types with more UMIs. We therefore implemented an iterative pipeline to

recover clusters of *bona fide* cells and remove cell doublets as well as degraded cells. We took

20,000 cells with the most UMIs in each cluster (twice the targeted number of cells, 100,000

overall), and processed them in *Monocle* (Ver 3) [46]. Default parameters were used, with the

exception that 100 dimensions were used for reduction, and batch was added as a covariate.

Leiden clustering identified a total of 154 clusters, and we used the *top_markers* function in

Monocle to identify the genes upregulated in each. We then removed clusters whose top genes

included any ribosomal genes or the mitochondrial genes *ndfl-4*, *nduo-6*, *atp-6*, *ctc-2*, *ctc-3*, *ctc-*

1, which we noticed were usually found together as the most upregulated genes in clusters that

did not specifically express any known markers for *C. elegans* cell-types. This removed a total

of 30,980 cells (31%). For the remaining 69,020 cells, raw counts were processed using the R

package *SoupX* to reduce ambient RNA contamination [47]. We then normalized, reduced

dimensions and clustered the background corrected cell profiles in *Monocle* using the same

parameters as above.

To annotate cell-types, we used the markers described in a previous study (Table S12 in

Ref [22]) that reanalyzed a previous L2 single-cell dataset [21]. Our cell-type annotation corresponds

to the "UMAP" column in that table, with the following exceptions: (a) we separated hypodermis

from seam cells, somatic gonad from sex muscle cells, and glia from excretory cells, since those

groups were not clustering together in our data. (b) cells identified as "Miscellaneous" in that

table were annotated as individual cell-type identifications in our data, with the exception of the

sphincter and anal muscles which were not differentiated from each other in our data. Finally,

we re-evaluated our cell type identifications, and filtered cell doublets as well as dead cell or

debris that may still contaminate *bona fide* cell-type clusters. We trained a classifier using our

manually curated cell-type classifications with a L2-penalized multinomial logistic regression

framework, as implemented in the *Scikit-learn* Python package (v0.22) [48]. We read the raw gene

expression matrices into python using scanpy (v1.4.2). We removed 2,582 genes that were

expressed in less than 10 cells. The gene expression levels of each cell were corrected so that

the total gene expression counts added up to 10,000. Per gene, these corrected counts were

normalized using a log(1+x) transformation. To speed up the computation of the multinomial

logistic regression, we only used the 2,037 genes with a mean expression between 0.0125 and

3, and a minimum dispersion of 0.5. We scaled the gene expression matrix so that the

expression level of each gene across cells had a mean of 0 and a variance of 1, after scaling

expression values over 20 were set to 20. We fit a multinomial logistic regression model using

the scaled gene expression values for the 2,037 highly variable genes from the complete set of

69,020 cells to obtain an estimate for the inverse regularization strength (C). Using the estimated

C of 7.74×10^{-04}, we performed 5 fold cross validation to estimate the probability that each cell

belongs to one of the manually curated cell-type classifications. Any cell with a probability higher

than 0.2 of belonging to 2 or more cell types (9,198) was classified as a doublet. Any cell which

did not belong to a cell-type with probability >= 0.4 (5,547) was classified as low quality. In total,

we removed 11,398 cells that were classified as a doublet or low quality. We removed an

additional 2,114 cells classified as Neurons as described in the section "Neuronal cell-type

classification". For the remaining final list of 55,508 cells, we used the output of the classifier as

the final cell-type classification. The final classification is shown in Figure 1. For display purposes

the plot in Figure 1 was generated by rerunning umap on the finalized dataset with *euclidean*

distance metric and *umap.min_dist*=0.5, resulting in a more compressed visualization of the

dataset.

Estimating the number of unique genotypes

We note that calculating the number of expected unique genotypes is akin to the well-known "Birthday problem" in statistics. Given C cells sampled from I individuals, the expected number of cells with a unique genotype is $C(1-1/I)^{C-1}$. Assuming 50%-90% of worms were successfully dissociated (a conservative range), we expect 31,134-40,257 unique genotypes.

Single-nucleotide variant counting

We used a list of single-nucleotide variants (SNVs) we previously curated for CB4856 compared to the N2 reference [17]. We derived genotype informative UMI counts for N2 and CB4856 variants using *Vartrix version 1.0* (https://github.com/10XGenomics/vartrix) directly on the output of *CellRanger*. To reduce SNV counts that result from SNVs in the ambient RNA background, we only kept SNVs that resided in genes with positive counts in the background corrected matrix.

Genotype inference using a hidden Markov model

We set up a hidden Markov model (HMM) to infer the genotypes of the recombinant progeny [49,50]. The HMM is used to calculate the probability of underlying genotypes for each individual and requires three components: (1) prior probabilities for each of the possible genotypes, (2) emission probabilities for observing variant informative reads given each of the possible genotypes, (3) and transition probabilities - the probabilities of recombination occurring between adjacent genotype informative sites.

For the autosomal chromosomes we defined prior genotype probabilities as 0.25 for homozygote N2, 0.5 for heterozygote CB4856/N2, and 0.25 for homozygote CB4856. For the

sex chromosome we defined prior genotype probabilities as 0.44 for homozygotes N2, 0.44 for

the heterozygote CB4856/N2, and 0.11 for homozygote CB4856. These values were chosen

because to generate the segregant population, N2 hermaphrodites were crossed to CB4856

males, and thus contributed twice as many X chromosomes to the progeny as CB4856.

Emission probabilities were calculated as previously described for low coverage

sequencing data [51,52] under the assumption that the observed counts of reads for both possible

variants (Y) at a genotype informative site (g) arise from a random binomial sampling of the

alleles present at that site and that sequencing errors (e) occur independently between reads at

a rate of 0.002:

$p(Y|g = NN) = \binom{D}{r}(1 - e)^r(1 - (1 - e))^{D-r}$

$p(Y|g = NC) = \binom{D}{r}\frac{1}{2}^D$

$p(Y|g = CC) = \binom{D}{r}(e)^r(1 - e)^{D-r}$

Where (D) is the total read depth at a genotype informative site for a given individual, (r) is the

total read depth for the N2 variant at that site, and N represents the N2 variant and C represents

the CB4856 variant.

Transition probabilities were derived from an existing N2 x CB4856 genetic map [53]. We

linearly interpolated genetic map distances from the existing map to all genotype informative

sites in our cross progeny. We scaled these genetic map distances, multiplying them by a factor

of 0.4, to account for the fact that the previous genetic map was built using ten generations of

intercrossing whereas progeny from our cross are derived from four generations of intercrossing

[54].

For QTL mapping we used an additive coding, summing the probability that the genotype

was homozygote CB4856 with one half the probability that the genotype was heterozygote

N2/CB4856.

eQTL mapping

Genotype probabilities were standardized, and markers in very high LD (r>.9999) were pruned.

This LD pruning is approximately equivalent to using markers spaced 5 centimorgans (cm) apart.

For each transcript, we counted the number of cells for which at least one UMI count was

detected in each cell type. Transcripts with non-zero counts in at least 20 cells in a cell type were

considered expressed in that cell type and used for downstream analyses.

As has been previously described for droplet scRNA-seq, counts of UMIs can be

adequately parameterized by a gamma-poisson distribution, which is also known as the negative

binomial distribution [55]. Thus we used a negative binomial regression framework for eQTL

mapping here. We also note that simpler approaches using log(counts+1) with ordinary least

squares behave pathologically, especially in regard to behavior with multiple partially correlated

covariates, and simulations (not shown) showed such models lead to inflated false positive rates.

For each expressed transcript in each cell type we first fit the negative binomial

generalized linear model:

$E[Y] = \mu$ (1)

$Var(Y) = \mu + \dfrac{1}{\theta}\mu^2$ (2)

$\mu = exp(\beta_i + X_t\beta_t + \mathbf{X_b}\beta_b + X_c\beta_c)$ (3)

Which has the following log-likelihood:

$$\ell(\beta, \theta) = -\sum_{n=1}^{N}[(y_n + \theta)log(\mu_n + \theta) - y_nlog(\mu_n) + log(| \ \Gamma(y_n + 1) \ |) -$$

$$\theta log(\theta) + log(| \ \Gamma(\theta) \ |) - log(| \ \Gamma(\theta + y_n) \ |)]$$ (4)

And where Y is a vector of UMI counts per cell, X_t is a vector of the log(total UMIs per cell) and

controls for compositional effects, $\mathbf{X_b}$ is an indicator matrix assigning cells to batches, and X_c is

the vector of standardized genotype probabilities across cells for the closest genotypic marker

to each transcript from the pruned marker set. In addition, β is a vector of estimated coefficients

from the model, μ_n is the expected value of Y for a given cell n, N is the total number of cells in

the given cell type, and θ is a negative binomial overdispersion parameter. Model parameters

were estimated using iteratively-reweighted least squares as implemented in the *negbin.reg*

function in the *Rfast2* R package. If the model did not converge, model parameters were

estimated with the *gam* function in the *mgcv* package [56], which opts for certainty of convergence

over speed. We note that due to the computational burden of fitting so many GLMs in the context

of sc-eQTL mapping, we chose to estimate θ once for each transcript in each cell type for this

model, and use that estimate of θ in the additional models for that transcript within the cell type,

as described below. This approach is conservative, as the effects of unmodeled factors (for

example *trans* eQTLs) will be absorbed into the estimate of overdispersion, resulting in larger

estimated overdispersion $(\frac{1}{\theta})$ and lower model likelihoods. Computational approaches that re-

estimate θ for each model, that jointly model all additive genetic effects, or that regularize θ

across models and transcripts [57], may further increase statistical power to identify linkages.

To evaluate the statistical significance of *cis* eQTLs, a likelihood ratio statistic

$-2(\ell_{nc} - \ell_{fc})$, was calculated comparing the log-likelihood of this model described above (ℓ_{fc})

to the log-likelihood of the model where β is re-estimated while leaving out the covariate X_c for

the *cis* eQTL marker (ℓ_{nc}). A p-value was derived under the assumption that this statistic is χ^2

distributed with one degree of freedom. This p-value was used for the evaluation of significance

of *cis* eQTLs for the neuronal subtypes. Within each neuronal subtype, FDR adjusted p-values

were calculated using the method of Benjamini and Hochberg [58]. For the other cell types (with

typically much larger cell numbers) and for the genome-wide scans for eQTLs, a permutation

procedure was used to calculate FDR adjusted p-values, and is described further below.

For each expressed transcript in each cell type we also scanned the entire genome for

eQTLs, enabling detection of *trans* eQTLs. A similar procedure was used as for *cis* eQTL except

that equation (3) was replaced with:

$\mu = exp(\beta_i + X_t\beta_t + \mathbf{X_b}\beta_b + X_g\beta_g)$ (5)

where X_g is a vector of the scaled genotype probabilities at the gth genotypic marker, and the

model is fit separately, one at a time, for each marker across the genome for each transcript. A

likelihood ratio statistic for each transcript, within each cell type, for each genotypic marker is

calculated by comparing this model to the model where β is re-estimated while leaving out the

covariate X_g. The likelihood ratio statistic was transformed into a LOD score, by dividing it by

$2log_e(10)$. We also used functions in the *fastglm* R package for this scan, again re-using

estimates of θ obtained as described above for each transcript for each cell type. For each

transcript and each chromosome, QTL peak markers were identified as the marker with the

highest LOD score. The 1.5 LOD-drop procedure was used to define approximate 95%

confidence intervals for QTL peaks [59].

FDR-adjusted p-values were calculated for QTL peaks. They were calculated as the ratio

of the number of transcripts expected by chance to show a maximum LOD score greater than a

particular LOD threshold vs. the number of transcripts observed in the real data with a maximum

LOD score greater than that threshold, for a series of LOD thresholds ranging from 0.1 to 0.1+the

maximum observed LOD for all transcripts within a cell type, with equal-sized steps of 0.01. Per

chromosome, the number of transcripts expected by chance at a given threshold was calculated

by permuting the assignments of segregant identity within each batch relative to segregant

genotypes, calculating LOD scores for all transcripts across the chromosome as described

above, and recording the maximum LOD score for each transcript. In each permutation instance,

the permutation ordering was the same across all transcripts. We repeated this permutation

procedure 10 times. Then, for each of the LOD thresholds, we calculated the average number

of transcripts with maximum LOD greater than the given threshold across the 10 permutations.

We used the *approxfun* function in R to interpolate the mapping between LOD thresholds and

FDR and estimate an FDR-adjusted p-value for each QTL peak [60].

The same procedure was performed for *cis* eQTL analysis, with the difference being that

the expected and observed number of transcripts at a given LOD threshold were calculated only

at the marker closest to the transcript. We note that, as expected, Benjamini and Hochberg

adjusted p-values, and FDR adjusted p-values from this permutation procedure for *cis* eQTLs

were nearly identical.

scRNA-seq in the parental strains

The parental QX2314 and PTM299 strains were grown separately for 4 generations on 10cm

plates, with recurrent cycles of bleaching and synchronization as was done for the segregant

population. For single cell preparation, synchronized L1 from both strains were seeded together

in equal numbers on 10cm plates, and they were processed together from that point onwards,

to limit any environmental effects. We believe that differences in efficiency of the cell preparation

procedure between N2 and CB4856 could explain the imbalanced representation in the final

dataset (6,721 N2 and 3,104 CB4856 cells). We took advantage of the different parental

genotypes when processing the cells, and called cells as those with at least 50 SNV counts

supporting one genotype, and less than 50 supporting the other. Cell-type identification was

automated by using the logistic regression model trained on the segregants which is discussed

above. Differential expression analysis was carried out using the *DEsingle* R package in each

cell type, as well as globally in all cells combined [61].

To compare differential expression results with our *cis* eQTL results, we first normalized

the effect size of each *cis* eQTL by its standard error. Those were used directly in comparisons

done within each cell type. To compare with global differential expression, those standardized

effects were combined across all cell types in which an eQTL was identified using Stouffer's

weighted-Z method [62].

Processing whole-worm eQTL data

Microarray genotype and gene expression data for our published expression QTL data were

acquired from the gene expression omnibus (GEO) [24]. Probe sequences were realigned to the

WBcel235 transcriptome using BWA, and uniquely mapping probes were used. Expression

probes that were present in less than 2/3 of the sample were removed. The genotype and

expression matrices were standardized. To map eQTLs, we calculated the Pearson correlation

between each probe and every genotype. Correlation coefficients were transformed to LOD

scores using $-n \cdot \frac{ln(1 - R^2)}{2ln(10)}$. To assess significance and account for multiple testing, we

permuted the sample identities 100 times and calculated the average number of transcripts with

an identified eQTL at different LOD scores. We compared these results to the unpermuted LOD

scores to estimate the false-discovery rate (FDR), and selected a cutoff corresponding to a rate

of 10% (LOD = 4.2), equivalent to the single-cell mapping. *cis* eQTLs were derived by calculating

the Pearson correlation between transcript expression and the normalized genotypes in the

variant nearest to a given transcript, transforming to LOD score and comparing against the global

threshold.

Hotspot analysis

To discover hotspots, we split the genome into 130 bins of 5 centimorgans each. We then

counted the number of eQTLs in each bin identified in each cell type (applying a 10% FDR

significance threshold), after removing all *cis* linkages. *Cis* linkages were defined here as those

where the transcribed gene falls within the 95% eQTL confidence-interval range extended by

1MB on both sides. A bin was considered to have an excess of linkages if the number of linkages

exceeded the number expected by chance from a Poisson distribution, given the average

number of linkages per bin for that cell type and a Bonferroni correction for the total number of

bins (p<3.8e-4) [63]. The *findpeaks* function in the *pracma* R package was used to identify peak

hotspot bins and prevent identifying sets of adjacent bins as hotspots.

For gene ontology (GO) analysis, we identified hotspot targets using the same procedure

above, but relaxed the significance threshold to 20% FDR. We then used the R package *topGO*

to identify enriched terms, with the genes expressed in the cell type used as background.

Neuronal cell-type classification

Neuronal classification was carried out using a combination of available *C. elegans* scRNA-seq

datasets, including a published L2 dataset [21,22], and the *C. elegans* Neuronal Gene Expression

Map & Network (CeNGEN) project [31]. Neuronal cells were processed separately using *monocle3*

with default parameters, with the exception that 100 dimensions were specified for the

preprocess_cds step. The analysis was carried out in two passes. In the first pass, we processed

all cells identified by our classifier as neurons. Following Leiden clustering, we removed 2,114

cells that were in clusters whose top genes were mostly mitochondrial and ribosomal genes,

similar to the analysis described above for the global dataset. We then processed the pruned

dataset in *monocle3* as described above. To annotate the final neuronal clusters, we first used

the list of marker genes from two previous publications [22,30], to derive candidate clusters that

uniquely express marker genes. We next used the *top_markers* function in *monocle3* to identify

upregulated genes in each cluster compared to the rest. These were compared with the data

available in the online SCeNGEN *Shiny* application (https://cengen.shinyapps.io/SCeNGEA) for

the candidate cluster. The full list of genes used for classification is found in Table S5. In the

final dataset, clusters Unknown_1 - Unknown_4 are of unknown identity and do not correspond

to the clusters of the same name in CeNGEN, while the clusters Unknown_touch and

Unknown_glut_2 do correspond to cell clusters of the same names in CeNGEN.

Single neuronal subtype eQTL (sn-eQTL) analysis

sn-eQTL mapping is described above (section "eQTL mapping"). GO annotation of genes with sn-eQTL was done in *topGO*, with the genes expressed in neurons (determined using the criteria for inclusion in eQTL mapping) used as background. A heatmap was plotted using the *ComplexHeatmap* package [64]. To determine the consistency in effect direction between the sn-eQTL neuron and the rest of the neurons, we repeated the eQTL mapping, aggregating cells from all neuronal cell-types but omitting the neuron with the sn-eQTL. The RIC and RIM neurons in the parental datasets were identified using the same gene markers used in the segregant eQTL dataset, as described in Table S5.

Acknowledgments

The work was supported by the Howard Hughes Medical Institute and NIH grants R01 HG004321 (L.K.) and K99 HG010369 (E.B-D). Figure 1A was created with Biorender.com. We thank Michael Mashock, Marco De Simone, and other members of the Technology Center for Genomics & Bioinformatics (TCGB) at UCLA for preparing and sequencing 10x 3' chromium single-cell libraries.

Author contributions

Conceptualization: E.B-D, J.B., J.S.B, and L.K.; Methodology: E.B-D, J.B. and J.S.B.; Investigation: E.B-D, J.B., L.G., S.Z. and J.S.B.; Formal analysis: E.B-D, J.B., and J.S.B; Writing – original draft: E.B-D and L.K.; Supervision: J.S.B and L.K; Funding acquisition: E.B-D. and L.K. All authors reviewed the manuscript.

Data availability

Raw sequencing data are available under NCBI bioproject PRJNA658829.

Code availability

Code for the HMM and eQTL mapping can be found at

https://github.com/joshsbloom/single_cell_eQTL. Data and code to recreate the figures in the

manuscript can be found at https://github.com/eyalbenda/worm_sceQTL.

Competing interests

The authors declare no competing interests

Materials & Correspondence

Should be addressed to E.B-D, JS.B or L.K

Supplementary Figures

Supplementary Figure 1. Observed representation of cell-types in our dataset compared

to expected. The expected number of cells was calculated by manually curating the cellular

lineage information available at https://www.wormatlas.org/ for the L2 stage.

Supplementary Figure 2. Probabilistic genotyping using a hidden markov model (HMM).

A cell with the median (69) number of unique genotype-informative SNV UMI counts is shown

for illustration. The trace is a summation of the probability of a CB4856 homozygous genotype

and half the probability of CB4856 heterozygous genotype at each position. Each vertical line is

a count for an SNV, and colors correspond to the count depth. Vertical lines pointing upwards

denote counts supporting the CB4856 variant, while lines pointing downwards are counts

supporting the N2 variant.

Supplementary Figure 3. *cis* **eQTLs reflect gene expression differences in the parent strains.** Comparison between *cis* eQTLs mapped across all cell types and gene expression differences in a dataset of 6,721 N2 and 3,104 CB4856 cells. For *cis* eQTLs mapped in multiple cell-types, effect sizes were combined using Stouffer's weighted-Z method [62]. Differential gene expression between N2 and CB4856 was calculated using R package DEsingle .

Supplementary Figure 4. Correlation between cell-type *cis*-eQTL signal and differential gene expression between the parental strains. The data corresponds to Figure S3, split out by each cell type separately.

Supplementary Figure 5. cell-type-specific *trans*-eQTL Hotspots. A genome-wide map of eQTLs in seam cells (A), neurons (B), body-wall muscle cells (C) and intestinal cells (D) is shown. The position of the eQTLs is shown on the x-axis, while the y-axis shows the position of the associated transcripts. The dotted line marks the peak position of the hotspot, while targets of each hotspot are colored.

Supplementary Figure 6. Cell-type expression of genes that form troponin complexes.

Size of circles corresponds to the percentage of cells expressing each gene in each cell-type,

and the color corresponds to average expression. Of the four troponin genes strongly expressed

in the body wall muscle, three are affected by a *trans*-eQTL hotspot on Chr. I.

Supplementary Figure 7. UMAP projection of 12,468 neurons. Each cluster is labeled based

on the neuronal identity. Clusters represent either single neurons, or few neurons with shared

function.

References

1. Albert, F. W. & Kruglyak, L. The role of regulatory variation in complex traits and disease. *Nat. Rev.*

Genet. **16**, 197–212 (2015).

2. Gusev, A. *et al.* Partitioning heritability of regulatory and cell-type-specific variants across 11

common diseases. *Am. J. Hum. Genet.* **95**, 535–552 (2014).

3. Hormozdiari, F. *et al.* Leveraging molecular quantitative trait loci to understand the genetic

architecture of diseases and complex traits. *Nat. Genet.* **50**, 1041–1047 (2018).

4. Raj, T. *et al.* Polarization of the Effects of Autoimmune and Neurodegenerative Risk Alleles in

Leukocytes. *Science* **344**, 519–523 (2014).

5. Fairfax, B. P. *et al.* Genetics of gene expression in primary immune cells identifies cell type–specific

master regulators and roles of HLA alleles. *Nat. Genet.* **44**, 502–510 (2012).

6. Ishigaki, K. *et al.* Polygenic burdens on cell-specific pathways underlie the risk of rheumatoid

arthritis. *Nat. Genet.* **49**, 1120–1125 (2017).

7. Donovan, M. K. R., D'Antonio-Chronowska, A., D'Antonio, M. & Frazer, K. A. Cellular deconvolution

of GTEx tissues powers discovery of disease and cell-type associated regulatory variants. *Nat.*

Commun. **11**, 955 (2020).

8. Kim-Hellmuth, S. *et al.* Cell type–specific genetic regulation of gene expression across human

tissues. *Science* **369**, (2020).

9. Yao, D. W., O'Connor, L. J., Price, A. L. & Gusev, A. Quantifying genetic effects on disease

mediated by assayed gene expression levels. *Nat. Genet.* **52**, 626–633 (2020).

10. Consortium, T. Gte. The GTEx Consortium atlas of genetic regulatory effects across human

tissues. *Science* **369**, 1318–1330 (2020).

11. van der Wijst, M. *et al.* The single-cell eQTLGen consortium. *eLife* **9**, e52155 (2020).

12. Cuomo, A. S. E. *et al.* Single-cell RNA-sequencing of differentiating iPS cells reveals dynamic

genetic effects on gene expression. *Nat. Commun.* **11**, 1–14 (2020).

13. van der Wijst, M. G. P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-eQTLs and

co-expression QTLs. *Nat. Genet.* **50**, 493–497 (2018).

14. Sulston, J. E. & Horvitz, H. R. Post-embryonic cell lineages of the nematode, Caenorhabditis

elegans. *Dev. Biol.* **56**, 110–156 (1977).

15. Hall, D. H. & Altun, Z. F. *C. elegans atlas*. (Cold Spring Harbor Laboratory Press, 2007).

16. Hobert, O., Glenwinkel, L. & White, J. Revisiting Neuronal Cell Type Classification in

Caenorhabditis elegans. *Curr. Biol. CB* **26**, R1197–R1203 (2016).

17. Burga, A., Ben-David, E., Vergara, T. L., Boocock, J. & Kruglyak, L. Fast genetic mapping of

complex traits in C. elegans using millions of individuals in bulk. *Nat. Commun.* **10**, 2680 (2019).

18. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for

Dimension Reduction. *ArXiv180203426 Cs Stat* (2018).

19. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected

communities. *Sci. Rep.* **9**, 1–12 (2019).

20. Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, 381–386 (2014).

21. Cao, J. *et al.* Comprehensive single-cell transcriptional profiling of a multicellular organism. *Science*

357, 661–667 (2017).

22. Packer, J. S. *et al.* A lineage-resolved molecular atlas of C. elegans embryogenesis at single-cell

resolution. *Science* **365**, (2019).

23. Mandric, I. *et al.* Optimized design of single-cell RNA sequencing experiments for cell-type-specific

eQTL analysis. *Nat. Commun.* **11**, 5504 (2020).

24. Rockman, M. V., Skrovanek, S. S. & Kruglyak, L. Selection at linked sites shapes heritable

phenotypic variation in C. elegans. *Science* **330**, 372–376 (2010).

25. Pukkila-Worley, R. & Ausubel, F. M. Immune defense mechanisms in the Caenorhabditis elegans

intestinal epithelium. *Curr. Opin. Immunol.* **24**, 3–9 (2012).

26. Ono, K. & Ono, S. Tropomyosin and Troponin Are Required for Ovarian Contraction in the

Caenorhabditis elegans Reproductive System. *Mol. Biol. Cell* **15**, 2782–2793 (2004).

27. Gumienny, T. L. TGF-β signaling in C. elegans. *WormBook* 1–34 (2013)

doi:10.1895/wormbook.1.22.2.

28. Suzuki, Y. *et al.* A BMP homolog acts as a dose-dependent regulator of body size and male tail

patterning in Caenorhabditis elegans. *Dev. Camb. Engl.* **126**, 241–250 (1999).

29. Andersen, E. C. *et al.* A Powerful New Quantitative Genetics Platform, Combining Caenorhabditis

elegans High-Throughput Fitness Assays with a Large Collection of Recombinant Strains. *G3*

Genes Genomes Genet. **5**, 911–920 (2015).

30. Hammarlund, M., Hobert, O., Miller, D. M. & Sestan, N. The CeNGEN Project: The Complete Gene

Expression Map of an Entire Nervous System. *Neuron* **99**, 430–433 (2018).

31. Taylor, S. R. *et al.* Expression profiling of the mature C. elegans nervous system by single-cell

RNA-Sequencing. *bioRxiv* 737577 (2019) doi:10.1101/737577.

32. Portman, D. S. & Emmons, S. W. Identification of C. elegans sensory ray genes using whole-

genome expression profiling. *Dev. Biol.* **270**, 499–512 (2004).

33. Sarkar, A. K. *et al.* Discovery and characterization of variance QTLs in human induced pluripotent

stem cells. *PLOS Genet.* **15**, e1008045 (2019).

34. Francesconi, M. & Lehner, B. The effects of genetic variation on gene expression dynamics during

development. *Nature* **505**, 208–211 (2014).

35. Yao, C. *et al.* Dynamic Role of trans Regulation of Gene Expression in Relation to Complex Traits.

Am. J. Hum. Genet. **100**, 571–580 (2017).

36. Kolberg, L., Kerimov, N., Peterson, H. & Alasoo, K. Co-expression analysis reveals interpretable

gene modules controlled by trans-acting genetic variants. *eLife* **9**, e58705 (2020).

37. Brynedal, B. *et al.* Large-Scale trans-eQTLs Affect Hundreds of Transcripts and Mediate Patterns of

Transcriptional Co-regulation. *Am. J. Hum. Genet.* **100**, 581–591 (2017).

38. Smith, E. N. & Kruglyak, L. Gene–Environment Interaction in Yeast Gene Expression. *PLOS Biol.* **6**,

e83 (2008).

39. Zhang, T. *et al.* Cell-type–specific eQTL of primary melanocytes facilitates identification of

melanoma susceptibility genes. *Genome Res.* **28**, 1621–1635 (2018).

40. Westra, H.-J. *et al.* Cell Specific eQTL Analysis without Sorting Cells. *PLOS Genet.* **11**, e1005223

(2015).

41. Orozco, L. D. *et al.* Unraveling Inflammatory Responses using Systems Genetics and Gene-

Environment Interactions in Macrophages. *Cell* **151**, 658–670 (2012).

42. Fairfax, B. P. & Knight, J. C. Genetics of gene expression in immunity to infection. *Curr. Opin.*

Immunol. **30**, 63–71 (2014).

43. Kim-Hellmuth, S. *et al.* Genetic regulatory effects modified by immune activation contribute to

autoimmune disease associations. *Nat. Commun.* **8**, 266 (2017).

44. Zhang, S., Banerjee, D. & Kuhn, J. R. Isolation and Culture of Larval Cells from C. elegans. *PLOS*

ONE **6**, e19505 (2011).

45. Kaletsky, R. *et al.* The C. elegans adult neuronal IIS/FOXO transcriptome reveals adult phenotype

regulators. *Nature* **529**, 92–96 (2016).

46. Qiu, X. *et al.* Single-cell mRNA quantification and differential analysis with Census. *Nat. Methods*

14, 309–315 (2017).

47. Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet based single-

cell RNA sequencing data. *bioRxiv* 303727 (2020) doi:10.1101/303727.

48. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830

(2011).

49. Broman, K. W. The Genomes of Recombinant Inbred Lines. *Genetics* **169**, 1133–1146 (2005).

50. Arends, D., Prins, P., Jansen, R. C. & Broman, K. W. R/qtl: high-throughput multiple QTL mapping.

Bioinforma. Oxf. Engl. **26**, 2990–2992 (2010).

51. Dodds, K. G. *et al.* Construction of relatedness matrices using genotyping-by-sequencing data.

BMC Genomics **16**, 1047 (2015).

52. Bilton, T. P. *et al.* Accounting for Errors in Low Coverage High-Throughput Sequencing Data When

Constructing Genetic Maps Using Biparental Outcrossed Populations. *Genetics* **209**, 65–76 (2018).

53. Rockman, M. V. & Kruglyak, L. Recombinational Landscape and Population Genomics of

Caenorhabditis elegans. *PLOS Genet.* **5**, e1000419 (2009).

54. Rockman, M. V. & Kruglyak, L. Breeding Designs for Recombinant Inbred Advanced Intercross

Lines. *Genetics* **179**, 1069–1078 (2008).

55. Svensson, V. Droplet scRNA-seq is not zero-inflated. *Nat Biotechnol* **38**, 147–150 (2020).

56. Wood, S. N. *Generalized Additive Models: An Introduction with R*. (Chapman and Hall/CRC, 2017).

57. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor RNA-Seq

experiments with respect to biological variation. *Nucleic Acids Res* **40**, 4288–4297 (2012).

58. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful

approach to multiple testing. *J R Stat Soc* (1995).

59. Dupuis, J. & Siegmund, D. Statistical methods for mapping quantitative trait loci from a dense set of

markers. *Genetics* **151**, 373–386 (1999).

60. Albert, F. W., Bloom, J. S., Siegel, J., Day, L. & Kruglyak, L. Genetics of trans-regulatory variation

in gene expression. *eLife* **7**, e35471 (2018).

61. Miao, Z., Deng, K., Wang, X. & Zhang, X. DEsingle for detecting three types of differential

expression in single-cell RNA-seq data. *Bioinformatics* **34**, 3223–3224 (2018).

62. Whitlock, M. C. Combining probability from independent tests: the weighted Z-method is superior to

Fisher's approach. *J. Evol. Biol.* **18**, 1368–1373 (2005).

63. Smith, E. N. & Kruglyak, L. Gene–Environment Interaction in Yeast Gene Expression. *PLOS Biol.* **6**,

e83 (2008).

64. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. *Bioinforma. Oxf. Engl.* **32**, 2847–2849 (2016).