# Avalanche criticality in individuals, fluid intelligence and working memory

Longzhou Xu (徐龙洲) [a, 1], Lianchun Yu (俞连春) [a, b, c, 1, *], Jianfeng Feng (冯建峰) [d, e, f, *]

[a] School of Physical Science and Technology, and Key Laboratory for Magnetism and Magnetic Materials of MOE, Lanzhou University, Lanzhou, 730000, China

[b] Lanzhou Center for Theoretical Physics, Key Laboratory of Theoretical Physics of Gansu Province, Lanzhou University, Lanzhou, Gansu 730000, China

[c] The School of Nationalities' Educators, Qinghai Normal University, Xining, 810000, China

[d] Institute of Science and Technology for Brain Inspired Intelligence, Fudan University, Shanghai, 200433, China

[e] Department of Computer Science, University of Warwick, Coventry, CV4 7AL, UK

[f] School of Mathematical Sciences, School of Life Science and the Collaborative Innovation Center for Brain Science, Fudan University, Shanghai, 200433, PR China

[*] Corresponding authors: Lianchun Yu (yulch@lzu.edu.cn), Jianfeng Feng (Jianfeng.Feng@warwick.ac.uk).

[1] These authors contributed equally to this work: Longzhou Xu, Lianchun Yu.

# Abstract

The critical brain hypothesis suggests that efficient neural computation can be realized through dynamics of the brain characterized by scale-free avalanche activities. However, the relation between human cognitive performance and the avalanche criticality in large-scale brain networks remains unclear. In this study, we analyzed the mean synchronization and synchronization entropy of blood oxygenation level signals from resting-state fMRI. We found that the scale-free avalanche activity was associated with intermediate synchronization and maximal synchronization entropy. The complexity of functional connectivity, as well as structure-function coupling, is maximized by criticality, which is consistent with theoretical predictions. We observed order-disorder phase transitions in resting-state brain dynamics and found that there were longer times spent in the subcritical regime. These results support the hypothesis that large-scale brain networks lie in the vicinity of a critical point. Finally, we showed evidence that the neural dynamics of human participants with higher fluid intelligence and working memory scores are closer to criticality. We identified brain regions whose critical dynamics showed significant positive correlations with fluid intelligence performance, and found these regions were located in the prefrontal cortex and inferior parietal cortex, which are believed to be important nodes of brain networks underlying human intelligence. Our results reveal the role that avalanche criticality plays in cognitive performance,

39 and provide a simple method to identify the critical point and map cortical states

40 on a spectrum of neural dynamics, ranging from subcriticality to supercriticality.

41 KEY WORDS: avalanche criticality, fluid intelligence, resting-state fMRI,

42 large-scale brain network, phase transition, working memory

# Introduction

The critical brain hypothesis states that the brain operates in close vicinity to a critical point that lies between order and disorder. This is characterized by a power-law form of the event size distribution (Cocchi, Gollo, Zalesky, & Breakspear, 2017; Hesse & Gross, 2014). This hypothesis is supported by a set of observations of power-law scaling in many different neural systems using various approaches (J. M. Beggs & Plenz, 2003; Gal & Marom, 2013; Meisel, Olbrich, Shriki, & Achermann, 2013; Plenz, 2012; Shriki et al., 2013; Solovey, Miller, Ojemann, Magnasco, & Cecchi, 2012; Enzo Tagliazucchi, Balenzuela, Fraiman, & Chialvo, 2012). Arguments in favour of this hypothesis have been strengthened by advantages in information transmission, information storage, and dynamic range, in neural systems operating near criticality (Shew, Yang, Petermann, Roy, & Plenz, 2009; Shew, Yang, Yu, Roy, & Plenz, 2011; Yang, Shew, Roy, & Plenz, 2012), with evidence arising in both theoretical and experimental work (Shew & Plenz, 2012). Meanwhile, this hypothesis still faces challenges from several perspectives (J. Beggs & Timme, 2012). For example, computational studies suggested that power laws may emerge from simple stochastic processes or noncritical neuronal systems (Touboul & Destexhe, 2010), so power laws alone are prerequisite but not sufficient evidence for criticality. Meanwhile, it has been asked: "If the brain is critical, what is the phase transition (Fontenele et al., 2019)?" Indeed, the observation of power law

avalanche activity along with a phase transition between order and disorder would be more persuasive for criticality. Furthermore, though previous studies have associated supercriticality with reduced consciousness (Meisel et al., 2013; Scott et al., 2014), near-critical dynamics with rest (Priesemann et al., 2014), and subcriticality with focused cognitive states (Fagerholm et al., 2015), there remains a gap between the specific brain state and efficient information processing endowed by criticality as predicted by theory (He, 2011). To fully understand the functional roles of critical and non-critical dynamics, more research is required to relate brain states and cognitive performance to neural dynamics that lie on a spectrum, ranging from subcriticality to supercriticality. To obtain a deeper understanding of this phenomenon, it is necessary to develop data analysis methods to represent this phase spectrum with high resolution and characterize the subsequent reorganization of brains with the transition in this spectrum (Fontenele et al., 2019).

With advances in brain imaging techniques such as functional magnetic resonance imaging (fMRI), the critical brain hypothesis has found roles in interpreting fundamental properties of large-scale brain networks in the context of structure-dynamics-function relationships (Karahanoğlu & Van De Ville, 2017; Lee et al., 2019). For example, it has been shown that structural connections of brains are mostly reflected in functional networks, and this structure-function coupling is disrupted when brains move away from criticality

during anesthesia (Enzo Tagliazucchi et al., 2016). Another application of

criticality is to explain the dynamic basis of brain complexity (Popiel, 2020; E.

Tagliazucchi & Chialvo, 2013; Timme et al., 2016). In particular, functional

connectivity (FC) complexity, which is an umbrella term describing the

variability, diversity, or flexibility of functional connections in brain networks,

has been associated with cognitive performance from many perspectives, such

as high-order cognition, aging, and cognitive impairment in brain disorders

(Ahmadlou, Adeli, Bajo, & Adeli, 2014; Anokhin, Birbaumer, Lutzenberger,

Nikolaev, & Vogel, 1996; Omidvarnia, Zalesky, Van De Ville, Jackson, &

Pedersen, 2019; Smyser et al., 2016; B. Wang et al., 2017). Studies have

suggested that the FC complexity may possibly be at its maximum at the critical

point, while the FC capacity arises from special topological properties of the

structural network, such as hierarchical modular organization (Song et al., 2019;

R. Wang et al., 2019).

To validate these applications, both computer modeling methods and

experimental data analysis methods were used. Computer modeling utilizes

structural imaging data to model large-scale brain dynamics and functional

networks (Deco, Jirsa, & McIntosh, 2011; Nakagawa, Jirsa, Spiegler, McIntosh,

& Deco, 2013). However, there is still disagreement on which type of phase

transition should be adopted for large-scale brain networks, e.g., first-order

discontinuous vs. second-order continuous phase transitions and edge of chaos

criticality vs. avalanche criticality (Kanders, Lorimer, & Stoop, 2017; Scarpetta,

Apicella, Minati, & de Candia, 2018). Experimental studies usually take

advantage of dynamic changes caused by interventions, such as deprived sleep,

anesthesia, or brain diseases, to show deviations from criticality and subsequent

reorganization of FC networks (Hobbs, Smith, & Beggs, 2010; Meisel et al.,

2013; Meisel, Storch, Hallmeyer-Elgner, Bullmore, & Gross, 2012; Enzo

Tagliazucchi et al., 2016). However, deviations caused by these interventions

are usually unidirectional, either in the sub- or supercritical directions.

Furthermore, it is not clear whether deviations from criticality caused by

different intervention methods follow an identical phase transition trajectory.

Recent studies have proposed the concept of a "critical line" instead of a

"critical point" and suggested that multiple phase transition trajectories may

exist (Kanders, Lee, Hong, Nam, & Stoop, 2020). Therefore, the successful

retrieval of the phase transition trajectory from the large-scale brain networks

will not only help to answer key questions regarding what the phase transition is

if the brain is critical but also have important implications in brain functional

imaging and large-scale brain modeling.

In this study we applied a large number of criticality-related metrics to

investigate the spatio-temporal dynamics of large scale brain networks as well

as their variation in individuals, and explored associations with three different

cognitive abilities in a sample of 295 healthy young adults from the Human

Connectome Project (HCP) 1200-subject release (Van Essen et al., 2013). We retrieved an inverted-U curve by mapping individuals' brains onto the phase plane between the mean synchronization (MS) and synchronization entropy (SE) of blood oxygenation level-dependent (BOLD) signals. We then performed the classic avalanche criticality analysis method on these data (J. M. Beggs & Plenz, 2003; Enzo Tagliazucchi et al., 2012), and found that subjects who were closest to the critical point exhibited moderate mean and maximal variability in synchrony of BOLD signals (i.e., located around the tipping point of the inverted-U curve). This is consistent with previous findings that the neural systems operate neither at the synchronous nor the asynchronous ends of the spectrum, but rather near the critical point between them (Fontenele et al., 2019). We also showed that the large individual variation around the critical point could be the cause of the mismatch between group-level data analysis and theoretical prediction. However, this individual variation also gave us a chance to examine previous conjectures on criticality in large-scale brain networks. And we indeed found that both FC complexity and structure-function coupling were maximized around the criticality. We also utilized a sliding window approach to observe "instantaneous" phase transition occurring in individual brains. We found that brains persisting in the subcritical regime exhibited longer dwell times than those in other regimes. Finally, we found the critical dynamics were associated with high scores in fluid intelligence and working memory tests, but

not with crystallized intelligence scores. Additionally, the critical dynamics in the frontal cortex, superior parietal lobule, angular gyrus and supramarginal gyrus et al, which were believed as vital regions in the networks of Parieto-Frontal Integration Theory (P-FIT) for intelligence (Jung & Haier, 2007), exhibited significant correlations with fluid intelligence performance.

# Material and methods

## Data acquisition and preprocessing

*fMRI data acquisition and preprocessing*

We used rfMRI data from the Human Connectome Project (HCP) 1200-subject release (Van Essen et al., 2013). Each subject underwent two sessions of rfMRI on separate days, each session with two separate 14 min 24s acquisitions generating 1200 volumes on a customized Siemens 3T Skyra scanner ( $TR = 720 \, ms$ , $TE = 33 \, ms$ , flip angle $= 52°$ , voxel size $= 2 \, mm$ isotropic , 72 slices , $FOV = 208 \times 180 mm$, matrix $= 104 \times 90 \, mm$ , multiband accelaration factor $= 8$ , echo spacing $= 0.58 \, ms$ ). The rfMRI data used for our analysis were processed according to the HCP minimal preprocessing pipeline (Glasser et al., 2016; Glasser et al., 2013) and denoising procedures. The denoising procedure pairs the independent component analysis with the FSL tool FIX to remove non-neural spatiotemporal components (Smith et al., 2015). And as a part of cleanup, HCP used 24 confound time series

derived from the motion estimation (the 6 rigid-body parameter time series, their backwards-look temporal derivatives, plus all 12 resulting regressors squared). Note that the global component of the fMRI fluctuations measured during the resting state is tightly coupled with the underlying neural activity, and the use of global signal regression as a pre-processing step in resting-state fMRI analyses remains controversial and is not universally recommended (Liu, Nalci, & Falahpour, 2017). Therefore, the global whole-brain signal was not removed in this work. We used the left-to-right acquisitions from the first resting-state dataset (i.e., resting state fMRI 1 FIX-denoised package).

The first 324 subjects in the dataset entered into our study, and we excluded 29 subjects for missing data. This left us with 295 subjects for further analysis, and 162 of them were females. All the participants were between the ages of 22-36, 58 were between the ages of 22-25, 130 were between the ages of 26-30, 104 were between the ages of 31-35, and 3 were 36 years old.

For further analysis, the whole cortex was parcellated into 96 regions using the Harvard-Oxford atlas (Makris et al., 2006), and the details are provided at https://identifiers.org/neurovault.image:1699, and from each region the time series averaged across the voxels were extracted and Z-normalized to construct region of interest (ROI) signals (atlas$_{96}$ signals). To test the results for different parcellation, we also used the Human Brainnetome atlas that comprised 246 regions (Fan et al., 2016), and Zalesky atlas that comprised 1024 regions

(Zalesky et al., 2010), the resulting signals were termed as atlas$_{246}$ and atlas$_{1024}$

signals, respectively.

*Diffusion tensor imaging (DTI) data acquisition and preprocessing*

The diffusion MRI images used in this study were also from the HCP

1200-subject release (Sotiropoulos et al., 2013). Briefly, the diffusion data were

collected using a single-shot, single refocusing spin-echo, echo-planar imaging

sequence ($\text{TR} = 5520\,\text{ms}$, $\text{TE} = 89.5\,\text{ms}$, flip angle $= 78\,°$, $\text{FOV} = 210 \times$

$180\,\text{mm}$, $\text{matrix} = 168 \times 144\,\text{mm}$, $\text{voxel size} = 1.25\,\text{mm istropic}$,

$\text{slices} = 111$, multiband acceleration factor $= 3$, echo spacing $= 0.78\,\text{ms}$).

Three gradient tables of 90 diffusion-weighted directions and six b=0 images,

each were collected with right-to-left and left-to-right phase encoding polarities

for each of the three diffusion weightings (b=1000, 2000, and 3000 s/mm$^2$). All

diffusion data were preprocessed with the HCP diffusion pipeline updated with

EDDY 5.0.10 (Sotiropoulos et al., 2013), and the details are provided at

https://www.humanconnectome.org. In this study, from the 295 selected subjects,

only 284 subjects were entered into our DTI data analysis because 11 of them

were missing the corresponding DTI data.

*Cognition measures*

We examined associations between cognitive ability and critical dynamics

conducted in our rfMRI analysis. Three relevant behavioral tasks were used as a

measure of cognitive ability, including fluid intelligence, working memory, and crystallized intelligence. The same 295 subjects were also entered into our cognitive ability analysis, but for the fluid intelligence analysis, 5 subjects were excluded due to missing their intelligence scores, or information about age and education.

The fluid intelligence scores in the HCP data release were measured using the number of correct responses on form A of the Penn Matrix Reasoning Test (PMAT, mean $= 17.0034$, standard deviation (SD) $= 4.9106$, range $= 4 - 24$), which had 24 items and 3 bonus items, using nonverbal visual geometric designs with pieces to assess reasoning abilities that can be administered in under 10 min (Barch et al., 2013; Hearne, Mattingley, & Cocchi, 2016). The PMAT (Bilker et al., 2012) is an abbreviated version of Raven's Standard Progressive Matrices test (Wendelken, Nakhabenko, Donohue, Carter, & Bunge, 2007), which comprises 60 items.

Crystallized intelligence was measured using the picture vocabulary test (Picture vocabulary, mean=116.8205, SD=10.1977, range=92.3914-153.0889) from the National Institutes of Health (NIH) toolbox (Barch et al., 2013; Hearne et al., 2016). This measure of receptive vocabulary was administered in a computer-adaptive testing (CAT) format. The participant was presented with four pictures and heard an audio recording saying a word, and was instructed to select the picture that most closely showed the meaning of the word. Because

the test used a variable length CAT with a maximum of twenty-five items, some

participants had fewer items, and the presented words depended on the

participant's performance.

Working memory was assessed using the List Sorting Working Memory test

(List sorting, mean=111.2075, SD=12.0946, range=84.63-144.50) from the NIH

Toolbox (Barch et al., 2013), in which the participants were required to

sequence sets of visually and a small number of orally presented stimuli in size

order from smallest to biggest. Pictures of different foods and animals were

displayed with both a sound clip and a written test that names them, and

involved two different conditions. In the 1-list condition, participants ordered a

series of objects, either food or animals, but in the 2-list condition, participants

were presented with both animal and food lists and asked to order each list by

increasing size. The number of list items increased in subsequent trials, and the

task was discontinued after 2 consecutive incorrect trials.

**Data analysis methods**

*Synchrony and variability in synchrony*

We measured the mean and variability in synchronization with a previously

described approach (Meisel et al., 2013; Yang et al., 2012). First, we obtained

the phase trace $\theta_j(t)$ from the signal $F_j(t)$ using its Hilbert transform

$H[F_j(t)]$:

$$\theta_j(t) = arctan \frac{H[F_j(t)]}{F_j(t)}. \qquad (1)$$

Next, we calculated the Kuramoto order parameter as follows:

$$r(t) = \frac{1}{u}\left|\sum_{j=1}^{u} e^{i\theta_j(t)}\right|, \qquad (2)$$

in which u is the number of ROIs in global network analysis, or the number of voxels in a particular region in regional analysis. The Kuramoto order parameter $r(t)$ was used as a time-dependent measure of phase synchrony of a system. The MS of a time period was calculated as

$$\langle r \rangle = \frac{1}{V}\sum_{t=1}^{V} r(t), \qquad (3)$$

where $V$ is the length of the time period. In this study, we calculated static MS of the entire scan period with $V = 1200$ time points. We derived the entropy of $r(t)$ as the measure of variability in synchronization (synchronization entropy, SE):

$$H(r) = -\sum_{n=1}^{N} p_n \, log_2 \, p_n, \qquad (4)$$

where $p_n$ is the probability that $r(t)$ falls into a bin between $min\,(r(t)) \leq b_n < r(t) < b_{n+1} \leq max\,(r(t))$ . In this study, we chose the number of bins $N = 30$, and the robustness of our results was also tested within an interval between 5 and 100.

*Avalanche analysis*

In our avalanche analysis, the ROI signals were reduced to a spatiotemporal

point process by detecting the suprathreshold peak positions intermediate

between two above-threshold time points, as shown in the example in Fig. 1A.

By binning the binary sequences with appropriate time resolution (time bin), we

obtained a spatial pattern of active ROIs within consecutive time bins. An

avalanche was defined as a series of consecutively active bins, which were led

and followed by blank bins without activation. The size S and duration T of the

avalanches were then defined as the total number of activations and total

number of time bins during this avalanche, respectively (J. M. Beggs & Plenz,

2003; Enzo Tagliazucchi et al., 2012).

If a system operates near a critical point, the size distribution ($P(S)$), duration

distribution ($P(T)$), and average size for a given duration ($\langle S \rangle(T)$) should be

fitted into power laws:

282
$$P(S) \sim S^{-\alpha}, \tag{5}$$

283
$$P(T) \sim T^{-\tau}, \tag{6}$$

284
$$\langle S \rangle(T) \sim T^{\gamma}, \tag{7}$$

285     where $\alpha$, $\tau$, and $\gamma$ are critical exponents of the system (Friedman et al., 2012;

286     Sethna, Dahmen, & Myers, 2001). Furthermore, the following scaling relation

287     was proposed as an important evaluation of the criticality (Fontenele et al., 2019;

288     Friedman et al., 2012), namely,

289
$$\frac{\tau - 1}{\alpha - 1} = \gamma. \tag{8}$$

In this study, we defined

$$\delta = \left| \frac{\tau-1}{\alpha-1} - \gamma \right| \tag{9}$$

to measure the distances of the systems from the critical point, so the smaller $\delta$ is, the closer the systems are to the critical point.

The scaling exponents governing the power-law distribution were estimated using the maximum likelihood estimator (MLE) (Clauset, Shalizi, & Newman, 2009; Marshall et al., 2016). Briefly, the MLE procedure supposes that the empirical data sample is from a power-law function in the range $(x_{min}, x_{max})$, with probability density $\frac{1}{\sum_{x=x_{min}}^{x_{max}} (\frac{1}{x})^\alpha} \left(\frac{1}{x}\right)^\alpha$ (Fontenele et al., 2019; Marshall et al., 2016). We estimated critical exponents $\alpha$ and $\tau$ by maximizing the likelihood function and via a lattice search algorithm (Marshall et al., 2016). We then used Clauset's goodness-of-fit test to quantify the plausibility of fits (Clauset et al., 2009; Deluca & Corral, 2013; Marshall et al., 2016). We used a power-law model to produce data sets over the fit range and compared the Kolmogorov–Smirnov (KS) statistics between (1) the real data and the fit against (2) the model data and the fit. If the real data produced a KS-statistic that was less than the KS-statistic found for at least 10% of the power-law models (i.e., p ≥ 0.1), we accepted the data as being fit by the truncated power law because the fluctuations of the real data from the power law were similar in the KS sense to random fluctuations in a perfect power-law model.

*Surrogate data*

To assess the statistical significance of the avalanche analysis results and MS-SE relationship, we generated comparable surrogate data and applied the analyses above to these data. Phase-shuffling is often used in hypothesis testing for avalanche size distribution (Gireesh & Plenz, 2008; Shriki et al., 2013). Phase-shuffling disrupts temporal as well as spatial correlations in multichannel time series.

Herein phase shuffling was done on the atlas$_{96}$ signals. The phase randomization procedures were as follows (Prichard & Theiler, 1994): (1) the discrete Fourier transformation was taken to of each subject; (2) rotating the phase at each frequency by an independent random variable that was uniformly chosen in the range $[0,2\pi]$. Crucially, the different time series were rotated by the different phases to randomize the phase information; (3) the inverse discrete Fourier transformation was applied to these time series to yield surrogate data.

*Branching parameter*

The branching parameter $\sigma$, which is defined as the average number of subsequent events that a single preceding event in an avalanche triggers, is a convenient measure to identify criticality (J. M. Beggs & Plenz, 2003). In theory, the system is critical for $\sigma = 1$ and sub- (super) critical for $\sigma < 1$ ($\sigma > 1$). In this study, σ was calculated as

$$\sigma = \frac{1}{N}\sum_{i=1}^{N}\frac{n_i^*}{n_i},$$ (10)

where $n_i$ is the number of ancestors, $n_i^*$ is the number of descendants in the next time bin, and $N$ is the total number of time bins with activations.

*Definition of kappa*

A nonparametric measure, $\kappa$, for neuronal avalanches was introduced by Shew and his colleagues (Shew et al., 2009). It quantifies the difference between an experimental cumulative density function (CDF) of the avalanche size, $F(\beta_k)$, and the theoretical reference CDF, $F^{NA}(\beta_k)$, which is a power-law function with theoretical expected exponent $\alpha = 1.5$:

$$\kappa = 1 + \frac{1}{m}\sum_{k=1}^{m}(F^{NA}(\beta_k) - F(\beta_k)),$$ (11)

where $\beta_k$ are avalanche sizes logarithmically spaced between the minimum and maximum observed avalanche sizes, and $m$ is the number of histogram bins. The unit value of $\kappa$ is characteristic of the system in a critical state, whereas values below and above 1 suggest sub- and supercritical states, respectively.

*Functional and structure connectivity matrix*

We constructed an FC matrix from atlas$_{96}$ signals by computing the Pearson correlation $C_{ij}$ between ROI $i$ and ROI $j$, and the mean FC strength $\langle FC \rangle$ was obtained by

$$\langle FC \rangle = \langle |C_{ij}| \rangle, \tag{12}$$

where $|\cdot|$ means the absolute value.

The structure connectivity (SC) matrix was constructed using DSI Studio (http://dsi-studio.labsolver.org) from DTI data. The DTI data were reconstructed in the Montreal Neurological Institute (MNI) space using q-space diffeomorphic reconstruction (F.-C. Yeh & Tseng, 2011) to obtain the spin distribution function (F. Yeh, Wedeen, & Tseng, 2010). A diffusion sampling length ratio of 1.25 was used. The restricted diffusion was quantified using restricted diffusion imaging (F.-C. Yeh, Liu, Hitchens, & Wu, 2017), and a deterministic fiber tracking algorithm (F.-C. Yeh, Verstynen, Wang, Fernández-Miranda, & Tseng, 2013) was used to obtain one million fibers with whole-brain seeding. The angular threshold was randomly selected from 15 degrees to 90 degrees. The step size was randomly selected from 0.1 voxels to 3 voxels. The anisotropy threshold was automatically determined by DSI Studio. The fiber trajectories were smoothed by averaging the propagation direction with a percentage of the previous direction. The percentage was randomly selected from 0% to 95%. Tracks with a length shorter than 5 or longer than 300 mm were discarded. The SC matrix was calculated by using the count of the connecting tracks using 96-region Harvard-Oxford atlas.

*Functional connectivity entropy*

The FC entropy $H(FC)$ is calculated by

$$H(FC) = \sum_i - p_i log_2(p_i), \qquad (13)$$

where $p_i$ is the probability distribution of $|C_{ij}|$, i.e., $\sum_i p_i = 1$ (Yao et al., 2013). In the calculation, the probability distribution was obtained by discretizing the interval (0, 1) into 30 bins.

*Functional connectivity diversity*

The functional diversity ($D(FC)$) of the FC matrix is measured by the similarity of the distribution to the uniform distribution (R. Wang et al., 2019):

$$D(FC) = 1 - \frac{1}{N_M} \sum_{i=1}^{M} \left| p_i - \frac{1}{M} \right|, \qquad (14)$$

where $N_M = 2\frac{M-1}{M}$ is a normalization factor, $D(FC)$ is in the range [0, 1], and $p_i$ is the probability distribution of $|C_{ij}|$, which was obtained by discretizing the interval (0, 1) into M bins ($M = 30$ in this work). For completely asynchronous or synchronized states, the correlation values fall into one bin at 0 or 1, where $D(FC) = 0$ reflects the simple dynamic interaction pattern. In an extreme case where all types of FC equivalently exist, $p_i$ would ideally follow a uniform distribution (i.e., probability in each bin $= \frac{1}{M}$) and $D(FC) = 1$.

*Functional connectivity flexibility*

To obtain the flexibility of FC in the whole brain, we utilized the sliding

window method to calculate connectivity number entropy ($CNE$) for each

region (Lei et al., 2020; Song et al., 2019). A non-overlapping sliding window

method was applied to the atlas$_{96}$ signals. The choice of window size must be

sufficient to yield a stable Pearson's correlation coefficient within each window,

yet small enough to reveal the temporal-dependent variation in FC (Lei et al.,

2020; Sakoğlu et al., 2010). We chose a window size in the range of 20-30,

corresponding to the number of windows ($n_{win}$) in the range of 40-60.

Within each time window, we first acquired the FC matrix via their time series

in this window. Then, the binary network matrix was obtained by binarizing the

FC matrix with a threshold $THR_{FC}$. Subsequently, we calculated the number of

regions connected to a particular region $k$ ($k = 1, 2, \ldots 96$) in each time

window. Therefore, we could obtain $p_i$, the probability for a particular

connection number occurring, where $i$ indicated the $i\text{-}th$ connection number

among all possible connection numbers. Then, for the region $k$, $CNE_k$ is a

complexity measure (i.e., Shannon entropy) for the disorder in the connection

numbers over time:

$$CNE_k = -\sum_{i=1}^{95} p_i log_2 p_i, \tag{15}$$

404   where the summation index runs from 1 to the number of all possible

405   connection numbers.

406   For each subject, the $CNE$ at the whole-brain level was obtained by simply

407  averaging the regional $CNE_k$ values over 96 regions:

$$CNE = \frac{1}{96}\sum_{k=1}^{96} CNE_k. \tag{16}$$

*Similarity between functional and structural networks*

To measure the similarity between functional and structural connection

networks, the FC matrices $C_{ij}$ were thresholded by $THR_{FC}$ to yield binary

adjacency matrices $A_{ij}$ such that $A_{ij} = 1$ if $C_{ij} \geq THR_{FC}$, and $A_{ij} = 0$

otherwise. The parameter $THR_{FC}$ was chosen to fix link density $\rho_{FC}$, which

was defined as the ratio of the connections in the network ($\sum_{i>j} A_{ij}$) to the total

possible number of connections. It is important to fix the link density when

comparing networks, as otherwise, differences could arise because the average

of the respective $C_{ij}$ are different (and therefore the number of nonzero entries

in $A_{ij}$) but not because connections are topologically reorganized across

conditions (Enzo Tagliazucchi et al., 2016).

The binary FC networks for each subject were compared with the

group-aggregated binary SC network but not with the individual's SC network

to avoid fluctuations in individual SC networks. First, the binary adjacency

matrices $B_{ij}$ of SC matrices were obtained for each subject such that $B_{ij} = 1$ if

there were tracked fiber links; otherwise, $B_{ij} = 0$. Then, the binary adjacency

structural connection matrices were summed up and again thresholded by a

thresholding value $THR_{SC}$ to yield a group-aggregated binary SC network. In

this way, high $THR_{SC}$ values would exclude connections that were shared by

fewer subjects but preserve connections that were common in most subjects.

To estimate the similarity between the binary FC network of each subject and

the group-aggregated binary SC network, we computed the Pearson correlation

$R(FC - SC)$ and Hamming distance $HD(FC - SC)$ between these two

networks (Enzo Tagliazucchi et al., 2016). Specifically, the Hamming distance

is defined as the number of symbol substitutions (in this case 0 or 1) needed to

transform one sequence into another and vice versa, and in this case, it is equal

to the number of connections that must be rewired to turn the functional

network connection into the structural network connection.

*Dynamic analysis of phase transitions*

We used the sliding window approach to capture the time-dependent changes in

measures used in this study. In the calculation, for the atlas$_{96}$ signals, the length

of the sliding window was set to $V = 200$ (volumes), and the sliding step was

set to $\Delta n = 10$ (volumes). In each window, we calculated the corresponding

dynamic measures, including dynamic MS $\langle r \rangle_n$, dynamic SE $H(r)_n$, and

dynamic FC matrix $\left( C_{ij} \right)_n$. From the dynamic FC matrix $\left( C_{ij} \right)_n$, we further

obtained dynamic FC entropy $H(FC)_n$, FC diversity $D(FC)_n$, Pearson

correlation between FC and SC $R(FC - SC)_n$, and Hamming distance

$HD(FC - SC)_n$.

*Data and code availability*

The rfMRI data, DTI data and cognitive data are available from the Human

Connectome Project at humanconnectome.org, WU-Minn Consortium. The

WU-Minn HCP Consortium obtained full informed consent from all participants,

and research procedures and ethical guidelines were followed in accordance

with the Washington University Institutional Review Boards (IRB #201204036;

Title: 'Mapping the Human Connectome: Structure, Function, and Heritability').

MATLAB (https://www.mathworks.com/) and SPSS

(https://www.ibm.com/analytics/spss-statistics-software) were used to conduct

the experiment's reported in this study. The datasets supporting this article and

the codes required to reproduce them can be found at

https://github.com/longzhou-xu/data_and_code_sort.git .

# Results

## The signature of critical dynamics in the cortical network

**Figure 1. Avalanche statistics obtained from group-level analysis. A.** Example of a point

process (red triangles) extracted from one normalized ROI BOLD signal. **B.** The probability

distributions of group-aggregated avalanche sizes for the threshold 1.4 SD and the time bin

width of 1 volume (vol.) in the fMRI data. The distributions are well approximated by power

law with an exponent of $\alpha = 1.56$ with Clauset's test $p = 0.99$, corresponding to $\kappa =$

0.9912 (histogram bins=40) and branching parameter $\sigma = 0.9097$. **C.** The distribution of

avalanche durations can be fitted well by a power law with an exponent of $\tau = 1.84$ under

the condition described in **B**, with Clauset's test $p = 0.3$. **D.** There is a relation between the

sizes and duration of the avalanches with a positive index $\gamma = 1.59$, which is close to $\frac{1-\tau}{1-\alpha} =$

1.51. In **B-D**, the grey open triangles were calculated from the surrogate data. **E.** The

branching ratio and power-law scaling exponents $\alpha$ of avalanche sizes for different

thresholds used to define the point process.

For the 295 available subjects, we first investigated the power-law distribution

of avalanche size at the population level. Here, we defined the activation as the

time point when the BOLD signals reached their peak value, while the signals

one step before and after this time point were above the chosen threshold (Fig.

1A). After preprocessing, the atlas$_{96}$ signals were converted into point processes

in which each time point represented an activation. We then calculated the

avalanche size distribution $P(S) \sim S^{-\alpha}$ (Fig. 1B), as well as the avalanche

duration distribution $P(T) \sim T^{-\tau}$ (Fig. 1C). First, from the estimated $\alpha$ and $\tau$

values, we tested whether the relationship between the scaling exponents holds

for different thresholds of ROI signals (Fontenele et al., 2019; Friedman et al.,

2012). We found that the closest matching occurred when the chosen threshold

was around 1.4 SD (Fig. 1D). Second, the power-law distribution of avalanche

sizes with a slope of $\alpha = 1.5$ could be predicted by theory for a critical

branching process with branching parameter $\sigma = 1$ (Harris, 1964; Zapperi,

Lauritsen, & Stanley, 1995). However, we found that the threshold of 1.4 SD

yielded $\sigma = 0.91$ and $\alpha = 1.56$ (Fig. 1E), which did not match well with the

theoretical prediction. We ran the same analysis on both atlas$_{246}$ signals (Fig.

S1a) and atlas$_{1024}$ signals (Fig. S1b) to find the mismatch still exists (Fig.

S1c-d).

**Figure 2. Signatures of criticality as a function of MS in resting-state brain networks. A.** Top panel: The inverted-U trajectory of the MS $\langle r \rangle$ vs. SE $H(r)$. The red dash line represents the quadratic fit of the data ($F$=188.758, $p$<0.001, adjusted $R^2$=0.561). Bottom panel: The frequency count for the distribution of MS $\langle r \rangle$. **B.** The branching parameters $\sigma$ vs. $\langle r \rangle$ for each subject. The green dashed line indicates $\sigma = 1$. The Pearson correlation value $R$ and the $p$ value are shown in the figure. The red dashed line represents the linear regression. For further analysis, we selected three representative groups of subjects according to their synchronization level: namely, LMS group ($\langle r \rangle = 0.2824 \pm 0.0219$, blue open circles in A and B), MMS group ($\langle r \rangle = 0.5041 \pm 0.0042$, green open circles in A and B) and HMS group ($\langle r \rangle = 0.6304 \pm 0.0246$, red open circles in A and B). **C.** Avalanche size distributions for the LMS group, MMS group, and HMS group. To show the difference between these groups, we used gray lines with $\alpha = 1.47$ to guide the eyes. The

corresponding group-aggregated branching parameters are $\sigma_{LMS} = 0.7237$ for the LMS

group, $\sigma_{MMS} = 1.0123$ for the MMS group, and $\sigma_{HMS} = 1.2023$ for the HMS group. **D.**

Avalanche duration distributions for three groups in **C**. To show the difference between these

groups, we used gray lines with $\tau = 1.7$ to guide the eyes. **E.** Scaling relations for the three

groups. The blue line and purple line correspond to $\gamma$ and $\frac{\tau-1}{\alpha-1}$, respectively. In the inset,

$\delta = \left| \gamma - \frac{\tau-1}{\alpha-1} \right|$ indicates the distance to the critical point. **F.** The dependence of $\kappa$ on the

numbers $m$ of histogram bins for the three groups.

As the above close check of hallmarks of criticality did not agree with each

other well, we moved forward to investigate whether this mismatch could be a

result of intersubject variability. We calculated both the MS and SE (see

Synchrony and variability in synchrony) using the atlas$_{96}$ signals for each of the

295 subjects and characterized the brain states of each subject with points in the

MS vs. SE phase plane, as seen in the top panel of Fig. 2A. We found that the

value of MS from these subjects extended from 0.2 to 0.7, and the distribution

of subjects was not even but exhibited a greater tendency to the low MS range

(Fig. 2A, bottom panel). This result suggested that even in the resting state,

there is significant variability among the subjects' brain states. It is clearly seen

that these state points formed an inverted-U trajectory in the phase plane. The

SE exhibited a maximum at the moderate value of MS, which implied the

existence of a state with dynamic richness between order and disorder. We

found the inverted-U curves and the calculation of SE were robust against

different parcellation (Fig. S2). We also performed a phase randomization

method on the fMRI data and found that this inverted-U curve disappeared in

the randomized surrogate datasets (size=500, identified by visual inspection;

examples can be seen in Fig. S3). Therefore, we argued that this inverted-U

curve reflected a special spatiotemporal structure of brain dynamics that did not

exist in randomized data.

As expected, with the increasing of MS, the spatiotemporal activation pattern

defined before exhibited transitions from random states to ordered states (Fig.

S4). We then calculated the branching parameter $\sigma$ for each subject. We found

that with increasing MS, the branching parameter increased from less than 1 to

higher than 1, crossing 1 at a moderate value of MS (Fig. 2B, and Fig. S5 for

different parcellation).

Furthermore, we selected three groups from the above subjects: the low mean

synchronization group (LMS group; the 20 most left subjects in Fig. 2A with an

MS value of $\langle r \rangle = 0.2824 \pm 0.0219$ ), the moderate mean synchronization

group (MMS group; the 20 subjects located near the peak of curve in Fig. 2A

with an MS value of $\langle r \rangle = 0.5041 \pm 0.0042$ ), and the high mean

synchronization group (HMS group; the 20 most right subjects in Fig. 2A with

an MS value of $\langle r \rangle = 0.6304 \pm 0.0246$ ). For each group, we performed

avalanche distribution analysis to identify which group was closest to the

critical point (Fig. 2C-F). After obtaining scaling exponents $\alpha$ and $\tau$ for each

group with a threshold of 1.4 SD (Fig. 2C-D), the scaling relationship showed

the best match for the MMS group (Fig. 2E), and more detailed analysis results

can be found in Fig. S6 in the Supplementary Materials.

Previous study showed that the truncations of power-law fit have a dramatic

impact on power-law exponents, particularly on the ratio $\frac{1-\tau}{1-\alpha}$, while $\gamma$ barely

changes (Destexhe & Touboul, 2020). To test the robustness of our results, we

performed analysis in Fig 2c-e with different fitting windows for avalanche size

S ( $S_{min} \in [1, 10]$ , and $S_{max} \in [30, 60]$ ) and avalanche duration T ( $T_{min} \in$

$[1, 5]$ , and $T_{max} \in [9, 20]$ ). We found that the ratio of the number of fittings

that met the critical criterion ( $|(1 - \tau)/(1 - \alpha) - \gamma| < 0.1$ and Clauset's

goodness-of-fit test $p > 0.1$) to all power-law-fit samples is highest for MMS

group ($\varphi = 0.0346$ for LMS, $\varphi = 0.1103$ for MMS, $\varphi = 0.0266$ for MMS).

We also calculated $\kappa$ , an often-used parameter that could distinguish the

difference between data and the theoretically suggested power-law distribution

(Fagerholm et al., 2015; Palva et al., 2013; Poil, Hardstone, Mansvelder, &

Linkenkaer-Hansen, 2012; Shew et al., 2009; Shew et al., 2011). As shown in

Fig. 2F, as the discrete bin number $m$ increases, the $\kappa$ values become stable.

The stabilized $\kappa$ is smaller than 1 for the LMS group but larger than 1 for the

HMS and MMS groups. The $\kappa$ value for MMS group was closest to 1.

Therefore, the above results suggested that subjects' brains with moderate MS

and maximal SE are poised closest to the critical point, supported by consistent

569 hallmarks of criticality. On the other hand, the large dispersion of subjects

570 among the phase space between asynchronous (subcritical) and synchronous

571 (supercritical) states also provides an opportunity to investigate the phase

572 transition in brains.

**The complexity in the FC network is maximized by criticality**

**Figure 3. Dependence of complexity in the FC network on the MS of BOLD signals. A.**

FC entropy $H(FC)$ as a function of MS $\langle r \rangle$. Red dashed line: quadratic fitting ($F =$

2287.892, $p$<0.001, adjusted $R^2 = 0.940$). **B.** FC diversity $D(FC)$ as a function of MS $\langle r \rangle$.

Red dashed line: quadratic fitting ($F = 1226.057$, $p$<0.001, adjusted $R^2 = 0.894$). **C.** FC

flexibility $CNE$ as a function of MS $\langle r \rangle$. Red dashed line: quadratic fitting ($F = 366.851$,

*p<0.001*, adjusted $R^2 = 0.715$). The red open triangles represent participants with outliers in

the quadratic fittings in **A-B.**

Since the disorder-order phase transition could be observed, we investigated

how this phase transition could impact the organization of FC networks. For

convenience, we used MS to indicate this transition. We assessed how the

variousness in FC strength changes as the brain undergoes a phase transition

from the sub- to supercritical states. We used FC entropy and FC diversity as

measures of variousness in FC strength in the brain networks. FC entropy is a

direct measure of Shannon entropy from the probability distribution of FC

strength obtained from the FC matrix, whereas FC diversity measures the

similarity between the distribution of real FC matrix elements and uniform

distribution. In previous studies, the former had been associated with healthy

aging (Yao et al., 2013), and the latter is predicted to be maximized at the

critical point by a computer model with Ginzburg-Landau equations (R. Wang

et al., 2019). We found that both FC entropy (Fig. 3A) and FC diversity (Fig. 3B)

peaked at the moderate value of MS; however, the peak position for these two

measures was more rightward than that of SE.

The flexibility in dynamic FC reflects the extent of abundant connection

patterns among regions and how frequent switching may occur between

different patterns. In this work, we adopted the connection number entropy

($CNE$) as a measure of flexibility in FC networks. Our previous study showed

that this measure was maximized at the critical point in a large-scale brain

network model that combined DTI structural data and excitable cellular

automaton (Song et al., 2019), and this measure could be reduced in the brains

of patients with moyamoya disease (Lei et al., 2020). In this study, we found

that the flexibility in FC was maximized with a moderate value of MS (Fig. 3C).

The maximization was robust in a wide range of $THR_{FC}$ thresholds and sliding

window lengths (Fig. S7). This result supported our previous conclusion (Lei et

al., 2020; Song et al., 2019). Compared with FC entropy and FC diversity, the

peak position for FC flexibility was nearer to the critical point.

FC entropy, diversity, and flexibility are often used in rfMRI studies to measure

the complexity in the structure and dynamic reconfiguration of FC networks.

Here, the study suggested that the complexity in FC networks is maximized by

criticality.

**The maximized structure-function coupling around the critical point**

**Figure 4. The dependence of structure-function coupling on the MS of brain networks.**

**A.** Pearson correlation between anatomical and functional networks as a function of MS $\langle r \rangle$.

The link density in the FC network $\rho_{FC} = 0.7$ and threshold in the group-aggregated SC

network $THR_{SC}=40$ (corresponding to $\rho_{SC} = 0.4836$) are shown in the figure. Red dashed

line: quadratic fitting ($F = 36.997$, $p < 0.001$, adjusted $R^2 = 0.197$), which is better than

linear fitting ( $F = 39.346$ , $p < 0.001$ , adjusted $R^2 = 0.115$ ). **B.** Hamming distance

$HD(FC - SC)$ between anatomical and functional networks as a function of MS $\langle r \rangle$. Red

dashed line: quadratic fitting ($F = 36.997$ , $p < 0.001$, adjusted $R^2=0.197$), which is better

than linear fitting ($F = 39.346$, $p<0.001$, adjusted $R^2 = 0.115$). **C.** The Pearson correlation

between anatomical and functional networks as a function of FC density $\rho_{FC}$   ($THR_{SC} = 40$

corresponding to $\rho_{SC} = 0.4826$) for the HMS, MMS, and LMS groups. **D**. The Hamming

distance between anatomical and functional networks as a function of FC density $\rho_{FC}$

($THR_{SC} = 40$ corresponding to $\rho_{SC} = 0.4826$) for the HMS, MMS, and LMS groups. In

**C-D**, the stars indicate significant differences between the HMS and MMS groups (two-tails

two-sample t-test, $p < 0.05$, uncorrected); the open triangles indicate significant differences

between the LMS and MMS groups (two-tails two-sample t-test, $p < 0.05$, uncorrected).

From the obtained FC matrix and SC matrix for each subject, we constructed

the FC networks for each subject and a group-aggregated SC network (see

Similarity between functional and structural networks). We used $THR_{FC}$ and

$THR_{SC}$ to control the link density in the FC networks and the group-aggregated

SC network, respectively. We measured the similarity between the FC network

and group-aggregated SC network with Pearson correlation and Hamming

distance. Fig. 4A and B demonstrate the dependence of similarity on the MS of

each subject with a link density of 0.7 in the FC networks. The similarity

between the FC and SC was maximal for subjects with moderate synchrony, as

the Pearson correlation was maximized (Fig. 4A), while the Hamming distance

was minimized (Fig. 4B) for these subjects. This maximization of similarity

between the FC and SC could be observed in a wide range of link densities in

the FC and SC networks. To further consolidate the above results, we measured

the similarity between the FC and SC for the three groups (LMS MMS and

HMS) defined above as a function of the FC network link density. Fig. 4C-D

show that as the FC network link density increased, the correlation coefficient

between the FC and SC matrices first increased and then decreased, and

consistently, the Hamming distance exhibited the opposite tendency. When the

FC link density was large, the MMS group showed a significantly higher

correlation and a lower Hamming distance between the FC and SC networks

than the other two groups. Similarly, by varying $THR_{SC}$, we found that the

maximized similarity in the FC and SC at the critical point was robust in the

wide range of link densities in the SC network (Figs. S8, S9 and S10).

We noticed that for a large link density of the FC network, the dependence of

similarity on link density monotonically decreased (Fig. 4C-D; Fig. S8a-b).

Since lower link density conserved only stronger links in FC networks, we

deduced that structural connections were mostly reflected in the strong

functional connections. Meanwhile, the similarity also decreased as the

threshold $THR_{SC}$ in SC networks decreased (Fig. S9a-b), suggesting that the

structural connections that were mostly reflected in the functional connections

were those shared by most subjects because structural connections specified to

individuals would be excluded with high $THR_{SC}$.

**The dynamic phase transition in individual subjects' brains**

**Figure 5. The dynamic phase transition in individual subjects' brains. A.** The dependence

of dynamic SE $H(r)_n$ on dynamic MS $\langle r \rangle_n$ from six subjects selected randomly from the

LMS, MMS, and HMS groups. The enlarged dark markers indicate the mean position for

corresponding subjects (markers with the same shape). **B.** The time-dependent changes in the

Kuramoto order parameter $r(t)$ for six subjects as demonstrated in figure a (with the same

color). **C.** The normalized frequency count of $r(t)$ for different levels of $\langle r \rangle$, indicated by

lines with different colors. **D.** The dwell time (the time interval between two successive

critical point crossing events) distribution for different levels of $\langle r \rangle$. **(E, F).** The distribution

of vertical and horizontal moving distances of phase points in one step of the sliding window.

**(G, H).** The vertical and horizontal velocities of state points of each subject as a function of

their MS $\langle r \rangle$. The vertical and horizontal velocities were calculated by $\frac{\langle |\Delta \langle r \rangle_n| \rangle}{\Delta n}$ and $\frac{\langle |\Delta \langle r \rangle_n| \rangle}{\Delta n}$,

where the symbol $|\cdot|$ indicates the absolute value, and $\langle \cdot \rangle$ was the average across all the

windows. $\Delta n$ is the step used to slide the windows. Here, $\Delta n = 10$ time points (volumes). Red

dashed lines in **G-H**: quadratic fitting ($F = 139.316$, $p < 0.001$, adjusted $R^2 = 0.485$ in

(G); $F = 81.181$, $p < 0.001$, adjusted $R^2 = 0.353$ in figure h). Both quadratic fittings were

better than linear fitting (adjusted $R^2 = 0.407$ in **G** and adjusted $R^2 = 0.173$ in **H**).

The observed individual brain states dispersed around the critical point provided

an opportunity to investigate the dynamic phase transition in individual brains.

To this end, for the LMS, MMS, and HMS groups defined above, we randomly

selected two subjects from each group. We calculated the dynamical MS

$\langle r \rangle_n$ and SE $H(r)_n$ for these six subjects with the sliding window approach (Fig.

5A, s1-s6) from their Kuramoto order parameters r(t) (Fig. 5B). We observed a

time-dependent change in individuals' brain states in the state space following

the inverted-U trajectory, as shown in the top panel of Fig. 2A. In the time

period limited by scan duration, we observed that subjects who were farther

away from the critical point tended to stay in the regime decided by MS, and

events of crossing the critical point (black lines at r(t) = 0.5 ) to the other

regime seldom occurred (s1, s2, s5, and s6 in Fig. 5A, or Fig. 5B, top and

bottom panel). Subjects who were nearer the critical point were more likely to cross the critical point (s3 and s4 in Fig. 5A, or Fig. 5B, middle panel).

To validate the above observation at the population level, we divided the 295 subjects at hand into eight groups with different levels of synchrony and calculated the corresponding probability distribution of the Kuramoto order parameter $r(t)$. It is seen clearly from Fig. 5C that as the synchrony level decreases, the distribution of the Kuramoto order parameter becomes narrow and less tilted. Meanwhile, we found that the dwell time, which referred to the time interval between two successive critical point crossing events, exhibited heavier tails in its distribution for low synchrony groups (Fig. 5D). These results implied the higher inertness in the subcritical regime than others, and brains were more likely to stay in this regime with longer dwell times.

Next, we calculated the distribution of vertical and horizontal moving distances in state space in a fixed time interval $\Delta n$ (the time points or volumes of one step of the sliding window) for all subjects. We found that the distributions of vertical and horizontal moving distances were both symmetrical with a mean of zero (Fig. 5E-F), suggesting that the inverted-U trajectory in the state space was stable and unlikely to change its shape as time progressed. Furthermore, the position ($\langle r \rangle$) dependent velocity distribution is maximal for horizontal velocity ($\frac{\langle |\Delta \langle r \rangle_n| \rangle}{\Delta n}$) and minimal for vertical velocity ($\frac{\langle |\Delta \langle r \rangle_n| \rangle}{\Delta n}$) near the critical point (Fig. 5G-H). The maximal horizontal velocity around the critical point implied that at

this point, the systems were most sensitive to the perturbations due to internal fluctuations or external modulations. Meanwhile, the lower vertical and horizontal velocities in the subcritical regime compared to the supercritical regime also reflected the high inertness in the subcritical regime.

**Figure 6. Dynamic modulations of FC complexity and structure-function coupling during the phase transition of brains. A.** The dependence of dynamic FC entropy as a function of instantaneous MS; thick dashed white line: quadratic fitting ($F = 106350.82$, $p < 0.001$, adjusted $R^2 = 0.877$). **B.** The dependence of dynamic FC diversity as a function of instantaneous MS. Thick dashed white line: quadratic fitting ($F = 80261.492$, $p < 0.001$, adjusted $R^2 = 0.843$). **C.** The dependence of dynamic FC-SC correlation as a

function of instantaneous MS; thick dashed white line: quadratic fitting ( $F = 5519.072$ ,

$p < 0.001$ , adjusted $R^2 = 0.270$ ), which is better than linear fitting (adjusted $R^2 = 0.225$ ).

**D.** The dependence of dynamic FC-SC Hamming distance as a function of instantaneous MS;

thick dashed white line: quadratic fitting ( $F = 5519.072$ , $p < 0.001$ , adjusted

$R^2 = 0.270$ ), which is better than linear fitting (adjusted $R^2 = 0.225$ ). In **A-D**, each dot

represents a calculation from one window. The dots with the same color represent the

calculation for one subject. However, due to the limited number of colors used, different

subjects may share the same color. In **C-D**, a link density of 0.7 was used to obtain the binary

FC network, and a threshold of 40 was used to obtain the group-aggregated structural

network.

It was of interest to determine whether the maximization of FC complexity, as

well as function-structure coupling, around the critical point could be realized

dynamically when the individual brains endured phase transition. To this end,

we obtained the time-dependent FC matrices with the sliding window method

and calculated FC entropy (Fig. 6A), FC diversity (Fig. 6B), and two measures

for similarity between FC and SC (Fig. 6C-D) as a function of instantaneous

MS in each time window. The time-dependent complexity and similarity

measures followed almost the exact trajectories as those in the static

measurements shown in Fig. 3A-B, as well as Fig. 4A-B. This result implied

that FC complexity and similarity between FC and SC were indeed modulated

by phase transition in brains, and their maximization could be realized dynamically by positioning the system around the critical point.

**High fluid intelligence and working memory capacity were associated with critical dynamics**

**Figure 7. Correlations between cognitive performance scores and SE, as well as MS.** **A-C.** Correlation between SE and PMAT scores, picture vocabulary test scores, as well as the list sorting working memory test scores. Red dashed lines in **A-C**: linear fitting. **D.** Scatterplot of the PMAT scores against the MS. The red dashed line represents the significant quadratic fit of the data ($F = 3.145$, $p = 0.045$, adjusted $R^2 = 0.015$), which is better than the linear fitting (adjusted $R^2 = 0.004$). **E.** Scatterplot of the picture vocabulary test scores against the MS. Both the linear and quadratic regressions are not significant (linear: $p = 0.991$; quadratic: $p = 0.988$). **F.** Scatterplot of the list sorting working memory test scores

against the MS. The red dashed line represents the significant quadratic fit of the data ($F =$ 4.376 , $p = 0.013$, adjusted $R^2 = 0.023$), which is better than linear fitting (adjusted $R^2 = 0.008$ ).

The results above support the hypothesis that large-scale brain networks lie in the vicinity of a critical point which is associated with moderate MS and maximal SE. Another key prediction from the critical brain hypothesis is that brains that are closer to criticality should be better in cognitive performance. Here, to address this prediction, we assessed linear relationships between SE and intelligence scores of the available subjects. We found that SE values were significantly correlated with fluid intelligence scores (PMAT; Fig. 7A) but not with crystallized intelligence scores (picture vocabulary; Fig. 7B). Meanwhile, we found that working memory scores, which were assessed using the Listing Sorting Working Memory test from the NIH Toolbox, were significantly correlated with SE (List sorting; Fig. 7C). We also noted here that these scores were significantly correlated with many other measures that were found to be maximized at the criticality, namely, FC entropy, FC diversity, and FC flexibility (Fig. S11 in Supplementary Materials). Meanwhile, there were significant quadratic relationships between MS and fluid intelligence, as well as working memory scores, but not for crystallized intelligence scores (Fig. 7D-F). We also found these results still held when potential confounds such as age and education achievements were regressed out (Fig. S12 and S13). Therefore, these

results imply that brains that are closer to criticality are associated with higher

fluid intelligence and working memory scores.

**Figure 8. The brain map for correlations between regional SE and fluid intelligence.** The

color bar indicated the Pearson correlation value between regional SE and PMAT. The

cortical and subcortical regions were defined by the Human Brainnetome Atlas

(http://atlas.brainnetome.org/bnatlas.html). Data was visualized using BrainNet Viewer (Xia,

Wang, & He, 2013).

Since a wide variety of experiments have demonstrated that fluid intelligence is

associated with a distributed network of regions in the Parieto-Frontal

Integration Theory (P-FIT), including frontal areas (Brodmann areas (BAs) 6, 9, 10, 45-47), parietal areas (BA 7, 39, 40), visual cortex (BAs 18, 19), fusiform gyrus (BA 37), Wernicke's area (BA 22) and dorsal anterior cingulate cortex (BA 32) (Jung & Haier, 2007; Nikolaidis et al., 2017), we decided to find more relationships between these regions with critical dynamics indicated by maximized SE.

To obtain the relevant regions in a fine-grained division of the brain, here we used the Human Brainnetome Atlas, which contains 210 cortical and 36 subcortical subregions (Fan et al., 2016). We extracted from each brain region the voxel-level BOLD signals and calculated the regional SE for these 246 regions. We found that regions whose SE exhibited significant ($p < 0.05$, FDR corrected) positive correlations with PMAT scores were located in the frontal areas (i.e., bilateral SFG, MFG, PrG, right IFG and PCL), parietal areas (i.e., bilateral AG, SMG, Pcun, right SPL), right inferior temporal gyrus (ITG), superior occipital gyrus (sOcG) and left Cingulate gyrus (CG) (Fig 8 and Table 1).

 **Table 1. The brain regions exhibited significant correlation between SE and fluid intelligence.**

| Lobe | Gyrus | Left/Right | Brodmann area | MNI coordinate | R value | FDR |
|---|---|---|---|---|---|---|
| Frontal Lobe | Superior Frontal Gyrus (SFG) | SFG_L_7_3 | 8, 9, 10 | [-11, 49, 40] | 0.2058 | 0.0120 |
| | | SFG_R_7_3 | 8, 9, 10 | [13, 48, 40] | 0.1795 | 0.0264 |
| | | SFG_R_7_7 | 10 | [8, 58, 13] | 0.1784 | 0.0268 |
| | Middle Frontal Gyrus (MFG) | MFG_L_7_1 | 9, 46 | [-27, 43, 31] | 0.1827 | 0.0264 |
| | | MFG_L_7_3 | 10, 11, 46 | [-28, 56, 12] | 0.1947 | 0.0176 |
| | | MFG_R_7_3 | 9, 10, 46 | [28, 55, 17] | 0.2706 | 0.0007 |
| | | MFG_L_7_5 | 8, 9, 44, 46 | [-33, 23, 45] | 0.1871 | 0.0229 |
| | | MFG_R_7_5 | 9, 44, 45, 46 | [42, 27, 39] | 0.2522 | 0.0017 |
| | | MFG_L_7_7 | 10, 11, 47 | [-26, 60, -6] | 0.1662 | 0.0429 |
| | Inferior Frontal Gyrus (IFG) | IFG_R_6_5 | 45, 47 | [42, 22, 3] | 0.1820 | 0.0264 |
| | Precentral Gyrus (PrG) | PrG_L_6_1 | 4, 6 | [-49, -8, 39] | 0.1797 | 0.0264 |
| | | PrG_R_6_1 | 4, 6 | [55, -2, 33] | 0.1718 | 0.0344 |
| | Paracentral Lobule (PCL) | PCL_R_2_2 | 4, 6 | [5, -21, 61] | 0.1728 | 0.0344 |
| Temporal Lobe | Inferior Temporal Gyrus (ITG) | ITG_R_7_5 | 20, 21, 37 | [54, -57, -8] | 0.2066 | 0.0120 |
| Parietal Lobe | Superior Parietal Lobule (SPL) | SPL_R_5_5 | 7, 40 | [31, -54, 53] | 0.1717 | 0.0344 |
| | Angular Gyrus (AG) | IPL_R_6_1 | 39 | [45, -71, 20] | 0.1687 | 0.0390 |
| | | IPL_R_6_2 | 7, 39 | [39, -65, 44] | 0.2373 | 0.0028 |
| | | IPL_L_6_5 | 39 | [-47, -65, 26] | 0.2011 | 0.0140 |
| | | IPL_R_6_5 | 39 | [53, -54, 25] | 0.2284 | 0.0043 |
| | Supramarginal Gyrus (SG) | IPL_L_6_3 | 2, 3, 40 | [-51, -33, 42] | 0.1997 | 0.0140 |
| | | IPL_R_6_3 | 2, 3, 40 | [47, -35, 45] | 0.2061 | 0.0120 |
| | | IPL_R_6_4 | 2, 40 | [57, -44, 38] | 0.2369 | 0.0028 |
| | Precuneus (Pcun) | Pcun_L_4_4 | 7 | [-6, -55, 34] | 0.1810 | 0.0264 |
| | | Pcun_R_4_4 | 7 | [6, -54, 35] | 0.1899 | 0.0219 |
| Limbic Lobe | Cingulate Gyrus (CG) | CG_L_7_6 | 23 | [-7, -23, 41] | 0.1868 | 0.0229 |
| Occipital Lobe | Superior Occipital Gyrus (sOcG) | sOcG_R_2_1 | 18, 19 | [16, -85, 34] | 0.2051 | 0.0120 |

812    Abbreviation: IPL, Inferior Parietal Lobule.

# **Discussion**

In this study, using the large scale WU-Minn HCP dataset and a large number

of criticality-inspired metrics, we provided evidence that though healthy young

brains at rest are on average close to the critical point, as critical brain

hypothesis has suggested, there is a considerable individual variation around

this point. This gave us a chance to validate previous theoretical predictions of

criticality on large scale brain networks, and indeed we observed that the

complexity of brain FC, as well as the structure-function coupling, was

maximized around the critical point. We proceeded to observe a dynamic phase

transition in individual subjects, and found that their brains tended to stay

subcritical, as indicated by a longer dwell time in this parameter region. Finally,

we found that high fluid intelligence and working memory capacity were

associated with critical dynamics rather than noncritical dynamics, not only

globally but also regionally, suggesting the functional advantages of critical

dynamics in resting-state brains.

Balance between functional segregation and functional integration is a central

organizing principle of the cerebral cortex. It has been argued that FC

complexity characterizes the interplay of functional segregation and functional

integration (Sporns, 2013). A comparison between simulated and empirically

obtained resting-state FC indicates that the human brain at rest lies in a dynamic

state that reflects the largest complexity its anatomical connectome can host (G.

Tononi, Sporns, & Edelman, 1994). Recently, many studies have tried to link

complexity with cognitive performance, human intelligence, and even

consciousness, either measured by Φ (big phi) in integrated information theory

or discriminated between levels of sedation (Ahmadlou et al., 2014; Duncan,

Chylinski, Mitchell, & Bhandari, 2017; Saxe, Calderone, & Morales, 2018;

Giulio Tononi, Edelman, & Sporns, 1998). Meanwhile, there is a growing

awareness that complexity is strongly related to criticality. A recent study

showed that criticality maximized complexity in dissociated hippocampal

cultures produced from rats (Timme et al., 2016). Here, in this study, we

measured FC complexity from different perspectives, either on its strength

diversity or on its dynamic flexibility (Fig. 3A-C, Fig. 6A-B). With the

observation of the phase transition trajectory, we demonstrated that these

measures of FC complexity were maximized around the critical point. Therefore,

the formulation that criticality maximizes complexity was supported in our

work empirically with fMRI data at the whole-brain network level.

It has been shown that human brains possess a stable set of functionally coupled

networks that echo many known features of anatomical organization (Krienen,

Yeo, & Buckner, 2014). Several computational modeling studies have

demonstrated that critical dynamics could best explore the repertoire provided

by the structural connectome (Deco & Jirsa, 2012; Enzo Tagliazucchi et al.,

2016). Recent studies also suggested the capacity of repertoire provided by the

structural connectome could be extended by the hierarchical modular structural

organization (R. Wang et al., 2019). Therefore, structure-function coupling was

believed to be at its maximal when the system is at criticality (R. Wang et al., 2019), and it could be disrupted by losing criticality (Cocchi et al., 2014), or disruption of hierarchical organization of structural networks. Previous studies in anesthetized human brains have found structure-function decoupling accompanied by unidirectional departure from a critical point (Enzo Tagliazucchi et al., 2016). It is possible that functional connectivity flexibility could be used as a measure of the extent that functional connectivity explores the repertoire provided by structural connectome, and the highest functional connectivity flexibility occurs when the system is at criticality (Song et al., 2019) (Fig. 3C). Our work demonstrated the maximal exploration of structural connections at the critical point occurs in resting state brains (Fig. 4). However, since we used a group aggregated structural connection networks, we did not investigate how organization of structural connections could impact on the capacity of network repertoire. This issue will be investigated in the future.

Interestingly, although the brain hovers around the critical point, the brain prefers to stay in the subcritical region, as the subject distribution was skewed toward a disordered state, and the dwell time in the subcritical state was longer (Fig. 5). Previous analysis of in vivo data has argued that the mammalian brain self-organizes to a slightly subcritical regime (Priesemann et al., 2014). It was suggested that operating in a slightly subcritical regime may prevent the brain from tipping over to supercriticality, which has been linked to epilepsy.

Meanwhile, with a slightly subcritical regime deviates only little from criticality, the computational capabilities may still be close to optimal. However, our results showed that the resting state brains could actually stay in the supercritical regimes. So, the preference of brains for subcritical regime may not because of prevention of too ordered states. In another study, by relating the EEG-domain cascades to spatial BOLD patterns in simultaneously recorded fMRI data, the researchers found that while resting-state cascades were associated with an approximate power-law form, the task state was associated with subcritical dynamics (Fagerholm et al., 2015). They argued that while a high dynamic range and a large repertoire of brain states may be advantageous for the resting state with near-critical dynamics, a lower dynamic range may reduce elements of interference affecting task performance in a focused cognitive task with subcritical dynamics (Fagerholm et al., 2015). Therefore, there remains a possibility that the resting state is not "pure resting state", but mixed with some occasional "task state" for some subjects. However, further delicately designed experimental studies are required to test this conjecture. It remains to uncover the relationship between cognitive states and neural dynamics that lies on a spectrum. The method proposed in this study may be useful in future studies of this topic.

Recently, Ezaki et al. used the Ising model to map BOLD signals on a two-dimensional phase space and found that human fMRI data were in the

paramagnetic phase and were close to the boundary with the spin-glass phase but not to the boundary with the ferromagnetic phase (Ezaki, Fonseca dos Reis, Watanabe, Sakaki, & Masuda, 2020). Since the spin-glass phase usually yields chaotic dynamics whereas the ferromagnetic phase is nonchaotic, their results suggested that the brain is around the "edge of chaos criticality" instead of "avalanche criticality". However, our findings support that avalanche criticality occur in large-scale brain networks. Therefore, it is interesting to investigate whether both kinds of criticality could co-occur in large-scale brain networks (Kanders et al., 2017). Ezaki et al. also found that criticality of brain dynamics was associated with human fluid intelligence, though they used performance IQ to reflect fluid intelligence, which refers to active or effortful problem solving and maintenance of information. In our work, we assessed the correlation between fluid intelligence and the critical dynamics indicated by synchronization entropy for brain regions, and found regions showed significant positive correlations were located in parietal-frontal network (Fig. 8 and Table 1). These regions were most frequently reported in studies of intelligence and its biological basis, including structural neuroimaging studies using voxel-based morphometry, magnetic resonance spectroscopy, and DTI, as well as functional imaging studies using positron emission tomography (PET) or fMRI (Jung & Haier, 2007). Also, in the Parieto-Frontal Integration Theory of intelligence, these regions are considered as the most crucial nodes of the brain network

underlying human intelligence (Jung & Haier, 2007; Nikolaidis et al., 2017).

Our study suggested that not only fluid intelligence, but also working memory

capacity was associated with critical dynamics. This is possibly because

working memory may share the same capacity constraint through similar neural

networks with fluid intelligence (Halford, Cowan, & Andrews, 2007; Jaeggi,

Buschkuehl, Jonides, & Perrig, 2008; Kane & Engle, 2002). In our study, the

critical dynamics in the frontal and parietal network also exhibited significant

correlation with working memory capacity (Fig. S14 and Table S1).

Furthermore, it has been well established that working memory is strongly

modulated by dopamine, and too strong or too weak dopamine D1 activation is

detrimental for working memory, with the optimal performance achieved at an

intermediate level (Cools & D'Esposito, 2011; Vijayraghavan, Wang, Birnbaum,

Williams, & Arnsten, 2007; Zahrt, Taylor, Mathew, & Arnsten, 1997). This

inverted-U dose-response has been observed in mice (Lidow, Koh, & Arnsten,

2003), rats (Zahrt et al., 1997), monkeys (Cai & Arnsten, 1997) and humans

(Gibbs & D'Esposito, 2005). Recent studies on neural network models have

shown that the optimal performance of working memory co-occurs with critical

dynamics at the network level and the excitation-inhibition balance at the level

of individual neurons and is modulated by dopamine at the synaptic level

through a series of U or inverted-U profiles (Hu, Huang, Jiang, & Yu, 2019).

Here in this study, we demonstrated that the optimal performance of working

memory and criticality co-occurs at the system level.

However, our study had several limitations. Firstly, the surrogate data test used

in this study ruled out the possibility that the results we obtained could be

explained by autocorrelations in the data. However, the long-range spatial

correlation of criticality cannot allow one to test the results by ruling out the

effects of correlation across the time series. Secondly, though we used the

denoising fMRI data from HCP with standard data pre-processing procedure, it

is still interesting to investigate how the pre-processing procedure affects the

results. Thirdly, in the avalanche analysis, the activation events defined in this

study were slightly different from definition used by others, such as

threshold-crossing events (Enzo Tagliazucchi et al., 2012) or above-threshold

events (Bocaccio et al., 2019; R. Wang et al., 2019). We compared these

different methods and found all these methods could generate scale free

avalanche activities, but unlike our method, the other two methods failed to

generate critical branching process (See Section II in Supplementary Materials).

Therefore, it is interesting to investigate the correlations between neural

activities and events detected by different detection methods from BOLD

signals. Finally, in this study we only focused on the cognitive abilities that are

associated with critical dynamics, and found significant but not strong

correlations between fluid intelligence, working memory and critical dynamics.

Recent works demonstrated that the functional network segregation, integration

and their balance could predict different cognitive abilities (R. Wang et al.,

2021). Therefore, future investigation on the relationship between this

functional balance and criticality across individuals may reveal various

associations between diverse cognitive abilities with not only critical dynamics,

but also non-critical dynamics. Despite all these shortages, we hope this study

may inspire future research work to validate our findings, e.g., through

observing not only the association between the departure of criticality and the

decline of cognitive performance, either in aging or brain disease, but also the

restore of criticality and the improvement of cognitive performance with

pharmacological or noninvasive brain stimulation (Barch, 2004; Reinhart &

Nguyen, 2019).

## Conclusions

In conclusion, we mapped individuals' brain dynamics from resting-state fMRI

scans on the phase transition trajectory and identify subjects who are close to

the critical point. With this approach, we validated two predictions of critical

brain hypothesis on large-scale brain networks, i.e., maximized FC complexity

and maximized structure-function coupling around the critical point. We also

observed the tendency of brain to stay in subcritical regime. Finally, we found

that the critical dynamics in large-scale brain networks were associated with

high scores in fluid intelligence and working memory, implying the vital role of

large-scale critical dynamics in cognitive performance. We also identified key

brain regions whose critical dynamics was highly correlated with human

intelligence. Our findings support the critical brain hypothesis that neural

computation is optimized by critical brain dynamics, as characterized by

scale-free avalanche activity, and could provide a solution for improving the

effects of future interventions targeting aspects of cognitive decline (Reinhart &

Nguyen, 2019), possibly by controlling the criticality through non-invasive

stimulation (Chialvo, Cannas, Grigera, Martin, & Plenz, 2020).

## Conflicts of interest

991 The authors declare no competing interests.

## Funding

This study was funded by the National Natural Science Foundation of China

(Grants No. 11105062) and the Fundamental Research Funds for the Central

Universities (Grant No. lzujbky-2021-62). J. F. is supported by the 111 Project

(Grant No. B18015), the National Key R&D Program of China

(No.2018YFC1312904; No.2019YFA0709502), the Shanghai Municipal

Science and Technology Major Project (Grant No. 2018SHZDZX01), ZJLab,

and Shanghai Center for Brain Science and Brain-Inspired Technology.

# References

1001 Ahmadlou, M., Adeli, A., Bajo, R., & Adeli, H. (2014). Complexity of functional
1002 connectivity networks in mild cognitive impairment subjects during a working memory
1003 task. *Clinical Neurophysiology, 125*(4), 694-702.
1004 doi:https://doi.org/10.1016/j.clinph.2013.08.033

Anokhin, A. P., Birbaumer, N., Lutzenberger, W., Nikolaev, A., & Vogel, F. (1996). Age increases brain complexity. *Electroencephalography and Clinical Neurophysiology, 99*(1), 63-68. doi:https://doi.org/10.1016/0921-884X(96)95573-3

Barch, D. M. (2004). Pharmacological manipulation of human working memory. *Psychopharmacology, 174*(1), 126-135. doi:10.1007/s00213-003-1732-3

Barch, D. M., Burgess, G. C., Harms, M. P., Petersen, S. E., Schlaggar, B. L., Corbetta, M., . . . Consortium, W. U.-M. H. (2013). Function in the human connectome: task-fMRI and individual differences in behavior. *NeuroImage, 80*, 169-189. doi:10.1016/j.neuroimage.2013.05.033

Beggs, J., & Timme, N. (2012). Being Critical of Criticality in the Brain. *Frontiers in Physiology, 3*, 163.

Beggs, J. M., & Plenz, D. (2003). Neuronal avalanches in neocortical circuits. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 23*(35), 11167-11177. doi:10.1523/JNEUROSCI.23-35-11167.2003

Bilker, W. B., Hansen, J. A., Brensinger, C. M., Richard, J., Gur, R. E., & Gur, R. C. (2012). Development of Abbreviated Nine-Item Forms of the Raven's Standard Progressive Matrices Test. *Assessment, 19*(3), 354-369. doi:10.1177/1073191112446655

Bocaccio, H., Pallavicini, C., Castro, M. N., Sánchez, S. M., De Pino, G., Laufs, H., . . . Tagliazucchi, E. (2019). The avalanche-like behaviour of large-scale haemodynamic activity from wakefulness to deep sleep. *Journal of The Royal Society Interface, 16*(158), 20190262. doi:10.1098/rsif.2019.0262

Cai, J. X., & Arnsten, A. F. T. (1997). Dose-Dependent Effects of the Dopamine D1 Receptor Agonists A77636 or SKF81297 On Spatial Working Memory in Aged Monkeys. *Journal of Pharmacology and Experimental Therapeutics, 283*(1), 183.

Chialvo, D. R., Cannas, S. A., Grigera, T. S., Martin, D. A., & Plenz, D. (2020). Controlling a complex system near its critical point via temporal correlations. *Scientific Reports, 10*(1), 12145. doi:10.1038/s41598-020-69154-0

Clauset, A., Shalizi, C. R., & Newman, M. E. J. (2009). Power-Law Distributions in Empirical Data. *SIAM Review, 51*(4), 661-703.

Cocchi, L., Gollo, L. L., Zalesky, A., & Breakspear, M. (2017). Criticality in the brain: A synthesis of neurobiology, models and cognition. *Progress in Neurobiology, 158*, 132-152. doi:https://doi.org/10.1016/j.pneurobio.2017.07.002

Cocchi, L., Harding, I. H., Lord, A., Pantelis, C., Yucel, M., & Zalesky, A. (2014). Disruption of structure–function coupling in the schizophrenia connectome. *NeuroImage: Clinical, 4*, 779-787. doi:https://doi.org/10.1016/j.nicl.2014.05.004

Cools, R., & D'Esposito, M. (2011). Inverted-U Shaped Dopamine Actions on Human Working Memory and Cognitive Control. *Biological Psychiatry, 69*(12), e113-e125. doi:10.1016/j.biopsych.2011.03.028

Deco, G., & Jirsa, V. K. (2012). Ongoing Cortical Activity at Rest: Criticality, Multistability, and Ghost Attractors. *The Journal of Neuroscience, 32*(10), 3366. doi:10.1523/JNEUROSCI.2523-11.2012

Deco, G., Jirsa, V. K., & McIntosh, A. R. (2011). Emerging concepts for the dynamical organization of resting-state activity in the brain. *Nature Reviews Neuroscience, 12*(1), 43-56. doi:10.1038/nrn2961

Deluca, A., & Corral, Á. (2013). Fitting and goodness-of-fit test of non-truncated and truncated power-law distributions. *Acta Geophysica, 61*(6), 1351-1394. doi:10.2478/s11600-013-0154-9

Destexhe, A., & Touboul, J. (2020). Comment on "Criticality Between Cortical States". *arXiv: Neurons and Cognition*.

Duncan, J., Chylinski, D., Mitchell, D. J., & Bhandari, A. (2017). Complexity and compositionality in fluid intelligence. *Proceedings of the National Academy of Sciences, 114*(20), 5295. doi:10.1073/pnas.1621147114

Ezaki, T., Fonseca dos Reis, E., Watanabe, T., Sakaki, M., & Masuda, N. (2020). Closer to critical resting-state neural dynamics in individuals with higher fluid intelligence. *Communications Biology, 3*(1), 52. doi:10.1038/s42003-020-0774-y

Fagerholm, E. D., Lorenz, R., Scott, G., Dinov, M., Hellyer, P. J., Mirzaei, N., . . . Leech, R. (2015). Cascades and cognitive state: focused attention incurs subcritical dynamics. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 35*(11), 4626-4634. doi:10.1523/JNEUROSCI.3694-14.2015

Fan, L., Li, H., Zhuo, J., Zhang, Y., Wang, J., Chen, L., . . . Jiang, T. (2016). The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cerebral Cortex, 26*(8), 3508-3526. doi:10.1093/cercor/bhw157

Fontenele, A. J., de Vasconcelos, N. A. P., Feliciano, T., Aguiar, L. A. A., Soares-Cunha, C., Coimbra, B., . . . Copelli, M. (2019). Criticality between Cortical States. *Physical Review Letters, 122*(20), 208101. doi:10.1103/PhysRevLett.122.208101

Friedman, N., Ito, S., Brinkman, B. A. W., Shimono, M., DeVille, R. E. L., Dahmen, K. A., . . . Butler, T. C. (2012). Universal Critical Dynamics in High Resolution Neuronal Avalanche Data. *Physical Review Letters, 108*(20), 208102. doi:10.1103/PhysRevLett.108.208102

Gal, A., & Marom, S. (2013). Self-organized criticality in single-neuron excitability. *Physical Review E, 88*(6), 062717. doi:10.1103/PhysRevE.88.062717

Gibbs, S. E. B., & D'Esposito, M. (2005). A functional MRI study of the effects of bromocriptine, a dopamine receptor agonist, on component processes of working memory. *Psychopharmacology, 180*(4), 1-10. doi:10.1007/s00213-005-0077-5

Gireesh, E. D., & Plenz, D. (2008). Neuronal avalanches organize as nested theta- and beta/gamma-oscillations during development of cortical layer 2/3. *Proceedings of the National Academy of Sciences, 105*(21), 7576. doi:10.1073/pnas.0800537105

Glasser, M. F., Smith, S. M., Marcus, D. S., Andersson, J. L. R., Auerbach, E. J., Behrens, T. E. J., . . . Van Essen, D. C. (2016). The Human Connectome Project's neuroimaging approach. *Nature Neuroscience, 19*(9), 1175-1187. doi:10.1038/nn.4361

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., . . . Jenkinson, M. (2013). The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage, 80*, 105-124. doi:https://doi.org/10.1016/j.neuroimage.2013.04.127

Halford, G. S., Cowan, N., & Andrews, G. (2007). Separating cognitive capacity from knowledge: a new hypothesis. *Trends in Cognitive Sciences, 11*(6), 236-242. doi:10.1016/j.tics.2007.04.001

Harris, T. E. (1964). The theory of branching process. *New York: Dover*.

He, B. J. (2011). Scale-free properties of the functional magnetic resonance imaging signal during rest and task. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 31*(39), 13786-13795. doi:10.1523/JNEUROSCI.2111-11.2011

Hearne, L. J., Mattingley, J. B., & Cocchi, L. (2016). Functional brain networks related to individual differences in human intelligence at rest. *Scientific Reports, 6*(1), 32328. doi:10.1038/srep32328

Hesse, J., & Gross, T. (2014). Self-organized criticality as a fundamental property of neural systems. *Frontiers in Systems Neuroscience, 8*, 166-166. doi:10.3389/fnsys.2014.00166

Hobbs, J. P., Smith, J. L., & Beggs, J. M. (2010). Aberrant Neuronal Avalanches in Cortical Tissue Removed From Juvenile Epilepsy Patients. *Journal of Clinical Neurophysiology, 27*(6).

Hu, G., Huang, X., Jiang, T., & Yu, S. (2019). Multi-Scale Expressions of One Optimal State Regulated by Dopamine in the Prefrontal Cortex. *Frontiers in Physiology, 10*, 113-113. doi:10.3389/fphys.2019.00113

Jaeggi, S. M., Buschkuehl, M., Jonides, J., & Perrig, W. J. (2008). Improving fluid
intelligence with training on working memory. *Proceedings of the National Academy of*
*Sciences, 105*(19), 6829. doi:10.1073/pnas.0801268105

Jung, R. E., & Haier, R. J. (2007). The Parieto-Frontal Integration Theory (P-FIT) of
intelligence: Converging neuroimaging evidence. *Behavioral and Brain Sciences, 30*(2),
135-154. doi:10.1017/S0140525X07001185

Kanders, K., Lee, H., Hong, N., Nam, Y., & Stoop, R. (2020). Fingerprints of a second order
critical line in developing neural networks. *Communications Physics, 3*(1), 13.
doi:10.1038/s42005-019-0276-8

Kanders, K., Lorimer, T., & Stoop, R. (2017). Avalanche and edge-of-chaos criticality do not
necessarily co-occur in neural networks. *Chaos: An Interdisciplinary Journal of Nonlinear*
*Science, 27*(4), 047408. doi:10.1063/1.4978998

Kane, M. J., & Engle, R. W. (2002). The role of prefrontal cortex in working-memory
capacity, executive attention, and general fluid intelligence: An individual-differences
perspective. *Psychonomic Bulletin & Review, 9*(4), 637-671. doi:10.3758/BF03196323

Karahanoğlu, F. I., & Van De Ville, D. (2017). Dynamics of large-scale fMRI networks:
Deconstruct brain activity to build better models of brain function. *Current Opinion in*
*Biomedical Engineering, 3*, 28-36. doi:https://doi.org/10.1016/j.cobme.2017.09.008

Krienen, F. M., Yeo, B. T. T., & Buckner, R. L. (2014). Reconfigurable task-dependent
functional coupling modes cluster around a core functional architecture. *Philosophical*
*transactions of the Royal Society of London. Series B, Biological sciences, 369*(1653),
20130526. doi:10.1098/rstb.2013.0526

Lee, H., Golkowski, D., Jordan, D., Berger, S., Ilg, R., Lee, J., . . . Vlisides, P. E. (2019).
Relationship of critical dynamics, functional connectivity, and states of consciousness in
large-scale human brain networks. *NeuroImage, 188*, 228-238.
doi:https://doi.org/10.1016/j.neuroimage.2018.12.011

Lei, Y., Song, B., Chen, L., Su, J., Zhang, X., Ni, W., . . . Mao, Y. (2020). Reconfigured
functional network dynamics in adult moyamoya disease: a resting-state fMRI study.
*Brain Imaging and Behavior, 14*(3), 715-727. doi:10.1007/s11682-018-0009-8

Lidow, M. S., Koh, P.-O., & Arnsten, A. F. T. (2003). D1 dopamine receptors in the mouse
prefrontal cortex: Immunocytochemical and cognitive neuropharmacological analyses.
*Synapse, 47*(2), 101-108. doi:10.1002/syn.10143

Liu, T. T., Nalci, A., & Falahpour, M. (2017). The global signal in fMRI: Nuisance or
Information? *NeuroImage, 150*, 213-229.
doi:https://doi.org/10.1016/j.neuroimage.2017.02.036

Makris, N., Goldstein, J. M., Kennedy, D., Hodge, S. M., Caviness, V. S., Faraone, S. V., . . . Seidman, L. J. (2006). Decreased volume of left and total anterior insular lobule in schizophrenia. *Schizophrenia Research, 83*(2), 155-171. doi:https://doi.org/10.1016/j.schres.2005.11.020

Marshall, N., Timme, N. M., Bennett, N., Ripp, M., Lautzenhiser, E., & Beggs, J. M. (2016). Analysis of Power Laws, Shape Collapses, and Neural Complexity: New Techniques and MATLAB Support via the NCC Toolbox. *Frontiers in Physiology, 7*, 250.

Meisel, C., Olbrich, E., Shriki, O., & Achermann, P. (2013). Fading Signatures of Critical Brain Dynamics during Sustained Wakefulness in Humans. *The Journal of Neuroscience, 33*(44), 17363. doi:10.1523/JNEUROSCI.1516-13.2013

Meisel, C., Storch, A., Hallmeyer-Elgner, S., Bullmore, E., & Gross, T. (2012). Failure of Adaptive Self-Organized Criticality during Epileptic Seizure Attacks. *PLoS Computational Biology, 8*(1), e1002312. doi:10.1371/journal.pcbi.1002312

Nakagawa, T. T., Jirsa, V. K., Spiegler, A., McIntosh, A. R., & Deco, G. (2013). Bottom up modeling of the connectome: Linking structure and function in the resting brain and their changes in aging. *NeuroImage, 80*, 318-329. doi:https://doi.org/10.1016/j.neuroimage.2013.04.055

Nikolaidis, A., Baniqued, P. L., Kranz, M. B., Scavuzzo, C. J., Barbey, A. K., Kramer, A. F., & Larsen, R. J. (2017). Multivariate Associations of Fluid Intelligence and NAA. *Cerebral Cortex, 27*(4), 2607-2616. doi:10.1093/cercor/bhw070

Omidvarnia, A., Zalesky, A., Van De Ville, D., Jackson, G. D., & Pedersen, M. (2019). Temporal complexity of fMRI is reproducible and correlates with higher order cognition. *bioRxiv*, 770826. doi:10.1101/770826

Palva, J. M., Zhigalov, A., Hirvonen, J., Korhonen, O., Linkenkaer-Hansen, K., & Palva, S. (2013). Neuronal long-range temporal correlations and avalanche dynamics are correlated with behavioral scaling laws. *Proceedings of the National Academy of Sciences of the United States of America, 110*(9), 3585-3590. doi:10.1073/pnas.1216855110

Plenz, D. (2012). Neuronal avalanches and coherence potentials. *The European Physical Journal Special Topics, 205*(1), 259-301. doi:10.1140/epjst/e2012-01575-5

Poil, S.-S., Hardstone, R., Mansvelder, H. D., & Linkenkaer-Hansen, K. (2012). Critical-State Dynamics of Avalanches and Oscillations Jointly Emerge from Balanced Excitation/Inhibition in Neuronal Networks. *The Journal of Neuroscience, 32*(29), 9817. doi:10.1523/JNEUROSCI.5990-11.2012

Popiel, N. J. K., S.; Abeyasinghe, P.M.; Riganello, F.; Nichols, E.S.; Owen, A.M.; Soddu, A. (2020). The Emergence of Integrated Information, Complexity, and 'Consciousness' at Criticality. *Entropy, 22*.

Prichard, D., & Theiler, J. (1994). Generating surrogate data for time series with several simultaneously measured variables. *Physical Review Letters, 73*(7), 951-954. doi:10.1103/PhysRevLett.73.951

Priesemann, V., Wibral, M., Valderrama, M., Pröpper, R., Le Van Quyen, M., Geisel, T., . . . Munk, M. H. J. (2014). Spike avalanches in vivo suggest a driven, slightly subcritical brain state. *Frontiers in Systems Neuroscience, 8*, 108-108. doi:10.3389/fnsys.2014.00108

Reinhart, R. M. G., & Nguyen, J. A. (2019). Working memory revived in older adults by synchronizing rhythmic brain circuits. *Nature Neuroscience, 22*(5), 820-827. doi:10.1038/s41593-019-0371-x

Sakoğlu, Ü., Pearlson, G. D., Kiehl, K. A., Wang, Y. M., Michael, A. M., & Calhoun, V. D. (2010). A method for evaluating dynamic functional network connectivity and task-modulation: application to schizophrenia. *Magnetic Resonance Materials in Physics, Biology and Medicine, 23*(5), 351-366. doi:10.1007/s10334-010-0197-8

Saxe, G. N., Calderone, D., & Morales, L. J. (2018). Brain entropy and human intelligence: A resting-state fMRI study. *PloS One, 13*(2), e0191582. doi:10.1371/journal.pone.0191582

Scarpetta, S., Apicella, I., Minati, L., & de Candia, A. (2018). Hysteresis, neural avalanches, and critical behavior near a first-order transition of a spiking neural network. *Physical Review E, 97*(6), 062305. doi:10.1103/PhysRevE.97.062305

Scott, G., Fagerholm, E. D., Mutoh, H., Leech, R., Sharp, D. J., Shew, W. L., & Knöpfel, T. (2014). Voltage Imaging of Waking Mouse Cortex Reveals Emergence of Critical Neuronal Dynamics. *The Journal of Neuroscience, 34*(50), 16611. doi:10.1523/JNEUROSCI.3474-14.2014

Sethna, J. P., Dahmen, K. A., & Myers, C. R. (2001). Crackling noise. *Nature, 410*(6825), 242-250. doi:10.1038/35065675

Shew, W. L., & Plenz, D. (2012). The Functional Benefits of Criticality in the Cortex. *The Neuroscientist, 19*(1), 88-100. doi:10.1177/1073858412445487

Shew, W. L., Yang, H., Petermann, T., Roy, R., & Plenz, D. (2009). Neuronal Avalanches Imply Maximum Dynamic Range in Cortical Networks at Criticality. *The Journal of Neuroscience, 29*(49), 15595. doi:10.1523/JNEUROSCI.3864-09.2009

Shew, W. L., Yang, H., Yu, S., Roy, R., & Plenz, D. (2011). Information capacity and transmission are maximized in balanced cortical networks with neuronal avalanches. *The*

*Journal of neuroscience : the official journal of the Society for Neuroscience, 31*(1), 55-63. doi:10.1523/JNEUROSCI.4637-10.2011

Shriki, O., Alstott, J., Carver, F., Holroyd, T., Henson, R. N. A., Smith, M. L., . . . Plenz, D. (2013). Neuronal avalanches in the resting MEG of the human brain. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 33*(16), 7079-7090. doi:10.1523/JNEUROSCI.4286-12.2013

Smith, S. M., Nichols, T. E., Vidaurre, D., Winkler, A. M., Behrens, T. E. J., Glasser, M. F., . . . Miller, K. L. (2015). A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nature Neuroscience, 18*(11), 1565-1567. doi:10.1038/nn.4125

Smyser, C. D., Snyder, A. Z., Shimony, J. S., Mitra, A., Inder, T. E., & Neil, J. J. (2016). Resting-State Network Complexity and Magnitude Are Reduced in Prematurely Born Infants. *Cerebral cortex (New York, N.Y. : 1991), 26*(1), 322-333. doi:10.1093/cercor/bhu251

Solovey, G., Miller, K., Ojemann, J., Magnasco, M., & Cecchi, G. (2012). Self-Regulated Dynamical Criticality in Human ECoG. *Frontiers in Integrative Neuroscience, 6*, 44.

Song, B., Ma, N., Liu, G., Zhang, H., Yu, L., Liu, L., & Zhang, J. (2019). Maximal flexibility in dynamic functional connectivity with critical dynamics revealed by fMRI data analysis and brain network modelling. *Journal of Neural Engineering, 16*(5), 056002. doi:10.1088/1741-2552/ab20bc

Sotiropoulos, S. N., Jbabdi, S., Xu, J., Andersson, J. L., Moeller, S., Auerbach, E. J., . . . Consortium, W. U.-M. H. (2013). Advances in diffusion MRI acquisition and processing in the Human Connectome Project. *NeuroImage, 80*, 125-143. doi:10.1016/j.neuroimage.2013.05.057

Sporns, O. (2013). Network attributes for segregation and integration in the human brain. *Current Opinion in Neurobiology, 23*(2), 162-171. doi:https://doi.org/10.1016/j.conb.2012.11.015

Tagliazucchi, E., Balenzuela, P., Fraiman, D., & Chialvo, D. (2012). Criticality in Large-Scale Brain fMRI Dynamics Unveiled by a Novel Point Process Analysis. *Frontiers in Physiology, 3*, 15.

Tagliazucchi, E., & Chialvo, D. R. (2013). Brain complexity born out of criticality. *AIP Conference Proceedings, 1510*(1), 4-13. doi:10.1063/1.4776495

Tagliazucchi, E., Chialvo, D. R., Siniatchkin, M., Amico, E., Brichant, J.-F., Bonhomme, V., . . . Laureys, S. (2016). Large-scale signatures of unconsciousness are consistent with a

departure from critical dynamics. *Journal of the Royal Society, Interface, 13*(114), 20151027-20151027. doi:10.1098/rsif.2015.1027

Timme, N. M., Marshall, N. J., Bennett, N., Ripp, M., Lautzenhiser, E., & Beggs, J. M. (2016). Criticality Maximizes Complexity in Neural Tissue. *Frontiers in Physiology, 7*, 425-425. doi:10.3389/fphys.2016.00425

Tononi, G., Edelman, G. M., & Sporns, O. (1998). Complexity and coherency: integrating information in the brain. *Trends in Cognitive Sciences, 2*(12), 474-484. doi:10.1016/S1364-6613(98)01259-5

Tononi, G., Sporns, O., & Edelman, G. M. (1994). A measure for brain complexity: relating functional segregation and integration in the nervous system. *Proceedings of the National Academy of Sciences, 91*(11), 5033. doi:10.1073/pnas.91.11.5033

Touboul, J., & Destexhe, A. (2010). Can power-law scaling and neuronal avalanches arise from stochastic dynamics? *PloS One, 5*(2), e8982-e8982. doi:10.1371/journal.pone.0008982

Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E. J., Yacoub, E., Ugurbil, K., & Consortium, W. U.-M. H. (2013). The WU-Minn Human Connectome Project: an overview. *NeuroImage, 80*, 62-79. doi:10.1016/j.neuroimage.2013.05.041

Vijayraghavan, S., Wang, M., Birnbaum, S. G., Williams, G. V., & Arnsten, A. F. T. (2007). Inverted-U dopamine D1 receptor actions on prefrontal neurons engaged in working memory. *Nature Neuroscience, 10*(3), 376-384. doi:10.1038/nn1846

Wang, B., Niu, Y., Miao, L., Cao, R., Yan, P., Guo, H., . . . Zhang, H. (2017). Decreased Complexity in Alzheimer's Disease: Resting-State fMRI Evidence of Brain Entropy Mapping. *Frontiers in Aging Neuroscience, 9*, 378-378. doi:10.3389/fnagi.2017.00378

Wang, R., Lin, P., Liu, M., Wu, Y., Zhou, T., & Zhou, C. (2019). Hierarchical Connectome Modes and Critical State Jointly Maximize Human Brain Functional Diversity. *Physical Review Letters, 123*(3), 038301. doi:10.1103/PhysRevLett.123.038301

Wang, R., Liu, M., Cheng, X., Wu, Y., Hildebrandt, A., & Zhou, C. (2021). Segregation, integration, and balance of large-scale resting brain networks configure different cognitive abilities. *Proceedings of the National Academy of Sciences, 118*(23), e2022288118. doi:10.1073/pnas.2022288118

Wendelken, C., Nakhabenko, D., Donohue, S. E., Carter, C. S., & Bunge, S. A. (2007). "Brain Is to Thought as Stomach Is to ??": Investigating the Role of Rostrolateral Prefrontal Cortex in Relational Reasoning. *Journal of Cognitive Neuroscience, 20*(4), 682-693. doi:10.1162/jocn.2008.20055

Xia, M., Wang, J., & He, Y. (2013). BrainNet Viewer: A Network Visualization Tool for Human Brain Connectomics. *PloS One, 8*(7), e68910. doi:10.1371/journal.pone.0068910

Yang, H., Shew, W. L., Roy, R., & Plenz, D. (2012). Maximal Variability of Phase Synchrony in Cortical Networks with Neuronal Avalanches. *The Journal of Neuroscience, 32*(3), 1061. doi:10.1523/JNEUROSCI.2771-11.2012

Yao, Y., Lu, W. L., Xu, B., Li, C. B., Lin, C. P., Waxman, D., & Feng, J. F. (2013). The Increase of the Functional Entropy of the Human Brain with Age. *Scientific Reports, 3*(1), 2853. doi:10.1038/srep02853

Yeh, F.-C., Liu, L., Hitchens, T. K., & Wu, Y. L. (2017). Mapping immune cell infiltration using restricted diffusion MRI. *Magnetic Resonance in Medicine, 77*(2), 603-612. doi:10.1002/mrm.26143

Yeh, F.-C., & Tseng, W.-Y. I. (2011). NTU-90: A high angular resolution brain atlas constructed by q-space diffeomorphic reconstruction. *NeuroImage, 58*(1), 91-99. doi:https://doi.org/10.1016/j.neuroimage.2011.06.021

Yeh, F.-C., Verstynen, T. D., Wang, Y., Fernández-Miranda, J. C., & Tseng, W.-Y. I. (2013). Deterministic Diffusion Fiber Tracking Improved by Quantitative Anisotropy. *PloS One, 8*(11), e80713. doi:10.1371/journal.pone.0080713

Yeh, F., Wedeen, V. J., & Tseng, W. I. (2010). Generalized q-Sampling Imaging. *IEEE Transactions on Medical Imaging, 29*(9), 1626-1635. doi:10.1109/TMI.2010.2045126

Zahrt, J., Taylor, J. R., Mathew, R. G., & Arnsten, A. F. (1997). Supranormal stimulation of D1 dopamine receptors in the rodent prefrontal cortex impairs spatial working memory performance. *The Journal of neuroscience : the official journal of the Society for Neuroscience, 17*(21), 8528-8535. doi:10.1523/JNEUROSCI.17-21-08528.1997

Zalesky, A., Fornito, A., Harding, I. H., Cocchi, L., Yücel, M., Pantelis, C., & Bullmore, E. T. (2010). Whole-brain anatomical networks: Does the choice of nodes matter? *NeuroImage, 50*(3), 970-983. doi:https://doi.org/10.1016/j.neuroimage.2009.12.027

Zapperi, S., Lauritsen, K. B., & Stanley, H. E. (1995). Self-Organized Branching Processes: Mean-Field Theory for Avalanches. *Physical Review Letters, 75*(22), 4071-4074. doi:10.1103/PhysRevLett.75.4071

1307    **Table 1. The brain regions exhibited significant correlation between SE and fluid intelligence.**

| Lobe | Gyrus | Left/Right | Brodmann area | MNI coordinate | R value | FDR |
|---|---|---|---|---|---|---|
| Frontal Lobe | Superior Frontal Gyrus (SFG) | SFG_L_7_3 | 8, 9, 10 | [-11, 49, 40] | 0.2058 | 0.0120 |
| | | SFG_R_7_3 | 8, 9, 10 | [13, 48, 40] | 0.1795 | 0.0264 |
| | | SFG_R_7_7 | 10 | [8, 58, 13] | 0.1784 | 0.0268 |
| | Middle Frontal Gyrus (MFG) | MFG_L_7_1 | 9, 46 | [-27, 43, 31] | 0.1827 | 0.0264 |
| | | MFG_L_7_3 | 10, 11, 46 | [-28, 56, 12] | 0.1947 | 0.0176 |
| | | MFG_R_7_3 | 9, 10, 46 | [28, 55, 17] | 0.2706 | 0.0007 |
| | | MFG_L_7_5 | 8, 9, 44, 46 | [-33, 23, 45] | 0.1871 | 0.0229 |
| | | MFG_R_7_5 | 9, 44, 45, 46 | [42, 27, 39] | 0.2522 | 0.0017 |
| | | MFG_L_7_7 | 10, 11, 47 | [-26, 60, -6] | 0.1662 | 0.0429 |
| | Inferior Frontal Gyrus (IFG) | IFG_R_6_5 | 45, 47 | [42, 22, 3] | 0.1820 | 0.0264 |
| | Precentral Gyrus (PrG) | PrG_L_6_1 | 4, 6 | [-49, -8, 39] | 0.1797 | 0.0264 |
| | | PrG_R_6_1 | 4, 6 | [55, -2, 33] | 0.1718 | 0.0344 |
| | Paracentral Lobule (PCL) | PCL_R_2_2 | 4, 6 | [5, -21, 61] | 0.1728 | 0.0344 |
| Temporal Lobe | Inferior Temporal Gyrus (ITG) | ITG_R_7_5 | 20, 21, 37 | [54, -57, -8] | 0.2066 | 0.0120 |
| Parietal Lobe | Superior Parietal Lobule (SPL) | SPL_R_5_5 | 7, 40 | [31, -54, 53] | 0.1717 | 0.0344 |
| | Angular Gyrus (AG) | IPL_R_6_1 | 39 | [45, -71, 20] | 0.1687 | 0.0390 |
| | | IPL_R_6_2 | 7, 39 | [39, -65, 44] | 0.2373 | 0.0028 |
| | | IPL_L_6_5 | 39 | [-47, -65, 26] | 0.2011 | 0.0140 |
| | | IPL_R_6_5 | 39 | [53, -54, 25] | 0.2284 | 0.0043 |
| | Supramarginal Gyrus (SG) | IPL_L_6_3 | 2, 3, 40 | [-51, -33, 42] | 0.1997 | 0.0140 |
| | | IPL_R_6_3 | 2, 3, 40 | [47, -35, 45] | 0.2061 | 0.0120 |
| | | IPL_R_6_4 | 2, 40 | [57, -44, 38] | 0.2369 | 0.0028 |
| | Precuneus (Pcun) | Pcun_L_4_4 | 7 | [-6, -55, 34] | 0.1810 | 0.0264 |
| | | Pcun_R_4_4 | 7 | [6, -54, 35] | 0.1899 | 0.0219 |
| Limbic Lobe | Cingulate Gyrus (CG) | CG_L_7_6 | 23 | [-7, -23, 41] | 0.1868 | 0.0229 |
| Occipital Lobe | Superior Occipital Gyrus (sOcG) | sOcG_R_2_1 | 18, 19 | [16, -85, 34] | 0.2051 | 0.0120 |

1308    Abbreviation: IPL, Inferior Parietal Lobule.

1309

1310

1311

**Figure 1. Avalanche statistics obtained from group-level analysis. A.** Example of a point

process (red triangles) extracted from one normalized ROI BOLD signal. **B.** The probability

distributions of group-aggregated avalanche sizes for the threshold 1.4 SD and the time bin

width of 1 volume (vol.) in the fMRI data. The distributions are well approximated by power

law with an exponent of $\alpha = 1.56$ with Clauset's test $p = 0.99$, corresponding to $\kappa =$

0.9912 (histogram bins=40) and branching parameter $\sigma = 0.9097$. **C.** The distribution of

avalanche durations can be fitted well by a power law with an exponent of $\tau = 1.84$ under

the condition described in **B**, with Clauset's test $p = 0.3$. **D.** There is a relation between the

sizes and duration of the avalanches with a positive index $\gamma = 1.59$, which is close to $\frac{1-\tau}{1-\alpha} =$

1.51. In **B-D**, the grey open triangles were calculated from the surrogate data. **E.** The

branching ratio and power-law scaling exponents $\alpha$ of avalanche sizes for different

thresholds used to define the point process.

1324

**Figure 2. Signatures of criticality as a function of MS in resting-state brain networks. A.**

Top panel: The inverted-U trajectory of the MS $\langle r \rangle$ vs. SE $H(r)$. The red dash line

represents the quadratic fit of the data ($F$=188.758, $p$<0.001, adjusted $R^2$=0.561). Bottom

panel: The frequency count for the distribution of MS $\langle r \rangle$. **B.** The branching parameters $\sigma$

vs. $\langle r \rangle$ for each subject. The green dashed line indicates $\sigma = 1$. The Pearson correlation

value $R$ and the $p$ value are shown in the figure. The red dashed line represents the linear

regression. For further analysis, we selected three representative groups of subjects according

to their synchronization level: namely, LMS group ($\langle r \rangle = 0.2824 \pm 0.0219$, blue open

circles in A and B), MMS group ($\langle r \rangle = 0.5041 \pm 0.0042$, green open circles in A and B) and

HMS group ($\langle r \rangle = 0.6304 \pm 0.0246$, red open circles in A and B). **C.** Avalanche size

distributions for the LMS group, MMS group, and HMS group. To show the difference

between these groups, we used gray lines with $\alpha = 1.47$ to guide the eyes. The

corresponding group-aggregated branching parameters are $\sigma_{LMS} = 0.7237$ for the LMS

group, $\sigma_{MMS} = 1.0123$ for the MMS group, and $\sigma_{HMS} = 1.2023$ for the HMS group. **D.**

Avalanche duration distributions for three groups in **C**. To show the difference between these

groups, we used gray lines with $\tau = 1.7$ to guide the eyes. **E.** Scaling relations for the three

groups. The blue line and purple line correspond to $\gamma$ and $\frac{\tau-1}{\alpha-1}$, respectively. In the inset,

$\delta = \left| \gamma - \frac{\tau-1}{\alpha-1} \right|$ indicates the distance to the critical point. **F.** The dependence of $\kappa$ on the

numbers $m$ of histogram bins for the three groups.

1345

1346

1347

1348

1349

1350

1351

1352

1353

1354

1355

**Figure 3. Dependence of complexity in the FC network on the MS of BOLD signals. A.** FC entropy $H(FC)$ as a function of MS $\langle r \rangle$. Red dashed line: quadratic fitting ($F = 2287.892$, $p<0.001$, adjusted $R^2 = 0.940$). **B.** FC diversity $D(FC)$ as a function of MS $\langle r \rangle$. Red dashed line: quadratic fitting ($F = 1226.057$, $p<0.001$, adjusted $R^2 = 0.894$). **C.** FC flexibility $CNE$ as a function of MS $\langle r \rangle$. Red dashed line: quadratic fitting ($F = 366.851$, $p<0.001$, adjusted $R^2 = 0.715$). The red open triangles represent participants with outliers in the quadratic fittings in **A-B**.

**Figure 4. The dependence of structure-function coupling on the MS of brain networks.**

**A.** Pearson correlation between anatomical and functional networks as a function of MS $\langle r \rangle$. The link density in the FC network $\rho_{FC} = 0.7$ and threshold in the group-aggregated SC network $THR_{SC}$=40 (corresponding to $\rho_{SC} = 0.4836$) are shown in the figure. Red dashed line: quadratic fitting ($F = 36.997$, $p < 0.001$, adjusted $R^2 = 0.197$), which is better than linear fitting ($F = 39.346$, $p < 0.001$, adjusted $R^2 = 0.115$). **B.** Hamming distance $HD(FC - SC)$ between anatomical and functional networks as a function of MS $\langle r \rangle$. Red dashed line: quadratic fitting ($F = 36.997$, $p < 0.001$, adjusted $R^2$=0.197), which is better than linear fitting ($F = 39.346$, $p$<0.001, adjusted $R^2 = 0.115$). **C.** The Pearson correlation between anatomical and functional networks as a function of FC density $\rho_{FC}$ ($THR_{SC} = 40$ corresponding to $\rho_{SC} = 0.4826$) for the HMS, MMS, and LMS groups. **D.** The Hamming

1376     distance between anatomical and functional networks as a function of FC density $\rho_{FC}$

1377     ($THR_{SC} = 40$ corresponding to $\rho_{SC} = 0.4826$) for the HMS, MMS, and LMS groups. In

1378     **C-D**, the stars indicate significant differences between the HMS and MMS groups (two-tails

1379     two-sample t-test, $p < 0.05$, uncorrected); the open triangles indicate significant differences

1380     between the LMS and MMS groups (two-tails two-sample t-test, $p < 0.05$, uncorrected).

1381

**Figure 5. The dynamic phase transition in individual subjects' brains. A.** The dependence

of dynamic SE $H(r)_n$ on dynamic MS $\langle r \rangle_n$ from six subjects selected randomly from the

LMS, MMS, and HMS groups. The enlarged dark markers indicate the mean position for

corresponding subjects (markers with the same shape). **B.** The time-dependent changes in the

Kuramoto order parameter $r(t)$ for six subjects as demonstrated in figure a (with the same

color). **C.** The normalized frequency count of $r(t)$ for different levels of $\langle r \rangle$, indicated by

lines with different colors. **D.** The dwell time (the time interval between two successive

critical point crossing events) distribution for different levels of $\langle r \rangle$. **(E, F).** The distribution

of vertical and horizontal moving distances of phase points in one step of the sliding window.

**(G, H).** The vertical and horizontal velocities of state points of each subject as a function of

their MS $\langle r \rangle$. The vertical and horizontal velocities were calculated by $\frac{\langle |\Delta \langle r \rangle_n| \rangle}{\Delta n}$ and $\frac{\langle |\Delta \langle r \rangle_n| \rangle}{\Delta n}$,

where the symbol $|\cdot|$ indicates the absolute value, and $\langle \cdot \rangle$ was the average across all the

windows. $\Delta n$ is the step used to slide the windows. Here, $\Delta n=10$ time points (volumes). Red

1396 dashed lines in **G-H**: quadratic fitting ($F = 139.316$, $p < 0.001$, adjusted $R^2 = 0.485$ in

1397 (G); $F = 81.181$, $p < 0.001$, adjusted $R^2 = 0.353$ in figure h). Both quadratic fittings were

1398 better than linear fitting (adjusted $R^2 = 0.407$ in **G** and adjusted $R^2 = 0.173$ in **H**).

1399

1400

1401

1402

1403

1404

**Figure 6. Dynamic modulations of FC complexity and structure-function coupling during the phase transition of brains. A.** The dependence of dynamic FC entropy as a function of instantaneous MS; thick dashed white line: quadratic fitting ($F = 106350.82$, $p < 0.001$, adjusted $R^2 = 0.877$). **B.** The dependence of dynamic FC diversity as a function of instantaneous MS. Thick dashed white line: quadratic fitting ($F = 80261.492$, $p < 0.001$, adjusted $R^2 = 0.843$). **C.** The dependence of dynamic FC-SC correlation as a function of instantaneous MS; thick dashed white line: quadratic fitting ($F = 5519.072$, $p < 0.001$, adjusted $R^2 = 0.270$), which is better than linear fitting (adjusted $R^2 = 0.225$). **D.** The dependence of dynamic FC-SC Hamming distance as a function of instantaneous MS; thick dashed white line: quadratic fitting ($F = 5519.072$, $p < 0.001$, adjusted

$R^2 = 0.270$ ), which is better than linear fitting (adjusted $R^2 = 0.225$ ). In **A-D**, each dot

represents a calculation from one window. The dots with the same color represent the

calculation for one subject. However, due to the limited number of colors used, different

subjects may share the same color. In **C-D**, a link density of 0.7 was used to obtain the binary

FC network, and a threshold of 40 was used to obtain the group-aggregated structural

network.

1421

1422

1423

**Figure 7. Correlations between cognitive performance scores and SE, as well as MS.**

**A-C.** Correlation between SE and PMAT scores, picture vocabulary test scores, as well as the

list sorting working memory test scores. Red dashed lines in **A-C**: linear fitting. **D.**

Scatterplot of the PMAT scores against the MS. The red dashed line represents the significant

quadratic fit of the data ($F = 3.145$, $p = 0.045$, adjusted $R^2 = 0.015$), which is better than

the linear fitting (adjusted $R^2 = 0.004$). **E.** Scatterplot of the picture vocabulary test scores

against the MS. Both the linear and quadratic regressions are not significant (linear: $p =$

$0.991$; quadratic: $p = 0.988$). **F.** Scatterplot of the list sorting working memory test scores

against the MS. The red dashed line represents the significant quadratic fit of the data ($F =$

$4.376$, $p = 0.013$, adjusted $R^2 = 0.023$), which is better than linear fitting (adjusted $R^2 =$

$0.008$).

1435

**Figure 8. The brain map for correlations between regional SE and fluid intelligence.** The color bar indicated the Pearson correlation value between regional SE and PMAT. The cortical and subcortical regions were defined by the Human Brainnetome Atlas (http://atlas.brainnetome.org/bnatlas.html). Data was visualized using BrainNet Viewer (Xia, Wang, & He, 2013).

1441