Gradual evolution of allopolyploidy in *Arabidopsis suecica*

Robin Burns[1], Terezie Mandáková[2], Joanna Jagoda[1], Luz Mayela Soto-Jiménez[1], Chang Liu[3], Martin A. Lysak[2], Polina Yu. Novikova[4,5*] and Magnus Nordborg[1*]

[1]Gregor Mendel Institute, Austrian Academy of Sciences, Vienna BioCenter, Vienna, Austria.
[2]CEITEC - Central European Institute of Technology, and Faculty of Science, Masaryk University, Brno, Czech Republic.
[3]Center for Plant Molecular Biology (ZMBP), University of Tübingen, Auf der Morgenstelle 32, 72076 Tübingen, Germany.
[4]VIB-UGent Center for Plant Systems Biology, Ghent, Belgium.
[5]Department of Chromosome Biology, Max Planck Institute for Plant Breeding Research, Cologne, Germany.

*Corresponding authors: pnovikova@mpipz.mpg.de, magnus.nordborg@gmi.oeaw.ac.at

Abstract

The majority of diploid organisms have polyploid ancestors. The evolutionary process of polyploidization (and subsequent re-diploidization) is poorly understood, but has frequently been conjectured to involve some form of "genome shock" — partly inspired by studies in crops, many of which are polyploid, and in which polyploidy has frequently been linked to dramatic genomic changes such as subgenome expression dominance and genome reorganization. It is unclear, however, whether domesticated polyploids are representative of natural ones. Here, we study polyploidization in *Arabidopsis suecica* (n = 13), a post-glacial allopolyploid species formed via hybridization of *A. thaliana* (n = 5) and *A. arenosa* (n = 8). We generated a chromosome-level genome assembly of *A. suecica* and complemented it with polymorphism and transcriptome data from multiple individuals of all species. Despite a divergence of ~6 Mya between the two ancestral species and appreciable differences in their genome composition, we see no evidence of a genome shock: the *A. suecica* genome is highly colinear with the ancestral genomes, there is no subgenome dominance in expression, and transposable element dynamics appear to be stable. We do, however, find strong evidence for changes suggesting gradual adaptation to polyploidy. In particular, the *A. thaliana* subgenome shows upregulation of meiosis-related genes, possibly in order to prevent aneuploidy and undesirable homeologous exchanges that are frequently observed in experimentally generated *A. suecica*, and the *A. arenosa* subgenome shows upregulation of cyto-nuclear related processes, possibly in response to the new cytoplasmic environment of *A. suecica,* with plastids maternally inherited from *A. thaliana*.

Introduction

Ancient polyploidization or whole-genome duplication is a hallmark of most higher-organism genomes[1,2], including our own[3,4]. While most of these organisms are now diploid and show only traces of polyploidy, there are many examples of recent polyploidization, in particular among flowering plants[5–9]. These examples are important because they allow us to study the process of polyploidization, rather than just inferring that it happened and trying to understand its evolutionary importance.

The existence of wide-spread recent polyploids, like *Capsella bursa-pastoris* (Shepherd's Purse)[10–12], *Trifolium repens* (white clover)[13], and *Brachypodium hybridum*[14,15], demonstrates that polyploid species can quickly become very successful. That said, new polyploid species face numerous challenges, ranging from those on a population level, such as bottlenecks[13,16] and competition with their diploid progenitors[17], to those on a cellular level, such as chromosome segregation[18–20] and changes to genome structure and regulation[21] — the so-called "genome shock"[22]. However, genomic changes resulting from the hybridization of two (or more) diverged genomes have primarily been studied in crops, such as cotton[5], strawberry[6], wheat[7] and peanut[8,9], which confounds polyploidization with domestication.

Here we focus on a young, natural allopolyploid species, *Arabidopsis suecica* (2n = 4x = 26), formed through the hybridization of *A. thaliana* (2n = 10) and *A. arenosa* (2n = 2x/4x = 16/32), circa 16 kya, during the Last Glacial Maximum[16] and now widely established in northern Fennoscandia (Fig. 1a). The ancestral species diverged around 6 Mya[23], and, based on mitochondrial and chloroplast sequences, it is clear that *A. thaliana* is the maternal and *A. arenosa* the paternal parent of the hybrid[24], a scenario also supported by the fact that *A. arenosa* itself is a ploidy-variable species, so that *A. suecica* could readily be generated through the fertilization of an unreduced egg cell (2n = 2x) from *A. thaliana* by a sperm cell (n = 2x) from autotetraploid *A. arenosa*[16,25]. We have previously shown that, although *A. suecica* shows clear evidence of a genetic bottleneck[16], it shares most of its variation with the ancestral species, demonstrating that the species was formed through a hybridization and polyploidization process that involved many crosses and individuals.In order to study genomic change in *A. suecica*, we used long-read sequencing to generate a high-quality, chromosome-level assembly of a single individual, taking advantage of the fact that *A. suecica*, like *A. thaliana*, is highly selfing, making it possible to sequence naturally inbred individuals. The genome sequence was complemented by a partial assembly of a tetraploid outcrosser *A. arenosa*, and by short-read genome and transcriptome sequencing data from many individuals of all three species — including "synthetic" *A. suecica* generated *de novo* in laboratory crosses.

Results and discussion

1. The genome is conserved

We assembled a reference genome from a naturally inbred *A. suecica* accession ("ASS3"), using 50x long-read PacBio sequencing (PacBio RS II). The absence of heterozygosity and the substantial (~11,6%) divergence between the subgenomes greatly facilitated the assembly. In contrast, assembling even a diploid *A. arenosa* genome is complicated by high heterozygosity (nucleotide diversity around 3.5%[26]) coupled with a relatively high level of repetitive sequences (compared to the gene-rich *A. thaliana* genome). Our attempt to assemble a tetraploid *A. arenosa*, which is also included here, led to a very fragmented assembly of 3,629 contigs with an N50 of 331 Kb. In contrast, the *A. suecica* assembly has an N50 contig size of 9.02 Mb. The assembled contigs totaled 276 Mb (~90% of the 305 Mb genome size estimated by flow cytometry — see Supplementary Fig. 1; ~88% of the 312Mb genome size estimated by kmer analysis). Contigs were placed into scaffolds using high-coverage chromosome conformation capture (HiC) data and by using the reference genomes of *A. thaliana* and *A. lyrata* (here the closest substitute for *A. arenosa*) as guides. This resulted in 13 chromosome-scale scaffolds (Supplementary Fig. 2a). The placement and orientation of each contig within a scaffold was confirmed and corrected using a genetic map for *A. suecica* (see Methods, Supplementary Fig. 3, Supplementary Fig. 4). The resulting chromosome-level assembly (Fig. 1b) contains 258 Mb, and has an N50 scaffold size of 19.59 Mb. The five chromosomes of the *A. thaliana* subgenome and the eight chromosomes of the *A. arenosa* subgenome sum to 119 Mb and 139 Mb, respectively.

101
Figure 1. The genome of *A. suecica* is largely colinear with the ancestral genomes. **a** Schematic
depicting the origin of *A. suecica* and its current distribution in the relation to the last glacial maximum
(LGM). **b** The chromosome-level assembly of the *A. suecica* genome with inner links depicting syntenic
blocks between the *A. thaliana* and *A. arenosa* subgenomes of *A. suecica*. The blue histogram
represents the distribution of TEs along the genome and the green histogram corresponds to the
distribution of protein-coding genes. **c** Synteny of the *A. thaliana* subgenome of *A. suecica* to the *A.*
thaliana TAIR10 reference. In total 13 colinear synteny blocks were found. **d** Synteny of the *A. arenosa*
subgenome to *A. lyrata*. In total 37 synteny blocks were found, 28 of which were colinear. Of the
remaining 9 blocks, 7 represent inversions in the *A. arenosa* subgenome of *A. suecica* compared to *A.*
lyrata, 1 is a translocation, and one corresponds to a previously reported mis-assembly in the *A. lyrata*
genome[27]. Orange bars represent a density plot of missing regions ("N" bases) in the *A. lyrata* genome.
113
114

Approximately 108 and 131 Mb of the *A. thaliana* and *A. arenosa* subgenomes of *A. suecica* are in large syntenic blocks to the genomes of the ancestral species: 13 and 37 blocks, respectively (Fig. 1c,d). The vast majority of these syntenic blocks are themselves also colinear, with the exception of five small-scale inversions (~3.7 Mb) and one translocation (~1Mb) on the *A. arenosa* subgenome— which may well (indeed probably do) reflect differences between *A. lyrata* and *A. arenosa*, two species separated by about a million years[23,26]. We also corrected for the described[27] mis-assembly in the *A. lyrata* reference genome using our genetic map. Overall we find that approximately 92% of the *A. suecica* genome is syntenic to the ancestral genomes, the 13 chromosomes of *A. suecica* having remained almost completely colinear (Fig. 1c,d). This highlights the conservation of the *A. suecica* genome and contrasts with the major rearrangements that have been observed in several polyploid crops[8,9,28–30] and which have often been associated with a "genome shock". Interestingly, major rearrangements have also been observed in synthetic *A. suecica*[31], and we see clear evidence of aneuploidy in ours — a topic to which we shall return below.

A total of 45,048 protein-coding genes were annotated for the *A. suecica* reference, of which 22,383 and 22,665 are located on the *A. thaliana* and *A. arenosa* subgenomes, respectively. We assessed completeness of the genome assembly and annotation with the BUSCO set for eudicots and found 2036 (96%) and 2007 (94.6%) complete genes for the *A. thaliana* and *A. arenosa* subgenomes, respectively (Supplementary Fig. 5c,d). Of the protein-coding genes, 17,814 had a one-to-one orthology between the subgenomes of *A. suecica* and 16,781 genes were conserved in a 1:1:1:1 relationship for both subgenomes of *A. suecica* and the ancestral species (using *A. lyrata* as a substitute for *A. arenosa*) (Supplementary Data 2, Supplementary Fig. 5b). We functionally annotated lineage-specific genes in *A. suecica* using InterPro, and only found significant enrichment in *A. thaliana* subgenome of *A. suecica* for two GO terms (GO:0008234 and GO:0015074), both of which are associated with repeat content (Supplementary Data 2). Ancestral genes not found in the *A. suecica* genome annotation were overrepresented for functional categories of plant defense response. However, checking coverage for these genes by mapping the raw *A. suecica* whole-genome resequencing data to the ancestral genomes did not confirm their loss, suggesting rather misassembly or misannotation, which is expected due to the repetitive and highly polymorphic nature of R-genes in plants.

2. The rDNA clusters are highly variable

In eukaryotic genomes, genes encoding ribosomal RNA (rRNA) occur as tandem arrays in rDNA clusters. The 45S rDNA clusters are particularly large, containing hundreds or thousands of copies, spanning millions of base pairs[32]. The nucleolus, the site of pre-ribosome assembly, forms at these clusters, but only if they are actively transcribed, and it was observed long ago that only one parent's rDNA tended to be involved in nucleolus formation in inter-specific hybrids, a phenomenon known as "nucleolar dominance"[33,34]. In *A. suecica,* it was observed that the rDNA clusters inherited from *A. thaliana* were silenced, and structural changes associated with these clusters were also suggested[35].

Given this, we examined the composition of 45S rDNA repeats as well as their transcription. While the large and highly repetitive 45S rDNA clusters are not part of the genome assembly, it is possible to measure the copy number of *A. thaliana* and *A. arenosa* 45S rRNA genes using sequencing coverage (see Methods), and we find three accessions to have experienced massive loss of the *A. thaliana* rDNA loci (Fig. 2a), which we confirmed for

one of the accessions ("AS90a") by FISH analysis (Fig. 2d,e). However, there is massive copy
number variation for 45S rRNA genes in *A. suecica* (Fig. 2a), and some accessions (e.g., the
reference accession "ASS3") have higher *A. thaliana* than *A. arenosa* 45S rRNA copy number
(Fig. 2b,c).
Turning to expression, we also find nucleolar dominance to be segregating in *A. suecica*
(see Methods and Supplementary Fig. 6), with the majority of accessions expressing both 45S
rRNA alleles, five exclusively expressing *A. arenosa* 45S rRNA, and one exclusively
expressing *A. thaliana* 45S rRNA (Fig. 2a).
This extensive variation in 45S cluster size and expression is reminiscent of the
intraspecific variation seen in *A. thaliana* (where different accessions express either the
chromosome 2 or chromosome 4 rDNA cluster, or both[36,37]), and suggests that the
phenomenon of nucleolar dominance can at least partly be explained by retained ancestral
variation. However, the large-scale decrease in rDNA cluster size observed in some
accessions may be a direct consequence of allopolyploidization itself, as synthetic *A. suecica*
sometimes shows immediate loss of 45S rDNA (even as early as the F1 stage) and this too
varies between siblings and generations (Supplementary Fig. 6a).

Figure 2. Expression and copy number variation of 45S rDNA in *A. suecica*. a The relationship
between expression levels (log$_2$ CPM) and copy number of 45S rRNA shows extensive variation of 45S
rDNA copy number and varying direction of "nucleolar dominance". Grey lines connect subgenomes of
the same accession. Values above the dashed line are taken as evidence for the expression of a
particular 45S rDNA allele, as this is above the maximum level of mis-mapping seen in the ancestral
species here used as a control (see Supplementary Figure 6b). **b** and **c** FISH results of a natural *A.*
suecica accession "ASS3" that has maintained both ancestral rDNA loci (174 copies calculated for the
A. thaliana 45S rDNA and 104 copies of the *A. arenosa* 45S rDNA). **d** and **e** FISH result of a natural
accession "AS90a" that has largely lost the rDNA cluster of the *A.thaliana* subgenome (8 copies
calculated for the *A. thaliana* 45S rDNA and 159 copies of the *A. arenosa* 45S rDNA).

3. No evidence for abnormal transposon activity

The possibility that hybridization and polyploidization leads to a "genome shock" in the form of increased transposon activity has been much discussed [21,22,38,39]. Certainly the two subgenomes of *A. suecica* differ massively in transposon content: there are almost twice as many annotated transposons in the *A. arenosa* as in the *A. thaliana* subgenome (64,409 vs 33,420; see Supplementary Figs. 5a and 7), and the true difference is almost certainly greater given that the *A. arenosa* subgenome assembly is less complete (and many of the missing regions are likely to be repeat-rich) and that the transposon annotation is biased towards *A. thaliana*. Has the combination of two such different genomes lead to increased transposon activity?

Our assembled *A. thaliana* subgenome does contain roughly 3,000 more annotated transposons than the TAIR10 *A. thaliana* reference genome, but this could reflect greater transposon number in the *A. thaliana* ancestors of this genome rather than increased transposon activity in *A. suecica.* In order to gain insight into transposon activity in *A. suecica*, we need to identify jumps that occurred after the species separated (and are thus only found in this species). We used the software PopoolationTE2[40] to call presence-absence variation on a population-scale level using genome re-sequencing datasets for 15 natural *A. suecica* accessions, 18 *A. thaliana* accessions genetically close to *A. suecica*, and 9 *A. arenosa* lines. Of the 25,677 insertion polymorphisms called with respect to the *A. thaliana* subgenome, 9,354 were shared between *A. thaliana* and *A. suecica*, 8,068 were only found in *A. thaliana*, and 8,255 were only found in *A. suecica*. Of the 100,849 insertions on the *A. arenosa* subgenome of *A. suecica,* 11,024 were shared with *A. arenosa*, 69,041 were private to *A. arenosa*, and 20,784 were private to *A. suecica* (Supplementary Data 1a,b; Supplementary Figs. 8,9). Considering the number of transposons per individual genome (Fig 4a), we see that most transposon insertions in a typical *A. thaliana* subgenome are also found in *A. thaliana*, and that the slightly higher transposon load in the *A. thaliana* subgenome is mainly due to these. The reason for this is likely a population bottleneck, as we shall see below. In contrast, the number of recent insertions (that are private to the species) is not higher in the *A. thaliana* subgenome, suggesting that transposon activity in this subgenome is not increased.

Turning to the *A. arenosa* subgenome, we see that a typical *A. suecica* contains only about half the number of transposons of a typical *A. arenosa* individual (Fig 4a). However, the latter is an outcrossing tetraploid, and it is thus fairer to compare with the number of transposons in four randomly chosen *A. arenosa* subgenomes. This largely accounts for the observed difference, but there are still clearly fewer transposons in *A. suecica*. A population bottleneck (see below) likely explains much of this, but it is impossible to rule out a contribution of decreased transposon activity in *A. suecica* as well.

To sum up, we see no evidence for a burst of transposon activity accompanying polyploidization in *A. suecica*, a conclusion also supported by a lack of increase in transposon expression for both synthetic and natural *A. suecica* compared to the *A. thaliana* and *A. arenosa* on both subgenomes (Supplementary Fig. 9). We do see clear traces of the population bottleneck accompanying the origin of *A. suecica*, however. The frequency distribution of polymorphic transposon insertions private in *A. suecica* is skewed towards zero — almost certainly because of purifying selection because the distribution is more similar to that of non-synonymous SNPs than to that of synonymous SNPs (Fig 4b and c). However, for both subgenomes, *A. suecica* also contains a large number of fixed or nearly-fixed insertions that are present in the ancestral species at lower frequency (Fig 4d and e). These are likely to have reached high-frequency as a result of a bottleneck. Shared transposons are

enriched in the pericentromeric regions of the genome depleted of protein-coding genes, while private transposons insertions, which are generally at low frequency, show a more uniform distribution across the genome, consistent with evidence for stronger selection against transposon insertion in the relatively gene-dense chromosome arms[41,42] (Supplementary Fig. 10).

Figure 3. TE dynamics in *A. suecica* reveal no evidence for abnormal transposon activity. a Median TE insertions per genome. As the *A. arenosa* population is an autotetraploid outcrosser, 4 randomly chosen haploid *A. arenosa* subgenomes of *A. suecica* were combined to make a 4n *A. suecica*. *A. suecica* does not show an increase in private TE insertions compared with the ancestral species for both subgenomes, and shared TEs constitute a higher fraction of TEs in *A. suecica* reflecting the strong population bottleneck at its origin. Site-frequency spectra of non-synonymous SNPs, synonymous SNPs and TEs in the **b** *A. thaliana* and **c** *A. arenosa* subgenomes of *A. suecica* suggest that TEs are under purifying selection on both subgenomes. **d** 3D histogram of a joint TE frequency spectrum for *A. thaliana* on the x-axis and the *A. thaliana* subgenome of *A. suecica* on the y-axis **e** 3D histogram of a joint TE frequency spectrum for *A. arenosa* on the x-axis and the *A. arenosa* subgenome of *A. suecica* on the y-axis. **d** and **e** show stable dynamics of private TEs in *A. suecica* and a bottleneck effect on the ancestral TEs (shared) at the origin of the *A. suecica* species.

An interesting subset of recent transposon insertions private to *A. suecica* are those that have jumped between the two subgenomes. We searched for full-length transposon copies that are present in both subgenomes of *A. suecica* and then assigned the resulting consensus sequences to either the *A. thaliana* or the *A. arenosa* ancestral genomes using BLAST (see Methods). We were able to assign 15 and 55 consensus sequences as being specific to the *A. thaliana* and *A. arenosa* ancestral genomes, respectively. Using these sequences, we searched our transposon polymorphism data for corresponding polymorphisms, and identified 1,823 *A. arenosa* transposon polymorphisms on the *A. thaliana* subgenome, and 461 *A.*

thaliana transposon polymorphisms on the *A. arenosa* subgenome. Like other private polymorphisms, these are skewed towards rare frequencies, and are uniformly distributed across the (sub-)genome. Most of the transposons that have jumped into the *A. thaliana* subgenome are helitrons and LTR elements (Supplementary Figure 12). LTR (copia) elements also make up most of the *A. thaliana* transposons segregating in the *A. arenosa* subgenome, with the most abundant element having a closest match in the *A. thaliana* mitochondrial genome. The fact that nearly four times as many new insertions appear to have resulted from jumps from *A. arenosa* to *A. thaliana* than the other way around is notable. It could reflect higher transposon activity in the *A. arenosa* subgenome, but can mostly be explained by differences in genome size and transposon number. If there were no differences in activity, we would expect the number of cross-subgenome jumps to be proportional to the number of potential source elements and the size of the target genome. As we have seen, the *A. arenosa* subgenome contains roughly twice as many transposons as the *A. thaliana* subgenome, and is about 17% larger. These factors alone would account for a 3.4-fold difference in cross-subgenome jumping.

In conclusion, transposon activity in *A. suecica* appears to be governed largely by the same processes that governed it in the ancestral species.

4. No bias in expression between subgenomes

Over time the traces of polyploidy are erased through an evolutionary process involving gene loss, often referred to as fractionation or re-diploidization[43,44][45–47]. Analyses of retained homeologs in ancient allopolyploids such as *A. thaliana*[48], *Zea mays*[49], *Brassica rapa*[50] and *Gossypium raimondii*[51] have revealed that one "dominant" subgenome remains more intact, with more highly expressed homeologs compared to the "submissive" genome(s)[48]. This pattern of "biased fractionation" has not been observed in ancient autopolyploids[52], such as pear[53], and is believed to be allopolyploid-specific. Subgenome dominance in expression has been reported for a number of more recent allopolyploids such as strawberry[6] and both natural and synthetic allopolyploids such as *Brassica napus*[54], and monkeyflower[55]. However, some allopolyploids display even subgenome expression, such as *C. bursa-pastoris*[10,12] and *B. hybridum*[14].

Subgenome dominance is often linked to differences in transposon content[6] and/or large genetic differences between subgenomes[56]. This makes *A. suecica*, with 6 Mya divergence between the gene-dense *A. thaliana* and the transposon-rich *A. arenosa*, a promising candidate to study this phenomenon at unprecedented resolution. Previous reports on subgenome dominance in *A. suecica* are conflicting, suggesting a bias to either the *A. thaliana*[57] or the *A. arenosa*[58] subgenome.

To investigate the evolution of gene expression in *A. suecica*, we generated RNA-seq data for 15 natural *A. suecica* accessions, 15 closely related *A. thaliana* accessions, 4 *A. arenosa* individuals, a synthetically generated *A. suecica* from a lab cross (the 2nd and 3rd hybrid generations) and the parental lines of this cross. Each sample had 2-3 biological replicates (Supplementary Data 2). On average, we obtained 10.6 million raw reads per replicate, of which 7.6 million reads were uniquely mapped to the *A. suecica* reference genome and 13,647 homeologous gene pairs (see Methods, Supplementary Fig. 13).

Considering the difference in expression between homeologous genes, we found no bias towards one or the other subgenome of *A. suecica*, for any sample or tissue, including synthetic *A. suecica* (Fig. 4a and Supplementary Fig. 14a). This strongly suggests that the

expression differences between the subgenomes have not changed systematically through
polyploidization, and is in contrast to previous studies, which reported a bias towards the *A.*
thaliana[57] or the *A. arenosa*[58] subgenome, likely because RNA-seq reads were not mapped
to an appropriate reference genome.
The set of genes that show large expression differences between the subgenomes
appears not to be biased towards any particular gene ontology (GO) category, and is
furthermore not consistent between accessions and individuals (Fig. 4b, Supplementary Fig.
14b,c). This suggests that many large subgenome expression differences are due to genetic
polymorphisms within *A. suecica* rather than fixed differences relative to the ancestral species.
Levels of expression dominance were reported to vary across tissues in *C. bursa*
pastoris[11]. To test whether expression dominance can vary for tissue-specific genes, we
examined homeologous gene-pairs where at least one gene in the gene pair showed tissue
specific expression, in whole-rosettes and floral buds. We do not find evidence for dominance
between subgenomes in tissue specific expression either (Figure 4b). Interestingly, the 402
genes with significant expression in whole rosettes for both homeologs showed GO
overrepresentation that included both photosynthesis and chloroplast related functions
(Supplementary Table 1). This result suggests that the *A. arenosa* subgenome has
established important cyto-nuclear communication with the chloroplast inherited from *A.*
thaliana, rather than being silenced. 1009 genes with floral bud specific expression for both
homeologous gene pairs were overrepresented for GO terms related to responses to chemical
stimuli, such as auxin and jasmonic acid, which may reflect early developmental changes in
this young tissue (Supplementary Table 1).
In summary, we find no evidence that one subgenome is dominant and contributes more
to the functioning of *A. suecica*. On the contrary, homeologous gene pairs are strongly
correlated in expression across tissues.
332
333
334

Figure 4. Patterns of gene expression between the subgenomes of *A. suecica* in rosettes and floral buds. **a** Violin plots of the mean log fold-change between the subgenomes for the 15 natural *A. suecica* accessions and two synthetic lines for whole rosettes. Mean log fold-change for the two accessions ("ASS3" and "AS530") where transcriptome data for both whole rosettes and flower buds were available. All the distributions are centered around zero suggesting even subgenome expression. **b** Violin plots for the mean log fold-change between the subgenomes for genes with tissue-specific expression. At least one gene in a homeologous gene pair was required to show tissue-specific expression.

5. Evolving gene expression in *A. suecica*

The previous section focused on differences in expression between the subgenomes, between homeologous copies of the same gene within the same individual. This section will focus on differences between individuals, between homologous copies of genes that are part of the same (sub-)genome. To provide an overview of expression differences between individuals we performed a principal component analysis (PCA) on gene expression separately for each (sub-)genome. For both subgenomes, the first principal component separates *A. suecica* from the ancestral species and the synthetic hybrid (Fig. 5a,b), suggesting that hybridization does not automatically result in large-scale transcriptional changes, and that altered gene expression changes in natural *A. suecica* have evolved over time. Given the limited time involved, the lack of correlation between expression and sequence divergence within and between species (Supplementary Fig. 15), and the fact the genes that have changed expression are far from random with respect to function (see below), we suggest that the first principal component primarily captures trans-regulated expression changes in *A. suecica* that are likely adaptive.

Figure 5. Differential gene expression analysis in *A. suecica*. Patterns of differential gene expression in *A. suecica* support adaptation to the whole-genome duplication for the *A. thaliana* subgenome and adaptation to the new plastid environment for the *A. arenosa* subgenome. **a** PCA for *A. thaliana*, *A. thaliana* subgenome of natural and synthetic *A. suecica* lines. PC1 separates natural *A. suecica* from the ancestral species and the synthetic lines. **b** PCA for *A. arenosa*, *A. arenosa* subgenome of natural and synthetic *A. suecica* lines. PC1 separates natural *A. suecica* from the ancestral species and the synthetic lines, whereas PC2 identifies outlier accessions discussed further below (see Fig.6). **c, d** Heatmap of DEGs for the two subgenomes of *A. suecica*. Positive numbers (red color) indicate higher expression. Genes and individuals have been clustered based on similarity in expression, resulting in clusters discussed in text. **e** Gene ontology enrichment for each cluster in **c** and **d**. Categories discussed in the text are highlighted.

To further characterize expression changes in natural *A. suecica* we analyzed differentially
expressed genes (DEGs) on both subgenomes compared to the corresponding ancestral
species. The total number of DEGs was 3,854 and 4,136 genes for the *A. thaliana* and *A.*
arenosa subgenomes, respectively (see Methods). These genes were clustered based on the
pattern of change across individuals (Fig. 5c,d) and GO enrichment analysis was carried out
for each cluster (Fig. 5e, Supplementary Table 2).
For the *A. thaliana* subgenome, we identified three clusters. Cluster 1 comprised 2,041
genes that showed decreased expression in *A. suecica* compared to *A. thaliana*. These genes
are strongly enriched for transcriptional regulation, which may be expected as we are
examining DEGs between the species. Also notable are enrichments for circadian rhythm
function, and response to nitrate, both of which may be related to the ecology of *A. suecica*
and its post-glacial migration to the Fennoscandinavia region (Fig. 1a).
Cluster 2 consisted of 1,376 genes that show increased expression in *A. suecica*
compared to *A. thaliana*, and several of the enriched GO categories, such as microtubule-
based movement, cytokinesis, meiosis and cell division, suggest that the *A. thaliana*
subgenome of *A. suecica* is adapting to polyploidy at the level of basic cell biology. That there
has been strong selection for this seems likely given that aneuploidy is frequent in synthetic
A. suecica (Supplementary Fig. 16), while natural *A. suecica* has a stable and conserved
karyotype. Importantly, there is independent evidence for adaptation to polyploidy via
modifications of the meiotic machinery in the other ancestor of *A. suecica*, *A. arenosa*, as
well[19,59,60], although we see very little overlap in the genes involved (Supplementary Fig. 16).
The nature of these changes in the *A. suecica A. thaliana* subgenome will require further
investigation, but we note that there is enrichment (see Methods, Supplementary Data 3) for
Myb family transcription factor binding sites[61] among upregulated genes in cluster 2. There is
also enrichment for "response to cold" genes in cluster 2, a category which was also over-
expressed in rosettes relative to floral buds (Supplementary Table 1), and which could again
be related to the post-glacial migration of *A. suecica*.
Cluster 3, finally, consists of 437 genes that are over-expressed in both natural and
synthetic *A. suecica* relative to *A. thaliana*. These expression changes are thus likely an
immediate consequence of hybridization and must reflect trans-regulation. Genes in this
cluster are enriched for "response to cadmium", "double-strand break repair" and "pre-
replicative complex assembly". *A. arenosa* is known to be somewhat metal-tolerant[62], and it is
possible that the *A. arenosa* subgenome simply upregulates related genes on the *A. thaliana*
subgenome. Upregulation of double-strand break repair and pre-replicative complex assembly
could be an immediate response to genome-doubling, given the degree of aneuploidy we
detect in the synthetic *A. suecica* lines (Supplementary Fig. 16). Notably, the synthetic lines
used in the expression analysis were selected to be healthy-looking, and did not show signs
of aneuploidy (Supplementary Fig. 17).
For the *A. arenosa* subgenome, we also find three clusters of DEGs (Fig. 5d) with GO
enrichment for two (Fig. 5e, Supplementary Table 2). Cluster 1 consists of 1,165 genes that
show increased expression in natural *A. suecica* compared to *A. arenosa* and synthetic *A.*
suecica, and are enriched for plastid-related functions, including oxidation-reduction, the
reductive pentose-phosphate pathway and photosynthesis. We attribute this to selection on
the *A. arenosa* subgenome to restore communication with the new plastid environment as
plastid genomes were maternally inherited from *A. thaliana*. Cluster 2 consists of 2,849 genes
that show decreased gene expression in *A. suecica* compared to *A. arenosa* and synthetic *A.*
suecica. These genes were primarily enriched for mRNA processing and cell division. Cluster

3 (122 genes) did not have a GO overrepresentation and showed an intriguing pattern
discussed in the next section.

6. Homeologous exchange contributes to variation in gene expression

The second principal component for gene expression identified three outlier-accessions of *A.*
suecica, two for the *A. thaliana* subgenome (Fig. 5a) and one for the *A. arenosa* subgenome
(Fig. 5b). While closely examining the latter accession, "AS530", we realized that it is
responsible for the cluster of genes with distinct expression patterns but no GO enrichment
just mentioned (Fig. 5d, Cluster 3). Genes from this cluster were significantly downregulated
on the *A. arenosa* subgenome (Fig. 6a) and upregulated on the *A. thaliana* subgenome (Fig.
6b) — for AS530 only. The further observation that 97 of the 122 genes (Supplementary Figure
20a) in the cluster are located in close proximity in the genome, pointed to a structural
rearrangement. The lack of DNA sequencing coverage on the *A. arenosa* subgenome around
these 97 genes and the doubled coverage for their homeologs on the *A. thaliana* subgenome,
strongly suggested a homeologous exchange (HE) event resulting in AS530 carrying four
copies of the *A. thaliana* subgenome and zero copies of the *A. arenosa* genome with respect
to this this, roughly 2.5 Mb region of the genome (Fig. 6c). This explanation was further
supported by HiC data, which showed clear evidence for interchromosomal contacts between
A. thaliana subgenome chromosome 1 and *A. arenosa* subgenome chromosome 6 around
the breakpoints of the putative HE in AS530 (Fig. 6 d,e), and by multiple discordant Illumina
paired-end reads at the breakpoints between the homeologous chromosomes, which
independently support the HE event (Supplementary Fig. 19a-d).
Based on this we examined the two outlier *A. suecica* accessions for the *A. thaliana*
subgenome (Fig. 5a; "AS150" and "ASÖ5"), and found that they likely share a single HE event
in the opposite direction (four copies of the *A. arenosa* subgenome and no copies of the *A.*
thaliana subgenome for a region of roughly 1Mb in size, see Supplementary Figure 18). This
demonstrates that HE occurs in *A. suecica* and contributes to the intraspecific variation we
observed in gene expression (Fig 5a, b). However, the majority of HEs are probably
deleterious as they will lead to gene loss: although the *A. thaliana* and *A. arenosa* genomes
are largely syntenic, AS530 is missing 108 genes (Supplementary Figure 19) that are only
present on the *A. arenosa* subgenome segment that has been replaced by the homeologous
segment from the *A. thaliana* subgenome, and AS150/ASÖ5 are missing 53 genes that were
only present on the *A. thaliana* subgenome.

Figure 6. Homeologous exchange contributes to expression variance within *A. suecica*. a Cluster 3 of Fig. 5d explains the outlier accession AS530 which is not expressing a cluster of genes on the *A. arenosa* subgenome. **b** Homeologous genes of this cluster on the *A. thaliana* subgenome of *A. suecica* show the opposite pattern and are more highly expressed in AS530 compared to the rest of the population. **c** 97 of the 122 genes from cluster 3 are located in close proximity to each other on the reference genome but appear to be deleted in AS530 based on sequencing coverage. **d** The *A. thaliana* subgenome homeologs have twice the DNA coverage, suggesting they are duplicated. **e** HiC data show (spurious) interchromosomal contacts at 25 Kb resolution between chromosome 1 and chromosome 6 around the breakpoint of the cluster of 97 genes in AS530 but not in reference accession ASS3.

Conclusion

This study has focused on the process of polyploidization in a young allotetraploid species,
A. suecica, that was generated roughly 16 kya through the hybridization of two species, *A.*
thaliana and *A. arenosa*, which differ substantially in everything from genome size and
chromosome number to mating system and ecology. Our study is one of relatively few to
focus on a natural polyploid rather than a domesticated species, and is unparalleled in its
resolution thanks to one of the parents being a major model species.

The main conclusion of our study is that polyploid speciation, at least in this case, was
clearly a gradual process rather than some kind of "event". We confirmed previous results
that genetic polymorphism is largely shared with the ancestral species, demonstrating that *A.*
suecica did not originate through a single unique hybridization event, but rather through
multiple crosses[16]. We also find no evidence for the kind of dramatic "genome shock" that
has often been suggested to accompany polyploidization and hybridization. The genome has
not been massively rearranged, transposable elements are not out of control, and there is no
subgenome dominance in expression. On the contrary, we find evidence of genetic
adaptation to "stable" life as a polyploid, in particular changes to the meiotic machinery and
in interactions with the plastids. These findings, coupled with the observation that
experimentally generated *A. suecica* are often unviable and do exhibit evidence of genome
rearrangements, suggest that the most important bottleneck in polyploid speciation may be
selective, and that domesticated polyploids may not be representative of natural
polyploidization. Darwin famously argued that "Natura non facit saltum"[63] — we suggest that
polyploidization is no exception from this.

Supplementary figures

Supplementary Figure 1. Measuring genome sizes of *Arabidopsis* species using flow cytometry. a FACs sorting of *Solanum lycopersicum* cells from 3 week old leaf tissue for two replicates. G1 represents the peak denoting the G1 phase of the cell cycle. Cells in the G1 phase have 2C DNA content (i.e. a 2N genome). **b** *A. thaliana* "CVI" accession **c** *A. lyrata* "MN47" (the reference accession) **d** *A. suecica* "ASS3" (the reference accession) **e** autopolyploid *A. arenosa* accession "Aa4" **f** Bar chart shows calculated genome sizes (rounded to the nearest whole number) for each species using *Solanum lycopersicum* as the standard .

Supplementary Figure 2. HiC as a tool to investigate structural rearrangements. **a** HiC contact map for the full chromosome-level genome assembly of *A. suecica*. **b** Mixing of *A. thaliana* and *A. arenosa* HiC reads suggest interchromosomal contacts between homeologous chromosomes is a result of mis-mapping for HiC reads. Such mis-mapping is typically filtered out in short read DNA and RNA datasets using insert size and proper pairs mapping filters, however in HiC long range chromosomal contacts are not filtered out. **c** Accession "AS530" with the region of homeologous exchange highlighted with an arrow (figure 6), no other rearrangements were observed. **d** HiC of synthetic *A. suecica* (F3).

Supplementary Figure 3. Crossover counts in an *A. suecica* F2 population. Per chromosome crossover counts in our F2 population (N=185). Chromosome 2 had too few SNPs to be analysed in our cross due to the recent bottleneck in *A. suecica*[16].

529 **Supplementary Figure 4. A genetic map for *A. suecica*.** Physical distance (Mb) vs genetic
530 distance (cM) is plotted for each: **a** *A. thaliana* subgenome and; **b** *A. arenosa* subgenome
531 chromosome. Chromosome 2 is not plotted as there are too few SNPs on this chromosome in
532 our cross, due to the recent bottleneck in *A. suecica*[16]

Supplementary Figure 5. Genome composition and orthologous gene relationships in *A. suecica*. a Genome composition of the *A. suecica* subgenomes and the ancestral genomes of *A. thaliana* and *A.lyrata* (here a substitute reference for *A. arenosa* because it is annotated). Ler = Landsberg *Erecta* (*A. thaliana*)[64]. The difference in CDS between the Tair10 annotation with that of and Ler and the *A. thaliana* subgenome is most likely due to the manual curation of Tair10 and the de novo gene prediction pipeline used in this paper. **b** Counts of orthologous relationships between the subgenomes of the reference *A. suecica* genome and the reference *A. thaliana* and *A. lyrata* genome. Ancestrally segregating genes are genes shared between the *A. thaliana* reference and the *A. arenosa* subgenome or shared between the *A. lyrata* reference and the *A. thaliana* subgenome. Therefore they most likely represent genes ancestrally segregating in the ancestor of *A. thaliana* and *A. lyrata*. BUSCO analysis of *A. suecica* using the BUSCO set for eudicots for the **c** *A. thaliana* and **d** *A. arenosa* subgenome.

556
557
Supplementary figure 6. **rDNA copy number variation and expression. a** Copy number of
A. thaliana and *A. arenosa* rDNA in natural *A. suecica*, ancestral species and synthetic lines.
Blue triangles represent the *A. thaliana* and *A.arenosa* parent lines of the synthetic *A. suecica*
cross. AT represents results when mapping to the *A. thaliana* consensus sequence and AA to
the *A. arenosa* consensus sequences for the 45S rRNA **b** Expression (log2 CPM) of *A.*
thaliana and *A. arenosa* rDNA in natural *A. suecica*, ancestral species and synthetic lines.
Accessions with log2 CPM of >=15 was taken as evidence for expression for the *A. thaliana*
and *A. arenosa* 45S rRNA in *A. suecica*, as this CPM value was above the maximum level of
mis-mapping observed in the ancestral species (*A. thaliana* mapping to the *A. arenosa* 45S
rRNA).

568
569 **Supplementary Figure 7. TE-composition of the *A. suecica* reference genome.** TE
570 composition of the **a** *A. thaliana* and **b** *A. arenosa* subgenome of *A. suecica*.
571
572
573

574
575
576 **Supplementary Figure 8. Site frequency spectrum (SFS) of shared TEs and private TEs**
577 **in _A. suecica_ broken down by TE family.** Shared TE SFS for the **a** _A. thaliana_ and **b** _A._
578 _arenosa_ subgenome. Private TE SFS for the **c** _A. thaliana_ and **d** _A. arenosa_ subgenome.
579

Supplementary Figure 9. Analysis of TE expression in *A. suecica*. Patterns of TE expression in natural and synthetic *A. suecica* show that allopolyploidy is not accompanied by an overall up-regulation in TE expression as predicted by the "genome shock" hypothesis. **a** Heatmap of TE expression for the *A. thaliana* subgenome of *A. suecica* (dark green) synthetic *A. suecica* (cyan) and *A. thaliana* (light green). **b** Heatmap of TE expression for the *A. arenosa* subgenome of *A. suecica* (dark purple) synthetic *A. suecica* (pink) and *A. arenosa* (light purple). **c** and **d** the breakdown of TE families expressed in each cluster, with helitrons being the most abundant class on the *A. thaliana* subgenome and TEs of an unknown family being the most abundant in the *A. arenosa* subgenome.

Supplementary Figure 10. Genomic distribution of TEs in the *A. suecica* genome. a
Shared TEs in the population between *A. thaliana* and the *A. thaliana* subgenome of *A.*
suecica. Shared TEs are likely older than private TEs and are enriched around the
pericentromeric regions in the *A. thaliana* subgenome. Private TEs are enriched in the
chromosomal arms for both species, where protein coding gene density is higher (Fig. 1b). **b**
as in **a** but examining TEs in the population of *A. arenosa* and the *A. arenosa* part of *A. suecica*.
Note the region between 5 and 10 on chromosome 2 was not included in the analysis as this
region shows synteny with an unplaced contig.

Supplementary Fig 11. Patterns of selection in *A. suecica*. a Comparison of shared variation (Nonsense SNPs, synonymous SNPs, and TEs) population frequencies in the *A. thaliana* subgenome of 15 natural *A. suecica* accessions and the closest 31 *A. thaliana* accessions. **b** Comparison of shared variation (Nonsense SNPs, synonymous SNPs, and TEs) frequencies in *A. arenosa* subgenome of 15 *A. suecica* accessions and 11 Swedish *A. arenosa* lines. Although results may be affected by the sampling and potential misidentification of the ancestral populations, the current data suggests strong purifying selection on the *A. thaliana* subgenome of *A. suecica* for shared nonsense SNPs, due to the lower population frequency of nonsense SNPs (the similar pattern observed for synonymous SNPs could be explained by background selection in this case). The similar pattern on both of the subgenomes for TEs shows a bottleneck effect. **c** Plotting quantile pairs of the population frequencies of private nonsynonymous and synonymous SNPs in *A. suecica* and ancestral populations against each other, each species shows evidence of evolution under purifying selection, since population frequency quantiles of nonsynonymous SNPs are skewed to lower values than population frequency quantiles of synonymous SNPS.

Supplementary Figure 12. Population frequencies of presence-absence calls for TEs
that have mobilized between the subgenomes in *A. suecica*. a TEs ancestrally from *A.*
thaliana present in the *A. arenosa* subgenome of *A. suecica* and **b** TEs ancestrally from *A.*
arenosa present in the *A. thaliana* subgenome of *A. suecica*. Y axis represents the fraction of
total TEs that have mobilized between the subgenomes.

Supplementary Figure 13. Gene filtered for cross mapping in RNA-seq. a A histogram of
log2 read counts (*A. thaliana*/*A. arenosa*) of genes on the *A. thaliana* subgenome in *A. suecica.*
The green vertical line represents a log fold-change of 0 **b** A histogram of log2 read counts (*A.*
arenosa/*A. thaliana*) of genes on the *A. arenosa* subgenome in *A. suecica.* The purple vertical
line represents a log fold-change of 0

635
636

Supplementary figure 14. Expression differences between subgenomes in natural and
synthetic *A. suecica*. a The distribution of expression differences across homeologous gene
pairs in natural and synthetic *A. suecica*. **b** A heatmap of log fold change for genes in the top
5% biased toward the *A. arenosa* subgenome. The gene must be in the 5% quantile for at
least 1 accession. **c** The same as in **b** but for the *A. thaliana* subgenome. Correlations of log
fold change for genes in the tails of the distribution (top 5% quantile) for the *A. arenosa*
subgenome **d** and the *A. thaliana* subgenome **e**

659
Supplementary Figure 15. Comparison of genetic and expression distance. **a** PCA plot
of biallelic SNPs in the population of *A. thaliana* and *A. suecica* for the *A. thaliana* subgenome
of *A. suecica* (N=280,000 biallelic SNPs), of the analyzed 13,647 genes in gene expression in
addition to 500bp up and downstream of each gene sequence **b** Correlation of π (pairwise
genetic differences) and expression distance (i.e. euclidean distance) for 13,647 genes
(*=Bootstrapped 1000 times). **c** PCA plot of biallelic SNPs in the population of *A. arenosa*
(N.B. we had DNA sequencing for only 3 of the 4 accessions used in the expression analysis)
and *A. suecica* for the *A. arenosa* subgenome of *A. suecica* (N= 233,070 biallelic SNPs), of
the analyzed 13,647 genes in gene expression in addition to 500bp up and downstream of
each gene sequence **d** Correlation of Pi (pairwise genetic differences) and expression
distance (i.e. euclidean distance) for 13,647 genes (*=Bootstrapped 1000 times). *A. arenosa*
was too few samples to give reliable correlations and therefore is NA. Grey bars represent the
95 confidence intervals.

Supplementary Figure 16. **Aneuploidy is frequent in synthetic *A. suecica*. a** Comparison of FISH analyses of the reference natural *A. suecica* "ASS3 "and synthetic *A. suecica.* Synthetic *A. suecica* shows aneuploidy in both subgenomes in the F2 generation (gain of one chromosome on the *A. thaliana* subgenome (N=11) and loss of one chromosome on the *A. arenosa* subgenome (N=15)). Natural *A. suecica* shows a stable karyotype **b** DNA sequencing coverage in the reference natural *A. suecica* accession "ASS3" **c** and **d** DNA sequencing coverage in siblings of F1 synthetic *A. suecica* show different cases of aneuploidy in synthetic *A. suecica*, chromosome 4 in **c** and chromosome 11 in **d e** overlap of genes involved in cell division from figure 5e and genes previously shown to play a role in the adaptation to autopolyploidy in *A. arenosa*[59]. The little overlap in genes between *A. suecica* and *A. arenosa* highlights that successful meiosis in polyploids is likely a complex trait.

686
687

a Log fold change between Synthetic *A. suecica* F2 and parent lines

b Log fold change between Synthetic *A. suecica* F3 and parent lines

Supplementary Figure 17. No aneuploidy in synthetic *A. suecica* lines used for RNA seq based on log fold change to parent lines. Log fold change for gene expression in **a** the 2nd and **b** the 3rd generation of synthetic *A. suecica* compared to the parent lines (Chr1-5 represent comparisons with the *A. thaliana* parents and Chr6-13 represent comparisons with the *A. arenosa* parent). No clear signal of aneuploidy (i.e. an elevated increase in expression for a full chromosome) is evident.

Supplementary Figure 18. Homeologous exchange contributes to expression variance within *A. suecica* on the *A. thaliana* subgenome. A Taking the top 5% quantiles (N=646) for variation in gene expression for the *A. thaliana* subgenome we find a large cluster 7 (N=131) where the two outlier accessions in our PCA ("AS150" and "ASÖ5") are expressing these genes differently to the rest of the population. **b** Homeologous genes of this cluster on the *A. thaliana* subgenome of *A. suecica* show that these genes are not expressed in these two accessions while **c** shows the opposite pattern and are higher expressed in "AS150" and "ASÖ5" compared to the rest of the population. **d** 103 of the 131 genes in cluster 7 are located on chromosome 4 in close proximity to each other on the *A. thaliana* subgenome of the *A. suecica* reference genome and appear to be deleted in "AS5Ö5" and "AS150" as they do not have DNA sequencing coverage. The *A. arenosa* subgenome homeologs (located on chromosome 11) have twice the DNA coverage, suggesting they are duplicated, in agreement with the expectations of HE event.

728
Supplementary Figure 19 Discordant read analysis supports HE in *A. suecica* a IGV
screen grab of reads mapped to the beginning of the likely HE event in chromosome 6 (at ~
15.9Mb) before coverage depth decreases to 0 in "AS530". Arrows point to the direction of the
break along the chromosome. Discordant read pairs (cyan) map between the *A. arenosa*
subgenome on chromosome 6 and the read pair maps (green) maps to the homeologous
chromosome 1 on the *A. thaliana* subgenome (at ~5Mb) in **b**. The end of the likely HE event
in chromosome 6 (at ~18.4Mb). Discordant read pairs map (cyan) between the *A. arenosa*
subgenome in **c** and the read pair (green) maps to chromosome 1 (at ~2.8Mb) on the *A.*
thaliana subgenome in **d**. **e** Gene counts between the syntenic regions. 431 have a 1:1
relationship, 108 genes are specific to the *A. arenosa* subgenome in this region and 105 genes
are specific to the *A. thaliana* subgenome. **f** Composition of the syntenic regions between the
two subgenomes
741
742
743
744

Supplementary Figure 20 Genomic locations of genes investigated for HE signatures in
A. suecica.

a   Genes biased in expression in rosettes for both sub-genomes in *A. suecica*

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0009768	photosynthesis, light harvesting in phot...	19	17	0.59	2.5e−24
2	GO:0018298	protein−chromophore linkage	34	17	1.05	2.3e−17
3	GO:0015979	photosynthesis	181	72	5.61	2.1e−16
4	GO:0015995	chlorophyll biosynthetic process	52	18	1.61	3.0e−14
5	GO:0009735	response to cytokinin	174	28	5.39	6.7e−13
6	GO:0042742	defense response to bacterium	278	34	8.61	2.0e−12
7	GO:0055114	oxidation−reduction process	960	80	29.74	6.0e−10
8	GO:0019253	reductive pentose−phosphate cycle	15	9	0.46	2.0e−09
9	GO:0009773	photosynthetic electron transport in pho...	10	7	0.31	2.9e−09
10	GO:0009409	response to cold	281	32	8.71	1.8e−08
11	GO:0009645	response to low light intensity stimulus	14	7	0.43	2.2e−08
12	GO:0009416	response to light stimulus	533	58	16.51	5.6e−08
13	GO:0010114	response to red light	56	12	1.74	1.1e−07
14	GO:0032544	plastid translation	11	6	0.34	3.4e−07
15	GO:0019761	glucosinolate biosynthetic process	27	10	0.84	5.8e−07
16	GO:0010218	response to far red light	44	10	1.36	7.0e−07
17	GO:0010258	NADH dehydrogenase complex (plastoquinon...	4	4	0.12	9.1e−07
18	GO:0010206	photosystem II repair	13	6	0.40	1.2e−06
19	GO:0009767	photosynthetic electron transport chain	39	17	1.21	1.8e−06
20	GO:0009644	response to high light intensity	56	12	1.74	6.1e−06
21	GO:1901259	chloroplast rRNA processing	17	6	0.53	7.9e−06
22	GO:0009769	photosynthesis, light harvesting in phot...	3	3	0.09	3.0e−05
23	GO:0010207	photosystem II assembly	19	7	0.59	8.9e−05
24	GO:0009637	response to blue light	79	11	2.45	9.7e−05
25	GO:0009817	defense response to fungus, incompatible...	36	7	1.12	1.0e−04
26	GO:0010196	nonphotochemical quenching	11	4	0.34	0.00025
27	GO:0019464	glycine decarboxylation via glycine clea...	5	3	0.15	0.00028
28	GO:0006782	protoporphyrinogen IX biosynthetic proce...	12	4	0.37	0.00037
29	GO:0006412	translation	510	33	15.80	0.00051
30	GO:0009098	leucine biosynthetic process	13	4	0.40	0.00052
31	GO:0043489	RNA stabilization	7	3	0.22	0.00094

b   Genes biased in expression in floral buds for both sub-genomes in *A. suecica*

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0009733	response to auxin	250	47	18.63	1.2e−08
2	GO:0009753	response to jasmonic acid	181	35	13.49	3.5e−07
3	GO:0009751	response to salicylic acid	141	26	10.51	1.9e−06
4	GO:0080167	response to karrikin	87	20	6.48	4.5e−06
5	GO:0009737	response to abscisic acid	430	61	32.05	1.5e−05
6	GO:0010224	response to UV−B	49	13	3.65	4.2e−05
7	GO:0055085	transmembrane transport	785	93	58.50	6.2e−05
8	GO:0009739	response to gibberellin	98	19	7.30	6.5e−05
9	GO:0071456	cellular response to hypoxia	148	25	11.03	0.00017
10	GO:0006355	regulation of transcription, DNA−templat...	1356	137	101.06	0.00022
11	GO:1905582	response to mannose	6	4	0.45	0.00041
12	GO:0043496	regulation of protein homodimerization a...	3	3	0.22	0.00041
13	GO:0071492	cellular response to UV−A	3	3	0.22	0.00041
14	GO:0051603	proteolysis involved in cellular protein...	326	26	24.30	0.00059
15	GO:0010345	suberin biosynthetic process	11	5	0.82	0.00072
16	GO:0009723	response to ethylene	184	20	13.71	0.00076
17	GO:0043086	negative regulation of catalytic activit...	91	17	6.78	0.00080

c   Genes biased in expression in rosettes for the *A. arenosa* sub-genome of *A. suecica*

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0019253	reductive pentose−phosphate cycle	15	4	0.23	6.9e−05
2	GO:0009409	response to cold	281	13	4.39	0.00047
3	GO:0015700	arsenite transport	3	2	0.05	0.00072

d   Genes biased in expression in rosettes for the *A. thaliana* sub-genome of *A. suecica*

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0071456	cellular response to hypoxia	148	9	2.38	0.00065

e   Genes biased in expression in floral buds for the *A. arenosa* sub-genome of *A. suecica*

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0006032	chitin catabolic process	10	4	0.41	0.00049

755 **Supplementary Table 1. Gene ontology (GO) analysis for gene expression comparison**
756 **between whole rosettes and floral buds in *A. suecica*. a** GO enrichment for genes that
757 overlap in expression bias (log fold-change > 2 in whole rosette compared to floral buds) in
758 the *A. thaliana* and *A. arenosa* subgenome of *A. suecica*. No significant GO was found for
759 genes biased towards the *A. thaliana* subgenome of *A. suecica* for floral buds.

a

A. thaliana cluster1

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0006355	regulation of transcription, DNA-templat...	1356	264	205.59	4.8e-08
2	GO:0010167	response to nitrate	26	9	3.94	9.4e-06
3	GO:0007623	circadian rhythm	113	34	17.13	2.3e-05
4	GO:0035672	oligopeptide transmembrane transport	24	12	3.64	6.5e-05
5	GO:0046323	glucose import	28	13	4.25	8.5e-05
6	GO:0009741	response to brassinosteroid	80	26	12.13	0.00015
7	GO:0009638	phototropism	13	8	1.97	0.00017
8	GO:0009723	response to ethylene	184	52	27.90	0.00029
9	GO:0009739	response to gibberellin	98	31	14.86	0.00029
10	GO:1902600	proton transmembrane transport	125	28	18.95	0.00041
11	GO:0071577	zinc ion transmembrane transport	15	8	2.27	0.00065

b

A. thaliana cluster2

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0007018	microtubule-based movement	61	31	6.27	7.6e-16
2	GO:0009735	response to cytokinin	174	40	17.88	9.3e-09
3	GO:0009658	chloroplast organization	190	47	19.53	1.1e-07
4	GO:0000911	cytokinesis by cell plate formation	53	20	5.45	6.5e-07
5	GO:1901259	chloroplast rRNA processing	17	10	1.75	1.3e-06
6	GO:0006412	translation	510	78	52.42	1.8e-06
7	GO:0032544	plastid translation	11	7	1.13	2.7e-05
8	GO:0007088	regulation of mitotic nuclear division	49	14	5.04	7.2e-05
9	GO:0006268	DNA unwinding involved in DNA replicatio...	17	8	1.75	0.00013
10	GO:0051301	cell division	327	71	33.61	0.00017
11	GO:0006880	intracellular sequestering of iron ion	7	5	0.72	0.00020
12	GO:0009409	response to cold	281	49	28.88	0.00021
13	GO:0000413	protein peptidyl-prolyl isomerization	44	13	4.52	0.00033

c

A. thaliana cluster3

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0000727	double-strand break repair via break-ind...	9	5	0.29	3.9e-06
2	GO:0006267	pre-replicative complex assembly involve...	8	4	0.26	6.8e-05
3	GO:0046686	response to cadmium ion	256	21	8.27	8.6e-05

d

A. arenosa cluster1

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0055114	oxidation-reduction process	960	153	81.94	3.8e-13
2	GO:0098869	cellular oxidant detoxification	89	23	7.60	1.0e-06
3	GO:0019253	reductive pentose-phosphate cycle	15	8	1.28	1.0e-05
4	GO:0009854	oxidative photosynthetic carbon pathway	7	5	0.60	8.1e-05
5	GO:0051186	cofactor metabolic process	437	74	37.30	0.00010
6	GO:0009247	glycolipid biosynthetic process	43	12	3.67	0.00018
7	GO:0019682	glyceraldehyde-3-phosphate metabolic pro...	34	10	2.90	0.00038
8	GO:0009407	toxin catabolic process	23	8	1.96	0.00042
9	GO:0050665	hydrogen peroxide biosynthetic process	10	5	0.85	0.00078
10	GO:0006749	glutathione metabolic process	31	9	2.65	0.00083
11	GO:0015979	photosynthesis	181	41	15.45	0.00096

e

A. arenosa cluster2

	GO.ID	Term	Annotated	Significant	Expected	classic
1	GO:0031145	anaphase-promoting complex-dependent cat...	8	7	1.68	0.00012
2	GO:0006397	mRNA processing	311	119	65.22	0.00018
3	GO:0000245	spliceosomal complex assembly	19	11	3.98	0.00033
4	GO:0006606	protein import into nucleus	46	22	9.65	0.00060
5	GO:0051301	cell division	327	86	68.57	0.00087

760
761 **Supplementary Table 2. List of overrepresented gene ontologies on the Fig. 5e**

Materials & Methods

PacBio sequencing of *A. suecica*

We used genomic DNA from whole rosettes of one *A. suecica* ("ASS3") accession to generate PacBio sequencing data. DNA was extracted using a modified PacBio protocol for preparing *Arabidopsis* genomic DNA for size-selected ~20kb SMRTbell libraries. Briefly, whole genomic DNA was extracted from 32g of 3-4 week old plants, grown at 16°C and subjected to a 2-day dark treatment. This generated 23 micrograms of purified genomic DNA with a fragment length of >40Kb for *A. suecica*. We assessed DNA quality with a Qubit fluorometer and a Nanodrop analysis, and ran the DNA on a gel to visualize fragmentation. Genomic libraries and single-molecule real-time (SMRT) sequence data were generated at the Functional Genomics Center Zurich (FGCZ), in Switzerland. The Pacbio RSII instrument was used with P6/C4 chemistry and an average movie length of 6 hours. A total of 12 SMRT cells were processed generating 16.3Gb of DNA bases with an N50 read length of 20 Kbp and median read length of 14 Kbp. Using the same genomic library, an additional 3.3 Gbp of data was generated by a Pacbio Sequel instrument at the Vienna Biocenter Core Facilities (VBCF), in Austria, with a median read length of 10Kbp.

A. suecica genome assembly

To generate the *A. suecica* assembly we first used FALCON[65] (version 0.3.0) with a length cutoff for seed reads set to 1 Kb in size. The assembly produced 828 contigs with an N50 of 5.81 Mb and a total assembly size of 271 Mb. Additionally, we generated a Canu[66] (v.1.3.0) assembly using default settings, which resulted in 260 contigs with an N50 of 6.65 Mb and a total assembly size of 267 Mb. Then we merged the two assemblies using the software quickmerge[67]. The resulting merged assembly consisted of 929 contigs with an N50 of 9.02 Mb and a total draft assembly size of 276 Mb. We polished the assembly using Arrow[68] (smrtlink release 5.0.0.6792) and Pilon (version 1.22). For Pilon[69], 100bp (with PCR duplicates removed), and a second PCR-free 250bp, Illumina paired end reads were used that had been generated from the reference *A. suecica* accession "ASS3".

Pacbio sequencing of *A. arenosa*

A natural Swedish autotetraploid *A. arenosa* accession "Aa4" was inbred in a lab for two generations in order to reduce heterozygosity. We extracted 64 g of whole genomic DNA from three week old seedlings in the same way as described for *A. suecica* (above), generating 50 µg of purified genomic DNA with a fragment sizes longer than 40 Kb in length. The *A. arenosa* genomic libraries and SMRT sequence data were generated at the Vienna Biocenter Core Facilities (VBCF), in Austria. A Pacbio Sequel instrument was used to generate a total of 22 Gbp of data from five SMRT cells, with an N50 of 13 Kbp and median read length 10 Kbp. In addition, two runs of Oxford Nanopore sequencing were carried out at the VBCF producing 750 Mbp in 180,000 reads (median 5 Kbp and 2.6 Kbp; N50 8.7 and 6.7 Kbp, respectively).

Assembly of autotetraploid *A. arenosa*

We assembled a draft contig assembly for the autotetraploid *A. arenosa* accession "Aa4" using FALCON (version 0.3.0) as for *A. suecica*. The assembly produced 3,629 contigs with an N50 of 331 Kb, maximum contig size of 2.5 Mb and a total assembly size of 461 Mb. The assembly size is greater than the calculated haploid size of 330 Mb using FACs (see Supplementary Figure 2) probably because of the high levels of heterozygosity in *A. arenosa*. The resulting assembly was polished as described for *A. suecica*.

HiC tissue fixation and library preparation

To generate physical scaffolds for the *A. suecica* assembly we generated proximity-ligation HiC sequencing data. We collected approximately 0.5 gram of tissue from 3-week old seedlings of the same reference *A. suecica* accession. Freshly collected plant tissue was fixed in 1% formaldehyde. Cross-linking was stopped by the addition of 0.15 M Glycine. The fixed tissue was ground to a powder in liquid nitrogen and suspended in 10 ml of nuclei isolation buffer. Nuclei was digested by adding 50 U DpnII and the digested chromatin was blunt-ended by incubation with 25 µL of 0.4 mM biotin-14-dCTP and 40 U of Klenow enzyme, as described in [ref]. 20 U of T4 DNA ligase was then added to start proximity ligation. The extracted DNA was sheared by sonication with a Covaris S220 to produce 250-500bp fragments. This was followed by size fractionation using AMPure XP beads. Biotin was then removed from unligated ends. DNA fragments were blunt-end repaired and adaptors were ligated to the DNA products following the NEBNext Ultra II RNA Library Prep Kit for Illumina.

   To analyse structural rearrangements we collected tissue for 1 other natural *A. suecica* "AS530", 1 *A. thaliana* accession "6978", 1 *A. arenosa* "Aa6" and 1 synthetic *A. suecica* (F3). Each sample had two replicates. We collected tissue and prepared libraries in the same manner as described above. 125bp paired-end Illumina reads were mapped using HiCUP[70] (version 0.6.1).

Reference-guided scaffolding of the *A. suecica* genome with *LACHESIS*

We sequenced 207 million pairs of 125bp paired-end Illumina reads from the HiC library of the reference accession "ASS3". We mapped reads using HiCUP (version 0.6.1) to the draft *A. suecica* contig assembly. This resulted in ~137 million read pairs with a unique alignment.

   Setting an assembly threshold of >= 1 Kb in size, contigs of the draft *A. suecica* assembly were first assigned to the *A. thaliana* or *A. arenosa* subgenome. To do this, we used nucmer from the software MUMmer[71] (version 3.23) to perform whole-genome alignments. We aligned the draft *A. suecica* assembly to the *A. thaliana* TAIR10 reference and to our *A. arenosa* draft contig assembly, simultaneously. We used the MUMer command dnadiff to produce 1-to-1 alignments. As the subgenomes are only ~86% identical, the majority of contigs could be conclusively assigned to either subgenome by examining how similar the alignments were. Contigs that could not be assigned to a subgenome based on percentage identity were examined manually, and the length of the alignment was used to determine subgenome assignment.

   Finally, we used the software LACHESIS[72] (version 1.0.0) to scaffold our draft assembly, using the reference genomes of *A. thaliana* and *A. lyrata as* a guide to assist with scaffolding

841　the contigs (we used *A. lyrata* here instead of our draft *A. arenosa* contig assembly, as *A.*
842　*lyrata* is a chromosome-level assembly). This produced a 13-scaffold chromosome-level
843　assembly for *A. suecica*.

Construction of the *A. suecica* genetic map

We crossed natural *A. suecica* accession "AS150" with the reference accession "ASS3". The
cross was uni-directional with "AS150" as the maternal and "ASS3" as the paternal plant. F1
plants were grown, and F2 seeds were collected, from which we grew and collected 192 F2
plants. We multiplexed the samples on 96 well plates using 75bp paired end reads and
generated data of 1-2x coverage per sample. Samples were mapped to the repeat-masked
scaffolds of the reference *A. suecica* genome using BWA-MEM[73] (version 0.7.15). Samtools[74]
(version 0.1.19) was used to filter reads for proper pairs and a minimum mapping quality of 5
(-F 256 -f 3 -q 5). We called variants directly from samtools mpileup output on the sequenced
F2 individuals at known biallelic sites between the two accessions used to generate the cross
(a total of 590,537 SNPs). We required sites to have non-zero coverage in a minimum of 20
individuals and filtered SNPs to have frequency between 0.45-0.55 in our F2 population (as
the expectation is 50:50),. We removed F2 individuals that did not have genotype calls for
more than 90% of the data. This resulted in 183 individuals with genotype calls for 334,257
SNPs.
Since sequencing coverage for the F2s was low this meant we had a low probability of
calling heterozygous SNPs, and a higher probability of calling a SNP as homozygous.
Therefore, we applied a Hidden Markov Model implemented in R package HMM[75] to classify
SNPs as homozygous or heterozygous for each of our F2 lines. We then divided the genome
into 500Kb non-overlapping windows, and classified each window as homozygous (here 0 or
1, for the reference or alternate SNP) or heterozygous (here 0.5). If the frequency of 1, 0 or
0.5 represented more than 50% of the SNPs in a given window, and exceeded missing calls
(NA), the window was designated as 1, 0 or 0.5 (otherwise it was NA). This was done per
chromosome and the resulting file for each chromosome and their markers were processed in
the R package qtl[76], in order to generate a genetic map. Markers genotyped in less than 100
F2s were excluded from the analysis. Linkage groups were assigned with a minimum LOD
score of 8 and a maximum recombination fraction of 0.35. Each chromosome was assigned
to one linkage group. We defined the final marker order by the best LOD score and the lowest
number of crossover events.
Notably, the assistance of a genetic map corrected the erroneous placement of a contig
at the beginning of chromosome 1 of the *A. arenosa* subgenome. The misplaced contig was
relocated from chromosome 1 to the pericentromeric region of chromosome 2 of the *A.*
arenosa subgenome in *A. suecica*. This error was a result of a mis-assembly of chromosome
1 in the *A. lyrata* reference, as was previously pointed out [27]. Also of note, chromosome 2 of
the *A. thaliana* subgenome of *A. suecica* was previously shown to be largely devoid of
intraspecific variation, thus we had sparse marker information for this chromosome in the
genetic map. Therefore, this chromosome-scale scaffold was largely assembled by the
manual inspection of 3D-proximity information based on our HiC sequencing and reviewing
contig order using the software Juicebox[77].

Gene prediction and annotation of the *A. suecica* genome

We combined *de novo* and evidence-based approaches to predict protein coding genes. For *de novo* prediction, we trained AUGUSTUS[78] on the set of conserved single copy genes using BUSCO[79] separately on *A. thaliana* and *A. arenosa* subgenomes of *A. suecica*. The evidence-based approach included both homology to the protein sequences of the ancestral species and the transcriptome of *A. suecica*. We aligned the peptide sequences from TAIR10 *A. thaliana* assembly to the *A. thaliana* subgenome of *A. suecica*, while the peptides from *A. lyrata* from the second version of *A. lyrata* annotation[80] (Alyrata_384_v2.1) were aligned to the *A. arenosa* subgenome of *A. suecica* using GenomeThreader[81] (1.7.0). We mapped the RNAseq reads from the reference accession of *A. suecica* (ASS3) from the rosettes and flower buds tissues (see above) to the reference genome using tophat[82] and generated intron hints from the split reads using bam2hints extension of AUGUSTUS. We split the alignment into *A. thaliana* and *A. arenosa* subgenomes and assembled the transcriptome of *A. suecica* for each subgenome separately in the genome-guided mode with Trinity[83] (2.6.6). Separately for each of the subgenomes, we filtered the assembled transcripts using tpm cutoff set to 1, collapsed similar transcripts using CD-HIT[84,85] with sequence identity set to 90 percent, and chose the longest open reading frame from the six-frame translation. We then aligned the proteins from *A. thaliana* and *A. arenosa* parts of *A. suecica* to the corresponding subgenomes using GenomeThreader (1.7.0). We ran AUGUSTUS using retrained parameters from BUSCO and merged hints from all three sources, these being: (1) intron hints from *A. suecica* RNAseq, (2) homology hints from ancestral proteins and (3) hints from *A. suecica* proteins.

RepeatModeler[86] (version 1.0.11) was used in order to build a *de novo* TE consensus library for *A. suecica* and identify repetitive elements based on the genome sequence. Genome locations for the identified TE repeats were determined by using RepeatMasker[87] (version 4.0.7) and filtered for full length matches using a code described in Bailly-Bechet et. al[88]. Helitrons are the most abundant TE family in both subgenomes (Supplementary Fig. 7).

In order to compare differences between the manual curation of Tair10 and our de novo pipeline, we *de novo* annotated the long read reference genome of *Ler* [64] with RNA-seq from the same tissue and growth conditions[89].

Synthetic *A. suecica* lines

To generate synthetic *A. suecica* we crossed a natural tetraploid *A. thaliana* accession (6978 aka "Wa-1") to a natural Swedish autotetraploid *A. arenosa* ("Aa4") accession. Similar to the natural *A. suecica*, *A. thaliana* was the maternal and *A. arenosa* was the paternal plant in this cross. Crosses in the opposite direction were unsuccessful. We managed to obtain very few F1 hybrid plants, which after one round of selfing set higher levels of seed formation. The resulting synthetic line was able to self-fertilize. F2 seeds were descended from a common F1 and were similar to natural *A. suecica* in appearance. We further continued the synthetic line to F3 (selfed 3rd generation).

Synteny analysis

We performed all-against-all BLASTP search using CDS sequences for the reference *A. suecica* genome and the ancestral genomes, *A. thaliana* and *A. lyrata* (here the closest substitute reference genome for *A. arenosa*, with annotation). We used the SynMap tool[90] from the online CoGe portal[91]. We examined synteny using the default parameters for DAGChainer

926 (maximum distance between two matches = 20 genes; minimum number of aligned pairs = 5
927 genes).

Estimating copy number of rDNA repeats using short DNA reads

To measure copy number of 45S rRNA repeats in our populations of different species, we
aligned short DNA reads to a single reference 45S consensus sequence of *A. thaliana*[92]. An
A. arenosa 45S rRNA consensus sequence was constructed by finding the best hit using
BLAST in our draft *A. arenosa* contig assembly. This hit matched position 1571-8232 bp of
the *A. thaliana* consensus sequence, was 6,647 bp in length and is 97% identical to the *A.*
thaliana 45s rRNA consensus sequence. The aligned regions of these two 45S rRNA
consensus sequences, determined by BLAST, were used in copy number estimates, to ensure
that the size of the sequences were equal. The relative increase in sequence coverage of
these loci, when compared to the mean coverage for the reference genome, was used to
estimate copy number.

Plant material for RNA sequencing

Transcriptomic data generated in this study included 15 accessions of *A. suecica*, 16
accessions of *A. thaliana*, 4 accessions of *A. arenosa* and 2 generations of an artificial *A.*
suecica line (the 2nd and 3rd selfed-generation). The sibling of a paternal *A. arenosa* parent
(Aa4) and the maternal tetraploid *A. thaliana* parent (6978 aka "Wa-1") of our artificial *A.*
suecica line were included as part of our samples (Supplementary Data 2). Each accession
was replicated 3 times. Seeds were stratified in the dark for 4 days at 4°C in 1 ml of sterilised
water. Seeds were then transferred to pots in a controlled growth chamber at 21°C. Humidity
was kept constant at 60%. Pots were thinned to 2-3 seedlings after 1 week. Pots were re-
randomized each week in their trays. Whole rosettes were collected when plants reached the
7-9 true-leaf stage of development. Samples were collected between 14:00-17:00h and flash-
frozen in liquid nitrogen.

RNA extraction and library preparation

For each accession, 2-3 whole rosettes in each pot were pooled and total RNA was extracted
using the ZR Plant RNA MiniPrepTM kit. We treated the samples with DNAse, and performed
purification of mRNA and polyA selection using the AMPure XP magnetic beads and the
Poly(A) RNA Selection Kit from Lexogen. RNA quality and degradation were assessed using
the RNA Fragment Analyzer (DNF-471 stranded sensitivity RNA analysis kit, 15nt).
Concentration of RNA per sample was measured using the Qubit fluorometer. Library
preparation was carried out following the NEBNext Ultra II RNA Library Prep Kit for Illumina.
Barcoded adaptors were ligated using NEBNext Multiplex Oligos for Illumina (Index Primers
Set 1 and 2). The libraries were PCR amplified for 7 cycles. 125bp paired-end sequencing
was carried out at the VBCF on Illumina (HiSeq 2500) using multiplexing.

RNA-seq mapping and gene expression analysis

We mapped 125bp paired-end reads to the *de novo* assembled *A. suecica* reference using STAR[93] (version 2.7), we filtered for primary and uniquely aligned reads using the parameters --outfilterMultimapNmax 1 --outSamprimaryFlag OneBestScore. We quantified reads mapped to genes using --quantMode GeneCounts.

In order to reduce signals that are the result of cross mapping between the subgenomes of *A. suecica* we used *A. thaliana* and *A. arenosa* as a control. For each gene in the *A. thaliana* subgenome we compared log fold change of gene counts in our *A. thaliana* population to those in our *A. arenosa* population. We filtered for genes with a $\log_2$(*A. thaliana*/*A. arenosa*) below 0. We applied the same filters for genes on the *A. arenosa* subgenome, here a $\log_2$(*A. arenosa*/*A. thaliana*) below 0. (see Supplementary Figure 13). This reduced the number of genes analyzed from 22,383 to 22,193 on the *A. thaliana* subgenome, and 22,665 to 21,886 on the *A. arenosa* subgenome

Expression analysis was then further restricted to unique homeologous gene pairs between the subgenomes of *A. suecica* and that are 1:1 (16,742 gene pairs). Gene counts were normalized for gene size by calculating Transcripts Per Million (TPM). The effective library sizes were calculated by computing a scaling factor based on the trimmed mean of M-values (TMM) in edgeR[94], separately for each subgenome. Lowly expressed genes were removed from the analysis by keeping genes that were expressed in at least five accessions in *A. thaliana* and *A. suecica*, at least three accessions of *A. arenosa* and at least one accession of synthetic *A. suecica*. 13,647 homeologous gene pairs satisfied our expression criteria. Since *A. suecica* is expressing both subgenomes, in order to correctly normalize the effective library size in *A. suecica* accessions, the effective library size was calculated as a mean of TPM counts in both subgenomes. The effective library size of *A. thaliana* accessions was calculated for TPM counts using the *A. thaliana* subgenome of the reference genome, as genes from this subgenome will be expressed in *A. thaliana*, and the effective library size of *A. arenosa* lines using the *A. arenosa* subgenome of the reference *A. suecica* genome. Gene counts were transformed to count per million (CPM) with a prior count of 1, and were $\log_2$-transformed. We used the mean of replicates per accession for downstream analyses.

To compare homeologous genes between the subgenomes in *A. suecica* we computed a log-fold change using $\log_2$(*A. arenosa* homeolog/*A. thaliana* homeolog). For tissue-specific genes we took genes that showed a log-fold change >=2 in expression between two tissues.

For comparing homologous genes between the (sub-)genomes of *A. suecica* and the ancestral species *A. thaliana* and *A. arenosa*, we performed a Wilcoxon test independently for each of the 13,647 homeologous gene-pairs. Using the normalised CPM values, we compared the relative expression level of a gene on the *A. thaliana* subgenome between our population of *A. thaliana* and *A. suecica*. We performed the same test on the *A. arenosa* subgenome comparing relative expression of a gene between our population of *A. arenosa* and *A. suecica*. We filtered for genes with an adjusted p-value below <0.05 (using FDR correction). This amounted to 3,854 and 4,136 DEGs for the *A. thaliana* and *A. arenosa* subgenomes, respectively.

1004

Expression analysis of rRNA

1006   RNA reads were mapped in a similar manner as DNA reads for the analysis of rDNA copy
1007   number (above). Expression analysis was performed in a similar manner to protein coding
1008   genes, in edgeR. We defined the exclusive expression of a particular 45S rRNA gene by taking
1009   a cut-off of 15 for $\log_2$(CPM) as this was the maximum level of cross-mapping we observed
1010   for the ancestral species (see Supplementary Fig. 6).

Expression analysis of transposable elements

To analyse the expression of transposable elements between species, the annotated TE
consensus sequences in *A. suecica* were aligned using BLAST all vs all. Highly similar TE
sequences (more than 85% similar for more than 85% percent of the TE sequence length),
were removed, leaving 813 TE families out of 1213. Filtered *A. suecica* TEs were aligned to
annotated *A. thaliana* (TAIR10) and *A. arenosa* (the PacBio contig assembly presented in
this study) TE sequences to assign each family to an ancestral species using BLAST. 208
TE families were assigned to the *A. thaliana* parent and 171 TE families were assigned to
the *A. arenosa* parent.
RNA reads were mapped to TE sequences using a similar approach as for gene
expression analysis using edgeR. TEs that showed expression using a cut-off of $\log_2$CPM >
2 were kept. 121 *A. thaliana* TE sequences and 93 *A. arenosa* TE sequences passed this
threshold. We took the mean of replicates per accession for further downstream analyses.

Gene ontology (GO) enrichment analysis

We used the R package TopGO[95] to conduct gene ontology enrichment analysis. We used
the "weight01" algorithm when running TopGO which accounts for the hierarchical structure
of GO terms and thus implicitly corrects for multiple testing. GO annotations were based on
the *A. thaliana* ortholog of *A. suecica* genes. Gene annotations for *A. thaliana* were obtained
using the R package biomaRt[96] from Ensembl 'biomaRt::useMart(biomart = "plants_mart",
dataset = "athaliana_eg_gene", host = 'plants.ensembl.org').

Genome sizes measurements

We measured genome size for the reference *A. suecica* accession "ASS3" and the *A. arenosa*
accession used for PacBio "Aa4", using *Solanum lycopersicum* cv. Stupicke (2C = 1.96 pg
DNA) as the standard. The reference *A. lyrata* accession "MN47" and the *A. thaliana*
accession "CVI" were used as additional controls. Each sample had 2 replicates.
In brief, the leaves from three week old fresh tissue were chopped using a razor blade in 500
µl of UV Precise P extraction buffer + 10 µl mercaptoethanol per ml (kit PARTEC CyStain PI
Absolute P no. 05- 5022) to isolate nuclei. Instead of the Partec UV Precise P staining buffer,
however, 1 ml of a 5 mg DAPI solution was used, as DAPI provides DNA content histograms
with high resolution. The suspension was then passed through a 30 µm filter (Partec CellTrics
no. 04-0042-2316) and incubated for 15 minutes on ice before FACs.

Genome size was measured using flow cytometry and a FACS Aria III sorter with near UV
375nm laser for DAPI. Debris was excluded by selecting peaks when plotting DAPI-W against
DAPI-A for 20,000 events.
The data were analyzed using the flowCore[97] package in R. Genome size was estimated
by comparing the mean G1 of the standard *Solanum lycopersicum* to that of each sample to
calculate the 2C DNA content of that sample using the equation:

$$Sample\ 2C\ DNA\ content\ =\ [(sample\ G1\ peak\ mean)/(standard\ G1\ peak\ mean)]$$
$$*\ standard\ 2C\ DNA\ content$$

1050   We also measured genome size for the reference *A. suecica* accession "ASS3" using the
1051   software jellyfish[98] and findGSE[99] using kmers (21mers). The genome size estimated was
1052   312Mb, compared to the 305Mb estimated using FACs (see Supplementary Fig 1).

Mapping of TE insertions

We used PopoolationTE2[40] (version v1.10.04) to identify TE insertions. The advantage of this
TE-calling software to others is that it avoids a reference bias by treating all TEs as *de-novo*
insertions. Briefly, it works by using discordant read pairs to calculate the location and
abundance of a TE in the genome for an accession of interest.
We mapped 100 bp Illumina DNA reads from [16,26,100], in addition to our newly generated
synthetic *A. suecica* using BWA MEM[73] (version 0.7.15) to a repeat-masked version of the *A.*
suecica reference genome, concatenated with our annotated repeat sequences (see 'Genome
annotation'), as this is the data format required by PopoolationTE2. Reads were given an
increased penalty of 15 for being unpaired. Reads were de-duplicated using Samtools[74] rmdup
(version 1.9). The resulting bam files were then provided to PopoolationTE2 to identify TE
insertions in the genome of each of our *A. suecica*, *A. thaliana* and *A. arenosa* accessions.
We used a mapping quality of 10 for the read in the discordant read pair mapping to the
genome. We used the 'separate' mode in the 'identify TE signatures' step and a '--min-
distance -200 --max-distance 500' in the 'pairupsignatues' step of the pipeline. TE counts
within each accession were merged if they fell within 400 bp of each other and if they mapped
to the same TE sequence. All TE counts (i.e. the processed TE counts for each accession)
were then combined to produce a population-wide count estimate. Population wide TE
insertions were merged if they mapped to the same TE sequence and fell within 400 bp of
each other. Coverage of each TE insertion in the population was also calculated for each
accession. The final file was a list TE insertions present in the population and the presence or
absence (or "NA" if there was no coverage to support the presence or absence of a TE
insertion) in each accession analyzed (Supplementary Data 1a,b). To avoid insertions that
could be potential mismapping, we excluded the region between 5 and 10Mb on chromosome
2 of the *A. thaliana* subgenome as this area shows synteny with an unplaced contig.

Assigning ancestry to TE sequences

In order to examine TE consensus sequences that have mobilized between the subgenomes
of *A. suecica*, we first examined which of our TE consensus sequences (N=1152) have at
least the potential to mobilize (i.e. have full length TE copies in the genome of *A. suecica*).
We filtered for TE consensus sequences that had TE copies in the genome of *A. suecica*
that are more than 80% similar in identity for more than 80% of the consensus sequence
length (N=936). Of these, 188 consensus sequences were private to the *A. thaliana*

subgenome, 460 were private to the *A. arenosa* subgenome, and 288 TE consensus
sequences were present in both subgenomes of *A. suecica*. To determine if TEs have
jumped from the *A. thaliana* subgenome to the *A. arenosa* subgenome and vice versa we
next needed to assign ancestry to these 288 TE consensus sequences. To do this we used
BLAST to search for these consensus sequences in the ancestral genomes of *A. suecica*,
using the TAIR10 *A. thaliana* reference and our *A. arenosa* PacBio contig assembly. Using
the same 80%-80% rule we assigned 55 TEs to *A. arenosa* and 15 TEs to *A. thaliana*
ancestry.

Read mapping and SNP calling

To call biallelic SNPs we mapped reads to the *A. suecica* reference genome using the same
filtering parameters described in "Mapping of TE insertions". Biallelic SNPs were called using
HaplotypeCaller from GATK[101] (version 3.8) using default quality thresholds. SNPs were
annotated using SnpEff[102]. Biallelic SNPs on the *A. thaliana* subgenome were polarized
using 38 diploid *A. lyrata* lines[26] and biallelic SNPS on the *A. arenosa* subgenome were
polarized using 30 *A. thaliana* accessions[100] closely related to *A. suecica*[16].

Chromosome preparation and FISH

Whole inflorescences of *A. arenosa*, *A. suecica* and *A. thaliana* were fixed in freshly
prepared ethanol:acetic acid fixative (3:1) overnight, transferred into 70% ethanol and stored
at -20°C until use. Selected inflorescences were rinsed in distilled water and citrate buffer
(10 mM sodium citrate, pH 4.8), and digested by a 0.3% mix of pectolytic enzymes
(cellulase, cytohelicase, pectolyase; all from Sigma-Aldrich) in citrate buffer for c. 3 hrs.
Mitotic chromosome spreads were prepared from pistils as previously described[103] by
Mandáková and Lysak and suitable slides pretreated by RNase (100 µg/ml, AppliChem) and
pepsin (0.1 mg/ml, Sigma-Aldrich).
For identification of *A. thaliana* and *A. arenosa* subgenomes in the allotetraploid
genome of *A. suecica*, FISH probes were made from plasmids pARR20–1 or pAaCEN
containing 180 bp of *A. thaliana* (pAL; Vongs et al. 1993) or ~250 bp of *A. arenosa* (pAa;
Kamm et al. 1995) pericentromeric repeats, respectively. The *A. thaliana* BAC clone T15P10
(AF167571) bearing 45S rRNA gene repeats was used for in situ localization of NORs.
Individual probes were labeled with biotin-dUTP, digoxigenin-dUTP and Cy3-dUTP by nick
translation, pooled, precipitated, and resuspended in 20 µl of hybridization mixture [50%
formamide and 10% dextran sulfate in 2× saline sodium citrate (2× SSC)] per slide as
previously described[96].
Probes and chromosomes were denatured together on a hot plate at 80°C for 2 min
and incubated in a moist chamber at 37°C overnight. Post hybridization washing was
performed in 20% formamide in 2× SSC at 42°C. Fluorescent detection was as follows:
biotin-dUTP was detected by avidin–Texas Red (Vector Laboratories) and amplified by goat
anti-avidin–biotin (Vector Laboratories) and avidin–Texas Red; digoxigenin-dUTP was
detected by mouse anti-digoxigenin (Jackson ImmunoResearch) and goat anti-mouse Alexa
Fluor 488 (Molecular Probes). Chromosomes were counterstained with DAPI (4',6-
diamidino-2-phenylindole; 2 µg/ml) in Vectashield (Vector Laboratories). Fluorescent signals
were analyzed and photographed using a Zeiss Axioimager epifluorescence microscope and
a CoolCube camera (MetaSystems). Images were acquired separately for the four

fluorochromes using appropriate excitation and emission filters (AHF Analysentechnik). The monochromatic images were pseudo colored and merged using Adobe Photoshop CS6 software (Adobe Systems).

DAP-seq enrichment analysis for transcription factor target genes

We downloaded the target genes of transcription factors from the plant cistrome database (http://neomorph.salk.edu/dap_web/pages/index.php), which is a collection of transcription factor binding sites and their target genes, in *A. thaliana*, based on DAP-seq[104]. To test for enrichment of a gene set (for example the genes in *A. thaliana* cluster 2 on Fig. 3) for target genes of a particular transcription factor, we performed a hyper-geometric test in R. As a background we used the total 13,647 genes used in our gene expression analysis. We then performed FDR correction for multiple testing to calculate an accurate p-value of the enrichment.

Data Availability

The raw sequencing for genome assembly ####. Genome assembly and annotation ####. The raw RNA-seq reads SRA ####. TE presence/absence calls for *A. suecica* and ancestral species can be found in the Supplementary Data 1a,b.

Acknowledgments

This work was supported, in part, by DFG SPP 1529 to M.N. and Detlef Weigel. T.M. and M.A.L. were supported by the Czech Science Foundation (grant no. 19-03442S) and the CEITEC 2020 project (grant no. LQ1601). P.Y.N. acknowledges postdoctoral fellowship of the Research Foundation–Flanders (12S9618N). We thank the Next Generation Sequencing Unit of the Vienna Biocenter Core Facilities (VBCF) for assistance. We thank Svante Holm and Torbjörn Säll for material collections and helpful discussions. We also thank Yves Van de Peer for providing useful feedback on the manuscript.

References

1   Van de Peer Y, Mizrachi E, Marchal K. The evolutionary significance of polyploidy. *Nat Rev Genet* 2017; **18**: 411–424.

2   Soltis PS, Soltis DE. Ancient WGD events as drivers of key innovations in angiosperms. *Curr Opin Plant Biol* 2016; **30**: 159–165.

3   Dehal P, Boore JL. Two rounds of whole genome duplication in the ancestral vertebrate. *PLoS Biol* 2005; **3**: e314.

4   Li Z, Tiley GP, Galuska SR, Reardon CR, Kidder TI, Rundell RJ *et al.* Multiple large-scale gene and genome duplications during the evolution of hexapods. *Proc Natl Acad Sci U S A* 2018; **115**: 4713–4718.

5 Chen ZJ, Sreedasyam A, Ando A, Song Q, De Santiago LM, Hulse-Kemp AM *et*
al. Genomic diversifications of five Gossypium allopolyploid species and their
impact on cotton improvement. *Nat Genet* 2020; **52**: 525–533.

6 Edger PP, Poorten TJ, VanBuren R, Hardigan MA, Colle M, McKain MR *et al.*
Origin and evolution of the octoploid strawberry genome. *Nat Genet* 2019; **51**:
541–547.

7 Ramírez-González RH, Borrill P, Lang D, Harrington SA, Brinton J, Venturini L
et al. The transcriptional landscape of polyploid wheat. *Science* 2018; **361**.
doi:10.1126/science.aar6089.

8 Zhuang W, Chen H, Yang M, Wang J, Pandey MK, Zhang C *et al.* The genome
of cultivated peanut provides insight into legume karyotypes, polyploid evolution
and crop domestication. Nature Genetics. 2019; **51**: 865–876.

9 Bertioli DJ, Jenkins J, Clevenger J, Dudchenko O, Gao D, Seijo G *et al.* The
genome sequence of segmental allotetraploid peanut Arachis hypogaea. *Nat*
Genet 2019; **51**: 877–884.

10 Kasianov AS, Klepikova AV, Kulakovskiy IV, Gerasimov ES, Fedotova AV,
Besedina EG *et al.* High-quality genome assembly of Capsella bursa-pastoris
reveals asymmetry of regulatory elements at early stages of polyploid genome
evolution. *Plant J* 2017; **91**: 278–291.

11 Kryvokhyzha D, Milesi P, Duan T, Orsucci M, Wright SI, Glémin S *et al.* Towards
the new normal: Transcriptomic convergence and genomic legacy of the two
subgenomes of an allopolyploid weed (Capsella bursa-pastoris). *PLoS Genet*
2019; **15**: e1008131.

12 Douglas GM, Gos G, Steige KA, Salcedo A, Holm K, Josephs EB *et al.* Hybrid
origins and the earliest stages of diploidization in the highly successful recent
polyploid Capsella bursa-pastoris. *Proc Natl Acad Sci U S A* 2015; **112**: 2806–
2811.

13 Griffiths AG, Moraga R, Tausen M, Gupta V, Bilton TP, Campbell MA *et al.*
Breaking Free: The Genomics of Allopolyploidy-Facilitated Niche Expansion in
White Clover. *Plant Cell* 2019; **31**: 1466–1487.

14 Gordon SP, Contreras-Moreira B, Levy JJ, Djamei A, Czedik-Eysenberg A,
Tartaglio VS *et al.* Gradual polyploid genome evolution revealed by pan-
genomic analysis of Brachypodium hybridum and its diploid progenitors. *Nat*
Commun 2020; **11**: 3670.

15 Catalán P, López-Álvarez D, Bellosta C, Villar L. Updated taxonomic
descriptions, iconography, and habitat preferences of Brachypodium distachyon,
B. stacei , and B. hybridum (Poaceae). *An Jard Bot Madr* 2016; **73**: 028.

16 Novikova PY, Tsuchimatsu T, Simon S, Nizhynska V, Voronin V, Burns R *et al.*
Genome Sequencing Reveals the Origin of the Allotetraploid Arabidopsis
suecica. *Mol Biol Evol* 2017; **34**: 957–968.

17 Fowler NL, Levin DA. Ecological Constraints on the Establishment of a Novel
Polyploid in Competition with Its Diploid Progenitor. *Am Nat* 1984; **124**: 703–
711.

18 Bomblies K, Madlung A. Polyploidy in the Arabidopsis genus. *Chromosome Res*
2014; **22**: 117–134.

19 Hollister JD, Arnold BJ, Svedin E, Xue KS, Dilkes BP, Bomblies K. Genetic
adaptation associated with genome-doubling in autotetraploid Arabidopsis
arenosa. *PLoS Genet* 2012; **8**: e1003093.

20 Bomblies K, Jones G, Franklin C, Zickler D, Kleckner N. The challenge of
evolving stable polyploidy: could an increase in 'crossover interference distance'
play a central role? *Chromosoma* 2016; **125**: 287–300.

21 Parisod C, Alix K, Just J, Petit M, Sarilar V, Mhiri C *et al.* Impact of transposable
elements on the organization and function of allopolyploid genomes. *New Phytol*
2010; **186**: 37–45.

22 McClintock B. The significance of responses of the genome to challenge.
Science. 1984; **226**: 792–801.

23 Hohmann N, Wolf EM, Lysak MA, Koch MA. A Time-Calibrated Road Map of
Brassicaceae Species Radiation and Evolutionary History. *Plant Cell* 2015; **27**:
2770–2784.

24 O'Kane SL, Schaal BA, Al-Shehbaz IA. The Origins of Arabidopsis suecica
(Brassicaceae) as Indicated by Nuclear rDNA Sequences. *Syst Bot* 1996; **21**:
559–566.

25 Jakobsson M, Hagenblad J, Tavaré S, Säll T, Halldén C, Lind-Halldén C *et al.* A
unique recent origin of the allotetraploid species Arabidopsis suecica: Evidence
from nuclear DNA markers. *Mol Biol Evol* 2006; **23**: 1217–1231.

26 Novikova PY, Hohmann N, Nizhynska V, Tsuchimatsu T, Ali J, Muir G *et al.*
Sequencing of the genus Arabidopsis identifies a complex history of
nonbifurcating speciation and abundant trans-specific polymorphism. *Nat Genet*
2016; **48**: 1077–1082.

27 Slotte T, Hazzouri KM, Ågren JA, Koenig D, Maumus F, Guo Y-L *et al.* The
Capsella rubella genome and the genomic consequences of rapid mating
system evolution. *Nat Genet* 2013; **45**: 831–835.

28 Ozkan H, Levy AA, Feldman M. Allopolyploidy-Induced Rapid Genome
Evolution in the Wheat (Aegilops–Triticum) Group. *Plant Cell* 2001; **13**: 1735–
1747.

29 Shaked H, Kashkush K, Ozkan H, Feldman M, Levy AA. Sequence elimination
and cytosine methylation are rapid and reproducible responses of the genome
to wide hybridization and allopolyploidy in wheat. *Plant Cell* 2001; **13**: 1749–
1759.

30   Chalhoub B, Denoeud F, Liu S, Parkin IAP, Tang H, Wang X *et al.* Plant
1243   genetics. Early allopolyploid evolution in the post-Neolithic Brassica napus
1244   oilseed genome. *Science* 2014; **345**: 950–953.

1245   31   Madlung A, Tyagi AP, Watson B, Jiang H, Kagochi T, Doerge RW *et al.*
1246   Genomic changes in synthetic Arabidopsis polyploids. *Plant J* 2005; **41**: 221–
1247   230.

1248   32   Copenhaver GP, Pikaard CS. Two-dimensional RFLP analyses reveal
1249   megabase-sized clusters of rRNA gene variants in Arabidopsis thaliana,
1250   suggesting local spreading of variants as the mode for gene homogenization
1251   during concerted evolution. The Plant Journal. 1996; **9**: 273–282.

1252   33   Navashin M. Chromosome Alterations Caused by Hybridization and Their
1253   Bearing upon Certain General Genetic Problems. *Cytologia* 1934; **5**: 169–203.

1254   34   Tucker S, Vitins A, Pikaard CS. Nucleolar dominance and ribosomal RNA gene
1255   silencing. *Curr Opin Cell Biol* 2010; **22**: 351–356.

1256   35   Pontes O, Neves N, Silva M, Lewis MS, Madlung A, Comai L *et al.*
1257   Chromosomal locus rearrangements are a rapid response to formation of the
1258   allotetraploid Arabidopsis suecica genome. Proceedings of the National
1259   Academy of Sciences. 2004; **101**: 18240–18245.

1260   36   Long Q, Rabanal FA, Meng D, Huber CD, Farlow A, Platzer A *et al.* Massive
1261   genomic variation and strong selection in Arabidopsis thaliana lines from
1262   Sweden. *Nat Genet* 2013; **45**: 884–890.

1263   37   Rabanal FA, Mandáková T, Soto-Jiménez LM, Greenhalgh R, Parrott DL,
1264   Lutzmayer S *et al.* Epistatic and allelic interactions control expression of
1265   ribosomal RNA gene clusters in Arabidopsis thaliana. *Genome Biol* 2017; **18**:
1266   75.

1267   38   Steige KA, Slotte T. Genomic legacies of the progenitors and the evolutionary
1268   consequences of allopolyploidy. *Curr Opin Plant Biol* 2016; **30**: 88–93.

1269   39   Vicient CM, Casacuberta JM. Impact of transposable elements on polyploid
1270   plant genomes. *Ann Bot* 2017; **120**: 195–207.

1271   40   Kofler R, Gomez-Sanchez D, Schlotterer C. PoPoolationTE2: Comparative
1272   Population Genomics of Transposable Elements Using Pool-Seq. *Mol Biol Evol*
1273   2016; **33**: 2759–2764.

1274   41   Quadrana L, Bortolini Silveira A, Mayhew GF, LeBlanc C, Martienssen RA,
1275   Jeddeloh JA *et al.* The Arabidopsis thaliana mobilome and its impact at the
1276   species level. *Elife* 2016; **5**. doi:10.7554/eLife.15716.

1277   42   Stuart T, Eichten SR, Cahn J, Karpievitch YV, Borevitz JO, Lister R. Population
1278   scale mapping of transposable element diversity reveals links to gene regulation
1279   and epigenomic variation. *Elife* 2016; **5**. doi:10.7554/eLife.20777.

1280   43   Wolfe KH. Yesterday's polyploids and the mystery of diploidization. *Nat Rev*

1281   *Genet* 2001; **2**: 333–341.

Conant GC, Birchler JA, Pires JC. Dosage, duplication, and diploidization: clarifying the interplay of multiple models for duplicate gene evolution over time. *Curr Opin Plant Biol* 2014; **19**: 91–98.

Aköz G, Nordborg M. The Aquilegia genome reveals a hybrid origin of core eudicots. *Genome Biol* 2019; **20**: 256.

Jiao Y, Wickett NJ, Ayyampalayam S, Chanderbali AS, Landherr L, Ralph PE *et al.* Ancestral polyploidy in seed plants and angiosperms. *Nature* 2011; **473**: 97–100.

Soltis PS, Marchant DB, Van de Peer Y, Soltis DE. Polyploidy and genome evolution in plants. *Curr Opin Genet Dev* 2015; **35**: 119–125.

Thomas BC, Pedersen B, Freeling M. Following tetraploidy in an Arabidopsis ancestor, genes were removed preferentially from one homeolog leaving clusters enriched in dose-sensitive genes. *Genome Res* 2006; **16**: 934–946.

Schnable JC, Springer NM, Freeling M. Differentiation of the maize subgenomes by genome dominance and both ancient and ongoing gene loss. *Proc Natl Acad Sci U S A* 2011; **108**: 4069–4074.

Cheng F, Wu J, Fang L, Sun S, Liu B, Lin K *et al.* Biased gene fractionation and dominant gene expression among the subgenomes of Brassica rapa. *PLoS One* 2012; **7**: e36442.

Renny-Byfield S, Gong L, Gallagher JP, Wendel JF. Persistence of subgenomes in paleopolyploid cotton after 60 my of evolution. *Mol Biol Evol* 2015; **32**: 1063–1071.

Garsmeur O, Schnable JC, Almeida A, Jourda C, D'Hont A, Freeling M. Two evolutionarily distinct classes of paleopolyploidy. *Mol Biol Evol* 2014; **31**: 448–454.

Li Q, Qiao X, Yin H, Zhou Y, Dong H, Qi K *et al.* Unbiased subgenome evolution following a recent whole-genome duplication in pear (Pyrus bretschneideri Rehd.). *Hortic Res* 2019; **6**: 34.

Bird KA, Niederhuth C, Ou S, Gehan M, Chris Pires J, Xiong Z *et al.* Replaying the evolutionary tape to investigate subgenome dominance in allopolyploid Brassica napus. doi:10.1101/814491.

Edger PP, Smith R, McKain MR, Cooley AM, Vallejo-Marin M, Yuan Y *et al.* Subgenome Dominance in an Interspecific Hybrid, Synthetic Allopolyploid, and a 140-Year-Old Naturally Established Neo-Allopolyploid Monkeyflower. *Plant Cell* 2017; **29**: 2150–2167.

Alger EI, Edger PP. One subgenome to rule them all: underlying mechanisms of subgenome dominance. *Curr Opin Plant Biol* 2020; **54**: 108–113.

57　Carlson KD, Fernandez-Pozo N, Bombarely A, Pisupati R, Mueller LA, Madlung
A. Natural variation in stress response gene activity in the allopolyploid
Arabidopsis suecica. *BMC Genomics* 2017; **18**: 653.

58　Chang PL, Dilkes BP, McMahon M, Comai L, Nuzhdin SV. Homoeolog-specific
retention and use in allotetraploid Arabidopsis suecica depends on parent of
origin and network partners. *Genome Biol* 2010; **11**: R125.

59　Yant L, Hollister JD, Wright KM, Arnold BJ, Higgins JD, Franklin FC *et al.*
Meiotic adaptation to genome duplication in Arabidopsis arenosa. *Curr Biol*
2013; **23**: 2151–2156.

60　Morgan C, Zhang H, Henry CE, Franklin FCH, Bomblies K. Derived alleles of
two axis proteins affect meiotic traits in autotetraploid Arabidopsis arenosa. *Proc*
Natl Acad Sci U S A 2020; **117**: 8980–8988.

61　Haga N, Kobayashi K, Suzuki T, Maeo K, Kubo M, Ohtani M *et al.* Mutations in
MYB3R1 and MYB3R4 cause pleiotropic developmental defects and preferential
down-regulation of multiple G2/M-specific genes in Arabidopsis. *Plant Physiol*
2011; **157**: 706–717.

62　Preite V, Sailer C, Syllwasschy L, Bray S, Ahmadi H, Krämer U *et al.*
Convergent evolution in Arabidopsis halleri and Arabidopsis arenosa on
calamine metalliferous soils. *Philos Trans R Soc Lond B Biol Sci* 2019; **374**:
20180243.

63　Darwin C. The origin of species by means of natural selection : or The
preservation of favored races in the struggle for life / by Charles Darwin. 1872.
doi:10.5962/bhl.title.2106.

64　Zapata L, Ding J, Willing E-M, Hartwig B, Bezdan D, Jiao W-B *et al.*
Chromosome-level assembly of Arabidopsis thaliana Ler reveals the extent of
translocation and inversion polymorphisms. *Proc Natl Acad Sci U S A* 2016;
113: E4052–60.

65　Chin C-S, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, Clum A *et al.*
Phased diploid genome assembly with single-molecule real-time sequencing.
Nat Methods 2016; **13**: 1050–1054.

66　Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and
repeat separation. *Genome Res* 2017; **27**: 722–736.

67　Chakraborty M, Baldwin-Brown JG, Long AD, Emerson JJ. Contiguous and
accurate de novo assembly of metazoan genomes with modest long read
coverage. *Nucleic Acids Res* 2016; **44**: e147.

68　Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C *et al.*
Nonhybrid, finished microbial genome assemblies from long-read SMRT
sequencing data. Nature Methods. 2013; **10**: 563–569.

69　Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S *et al.* Pilon:

an integrated tool for comprehensive microbial variant detection and genome
assembly improvement. *PLoS One* 2014; **9**: e112963.

70 Wingett S, Ewels P, Furlan-Magaril M, Nagano T, Schoenfelder S, Fraser P *et*
al. HiCUP: pipeline for mapping and processing Hi-C data. *F1000Res* 2015; **4**:
1310.

71 Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A.
MUMmer4: A fast and versatile genome alignment system. *PLoS Comput Biol*
2018; **14**: e1005944.

72 Burton JN, Adey A, Patwardhan RP, Qiu R, Kitzman JO, Shendure J.
Chromosome-scale scaffolding of de novo genome assemblies based on
chromatin interactions. *Nat Biotechnol* 2013; **31**: 1119–1125.

73 Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* 2009; **25**: 1754–1760.

74 Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N *et al.* The
Sequence Alignment/Map format and SAMtools. *Bioinformatics* 2009; **25**: 2078–
2079.

75 Himmelmann L. HMM: Hidden Markov Models. *R package version* 2010; **1**.

76 Broman KW, Wu H, Sen S, Churchill GA. R/qtl: QTL mapping in experimental
crosses. *Bioinformatics* 2003; **19**: 889–890.

77 Durand NC, Shamim MS, Machol I, Rao SSP, Huntley MH, Lander ES *et al.*
Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C
Experiments. *Cell Syst* 2016; **3**: 95–98.

78 Stanke M, Morgenstern B. AUGUSTUS: a web server for gene prediction in
eukaryotes that allows user-defined constraints. *Nucleic Acids Res* 2005; **33**:
W465–7.

79 Seppey M, Manni M, Zdobnov EM. BUSCO: Assessing Genome Assembly and
Annotation Completeness. In: Kollmar M (ed). *Gene Prediction: Methods and*
Protocols. Springer New York: New York, NY, 2019, pp 227–245.

80 Rawat V, Abdelsamad A, Pietzenuk B, Seymour DK, Koenig D, Weigel D *et al.*
Improving the Annotation of Arabidopsis lyrata Using RNA-Seq Data. *PLoS One*
2015; **10**: e0137391.

81 Gremme G, Brendel V, Sparks ME, Kurtz S. Engineering a software tool for
gene structure prediction in higher organisms. *Information and Software*
Technology 2005; **47**: 965–978.

82 Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2:
accurate alignment of transcriptomes in the presence of insertions, deletions
and gene fusions. *Genome Biol* 2013; **14**: R36.

83 Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I *et al.* Full-

length transcriptome assembly from RNA-Seq data without a reference genome.
Nat Biotechnol 2011; **29**: 644–652.

84   Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* 2012; **28**: 3150–3152.

85   Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* 2006; **22**: 1658–1659.

86   Smit AFA, Hubley R. RepeatModeler Open-1.0 http://www.repeatmasker.org.
2008-2015.

87   Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0. . 2013-2015.

88   Bailly-Bechet M, Haudry A, Lerat E. 'One code to find them all': a perl tool to
conveniently parse RepeatMasker output files. *Mob DNA* 2014; **5**: 13.

89   Lei L, Steffen JG, Osborne EJ, Toomajian C. Plant organ evolution revealed by
phylotranscriptomics in Arabidopsis thaliana. *Sci Rep* 2017; **7**: 7567.

90   Lyons E, Pedersen B, Kane J, Freeling M. The value of nonmodel genomes and
an example using SynMap within CoGe to dissect the hexaploidy that predates
the rosids. *Trop Plant Biol* 2008; **1**: 181–190.

91   Lyons E, Freeling M. How to usefully compare homologous plant genes and
chromosomes as DNA sequences. *Plant J* 2008; **53**: 661–673.

92   Rabanal FA, Nizhynska V, Mandáková T, Novikova PY, Lysak MA, Mott R *et al.*
Unstable Inheritance of 45S rRNA Genes in Arabidopsis thaliana. *G3* 2017; **7**:
1201–1209.

93   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S *et al.* STAR:
ultrafast universal RNA-seq aligner. *Bioinformatics* 2013; **29**: 15–21.

94   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics*
2010; **26**: 139–140.

95   Alexa A, Rahnenfuhrer J. topGO: enrichment analysis for gene ontology. *R*
package version 2010; **2**: 2010.

96   Durinck S, Spellman PT, Birney E, Huber W. Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat*
Protoc 2009; **4**: 1184–1191.

97   Hahne F, LeMeur N, Brinkman RR, Ellis B, Haaland P, Sarkar D *et al.* flowCore:
a Bioconductor package for high throughput flow cytometry. *BMC Bioinformatics*
2009; **10**: 106.

98   Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting
of occurrences of k-mers. *Bioinformatics* 2011; **27**: 764–770.

99 Sun H, Ding J, Piednoël M, Schneeberger K. findGSE: estimating genome size
variation within human and Arabidopsis using k-mer frequencies. *Bioinformatics*
2018; **34**: 550–557.

100 Genomes Consortium. Electronic address, magnus nordborg gmi oeaw ac at,
Genomes, Consortium. 1,135 Genomes Reveal the Global Pattern of
Polymorphism in Arabidopsis thaliana. *Cell* 2016; **166**: 481–491.

101 McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A *et al.*
The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. *Genome Res* 2010; **20**: 1297–1303.

102 Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L *et al.* A program for
annotating and predicting the effects of single nucleotide polymorphisms,
SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2;
iso-3. *Fly* 2012; **6**: 80–92.

103 Mandáková T, Lysak MA. Chromosome Preparation for Cytogenetic Analyses in
Arabidopsis. *Curr Protoc Plant Biol* 2016; **1**: 43–51.

104 O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR *et al.*
Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. *Cell*
2016; **165**: 1280–1292.

1451