Cognitive deficits found in a pro-inflammatory state are independent of ERK 1/2 signaling in the murine brain hippocampus treated with Shiga toxin 2 from enterohemorrhagic *Escherichia Coli*

Clara Berdasco[1,2], Alipio Pinto[1,2], Mariano Blake[3], Fernando Correa[4], Nadia A. Longo Carbajosa[5], Patricia Geogeghan[6], Adriana Cangelosi[6], Mariela M. Gironacci[5], Jorge Goldstein[1,2*]

[1] Departamento de Fisiología, Facultad de Medicina, Universidad de Buenos Aires, Buenos Aires, Argentina

[2] Laboratorio de Neurofisiopatología, Instituto de Fisiología y Biofísica "Houssay", Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad de Buenos Aires, Buenos Aires, Argentina

[3] Instituto de Fisiología y Biofísica "Houssay", Consejo Nacional de Investigaciones Científicas y Técnicas, Facultad de Medicina, Universidad de Buenos Aires, Buenos Aires, Argentina

[4] Laboratorio de Fisiopatología de la Preñez y el Parto, Centro de Estudios Farmacológicos y Botánicos, Consejo Nacional de Investigaciones Científicas y Técnicas, Facultad de Medicina, Universidad de Buenos Aires, Buenos Aires, Argentina

[5] Departamento de Química Biológica, Instituto de Química y Fisicoquímica Biológicas-Consejo Nacional de Investigaciones Científicas y Técnicas, Facultad de Farmacia y Bioquímica, Universidad de Buenos Aires, Buenos Aires, Argentina

21   [6] Centro Nacional de Control de Calidad de Biológicos, Administración Nacional de

22   Laboratorios e Institutos de Salud "Dr. Carlos G. Malbrán", Buenos Aires, Argentina

23   * Corresponding author

24   E-mail:jogol@fmed.uba.ar ; jorgoldstein@gmail.com (JG)

Abstract

Shiga toxin 2 (Stx2) from enterohemorrhagic *Escherichia coli* (EHEC) produces

hemorrhagic colitis, hemolytic uremic syndrome (HUS) and acute encephalopathy. The

mortality rate in HUS increases significantly when the central nervous system (CNS) is

involved. Besides, EHEC also releases lipopolysaccharide (LPS). Many reports have

described cognitive dysfunctions in HUS patients, the hippocampus being one of the brain

areas targeted by EHEC infection. In this context, a translational murine model of

encephalopathy was employed to establish the deleterious effects of Stx2 and the

contribution of LPS in the hippocampus. Results demonstrate that systemic administration

of a sublethal dose of Stx2 reduced memory index and produced depression like behavior,

pro-inflammatory cytokine release and NF-kB activation independent of the ERK 1/2

signaling pathway. On the other hand, LPS activated NF-kB dependent on ERK 1/2

signaling pathway. Cotreatment of Stx2 with LPS aggravated the pathologic state, while

dexamethasone treatment succeeded in preventing behavioral alterations. Our present work

suggests that the use of drugs such as corticosteroids or NF-kB signaling inhibitors may

serve as neuroprotectors from EHEC infection.

Author Summary

Shiga toxin (Stx) from enterohemorrhagic Escherichia coli (EHEC) is one of the most virulent factors responsible for hemolytic uremic syndrome (HUS). Stx2, the endemic variant targets the brain, among other organs, thus inducing encephalopathies. Central nervous system (CNS) compromise was the main predictor of death in patients with HUS. Stx2 may exert a direct action in the CNS, by disrupting the neurovascular unit. In this context, we investigate the molecular signaling triggered by Stx2 in the murine brain hippocampus involved in inflammatory mechanisms that altered hippocampal-related cognitive behaviors. The present data underscore that the use of drugs such as dexamethasone or those blocking the cascade by preventing NF-kB translocation to the nucleus may serve as effective neuroprotectors with potentially beneficial use in the clinic.

Introduction

Shiga toxin 2 (Stx2) from Shiga toxin producing *Escherichia coli* (STEC) is one of the most virulent factors responsible for hemolytic uremic syndrome (HUS), a triad of clinical events that includes acute renal failure, nonimmune microangiopathic hemolytic anemia and thrombocytopenia. Prodromal bloody diarrhea may occur due to infection with diarrheagenic *Escherichia coli* pathotypes, mainly the serotype O157: H7 of enterohemorrhagic *Escherichia coli* (EHEC) [1-3].

The highest rates of HUS on the planet (more than 400 annual cases) are found in Argentina, where the incidence of pediatric cases reaches up to 17 cases per 100,000 children under 5 years of age [4]. The lethality reported in these patients is between 1 to 4% but it may rise up to 3-fold when the central nervous system (CNS) is involved [5-7].

Acute encephalopathy produced by STEC infection represents a critical episode in severe cases of disease [8]. We and other laboratories have demonstrated that Stx2 may exert a direct action in the CNS, in neurons and glial cells [9-12]. Furthermore, it has been reported that STEC may affect the CNS without renal involvement [13].

Stx2 produces its deleterious action through its canonical cell membrane receptor globotriaosylceramide (Gb3), a glycosphingolipid localized on the external bilayer lipid cell membrane. Once Stx2 binds to Gb3, the toxin reaches the ribosomes through a retrograde pathway and triggers ribotoxic stress leading to pro-apoptotic and pro-inflammatory mechanisms [14]. In addition to Stx2, the Gram-negative STEC also releases lipopolysaccharide (LPS), an endotoxin derived from its outer membrane [15]. It has been reported that patients with HUS/encephalopathy also present endotoxemia. Several animal models developed to study the effect of endotoxemia on STEC pathogenesis have shown that LPS amplifies the deleterious effects of STEC. In connection with this amplification, we have previously reported a relevant pro-inflammatory component in CNS damage produced by Stx2 [10, 11, 16].

Several reports have described cognitive dysfunction in HUS patients, the hippocampus being one of these damaged areas in the brain [17-20]. Alterations found in these patients include trouble finding words, severe alteration of consciousness, late memory decline, orientation deficits, seizures and coma. We have previously demonstrated morphological alterations induced by Stx2 and/or Stx2+LPS in the murine hippocampus [11]. In this regard, the aim of this work was to dig deeper in the understanding of the alterations mentioned above, which produce a cognitive decrease in memory and depression-like

behavior related to hippocampal damage. Furthermore, we sought to establish whether pro-inflammatory involvement played a relevant role in pathogenesis and elucidate the intracellular mechanism likely to be involved in these events.

Materials and Methods

Animals

NIH male Swiss mice of 4 weeks (approximately 25±3 g) were housed under 12 h-light/12 h-dark cycles. Food and water were provided *ad libitum* and the experimental protocols and euthanasia procedures were reviewed and approved by the Institutional Animal Care and Use Committee of the School of Medicine, Universidad de Buenos Aires, Argentina (Resolution Nº 046/2017). After an acclimatization period, animals were divided into four different groups according to intravenous (i.v.) treatment (the total amount of i.v. solution injected was 100 µl per mouse): Control (saline solution), LPS (800 ng), Stx2 (1 ng) and Stx2+LPS (1 ng and 800 ng, respectively). Stx2 dose was about 60% of the LD50 (1.6 ng per mouse). All the procedures were performed in accordance with the EEC guidelines for the care and use of experimental animals (EEC Council 86/609).

Nose-poke habituation task

The apparatus employed was a hole-board (Ugo Basile Mod. 6650, Comerio, Italy) made of a matte gray Perspex panel (40 cm× 40 cm× 22 cm) which has 16 flush mounted tubes of 3 cm (Figure 1A). Each tube has an infrared emitter and a diametrically opposed receiver connected to an automatic counter to register the number of nose-pokes into the holes. The assay consisted of two stages: training (TR) and retention test (TS). TR was performed 2

days after the i.v. treatment. During TR, each mouse was placed at the center of the

apparatus and the number of nose-pokes was automatically registered for 5 minutes (Fig

1A). During the TS stage, performed 30 minutes after TR, each mouse was placed again on

the apparatus and the nose-pokes were registered for 5 minutes. The apparatus was

carefully cleaned with ethanol 70% to avoid interference by previous animals'

performance. Exploratory activity was expressed as nose-pokes per 5 minutes. Differences

in the number of nose-pokes between TR and TS represent the development of memory. A

more significant difference between TR and TS means better memory formation. Results

were also represented using a memory index derived from the following mathematical

formula:

116
$$\text{Memory Index} = \frac{(\text{number of training visits} - \text{number of test visits})}{\text{number of training visits}}$$

Immunofluorescence assay

Two days after the i.v. treatments mice were anesthetized with sodium pentobarbital (100

mg/kg) and intracardiacally perfused with paraformaldehyde 4% diluted in 0.1 M

phosphate buffer saline (PBS), pH 7.4. The brains were removed from the skulls, post-fixed

overnight at 4 °C with the same fixative solution described above, cryopreserved daily with

solutions with increasing concentrations of sucrose in PBS (10, 20 and 30%), and left

overnight at 4 °C. Brain sections of 20 μm were obtained in a cryostat and stored at -20°C

in a cryopreservation solution (50% PBS, 30% ethylene glycol and 20% glycerol) until the

day of the immunofluorescence assay.

Immunofluorescence for ionized calcium binding adaptor molecule 1 (Iba1) was carried out

to evaluate microglia (MG) reactivity. For this purpose, brain sections were blocked with a

PBS solution containing 0.1% Triton X-100 and 10% goat serum to be incubated with a

monoclonal goat anti-Iba1 antibody overnight at 4 °C (1:250 - Millipore, Temecula, CA,

USA) followed by a donkey anti-goat Alexa Fluor 488 antibody (1:500 - Millipore,

Temecula, CA, USA) for one hour at room temperature. All brain sections were also

incubated with Hoechst 33342 (1:500 - Sigma, St. Louis, MO, USA) for 15 minutes at

room temperature to detect brain cell nuclei. Negative controls were carried out by omitting

the primary antibody. The hippocampal CA1 area (-1.70 and -1.82mm from bregma) was

observed on an Olympus BX50 epifluorescence microscope equipped with a Cool-Snap

digital camera and on an Olympus FV1000 confocal microscope.

Four distinct MG phenotypes immunolocalized in the hippocampal CA1 area were

classified according to morphology [21]: ramified, hyper-ramified, bushy, and ameboid. A

ramified morphology is associated with a resting microglial profile and was identified as a

small soma with multiple long thin processes. The hyper-ramified phenotype was identified

as a larger soma with thicker and branching processes. Bushy MG were also identified as

having a larger soma with thicker but fewer processes. The ameboid morphology was

identified as the largest soma with very few or absence of processes. The proportion of each

type of morphology was measured for each treatment. Microglial activation was quantified

using the following formula [22]:

Microglial activation score = $(0 \times n) + (1 \times n) + (2 \times n) + (3 \times n)$

The numbers 0, 1, 2 and 3 represent arbitrary factors for the four different microglial

morphologies (ramified=0, hyper-ramified=1, bushy=2, and amoeboid=3), and "n"

represents the number of microglial cells with each specific morphology and corresponding

arbitrary factor per micrograph.

Enzyme-linked immunosorbent assay (ELISA)

Animals were sacrificed by decapitation 1 or 2 days after the four i.v. treatments, and the

hippocampi were dissected and then homogenized on ice in PBS containing protease

inhibitor cocktail (Sigma, St. Louis, MO, USA). Next, samples were centrifuged at 10,000

g for 15 minutes at 4 °C. IL-6, TNFα and IL-10 concentrations were immediately

determined in the supernatant of murine hippocampal homogenates using the specific

ELISA with antibodies and standards obtained from BD Biosciences (San Diego, CA,

USA). The assay was performed according to the manufacturers' instructions.

Western blot

Animals were sacrificed by decapitation 2, 6, 12 and 24 hours after the i.v. treatments, and

the hippocampi were dissected and homogenized in ice-cold lysis buffer (pH 7.4)

containing 24 mmol/L HEPES, 1 mmol/L EDTA, 2 mmol/L tetrasodium pyrophosphate, 70

mmol/L sodium fluoride, 1 mmol/L β-glycerophosphate, 1% Triton X-100, 1 mmol/L

PMSF, 10 μg/ml aprotinin and 2 μg/ml leupeptin (Sigma, St. Louis, MO, USA). Protein

concentration in the homogenates was quantitatively determined through the Bradford

assay. Hippocampal samples with equal amounts of protein (40 μg) were separated by

electrophoresis in 10% sodium dodecyl sulfate polyacrylamide gels (SDS-PAGE) and

transferred onto PVDF membrane (Bio-Rad Labs Inc.). Nonspecific binding sites on the

membrane were blocked by incubation with 5% milk in Tris-buffered saline solution

containing 0.1% Tween 20 at room temperature for 1 hour. The membranes were then

incubated with primaries antibodies (1:500) overnight at 4 °C: rabbit anti-pERK1/2, rabbit

anti-ERK1/2, rabbit anti-pIκBα or rabbit anti-IκBα. To avoid inaccuracies in protein

loading, mouse anti-GAPDH was employed as an internal standard. All primary antibodies

were purchased from Cell Signaling Technology, Inc. (Danvers, MA, USA). After 1-hour

rinse, membranes were incubated with horseradish peroxidase-conjugated secondary

antibody at 37 °C for 1 hour. Protein loading was evaluated by stripping and reblotting

membranes with anti-GAPDH antibody (1:1000).

Forced swim test

The four i.v. treated mouse groups described in *Animals* received either saline solution

(without dexamethasone) or dexamethasone (7.5 mg/kg) by intraperitoneal (i.p. 100 µl)

administration once a day for two consecutive days. Day zero was considered the day of the

i.v. and i.p. treatments. Thus, the eight groups were divided as follows: Control with or

without dexamethasone, LPS with or without dexamethasone, Stx2 with or without

dexamethasone and Stx2+LPS with or without dexamethasone. After two days of

treatments, each mouse was placed for 6 minutes in 3-liter beakers containing a volume of

water at 25 °C that measured 15 cm height, which prevented mice from reaching the

bottom of the container. The 6-minute test was recorded, and the time of immobility was

measured only in the last 4 minutes [23]. After this time, each mouse was removed from the

container and placed temporarily in a drying cage with a heat lamp above it. The water was

changed after every session to avoid influence on the next mouse.

Statistical analysis

All experimental data are presented as the mean ± SEM. Particularly, in the forced swim

test, statistical significance was evaluated using two-way analysis of variance (ANOVA).

For all other assays, statistical significance was evaluated using one-way ANOVA followed

by Bonferroni's *post hoc* test (GraphPad Prism 4, GraphPad Software Inc., San Diego, CA,

USA). The criterion for significance was $p < 0.05$ for all experiments.

Results

Stx2+LPS reduced short term memory

The nose-poke habituation task was employed to establish whether the toxins change the

capability of mice to store and evoke short term memories [24]. While no significant

differences were found in exploratory activity among groups during TR (Control, LPS,

Stx2 and Stx2+LPS), mice treated with Control, LPS or Stx2 yielded a significant reduction

in the number of nose-pokes recorded between TR and TS (Fig 1B). These findings suggest

that mice in these groups were able to store and retrieve short-term memory of the

habituation task. Nevertheless, no significant differences were found between TR and TS

for Stx2+LPS-treated mice, which is indicative of amnesia. Consequently, the memory

index was significantly lower in Stx2+LPS-treated mice compared to all other treatments

(Fig 1C) ($p < 0.05$). No differences were observed in the memory index across Control, LPS

and Stx2-treated mice.

Toxins produced MG reactivity in the CA1 hippocampal area

Iba1 is a microglial/macrophagic-specific calcium-binding protein which participates in

membrane ruffling and phagocytosis in activated microglia [25]. An anti-Iba1 antibody was

employed to establish whether the toxins produced activated MG morphology, as this

molecule is upregulated in activated MG states [21, 22, 26].

After 2 days of treatment, the four MG morphological types were observed in the CA1

hippocampal area (Fig 2A). The MG activation score was significantly increased by both

toxins compared to Control (Fig 2B), and maximally increased in Stx2+LPS-treated mice.

No significant differences were found between LPS and Stx2-treated mice. In addition, the

toxins also produced a significant increase in hyper-ramified, bushy and amoeboid MG

activated morphologies as compared to Control (Fig 2C). In contrast, the toxins produced a

significant decrease in ramified morphology (Fig 2C).

Toxins increased pro-inflammatory cytokines and decreased anti-inflammatory

cytokines in the hippocampus

In order to establish whether these toxins altered the pro-inflammatory and anti-

inflammatory cytokine profile in the mouse hippocampus treated with LPS and/or Stx2, an

ELISA was performed to measure pro-inflammatory cytokines IL-6 and TNFα and anti-

inflammatory cytokine IL-10. The expressions of IL-6 and TNFα were both significantly

increased after 1 day of treatment in all toxin-treated mice (Fig 3A, B). On the other hand,

IL-10 was significantly decreased after 1 day of treatment only in Stx2+LPS-treated mice

(Fig 3C), but significantly decreased after 2 days of treatment in all toxin-treated mice (Fig

3D).

The pro-inflammatory profile induced by Stx2 was pIkB-dependent but ERK-

independent

Given that toxins induced the expression of pro-inflammatory cytokines, we next studied

the intracellular signaling involved in these processes. For this purpose, the ERK and IkB

phosphorylation profiles were determined by Western blot following treatment with

Stx2+LPS between 2 and 24 hours. As shown in figure 4A, upper panel, a significant

reduction in phosphorylated ERK (pERK) was observed after 12 hours of treatment, while

pIkB was increased after 2 hours (Fig 4B, upper panel).

To investigate whether the response elicited by Stx2+LPS may be due to a synergistic

effect of both compounds, we evaluated the individual effect of LPS or Stx2 on ERK and

IkB phosphorylation. After 12 hours, Stx2 but not LPS alone induced a decrease in ERK

phosphorylation (Fig 4A, lower panel), which suggests that the reduction in ERK activation

induced by Stx2+LPS resulted from a predominant effect of Stx2. Regarding Ikb, 2 hours

of treatment with Stx2 or LPS alone induced the same increase in IkB phosphorylation as

combined Stx2+LPS treatment (Fig 4B, lower panel).

Stx2 and Stx2+LPS induced depression-like behavior, which was attenuated by

dexamethasone

To evaluate whether the toxins produced depression-like behavior and whether this

behavior may involve an inflammatory component, toxin-treated mice were treated with

corticosteroid dexamethasone and then subjected to the forced swimming test. Immobility

time (considered when the animals were restricted to floating movements [23] was

significantly higher in Stx2+LPS-treated mice as compared to Stx2, and also significantly

increased in these two groups compared to LPS and Control (Fig 5). Treatment with

dexamethasone significantly reduced immobility time in Stx2 and Stx2+LPS-treated mice

(Fig 5). No significant differences were found between the i.p. treatments in Control and

LPS-treated mice.

Discussion

In vivo studies and clinical reports have shown the importance of neuroinflammation in the

pathogenesis of Stx2-induced encephalopathy [10, 16, 27, 28]. However, none of these

studies have shown the molecular pathways through which the inflammation is produced.

The current report presents for the first time an introduction to the molecular mechanisms

triggered by Stx2 in the murine brain hippocampus. Furthermore, it connects the

inflammatory process triggered by Stx2 at the molecular level with behavioral changes,

which confirms our previous results showing damage in the neurovascular unit produced by

Stx2 in the CA1 hippocampal area [11].

The alterations in memory index and depression-like behavior produced by the toxins in the

nose-poke habituation task and the forced swimming test, respectively, may be related with

damage in hippocampal neurons and correlate with the cognitive imbalance reported in

patients [17]. Considering previous reports [9-11], Stx2 may damage hippocampal neurons

in at least two ways: first, Stx2 binds to the canonical Gb3 receptor localized in CA1

hippocampal neurons; and, second, indirectly by an inflammatory process with MG playing

a fundamental role [10]. Indeed, we have recently reported in an *in vitro* model that Stx2 is

up-taken by MG through its receptor Gb3, which produces an increase in MG metabolism,

phagocytic capacity and pro-inflammatory cytokine release [29].

MG are mononuclear phagocytes resident in CNS parenchyma. MG are important sentinels

responsible for CNS homeostasis which play an important role in synaptic organization,

control of neuronal excitability, phagocytic removal of debris and trophic support leading

to brain protection and repair [30, 31]. As primary innate immune cells, MG actively move

through the CNS microenvironment while they sense disturbances in homeostasis by

extending and retracting their fine processes [31]. Once they find some homeostatic

imbalance through their wide range of receptors, MG change their morphology from a

highly ramified small cell soma (traditionally called "resting") to a variety of other distinct

morphologies (traditionally called "activated"), ultimately becoming a poorly ramified

amoeboid cell soma. These cells are also highly specialized in cytokine secretion [21, 22].

Stx2 and LPS produced a significant increase in the secretory MG morphology, with more

amoeboid than ramified cells. Furthermore, this event was accompanied by a significant

increase in pro-inflammatory IL-6 and TNFα and a significant reduction in anti-

inflammatory IL-10 secretory profile. Traditionally, cytokine secretion has been classified

according to its pro or anti-inflammatory profile. The M1 profile is responsible for the

secretion of pro-inflammatory cytokines like IL-6 and TNFα. In contrast, the M2 profile is

responsible for the secretion of anti-inflammatory cytokines such as IL-10 [32-34]. In

addition, secretory ameboid MG morphology may also be an important oxidative source of

reactive oxygen species (ROS) and nitric oxide (NO) radical derived products [35], as we

have previously shown in the current model of Stx2-produced encephalopathy [11].

Cytokine synthesis is regulated by many intracellular signaling pathways triggered by

different kinds of receptors that respond to different stimuli [36, 37]. For example, the

binding of LPS to toll-like receptor 4 (TLR4) leads to the activation of the intracellular

mitogen-activated protein kinase (MAPK) pathway and NF-κB nuclear translocation,

which results in the secretion of pro-inflammatory cytokines [38, 39]. However,

information regarding the intracellular signaling pathway triggered by Stx2 is very scarce,

and particularly much less is known about it in brain parenchymal cells.

To our knowledge, all that is known so far about damage produced by Stx via the MAPK

pathway is the activation of p38, which induces IL-8 secretion in the human colonic

epithelial cell line [40], IL-1 secretion by human macrophages [41] and TNF production by

a human adenocarcinoma-derived renal tubular epithelial cell line [42]. In addition, p38

activation has been also found partially responsible for Stx-induced cell death in Vero cells

[43] and in intestinal epithelial cells [44]. On the other hand, the inhibition of p38 is a

determinant of TNF up-regulation from Stx toxicity in brain microvascular endothelial cells

[45]. Furthermore, signaling cascades including JNK and p38 molecules have been related

to cell death by ribotoxic stress leading to apoptosis in intestinal epithelial cells [44].

MAPKs are a family of proteins which, in response to the activation of cell surface

receptors, phosphorylate specific serine or threonine residues preceded by a proline residue

on their substrates. Extracellular signals such as hormones, cytokines or growth factors

activate the MAPK cascade which consists in a linear signaling triad comprising a MAPK

kinase kinase (MAPKKK or MAP3K) and a MAPK kinase (MAPKK or MAP2K), ending

in the phosphorylation and activation of MAPKs, which include the extracellular signal

regulated kinases 1 and 2 (ERK1/2), p38 isoforms (α, β, γ, and δ) and c-Jun N-terminal

kinases (JNK1, JNK2, and JNK3). The activation of MAPKs culminates with the

phosphorylation of many cytosolic and nuclear substrates such as downstream kinases,

phosphatases or transcription factors. The dysregulation of this intracellular cascade is

implicated in many neurological disorders, including depression and Alzheimer's disease

[46]. Similar dysregulation was also found in our present work and could be compared to

these neurological disorders.

According to the present data, in our encephalopathy murine model, the deleterious effect

produced by Stx2 together with LPS [10, 11, 16, 29] suggests a potentially synergistic

action rather than an additive one (Fig 6). While both toxins induced the activation and

translocation of NF-κB to the nucleus, they performed it independently through two distinct

pathways: LPS activated the ERK1/2 pathway, and Stx2 activated an ERK1/2-independent

pathway. Furthermore, both pathways also produced the expression of genes related to a

pro-inflammatory state such as TNFα and IL-6 and reduced the expression of those

associated to anti-inflammatory cytokines such IL-10. Although further experiments are

necessary to determine whether p38, JNK or a third unknown pathway may be regulated by

Stx2 concerning NF-κB activation, it may be concluded that the activation of these

molecular cascades may also be implicated in the activation of MG and concomitant

cognitive deterioration observed in the current mouse model. Moreover, the fact that this

deterioration has also been reported in patients suggests that the intracellular mechanisms

studied in this work may also be at play in clinical scenarios. Since there is no consensus

therapy currently available to treat STEC-derived encephalopathy, it is essential to

elucidate the molecular mechanisms underlying the pathophysiological process of

encephalopathy produced by Stx2 to propose therapeutic strategies which may efficiently

protect the brain parenchyma [47]. Accordingly, our present work shows that the use of

drugs such as dexamethasone or those blocking the cascade by preventing NF-kB

translocation to the nucleus may serve as effective neuroprotectors with potentially

beneficial use in the clinic.

Funding

347 This work was supported by Agencia Nacional de Promoción Científica y Tecnológica

348 (ANPCyT) (PICT-2016-1175) and Universidad de Buenos Aires (UBACyT)

349 (20020160100135BA), Argentina (JG), and (ANPCyT) 2016-0129 & 2016-0803 (FC).

Acknowledgments

351 The authors are especially grateful to German Nicolás La Iacona for his technical assistance

352 in using the confocal microscope. We are also grateful to María Marta Rancez for her

353 special dedication and technical assistance.

References

1. Fakhouri, F., et al., *Haemolytic uraemic syndrome.* Lancet, 2017. **390**(10095): p. 681-696.

2. Karmali, M.A., et al., *The association between idiopathic hemolytic uremic syndrome and*
infection by verotoxin-producing Escherichia coli. J Infect Dis, 1985. **151**(5): p. 775-82.

3. Karmali, M.A., et al., *Sporadic cases of haemolytic-uraemic syndrome associated with*
faecal cytotoxin and cytotoxin-producing Escherichia coli in stools. Lancet, 1983. **1**(8325):
p. 619-20.

4. Rivas M, P.N., Luchessi PM, Masana M, Torres AG, *Diarrheogenic Escherichia coli in*
Argentina. Pathogenic Escherichia coli in Latin America. 2010. 142-161.

5. Upadhyaya, K., et al., *The importance of nonrenal involvement in hemolytic-uremic*
syndrome. Pediatrics, 1980. **65**(1): p. 115-20.

6. Sheth, K.J., H.M. Swick, and N. Haworth, *Neurological involvement in hemolytic-uremic*
syndrome. Ann Neurol, 1986. **19**(1): p. 90-3.

7. Hahn, J.S., et al., *Neurological complications of hemolytic-uremic syndrome.* J Child Neurol,
1989. **4**(2): p. 108-13.

8. Torres, A.G., et al., *Recent Advances in Shiga Toxin-Producing Escherichia coli Research in*
Latin America. Microorganisms, 2018. **6**(4).

9.   Obata, F. and T. Obrig, *Distribution of Gb(3) Immunoreactivity in the Mouse Central*
Nervous System. Toxins (Basel), 2010. **2**(8): p. 1997-2006.

10. Pinto, A., et al., *Dexamethasone prevents motor deficits and neurovascular damage*
produced by shiga toxin 2 and lipopolysaccharide in the mouse striatum. Neuroscience,
2017. **344**: p. 25-38.

11. Berdasco, C., et al., *Shiga toxin 2 from enterohemorrhagic Escherichia coli induces reactive*
glial cells and neurovascular disarrangements including edema and lipid peroxidation in
the murine brain hippocampus. J Biomed Sci, 2019. **26**(1): p. 16.

12. Obata, F. and T. Obrig, *Role of Shiga/Vero toxins in pathogenesis.* Microbiol Spectr, 2014.
2(3).

13. Matthies, J., et al., *Extrarenal Manifestations in Shigatoxin-associated Haemolytic Uremic*
Syndrome. Klin Padiatr, 2016. **228**(4): p. 181-8.

14. Hall, G., S. Kurosawa, and D.J. Stearns-Kurosawa, *Shiga Toxin Therapeutics: Beyond*
Neutralization. Toxins (Basel), 2017. **9**(9).

15. Zhang, H., et al., *Bacterial lipoprotein and lipopolysaccharide act synergistically to induce*
lethal shock and proinflammatory cytokine production. J Immunol, 1997. **159**(10): p. 4868-
78.

16. Pinto, A., et al., *Dexamethasone rescues neurovascular unit integrity from cell damage*
caused by systemic administration of shiga toxin 2 and lipopolysaccharide in mice motor
cortex. PLoS One, 2013. **8**(7): p. e70020.

17. Weissenborn, K., et al., *Neurologic manifestations of E coli infection-induced hemolytic-*
uremic syndrome in adults. Neurology, 2012. **79**(14): p. 1466-73.

18. Magnus, T., et al., *The neurological syndrome in adults during the 2011 northern German*
E. coli serotype O104:H4 outbreak. Brain, 2012. **135**(Pt 6): p. 1850-9.

19. Schuppner, R., et al., *Neurological Sequelae in Adults After E coli O104: H4 Infection-*
Induced Hemolytic-Uremic Syndrome. Medicine (Baltimore), 2016. **95**(6): p. e2337.

20. Garg, A.X., et al., *Long-term renal prognosis of diarrhea-associated hemolytic uremic*
syndrome: a systematic review, meta-analysis, and meta-regression. JAMA, 2003. **290**(10):
p. 1360-70.

21. Crews, F.T. and R.P. Vetreno, *Mechanisms of neuroimmune gene induction in alcoholism.*
Psychopharmacology (Berl), 2016. **233**(9): p. 1543-57.

22. Pomilio, C., et al., *Glial alterations from early to late stages in a model of Alzheimer's*
disease: Evidence of autophagy involvement in Abeta internalization. Hippocampus, 2016.
26(2): p. 194-210.

23. Yankelevitch-Yahav, R., et al., *The forced swim test as a model of depressive-like behavior.*
J Vis Exp, 2015(97).

24. Kuc, K.A., et al., *Holeboard discrimination learning in mice.* Genes Brain Behav, 2006. **5**(4):
p. 355-63.

25. Kanazawa, H., et al., *Macrophage/microglia-specific protein Iba1 enhances membrane*
ruffling and Rac activation via phospholipase C-gamma -dependent pathway. J Biol Chem,
2002. **277**(22): p. 20026-32.

26. Ito, D., et al., *Microglia-specific localisation of a novel calcium binding protein, Iba1.* Brain
Res Mol Brain Res, 1998. **57**(1): p. 1-9.

27. Kioka, N., et al., *Chemokine expression in human astrocytes in response to shiga toxin 2.* Int
J Inflam, 2012. **2012**: p. 135803.

28. Hosaka, T., K. Nakamagoe, and A. Tamaoka, *Hemolytic Uremic Syndrome-associated*
Encephalopathy Successfully Treated with Corticosteroids. Intern Med, 2017. **56**(21): p.
2937-2941.

29. Berdasco, C., et al., *Environmental Cues Modulate Microglial Cell Behavior Upon Shiga*
Toxin 2 From Enterohemorrhagic Escherichia coli Exposure. Front Cell Infect Microbiol,
2019. **9**: p. 442.

30. Denes, A., et al., *Proliferating resident microglia after focal cerebral ischaemia in mice.* J
Cereb Blood Flow Metab, 2007. **27**(12): p. 1941-53.

31. Nimmerjahn, A., F. Kirchhoff, and F. Helmchen, *Resting microglial cells are highly dynamic*
surveillants of brain parenchyma in vivo. Science, 2005. **308**(5726): p. 1314-8.

32. Orihuela, R., C.A. McPherson, and G.J. Harry, *Microglial M1/M2 polarization and metabolic*
states. Br J Pharmacol, 2016. **173**(4): p. 649-65.

33. Martinez, F.O. and S. Gordon, *The M1 and M2 paradigm of macrophage activation: time*
for reassessment. F1000Prime Rep, 2014. **6**: p. 13.

34. Cherry, J.D., J.A. Olschowka, and M.K. O'Banion, *Neuroinflammation and M2 microglia: the*
good, the bad, and the inflamed. J Neuroinflammation, 2014. **11**: p. 98.

35. Torre-Fuentes, L., et al., *Experimental models of demyelination and remyelination.*
Neurologia, 2020. **35**(1): p. 32-39.

36. Cavaillon, J.M., *Cytokines and macrophages.* Biomed Pharmacother, 1994. **48**(10): p. 445-
53.

37. Arango Duque, G. and A. Descoteaux, *Macrophage cytokines: involvement in immunity*
and infectious diseases. Front Immunol, 2014. **5**: p. 491.

38. Schnare, M., et al., *Toll-like receptors control activation of adaptive immune responses.* Nat
Immunol, 2001. **2**(10): p. 947-50.

39. Sabroe, I., et al., *Toll-like receptor (TLR)2 and TLR4 in human peripheral blood*
granulocytes: a critical role for monocytes in leukocyte lipopolysaccharide responses. J
Immunol, 2002. **168**(9): p. 4701-10.

40. Thorpe, C.M., et al., *Shiga toxins stimulate secretion of interleukin-8 from intestinal*
epithelial cells. Infect Immun, 1999. **67**(11): p. 5985-93.

41. Foster, G.H. and V.L. Tesh, *Shiga toxin 1-induced activation of c-Jun NH(2)-terminal kinase*
and p38 in the human monocytic cell line THP-1: possible involvement in the production of
TNF-alpha. J Leukoc Biol, 2002. **71**(1): p. 107-14.

42. Nakamura, A., et al., *Activation of beta(2)-adrenoceptor prevents shiga toxin 2-induced*
TNF-alpha gene transcription. J Am Soc Nephrol, 2001. **12**(11): p. 2288-99.

43. Ikeda, M., et al., *Shiga toxin activates p38 MAP kinase through cellular Ca(2+) increase in*
Vero cells. FEBS Lett, 2000. **485**(1): p. 94-8.

44. Smith, W.E., et al., *Shiga toxin 1 triggers a ribotoxic stress response leading to p38 and JNK*
activation and induction of apoptosis in intestinal epithelial cells. Infect Immun, 2003.
71(3): p. 1497-504.

45. Stricklett, P.K., A.K. Hughes, and D.E. Kohan, *Inhibition of p38 mitogen-activated protein*
kinase ameliorates cytokine up-regulated shigatoxin-1 toxicity in human brain
microvascular endothelial cells. J Infect Dis, 2005. **191**(3): p. 461-71.

46. Ghose, R., *Nature of the Pre-Chemistry Ensemble in Mitogen-Activated Protein Kinases.* J
Mol Biol, 2019. **431**(2): p. 145-157.

47. Goldstein, J., K. Nunez-Goluboay, and A. Pinto, *Therapeutic Strategies to Protect the*
Central Nervous System against Shiga Toxin from Enterohemorrhagic Escherichia coli. Curr
Neuropharmacol, 2020.

Figure legends

Figure 1. Short term memory. A: the hole-board apparatus (left), representative nose-
poke assay (right). B and C: tests measured after 2 days of treatment. TR: training test; TS:
retention test. Different letters (a, b) represent significant differences. Data were analyzed
by non-parametric one-way ANOVA, Bonferroni's *post hoc* test, $p<0.05$, n=15.

Figure 2. CA1 hippocampal MG response to toxins. A: Iba1 immunofluorescence of
representative images from all distinct morphological MG types. B: microglial activation
score after 2 days of treatment. Each letter (a, b, c) represents significant differences across
each treatment. C: percentage of the microglial morphology in each treatment. Asterisks (*)
represent significant differences across morphologies from all toxin-treated mice as
compared to Control. Data from figures B and C were analyzed by one-way ANOVA,
Bonferroni's *post hoc* test, $p<0.05$, n=4.

Figure 3. Pro- and anti-inflammatory cytokine measurements in the murine
hippocampus. A: IL-6 after 1 day of treatment. B: TNFα after 1 day of treatment. C: IL-10
after 1 day of treatment. D: IL-10 after 2 days of treatment. Each letter (a, b) represents
significant differences across treatments in all figures. Data were analyzed by one-way
ANOVA, Bonferroni's *post hoc* test, $p<0.05$, n=8.

Figure 4. Changes in intracellular signaling. A, upper panel: Western blot of pERK 1/2

expression after 2, 6, 12 and 24 hours of Control or Stx2+LPS treatments; A, lower panel:

Western blot of pERK 1/2 after 12 hours of the four treatments. B, upper panel: Western

blot of pIκBα after 2, 6, 12 and 24 hours of Control or Stx2+LPS treatments; B, lower

panel: Western blot of pIκBα after 2 hours of the four treatments. Each letter (a, b and c)

represents significant differences across treatments. All data were analyzed by one-way

ANOVA, Bonferroni´s *post hoc* test, p<0.05, n=8.

Figure 5. Depression-like behavior and the effects of dexamethasone. The effect of

toxins and dexamethasone on mouse behavior after 2 days of treatment. Each letter (a, b)

represents significant differences between i.p. treatments with or without dexamethasone. *[1]

represents a significant difference between Stx2-treated mice without dexamethasone and

the other treated mice without dexamethasone. *[2] represents significant differences between

Stx2+LPS-treated mice without dexamethasone and the other treated mice without

dexamethasone. Data were analyzed by non-parametric two-way ANOVA, Bonferroni's

post hoc test, p<0.05, n=12.

Figure 6: Suggested model of Stx2 and LPS intracellular deleterious actions in the

murine hippocampus: The toxins reach the hippocampus crossing the blood-brain barrier

that activate microglial cells. They activate NF-κB translocation to the nucleus through two

distinct pathways engaging inflammatory processes and cognitive deficits. Astrocyte

(green); MG cells (violet); neuron (yellow).

Figure 1

A

Microglial morphology

Ramified   Hiper-ramified   Bushy   Ameboid

B

Microglial activation score (A.U.)

Control   LPS   Stx2   LPS+Stx2

C

Ramified   Bushy
Hyper-ramified   Ameboid

Microglial morphology (%)

Control   LPS   Stx2   LPS+Stx2

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6