The recombination landscape and multiple QTL mapping in a *Solanum tuberosum* cv. 'Atlantic'-derived F$_1$ population

Guilherme da Silva Pereira[1,2*], Marcelo Mollinari[1,2], Mitchell J. Schumann[1], Mark E.

Clough[1,3], Zhao-Bang Zeng[1,2], G. Craig Yencho[1*]

[1]Department of Horticultural Science, North Carolina State University, Raleigh, NC, USA

[2]Bioinformatics Research Center, North Carolina State University, Raleigh, NC, USA

[3]Vernon G. James Research and Extension Center, Department of Horticultural Science, North

Carolina State University, Plymouth, NC, USA

* Correspondence:

Guilherme da Silva Pereira

University of São Paulo

Luiz de Queiroz College of Agriculture

Av. Pádua Dias, 11

Piracicaba, SP, Brazil 13418-900

email: gdasilv@ncsu.edu

G. Craig Yencho

North Carolina State University

Department of Horticultural Science

Room 214-A Kilgore Hall, Box 7609

2721 Founders Drive

Raleigh, NC, USA 27695-7609

craig_yencho@ncsu.edu

Current addresses:

Guilherme da Silva Pereira: Department of Genetics, Luiz de Queiroz College of Agriculture,

University of São Paulo, Piracicaba, Brazil

Mitchell J. Schumann: Bayer Crop Science, Chesterfield, MO, USA

Word counts:

Abstract = 215, Introduction = 1040, Materials and Methods = 1770, Results = 1763, Discussion =

2220, Final Considerations = 410, Total = 7203

Figures = 4 (color), Tables = 1, Supporting information = 23

Keywords: linkage map, mixed model, multiple interval mapping, polyploid, quality, tetraploid

potato, yield.

Abstract

There are many challenges involved with the genetic analyses of autopolyploid species, such as the

tetraploid potato, *Solanum tuberosum* ($2n = 4x = 48$). The development of new analytical methods

has made it valuable to re-analyze an F_1 population ($n = 156$) derived from a cross involving

'Atlantic', a widely grown chipping variety in the USA. A fully integrated genetic map with 4,285

single nucleotide polymorphisms, spanning 1,630 cM, was constructed with MAPpoly software. We

observed that bivalent configurations were the most abundant ones (51.0~72.4% depending on parent

and linkage group), though multivalent configurations were also observed (2.2~39.2%). Seven traits

were evaluated over four years (2006-8 and 2014) and quantitative trait loci (QTL) mapping was

carried out using QTLpoly software. Based on a multiple-QTL model approach, we detected 21 QTL

for 15 out of 27 trait-year combination phenotypes. A hotspot on linkage group 5 was identified as

QTL for maturity, plant yield, specific gravity and internal heat necrosis resistance over different

years were co-located. Additional QTL for specific gravity and dry matter were detected with

maturity-corrected phenotypes. Among the genes around QTL peaks, we found those on

chromosome 5 that have been previously implicated in maturity (*StCDF1*) and tuber formation

(*POTH1*). These analyses have the potential to provide insights into the biology and breeding of

tetraploid potato and other autopolyploid species.

1   Introduction

Polyploids can be classified based on the meiotic pairing dynamics of their duplicated sets of

chromosomes. Allopolyploids are formed by the genomes of more than one ancestor species, which

generally implies in disomic inheritance, where bivalent and preferential chromosome pairing is

observed. Autopolyploids are formed by the genome duplication of a single ancestor species, such

that polysomic inheritance is expected, with both bivalent and multivalent random chromosome

pairing (Comai, 2005; Zielinski and Scheid, 2012). In autopolyploids, multivalent formation is one of

the conditions for double reduction, a phenomenon in which sister chromatid segments end up in the

same gamete (Huang *et al.*, 2019). In order to investigate these complicated inheritance patterns,

genetic mapping studies in autopolyploid species still need to deal with the molecular marker system

limitation. Even though up to four different alleles (e.g. *abcd*) are expected in a locus of an

outcrossing, autotetraploid individual, biallelic single nucleotide polymorphisms (SNPs) can be

characterized, at best, by the dosage of an arbitrary allele, say *A*, as nulliplex (*aaaa*), simplex (*Aaaa*),

duplex (*AAaa*), triplex (*AAAa*) or quadriplex (*AAAA*).

Several polyploid species are agriculturally important (Bennett, 2004). Particularly, potato is the third

most cultivated food crop in the world after rice and wheat, with more than 17.6 million hectares

cultivated worldwide in 2018, and a production exceeding 368.2 million tons (FAO, 2020). Yet

advancement in the breeding of new varieties is still relatively slow. 'Russet Burbank', which still

accounts for a large amount of the acreage planted in the United States, was developed in the early

1900s (Bethke *et al.*, 2014), and 'Atlantic', a major chipping variety grown in the country, was

originally released in 1976 (Webb *et al.*, 1978). One of the reasons of such relatively slow breeding

process is the relative complexity of the potato genome. Cultivated potato, *Solanum tuberosum*, is a

highly heterozygous autotetraploid ($2n = 4x = 48$) outcrossing crop, which makes it very difficult

to combine potentially beneficial characteristics through conventional breeding methods. In fact,

many agronomic traits are polygenic, and show continuous distributions (Ghislain and Douches,

2020), where genotype-by-environment is usually important and need to be assessed.

The genome complexity has also impacted on the correct assessment of dosage from molecular

markers, impeding proper linkage map construction and quantitative trait loci (QTL) mapping.

Research has been conducted over the years to develop molecular techniques to facilitate potato

breeding. However, since the first molecular genetic maps of potato were published (reviewed by

Mann *et al.*, 2011), very few breeding programs have begun to employ marker-assisted selection

routinely (Slater *et al.*, 2014). Some of the exceptions to this are for genetically simpler traits such as

potato cyst nematode resistance (Schultz *et al.*, 2012) and potato virus Y (Hämäläinen *et al.*, 1997)

and other disease resistance loci (Ramakrishnan *et al.*, 2015). Genomics-assisted breeding for more

complex traits such as yield are still elusive, and more studies on the genetic architecture of these

traits as well as on genome-wide prediction (Endelman *et al.*, 2018) are needed.

The early linkage maps constructed for potato have incorporated qualitatively scored molecular

markers (Mann *et al.*, 2011). However, this traditional presence-absence based scoring limited not

only the ability to explore the full range of segregating markers, but also to build integrated genetic

maps for full-sib populations (Luo *et al.*, 2004). Advances in molecular technologies have been made

available to breeders with the publication of the diploid potato genome sequence (Xu *et al.*, 2011)

and the subsequent development and release of the Illumina Infinium® 8,303 Potato Array (Felcher *et*

al., 2012) through the USDA-NIFA Solanaceae Coordinated Agricultural Project (SolCAP). The

original potato array, which was subsequently upgraded to a 12k form, was made up of 8,303 SNPs

and has been shown to give very good coverage of the published potato genome (Felcher *et al.*, 2012;

Sharma *et al.*, 2013). Utilizing Illumina's technology of dual colored fluorescents to label the

different nucleotides allows for scoring allele dosage (Schmitz Carley *et al.*, 2017; Zych *et al.*, 2019).

Along with the development of an array that uses dosage-sensitive SNPs, new analytical methods

have recently been developed to effectively utilize this information in reconstructing linkage maps

(Hackett *et al.*, 2014; Bourke *et al.*, 2018; Mollinari and Garcia, 2019) and mapping QTL (Hackett *et*

al., 2016; Chen *et al.*, 2018) in autotetraploid species.

Novel linkage and QTL mapping approaches (Mollinari *et al.*, 2020; Pereira *et al.*, 2020) have

enabled the exploration of the meiotic configurations and the detection of multiple QTL for traits of

interest, offering breeding opportunities and insights into the biology of polyploid species.

Particularly, Mollinari *et al.* (2020) presented a detailed characterization of the polysomic inheritance

in an even more complex autopolyploid species, the hexaploid sweetpotato, *Ipomoea batatas* ($2n =$

$6x = 90$). In tetraploid potato, double reduction has been documented for several mapping

populations (e.g. Bourke *et al.*, 2015), but these studies lack a comprehensive assessment of

multivalent formation and preferential chromosome pairing. As haplotypes and thus the patterns of

genome transmission from parents to progeny had been inferred, Pereira *et al.* (2020) were able to

estimate additive relationship based on identity-by-descent along the genetic map. Such an

information was ultimately used in a random-effect model approach, facilitating multiple QTL

mapping. In fact, single-QTL models used so far may have hindered the discovery of putative loci

underlying the variation of agriculturally important traits in the species. Comparatively, multiple-

QTL models have been shown to have increased detection power and to provide the statistical basis

for separation of linked QTL (Pereira *et al.*, 2020).

The advancement of molecular technologies and analytical methods, coupled with newly collected

data, have made it valuable to re-analyze the B2721 potato mapping population, derived from the

‘Atlantic’ cultivar (McCord *et al.*, 2011a, 2011b; Schumann *et al.*, 2017). The aims of this study were

(*i*) to perform SNP dosage calling and create a fully phased, integrated linkage map in order to study

the meiotic configurations in the B2721 mapping population, and (*ii*) to carry out multiple QTL-

based analyses for yield, foliage maturity, dry matter, specific gravity, skin texture and internal heat

necrosis resistance evaluated over four years. This study has allowed us to infer the tetraploid

inheritance mechanisms from a linkage analysis perspective, and to better understand these traits at

the genetic level in order to bring breeders closer to marker-assisted selection.

2   Material and Methods

2.1   Mapping population and field experiment

The B2721 family consisted of 156 progenies and was created by crossing ‘Atlantic’ × B1829-5. This

bi-parental mapping population was chosen based on its segregation for internal heat necrosis (IHN),

a non-pathogenic physiological disorder characterized by brownish spots in the tuber parenchyma

(Yencho *et al.*, 2008). ‘Atlantic’ is susceptible to IHN, whereas B1829-5, an advanced round white

clone from the USDA-ARS Beltsville potato breeding program, is resistant. The B2721 population

also segregates for a wide range of agronomic traits of interest, and these were our focus in this work,

although IHN related traits will also be considered. In this study, we used B2721 data previously

collected in 2006, 2007 and 2008 (McCord *et al.*, 2011b, 2011a) as well as additional data collected

in 2014 (Schumann *et al.*, 2017).

The field designs for years 2006, 2007 and 2008 were described by McCord *et al.* (2011a,b). Briefly,

in 2006, the population was planted in an unreplicated trial with six plants per clone, whereas in 2007

and 2008, the population was planted in two replications with 10 plants per plot in a randomized

complete block design, the same design also used in 2014. All experiments were carried out at the

Tidewater Research Station (35°52'20" N, 76°39'33" W) in Plymouth, North Carolina, in a

Portsmouth fine sandy loam. In 2014, untreated seed pieces were planted on March 21 and the

harvest date was July 15 (a total of 116 days). Standard crop production methods used for potato

cultivation in eastern North Carolina were used for all experiments. At harvest, plots were dug using

a chain digger and harvested by hand. Tubers were then washed, culled for excessive rots and

defects, and phenotype measurements were obtained.

2.2   Phenotypic data and analyses

Phenotyping for all IHN-related and agronomic traits in 2006, 2007 and 2008 was described

previously (McCord *et al.*, 2011b). In 2014, in addition to IHN incidence (NI) and IHN severity (NS)

(Schumann *et al.*, 2017), the B2721 population was evaluated for five other traits: plant yield (PY),

foliage maturity (FM), dry matter (DM), specific gravity (SG) and skin texture (ST). PY was

measured as total plot weight minus the tubers culled divided by 10 (number of plants per plot) in

kg/plant. FM was measured as the area under the senescence progress curve (Shaner and Finney,

1977), based on foliage ratings taken on June 9, June 25 and July 2, using a scale of 0 (0%

yellowing) to 5 (100% senesced) with half point increments. For DM determinations, tubers were

quartered from stem end to bud end using an Easy Wedger, Model N55550-4 (NEMCO Inc.,

Hicksville OH). For each plot, one quarter from four different tubers was placed into plastic Whirl-

Pak® bags. Samples were then weighed, frozen at –20 °C, lyophilized, and weighed again. DM was

calculated as the proportion between dry and fresh weights for each sample, in percentage. SG was

calculated using the formula $SG = w_{air}/(w_{air} - w_{water})$ by using the air weight (w_{air}) and water

weight (w_{water}) for each plot. ST was rated on a 0 (smooth) to 3 (russeted) scale. Finally, NI was

calculated as the proportion of tubers showing any sign of IHN over all evaluated tubers, and NS was

rated on a 9 (no IHN signs) to 1 (completely necrotic) scale and averaged out for all evaluated tubers

(Schumann *et al.*, 2017).

In order to estimate mean-basis broad-sense heritability (H^2) (Holland *et al.*, 2003), a joint analysis

with data from 2007, 2008 and 2014 (replicated trials) was carried out, where years, blocks within

years, genotypes and genotype-by-year interactions were treated as random effects. Except for 2006

data (single observation), adjusted means were obtained separately for each trait-year combination

based on a mixed model with genotypes and blocks as fixed and random effects, respectively, plus

the random residual error using ASReml-R v. 4.1.0 (Butler *et al.*, 2018). The phenotype means were

named after a trait abbreviation followed by two-digit year (trait-year combination, e.g. PY06) and

later used for QTL mapping purposes. Progeny variation within each phenotype was explored with

boxplots from R package ggplot2 (Wickham, 2016). Pearson's correlations were computed using

Rpackage psych v. 1.9.12 (Revelle, 2018), and graphical visualization through network and

correlation plots were obtained using R packages corrr v. 0.4.0 (Ruiz *et al.*, 2019) and corrplot v.

0.84 (Wei *et al.*, 2017). As an approach to investigate genotype-by-environment (GE) interaction,

genotype (G) plus GE interaction (GGE) biplot analysis (Yan and Kang, 2003) was carried out using

R package GGEBiplots v. 0.1.1 (Dumble *et al.*, 2017) based on the standardized means. All analyses

were based on R software v. 3.6.0 (R Core Team, 2020).

2.3   Genotypic data, dosage calling and linkage map construction

Genotyping was performed using the Illumina Infinium® 8,303 Potato Array (Felcher *et al.*, 2012).

The intensity of each fluorescent labeled nucleotide was read using the Illumina iScan Reader and

imported into GenomeStudio. The ordered pair of normalized intensities (x, y) from the two allelic

variants was transformed into polar coordinates $\theta = \frac{2}{\pi} \operatorname{atan}\left(\frac{y}{x}\right)$ and $r = \sqrt{x^2 + y^2}$ to proceed with

the dosage calling of the parents and offspring using the R package ClusterCall (Schmitz Carley *et*

al., 2017). Uninformative markers as well as those resulting in unclassified parents were filtered out

resulting in 5,599 SNPs.

Allele dosage from these markers were imported into MAPpoly v. 0.1.0 (Mollinari and Garcia, 2019)

together with their respective chromosome positions (*S. tuberosum* genome v. 4.0.3). After testing for

segregation distortion ($\sim 8.9 \times 10^{-6}$ using Bonferroni correction) and filtering out redundant

markers, the pairwise recombination fraction of the 4,812 SNPs (~11.6 million pairs) were computed

for all possible linkage phases and the most likely configuration was selected. Markers were

assembled into 12 linkage groups (LGs) using the Unweighted Pair Group Method with Arithmetic

Mean (UPGMA) algorithm. Since 91.2% of markers within the groups coincided with their

respective chromosomes, groups were formed using exclusively the genome information. Markers

were ordered according to the reference genome (Xu *et al.*, 2011) and phased using function

'est_rf_hmm_sequential'. Briefly, this function sequentially inserts markers into the map using two-

point information to eliminate unlikely linkage phases (logarithm of the odds, LOD < 10.0), and for

the remaining configurations, the multipoint LOD score is used (LOD < 10.0). If the insertion of a

marker still results in less than 60 linkage phase configurations, a next marker was inserted and tested

201 in the multiple configurations until the map is completed. In the final step, the multipoint

202 recombination fraction was estimated allowing a global genotyping error of 5%.

203 Next, the conditional probabilities of the 36 possible genotypes were obtained for every cM of the

204 linkage map for all individuals in the offspring. These probabilities were used to assess the meiotic

205 process that formed the offspring and for further QTL analysis. Probabilistic profiles for pairing

206 behavior were obtained for all LGs for the three possible meiotic pairing configurations, i.e., *ab*/*cd*,

207 *ac*/*bd*, and *ad*/*bc*, where *a*, *b*, *c* and *d* denote the homologs in parent 'Atlantic'. The notation *ab*/*cd*

208 indicates that homolog *a* paired with *b*, and *c* paired with *d*, for example. The same reasoning applies

209 to parent B1829-5 with homologs *e*, *f*, *g* and *h*. Also, the probabilistic haplotypes of all individuals in

210 the offspring were reconstructed and the crossing-over points and respective homologs involved in

211 the exchange were inferred. Using this information and the heuristic algorithm presented by

212 Mollinari *et al.* (2020), recombination chains were assembled for all offspring individuals, and the

213 homologs involved in each meiosis were assessed. Recombination chains with more than two

214 homologs involved imply that a multivalent formation was present during the meiosis.

215 **2.4 QTL mapping and gene search**

216 The conditional genotype probabilities calculated from the inferred genetic map were used to a

217 compute sib-pair additive relationship based on identity-by-descent for every cM position in the map.

218 Using the R package QTLpoly v. 0.2.0 (Pereira *et al.*, 2020), we detected putative QTL positions

219 using the random-effect multiple interval mapping (REMIM) model for each phenotype (trait-year

220 combination) based on a forward-backward procedure. First, putative QTL were consecutively added

221 to the model based on a forward search with a relatively relaxed genome-wide significance threshold

222 ($\alpha = 0.20$). Then, a backward elimination step was carried out under a more stringent threshold ($\alpha =$

223 0.05). The variance components associated with the putative QTL were tested using linear score

224 statistics (Qu *et al.*, 2013), whose associated *P*-values were used in QTL profile visualization with

225 the logarithm of *P*-values $[LOP = -\log_{10}(P)]$. The genome-wide significance was assessed using a

226 score-based resampling method (Zou *et al.*, 2004). A final multiple QTL model was fitted with the

227 selected positions, and QTL genotype best linear unbiased predictions (BLUPs) were used to

228 compute additive allele effects (Kempthorne, 1955). QTL heritability (h^2_{QTL}) was calculated as the

229 ratio between the variance associated with the QTL and the total variance.

230 In order to compare different approaches, we also ran the fixed-effect interval mapping (FEIM)

231 model (Hackett *et al.*, 2014) as implemented in the QTLpoly package (Pereira *et al.*, 2020). Using the

232 same conditional genotype probabilities, a single-QTL model was used to fit six additive effects (as

233 one additive effect from each parent is taken as reference) at every cM position. The model with a

234 fitted QTL was compared to a null model (with no QTL) using likelihood ratio tests (LRT). LRT

235 statistics were converted into LOD scores, and QTL were declared when the LOD score reached a

236 threshold ($\alpha = 0.05$) based on 1,000 permutation tests (Churchill and Doerge, 1994). In both

237 REMIM and FEIM analyses, a window size of 20 cM was used to avoid that two closely linked

238 positions were declared as QTL. Approximate 95% QTL support intervals were estimated by

239 dropping 1.5 from the QTL peak *LOP* (REMIM) or *LOD* (FEIM) (Pereira *et al.*, 2020).

240 Based on the QTL peaks from REMIM, we searched for candidate genes in the region delimited by

241 markers on the left and on the right of the QTL peak or within 200 kbp each side from the QTL peak,

242 whichever was the largest, on the *S. tuberosum* v. 4.03 genome (ST4.03) using the PhytoMine tool

243 (https://phytozome.jgi.doe.gov/phytomine/) by Phytozome v. 12 (Goodstein *et al.*, 2012). Finally, we

244 used the Web Gene Ontology Annotation Plot (WEGO) v. 2.0 (Ye *et al.*, 2018;

245 http://wego.genomics.org.cn/) to count and plot classifications of Gene Ontology (GO) terms for

246 annotated genes within our QTL regions.

3   Results

3.1   Trait correlations and GE interactions

Based on the phenotype means (see Supplementary File S1), variation was found for all evaluated

traits over the four years of evaluation (Figure 1A), indicating that they were segregating in the

B2721 population. As the parental means were within the population phenotypic range, we could

observe transgressive segregation for all trait-year combinations. PY06 has shown quite distinct

median and broader variation in comparison to the following years, which is likely because there

were no replicates in 2006, so that the environmental error could not be modeled. In contrast, FM14

exhibited a narrower variation when compared to the previous years. This was rather expected, since

in 2014 maturity evaluation was carried out in a shorter time-window (24 days) in comparison to

2007 and 2008 (30 days). The overall change in ST14 scores when compared to the previous years

was likely due to differences in evaluator scoring. Individuals seemed to be less affected by IHN in

2008, as NS08 and NI08 have shown distinct variation and median in comparison to the other years.

Broad-sense heritability (H^2) ranged from 0.39 (DM) to 0.81 (SG), with similarly high values for PY

(0.78), FM (0.70) and ST (0.79). For IHN-related traits, NS (0.74) showed a relatively higher

heritability when compared to NI (0.66).

The correlation among variables was explored using a network plot (Figure 1B), where path color

and transparency represent the signal and strength of the correlation, respectively. The variable

positioning is given by multidimensional scaling that leverages the magnitude of the correlations, i.e.

the closer the variables, the higher the absolute values of their correlations (Ruiz *et al.*, 2019). In

general, the phenotype of the same trait evaluated in different years clustered together. As expected,

NS was close to NI as well as DM to SG, thus indicating high correlation, because each one of these

trait pairs are different ways of assessing the same characteristic (IHN response and solids content,

respectively). FM appeared centrally in this plot as it was correlated with most variables either

negatively (e.g. with PY and NI) or positively (e.g. with SG, DM and NS). Note that for this network

plot, a subset of 144 full-sibs with complete observations among all traits was ultimately used. For

correlation estimates based on pairwise complete observations, please see Supplementary Figure S1

and Supplementary File S2. Even though data from 2006 was originated from a single replication,

they were in general agreement with the adjusted means from the other years. DM observations were

the least correlated ones across years in comparison to all the other trait sets (see Supplementary

Figure S1).

GGE biplot was used to investigate the existence of GE interaction (Figure 1C), where each year was

considered an environment. The distribution of full-sibs along the first two principal components,

which accounted for 43% of the variation, did not show any clustering. This was expected as

mapping populations usually consists of non-selected individuals. However, some full-sibs were

found to be more scattered in the top-left (fourth) quadrant, indicating the most affected individuals

by IHN (i.e. high NI). In contrast, more individuals were found to be less scattered in the bottom-

right (second) quadrant, where NS vectors lie, indicating smaller differentiation when full-sibs were

less susceptible to IHN. Relatively shorter vector lengths of PY, ST and DM08 indicated limited

capacity to discriminate genotypes using these traits, in comparison to longer vectors. Only relatively

weak GE interaction was observed, since phenotypes from different years consistently exhibited the

same vector orientation for each trait, with small angles between vectors. Greater angles between

vectors of phenotypes from 2006 (such as DM06, SG06 and NS06) and from the remaining years

should be treated carefully, as these 2006 measurements were based on single plots. The same vector

orientation for SG and DM indicated that these traits were positively correlated, whereas the opposite

orientations for NS and NI vectors revealed their negative correlations.

Figure 1. B2721 mapping population phenotypes evaluated for four years (2006-8 and 2014). (A) Boxplots showing distribution of full-sib means along with parental means. (B) Network plot showing positive (blue) and negative (red) significant correlations ($|r| \geq 0.26$, $P < 0.001$). (C) GGE biplot showing the distribution of full-sibs (green) and phenotypes (blue) on the two first principal components. Traits: plant yield (PY), foliage maturity (FM), specific gravity (SG), dry matter (DM), skin texture (ST), and internal heat necrosis severity (NS) and intensity (NI). The phenotypes were named after a trait abbreviation followed by two-digit year (e.g. PY06).

3.2   Linkage analysis and inheritance mechanisms

The normalized intensities from the Illumina Infinium® 8,303 Potato Array were obtained for the B2721 population (see Supplementary File S3). The dosage calling procedure yielded 5,599 informative SNPs with 1.05% of missing data. Fourteen percent of the SNPs were filtered out due to segregation distortion and redundancy resulting in 4,812 informative markers: 1,311 (27.3%)

simplex, 1,234 (25.6%) double-simplex, and 2,267 (47.1%) multiplex (i.e. from duplex to quadriplex;

see Supplementary Figure S2). The complete linkage map (see Supplementary Table S1,

Supplementary Figure S3 and Supplementary File S5) consisted of 12 LGs with eight haplotypes

each, with four homologs per parent. A total of 4,285 markers spanned 1,629.99 cM in length, with

an average density of 2.64 SNPs/cM. Linkage group lengths ranged from 106.20 cM (LG 5) to

205.88 cM (LG 1), with average LG length of 135.83 cM. The scatterplots of the physical distance in

S. tuberosum genome v. 4.03 versus genetic distance in the 12 LGs in B2721 population map is

shown in Supplementary Figure S4.

The probabilistic pairing profiles showed no preferential pairing between homologs in both parents

(see Supplementary Figure S5). Among all meiotic configurations, only 7.3% and 9.8% were

inconclusive for parents 'Atlantic' and B1829-5, respectively. From the remaining configurations

(Figure 2), the percentage of cases with no crossing-over varied from 6.8% (LG 1, parent 'Atlantic')

to 40.7% (LG 11, parent B1829-5), with 24.2% on average. Configuration involving two

chromosomes with at least one crossing-over (i.e. bivalent configurations), were the most abundant

varying from 51.0% (LG 11, parent B1829-5) to 72.4% (LG 9, parent 'Atlantic'), with a mean of

62.3%. Multivalent configurations, i.e., involving three or four homologs, ranged from 2.2% (LG 8,

parent B1829-5) to 39.2% (LG 1, parent 'Atlantic'), with 13.5% on average.

Figure 2. Distribution of number of homologs involved in a recombination chain.

3.3 Multiple QTL mapping and candidate genes

From the REMIM analysis, a total of 21 QTL were mapped in 15 out of 27 evaluated trait-year combinations (Table 1; see Supplementary Figure S6). One QTL was detected for 10 phenotypes each (for PY06, PY08, SG07, SG08, ST07, ST08, NS06, NS08, NI06 and NI08). The other five phenotypes have shown two (for FM07, FM14, SG06 and NI07) or three (for FM08) QTL each. No QTL were mapped for 12 phenotypes (PY07, PY14, SG14, DM06, DM07, DM08, DM14, ST06, ST14, NS07, NS14 and NI14), mostly from years 2006 and 2014. Ten QTL were mapped on LG 5, four on LG 1, three on LG 7, and one on LGs 2, 3, 4 and 9 each. QTL heritability (h^2_{QTL}) ranged from 7.1% (FM08 on LG 7) to 64.0% (FM08 on LG 5), including five QTL with higher ($64.0 \geq h^2_{\mathrm{QTL}} \geq 36.0$), 10 with moderate ($29.8\% \geq h^2_{\mathrm{QTL}} \geq 20.4\%$), and the remaining six QTL with lower

(18.6% ≥ h^2_{QTL} ≥ 7.1%) heritability (Table 1; Figure 3). In comparison to REMIM results, FEIM

detected 17 QTL from 12 phenotypes (see Supplementary Figure S8). On the one hand, FEIM did not

identified QTL previously identified for PY06 (on LG 5), FM08 (on LGs 1 and 7), SG06 (on LG 3),

NI06 (on LG 1) and NI07 (on LGs 1 and 5). On the other hand, FEIM identified additional QTL for

SG06 (on LGs 8 and 10), ST08 (LG 9), NI14 (on LG 5) (see Supplementary Table S). The proportion

of variance explained (PVE) by FEIM-derived QTL ranged from 13.0% to 52.7%.

With 10 QTL on the proximal half of LG 5, this region was characterized as a QTL hotspot. The

major QTL on LG 5 for FM was detected for all three years of evaluation (2007-8 and 2014).

However, *LOP* profiles (see Supplementary Figure S6) showed suggestive QTL (at a lower *LOP* that

did not reach the threshold), for instance, on LG 5 for PY06, SG14 and NS14 and NI14. For the five

traits with QTL on LG 5 (PY, FM, SG, NS and NI), most QTL peaks were found at 29 cM and

support intervals ranged from 0 to 50 cM. Figure 4 depicts additive allele effects of co-located QTL

on LG 5 for these different traits evaluated in 2008. It is interesting to note that for the QTL sets

PY08.1/NI08.1 and FM08.2/SG08.1/NS08.1, major allele effects appeared in the same direction

within sets, but in opposite directions between them. As these QTL on LG 5 were responsible for

explaining significant portion of the variation for these traits (Table 1), the QTL-based predicted

means were highly correlated, either positively (PY with NI, and FM with SG and NS) or negatively

(PY/NI with FM/SG/NS) (see Supplementary Figure S11). It is worth mentioning that selecting

towards two out of four alleles per parent will tend to favor different sets of phenotypes depending on

the alleles. For example, selecting allele pair *ac* from 'Atlantic' and *eg* from B1829-5 will likely

increase PY and NI and decrease FM, SG and NS. Additive effects for all QTL can be found on

Supplementary File S6.

Table 1. Random-effect multiple interval mapping (REMIM) for B2721 traits evaluated for four

years (2006-8 and 2014).

Trait[a]	QTL	Linkage group	Position (cM)	SI (cM)[b]	Score	P-value	σ^2_{QTL}[c]	h^2_{QTL}[d] (%)
PY06	1	5	32	13-42	137.41	4.70e–05	1.11e–01	20.4
PY08	1	5	46	0-53	168.95	7.21e–06	2.17e–02	36.0
FM07	1	5	26	18-29	355.99	<2.22e–16	3.36e+02	57.7
	2	7	66	52-70	167.43	8.98e–07	7.28e+01	12.5
FM08	1	1	9	0-32	101.58	2.16e–04	1.17e+02	7.4
	2	5	27	0-41	425.04	<2.22e–16	1.01e+03	64.0
	3	7	65	35-71	104.85	2.00e–04	1.13e+02	7.1
FM14	1	5	18	0-29	197.01	9.77e–08	7.12e+00	29.8
	2	7	66	54-70	138.45	1.57e–05	4.01e+00	16.8
SG06	1	2	97	34-117	125.63	6.65e–05	6.69e–06	15.9
	2	3	44	31-50	109.89	1.49e–04	8.94e–06	21.2
SG07	1	5	19	4-32	135.93	3.26e–05	1.39e–05	27.6
SG08	1	5	28	15-32	240.07	1.47e–08	4.24e–05	48.9
*SG07	1	2	104	68-125	131.20	8.32e–05	8.35e–06	23.0
*SG08	1	2	115	60-124	143.00	5.49e–05	7.80e–06	20.9
*DM07	1	2	115	63-121	128.17	1.40e–04	2.62e–01	14.9
	2	6	36	31-73	134.38	3.58e–05	4.13e–01	23.3
ST07	1	9	41	27-82	109.68	1.97e–04	1.59e–01	21.9
ST08	1	4	2	0-4	234.05	1.08e–08	6.32e–01	48.1
*ST08	1	4	2	0-4	255.84	3.62e–09	5.45–e01	46.9
NS06	1	1	0	0-52	113.07	1.29e–04	1.17e–01	22.9
NS08	1	5	26	12-32	117.77	1.77e–04	2.84e–02	23.4
NI06	1	1	0	0-69	110.07	1.66e–04	9.28e+01	22.0
NI07	1	1	0	0-8	103.32	2.26e–04	1.39e+02	16.8
	2	5	25	8-34	134.43	3.35e–05	1.99e+02	24.2
NI08	1	5	27	12-32	131.28	6.54e–05	5.65e+01	28.5

*Maturity-corrected phenotypes.
[a]Traits: plant yield (PY), foliage maturity (FM), specific gravity (SG), dry matter (DM), skin texture (ST), and internal
heat necrosis severity (NS) and intensity (NI).
[b]~95% support interval (SI) based on $LOP - 1.5$, in centiMorgans.
[c]Variance component associated with the QTL (σ^2_{QTL}).
[d]QTL heritability (h^2_{QTL}), in percentage.

Figure 3. QTL identified for seven traits evaluated for four years (2006-8 and 2014). Dots are plotted according to QTL peak location along the linkage groups. Color scale represents P-values, and sizes are proportional to QTL heritability (h^2_{QTL}). Traits: plant yield (PY), foliage maturity (FM), specific gravity (SG), dry matter (DM), skin texture (ST), and internal heat necrosis severity (NS) and intensity (NI). * indicates maturity-corrected phenotypes.

Due to such a large QTL effect on LG 5 for FM and its consistent impact on most traits, we fitted simple linear regressions of the form $y = 1\beta_0 + \beta_1 x + e$ using the lm() function of R (R Core Team, 2020), with FM as the explanatory variable x and any other phenotype as dependent variable y from each respective year where FM was measured (i.e. 2007, 2008 and 2014). The vector of residuals e were used as maturity-corrected phenotypes (i.e. virtually free of FM influence; see Supplementary Figure S10) for new QTL mapping runs, as suggested by Bradshaw et $al.$ (2004). We found that the slopes β_1 were highly significant for all traits (***$P < 0.001$), except DM14 (*$P < 0.05$), and

DM08, PY14, SG14 and ST14 ($P > 0.1$) (see Supplementary Table S2). REMIM analyses on

maturity-corrected phenotypes resulted in five QTL (see Supplementary Fig. S7). A QTL for ST08

co-located with the one previously identified on LG 4. Other four QTL were newly identified: one

QTL from each SG07, SG08 and DM07 were closely located on LG 2, and another QTL for DM07

was detected on LG 6 (Table 1; Figure 3). FEIM analyses returned eight QTL, from which two were

previously identified for ST08 (LGs 4 and 9), and other six were newly detected for SG07 (LG 2),

SG 08 (LGs 2 and 5), DM07 (LG 6), DM08 (LG 11), and NI14 (LG 4) (see Supplementary Table S4

and Supplementary Figure S9).

Figure 4. Additive allele effects for co-located QTL on LG 5 for five traits evaluated in 2008. Letters

represent each parental haplotype from the linkage map ('Atlantic' = *a* through *d*, B1829-5 = *e*

through *h*). Traits: plant yield (PY), foliage maturity (FM), specific gravity (SG), and internal heat

necrosis severity (NS) and incidence (NI).

As the QTL support intervals comprised too large of a region to manually curate (>13 Mb per QTL,

on average), we focused our candidate gene search on the region within the markers flanking the

QTL peak (~400 kb, on average) (see Supplementary Table S3). A total of 533 annotated genes on *S.*

tuberosum v. 4.03 genome were within these QTL regions, with 342 genes assigned to at least one

GO term (see Supplementary File S7 and Supplementary Figure S12).

4   **Discussion**

4.1   **The recombination landscape of B2721 mapping population**

The ability to assess the intricate recombination landscape of an outcrossing, autopolyploid species

encompass the construction of a highly informative map. That is, the parental maps have to be

integrated between the two parent and must differentiate each one of the parental haplotypes. Two

genetic maps have been built previously for the B2721 population. The first map used amplified

length polymorphisms (AFLP) and a few simple sequence repeat (SSR) markers, and consisted of

two separate linkage maps, one for each parent, with limited integration within parents (McCord *et*

al., 2011b). The second map used dosage-sensitive SNP array, which led to full integration and

haplotyping in the B2721 population (Schumann *et al.*, 2017). However, this map contained 3,427

SNPs and spanned 1,397.86 cM, which is a relatively smaller but less saturated than our current map

with 4,285 SNPs distributed along 1,629.99 cM. The ability to incorporate more markers with

MAPpoly was possibly due to (*i*) the multipoint-based algorithm that considers all parental homologs

and marker dosages simultaneously, and (*ii*) the genome-assisted approach for marker ordering.

Schumann *et al.* (2017) used the software TetraploidSNPMap (Hackett *et al.*, 2016), which relies on

(*i*) simplex markers to assemble homologs, bridging these homologs afterwards using multidose

markers, and (*ii*) two-point genetic distances to obtain the marker order by using the

multidimensional scaling (MDS) algorithm (Preedy and Hackett, 2016). Increased marker saturation

provides better support to estimate the conditional probabilities, which are used to infer pairing

profiles during meiosis in a segregating population and to map QTL.

The probabilistic pairing profiles showed no preferential pairing between homologs in both parents

(see Supplementary Figure S5), likely due to the cultivated potato's autotetraploid nature.

Consequently, alleles present in different homologs in the same homology group had an equal chance

to recombine with alleles from other loci, amplifying the range of possible genotypes when compared

to diploids or allotetraploids. In addition, most of the parental meiotic configurations based on the

B2721 population were inferred as bivalents (62.3% on average) since they involved two homologs

exchanging segments during the metaphase I (Figure 2). These results are in accordance with recent

findings published by Choudhary *et al.* (2020). These authors presented a map of meiotic stages in *S.*

tuberosum in one diploid and three tetraploid varieties using fluorescence *in situ* hybridization

(FISH). They concluded that bivalent chromosome associations are the most common in metaphase I

of tetraploid potatoes. Likewise, the broad range of multivalent signature rates observed in our study

(2.2~39.2%) was also observed by Choudhary *et al.* (2020) (7~48%). Similarly, as their observations

differed depending on variety and chromosome, in our study, the variation occurred by parents and

by linkage groups. Thus, our meiotic assessment, although not as precise as a cytological analysis

and prone to sampling errors, can be very helpful, serving as a proxy to evaluate meiotic

configurations by using a straightforward expansion of linkage analysis.

4.2   Multiple QTL mapping for yield, quality and IHN related traits

Single-QTL models were previously applied to the first two linkage maps of B2721. These models

were based on an interval mapping model, and the 95$^{\text{th}}$ percentile ($\alpha = 0.05$) of maximum LOD

scores from 1,000 permutation tests per phenotype was used to declare QTL, with several additional

suggestive QTL (below this threshold) also being recorded (McCord *et al.*, 2011b, 2011a). One of the

issues with the lack of integration within and between parental maps is the same region being

declared as a QTL in separate parental haplotypes. For instance, for plant yield, McCord *et al.*

(2011b) listed a total of seven QTL across three years, PY06 (one), PY07 (five) and PY08 (two),

from both parental maps, in addition to 11 suggestive ones. Our multiple-QTL analyses resulted in a

single co-located QTL region on LG 5 for each PY06 and PY08 (Table 1; Figure 3), with suggestive

QTL for PY07 and PY14 (see Supplementary Figure S6). QTL on LG 5 for tuber yield have also

been reported in both diploid (DM 1-3 516 R44 × RH89-039-16, $n = 96$) (Manrique-Carpintero *et*

al., 2015) and tetraploid ('Liberator' × W4013-1, $n = 110$) (Rak *et al.*, 2017) potato mapping

populations.

For foliage maturity, McCord *et al.* (2011b) and Schumann *et al.* (2017) described major QTL for

years 2007-8 on LG 5 as well as the one on LG 7. In the current study, another QTL on LG 1 was

identified for FM08, likely due to increased detection power of REMIM. Despite the differences

regarding phenotypic distribution for FM14 (Figure 1A), we were able to find QTL on LGs 5 and 7

(Figure 3). This LG 5 region is well known as a main source of QTL in potato. Prior to the release of

the SNP potato array, a major QTL for plant maturity had already been reported on chromosome 5,

also linked to a late blight (*Phytophthora infestans*) resistance locus (reviewed by Danan *et al.*,

2011). More recently, additional studies noticed that the same region on LG 5 also underlies the

variation of several tuber quality-related traits in addition to resistance to early blight (*Altenaria*

tenuis) and Verticillium wilt (*Verticillium* spp.) (Massa *et al.*, 2018), and of tuber-Cd and Zn

concentration (Mengist *et al.*, 2018), for example. For IHN-related traits, three QTL were identified

on LG 5, one for each NS08, NI07 and NI08, using REMIM. FEIM identified an additional

significant QTL on LG 5 for NI14 (see Supplementary Table S4 and Supplementary Figure S8).

Previously, IHN-related QTL had been reported as suggestive only (Schumann *et al.*, 2017), and this

difference in significance is likely due to an improved map construction.

A major QTL on LG 5 was also found to co-locate for SG07 and SG08, while it appeared only as

suggestive for SG14 and with no evidence for SG06, which in turn showed one QTL on each LGs 2

and 3. The evaluation of maturity-corrected phenotypes helped to detect this co-located QTL for

DM07, and SG07 and SG08 on LG 2. An alternative to perform QTL discovery free of maturity

influence, though, would be to use materials that do not show much variation for this trait, i.e. crosses

between either late or early maturity cultivars. Specific gravity and dry matter are supposedly related

with solid content in the tuber. In fact, the correlation among their adjusted means were generally

high (up to 0.81 between DM07 and SG07; see Supplementary Figure S11 and Supplementary File

S2), but the only set of co-located QTL between any SG and DM measures was that one on LG 2. It

is worth to mention that their broad-sense heritability values (H^2) differed greatly (0.39 for DM and

0.81 for SG), so that the ability to detect QTL for DM may have been affected because smaller

proportion of the total variance could be attributed to genetic variance. Other mapping populations

have revealed QTL on LG 5 for DM (12601ab1 × 'Stirling', $n = 227$) (Bradshaw *et al.*, 2008) and

SG (Manrique-Carpintero *et al.*, 2015), where somewhat distinct heritability values were found (0.81

for DM and 0.46 for SG).

Skin texture was a trait which diverged the most from the other traits regarding QTL locations,

specifically for not showing enough statistical evidence for QTL on LG 5 (see Supplementary Figure

S6). In diploid potatoes, skin texture is believed to be controlled by three loci (Jong, 1981). In our

B2721 population, we detected two QTL (one for ST07 and ST08 each), in regions on LGs 4 and 9

where no other traits have shown evidence for QTL in our study. That is, the apparent strong

correlation observed between ST07 and FM07 ($r = -0.44$, $P < 0.001$), for example, could not be

attributed to genetics, as correlation between QTL-based predicted means for these two traits was

close to zero ($r = -0.07$, $P = 0.40$; see Supplementary Figure S11).

When performing QTL discovery in outbred polyploid populations, in addition to genome

complexity, one must keep two factors in mind. First, in F_1 populations, the fact that two parental

genotypes are contrasting does not directly imply that major QTL will be detected. For instance,

considering a biallelic QTL, a cross between two contrasting parents, *QQQQ* × *qqqq*, would not

segregate (i.e. 100% *QQqq*). In order to be detectable, alleles need to be contrasting within parent,

e.g. *Qqqq* × *qqqq* or *Qqqq* × *Qqqq* in the simplest cases, resulting in segregations such as 50% *Qqqq*

: 50% *qqqq* or 25% *QQqq* : 50% *Qqqq* : 25% *qqqq*, respectively. We speculate that the allele effects

of undetectable QTL towards IHN resistance, for example, did not differ significatively within each

parent. For the QTL on LG 5 for NI08, the allele contributions to the population mean (~6.1%) were

±2.5% at most (Figure 4). In other words, these effects would not change the fact that most

individuals would be less susceptible (NI < 10%) as resistance could have been already conferred by

these undetectable QTL.

If there are segregating QTL alleles, another factor that influences detection is small population sizes

that may not allow to separate signal from noise properly. Consequently, thresholds for declaring a

QTL are usually required to be more stringent in order to avoid false positives. In general, tetraploid

potato QTL mapping studies have been performed in relatively small population sizes (~150 or less

individuals). In these cases, even very low *p*-values and high QTL heritability estimates, such as

those found for FM07, FM08 and SG08, should be treated carefully as they can be relatively biased

due to sampling. Finally, reduced population sizes also limit the ability to test closely linked versus

pleiotropy for co-located QTL across different traits or environments accurately.

4.3   Candidate genes within QTL regions

Among the candidate genes (see Supplementary File S7), GO terms that overlapped DNA and

protein binding functions with regulation of biological process were of particular interest. In total, 19

genes were found to have the same two GO terms (GO:0003677 and GO:0006355) attributed to the

S. tuberosum CYCLING DOF (DNA-binding with one finger) FACTOR 1, *StCDF1*

(PGSC0003DMG400018408, ST4.03ch05:4539029..4541329), which has been described as the

transcription factor underlying the major QTL for maturity on LG 5 (Kloosterman *et al.*, 2013). The

SolCAP markers around the QTL peak for FM08 (c2_22959 at 4,434,048bp and c2_23052 at

4,906,728 bp) were found to be closer to *StCDF1* than the flanking markers for FM07 and FM14

(c2_11829 at 4,041,510 bp and c2_22986 at 4,279,075 bp), and the marker mentioned by Massa *et al.*

(2015) (c2_22986 at 4,105,142). The same LG 5 region also includes yield-related QTL, where we

found the *Potato homeobox 1* gene, *POTH1* (PGSC0003DMG400013493,

ST4.03ch05:9248372..9258054), that appears to regulate tuber development in potato via StBEL5-

POTH1 heterodimer (Rosin *et al.*, 2003). Finally, QTL underlying IHN-related traits were also found

in the same LG 5 hotspot, where a homeobox-associated leucine zipper, *StHOX40*

(PGSC0003DMG400030494, ST4.03ch05:4264626..4267698), was identified. This gene was found

to be either up or downregulated under heat and drought stresses in potato, respectively (Li *et al.*,

2019), which are believed to be two important triggers of IHN (Yencho *et al.*, 2008).

Other genes, such as *StERF6* (PGSC0003DMG400016651, ST4.03ch06:33827698..33832740) and

StNAC046 (PGSC0003DMG400031266, ST4.03ch05:4946232..4948779), were characterized as

transcription factors from AP2/ERF and NAC families, respectively, which have been implicated in a

wide range of regulation processes in plants (Olsen *et al.*, 2005; Mizoi *et al.*, 2012). A gene from the

NAC family, *StNAC032* (PGSC0003DMG400002824, ST4.03ch04:245125..246607), was retrieved

from the LG 4 region of a major QTL for skin texture. Several genes from the NAC family, including

StNAC032, were found to be expressed in the tuber skin and appeared to be involved in suberin and

associated wax biosynthesis (Soler *et al.*, 2020), which could contribute to skin texture, as well as be

involved with response to drought and other biotic and abiotic stresses (Singh *et al.*, 2013).

In plants, a single QTL region may be found to underlie the variation of more than one phenotype at

once, which can potentially explain the correlation among traits. A QTL hotspot has been identified

in the short arm of chromosome 5 of potato (Bradshaw *et al.*, 2004; Danan *et al.*, 2011; Li *et al.*,

2018), and our B2721 population stuck to the rule. A major QTL for maturity was found to overlap

loci underlying the variation for other traits such as yield, specific gravity and IHN-related resistance,

as described previously (McCord *et al.*, 2011a, 2011b; Schumann *et al.*, 2017) and corroborated with

newly analyzed phenotypic data. Pleiotropy is one possible explanation of physiologically correlated

traits, such as maturity, yield and specific gravity (as a proxy of solid content, thus contributing to

yield). Transcription factors are known to have this property in which a single molecular function

may be involved in multiple biological processes (He and Zhang, 2006). For instance, a DNA

binding element from the HAP complex, *Ghd8*, was found to play pleiotropic roles in regulating

yield, flowering and plant height in rice (Yan *et al.*, 2011). On the other hand, the presence of QTL

for yield-related traits and pathogenic (e.g. late blight) and non-pathogenic (e.g. IHN) potato

disorders in the same hotspot on LG 5 could well be due to closely linked genes. Another example

from rice has suggested the multigenic nature of a major QTL for grain yield under drought, where

the central role of a NAM transcription factor, *OsNAM$_{12.1}$*, in concert with extra co-localized genes,

influenced other traits such as root and panicle branching, and transpiration efficiency (Dixit *et al.*,

2015).

5   Final Considerations

'Atlantic' has been widely adopted as a chipping variety, mostly due to its high specific gravity and

yield, and in spite of its medium-late maturity and susceptibility to internal heat necrosis (Webb *et*

al., 1978). The genetic basis of the correlation between these traits was dissected here by means of

multiple QTL models in combination with a dense, fully phased genetic map. We found that not only

a major QTL on LG 5 underlies the variation of these traits concomitantly, but also that the additive

effects attributed to each parental haplotype appear in such a way where, in general, when an allele

increases SG and FM measurements (thus towards early maturity), it decreases PY and IHN

resistance (hence towards low NS and high NI), and vice-versa. *StCDF1* has been pointed out as

playing a central role in this major QTL for maturity, with its day-length signaling function that

regulates the tuberization pathway (Kloosterman *et al.*, 2013). In addition to low rainfall in the early

season and high temperatures, longer days to harvest are known to increase IHN incidence (reviewed

by Yencho *et al.*, 2008). Similarly, higher number of rain events seemed to contribute to the

occurrence of IHN, because rainfall interferes, among other variables, with light intensity (Sterrett *et*

al., 1991). Therefore, due to its circadian role, we believe that *StCDF1* indirectly participates in IHN

progression.

Our list of candidate genes can be used in further dissection of QTL regions. Ideally, factors other

than *StCDF1* could be targeted in benefit of yield and specific gravity increase at the expense of late

maturity. One way of doing so is to focus on linked genes (e.g. *POTH1* for yield) or other non-linked

QTL (e.g. the one on LG 2 for specific gravity). Moreover, the evaluation of germplasm collections

in order to assess haplotypic diversity could assist the development and implementation of markers to

monitor loci of interest, as previously performed for *StCDF1* in chromosome 5 (Gebhardt *et al.*,

2004; Hardigan *et al.*, 2017). Marker-assisted breeding for quantitative, more complex traits could be

implemented upon validation of these QTL neighboring markers in breeding populations of tetraploid

potato. DNA markers also can be employed in genome-wide prediction in this crop (Sverrisdóttir *et*

al., 2017; Enciso-Rodriguez *et al.*, 2018). In this case, both meiotic and trait genetic architecture

assessments derived from our analyses can be particularly useful in refining simulations of potato and

other autopolyploid crop breeding programs integrating genomics-assisted breeding (Gaynor *et al.*,

2020).

6   Author Contributions

GCY designed the experiments and supervised the project. MJS and MEC performed field

experiments and collected data. GSP, MM and ZBZ analyzed data. GSP drafted the manuscript with

major contributions from MM. All authors read and approved the manuscript.

7   Funding

This project was funded by USDA-ARS [58-1245-3-307, 2014-34141-22266] and USDA-NIFA

[2013-34141-21392]. Scholarships were awarded to MJS by Pioneer and Monsanto, and to GSP and

MM by Bill & Melinda Gates Foundation [OPP1052983]. Currently, GSP holds a scholarship from

PrInt-CAPES, Brazil [88887.369781/2019-00].

8   Acknowledgments

We would like to thank all the staff at the North Carolina Department of Agriculture and Consumer

Services Tidewater Research Station, Plymouth, NC, for their hard work and dedication to quality

research in aiding with the field aspect of this study. We also appreciate Michigan State University

Potato Breeding and Genetics Program for their assistance with the Illumina Infinium® 8,303 Potato

Array.

9   Competing Interests

610   The authors declare no competing financial interests.

10   Data Archiving

613   The datasets analyzed and scripts used in this study can be found in the supplementary material and

614   at the study GitHub page (https://github.com/mmollina/B2721_map). Both MAPpoly

615   (https://github.com/mmollina/MAPpoly) and QTLpoly (https://github.com/guilherme-

616   pereira/QTLpoly) R packages are available at GitHub.

11   References

Bennett MD (2004). Perspectives on polyploidy in plants - Ancient and neo. *Biol J Linn Soc* **82**: 411–

423.

Bethke PC, Nassar AMK, Kubow S, Leclerc YN, Li XQ, Haroon M, *et al.* (2014). History and

Origin of Russet Burbank (Netted Gem) a Sport of Burbank. *Am J Potato Res* **91**: 594–609.

Bourke PM, van Geest G, Voorrips RE, Jansen J, Kranenburg T, Shahin A, *et al.* (2018).

polymapR—linkage analysis and genetic map construction from F1 populations of outcrossing

polyploids. *Bioinformatics*: 1–7.

Bourke PM, Voorrips RE, Visser RGF, Maliepaard C (2015). The double-reduction landscape in

tetraploid potato as revealed by a high-density linkage map. *Genetics* **201**: 853–863.

Bradshaw JE, Hackett CA, Pande B, Waugh R, Bryan GJ (2008). QTL mapping of yield, agronomic

and quality traits in tetraploid potato (Solanum tuberosum subsp. tuberosum). *Theor Appl Genet*

116: 193–211.

Bradshaw JE, Pande B, Bryan GJ, Hackett CA, McLean K, Stewart HE, *et al.* (2004). Interval

mapping of quantitative trait loci for resistance to late blight [Phytophthora infestans (Mont.) de

bary], height and maturity in a tetraploid population of potato (Solanum tuberosum subsp.

tuberosum). *Genetics* **168**: 983–995.

Butler DG, Cullis BR, Gilmour AR, Gogel BJ, Thompson R (2018). *ASReml-R Reference Manual*

Version 4. Hemel Hempstead.

Chen J, Zhang F, Wang L, Leach L, Luo Z (2018). Orthogonal contrast based models for quantitative

genetic analysis in autotetraploid species. *New Phytol* **220**: 332–346.

Choudhary A, Wright L, Ponce O, Chen J, Prashar A, Sanchez-Moran E, *et al.* (2020). Varietal

variation and chromosome behaviour during meiosis in Solanum tuberosum. *Heredity (Edinb)*.

Churchill GA, Doerge RW (1994). Empirical threshold values for quantitative trait mapping.

Genetics **138**: 963–71.

Comai L (2005). The advantages and disadvantages of being polyploid. *Nat Rev Genet* **6**: 836–846.

Danan S, Veyrieras JB, Lefebvre V (2011). Construction of a potato consensus map and QTL meta-

analysis offer new insights into the genetic architecture of late blight resistance and plant

maturity traits. *BMC Plant Biol* **11**.

Dixit S, Kumar Biswal A, Min A, Henry A, Oane RH, Raorane ML, *et al.* (2015). Action of multiple

intra-QTL genes concerted around a co-localized transcription factor underpins a large effect

QTL. *Sci Rep* **5**: 1–16.

Dumble S, Bernal EF, Villardon PG (2017). GGEBiplots: GGE Biplots with 'ggplot2'.

Enciso-Rodriguez F, Douches D, Lopez-Cruz M, Coombs J, de los Campos G (2018). Genomic

Selection for Late Blight and Common Scab Resistance in Tetraploid Potato (Solanum

tuberosum). *G3 Genes|Genomes|Genetics* **8**: 2471–2481.

Endelman JB, Carley CAS, Bethke PC, Coombs JJ, Clough ME, Silva WL, *et al.* (2018). Genetic

Variance Partitioning and Genome-Wide Prediction with Allele Dosage Information in

Autotetraploid Potato. *Genetics* **209**: 77–87.

FAO (2020). FAOSTAT Crops.

Felcher KJ, Coombs JJ, Massa AN, Hansey CN, Hamilton JP, Veilleux RE, *et al.* (2012). Integration

of two diploid potato linkage maps with the potato genome sequence. *PLoS One* **7**: e36347.

Gaynor RC, Gorjanc G, Hickey JM (2020). AlphaSimR : Simulations An R-package for Breeding

Program. *bioRxiv*: 1–21.

Gebhardt C, Ballvora A, Walkemeier B, Oberhagemann P, Schüler K (2004). Assessing genetic

potential in germplasm collections of crop plants by marker-trait association: A case study for

potatoes with quantitative variation of resistance to late blight and maturity type. *Mol Breed* **13**:

93–102.

Ghislain M, Douches DS (2020). The Genes and Genomes of the Potato. In: Campos H, Ortiz O

(eds) *The Potato Crop*, Springer, pp 139–162.

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, *et al.* (2012). Phytozome: a

comparative platform for green plant genomics. *Nucleic Acids Res* **40**: D1178--D1186.

Hackett CA, Boskamp B, Vogogias A, Preedy KF, Milne I, Vogogias T, *et al.* (2016).

TetraploidSNPMap: software for linkage analysis and QTL mapping in autotetraploid

populations using SNP dosage data. *J Hered* **108**: 438–442.

Hackett CA, Bradshaw JE, Bryan GJ (2014). QTL mapping in autotetraploids using SNP dosage

information. *Theor Appl Genet* **127**: 1885–1904.

Hämäläinen JH, Watanabe KN, Valkonen JPT, Arihara A, Plaisted RL, Pehu E, *et al.* (1997).

Mapping and marker-assisted selection for a gene for extreme resistance to potato virus Y.

Theor Appl Genet **94**: 192–197.

Hardigan MA, Laimbeer FPE, Newton L, Crisovan E, Hamilton JP, Vaillancourt B, *et al.* (2017).

Genome diversity of tuber-bearing Solanum uncovers complex evolutionary history and targets

of domestication in the cultivated potato. *Proc Natl Acad Sci U S A* **114**: E9999–E10008.

He X, Zhang J (2006). Toward a molecular understanding of pleiotropy. *Genetics* **173**: 1885–1891.

Holland JB, Nyquist WE, Cervantes-Martínez CT (2003). Estimating and interpreting heritability for

plant breeding: an update. In: Janick J (ed) *Plant Breeding Reviews*, John Wiley & Sons: New

Jersey Vol 22, pp 9–112.

Huang K, Wang T, Dunn DW, Zhang P, Cao X, Liu R, *et al.* (2019). Genotypic Frequencies at

Equilibrium for Polysomic Inheritance Under Double-Reduction. *G3 Genes|Genomes|Genetics*

9: 1693–1706.

Jong H (1981). Inheritance of russeting in cultivated diploid potatoes. *Potato Res* **24**: 309–313.

Kempthorne O (1955). The correlation between relatives in a simple autotetraploid population.

Genetics **40**: 168–174.

Kloosterman B, Abelenda JA, Gomez MDMC, Oortwijn M, De Boer JM, Kowitwanich K, *et al.*

(2013). Naturally occurring allele diversity allows potato cultivation in northern latitudes.

Nature **495**: 246–250.

Li W, Dong J, Cao M, Gao X, Wang D, Liu B, *et al.* (2019). Genome-wide identification and

characterization of HD-ZIP genes in potato. *Gene* **697**: 103–117.

Li X, Xu J, Duan S, Zhang J, Bian C, Hu J, *et al.* (2018). Mapping and QTL analysis of early-

maturity traits in tetraploid potato (Solanum tuberosum L.). *Int J Mol Sci* **19**.

Luo ZW, Zhang RM, Kearsey MJ (2004). Theoretical basis for genetic linkage analysis in

autotetraploid species. *PNAS* **101**: 7040–7045.

Mann H, Iorizzo M, Gao L, D'Agostino N, Carputo D, Chiusano ML, *et al.* (2011). Molecular

Linkage Maps: Strategies, Resources and Achievements. In: Bradeen JM, Kole C (eds)

Genetics, Genomics and Breeding of Potato, CRC Press: Enfield, pp 68–89.

Manrique-Carpintero NC, Coombs JJ, Cui Y, Veilleux RE, Robin Buell C, Douches D (2015).

Genetic map and QTL analysis of agronomic traits in a diploid potato population using single

nucleotide polymorphism markers. *Crop Sci* **55**: 2566–2579.

Massa AN, Manrique-Carpintero NC, Coombs J, Haynes KG, Bethke PC, Brandt TL, *et al.* (2018).

Linkage analysis and QTL mapping in a tetraploid russet mapping population of potato. *BMC*

Genet **19**: 1–13.

Massa AN, Manrique-Carpintero NC, Coombs JJ, Zarka DG, Boone AE, Kirk WW, *et al.* (2015).

Genetic linkage mapping of economically important traits in cultivated tetraploid potato

(Solanum tuberosum L.). *G3 Genes, Genomes, Genet* **5**: 2357–2364.

McCord PH, Sosinski BR, Haynes KG, Clough ME, Yencho GC (2011a). QTL mapping of internal

heat necrosis in tetraploid potato. *Theor Appl Genet* **122**: 129–142.

McCord PH, Sosinski BR, Haynes KG, Clough ME, Yencho GC (2011b). Linkage mapping and

QTL analysis of agronomic traits in tetraploid potato (Solanum tuberosum subsp. tuberosum).

Crop Sci **51**: 771–785.

Mengist MF, Alves S, Griffin D, Creedon J, McLaughlin MJ, Jones PW, *et al.* (2018). Genetic

mapping of quantitative trait loci for tuber-cadmium and zinc concentration in potato reveals

associations with maturity and both overlapping and independent components of genetic control.

Theor Appl Genet **131**: 929–945.

Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K (2012). AP2/ERF family transcription factors in plant

abiotic stress responses. *Biochim Biophys Acta - Gene Regul Mech* **1819**: 86–96.

Mollinari M, Garcia AAF (2019). Linkage Analysis and Haplotype Phasing in Experimental

Autopolyploid Populations with High Ploidy Level Using Hidden Markov Models. *G3*

Genes|Genomes|Genetics **9**: 3297–3314.

Mollinari M, Olukolu BA, Pereira GS, Khan A, Gemenet D, Yencho GC, *et al.* (2020). Unraveling

the Hexaploid Sweetpotato Inheritance Using Ultra-Dense Multilocus Mapping. *G3*

Genes|Genomes|Genetics **10**: 281–292.

Olsen AN, Ernst HA, Leggio L Lo, Skriver K (2005). NAC transcription factors: structurally distinct,

functionally diverse. *Trends Plant Sci* **10**: 79–87.

Pereira G da S, Gemenet DC, Mollinari M, Olukolu BA, Wood JC, Diaz F, *et al.* (2020). Multiple

QTL Mapping in Autopolyploids: A Random-Effect Model Approach with Application in a

Hexaploid Sweetpotato Full-Sib Population. *Genetics* **215**: 579–595.

Preedy KF, Hackett CA (2016). A rapid marker ordering approach for high-density genetic linkage

maps in experimental autotetraploid populations using multidimensional scaling. *Theor Appl*

Genet **129**: 2117–2132.

Qu L, Guennel T, Marshall SL (2013). Linear score tests for variance components in linear mixed

models and applications to genetic association studies. *Biometrics* **69**: 883–892.

R Core Team (2020). R: A Language and Environment for Statistical Computing.

Rak K, Bethke PC, Palta JP (2017). QTL mapping of potato chip color and tuber traits within an

autotetraploid family. *Mol Breed* **37**.

Ramakrishnan AP, Ritland CE, Blas Sevillano RH, Riseman A (2015). Review of Potato Molecular

Markers to Enhance Trait Selection. *Am J Potato Res* **92**: 455–472.

Revelle W (2018). psych: Procedures for Psychological, Psychometric, and Personality Research.

Rosin FM, Hart JK, Horner HT, Davies PJ, Hannapel DJ (2003). Overexpression of a Knotted-Like

Homeobox Gene of Gibberellin Accumulation 1. *Plant Physiol* **132**: 106–117.

Ruiz E, Jackson S, Cimentada J (2019). corrr: Correlations in R.

Schmitz Carley CA, Coombs JJ, Douches DS, Bethke PC, Palta JP, Novy RG, *et al.* (2017).

Automated tetraploid genotype calling by hierarchical clustering. *Theor Appl Genet* **130**: 717–

726.

Schultz L, Cogan NOI, Mclean K, Dale MFB, Bryan GJ, Forster JW, *et al.* (2012). Evaluation and

implementation of a potential diagnostic molecular marker for H1-conferred potato cyst

nematode resistance in potato (Solanum tuberosum L.). *Plant Breed* **131**: 315–321.

Schumann MJ, Zeng ZB, Clough ME, Yencho GC (2017). Linkage map construction and QTL

analysis for internal heat necrosis in autotetraploid potato. *Theor Appl Genet* **130**: 2045–2056.

Shaner G, Finney RE (1977). The Effect of Nitrogen Fertilization on the Expression of Slow-

Mildewing Resistance in Knox Wheat. *Phytopathology* **67**: 1051–1056.

Sharma SK, Bolser D, de Boer J, Sonderkaer M, Amoros W, Carboni MF, *et al.* (2013). Construction

of reference chromosome-scale pseudomolecules for potato: integrating the potato genome with

genetic and physical maps. *G3 Genes|Genomes|Genetics* **3**: 2031–47.

Singh AK, Sharma V, Pal AK, Acharya V, Ahuja PS (2013). Genome-wide organization and

expression profiling of the NAC transcription factor family in potato (solanum tuberosum L.).

DNA Res **20**: 403–423.

Slater AT, Cogan NOI, Hayes BJ, Schultz L, Dale MFB, Bryan GJ, *et al.* (2014). Improving breeding

efficiency in potato using molecular and quantitative genetics. *Theor Appl Genet* **127**: 2279–

2292.

Soler M, Verdaguer R, Fernández-Piñán S, Company-Arumí D, Boher P, Góngora-Castillo E, *et al.*

(2020). Silencing against the conserved NAC domain of the potato StNAC103 reveals new

NAC candidates to repress the suberin associated waxes in phellem. *Plant Sci* **291**: 110360.

Sterrett SB, Lee GS, Henninger MR, Lentner M (1991). Predictive Model for Onset and

Development of Internal Heat Necrosis of 'Atlantic' Potato. *J Am Soc Hortic Sci* **116**: 701–705.

Sverrisdóttir E, Byrne S, Sundmark EHR, Johnsen HØ, Kirk HG, Asp T, *et al.* (2017). Genomic

prediction of starch content and chipping quality in tetraploid potato using genotyping-by-

sequencing. *Theor Appl Genet* **130**: 2091–2108.

Webb RE, Wilson DR, Shumaker JR, Graves B, Henninger MR, Watts J, *et al.* (1978). Atlantic: A

new potato variety with high solids, good processing quality, and resistance to pests. *Am Potato*

J **55**: 141–145.

Wei T, Simko V, Levy M, Xie Y, Jin Y, Zemla J (2017). corrplot: Visualization of a Correlation

Matrix.

Wickham H (2016). *ggplot2: elegant graphics for data analysis*. Springer.

Xu X, Pan S, Cheng S, Zhang B, Mu D, Ni P, *et al.* (2011). Genome sequence and analysis of the

tuber crop potato. *Nature* **475**: 189–195.

Yan W, Kang MS (2003). *GGE Biplot Analysis: a graphical tool for breeders, geneticists, and*

agronomists. CRC Press: Boca Raton.

Yan WH, Wang P, Chen HX, Zhou HJ, Li QP, Wang CR, *et al.* (2011). A major QTL, Ghd8, plays

pleiotropic roles in regulating grain productivity, plant height, and heading date in rice. *Mol*

Plant **4**: 319–330.

Ye J, Zhang Y, Cui H, Liu J, Wu Y, Cheng Y, *et al.* (2018). WEGO 2.0: A web tool for analyzing

and plotting GO annotations, 2018 update. *Nucleic Acids Res* **46**: W71–W75.

Yencho GC, McCord PH, Haynes KG, Sterrett SBR (2008). Internal heat necrosis of potato - A

review. *Am J Potato Res* **85**: 69–76.

Zielinski M-L, Scheid OM (2012). Meiosis in Polyploid Plants. In: Soltis PS, Soltis DE (eds)

Polyploidy and Genome Evolution, Springer: Berlin, Heidelberg, pp 33–55.

Zou F, Fine JP, Hu J, Lin DY (2004). An efficient resampling method for assessing genome-wide

statistical significance in mapping quantitative trait loci. *Genetics* **168**: 2307–2316.

Zych K, Gort G, Maliepaard CA, Jansen RC, Voorrips RE (2019). FitTetra 2.0 - Improved genotype

calling for tetraploids with multiple population and parental data support. *BMC Bioinformatics*

20: 1–8.