The recombination landscape and multiple QTL mapping in a *Solanum tuberosum* cv. 'Atlantic'-derived F$_1$ population

Guilherme da Silva Pereira[1,2*], Marcelo Mollinari[1,2], Mitchell J. Schumann[1], Mark E.

Clough[1,3], Zhao-Bang Zeng[1,2], G. Craig Yencho[1*]

[1]Department of Horticultural Science, North Carolina State University, Raleigh, NC, USA

[2]Bioinformatics Research Center, North Carolina State University, Raleigh, NC, USA

[3]Vernon G. James Research and Extension Center, Department of Horticultural Science, North

Carolina State University, Plymouth, NC, USA

Current addresses:

Guilherme da Silva Pereira: Department of Genetics, Luiz de Queiroz College of Agriculture,

University of São Paulo, Piracicaba, Brazil

Mitchell J. Schumann: Bayer Crop Science, Chesterfield, MO, USA

Word counts:

Abstract = 193, Introduction = 751, Materials and Methods = 1690, Results = 1574, Discussion =

2125, Total = 6140

Figures = 4 (color), Tables = 1, Supporting information = 19

Keywords: linkage map, mixed model, multiple interval mapping, polyploid, quality, tetraploid

potato, yield.

Abstract

There are many challenges involved with the genetic analyses of autopolyploid species, such as the

tetraploid potato, *Solanum tuberosum* ($2n = 4x = 48$). The development of new analytical methods

has made it valuable to re-analyze an F_1 population ($n = 156$) derived from a cross involving

'Atlantic', a widely grown chipping variety in the USA. A fully integrated genetic map with 4,285

single nucleotide polymorphisms, spanning 1,630 cM, was constructed with MAPpoly software. We

observed that bivalent configurations were the most abundant ones (51.0~72.4% depending on parent

and linkage group), though multivalent configurations were also observed (2.2~39.2%). Seven traits

were evaluated over four years (2006-8 and 2014) and quantitative trait loci (QTL) mapping was

carried out using QTLpoly software. We detected 21 QTL for 15 out of 27 trait-year combination

phenotypes. A hotspot on linkage group 5 was identified as QTL for maturity, plant yield, specific

gravity and internal heat necrosis resistance over different years were co-located. We found over 500

genes around QTL peaks, including those on chromosome 5 that have been previously implicated in

maturity (*StCDF1*) and tuber formation (*POTH1*). These QTL regions can be investigated further in

order to allow genomics-assisted breeding in tetraploid potato.

1   Introduction

Potato is the third most cultivated food crop in the world after rice and wheat, with more than 17.6

million hectares cultivated worldwide in 2018, and a production exceeding 368.2 million tons (FAO,

2020). Yet advancement in the breeding of new varieties is still relatively slow. 'Russet Burbank',

which still accounts for a large amount of the acreage planted in the United States, was developed in

the early 1900s (Bethke *et al.*, 2014), and 'Atlantic', a major chipping variety grown in the country,

was originally released in 1976 (Webb *et al.*, 1978). One of the reasons of such relatively slow

breeding process is the relative complexity of the potato genome. Cultivated potato, *Solanum*

tuberosum, is a highly heterozygous autotetraploid ($2n = 4x = 48$) outcrossing crop, which makes it

very difficult to align potentially beneficial characteristic combinations through conventional

breeding methods. In fact, many agronomic traits are polygenic, and show continuous distributions

(Ghislain and Douches, 2020).

Much of the genome complexity has also impacted on the correct assessment of allelism from

molecular markers, impeding linkage map construction and quantitative trait loci (QTL) mapping.

Research has been conducted over the years to develop molecular techniques to facilitate potato

breeding. However, since the first molecular genetic maps of potato were published (reviewed by

Mann *et al.*, 2011), very few breeding programs have begun to employ marker-assisted selection

(MAS) routinely (Slater *et al.*, 2014). Some of the exceptions to this are for genetically simpler traits

such as potato cyst nematode resistance (Schultz *et al.*, 2012) and potato virus Y (Hämäläinen *et al.*,

1997). More complex traits such as yield are still elusive, and much more work needs to be done.

The early linkage maps constructed for potato have incorporated qualitatively scored molecular

markers (Mann *et al.*, 2011). However, this traditional presence-absence based scoring limited not

only the ability to explore the full range of segregating markers, but also to build integrated genetic

maps for full-sib populations (Luo *et al.*, 2004). Advances in molecular technologies have been made

available to breeders with the publication of the diploid potato genome sequence (Xu *et al.*, 2011)

and the subsequent development and release of the Illumina Infinium® 8,303 Potato Array (Felcher *et*

al., 2012) through the USDA-NIFA Solanaceae Coordinated Agricultural Project (SolCAP). The

original potato array, which was subsequently upgraded to a 12k form, was made up of 8,303 single

nucleotide polymorphisms (SNPs) and has been shown to give very good coverage of the published

potato genome (Felcher *et al.*, 2012; Sharma *et al.*, 2013). Utilizing Illumina's technology of dual

colored fluorescents to label the different nucleotides allows for scoring allele dosage (Schmitz

Carley *et al.*, 2017; Zych *et al.*, 2019). Along with the development of an array that uses dosage-

sensitive SNPs, new analytical methods have recently been developed to effectively utilize this

information in reconstructing linkage maps (Hackett *et al.*, 2014; Bourke *et al.*, 2018; Mollinari and

Garcia, 2019) and mapping QTL (Hackett *et al.*, 2016; Chen *et al.*, 2018) in autotetraploid species.

Novel linkage and QTL mapping approaches (Mollinari *et al.*, 2020; da Silva Pereira *et al.*, 2020)

have enabled the exploration of new horizons when studying genetics of polyploid species.

Particularly, Mollinari *et al.* (2020) presented a detailed characterization of the inheritance pattern in

an even more complex autopolyploid species, the hexaploid sweetpotato, *Ipomoea batatas* ($2n =$

$6x = 90$). As the inheritance pattern from parents to progeny had been unraveled, da Silva Pereira *et*

al. (2020) were able to estimate identity-by-descent (IBD)-based additive relationship along the

genetic map, which was ultimately used in a random-effect model approach, facilitating multiple

QTL mapping. In tetraploid potato, double reduction has been documented for several mapping

populations (e.g. Bourke *et al.*, 2015), but these studies lack a comprehensive assessment of

multivalent formation and preferential chromosome pairing. In addition, single-QTL models used so

far may have hindered the discovery of putative loci underlying the variation of agriculturally

important traits in the species.

The advancement of molecular technologies and analytical methods, coupled with newly collected

data, have made it valuable to re-analyze the B2721 potato mapping population, derived from the

'Atlantic' cultivar (McCord *et al.*, 2011a, 2011b; Schumann *et al.*, 2017). We performed SNP dosage

calling, created a fully phased, integrated linkage map, and carried out multiple QTL-based analyses

for yield, foliage maturity, dry matter, specific gravity, and skin texture evaluated over four years.

This has allowed us to infer the tetraploid inheritance mechanisms from a linkage analysis

perspective and to better understand these traits at the genetic level as well as to bring breeders closer

to MAS.

2　Material and Methods

2.1　Mapping population and field experiment

The B2721 family consisted of 156 progenies and was created by crossing 'Atlantic' × B1829-5. This

bi-parental mapping population was chosen based on its segregation for internal heat necrosis (IHN),

a non-pathogenic physiological disorder characterized by brownish spots in the tuber parenchyma

(Yencho *et al.*, 2008). 'Atlantic' is susceptible to IHN, whereas B1829-5 is an advanced round white

resistant clone from the USDA-ARS Beltsville potato breeding program. The B2721 population also

segregates for a wide range of agronomic traits of interest, and these will be our focus in this work,

although IHN related traits will also be considered. B2721 was previously evaluated in 2006, 2007

and 2008 (McCord *et al.*, 2011a, 2011b) and additional data was collected in 2014 (Schumann *et al.*,

2017).

The field designs for years 2006, 2007 and 2008 were described by McCord *et al.* (2011a,b). Briefly,

in 2006, the population was planted in an unreplicated trial with six plants per clone, whereas in 2007

and 2008, the population was planted in two replications with 10 plants per plot in a randomized

complete block design, the same design also used in 2014. All experiments were carried out at the

Tidewater Research Station (35°52'20" N, 76°39'33" W) in Plymouth, North Carolina, in a

Portsmouth fine sandy loam. In 2014, untreated seed pieces were planted on March 21 and the

harvest date was July 15 (a total of 116 days). Standard crop production methods used for potato

cultivation in eastern North Carolina were used for all experiments. At harvest, plots were dug using

a chain digger and harvested by hand. Tubers were then washed, culled for excessive rots and

defects, and phenotype measurements were obtained as described next.

2.2 Phenotypic data and analyses

Phenotyping for all IHN-related and agronomic traits in 2006, 2007 and 2008 was described

previously (McCord *et al.*, 2011b). In 2014, in addition to IHN incidence (NI) and IHN severity (NS)

(Schumann *et al.*, 2017), the B2721 population was evaluated for five other traits: plant yield (PY),

foliage maturity (FM), dry matter (DM), specific gravity (SG) and skin texture (ST). PY was

measured as total plot weight minus the tubers culled divided by 10 (number of plants per plot) in

kg/plant. FM was measured as the area under the senescence progress curve (Shaner and Finney,

1977), based on foliage ratings taken on June 9, June 25 and July 2, using a scale of 0 (0%

yellowing) to 5 (100% senesced) with half point increments. For DM determinations, tubers were

quartered from stem end to bud end using an Easy Wedger, Model N55550-4 (NEMCO Inc.,

Hicksville OH). For each plot, one quarter from four different tubers was placed into plastic Whirl-

Pak® bags. Samples were then weighed, frozen at –20 °C, lyophilized, and weighed again. DM was

calculated as the proportion between dry and fresh weights for each sample, in percentage. SG

calculated using the formula $SG = w_{air}/(w_{air} - w_{water})$ by using the air weight (w_{air}) and water

weight (w_{water}) for each plot. ST was rated on a 0 (smooth) to 3 (russeted) scale. Finally, NI was

calculated as the proportion of tubers showing any sign of IHN over all evaluated tubers, and NS was

rated on a 9 (no IHN signs) to 1 (completely necrotic) scale and averaged out for all evaluated tubers

(Schumann *et al.*, 2017).

In order to estimate mean-basis broad-sense heritability (H^2) (Holland *et al.*, 2003), a joint analysis

with data from 2007, 2008 and 2014 (replicated trials) was carried out, where years, blocks within

years, genotypes and genotype-by-year interactions were treated as random effects. Except for 2006

data (single observation), adjusted means were obtained separately for each trait-year combination

based on a mixed model with genotypes and blocks as fixed and random effects, respectively, plus

the random residual error using ASReml-R v. 4.1.0 (Butler *et al.*, 2018). The phenotype means were

named after a trait abbreviation followed by two-digit year (trait-year combination, e.g. PY06) and

later used for QTL mapping purposes. Progeny variation within each phenotype was explored with

boxplots from R package ggplot2 (Wickham, 2016). Pearson's correlations were computed using R

v. 3.6.0 (R Core Team, 2019), and graphical visualization through network and correlation plots were

obtained using R packages corrr v. 0.4.0 (Ruiz *et al.*, 2019) and corrplot v. 0.84 (Wei *et al.*, 2017).

As an approach to investigate genotype-by-environment (GE) interaction, genotype (G) plus GE

interaction (GGE) biplot analysis (Yan and Kang, 2003) was carried out using R package

GGEBiplots v. 0.1.1 (Dumble *et al.*, 2017) based on the standardized means.

2.3   Genotypic data, dosage calling and linkage map construction

Genotyping was performed using the Illumina Infinium® 8,303 Potato Array (Felcher *et al.*, 2012).

The intensity of each fluorescent labeled nucleotide was read using the Illumina iScan Reader and

imported into GenomeStudio. The ordered pair of normalized intensities (x, y) from the two allelic

variants was transformed into polar coordinates $\theta = \frac{2}{\pi} \operatorname{atan}\left(\frac{y}{x}\right)$ and $r = \sqrt{x^2 + y^2}$ to proceed with

the dosage calling of the parents and offspring using the R package ClusterCall (Schmitz Carley *et*

al., 2017). Uninformative markers as well as those resulting in unclassified parents were filtered out

resulting in 5,599 SNPs.

Allele dosage from these markers were imported into MAPpoly v. 0.1.0 (Mollinari and Garcia, 2019)

together with their respective chromosome positions (*S. tuberosum* genome v. 4.0.3). After testing for

segregation distortion ($\sim 8.9 \times 10^{-6}$ using Bonferroni correction) and filtering out redundant

markers, the pairwise recombination fraction of the 4,812 SNPs (~11.6 million pairs) were computed

for all possible linkage phases and the most likely configuration was selected. Markers were

assembled into 12 linkage groups (LGs) using the Unweighted Pair Group Method with Arithmetic

Mean (UPGMA) algorithm. Since 91.2% of markers within the groups coincided with their

respective chromosomes, groups were formed using exclusively the genome information. Markers

were ordered according to the reference genome (Xu *et al.*, 2011) and phased using function

'est_rf_hmm_sequential'. Briefly, this function sequentially inserts markers into the map using two-

point information to eliminate unlikely linkage phases (LOD < 10.0), and for the remaining

configurations, the multipoint LOD score is used (LOD < 10.0). If the insertion of a marker still

results in less than 60 linkage phase configurations, a next marker was inserted and tested in the

multiple configurations until the map is completed. In the final step, the multipoint recombination

fraction was estimated allowing a global genotyping error of 5%.

Next, the conditional probabilities of the 36 possible genotypes were obtained for every cM of the

linkage map for all individuals in the offspring. These probabilities were used to assess the meiotic

process that formed the offspring and for further QTL analysis. Probabilistic profiles for pairing

behavior were obtained for all LGs for the three possible meiotic pairing configurations, i.e., *ab*/*cd*,

ac/*bd*, and *ad*/*bc*, where *a*, *b*, *c* and *d* denote the homologs in parent 'Atlantic'. The notation *ab*/*cd*

indicates that homolog *a* paired with *b*, and *c* paired with *d*, for example. The same reasoning applies

to parent B1829-5 with homologs *e*, *f*, *g* and *h*. Also, the probabilistic haplotypes of all individuals in

the offspring were reconstructed and the crossing-over points and respective homologs involved in

the exchange were inferred. Using this information and the heuristic algorithm presented by

Mollinari *et al.* (2020), recombination chains were assembled for all offspring individuals, and the

homologs involved in each meiosis were assessed. Recombination chains with more than two

homologs involved imply that a multivalent formation was present during the meiosis.

2.4   QTL mapping and gene search

The conditional genotype probabilities calculated from the inferred genetic map were used to a

compute sib-pair IBD-based additive relationship matrix for every cM position in the map. Using the

R package QTLpoly v. 0.2.0 (da Silva Pereira *et al.*, 2020), we detected putative QTL positions using

the random-effect multiple interval mapping (REMIM) model for each phenotype (trait-year

combination) based on a forward-backward procedure. First, putative QTL were consecutively added

to the model based on a forward search with a relatively relaxed genome-wide significance threshold

($\alpha = 0.20$). Then, a backward elimination step was carried out under a more stringent threshold ($\alpha =$

0.05). The variance components associated with the putative QTL were tested using linear score

statistics (Qu *et al.*, 2013), whose associated *P*-values were log-transformed to facilitate QTL profile

visualization with $LOP = -\log_{10}(P)$. The genome-wide significance was assessed using a score-

based resampling method (Zou *et al.*, 2004). A final multiple QTL model was fitted with the selected

positions, and QTL genotype best linear unbiased predictions (BLUPs) were used to compute

additive allele effects (Kempthorne, 1955). QTL heritability (h^2_{QTL}) was calculated as the ratio

between the variance associated with the QTL and the total variance.

In order to compare different approaches, we also ran the fixed-effect interval mapping (FEIM)

model (Hackett *et al.*, 2014) as implemented in the QTLpoly package (da Silva Pereira *et al.*, 2020).

Using the same conditional genotype probabilities, a single-QTL model was used to fit six additive

effects (as one additive effect from each parent is taken as reference) at every cM position. The

model with a fitted QTL was compared to a null model (with no QTL) using likelihood ratio tests

(LRT). LRT statistics were converted into LOD scores, and QTL were declared when the LOD score

reached a threshold ($\alpha = 0.05$) based on 1,000 permutation tests (Churchill and Doerge, 1994). In

both REMIM and FEIM analyses, a window size of 20 cM was used to avoid that two closely linked

positions were declared as QTL. Approximate 95% QTL support intervals were estimated by

dropping 1.5 from the QTL peak LOP (REMIM) or LOD (FEIM) (da Silva Pereira *et al.*, 2020).

Based on the QTL peaks from REMIM, we searched for candidate genes in the region delimited by

markers on the left and on the right of the QTL peak or within 200 kbp each side from the QTL peak,

whichever was the largest, on the *S. tuberosum* v. 4.03 genome (ST4.03) using the PhytoMine tool

(https://phytozome.jgi.doe.gov/phytomine/) by Phytozome v. 12 (Goodstein *et al.*, 2012). Finally, we

used the Web Gene Ontology Annotation Plot (WEGO) v. 2.0 (Ye *et al.*, 2018;

http://wego.genomics.org.cn/) to count and plot classifications of Gene Ontology (GO) terms for

annotated genes within our QTL regions.

3   Results

3.1   Trait correlations and GE interactions

Based on the phenotype means (see Supplementary File S1), variation was found for all evaluated

traits over the four years of evaluation (Figure 1A), indicating that they were segregating in the

B2721 population. As the parental means were within the population phenotypic range, we could

observe transgressive segregation for all trait-year combinations. PY06 has shown quite distinct

median and broader variation in comparison to the following years, which is likely because there

were no replicates in 2006, so that the environmental error could not be modeled. In contrast, FM14

exhibited a narrower variation when compared to the previous years. This was rather expected, since

in 2014 maturity evaluation was carried out in a shorter time-window (24 days) in comparison to

2007 and 2008 (30 days). The overall change in ST14 scores when compared to the previous years

was likely due to differences in using evaluator scoring. Individuals seemed to be less affected by

IHN in 2008, as NS08 and NI08 have shown distinct variation and median in comparison to the other

years. Broad-sense heritability (H^2) ranged from 0.39 (DM) to 0.81 (SG), with similarly high values

for PY (0.78), FM (0.70) and ST (0.79). For IHN-related traits, NS (0.74) showed a relatively higher

heritability when compared to NI (0.66).

The correlation among variables was explored using a network plot (Figure 1B), where path color

and transparency represent the signal and strength of the correlation, respectively. The variable

positioning is given by multidimensional scaling that leverages the magnitude of the correlations, i.e.

the closer the variables, the higher the absolute values of their correlations (Ruiz *et al.*, 2019). In

general, the phenotype of the same trait evaluated in different years clustered together. As expected,

NS was close to NI as well as DM to SG, because each one of these trait pairs are different ways of

assessing the same characteristic (IHN response and solids content, respectively). FM appeared

centrally in this plot as it was correlated with most variables either negatively (e.g. with PY and NI)

or positively (e.g. with SG and DM). Note that for this network plot, a subset of 144 full-sibs with

complete observations among all traits was ultimately used. For correlation estimates based on

pairwise complete observations, please see Supplementary Figure S1 and Supplementary File S2.

Even though data from 2006 was originated from a single replication, they were in general agreement

with the adjusted means from the other years. DM observations were the least correlated ones across

years in comparison to all the other trait sets (see Supplementary Figure S1).

GGE biplot was used to investigate the existence of GE interaction (Figure 1C), where each year was

considered an environment. The distribution of full-sibs along the first two principal components,

which accounted for 43% of the variation, did not show any clustering. This was expected as

mapping populations usually consists of non-selected individuals. However, some full-sibs were

found to be more scattered in the fourth quadrant, indicating the most affected individuals by IHN

(i.e. high NI). In contrast, more individuals were found to be less scattered in the second quadrant,

where NS vectors lie, indicating smaller differentiation when full-sibs were less susceptible to IHN.

Relatively shorter vector lengths of PY, ST and DM08 indicated limited capacity to discriminate

genotypes using these traits, in comparison to longer vectors. Only relatively weak GE interaction

was observed, since phenotypes from different years consistently exhibited the same vector

orientation for each trait, with small angles between vectors. Greater angles between vectors of

phenotypes from 2006 (such as DM06, SG06 and NS06) and from the remaining years should be

treated carefully, as these 2006 measurements were based on single plots. The same vector

orientation for SG and DM indicated that these traits were positively correlated, whereas the opposite

orientations for NS and NI vectors revealed their negative correlations, as already mentioned.

Figure 1. B2721 mapping population phenotypes evaluated for four years (2006-8 and 2014). (A)

Boxplots showing distribution of full-sib means along with parental means. (B) Network plot

showing positive (blue) and negative (red) significant correlations ($|r| \geq 0.26$, $p < 0.001$). (C) GGE

biplot showing the distribution of full-sibs (green) and phenotypes (blue) on the two first principal

components. Traits: plant yield (PY), foliage maturity (FM), specific gravity (SG), dry matter (DM),

skin texture (ST), and internal heat necrosis severity (NS) and intensity (NI).

3.2   Linkage analysis and inheritance mechanisms

The normalized intensities from the Illumina Infinium® 8,303 Potato Array were obtained for the

B2721 population (see Supplementary File S3). The dosage calling procedure yielded 5,599

informative SNPs with 1.05% of missing data. Fourteen percent of the SNPs were filtered out due to

segregation distortion and redundancy resulting in 4,812 informative markers: 1,311 (27.3%)

simplex, 1,234 (25.6%) double-simplex, and 2,267 (47.1%) multiplex (see Supplementary Figure

S2). The complete linkage map (see Supplementary Table S1, Supplementary Figure S3 and

Supplementary File S5) consisted of 12 LGs with eight haplotypes each, with four homologs per

parent. A total of 4,285 markers spanned 1,629.99 cM in length, with an average density of 2.64

SNPs/cM. Linkage group lengths ranged from 106.20 cM (LG 5) to 205.88 cM (LG 1), with average

LG length of 135.83 cM. The scatterplots of the physical distance in *S. tuberosum* genome v. 4.03

versus genetic distance in the 12 LGs in B2721 population map is shown in Supplementary Figure

S4.

The probabilistic pairing profiles showed no preferential pairing between homologs in both parents

(see Supplementary Figure S5). Among all meiotic configurations, only 7.3% and 9.8% were

inconclusive for parents 'Atlantic' and B1829-5, respectively. From the remaining configurations

(Figure 2), the percentage of cases with no crossing-over varied from 6.8% (LG 1, parent 'Atlantic')

to 40.7% (LG 11, parent B1829-5), with 24.2% on average. Configuration involving two

chromosomes with at least one crossing-over (i.e. bivalent configurations), were the most abundant

varying from 51.0% (LG 11, parent B1829-5) to 72.4% (LG 9, parent 'Atlantic'), with a mean of

62.3%. Multivalent configurations, i.e., involving three or four homologs, ranged from 2.2% (LG 8,

parent B1829-5) to 39.2% (LG 1, parent 'Atlantic'), with 13.5% on average.

Figure 2. Distribution of number of homologs involved in a recombination chain.

3.3 Multiple QTL mapping and candidate genes

From the REMIM analysis, a total of 21 QTL were mapped in 15 out of 27 evaluated trait-year

combinations (Table 1; see Supplementary Figure S6). One QTL was detected for 10 phenotypes

each (for PY06, PY08, SG07, SG08, ST07, ST08, NS06, NS08, NI06 and NI08). The other five

phenotypes have shown two (for FM07, FM14, SG06 and NI07) or three (for FM08) QTL each. No

QTL were mapped for 12 phenotypes (PY07, PY14, SG14, DM06, DM07, DM08, DM14, ST06,

ST14, NS07, NS14 and NI14), mostly from years 2006 and 2014. Ten QTL were mapped on LG 5,

four on LG 1, three on LG 7, and one on LGs 2, 3, 4 and 9 each. QTL heritability (h^2_{QTL}) ranged from

7.1% (FM08 on LG 7) to 64.0% (FM08 on LG 5), including five QTL with higher ($64.0 \geq h^2_{\mathrm{QTL}} \geq$

36.0), 10 with moderate (29.8% $\geq h^2_{QTL} \geq$ 20.4%), and the remaining six QTL with lower

(18.6% $\geq h^2_{QTL} \geq$ 7.1%) heritability (Table 1; Figure 3). In comparison to REMIM results, FEIM

detected 17 QTL from 12 phenotypes (see Supplementary Figure S7). On the one hand, FEIM did not

identified QTL previously identified for PY06 (on LG 5), FM08 (on LGs 1 and 7), SG06 (on LG 3),

NI06 (on LG 1) and NI07 (on LGs 1 and 5). On the other hand, FEIM identified additional QTL for

SG06 (on LGs 8 and 10), ST08 (LG 9), NI14 (on LG 5) (see Supplementary Table S3). The

proportion of variance explained (PVE) by FEIM-derived QTL ranged from 13.0% to 52.7%.

Table 1. Random-effect multiple interval mapping (REMIM) for B2721 traits evaluated for four

years (2006-8 and 2014).

Trait[a]	QTL	Linkage group	Position (cM)	SI (cM)[c]	Score	P-value	σ^2_{QTL}[d]	h^2_{QTL}[e] (%)
PY06	1	5	32	13-42	137.41	4.70e–05	1.11e–01	20.4
PY08	1	5	46	0-53	168.95	7.21e–06	2.17e–02	36.0
FM07	1	5	26	18-29	355.99	<2.22e–16	3.36e+02	57.7
	2	7	66	52-70	167.43	8.98e–07	7.28e+01	12.5
FM08	1	1	9	0-32	101.58	2.16e–04	1.17e+02	7.4
	2	5	27	0-41	425.04	<2.22e–16	1.01e+03	64.0
	3	7	65	35-71	104.85	2.00e–04	1.13e+02	7.1
FM14	1	5	18	0-29	197.01	9.77e–08	7.12e+00	29.8
	2	7	66	54-70	138.45	1.57e–05	4.01e+00	16.8
SG06	1	2	97	34-117	125.63	6.65e–05	6.69e–06	15.9
	2	3	44	31-50	109.89	1.49e–04	8.94e–06	21.2
SG07	1	5	19	4-32	135.93	3.26e–05	1.39e–05	27.6
SG08	1	5	28	15-32	240.07	1.47e–08	4.24e–05	48.9
ST07	1	9	41	27-82	109.68	1.97e–04	1.59e–01	21.9
ST08	1	4	2	0-4	234.05	1.08e–08	6.32e–01	48.1
NS06	1	1	0	0-52	113.07	1.29e–04	1.17e–01	22.9
NS08	1	5	26	12-32	117.77	1.77e–04	2.84e–02	23.4
NI06	1	1	0	0-69	110.07	1.66e–04	9.28e+01	22.0
NI07	1	1	0	0-8	103.32	2.26e–04	1.39e+02	16.8
	2	5	25	8-34	134.43	3.35e–05	1.99e+02	24.2
NI08	1	5	27	12-32	131.28	6.54e–05	5.65e+01	28.5

310 [a]Traits: plant yield (PY), foliage maturity (FM), specific gravity (SG), skin texture (ST), and internal heat necrosis

311 severity (NS) and intensity (NI).

312 [b]~95% support interval (SI) based on $LOP - 1.5$, in centiMorgans.

313 [c]Variance component associated with the QTL (σ^2_{QTL}).

314 [d]QTL heritability (h^2_{QTL}), in percentage.

Figure 3. QTL identified for six traits evaluated for four years (2006-8 and 2014). Dots are plotted

according to QTL peak location along the linkage groups. Color scale represents *P*-values, and sizes

are proportional to QTL heritability (h^2_{QTL}). Traits: plant yield (PY), foliage maturity (FM), specific

gravity (SG), skin texture (ST), and internal heat necrosis severity (NS) and intensity (NI).

With 10 QTL on the proximal half of LG 5, this region was characterized as a QTL hotspot. The

major QTL on LG 5 for FM was detected for all three years of evaluation (2007-8 and 2014).

However, *LOP* profiles (see Supplementary Figure S6) showed suggestive QTL (at a lower *LOP* that

did not reach the threshold), for instance, on LG 5 for PY06, SG14 and NS14 and NI14. For the five

traits with QTL on LG 5 (PY, FM, SG, NS and NI), most QTL peaks were found at 29 cM and

support intervals ranged from 0 to 50 cM. Figure 4 depicts additive allele effects of co-located QTL

on LG 5 for these different traits evaluated in 2008. It is interesting to note that for the QTL sets

PY08.1/NI08.1 and FM08.2/SG08.1/NS08.1, major allele effects appeared in the same direction

within sets, but in opposite directions between them. As these QTL on LG 5 were responsible for

explaining significant portion of the variation for these traits (Table 1), the QTL-based predicted

means were highly correlated, either positively (PY with NI, and FM with SG and NS) or negatively

(PY/NI with FM/SG/NS) (see Supplementary Figure S8). It is worth mentioning that selecting

towards two out of four alleles per parent will tend to favor different sets of phenotypes depending on

the alleles. For example, selecting allele pair *ac* from 'Atlantic' and *eg* from B1829-5 will likely

increase PY and NI and decrease FM, SG and NS. Additive effects for all QTL can be found on

Supplementary File S6.

Figure 4. Additive allele effects for co-located QTL on LG 5 for five traits evaluated in 2008. Letters

represent each parental haplotype from the linkage map ('Atlantic' = *a* through *d*, B1829-5 = *e*

through *h*). Traits: plant yield (PY), foliage maturity (FM), specific gravity (SG), and internal heat

necrosis severity (NS) and incidence (NI).

As the QTL support intervals comprised too large of a region to manually curate (>13 Mb per QTL,

on average), we focused our candidate gene search on the region within the markers flanking the

QTL peak (~400 kb, on average) (see Supplementary Table S2). A total of 533 annotated genes on *S.*

tuberosum v. 4.03 genome were within these QTL regions, with 342 genes assigned to at least one

GO term (see Supplementary File S7). From a total of 594 GO terms, 309 referred to a different set

of molecular functions (mostly related to catalytic activity or binding), 208 to a biological process

(greatly represented by metabolic or cellular processes), and 77 to a cellular component (such as

membrane or another cell part) (see Supplementary Figure S9).

4   Discussion

For the same B2721 population, two separate linkage maps, one for each parent, have been

previously built using amplified length polymorphisms (AFLP) and a few simple sequence repeat

(SSR) markers, allowing varied degrees of map integration within parents (McCord *et al.*, 2011b).

Subsequently, Schumann *et al.* (2017) have shown how the dosage-sensitive SNP array contributed

to map integration and haplotyping in the B2721 population. In comparison, their map contained

3,427 SNPs and spanned 1,397.86 cM, which is a relatively smaller but less saturated than the current

map with 4,285 SNPs distributed along 1,629.99 cM. The ability to incorporate more markers with

MAPpoly was possibly due to (*i*) the multipoint-based algorithm that considers all parental homologs

and marker dosages simultaneously, and (*ii*) the genome-assisted approach for marker ordering.

Schumann *et al.* (2017) used the software TetraploidSNPMap (Hackett *et al.*, 2016), which relies on

(*i*) simplex markers to assemble homologs, bridging these homologs afterwards using multidose

markers, and (*ii*) two-point genetic distances to obtain the marker order by using an algorithm such as

the Multidimensional Scaling (MDS) algorithm (Preedy and Hackett, 2016).

As expected, the probabilistic pairing profiles showed no preferential pairing between homologs in

both parents (see Supplementary Figure S5), likely due to the cultivated potato's autotetraploid

nature. Consequently, alleles present in different homologs in the same homology group had an equal

chance to recombine with alleles from other loci, amplifying the range of possible genotypes when

compared to diploids or allotetraploids. In addition, most of the parental meiotic configurations based

on the B2721 population (62.3% on average) were inferred as bivalents since they involved two

homologs exchanging segments during the metaphase I (Figure 2). These results are in accordance

with recent findings published by Choudhary *et al.* (2020). These authors presented a map of meiotic

stages in *S. tuberosum* in one diploid and three tetraploid varieties using fluorescence *in situ*

hybridization (FISH). They concluded that bivalent chromosome associations are the most common

in metaphase I of tetraploid potatoes. Likewise, the broad range of multivalent signature rates

observed in our study (2.2~39.2%) was also observed by Choudhary *et al.* (2020) (7~48%).

Similarly, as their observations differed depending on variety and chromosome, in our study, the

variation occurred by parents and by linkage groups. Thus, our meiotic assessment, although not as

precise as a cytological analysis and prone to sampling errors, can be very helpful, serving as a proxy

to evaluate meiotic configurations by using a straightforward expansion of linkage analysis.

Previous QTL mapping analyses in the B2721 mapping population were based on an interval

mapping model, and the 95th percentile ($\alpha = 0.05$) of maximum LOD scores from 1,000 permutation

tests per phenotype was used to declare QTL, with several additional suggestive QTL (below this

threshold) also being recorded (McCord *et al.*, 2011a, 2011b). The lack of integration within and

between parental maps may have resulted in the same region being declared as a QTL in separate

parental haplotypes. For instance, for plant yield, McCord *et al.* (2011b) listed a total of seven QTL

across three years, PY06 (one), PY07 (five) and PY08 (two), from both parental maps, in addition to

11 suggestive ones. The current REMIM analyses resulted in a single co-located QTL region on LG

5 for PY06 and PY08 (Table 1; Figure 3), with suggestive QTL PY07 and PY14 (see Supplementary

Figure S6). QTL on LG 5 for tuber yield have also been reported in both diploid (DM 1-3 516 R44 ×

RH89-039-16, $n = 96$) (Manrique-Carpintero *et al.*, 2015) and tetraploid ('Liberator' × W4013-1,

$n = 110$) (Rak *et al.*, 2017) potato mapping populations.

For foliage maturity, McCord *et al.* (2011b) and Schumann *et al.* (2017) described major QTL for

years 2007-8 on LG 5 as well as the one on LG 7. In the current study, another QTL on LG 1 was

identified for FM08. Despite the differences regarding phenotypic distribution for FM14 (Figure 1A),

we were able to find QTL on LGs 5 and 7 (Figure 3). This LG 5 region is well known as a main

source of QTL in potato. Prior to the release of the SNP potato array, a major QTL for plant maturity

had already been reported on chromosome 5, also linked to a late blight (*Phytophthora infestans*)

resistance locus (reviewed by Danan *et al.*, 2011). More recently, Massa *et al.* (2018) noticed that the

same region on LG 5 also underlies the variation of several tuber quality-related traits in addition to

resistance to early blight (*Altenaria tenuis*) and Verticillium wilt (*Verticillium* spp.) for a russet

mapping population ('Rio Grande Russet' × 'Premier Russet', $n = 162$). In fact, for IHN-related

traits, three QTL were identified on LG 5, one for each NS08, NI07 and NI08, using REMIM. FEIM

identified an additional significant QTL on LG 5 for NI14 (see Supplementary Table S3 and

Supplementary Figure S7). Previously, IHN-related QTL had been reported as suggestive only

(Schumann *et al.*, 2017), and this difference in significance is likely due to an improved map

construction.

A major QTL on LG 5 was also found to co-locate for SG07 and SG08, while it appeared only as

suggestive for SG14 and with no evidence for SG06. There were no QTL for dry matter either.

Specific gravity and dry matter are supposedly related with solid content in the tuber. In fact, the

correlation among their adjusted means were generally high (up to 0.81 between DM07 and SG07;

see Supplementary Figure S8 and Supplementary File S2), but the lack of co-located QTL limited

conclusions regarding any shared genetic control. It is worth to mention that their broad-sense

heritability values (H^2) differed greatly (0.39 for DM and 0.81 for SG), so that the ability to detect

QTL for DM may have been affected because smaller proportion of the total variance could be

attributed to genetic variance. Other mapping populations have revealed QTL on LG 5 for DM

(12601ab1 × 'Stirling', $n = 227$) (Bradshaw *et al.*, 2008) and SG (Manrique-Carpintero *et al.*, 2015),

where somewhat distinct heritability values were found (0.81 for DM and 0.46 for SG).

Skin texture was a trait which diverged the most from the other traits regarding QTL locations,

specifically for not showing enough statistical evidence for QTL on LG 5 (see Supplementary Figure

S6). In diploid potatoes, skin texture is believed to be controlled by three loci (Jong, 1981). In our

B2721 population, we detected two QTL (one for ST07 and ST08 each), in regions on LGs 4 and 9

where no other traits have shown evidence for QTL in our study. That is, the apparent strong

correlation observed between ST07 and FM07 ($r = -0.44, p < 0.001$), for example, could not be

attributed to genetics, as correlation between QTL-based predicted means for these two traits was

close to zero ($r = -0.07, p = 0.40$; see Supplementary Figure S8).

When performing QTL discovery in outbred polyploid populations, in addition to genome

complexity, one must keep two factors in mind. First, cultivars and elite clones, such as those used as

B2721 parents, have usually been selected such that alleles for characteristic of interest were led to

fixation. For instance, as observed by Schumann *et al.* (2017), IHN-related traits were concentrated

towards resistance (Figure 1A). It is likely that one or a few major loci underlying IHN resistance

have their alleles fixed within a parent (likely B1829-5, in our case) or, at least, that these allele

effects do not differ greatly, hence their contribution towards resistance is more or less the same. For

the QTL on LG 5 for NI08, the allele contributions to the population mean (~6.1%) were ±2.5% at

most (Figure 4). In other words, these effects would not change the fact that most individuals would

be less susceptible (NI < 10%) as resistance could have been already conferred by other undetectable

QTL. In F_1 populations, the fact that two parental genotypes are contrasting does not directly imply

that major QTL will be detected. For instance, considering a biallelic QTL, a cross between two

contrasting parents, $QQQQ \times qqqq$, would not segregate (i.e. 100% $QQqq$). In order to be detectable,

alleles need to be contrasting within parent, e.g. $Qqqq \times qqqq$ or $Qqqq \times Qqqq$ in the simplest cases,

resulting in segregations such as 50% $Qqqq$: 50% $qqqq$ or 25% $QQqq$: 50% $Qqqq$: 25% $qqqq$,

respectively.

If there are segregating QTL alleles, another factor that influences detection is small population sizes

that may not allow to separate signal from noise properly. Consequently, thresholds for declaring a

QTL are usually required to be more stringent in order to avoid false positives. In general, tetraploid

potato QTL mapping studies have been performed in relatively small population sizes (~150 or less

individuals). In these cases, even very low p-values and high QTL heritability estimates, such as

those found for FM07, FM08 and SG08, should be treated carefully as they can be relatively biased

due to sampling. Finally, reduced population sizes also limit the ability to test closely linked versus

pleiotropy for co-located QTL across different traits or environments accurately.

Among the candidate genes (see Supplementary File S7), GO terms that overlapped DNA and

protein binding functions with regulation of biological process were of particular interest. In total, 19

genes were found to have the same two GO terms (GO:0003677 and GO:0006355) attributed to the

S. tuberosum CYCLING DOF (DNA-binding with one finger) FACTOR 1, *StCDF1*

(PGSC0003DMG400018408, ST4.03ch05:4539029..4541329), which has been described as the

transcription factor underlying the major QTL for maturity on LG 5 (Kloosterman *et al.*, 2013). Such

a region also includes yield-related QTL, where we found the *Potato homeobox 1* gene, *POTH1*

(PGSC0003DMG400013493, ST4.03ch05:9248372..9258054), that appears to regulate tuber

development in potato via StBEL5-TOPH1 heterodimer (Rosin *et al.*, 2003). Finally, QTL

underlying IHN-related traits were also found in the same LG 5 hotspot, where a homeobox-

associated leucine zipper, *StHOX40* (PGSC0003DMG400030494, ST4.03ch05:4264626..4267698),

was identified. This gene was found to be either up or downregulated under heat and drought stresses

in potato, respectively (Li *et al.*, 2019), which are believed to be two important triggers of IHN

(Yencho *et al.*, 2008).

Other genes, such as *StERF6* (PGSC0003DMG400016651, ST4.03ch06:33827698..33832740) and

StNAC046 (PGSC0003DMG400031266, ST4.03ch05:4946232..4948779), were characterized as

transcription factors from AP2/ERF and NAC families, respectively, which have been implicated in a

wide range of regulation processes in plants (Olsen *et al.*, 2005; Mizoi *et al.*, 2012). A gene from the

NAC family, *StNAC032* (PGSC0003DMG400002824, ST4.03ch04:245125..246607), was retrieved

from the LG 4 region of a major QTL for skin texture. Several genes from the NAC family, including

StNAC032, were found to be expressed in the tuber skin and appeared to be involved in suberin and

associated wax biosynthesis (Soler *et al.*, 2020), which could contribute to skin texture, as well as be

involved with response to drought and other biotic and abiotic stresses (Singh *et al.*, 2013).

In plants, a single QTL region may be found to underlie the variation of more than one phenotype at

once, which can potentially explain the correlation among traits. A QTL hotspot has been identified

in the short arm of chromosome 5 of potato (Danan *et al.*, 2011; D'hoop *et al.*, 2014), and our B2721

population stuck to the rule. A major QTL for maturity was found to overlap loci underlying the

variation for other traits such as yield, specific gravity and IHN-related resistance, as described

previously (McCord *et al.*, 2011a, 2011b; Schumann *et al.*, 2017) and corroborated with newly

analyzed phenotypic data. Pleiotropy is one possible explanation of physiologically correlated traits,

such as maturity, yield and specific gravity (as a proxy of solid content, thus contributing to yield).

Transcription factors are known to have this property in which a single molecular function may be

involved in multiple biological processes (He and Zhang, 2006). For instance, a DNA binding

element from the HAP complex, *Ghd8*, was found to play pleiotropic roles in regulating yield,

flowering and plant height in rice (Yan *et al.*, 2011). On the other hand, the presence of QTL for

yield-related traits and pathogenic (e.g. late blight) and non-pathogenic (e.g. IHN) potato disorders in

the same hotspot on LG 5 could well be due to closely linked genes. Another example from rice has

suggested the multigenic nature of a major QTL for grain yield under drought, where the central role

of a NAM transcription factor, *OsNAM$_{12.1}$*, in concert with extra co-localized genes, influenced other

traits such as root and panicle branching, and transpiration efficiency (Dixit *et al.*, 2015). In potato,

our list of candidate genes can be used in further experiments to dissect such a QTL hotspot.

Genomics-assisted breeding for quantitative, more complex traits could be implemented upon

validation of these QTL neighboring markers in breeding populations of tetraploid potato. In

addition, both meiotic and trait genetic architecture assessments derived from our analyses can be

particularly useful in refining simulations of potato breeding programs (Gaynor *et al.*, 2020).

5   Author Contributions

501   GCY designed the experiments and supervised the project. MJS and MEC performed field

502   experiments and collected data. GSP, MM and ZBZ analyzed data. GSP drafted the manuscript with

503   major contributions from MM. All authors read and approved the manuscript.

6   Funding

506   This project was funded by USDA-ARS [58-1245-3-307, 2014-34141-22266] and USDA-NIFA

507   [2013-34141-21392]. Scholarships were awarded to MJS by Pioneer and Monsanto, and to GSP and

508   MM by Bill & Melinda Gates Foundation [OPP1052983]. Currently, GSP holds a scholarship from

509   PrInt-CAPES, Brazil [88887.369781/2019-00].

7   Acknowledgments

512   We would like to thank all the staff at the North Carolina Department of Agriculture and Consumer

513   Services Tidewater Research Station, Plymouth, NC, for their hard work and dedication to quality

514   research in aiding with the field aspect of this study. We also appreciate Michigan State University

515   Potato Breeding and Genetics Program for their assistance with the Illumina Infinium® 8,303 Potato

516   Array.

8   Competing Interests

519   The authors declare no competing financial interests.

9   Data Archiving

522   The datasets analyzed and scripts used in this study can be found in the supplementary material and

523   at the study GitHub page (https://github.com/mmollina/B2721_map). Both MAPpoly

524   (https://github.com/mmollina/MAPpoly) and QTLpoly (https://github.com/guilherme-

525   pereira/QTLpoly) R packages are available at GitHub.

10   References

Bethke PC, Nassar AMK, Kubow S, Leclerc YN, Li XQ, Haroon M, *et al.* (2014). History and

Origin of Russet Burbank (Netted Gem) a Sport of Burbank. *Am J Potato Res* **91**: 594–609.

Bourke PM, van Geest G, Voorrips RE, Jansen J, Kranenburg T, Shahin A, *et al.* (2018).

polymapR—linkage analysis and genetic map construction from F1 populations of outcrossing

polyploids. *Bioinformatics*: 1–7.

Bourke PM, Voorrips RE, Visser RGF, Maliepaard C (2015). The double-reduction landscape in

tetraploid potato as revealed by a high-density linkage map. *Genetics* **201**: 853–863.

Bradshaw JE, Hackett CA, Pande B, Waugh R, Bryan GJ (2008). QTL mapping of yield, agronomic

and quality traits in tetraploid potato (Solanum tuberosum subsp. tuberosum). *Theor Appl Genet*

116: 193–211.

Butler DG, Cullis BR, Gilmour AR, Gogel BJ, Thompson R (2018). *ASReml-R Reference Manual*

Version 4. Hemel Hempstead.

Chen J, Zhang F, Wang L, Leach L, Luo Z (2018). Orthogonal contrast based models for quantitative

genetic analysis in autotetraploid species. *New Phytol* **220**: 332–346.

Choudhary A, Wright L, Ponce O, Chen J, Prashar A, Sanchez-Moran E, *et al.* (2020). Varietal

variation and chromosome behaviour during meiosis in Solanum tuberosum. *Heredity (Edinb)*.

Churchill GA, Doerge RW (1994). Empirical threshold values for quantitative trait mapping.

Genetics **138**: 963–71.

D'hoop BB, Keizer PLC, Paulo MJ, Visser RGF, van Eeuwijk FA, van Eck HJ (2014). Identification

of agronomically important QTL in tetraploid potato cultivars using a marker-trait association

analysis. *Theor Appl Genet* **127**: 731–748.

Danan S, Veyrieras JB, Lefebvre V (2011). Construction of a potato consensus map and QTL meta-

analysis offer new insights into the genetic architecture of late blight resistance and plant

maturity traits. *BMC Plant Biol* **11**.

Dixit S, Kumar Biswal A, Min A, Henry A, Oane RH, Raorane ML, *et al.* (2015). Action of multiple

intra-QTL genes concerted around a co-localized transcription factor underpins a large effect

QTL. *Sci Rep* **5**: 1–16.

Dumble S, Bernal EF, Villardon PG (2017). GGEBiplots: GGE Biplots with 'ggplot2'.

FAO (2020). FAOSTAT Crops.

Felcher KJ, Coombs JJ, Massa AN, Hansey CN, Hamilton JP, Veilleux RE, *et al.* (2012). Integration

of two diploid potato linkage maps with the potato genome sequence. *PLoS One* **7**: e36347.

Gaynor RC, Gorjanc G, Hickey JM (2020). AlphaSimR : Simulations An R-package for Breeding

Program. *bioRxiv*: 1–21.

Ghislain M, Douches DS (2020). The Genes and Genomes of the Potato. In: Campos H, Ortiz O

(eds) *The Potato Crop*, Springer, pp 139–162.

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, *et al.* (2012). Phytozome: a

comparative platform for green plant genomics. *Nucleic Acids Res* **40**: D1178--D1186.

Hackett CA, Boskamp B, Vogogias A, Preedy KF, Milne I, Vogogias T, *et al.* (2016).

TetraploidSNPMap: software for linkage analysis and QTL mapping in autotetraploid

populations using SNP dosage data. *J Hered* **108**: 438–442.

Hackett CA, Bradshaw JE, Bryan GJ (2014). QTL mapping in autotetraploids using SNP dosage

information. *Theor Appl Genet*.

Hämäläinen JH, Watanabe KN, Valkonen JPT, Arihara A, Plaisted RL, Pehu E, *et al.* (1997).

Mapping and marker-assisted selection for a gene for extreme resistance to potato virus Y.

Theor Appl Genet **94**: 192–197.

He X, Zhang J (2006). Toward a molecular understanding of pleiotropy. *Genetics* **173**: 1885–1891.

Holland JB, Nyquist WE, Cervantes-Martínez CT (2003). Estimating and interpreting heritability for

plant breeding: an update. In: Janick J (ed) *Plant Breeding Reviews*, John Wiley & Sons: New

Jersey Vol 22, pp 9–112.

Jong H (1981). Inheritance of russeting in cultivated diploid potatoes. *Potato Res* **24**: 309–313.

Kempthorne O (1955). The correlation between relatives in a simple autotetraploid population.

Genetics **40**: 168–174.

Kloosterman B, Abelenda JA, Gomez MDMC, Oortwijn M, De Boer JM, Kowitwanich K, *et al.*

(2013). Naturally occurring allele diversity allows potato cultivation in northern latitudes.

Nature **495**: 246–250.

Li W, Dong J, Cao M, Gao X, Wang D, Liu B, *et al.* (2019). Genome-wide identification and

characterization of HD-ZIP genes in potato. *Gene* **697**: 103–117.

Luo ZW, Zhang RM, Kearsey MJ (2004). Theoretical basis for genetic linkage analysis in

autotetraploid species. *PNAS* **101**: 7040–7045.

Mann H, Iorizzo M, Gao L, D'Agostino N, Carputo D, Chiusano ML, *et al.* (2011). Molecular

Linkage Maps: Strategies, Resources and Achievements. In: Bradeen JM, Kole C (eds)

Genetics, Genomics and Breeding of Potato, CRC Press: Enfield, pp 68–89.

Manrique-Carpintero NC, Coombs JJ, Cui Y, Veilleux RE, Robin Buell C, Douches D (2015).

Genetic map and QTL analysis of agronomic traits in a diploid potato population using single

nucleotide polymorphism markers. *Crop Sci* **55**: 2566–2579.

Massa AN, Manrique-Carpintero NC, Coombs J, Haynes KG, Bethke PC, Brandt TL, *et al.* (2018).

Linkage analysis and QTL mapping in a tetraploid russet mapping population of potato. *BMC*

Genet **19**: 1–13.

McCord PH, Sosinski BR, Haynes KG, Clough ME, Yencho GC (2011a). QTL mapping of internal

heat necrosis in tetraploid potato. *Theor Appl Genet* **122**: 129–142.

McCord PH, Sosinski BR, Haynes KG, Clough ME, Yencho GC (2011b). Linkage mapping and

QTL analysis of agronomic traits in tetraploid potato (Solanum tuberosum subsp. tuberosum).

Crop Sci **51**: 771–785.

Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K (2012). AP2/ERF family transcription factors in plant

abiotic stress responses. *Biochim Biophys Acta - Gene Regul Mech* **1819**: 86–96.

Mollinari M, Garcia AAF (2019). Linkage Analysis and Haplotype Phasing in Experimental

Autopolyploid Populations with High Ploidy Level Using Hidden Markov Models. *G3*

Genes|Genomes|Genetics **9**: 3297–3314.

Mollinari M, Olukolu BA, Pereira GS, Khan A, Gemenet D, Yencho GC, *et al.* (2020). Unraveling

the Hexaploid Sweetpotato Inheritance Using Ultra-Dense Multilocus Mapping. *G3*

Genes|Genomes|Genetics **10**: 281–292.

Olsen AN, Ernst HA, Leggio L Lo, Skriver K (2005). NAC transcription factors: structurally distinct,

functionally diverse. *Trends Plant Sci* **10**: 79–87.

Preedy KF, Hackett CA (2016). A rapid marker ordering approach for high-density genetic linkage

maps in experimental autotetraploid populations using multidimensional scaling. *Theor Appl*

Genet **129**: 2117–2132.

Qu L, Guennel T, Marshall SL (2013). Linear score tests for variance components in linear mixed

models and applications to genetic association studies. *Biometrics* **69**: 883–892.

R Core Team (2019). R: A Language and Environment for Statistical Computing.

Rak K, Bethke PC, Palta JP (2017). QTL mapping of potato chip color and tuber traits within an

autotetraploid family. *Mol Breed* **37**.

Rosin FM, Hart JK, Horner HT, Davies PJ, Hannapel DJ (2003). Overexpression of a Knotted-Like

Homeobox Gene of Gibberellin Accumulation 1. *Plant Physiol* **132**: 106–117.

Ruiz E, Jackson S, Cimentada J (2019). corrr: Correlations in R.

Schmitz Carley CA, Coombs JJ, Douches DS, Bethke PC, Palta JP, Novy RG, *et al.* (2017).

Automated tetraploid genotype calling by hierarchical clustering. *Theor Appl Genet* **130**: 717–

726.

Schultz L, Cogan NOI, Mclean K, Dale MFB, Bryan GJ, Forster JW, *et al.* (2012). Evaluation and

implementation of a potential diagnostic molecular marker for H1-conferred potato cyst

nematode resistance in potato (Solanum tuberosum L.). *Plant Breed* **131**: 315–321.

Schumann MJ, Zeng ZB, Clough ME, Yencho GC (2017). Linkage map construction and QTL

analysis for internal heat necrosis in autotetraploid potato. *Theor Appl Genet* **130**: 2045–2056.

Shaner G, Finney RE (1977). The Effect of Nitrogen Fertilization on the Expression of Slow-

Mildewing Resistance in Knox Wheat. *Phytopathology* **67**: 1051–1056.

Sharma SK, Bolser D, de Boer J, Sonderkaer M, Amoros W, Carboni MF, *et al.* (2013). Construction

of reference chromosome-scale pseudomolecules for potato: integrating the potato genome with

genetic and physical maps. *G3 Genes|Genomes|Genetics* **3**: 2031–47.

da Silva Pereira G, Gemenet DC, Mollinari M, Olukolu BA, Wood JC, Diaz F, *et al.* (2020). Multiple

QTL Mapping in Autopolyploids: A Random-Effect Model Approach with Application in a

Hexaploid Sweetpotato Full-Sib Population. *Genetics* **215**: 579–595.

Singh AK, Sharma V, Pal AK, Acharya V, Ahuja PS (2013). Genome-wide organization and

expression profiling of the NAC transcription factor family in potato (solanum tuberosum L.).

DNA Res **20**: 403–423.

Slater AT, Cogan NOI, Hayes BJ, Schultz L, Dale MFB, Bryan GJ, *et al.* (2014). Improving breeding

efficiency in potato using molecular and quantitative genetics. *Theor Appl Genet* **127**: 2279–

2292.

Soler M, Verdaguer R, Fernández-Piñán S, Company-Arumí D, Boher P, Góngora-Castillo E, *et al.*

(2020). Silencing against the conserved NAC domain of the potato StNAC103 reveals new

NAC candidates to repress the suberin associated waxes in phellem. *Plant Sci* **291**: 110360.

Webb RE, Wilson DR, Shumaker JR, Graves B, Henninger MR, Watts J, *et al.* (1978). Atlantic: A

new potato variety with high solids, good processing quality, and resistance to pests. *Am Potato*

J **55**: 141–145.

Wei T, Simko V, Levy M, Xie Y, Jin Y, Zemla J (2017). corrplot: Visualization of a Correlation

Matrix.

Wickham H (2016). *ggplot2: elegant graphics for data analysis*. Springer.

Xu X, Pan S, Cheng S, Zhang B, Mu D, Ni P, *et al.* (2011). Genome sequence and analysis of the

tuber crop potato. *Nature* **475**: 189–195.

Yan W, Kang MS (2003). *GGE Biplot Analysis: a graphical tool for breeders, geneticists, and*

agronomists. CRC Press: Boca Raton.

Yan WH, Wang P, Chen HX, Zhou HJ, Li QP, Wang CR, *et al.* (2011). A major QTL, Ghd8, plays

pleiotropic roles in regulating grain productivity, plant height, and heading date in rice. *Mol*

Plant **4**: 319–330.

Ye J, Zhang Y, Cui H, Liu J, Wu Y, Cheng Y, *et al.* (2018). WEGO 2.0: A web tool for analyzing

and plotting GO annotations, 2018 update. *Nucleic Acids Res* **46**: W71–W75.

Yencho GC, McCord PH, Haynes KG, Sterrett SBR (2008). Internal heat necrosis of potato - A

review. *Am J Potato Res* **85**: 69–76.

Zou F, Fine JP, Hu J, Lin DY (2004). An efficient resampling method for assessing genome-wide

statistical significance in mapping quantitative trait loci. *Genetics* **168**: 2307–2316.

Zych K, Gort G, Maliepaard CA, Jansen RC, Voorrips RE (2019). FitTetra 2.0 - Improved genotype

calling for tetraploids with multiple population and parental data support. *BMC Bioinformatics*

20: 1–8.