Effects of a combined enrichment intervention on the behavioural and physiological welfare of

captive Asiatic lions (*Panthera leo persica*)

Sitendu Goswami[1*], Shiv Kumari Patel[1], Shwetank Pandit[2], Riyaz Kadivar[2], Praveen Chandra

Tyagi[1], Pradeep Kumar Malik[1], Samrat Mondol[1*]

[1] Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand, India.

[2] Sakkarbaug Zoological Garden, Junagadh, Gujarat, India

[*] Corresponding authors: Samrat Mondol, Ph.D., Animal Ecology and Conservation Biology

Department, Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand 248001. Email-

samrat@wii.gov.in

Sitendu Goswami, Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand 248001.

Email- sitendu@wii.gov.in

23   **Running head:** Impacts of enrichment on Asiatic lions.

Abstract

The endangered Asiatic lion (*Panthera leo persica*) is currently distributed as a single wild

population of 670 individuals and ~400 captive animals globally. Although the captive lions are

major hope for the species' long-term conservation through repatriation, their welfare status and

management practises need research attention. To this end, we tested the efficacy of feeding,

sensory and manipulable enrichment interventions on the welfare of Asiatic lions at the

conservation breeding centre of Sakkarbaug Zoological Garden, Gujarat. We adopted a holistic

approach by measuring physiological and behavioural responses of 35 captive Asiatic lions,

divided into control (n=16) and test (n=19) groups. The test subjects approached feeding devices

first and manipulable devices for a longer duration. Manipulable devices were used

homogenously with two significant time peaks, but sensory devices were used sporadically

throughout the day with no discernible peak usage. The control subjects remained unchanged in

all welfare parameters compared to their pre-treatment levels. However, post-enrichment

behavioural assessments showed higher behaviour diversity (95% increase from the baseline

period), reduced enclosure zone bias (40.25% reduction) and aberrant repetitive behaviours

(80.68%) in test samples. Similarly, faecal corticosterone measures showed lower stress levels in

test samples (58% decrease), confirming significant improvement in all welfare indices than

control groups. These results have universal applicability to assess welfare indices of other

captive species in Indian zoos. We hope that the results will encourage zoo managers and

regulatory agencies to improve animal welfare practices.

Keywords: Animal welfare, conservation breeding initiative, captive animal management,

behaviour diversity, space usage, stereotypy, faecal corticosterone.

1. Introduction

Since the early Palaeolithic depictions of the "lowenmensch" by the Aurignacians (Chauvet et al., 1996), lions have been heralded across several cultures as emblems of man's relationship with nature (McCall, 1973). Once ubiquitous across south-western Asia between Syria and parts of eastern India (Ball, 1880; Joslin, 1984), they were hunted to extinction from large parts of their range by the late 19th century (Jhala et al., 2019; MacKenzie, 2017; Storey, 1991). During the 1860s, the Asiatic lion population in Gir forests, India was struggling to survive. Timely conservation measures along with a hunting ban were instrumental to the recovery of the species. Sakkarbaug zoo, founded in 1863, played a definitive role in the conservation of Asiatic lions by treating injured and diseased individuals and maintaining a viable captive stock for future repatriation. Subsequent affirmative conservation actions including an ex-situ conservation breeding programme (Smith, 1984) led to the species revival from the edge of extinction. Today the extant population of more than 500 wild Asiatic lions(Gujarat Forest Department, 2015) inhabit fragmented habitats scattered over a human dominated landscape in Gujarat(Gogoi et al., 2020), Western India, along with approximately 400 Asiatic lions spread across zoos in several countries, with more than 60% of the captive population residing in Indian zoos and conservation breeding centres (Srivastav et al., 2018). The survival and proliferation of this species relies on repatriation to insulate the extant population from future stochastic extinction events (Jhala et al., 2019). Although the conservation breeding programme has been successful in maintaining a genetically diverse stock, the welfare status and incumbent management practices for the animals has not received adequate research focus (Pastorino et al., 2017; Goswami et al., 2020).

Conservation breeding programmes for endangered species should be designed to promote species-typical behaviours and cognitive plasticity for better post-release fitness (Rabin, 2003).

The deleterious impacts of sterile captive environments manifest with loss of species-typical

behaviours and increase in psychosomatic disorders in animals(Broom, 2011; Dawkins, 2004;

Fraser, 1999; Rabin, 2003). Therefore, welfare-based management practices are vital for any

successful conservation breeding programme (Swaisgood, 2010). The benefits of an individual-

centric welfare evaluation (Joslin, 1984) complemented with targeted enrichment interventions

has been successfully demonstrated for several captive animals viz., ursids (Carlstead et al.,

1991; McGowan et al., 2010), felids (Powell, 1995; Suárez et al., 2017), canids (Cloutier and

Packard, 2014; Leonard, 2008), equids (Bulens et al., 2013), small mammals (Clark and Melfi,

2012; Vargas and Anderson, 1999), reptiles (DeGregorio et al., 2017), and amphibians (Michaels

et al., 2014). Apart from improving welfare, enrichment interventions have been shown to play

an important positive role in increasing post-release fitness in several species (Brown et al.,

2003; Rabin, 2003; Reading et al., 2013). Earlier study on welfare status of captive Asiatic lions

reported that individual variations (personality and rearing history) are associated with

differential welfare outcomes in lions housed under similar captive environments (Goswami et

al., 2020), necessitating individually tailored husbandry regimen for the animals. This study

focuses on identifying the areas for improvement of the current husbandry and management

practices for Asiatic lions.

Feeding, sensory and manipulable enrichments have been shown to improve the welfare of

captive felids (Powell, 1995; Van Metter et al., 2008). We tested the efficacy of a combined

enrichment intervention on the welfare of captive Asiatic lions at Sakkarbaug Zoological Garden

(SZG). We measured several behavioural (species-typical behaviour diversity, enclosure usage

and aberrant repetitive behaviours) and physiological (faecal corticosterone metabolites) welfare

indices as a response to enrichment interventions. This is the first controlled trial study to

incorporate both behavioural and physiological tools to measure the welfare status change in

response to enrichment interventions of Asiatic lions housed at a conservation breeding centre.

We hope that our findings assist in the improvement of incumbent husbandry and management

practices for the species.

2. Methods

2.1. Study area

We conducted this study at the Asiatic lion conservation breeding centre at SZG. The

conservation breeding programme of Asiatic lions was initiated in 1958 with nine founders and

has since proliferated to house more than 47 individuals housed at SZG with several breeding

pairs in other participating zoos across the world. Since SZG houses both captive-born and wild-

rescued individuals, it holds intrinsic value for the conservation of Asiatic lions. During the study

period, the off-display conservation breeding facility housed 47 individuals in 20 large

naturalistic enclosures. A schematic map of the facility is provided in Supplementary Figure 1.

2.2. Subjects and housing

We selected 15 enclosures housing 35 Asiatic lions for the present study. Some of the subjects

were wild-rescued (N = 19, Male = 11, Female = 8), while the rest were born in captivity (N =

16, Male = 3, Female = 13). Subjects were primarily adult animals (N = 31) with a few sub-

adults (N = 4) and were randomly assigned to test and control groups to ensure uniformity of

treatment. The test group consisted 19 subjects (Male = 7, Female = 12) the control group

consisted 16 subjects (Male = 7, Female = 9), respectively.

Asiatic lions were housed in 15 enclosures in pair (1:1 and 0:2) (N = 9) or heterosexual (1:2) (N

=6) configurations (Supplementary Table 1). All subjects were housed in the same enclosure for

at least a year with the same enclosure mate and were accustomed to the enclosure and

management practices. In most cases subjects were housed with related individuals. All

enclosures provided adequate space (above 400m^2/animal in general with some having

1500m^2/animal) for the animals. The naturalistic enclosures provided subjects with sufficient

natural vegetation cover, simulating the habitat of Asiatic lions while protecting subjects from

visitor disturbance. Enclosures were well covered with trees and shrubs, which provided subjects

with a good combination of shade and sun throughout the day. Fresh drinking water was made

available inside retiring cells adjoining the paddock area. Subjects were allowed free access to

the paddock area and retiring cubicles throughout the day and night. However, these enclosures

offered little in terms of novelty and cognitive enrichment. Our earlier study established low

enclosure space utilization, low behaviour diversity, and a high incidence of aberrant repetitive

behaviours prevalent in Asiatic lions housed at SZG (Goswami et al., 2020).

A group of 10 animal keepers and contractual workers carried out all husbandry work for the

lions. Keepers typically reported for duty at 0700 hours and cleaned all leftover food from

retiring cells during 0730-1100 hours. The leftover food items were weighed to ascertain the

food consumption by each individual. The keepers would provide the daily ration of buffalo

meat to the subjects between 1700-1800 hours at the feeding cubicle, where each lion was fed

separately. Keepers were also tasked with behavioural monitoring of the subjects and recorded

the commencement and cessation of mating events between subjects. The zoo managers and the

animal keepers met regularly to ascertain the welfare need of the captive animals based on

subjective evaluations. The approach to welfare assessment was preventative and based solely on

incidental keeper observations hence difficult to quantify and address. We expected to observe

the effect of novel enrichment devices on the behavioural and physiological welfare indices of

these captive animals.

2.3. Study design

We hypothesized that enrichment interventions would lead to significant improvement in

behavioural (Goswami et al., 2020)and physiological welfare measures for all test subjects in

contrast to the controls. We divided the subjects into control (N = 16) and treatment (N = 19)

groups by a blind draw of enclosures for enrichment interventions. We conducted the study in

two phases: baseline and post-enrichment. During the baseline period, we measured behaviour

for 14 days/subject and collected two fresh faecal samples/subject/week.   This was followed by

a one-week exposure to the enrichment intervention to get test subjects accustomed to the new

devices. During the testing period (14 days) we recorded the behaviour of test subjects as a

response to the daily enrichment intervention. Apart from the enrichment interventions, housing

and husbandry conditions remained unchanged for both groups throughout the course of the

study.

2.4. Data collection

2.4.1.   Behaviour observations

Two observers (viz., first author and field assistant) collected all behaviour data, which

necessitated accounting for inter-observer reliability. We created a detailed ethogram of the

study subjects from one month of ad-libitum behaviour sampling and compared the behaviour

data recorded from the same subject by both observers to ensure consistency (Goswami et al.,

2020). We commenced data collection only after inter-observer reliability levels remained

consistent across three consecutive sessions (Cronbach's alpha >0.9). We simultaneously video

recorded all behaviour sessions to aid in data entry and reduce errors. We recorded behaviour

during 0500-1100, 1200-1800, and 2200-0500 hours. We used instantaneous scan sampling

(Altmann, 1974)at 1-minute intervals to record all behavioural states and all occurrences of

behavioural events (Goswami et al., 2020). During every six-hour data collection period, we

recorded four hours of behaviour data with 30 minutes rest for observers after every hour. Data

for the observer-resting period was later tabulated from the video recordings. We collected 195 ±

12.3 hours of behaviour data/subject during the study period.

We measured three behavioural welfare indices: species-typical behaviour diversity

(Wemelsfelder et al., 2000), spread of participation index (Plowman, 2003) and aberrant

repetitive behaviours or stereotypy (Mason & Latham, 2004) to measure the welfare of Asiatic

lions (Goswami et al., 2020). We compared the behavioural welfare indices of all test and control

subjects across the baseline and enrichment period. Baseline information for diversity of

behaviour repertoire in control and test subjects was collected using Shannon-Weiner diversity

index (SWI) (Goswami et al., 2020). We measured enclosure usage patterns of subjects based on

the spread of participation index (SPI) (Plowman, 2003)to understand if the enclosures provided

enough complexity to meet their welfare requirement (Plowman, 2003; Ross and Shender, 2016).

We divided each enclosure (N = 15) into ten zones, i.e., three primary zones (proximal, medial,

and distal), which were subdivided into three secondary zones (left, middle, right), and the tenth

zone was the area adjacent to the entrance to feeding cubicles (Goswami et al., 2020). We

measured the proportion of aberrant repetitive behaviours (ARBs) or stereotypy as an indicator

of poor welfare and compared their prevalence between control and test subjects across

treatments.

2.4.2.   Physiological measures

We measured faecal corticosterone metabolites of all subjects to contrast the physiological

impacts of the enrichment interventions on control and test subjects. Faecal corticosterone is a

reliable physiological indicator of stress and compromised welfare in captive felids (Ruskell et

al., 2015; Schildkraut, 2016; Vaz et al., 2017; Young et al., 2004). We collected two fresh faecal

samples/week from each subject (both control and test) during the entire study period (pre-

enrichment and post-enrichment).

We collected all fresh faecal samples using the dry sampling approach (Biswas et al., 2019) and

stored in -20^{o}C freezer onsite and later transported to the Wildlife Institute of India in dry ice.

The samples were stored in the laboratory at -20^{o}C freezer until further processing. To control

for effects of moisture and diet we pulverized the frozen samples before lyophilizing (#FD-5,

Allied Frost, New Delhi, India) them for 72 hours prior to hormone extraction (Mondol et al.,

2020). Subsequently, we sieved lyophilized samples through a 0.5 mm stainless steel mesh to

obtain homogenized faecal powder. We thoroughly mixed the dried faecal powder and extracted

hormone by pulse-vortexing 0.1 grams of powder in 15 ml of 70% ethanol, followed by

centrifugation at 2200 rpm for 20 min (Mondol et al., 2020; Wasser et al., 2010). We collected

and stored the hormone extracts in 2 ml cryochill vials (1:15 dilution) and stored in -20^{o}C freezer

till further analyses.

We used corticosterone EIA kit (#K014, Arbor Assays, MI, USA) for corticosterone metabolite

estimation in faecal hormone extracts. Sample extracts were air-dried inside an incubator and

resuspended in assay buffer as per required dilutions. Each sample was assayed in duplicate

using kit protocol and the optical density was measured at 450 nm using an ELISA plate reader

(#GMB-580, Genetix Biotech Asia Pvt. Ltd., New Delhi, India). Hormone metabolite

concentration is interpolated using four parametric logistic (4PL) regression function in

GraphPad prism software version 5 (GraphPad Software, California, USA). Cross-reactivities of

the antibody are listed in Supplementary Table 2.

We tested for parallelism and accuracy to validate the corticosterone assay. We used dilutions of

pooled extracts from a random combination of male and female samples (N=20) to assess

reliable quantification of corticosterone at different concentrations and find optimal dilutions for

final assays (at 50% binding). We plotted the relative dose against percent bound hormones for

the pools and the standards and generated best-fit curve using 4PL regression, where parallel

slopes indicate similar immunoreactivity at different concentrations. For accuracy test, we spiked

corticosterone standards with equal volumes of diluted faecal extract of known concentration

(dilution level close to 50% bound from parallelism test) and assayed with standards. We plotted

the results as regression lines using observed and expected concentrations to show that faecal

contaminants were not interfering with assay accuracy at the tested dilution. We calculated inter-

and intra- assay coefficients of variation using repeated measures of same-pooled extract.

2.4.3.   Enrichment interventions

Enrichment interventions add cognitive complexity to enclosures and provide animals the

opportunity to express species-typical behaviours (Mellen and Shepherdson, 1997; Skibiel et al.,

2007). We used three types of enrichment devices: manipulable (Mellen and Shepherdson, 1997;

Powell, 1995), sensory (Skibiel et al., 2007), and feed (Powell, 1995; Skibiel et al., 2007) (Table

1). Manipulable enrichments included hanging lion-sam balls, burlap bags, wooden perches/

platforms, wooden planks etc., whereas the sensory enrichments included olfactory

augmentations such as scent trails made from the blood of buffalo, urine of unknown

conspecifics, and dung of other prey species (Sambar, *Rusa unicolor*) etc. We also installed

sensory enrichment (tactile) by wrapping rough coir rope on the bark of trees inside enclosures to

promote auto-grooming and scent-marking behaviours in the subjects. Nutritional enrichment

has been shown to garner the highest amount of attention in captive animals and can be useful

for addressing neophobia to enrichment devices in shy individuals (Powell, 1995; Resende et al.,

2009; Skibiel et al., 2007). We provided a range of nutritional enrichment devices, which

included frozen dressed chicken, buffalo tails that were suspended by natural fibre ropes from

trees at different parts of the enclosure. While manipulable and sensory enrichment devices were

quasi-permanent and required little daily care, nutritional enrichment devices needed to be

replenished every day. We installed and replenished enrichment devices between 0600-0700

hours at four enclosures every day. Keepers would arrive early and confine the subjects inside

retiring cells with food rewards, while the enrichment interventions were placed at designated

enclosures. We furnished all enclosure zones with the same combination of manipulable, sensory

and nutritional enrichment.

To calculate enrichment device preferences, we collated all scans during which Asiatic lions

interacted with the enrichment devices. We also recorded the type of behaviours that were

performed by the subjects while interacting with the enrichment devices viz., exploration, social

play, aggression, foraging and fear (Table 2). We counted the different types of behaviours

performed per enrichment device by all subjects.

2.5. Data analysis

We analysed all data in R v3.6.2 (R Core Team, 2020) with R studio(RStudio, 2020). We used

packages "tidyverse" (Wickham et al., 2019), "psych" (Revelle, 2019)and "dplyr(Wickham et

al., 2020),"ggpubr"(Kassambara, 2020), "sur"(Harel, 2020) and "ggplot2" (Wickham, 2011) to

calculate statistical outputs and create graphical summaries of the findings. We conducted tests

for normality and ensured that variances were similar between groups to select appropriate

parametrical and non-parametrical statistical analysis. We used unpaired t-test to compare the

welfare indices between test and control groups during baseline and post-enrichment

interventions. We used paired t-tests, to compare the differences in welfare indices within test

and control groups and measured the amount of time spent by test subjects on different types of

enrichment devices. Additionally, we calculated the effect size (Cohen, 1992, p. 199)of the

enrichment intervention for the test group, which is scale-independent and gives a measure of the

magnitude of difference between baseline and post-treatment conditions.

3. Results

3.1. Enrichment preferences of study subjects

During 14 days of enrichment intervention, we recorded 1,09,238 scans of test subjects (n=19)

using enrichment devices. The test subjects used manipulable enrichment devices for longer

durations (56.8%) compared to all other types viz., sensory (12.99%) and feeding (30.20%)

(Table 3). It was challenging to measure the usage levels of sensory enrichments accurately and

hence these values may have been under-reported in our results. When subjects were released

inside the enclosures, they approached the feed-based enrichment devices first followed by

manipulable and sensory devices. This trend continued throughout the intervention period. Feed-

based enrichment devices were extensively used during the first two hours of the day between

0700-0900 hours (Figure 1, Table 4a) and garnered little or no interest after they were depleted.

On the other hand, manipulable and sensory enrichment device usage peaked twice daily at 0700

and 1600 hours (Figure 1, Table 4a) with continued usage throughout the day.

In terms of enrichment directed behaviours, we found that the peak of aggression (6.78%), chase

(7.69%), exploratory (6.35%) and forage (7.66%) behaviours in test subjects occurred between

0700-0900 hours (Table 4b). The next peak of enrichment-directed behaviours occurred between

1300-1700 hours constituting mostly of play (20.66%) and exploratory behaviours (4.11%)

(Table 4b). Early in the day (0700-0900 hours), when subjects were actively exploring around

the enclosure for enrichment devices, aggressive dominant and displacement behaviours were

common but subsided as the day progressed (Figure 1, Table 4b). We found that manipulable

devices (11.51%) were more commonly associated with aggressive behaviours compared to

food-based enrichment (2.94%) (Table 3). We also observed that positive social behaviours were

more commonly associated with food-based enrichment devices (1.16%) rather than the

manipulable devices (0.35%) (Table 3). Subjects usually monopolized the manipulable

enrichment devices and showed high fidelity towards specific items. The feed-based enrichment

devices were associated with the highest proportion of chase behaviour (11.54%), resulting in

one animal monopolizing a device and chasing away others (Table 3). Some feed-based

enrichment devices were consumed within minutes (e.g. dressed chicken) while others required

more processing time (e.g. buffalo tail, frozen blood cubes). Once the feed enrichment devices

were completely utilized, the lions concentrated on the manipulable and sensory enrichments and

engaged in social and exploratory behaviours throughout the later parts of the day (Table 3,

Figure 1). The manipulable enrichment devices were associated with the largest diversity of

species-typical behaviours, which included 35.16% play behaviour (Table 3) and were more

commonly associated with aggressive interactions (11.51%) compared to all other enrichment

types. Aggressive behaviours occurred when enclosure mates tried to steal or use a certain

manipulable device from the lion that was playing with it. Since we installed more enrichment

devices than subjects in every enclosure, displaced individuals had the opportunity to interact

with other devices when displaced by conspecifics. Sensory enrichment devices contributed

primarily to exploratory behaviours (11.7%) that led to increased enclosure usage depicted by the

lowering of SPI levels (Table 3).

However, it is important to point out that the true impacts of different types of enrichment

devices cannot be assessed merely through the measuring the amount of time the animal spent

with it, but by comparing the welfare indices of test subjects with control and their own baseline

status. The results clearly demonstrate that the presence of different types of enrichment devices

was instrumental in mitigating monopolization of resources by dominant individuals.

3.2. Behavioural welfare indices

During pre-enrichment period, behaviour diversity levels of control subjects (M = 0.9, SD = 0.3)

were similar to that of the test subjects (M = 0.83, SD = 0.35). After enrichment interventions,

the behaviour diversity of control subjects remained unchanged (M = 0.77, SD = 0.35, t(30)=

1.11, p = 0.29) while that of the test subjects increased significantly (M = 1.62, SD = 0.18, t(36)

= -8.8, p <0.001, Cohen's d = 2.83) (Table 5 Figure 2a).

During the baseline, we observed high SPI values in test (M = 0.77, SD = 0.12) and control

groups (M = 0.82, SD = 0.09) indicating enclosure zone use bias. However, after enrichment

intervention the SPI values decreased significantly indicating, enclosure usage homogeneity (M

= 0.46, SD = 0.08) for test subjects (t(36) = 9.37, p <0.0001, Cohen's d = 3.03), while it

remained unchanged for control subjects (M = 0.8, SD = 0.13, t(30)= 0.5, p = 0.61) (Table 5,

Figure 2b).

During baseline, test (M = 12.32, SD = 6.03), and control subjects (M = 12.8, SD = 6) showed

similar levels of aberrant repetitive behaviours. After enrichment interventions, we observed a

significant decrease in ARBs in test subjects (M = 2.38, SD = 2.46, t (36) = 6.65, p < 0.0001,

Cohen's d= 2.15) while the levels remained unchanged for the control group (M = 13.31, SD = 6,

t(30) = 0.23, p = 0.81) (Table 5, Figure 2c).

3.3. Faecal corticosterone level

Parallelism and accuracy tests for corticosterone metabolites showed reliable measures from lion

faeces across different concentration ranges. Serial dilutions of faecal extracts paralleled the

standard curves (Figure 3a). There were no differences between slopes of standard and pooled

extract curves for corticosterone ($F(1,10) = 2.06$, $P = 0.182$), and the curves were significantly

different in their elevation ($F(1,11) = 66.25$, $P = 0.0001$). Accuracy tests produced slopes of 0.92

at working dilution of 1:60 (Figure 3b), suggesting that faecal extracts did not interfere with their

metabolite measurement precisions. Intra-assay coefficient of variation (CV) was 9.8%, whereas

inter-assay CV was 8.99% (Supplementary Table 2).

We compared the average faecal corticosterone levels of each subject across baseline and post-

enrichment phases. Compared to baseline levels (M = 7.74 μg/gm, SD = 3.2), the faecal

corticosterone measures of test subjects decreased significantly (M = 2.46 μg/gm, SD= 0.85,

$t(36) = 5.33$, p<0.0001, Cohen's d = 2.22), while it remained unchanged from baseline (M = 7.1

μg/gm, SD = 4.08) for control subjects (M= 6.96 μg/gm, SD = 3.93, $t(30) = 0.093$, p = 0.92)

(Table 5 , Figure 2d). These results confirm that the enrichment interventions led to significant

improvement in behavioural and physiological welfare indices of test subjects as compared to the

control group during pre-enrichment period.

4. Discussion

Animal welfare is not limited to the prevention of cruelty and amelioration of symptoms of

stress, but also accords the opportunity to express species-typical behaviour patterns (Broom,

2011). The National Zoo Policy of India (MoEF,India, 1998) espouses animal welfare with

captive animal management, yet welfare-centric management protocols remain divorced from

husbandry practices across several zoos. Although few and far between, welfare research from

Indian zoos report a prevalence of stereotypic behaviours (Goswami et al., 2020; Mallapur et al.,

2007; Sanyal, 1892; Vaz et al., 2017) and elevated faecal corticosterone levels (Vaz et al., 2017)

across several captive felids, and prescribe specie-appropriate enclosure design and enrichment

interventions as remedial measures (Goswami et al., 2020; Mallapur et al., 2007; Vaz et al.,

2017). Enrichment interventions have been shown to improve welfare conditions in several

captive felids including African lions (Mellen and Shepherdson, 1997; Powell, 1995; Skibiel et

al., 2007), yet enrichment protocols are seldom incorporated in the daily management of captive

Asiatic lions at Indian zoos. The Asiatic lion conservation breeding programme has successfully

maintained a viable captive population insulated from stochastic ecological events. Most

research pertaining to captive Asiatic lions investigate their genetic and demographic status

(Bagatharia et al., 2013; Pastorino et al., 2017) with only a handful mentioning their welfare

(Goswami et al., 2020; Mallapur et al., 2002; Pastorino et al., 2017; Vaz et al., 2017). Our study

is the first to assess the effects of enrichment protocols on the welfare of captive Asiatic lions.

Welfare encompasses both internal and external conditions affecting an animal and hence must

be measured using a multifarious approach. As highlighted by Miller et al. (2020) the focus on

negative welfare indicators has led to zoos adhering to minimum husbandry guidelines that

attempt to suppress symptoms of poor welfare rather than trying to improve existing conditions.

In this study, we measured two traditional negative welfare parameters like ARB (Mason and

Latham, 2004) and faecal corticosterone levels (Schildkraut, 2016; Vaz et al., 2017) along with

two positive welfare indicators (Miller et al., 2020) viz., spread of participation index (SPI)

(Cabana et al., 2018; Powell, 1995) and behaviour diversity (Pastorino et al., 2017). Previous

studies on captive African lions that establish the positive effects of enrichment interventions

have primarily relied on behavioural welfare indices (Martínez-Macipe et al., 2015; Ncube and

Ndagurwa, 2010; Powell, 1995; Regaiolli et al., 2019; Van Metter et al., 2008). Our study is the

first to showcase both the behavioural and physiological impacts of enrichment interventions in

captive Asiatic lions housed in a conservation breeding programme. The welfare evaluation

framework used in this study can be applied universally (for other species) to measure the

impacts of goal-oriented enrichment interventions.

While previous studies have tested the welfare impacts of food (Powell, 1995; Van Metter et al.,

2008), manipulable (Powell, 1995), sensory (Martínez-Macipe et al., 2015; Regaiolli et al., 2019)

and social stimulation (Leonard, 2008; Ncube and Ndagurwa, 2010) individually. This is the first

control trial study to test the efficacy of a combined enrichment strategy on the welfare of Asiatic

lions. This has allowed us to understand the temporal usage pattern of enrichment devices in

Asiatic lions, which can prove useful for tailoring enrichment strategies. For example, individual

test subjects showed high fidelity towards manipulable enrichment devices, which were also

associated with the highest amount of conspecific-directed aggression. Therefore, to reduce

stress and aggressive behaviours we recommend providing more manipulable devices than the

number of animals in an enclosure. Upon release, subjects rushed to the feeding enrichment

devices, therefore to prevent aggression and injury, it is important to spread such devices far

apart, which will reduce chances of monopolization and food-based aggression. While sensory

enrichment devices are designed to encourage exploratory behaviours in lions, the feed-based

enrichment devices create a positive association for these novel objects and reduce neophobia.

Finally, we found that manipulable enrichment devices allow subjects to interact and exert

control over the captive environment while engaging in species-typical behaviours. Our

combined enrichment strategy brought novelty in a captive environment and presented animals

with the choice to express species-typical behaviour patterns. All enrichment devices used in this

study can be locally sourced or fabricated with minimal effort and require less than thirty

minutes to install or replenish with a team of three people.

Post-enrichment intervention, we recorded a significant improvement in the welfare indices of

test subjects compared to the control subjects. Subjects that would normally stay near the retiring

cells and remain inactive all day started exploring and utilizing different parts of the enclosure

and followed by a significant increase in behaviour diversity levels. Faecal corticosterone levels

of test subjects decreased significantly from the baseline, while it remained unchanged for

control subjects, showing the inter-linkages between behavioural and physiological indices of

welfare. Our findings agree with existing research that associate enrichment interventions with

increased behavioural diversity (Rabin, 2003; Wemelsfelder et al., 2000), lowered SPI (Nogueira

et al., 2004; Rose and Robert, 2013; Traylor-Holzer and Fritz, 1985), reduction of ARB (Clark

and Melfi, 2012; Mallapur et al., 2007; Vaz et al., 2017), and reduction of stress levels

(Hutchinson et al., 2012; Marcon et al., 2018; Mitra and Sapolsky, 2009; Nazar and Marin,

2011). However, it is important to point out that due to logistic constraints (sample size, time

etc.) we did not address the variable responses of subjects with different personality types on

enrichment interventions. Earlier work on captive Asiatic lions indicated that lions with different

personality traits are likely to react differently to novel enrichment devices (Goswami et al.,

2020). We suggest that future studies should focus on addressing impacts of such individual

variations on differential responses and enrichment preferences with larger sample size. We

studied the largest captive population of Asiatic lions, but the methodologies of this study can be

scaled for smaller zoos housing fewer animals. Although we used at least three types of

enrichment devices per intervention at multiple enclosure zones for experimental uniformity, at a

smaller scale, zoo managers can try one enrichment device type per enclosure and rotate them

with a new one every week. Our results conclusively show that enrichment interventions can

lead to better behavioural and physiological welfare for Asiatic lions. We have delineated a

practical but holistic approach to enrichment interventions that can be implemented and tested

with little effort and manpower. With this study, we hope to empower and enable zoo managers

and regulatory agencies to mandate the incorporation of enrichment interventions into daily

husbandry regimen and proactively improve the welfare status of the animals under their care.

We sincerely hope that the results of this study contribute to the improvement in animal welfare

practices across ex-situ institutions at Indian zoos and conservation breeding programs.

5. Conclusion

The success of conservation breeding program depends on repatriating endangered animals to

reclaimed habitats, hence housing the animals in appropriate welfare conditions while

persevering behaviour diversity is of utmost importance. The Asiatic lion conservation breeding

program has successfully maintained a genetically diverse stock of the species in captivity that

has proliferated to an appreciable number. With the incorporation of welfare-centric management

practices, the Asiatic lions housed in the program can be ideal representatives of the species and

great candidates for future reintroduction programmes. Until that day of repatriation, we should

strive to create a cognitively enriching environment that preserves the species-specific traits of

the Asiatic lions. Like the endangered Asiatic lion, the Indian ex-situ institutions run

conservation breeding programmes for several endemic species. So far, the welfare status of

animals housed under conservation breeding programmes has received little research attention.

We hope this study encourages zoo managers and regulators to incorporate enrichment

interventions with animal management protocols.

Ethics statement

This study was conducted in accordance with CZA norms on the welfare of animals housed in conservation breeding centres. We did not conduct any invasive sampling procedures on the subjects for experimental setup and data collection. All necessary precautions were taken to ensure that enrichment interventions were integrated with the daily husbandry regimens of the test subjects.

Acknowledgements

We thank Dr. S.C. Pant (Principle Chief conservator of forests, Gujarat) for providing necessary permits and logistical support for this study. We acknowledge the staff members of the Sakkarbaug zoo for their help at every stage of the study. Our thanks to Mr. Salim Chuvan, head-keeper of Asiatic lions for his insights on the unique behavioural traits of the study subjects and Mr. Ashkar Bloch for his assistance during field data collection. We thank Dr. V. K. Varshney of the Chemistry and Bioprospecting Division, Forest research Institute for his laboratory facilities and Mr. Shubham Kumar for laboratory work. We also thank the Director, Dean and Research Coordinator for their continuous support. Zoological Society of London funded this study. Samrat Mondol was supported by Department of Science and Technology INSPIRE Faculty Award (IFA12-LSBM-47).

References

Altmann, J., 1974. Observational study of behavior: sampling methods. Behaviour 49, 227–266.

Bagatharia, S.B., Joshi, M.N., Pandya, R.V., Pandit, A.S., Patel, R.P., Desai, S.M., Sharma, A., Panchal, O., Jasmani, F.P., Saxena, A.K., 2013. Complete mitogenome of Asiatic lion resolves phylogenetic status within Panthera. BMC Genomics 14, 572. https://doi.org/10.1186/1471-2164-14-572

Ball, V., 1880. Observations on Lion-Breeding in the Gardens of the Royal Zoological Society of

Ireland. The Transactions of the Royal Irish Academy 28, 723–758.

Biswas, S., Bhatt, S., Paul, S., Modi, S., Ghosh, T., Habib, B., Nigam, P., Talukdar, G., Pandav,

B., Mondol, S., 2019. A practive faeces collection protocol for multidisciplinary research

in wildlife science. Current Science 116, 1878–1885.

https://doi.org/10.18520/cs/v116/i11/1878-1885

Broom, D.M., 2011. A history of animal welfare science. Acta Biotheoretica 59, 121–37.

https://doi.org/10.1007/s10441-011-9123-3

Brown, C., Davidson, T., Laland, K., 2003. Environmental enrichment and prior experience of

live prey improve foraging behaviour in hatchery reared Atlantic salmon. Journal of

Fish Biology 63, 187–196.

Bulens, A., Van Beirendonck, S., Van Thielen, J., Driessen, B., 2013. The enriching effect of

non-commercial items in stabled horses. Applied Animal Behaviour Science 143, 46–51.

Cabana, F., Maguire, R., Hsu, C.-D., Plowman, A., 2018. Identification of possible nutritional

and stress risk factors in the development of marmoset wasting syndrome. Zoo Biology

37, 98–106. https://doi.org/10.1002/zoo.21398

Carlstead, K., Seidensticker, J., Baldwin, R., Park, N.Z., 1991. Environmental Enrichment for

Zoo Bears. Zoo Biology 16, 3–16.

Chauvet, J.-M., Brunel Deschamps, E., Hillaire, C., 1996. Dawn of art: the Chauvet Cave: the

oldest known paintings in the world.

Clark, F.E., Melfi, V.A., 2012. Environmental enrichment for a mixed-species nocturnal

mammal exhibit. Zoo Biology 31, 397–413. https://doi.org/10.1002/zoo.20380

Cloutier, T.L., Packard, J.M., 2014. Enrichment options for African painted dogs (*Lycaon*

pictus). Zoo biology 33, 475–480.

Cohen, J., 1992. A power primer. Psychological bulletin 112, 155.

Dawkins, M.S., 2004. Using behaviour to assess animal welfare. Animal welfare 13, 3–7.

DeGregorio, B.A., Sperry, J.H., Tuberville, T.D., Weatherhead, P.J., 2017. Translocating

ratsnakes: does enrichment offset negative effects of time in captivity? Wildlife Research

44, 438–448.

Fraser, D., 1999. Animal ethics and animal welfare science: bridging the two cultures. Applied

Animal Behaviour Science 65, 171–189.

Gogoi, K., Kumar, U., Banerjee, K., Jhala, Y.V., 2020. Spatially explicit density and its

determinants for Asiatic lions in the Gir forests. PloS one 15, e0228374.

Goswami, S., Tyagi, P.C., Malik, P.K., Pandit, S.J., Kadivar, R.F., Fitzpatrick, M., Mondol, S.,

2020. Effects of personality and rearing-history on the welfare of captive Asiatic lions

(*Panthera leo persica*). PeerJ 8, e8425. https://doi.org/10.7717/peerj.8425

Gujarat Forest Department, S. c., 2015. 14th Lion Population Estimation Report – 2015. Gujarat

Forest Department, Sasan Gir, Junagadh, India.

Harel, D., 2020. sur: Companion to "Statistics   Using R: An Integrative Approach.

Hutchinson, K.M., McLaughlin, K.J., Wright, R.L., Ortiz, J.B., Anouti, D.P., Mika, A.,

Diamond, D.M., Conrad, C.D., 2012. Environmental enrichment protects against the

effects of chronic stress on cognitive and morphological measures of hippocampal

integrity. Neurobiology of learning and memory 97, 250–260.

Jhala, Y.V., Banerjee, K., Chakrabarti, S., Basu, P., Singh, K., Dave, C., Gogoi, K., 2019. Asiatic

lion: ecology, economics and politics of conservation. Frontiers in Ecology and Evolution

7, 312.

Joslin, P., 1984. The environmental limitations and future of the Asiatic lion. Journal of the

Bombay Natural History Society. Bombay 81, 648–664.

Kassambara, A., 2020. ggpubr: "ggplot2" Based Publication Ready Plots.

Leonard, D.A., 2008. An evaluation of environmental enrichment for two highly social and

endangered canid species, the African wild dog (*Lycaon pictus*) and Mexican wolf (*Canis*

lupus baileyi). Saint Louis University.

MacKenzie, J.M., 2017. The empire of nature: Hunting, conservation and British imperialism.

Mallapur, A., Qureshi, Q., Chellam, R., 2002. Enclosure Design and Space Utilization by Indian

Leopards (*Panthera pardus*) in Four Zoos in Southern India. Animal Welfare 5, 111–

124.

Mallapur, A., Waran, N., Sinha, A., 2007. A note on enrichment for captive lion-tailed macaques

(*Macaca silenus*). Applied Animal Behaviour Science 108, 191–195.

https://doi.org/10.1016/j.applanim.2006.11.020

Marcon, M., Mocelin, R., Benvenutti, R., Costa, T., Herrmann, A.P., de Oliveira, D.L.,

Koakoski, G., Barcellos, L.J., Piato, A., 2018. Environmental enrichment modulates the

response to chronic stress in zebrafish. Journal of Experimental Biology 221.

Martínez-Macipe, M., Lafont-Lecuelle, C., Manteca, X., Pageat, P., Cozzi, A., 2015. Evaluation

of an innovative approach for sensory enrichment in zoos: semiochemical stimulation for

captive lions (*Panthera leo*). Animal Welfare 24, 455–461.

Mason, G.J., Latham, N.R., 2004. Can't stop, won't stop: Is stereotypy a reliable animal welfare

indicator? Animal Welfare 13, 57–69. https://doi.org/10.2307/4493573

McCall, D.F., 1973. The prevalence of lions: kings, deities and feline symbolism in Africa and

Elsewhere. Paideuma 130–145.

McGowan, R.T.S., Robbins, C.T., Alldredge, J.R., Newberry, R.C., 2010. Contrafreeloading in

grizzly bears: implications for captive foraging enrichment. Zoo Biology 29, 484–502.

Mellen, J.D., Shepherdson, D.J., 1997. Environmental enrichment for felids: an integrated

approach. International Zoo Yearbook 35, 191–197.

Michaels, C.J., Downie, J.R., Campbell-Palmer, R., 2014. The importance of enrichment for

advancing amphibian welfare and conservation goals. Amphibian Reptile Conservation 8,

7–23.

Miller, L.J., Vicino, G.A., Sheftel, J., Lauderdale, L.K., 2020. Behavioral Diversity as a Potential

Indicator of Positive Animal Welfare. Animals 10, 1211.

https://doi.org/10.3390/ani10071211

Mitra, R., Sapolsky, R.M., 2009. Effects of enrichment predominate over those of chronic stress

on fear-related behavior in male rats. Stress 12, 305–312.

MoEF,India, 1998. National Zoo Policy.

Mondol, S., Booth, R.K., Wasser, S.K., 2020. Fecal stress, nutrition and reproductive hormones

for monitoring environmental impacts on tigers (*Panthera tigris*). Conservation

physiology 8, coz091.

Nazar, F.N., Marin, R.H., 2011. Chronic stress and environmental enrichment as opposite factors

affecting the immune response in Japanese quail (*Coturnix coturnix japonica*). Stress 14,

166–173.

Ncube, S., Ndagurwa, H.G.T., 2010. Influence of social upbringing on the activity pattern of

captive lion (*Panthera leo*)cubs: Benefits of behavior enrichment. Current Zoology 56,

389–394.

Nogueira, S.S., Bernardi, L.G., Nogueira-Filho, S.L., 2004. A note on comparative enclosure

facility usage by wild and captive-born capybaras (*Hydrochoerus hydrochaeris*). Applied

Animal Behaviour Science 89, 139–143.

Pastorino, G.Q., Viau, A., Curone, G., Pearce-Kelly, P., Faustini, M., Vigo, D., Mazzola, S.M.,

Preziosi, R., 2017. Role of personality in behavioral responses to new environments in

captive Asiatic lions (*Panthera leo persica*). Veterinary medicine international 2017.

Plowman, A., 2003. A note on a modification of the spread of participation index allowing for

unequal zones. https://doi.org/10.1016/S0168-1591(03)00142-4

Powell, D., 1995. Preliminary evaluation of environmental enrichment techniques for African

lions (*Panthera Leo*). Animal Welfare 4, 361–370.

R Core Team, 2020. R: A language and environment for statistical computing. R   Foundation

for Statistical Computing, Vienna, Austria.

Rabin, L.A., 2003. Maintaining behavioural diversity in captivity for conservation: natural

behaviour management. Animal Welfare 12, 85–94.

Reading, R.P., Miller, B., Shepherdson, D., 2013. The value of enrichment to reintroduction

success. Zoo Biology 32, 332–341. https://doi.org/10.1002/zoo.21054

Regaiolli, B., Rizzo, A., Ottolini, G., Miletto Petrazzini, M.E., Spiezio, C., Agrillo, C., 2019.

Motion illusions as environmental enrichment for zoo animals: a preliminary

investigation on lions (Panthera leo). Frontiers in Psychology 10, 2220.

Resende, L.S., Remy, G.L., Ramos Jr, V. de A., Andriolo, A., 2009. The influence of feeding

enrichment on the behavior of small felids (Carnivora: Felidae) in captivity. Zoologia

(Curitiba) 26, 601–605.

Revelle, W., 2019. psych: Procedures for Psychological, Psychometric, and Personality

Research. Evanston, Illinois.

Rose, P., Robert, R., 2013. Evaluating the activity patterns and enclosure usage of a little-studied

zoo species, the sitatunga (*Tragelaphus spekii*). Journal of Zoo and Aquarium Research

1, 14–19.

Ross, S.R., Shender, M.A., 2016. Daily travel distances of zoo-housed chimpanzees and gorillas:

implications for welfare assessments and space requirements. Primates 57, 395–401.

https://doi.org/10.1007/s10329-016-0530-6

RStudio, 2020. RStudio: An integrated development environment for R.

Ruskell, A.D., Meiers, S.T., Jenkins, S.E., Santymire, R.M., 2015. Effect of bungee-carcass

enrichment on behavior and fecal glucocorticoid metabolites in two species of zoo-

housed felids. Zoo Biology 34, 170–177. https://doi.org/10.1002/zoo.21192

Sanyal, R.B., 1892. A Hand-book of the Management of Animals in Captivity in Lower Bengal.

Printed at the Bengal Secretariat Press.

Schildkraut, R., 2016. Characteristation of positive welfare indices in captive african lions

(*Panthera leo*) (Master's Thesis). University of Sydney.

Skibiel, A.L., Trevino, H.S., Naugher, K., 2007. Comparison of several types of enrichment for

captive felids. Zoo Biology: Published in affiliation with the American Zoo and

Aquarium Association 26, 371–381.

Smith, G.L., 1984. International Studbook: Indian Lion, *Panthera leo persica* (Meyer, 1826).

Knoxville Zoological Park.

Srivastav, A., Nigam, P., Begum, N., 2018. Indian National Studbook of Asiatic Lion (*Panthera*

leo persica).

Storey, W.K., 1991. Big cats and imperialism: Lion and tiger hunting in Kenya and Northern

India, 1898-1930. Journal of World History 2, 135–173.

Suárez, P., Recuerda, P., Arias-de-Reyna, L., 2017. Behaviour and welfare: The visitor effect in

captive felids. Animal Welfare 26, 25–34.

Swaisgood, R.R., 2010. The conservation-welfare nexus in reintroduction programmes: a role for

sensory ecology. Animal Welfare 19, 125–137.

Traylor-Holzer, K., Fritz, P., 1985. Utilization of space by adult and juvenile groups of captive

chimpanzees (*Pan troglodytes*). Zoo Biology 4, 115–127.

https://doi.org/10.1002/zoo.1430040205

Van Metter, J.E., Harriger, M.D., Bolen, R.H., 2008. Environmental enrichment utilizing

stimulus objects for African lions (*Panthera leo leo*) and Sumatran tigers (*Panthera*

tigris sumatrae). Bios 79, 7–16.

Vargas, A., Anderson, S.H., 1999. Effects of experience and cage enrichment on predatory skills

of black-footed ferrets (*Mustela nigripes*). Journal of Mammalogy 80, 263–269.

Vaz, J., Narayan, E.J., Dileep Kumar, R., Thenmozhi, K., Thiyagesan, K., Baskaran, N., 2017.

Prevalence and determinants of stereotypic behaviours and physiological stress among

tigers and leopards in Indian zoos. PloS one 12, e0174711.

Wasser, S.K., Azkarate, J.C., Booth, R.K., Hayward, L., Hunt, K., Ayres, K., Vynne, C.,

Gobush, K., Canales-Espinosa, D., Rodríguez-Luna, E., 2010. Non-invasive

measurement of thyroid hormone in feces of a diverse array of avian and mammalian

species. General and comparative endocrinology 168, 1–7.

Wemelsfelder, F., Haskell, M., Mendl, M.T., Calvert, S., Lawrence, A.B., 2000. Diversity of

behaviour during novel object tests is reduced in pigs housed in substrate-impoverished

conditions.

Wickham, H., 2011. ggplot2. Wiley Interdisciplinary Reviews: Computational Statistics 3, 180–

185.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., 2019. Welcome to the Tidyverse. Journal of Open

Source Software 4, 1686.

Wickham, H., Henry, L., Miller, K., 2020. dplyr: A Grammar of Data Manipulation.

Young, K.M., Walker, S.L., Lanthier, C., Waddell, W.T., Monfort, S.L., Brown, J.L., 2004.

Noninvasive monitoring of adrenocortical activity in carnivores by fecal glucocorticoid

analyses. General and comparative endocrinology 137, 148–165.

Figure legends

Figure 1: Temporal pattern of enrichment device use and behavioural patterns by Asiatic lions

on (a) Feeding, (b) Manipulable, and (c) Sensory enrichment devices. Primary Y- axis represents

the observation period from 0700-1800 hours. The primary x-axis represents the scan counts

(log10 scale) of specific behaviours on each type of enrichment device. Hourly counts of

enrichment-directed behaviours are represented as colour bars on the horizontal axis.

Figure 2: Comparison of the welfare indices between test and control groups across pre- and

post-enrichment conditions. The four different welfare indices used in this study are (a)

Behaviour diversity; (b) Spread of participation index; (c) Aberrant repetitive behaviours (ARB)

and (d) Faecal corticosterone levels.

Figure 3: Standardization of lion faecal corticosterone through (a) parallelism and (b) accuracy

tests. The graphs show accurate measures of corticosterone from captive lion faecal samples.

Supplementary Figure 1: Schematic representation of enclosures at the SZG conservation

breeding center for Asiatic lions.

Table 1: Description of enrichment devices used in this study at enclosures

Sl. no	Enrichment device	Enrichment type	Description
1	Platform	Manipulable	A log platform made to enable subjects to access vantage points within the enclosure
2	Sam ball	Manipulable	A hard-plastic ball with no holes or perforations
3	Log branches	Manipulable	Tree logs sections with bark suspended from trees
4	Lion bite ball	Manipulable	A hard-plastic ball with a rattle inside suspended by a bungee rope with canvas sheet that acts as a bite
5	Buffalo shank	Feeding	Buffalo shank suspended from a zipline
6	Chicken carcass	Feeding	Frozen Chicken carcass suspended from a tree by natural fibers
7	Buffalo tail	Feeding	Skinned buffalo tail suspended from tree branches
8	Buffalo blood	Feeding	Frozen buffalo blood cubes, provided on a clean concrete platform
9	Star anise oil mixed with garlic	Sensory	Sprayed on tree trunks or used to create scent trails to other enrichment devices
10	Prey dung	Sensory	Sambar (*Rusa unicolor*) dung smeared on tree trunks or mixed with enclosure soil at certain zones
11	Conspecific urine	Sensory	Earth drenched in the urine conspecifics from nearby enclosures smeared on tree trunks, or mixed with enclosure substrate at certain zones

Table 1: Behaviours observed that are associated with different enrichment devices

Sl. no	Behaviour	Description	Category
1	Aggressive Growl	Baring teeth with short vocalizations directed at conspecifics or humans, while interacting with enrichment	Aggression
2	Charge	Run up to a conspecific at a rapid short charge as a result of being challenged while suing an enrichment device	Aggression
3	Strike	Hit conspecific with one paw, while defending an enrichment device	Aggression
4	Bite	Bite conspecific, while asserting control over an enrichment device	Aggression
5	Allogrooming	Lick conspecific, while interacting with enrichment device	Social
6	Nuzzling	Rub noses with conspecifics with soft grunts, and inviting them to play with enrichment device	Social
7	Roll	Rolling in the dirt with conspecific while interacting with a enrichment device	Social
8	Rub	Rub body or neck against the enrichment device	Explore
9	Sniffing	Raise head and sniff near a sensory enrichment	Explore
10	Flehmen	Bare teeth and sniff the air near an enrichment device	Explore
11	Scent mark	Spray urine or scratch the tree next to an enrichment device	Explore
12	Digging	Dig near an enrichment device	Explore
13	Scratching	Scratch the trees where enrichment devices are installed	Explore
14	Chase	Chase away conspecifics trying to use or steal enrichment device successfully or unsuccessfully. Does not result in an overtly aggressive interaction	Chase
15	Run	Run around the enclosure with the enrichment device in mouth. May or May not be chased by conspecific	Play
16	Stalk	Stalk an enrichment device	Play
17	Play	Kick around an enrichment device	Play
18	Drag	Drag enrichment device along the enclosure	Play
19	Bite	Bite enrichment device	Play
20	Eat	Eat a food based or sensory enrichment device	Forage
21	Lick	Lick enrichment devices intently	Forage
22	Neophobia	Approach enrichment device slowly and run away	Fear
23	Fearful growl	Growling at enrichment device, ears back and cowering	Fear
24	Circling	Warily circling around the enrichment device	Fear

Table 1: Variations in behavioural performances in different enrichment devices used in this study.

Behaviour category	Feed enrichment (%)	Manipulable enrichment (%)	Sensory enrichment (%)
Aggression	2.94	11.51	0
Chase	11.54	1.89	0
Explore	3.05	5.95	11.77
Fear	0	1.96	0
Forage	11.48	0	0
Play	0	35.16	1.21
Social	1.16	0.35	0
Total	30.20	56.8	12.99

Table 4: Temporal patterns of enrichment usage and behavioural categories in the test subjects (n=35) during enrichment intervention experiments. The upper part of the table presents the enrichment use pattern, whereas the lower part of the table shows behaviour categories.

Enrichment use pattern at temporal scale (%)							
Enrichment	**0700-0900**	**0900-1100**	**1100-1300**	**1300-1500**	**1500-1700**	**1700-1900**	**Total**
Feed	22.13	8.06	0	0	0	0	30.2
Manipulable	11.269	6.94	7.02	23.44	5.66	2.44	56.8
Sensory	1.9	2.0	3.18	0.99	3.81	1.09	12.99
Behaviour categories at temporal scale (%)							
Behaviours	**0700-0900**	**0900-1100**	**1100-1300**	**1300-1500**	**1500-1700**	**1700-1900**	**Total**
Aggression	6.78	1.91	1.97	0.95	1.91	1	14.57
Chase	7.69	4.31	0.92	0	0.15	0.30	13.40
Explore	6.35	2.8	3.71	2.39	4.11	1.4	20.78
Fear	0	0.49	0.16	0.48	0.16	0.65	1.96
Forage	7.66	3.83	0	0	0	0	11.48
Play	5.95	3.11	3.39	20.6	3.11	0.18	36.29
Social	0.86	0.53	0.08	0	0	0.03	1.52

Table 5: Summary statistics of welfare indices within control and test groups

Welfare indices	Pre-enrichment	Post-enrichment	t-value*	Significance (p-value)	Effect size (Cohen's d)
Control group (N=16)					
Behaviour diversity (SWI)	0.9±0.31	0.77±0.35	1.11	0.29	0.39
Spread of participation index (SPI)	0.82±0.09	0.8±0.13	0.51	0.61	0.17
Aberrant repetitive behaviours (ARB)	12.81±6	13.31±6.07	0.23	0.81	0.08
Faecal corticosterone measure	7.1±4.08	6.96±3.93	0.09	0.92	0.03
Test group (N = 19)					
Behaviour diversity (SWI)	0.83±0.35	1.62±0.18	-8.8	0.0001	2.83
Spread of participation index (SPI)	0.77±0.12	0.46±0.08	9.37	0.0001	3.03
Aberrant repetitive behaviours (ARB)	12.32±6.03	2.38±2.46	6.65	0.0001	2.15
Faecal corticosterone measure	7.74±3.25	2.46±0.85	6.85	0.0001	2.22

* For control group t(30) and test group t(36).

a.Behaviour diversity (SWI)

b. Spread of participation index (SPI)

c. Aberrant repetitive behaviour (ARB)

d. Faecal corticosterone

Pre-enrichment

Post-enrichment

a. Corticosterone Parallelism

b. Corticosterone Accuracy (1:60)

y=0.9229x + 11.75, R^2=0.9989

Table 1: Details of subjects included in the study, viz., house name, sex (M=Male, F=Female), origin (C=Captive, W=Wild), sex ratio for housing, enclosure size, space per animal, age in days, and group.

Sl no	Name	Sex	Origin	Sex ratio (M:F)	Enclosure size (m²)	Space/animal (m²)	Age (in days)	Group
1	A1	M	W	1:1	1300	650	3414.00	Control
2	Aftab	M	W	1:1	1200	600	3500.00	Control
3	Amal	M	C	1:2	1576	788	2214.74	Test
4	Ambica	F	C	0:2	1123	561.5	3984.56	Test
5	Amiya	F	C	0:2	1123	561.5	2037.56	Test
6	Ani	F	W	1:1	1600	800	1946.00	Test
7	Bahadur	M	C	1:3	1600	533.33	653.33	Test
8	AmrapurI	F	W	1:3	1600	533.33	800.33	Test
9	AmrapurII	F	W	1:3	1600	533.3	800.33	Test
10	Dhari	M	W	1:3	1100	366.6	619.50	Test
11	Dheer	M	W	2:0	1600	800	5389.08	Control
12	Gina	F	C	1:1	1100	550	1316.90	Control
13	Girm	M	W	1:1	1123	562.5	3337.92	Control
14	Hemal	M	W	1:2	1700	566.6	2029.00	Control
15	Hemali	F	W	1:2	1271	635.5	2149.00	Control
16	Jenifer	F	C	1:1	1700	850	1491.33	Test
17	Jesal	M	W	1:2	1300	433.3	5722.18	Test
18	Maheswari	F	C	1:2	1300	433.3	3411.26	Test
19	Mariyam	F	C	1:2	1300	433.3	3423.35	Test
20	Maytri	F	C	1:1	1123	562.5	3072.90	Control
21	Nagraj	M	W	1:2	6542	2180	3227.00	Control
22	Patvad	F	W	1:1	1200	600	3729.00	Control
23	Patvadm	M	W	1:1	1271	635.5	1537.00	Test
24	Radha	F	C	1:2	1576	525.3	1291.30	Test
25	Rani	F	C	1:2	1576	525.3	1291.30	Test
26	Ranita	F	C	1:2	6542	2180	2119.00	Control
27	Ranshi	F	W	1:2	6542	2180	5109.00	Control
28	Rudi	F	W	1:1	1600	800	4700.00	Control
29	Subhi	F	C	1:1	1700	850	2240.33	Test
30	Sujan	F	W	1:1	1300	650	2790.80	Control
31	Taukir	M	C	1:1	1303	651.5	2448.00	Control
32	Teeta	F	C	1:2	1700	566.6	2166.93	Test
33	Tejaswini	F	C	1:2	1700	566.6	4320.93	Test
34	Trakuda	M	W	1:2	1700	566.6	3956.93	Test
35	Veer	M	W	2:0	1600	800	5387.30	Control

Supplementary Table 2: Details of the lion faecal corticosterone assay conducted in this study.

Hormone	Assay method	Dilution	Slope (R^2)	Inter-assay CV	Intra-assay CV	Cross-reactivity
Corticosterone	EIA	1:60	0.92 (0.99)	9.0	9.8	100% with corticosterone, 12.3% with Desoxycorticosterone, 2.3% with Tetrahydrocorticosterone and <1% with Aldosterone, Cortisol, Progesterone, Dexamethasone, Corticosterone-21-Hemisuccinate, Cortisone and Estradiol