**Age-associated changes in lineage composition of the enteric nervous system regulate gut health and disease**

Subhash Kulkarni[1#], Monalee Saha[1], Alpana Singh[1], Jared Slosberg[2], Sushma Nagaraj[1], Laren Becker[3], Chengxiu Zhang[1], Alicia Bukowski[1], Zhuolun Wang[1], Guosheng Liu[1], Jenna Leser[1,] Mithra Kumar[1], Shriya Bakhshi[1], Matthew Anderson[4], Mark Lewandoski[4], Elizabeth Vincent[3], Loyal A. Goff [2, 5, 6], Pankaj Jay Pasricha[1]

[1]Center for Neurogastroenterology, Department of Medicine, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[2]Department of Genetic Medicine, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[3]Division of Gastroenterology, Stanford University – School of Medicine, Stanford CA 94301

[4]Center for Cancer Research, National Cancer Institute, Frederick, MD 21702

[5]Department of Neuroscience, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[6]Kavli Neurodiscovery Institute, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[#]Address all correspondence to:

Subhash Kulkarni, Ph.D.

Assistant Professor

Department of Medicine

skulkar9@jh.edu

**Abstract**

The enteric nervous system (ENS), a collection of neurons contained in the wall of the gut, is of fundamental importance to gastrointestinal and systemic health. According to the prevailing paradigm, the ENS arises from progenitor cells migrating from the neural crest and remains largely unchanged thereafter. Here, we show that the lineage composition of maturing ENS changes with time, with a decline in the canonical lineage of neural-crest derived neurons and their replacement by a newly identified lineage of mesoderm-derived neurons. Single cell transcriptomics and immunochemical approaches establish a distinct expression profile of mesoderm-derived neurons. The dynamic balance between the proportions of neurons from these two different lineages in the post-natal gut is dependent on the availability of their respective trophic signals, GDNF-RET and HGF-MET. With increasing age, the mesoderm-derived neurons become the dominant form of neurons in the ENS, a change associated with significant functional effects on intestinal motility which can be reversed by GDNF supplementation. Transcriptomic analyses of human gut tissues show reduced GDNF-RET signaling in patients with intestinal dysmotility which is associated with reduction in neural crest-derived neuronal markers and concomitant increase in transcriptional patterns specific to mesoderm-derived neurons. Normal intestinal function in the adult gastrointestinal tract therefore appears to require an optimal balance between these two distinct lineages within the ENS.

**Introduction**

The enteric nervous system (ENS) is a large collection of neurons, glial, and precursor cells that resides within the gastrointestinal wall and regulates gut motility and secretion along with modulating epithelial and immune cell function[1,2]. During fetal development, the mammalian ENS is populated by neurons and glia derived from neural crest (NC)-derived precursors[3-9]. These precursors follow diverse migratory routes to colonize and innervate various parts of the gut before birth[10-12]. It is not clear, however, that this lineage persists in its entirety in the adult gut, as indicated by the observed lack of expression of fluorescent reporter protein in a subpopulation of adult enteric neurons in NC-lineage-traced mice[13,14]. Alternative sources of enteric neurons that have been proposed in the literature include the ventral neural tube (VENT)[15], or the Pdx1-expressing pancreatic endoderm[14], but the interpretation of these studies has been limited by the lack of robust lineage markers for non-NC derived neurons[16]. In addition, while prior studies have documented cellular changes to the ageing ENS[17], the developmental mechanisms behind these changes are unknown. Thus, confirmation of a second, distinct lineage of enteric neurons in adults is important for our understanding of the healthy post-natal development and aging of the ENS, as well as for the pathogenesis of acquired disorders of the ENS.

In this study, we found that while the early post-natal ENS is derived from the canonical NC-lineage, this pattern changes rapidly as the ENS matures, due to the arrival and continual expansion of a novel population of <u>M</u>esoderm-derived <u>E</u>nteric <u>N</u>euron<u>s</u> (MENs) which represent an equal proportion of the ENS in young adulthood and with increasing age, eventually outnumber the <u>N</u>C-derived <u>E</u>nteric <u>N</u>euron<u>s</u> (NENs). We also found that, while the NEN population is regulated by glial derived neurotrophic factor (GDNF) signaling through its receptor RET, the MEN population is regulated by hepatocyte growth factor (HGF) signaling. Increasing HGF levels during maturation or by pharmacological dosing increase proportions of MENs. Similarly,

decrease in GDNF with age decrease NENs; and increasing GDNF levels by pharmacological

dosing increase NENs proportions in the adult ENS to impact intestinal motility.

These results indicate for the first time that the mesoderm is an important source of neurons in

the second largest nervous system of the body. The increasing proportion of neurons of meso-

dermal lineage is a natural consequence of maturation and aging; further, this lineage can be

expected to have vulnerabilities to disease that are distinct from those affecting the NEN popu-

lation. These findings therefore provide a new paradigm for understanding the structure and

function of the adult and aging ENS in health, age-related gut dysfunction and other acquired

disorders of gastrointestinal motility.

**Results**

**Only half of all mid-age adult enteric neurons are derived from the neural crest.** We ana-

lyzed small intestinal longitudinal muscle-myenteric plexus (LM-MP) from adult (post-natal day

60; P60) Wnt1-cre:*Rosa26*-tdTomato mice, in which tdTomato is expressed by all derivatives of

Wnt1$^+$ NC-cells[18]. In these tissues, while GFAP, a glial marker, was always co-expressed with

tdTomato (Fig 1a), tdTomato-expression was absent in many myenteric neurons (Fig 1b, c). By

careful enumeration, almost half of all myenteric neurons expressing the pan-neuronal marker

HuC/D were found to not express the reporter (Fig 1d, percent tdTomato$^+$ neurons: 55.73 ±

2.404 SEM; percent tdTomato$^-$ neurons: 44.27 ± 2.404 SEM; enumerated from 2216 neurons

from 6 mice). In these lineage-traced mice, myenteric ganglia were found to contain tdToma-

to$^{high}$ and tdTomato$^{low}$ neurons and due care was taken to image subsets (Suppl. Fig 1a). Both

tdTomato$^{high}$ and tdTomato$^{low}$ neurons were classified as tdTomato$^+$ and only neurons that did

not show any tdTomato-expression were classified as tdTomato$^-$ neurons. Immunofluorescence

staining of small intestinal LM-MP tissue from adult Wnt1-cre:*Rosa26*-tdTomato mice using an-

tibodies against tdTomato protein showed a consistent lack of reporter expression in a popula-

tion of myenteric neurons (Suppl. Fig 1b). This lack of reporter expression is not a function of

tissue peeling or fixation, as the tdTomato⁻ cells within the myenteric ganglia were also ob-

served in freshly harvested unpeeled gut tissue, when imaged using a live tissue fluorescence

microscope (Suppl. Fig 1c). In addition, we noticed that myenteric ganglia of the Wnt1-

cre:*Rosa26*-tdTomato transgenic mouse show similar tdTomato aggregation in cells of the

myenteric plexus from both freshly harvested and immediately fixed tissue (Suppl. Fig 1d, e).

Since we previously showed ongoing neuronal loss in healthy ENS[19], we reasoned that cells

that show hyper-aggregation of tdTomato are in advanced stages of cell death. Cells with hyper-

aggregation of tdTomato showed a lack of staining for the nuclear marker DAPI (Suppl. Fig 1e),

suggesting that these are indeed cells in the advanced stages of death, and given that they are

not labeled with antibodies against HuC/D, their presence does not alter our estimation of

tdTomato-expressing neurons (Suppl. Fig 1f). Thus, the absence of reporter expression in

myenteric ganglia of freshly harvested tissue and the presence of tdTomato aggregation in both

freshly harvested and fixed tissues shows that our observations are not caused due to any

technical issues in tissue isolation and preservation.

Additional confirmation was provided by an analogous Wnt1-cre:*Hprt*-tdTomato lineage-traced

mouse line, in which tdTomato was expressed from the *Hprt* locus in a Wnt1-cre-dependent

manner (*Hprt* locus is X-inactivated in females, hence only adult male mice were used)[20]; and

by the Pax3-cre:*Rosa26*-tdTomato lineage-traced mouse line, where *Rosa26*-tdTomato driven

by the Pax3-cre transgene labels the derivatives of the neural tube and pre-migratory NC

(Suppl. Fig 1g, h)[21]. Similar lack of reporter expression was previously observed in adult myen-

teric neurons from Sox10-cre NC-lineage-traced mice, and more recently in significant numbers

of myenteric neurons from Wnt1-cre:*Rosa26*-H2B-mCherry mice, which expresses nuclear lo-

calized reporter mCherry in NC-lineage cells, further confirming our observations of the pres-

ence of non-NC-derived neurons in the adult ENS[13,22].

**Lineage-tracing confirms a mesodermal derivation for half of all adult myenteric neurons.**

Alternative sources of enteric neurons proposed previously include the ventral neural tube[15], and the pancreatic endoderm[14], but the interpretation of these studies was limited by the lack of robust lineage markers[16]. Brokhman et al.[14] found evidence of labeled neurons in inducible Pdx1-cre, Foxa2-cre, and Sox17-cre lineage-traced mouse lines and inferred their derivation from pancreatic endoderm. However, in a Pdx1-cre lineage-traced mouse line, many neu-roectoderm-derived neurons of the central nervous system have also been previously shown to be derived from Pdx1-expressing cells[23], suggesting that Pdx1 is not an exclusive endodermal marker. Foxa2 is expressed by mesoderm-derived cells in the heart[24] and Sox17 also labels mesoderm-derived cells, especially cells of the intestinal vasculature[25-28]. We therefore hypothe-sized that the mesoderm may be the true developmental source of the non-NC enteric neurons.

Mesoderm-derived cells during embryogenesis express Tek[24] and analysis of LM-MP tissues from adult male Tek-cre:*Hprt*-tdTomato lineage-traced mice revealed the presence of a popula-tion of tdTomato$^+$ neurons (Fig 2a). Since Tek is not expressed in the adult ENS (Suppl. Fig 2a), the presence of Tek-derived adult myenteric neurons suggested their mesodermal origin. We then used *mesoderm posterior 1* or Mesp1 as a definitive developmental marker for the meso-derm[29-33] to confirm the mesodermal derivation of the non-NC derived adult enteric neurons. Reporter expression in this mouse line has been previously studied in cardiac development, which we confirmed by observing its expression in the adult heart tissue (Fig 2b). Expression of tdTomato in vascular cells of both Tek-cre and Mesp1-cre mice was more pronounced than in myenteric neurons (Fig 2a, c) and in cardiac myocytes of Mesp1-cre mice (Fig 2b, Suppl. Fig 2b), which reflects the variable expression of the reporter gene in both the neural crest-lineage and mesoderm-lineage specific transgenic mouse lines (Fig 2b, c; Suppl. Figs 1a, 2c). Using an analogous Mesp1-cre:*Rosa26*-EGFP lineage-traced mouse line, in which EGFP was expressed from the *Rosa26* locus in a Mesp1-cre-dependent manner, we next confirmed the derivation of a

population of myenteric neurons from Mesp1-expressing mesoderm (Fig 2e, Suppl. Fig 2d, e).

Similar to our observations with the tdTomato reporter (Fig 2a-d), we observed lower expression

of the reporter EGFP in myenteric neurons as compared to vascular cells (Fig 2e; Suppl. Fig 2d,

e). Variable expression of CAG and other pan-cellular promoters in various cell-types and dur-

ing processes of maturation have previously been reported[34-37]. In addition, an earlier study by

Agah et al showed differing degrees of CAG-driven LacZ reporter activation in a cardiac-specific

transgenic cre mouse line, which were unrelated to copy number, suggesting insertional and

positional effects or, potentially, differential methylation[38]. These reports are consistent with our

observation and could potentially help explain the observed variable expression of CAG and

CMV promoter-driven reporter genes in our study.

Analysis of small intestinal LM-MP from P60 Mesp1-cre:*Rosa26*-tdTomato lineage-traced mice

showed that tdTomato-expression was observed in about half of all myenteric neurons (Fig 2f;

percent tdTomato$^+$ neurons: 50.28 ± 2.89 SEM; percent tdTomato$^-$ neurons: 49.72 ± 2.89 SEM;

enumerated from 484 neurons from 3 mice). Because of the variable expression of the reporter

gene in the transgenic mice described above, we also labeled Mesp1-cre:*Rosa26*-tdTomato$^+$

neurons with the S46 antibody, which was raised against the slow tonic myosin heavy chain

protein MHCst derived from avian embryonic upper leg muscle, and is expressed exclusively by

mesoderm-derived cell populations (Fig 2g)[39-42]. MHCst immunostaining was exclusively ob-

served in all Tek-cre:*Rosa26*-tdTomato$^+$ MENs (Suppl. Fig 2f). None of the 107 MHCst$^+$ intra-

ganglionic cells observed (35.67 ± 6.17 SEM MHCst$^+$ cells per mouse) across 3 Wnt1-

cre:*Rosa26*-tdTomato mice expressed tdTomato, suggesting that MHCst immunostaining was

exclusive to non-NC lineage neurons (Fig 2h, Suppl. Fig 3a). Thus, both lineage tracing and

protein biomarker expression provides strong support for their mesodermal origin (Fig 2g,

Suppl. Fig 3a). Along with our observations on Wnt1-cre:*Rosa26*-tdTomato mice (Fig 1c), our

results indicate that these two distinct lineages together account for all the adult small intestinal

myenteric neurons.

In a prior report, Avetisyan et al. showed that the proteins RET, a receptor tyrosine kinase that

transduces GDNF signaling in NC-derived enteric neuronal precursors, and MET, a receptor for

hepatocyte growth factor (HGF), are expressed by different subsets of adult myenteric

neurons[43]. MET is classically expressed by mesoderm-derived cells[44], and by using im-

munostaining of small intestinal LM-MP tissues from Wnt1-cre:*Rosa26*-tdTomato and Mesp1-

cre:*Rosa26*-tdTomato mice, we found that the expression of MET was restricted to a sub-

population of adult MENs (Fig 3a, b). By contrast, RET expression was confined to NENs (Fig

3c).

We then studied whether MENs and NENs differed phenotypically. In the adult murine ENS, in-

hibitory enteric motor neurons express the nitric oxide-producing enzyme nitric oxide synthase 1

(NOS1); and enteric sensory neurons, called intrinsic primary afferent neurons (IPANs) express

the neuropeptide calcitonin gene related peptide (CGRP)[9,43]. Immunostaining using antibodies

against NOS1 (expressed by nitrergic neurons) and CGRP (sensory neuronal marker) expres-

sion in the LM-MP from P60 Wnt1-cre:*Rosa26*-tdTomato mice showed that both tdTomato$^+$

NENs and tdTomato$^-$ MENs express these two neuronal markers (Fig 3d, e). However, quantifi-

cation of proportions in these two neuronal lineages shows that while the majority (~73%) of

NOS1$^+$ inhibitory neurons are NENs (Fig 3f; of the 616 NOS1$^+$ neurons observed across 3 mice,

NOS1$^+$ tdTomato$^+$: 72.94% ± 4.46 SEM; NOS1$^+$ tdTomato$^-$: 27.06% ± 4.46 SEM; p = 0.0019;

Student"s t-test), the majority (~75%) of CGRP$^+$ neurons are MENs (Fig 3g; of the 146 CGRP$^+$

neurons observed across 3 mice, CGRP$^+$ tdTomato$^-$: 75.76% ± 0.63 SEM; CGRP$^+$ tdTomato$^+$:

24.24% ± 0.63 SEM; p < 0.0001; Student"s t-test). These results are in keeping with a previous

report that show low expression of NOS1 and abundant expression of CGRP by MET[+]

neurons[43], previously not known to be derived from a different lineage. In addition, MENs also

express Cadherin-3 (CDH3, Fig 3h, Suppl. Fig 3b), which is known to also mark a sub-

population of mechanosensory spinal neurons[45]. Apart from their derivation from a distinct de-

velopmental lineage, the mean cell size of MENs was significantly larger than that of NENs

(Suppl. Fig 4a; n = 143 neurons/group observed across 3 mice; Feret diameter (µm) MENs:

17.47 ± 0.50 SEM, NENs: 13.03 ± 0.36 SEM, p = 0.002; Student"s t-test).

**An expanded molecular characterization of MENs using unbiased single cell RNA se-**

**quencing (scRNAseq)-based analyses.** The distinct nature of the MENs and NENs was fur-

ther ascertained by scRNAseq-based analyses on all cells isolated from ileal LM-MP. Unlike

other studies that analyzed either FACS-sorted Wnt1-cre:*Rosa26*-tdTomato[+] ENS-cells or used

classical NENs markers such as Snap25 to drive identification and/or enrichment of enteric neu-

rons[46,47], we performed agnostic clustering of the scRNAseq data from tissues of two 6-month-

old adult male C57BL/6 wildtype mice (Fig 4a; Suppl. Fig 4b, c). We identified clusters of NC-

derived cells by exclusive expression of canonical NC markers Ret and Sox10 (which include

the clusters of Ret-expressing NENs and Sox10-expressing Neuroglia); and the MENs cluster

by its co-expression of the genes Calcb (CGRP), Met, and Cdh3 (Fig 4a, b). With cells from

both samples pooled together, we compared 1,737 NC-derived cells across the NENs (77 cells)

and Neuroglia (1660 cells) clusters with 2,223 cells in the MENs cluster. A list of top differential-

ly expressed genes for every cluster are provided in Suppl. Table 2.

The neuronal nature of MENs was first established by testing our scRNAseq dataset for expres-

sion of genes known to be expressed by enteric or other neurons. We found detectable expres-

sion of genes Elavl2 (HuB), Hand2, Pde10a, Vsnl1, Tubb2b, Stmn2, Stx3, and Gpr88 in the

MENs as well as in the NC-derived cell clusters (Fig 4b). While expression of Elavl2, Stmn2 and

Hand2 has been previously observed in enteric neurons[48,49], the expression of neuronal marker genes Pde10a (expressed by medium spiny neurons in the striatum[50]), Vsnl1 (expressed by hippocampal neurons[51]), Gpr88 (expressed by striatal neurons[52]), Stx3 (expressed by hippocampal neurons[53]), and Tubb2b (expressed by adult retinal neurons[54]) has not been studied. Using antibodies against these markers along with co-staining with ANNA1 antisera to mark all HuC/D[+] neurons, we visualized the expression of these neuronal markers in both tdTomato[+] NENs and tdTomato[-] MENs of the small intestinal myenteric plexus tissue from adult Wnt1-cre:*Rosa26*-tdTomato mice (Fig 4c, Suppl. Fig 3l-q). In addition, the MENs cluster was also enriched in previously characterized ENS markers, such as Ntf3 and Il18[2,55] (Fig 4b). We validated the MEN-specificity of these markers using immunochemical analyses (Fig 4c; Suppl. Fig 3b - k).

Examination of the MENs cluster yielded additional MENs-specific marker genes Slpi, Aebp1, Clic3, Fmo2, Smo, Myl7, and Slc17a9, whose expression by adult enteric neurons has not been previously described and which we also validated (Fig 4b, c; Suppl. Fig 3b - k). Next, we found that the mean UMI per cluster in MENs (n = 2223 cells, 4023.91 ± 85.35 SEM) was significantly greater than in Neuroglia (n = 1660 cells, 1137.93 ± 27.93 SEM, p < 0.0001) and while it was greater than in NENs (n = 77 cells, 3391.71 ± 386.52 SEM), this difference did not meet statistical threshold (p = 0.11, One way ANOVA).

These data help show the neuronal nature of MENs while distinguishing them from NENs at the molecular level. In addition, cells within the Neuroglia cluster showed expression of glial genes such as Sox10 as well as neuronal genes such as Ncam1, Hand2 and Stmn2, suggesting that NC-derived neurons may be present within both the NENs cluster as well as the Neuroglia cluster (Fig 4b).

**Detection of MENs in other murine and human datasets.** Recently, by performing single nu-

clei RNA sequencing (snRNAseq) on flow sorted Phox2b-expressing nuclei from the adult mu-

rine LM-MP tissue, May-Zhang et al[56] detected the presence of a cluster of neurons in the adult

murine gut that expressed mesenchymal markers. We tested whether the MENs-expressed

genes that we validated earlier (Fig 4c, Suppl. Fig 3) were also expressed by the „mesenchymal"

neurons in the May-Zhang et al dataset. We found a high degree of similarity between the top

genes expressed by the MENs cluster in our scRNAseq dataset and the „mesenchymal" neu-

rons in the May-Zhang et al dataset (Table 1, Suppl. Fig 4d). Furthermore, examination of the

top genes shows that this neuronal sub-population is significantly different from the other Calcb-

expressing neuronal populations, which we presume to be the CGRP-expressing NENs (Table

1).

To confirm that the „mesenchymal" neurons identified by May-Zhang et al are MENs, we used

an agnostic bioinformatics projection-based approach which we have previously published[57].

This approach allows us to learn latent space representations of gene expression patterns using

single cell transcriptomic data, which include patterns that correspond to cell-type-specific gene

expression signatures[57]. We can then project other single cell and bulk RNA transcriptomic da-

tasets into these learned patterns to accurately quantify the differential use of these transcrip-

tional signatures in target data across platforms, tissues, and species[57]. In this instance, gene

expression patterns were learned on our murine single cell transcriptomic data using non-

negative matrix factorization (NMF) dimensionality reduction and the data was decomposed into

50 distinct NMF-patterns[57,58] (Fig 5a). Four NMF patterns were identified that were specific to

MENs (Fig 5a, b; Suppl. Fig 4e). We next projected the snRNAseq dataset from May-Zhang et

al[56] into the four MEN-specific patterns and found that all these patterns specifically labeled the

cluster of „mesenchymal" neurons in the May-Zhang et al dataset of the adult ileal ENS (Fig 5b).

Further, upon clustering the projection weights from the four MEN-specific patterns, we identi-

fied a cluster of cells that showed higher usage of all four patterns, when compared to all the

other cells in the dataset (Suppl. Fig 4f). The genes that define this cluster of cells are similar to

the top-expressed genes that define our MENs cluster (Suppl. Fig 4g, Table 1). This approach,

together with the high similarities in the top-expressed genes, establishes that cluster of neu-

rons annotated as „mesenchymal" neurons by May-Zhang et al are indeed MENs.

In addition, our results are supported by a recent report by Elmentaite et al.[59] who performed

scRNAseq on dissociated single cells from the human gut at fetal, juvenile, and adult ages and

catalogued multiple intestinal cell-types to generate a comprehensive gut cell atlas. Canonical

enteric neurons, while adequately represented in the fetal ages, were not found in the juvenile

and adult ages which were dominated by mesenchymal and other cell types[59]. Given that our

scRNAseq approach of sequencing diverse cell types from the murine gut wall without any

marker-based pre-selection was similar to this approach in human tissue, we reasoned that

MENs would also be represented in Elmentaite et al"s description of mesenchymal cell clusters,

especially in the juvenile and adult ages. To detect putative MENs in this human gut cell atlas

data, we projected the entire human mesenchymal cell scRNAseq dataset into the four murine

MEN-specific NMF patterns and detected putative human MENs clusters (Suppl. Fig 4h). Since

our identification of MENs and the generation of MEN-specific NMF patterns was based on

post-natal healthy murine tissue, we next projected data from only post-natal healthy mesen-

chymal clusters into the four MEN-specific NMF patterns (Fig 5b). The four patterns identified a

population of cells within the post-natal intestinal mesenchymal cell clusters suggesting that

these cells are putative MENs in the human gut (Fig 5b). These putative MENs, which map to

Cluster 1 in our representation of the intestinal mesenchymal cells from the post-natal gut cell

atlas data (Fig 6a) were annotated previously by Elmentaite et al. as transitional stromal cells

(Fig 6b). We next tested whether these clusters expressed neuronal markers expressed by en-

teric neurons and found that the pan-neuronal marker Uchl1 as well as neuronal markers

Hand2, Stx3, Slc17a9, Pde10a and Tubb2b that we earlier showed to be expressed by MENs

(Fig 6c) were expressed by the putative MENs in the adult human gut.

We next hypothesized that the reason these cells were not annotated as enteric neurons was

due to the use of a NENs-restricted gene list. Apart from the NENs marker Ret, such gene lists

often contain the pre-synaptic gene Snap25, a component of the SNARE complex, which is

widely assumed to be a pan-neuronal marker in the adult ENS[46,47,60,61]. While we have previous-

ly observed the NENs-specific nature of RET, transcriptomic data suggests that MENs either do

not express Snap25 or do so in very small amounts (Suppl. Fig 4d). We tested the expression of

SNAP-25 by NENs and MENs in the myenteric ganglia of an adult Wnt1-cre:*Rosa26*-tdTomato

mouse and found that SNAP-25 expressing neurons expressed tdTomato, but not the MENs-

marker MHCst, suggesting the NEN-specific expression of this canonical marker for synaptic

neurons (Fig 7a). The low or lack of expression of Ret, Snap-25, Elavl3 and Elavl4 in MENs as

observed both by us and May-Zhang et al[56] (Suppl. Fig 4d) may help explain why these canoni-

cal gene lists missed the correct identification and annotation of MENs" neuronal nature in hu-

man datasets[47,59]. Next, we examined the expression of MENs markers MHCst, MET, and

SLC17A9 in LM-MP tissues from adult humans with no known gut motility disorder and found

them to be expressed by a population of myenteric neurons in normal adult human ENS (Fig 7b,

c; Suppl. Fig 5a, b). Thus, our bioinformatic approaches and immunofluorescence analyses of

MEN-specific markers in adult human gut together shows that the human gut similarly contains

a population of MENs.

We next tried sub-clustering the MENs cell cluster in our scRNAseq data to study whether indi-

vidual subclusters could be discriminated on the basis of MENs markers or other genes. How-

ever, these clusters did not yield meaningful information on specific marker genes that could be

used to define them (Suppl. Fig 6a, b).

**The proportion of mesoderm-derived neurons expands with age to become the dominant population in the aging ENS**. Since the ENS at birth consists solely of NENs[13], we next studied the birth-date and eventual expansion of the MEN lineage in the post-natal gut. Using Wnt1-cre:*Rosa26*-tdTomato mouse line, we enumerated the tdTomato⁻ MENs in LM-MP at different ages and found a significant age-associated increase in MENs proportions (Fig 8a, b, c; One-way ANOVA, F = 117.6, DFn, DFd = 5, 12; p < 0.0001). At P11, MENs were found only in few isolated myenteric ganglia (Fig 8a) and together represented only ~5% of all myenteric neurons (tdTomato⁻ neurons: 4.12% ± 2.98 SEM; n = 1327 HuC/D⁺ neurons counted from 3 mice), suggesting that they originate shortly before this age. The proportion of MENs in myenteric ganglia rises sharply thereafter: by P22, they account for ~30% of all neurons (tdTomato⁻ neurons: 29.63% ± 1.229 SEM; n = 742 neurons counted from 3 mice); and at P60 they represent ~45% of all neurons (tdTomato⁻ neurons: 46.38% ± 4.62 SEM; n = 700 neurons counted from 3 mice). During the adult ages, the proportions of MENs in the adult ENS remains relatively stable as at P180, MENs continue to represent roughly half of all myenteric neurons (tdTomato⁻ neurons: 57.29% ± 3.62 SEM; n = 586 neurons counted from 3 mice). However, MENs dominate the proportions of the aging ENS as at the very old age of 17 months (P510), the ENS is populated almost exclusively by MENs (Fig 8b; tdTomato⁻ neurons: 95.99% ± 1.62 SEM; n = 996 neurons counted from 3 mice). Thus, the proportions of NENs and MENs in the myenteric ganglia can be used as a biomarker for deducing ENS age, as a healthy ENS dominant in NENs would be juvenile, one with roughly equal proportions would be adult, and an aging ENS is dominated by MENs.

We next tested whether the representation of MEN-specific transcriptomic signatures show a similar increase during the maturation and aging of the human gut. We tested whether the representation of the four MEN-specific NMF patterns (Fig 5a) increased with age in the human

transcriptomic data from Elmentaite et al[59]. We grouped the transcriptomic data from the wide-

range of healthy post-natal specimens of the gut cell atlas into three groups: Juvenile (contain-

ing data on 2910 cells represented by 10 samples within the age range of 4 – 12 years), Adult

(containing data on 3696 cells represented by 36 samples within the age range of 20 – 50

years), and Aging (containing data on 2848 cells represented by 37 samples within age range of

60 – 75 years) and tested whether the representation of the four MEN-specific patterns changed

significantly between these age groups. Barring pattern 16 which showed a non-significant in-

crease in projection weights between the Juvenile and Aging samples (One-way ANOVA, F =

2.10, DFn, DFd = 2, 8, p = 0.18), projection weights of MEN-specific pattern 27 (One-way ANO-

VA, F = 16.08, DFn, DFd = 2, 8, p = 0.0016), pattern 32 (One-way ANOVA, F = 8.13, DFn, DFd

= 2, 8, p = 0.01) and pattern 41 (One-way ANOVA, F = 9.818, DFn, DFd = 2, 8, p = 0.007) all

show significant age-associated increase in the human gut tissue (Fig 8d, every datapoint refers

to mean projected pattern weight for cells within a defined age or age range). This suggests that

analogous to the age-associated increase in MENs we observed in murine gut (Fig 8b), the pro-

portions of MENs in the human gut also might increase with age. We also found that the repre-

sentation of MENs cluster (Cluster 1) in the post-natal human gut cell atlas data expanded with

maturation and age (Suppl. Fig 5c)

**GDNF and HGF levels regulate the populations of the neural crest-derived and the meso-**

**derm-derived neurons, respectively, in the maturing ENS.** GDNF-RET signaling is respon-

sible for proliferation and neurogenesis from NC-derived ENS precursor cells during develop-

ment as well as for the survival of Ret-expressing enteric neurons[62-65]. Similarly, HGF signaling

has been shown to be essential for the proliferation of mesoderm-derived cells[66]. Since the ex-

pression of the HGF receptor MET and the GDNF receptor RET is exclusive to MENs and

NENs respectively, we studied the correlation between age and levels of HGF and GDNF in LM-

MP (Fig 8e, f). Given that both GDNF, as well as HGF expression is found in the LM-MP

layer[43,67,68], we used LM-MP tissue to study how the levels of these two trophic factors change

in and around the ENS tissue during maturation. We found that in agreement with a previous

report[66], GDNF protein levels are markedly reduced between the P10 and the P30 ages and

remains reduced thereafter up to the P90 age (Fig 8g; n = 3 mice/age-group, F = 6.821, DFn,

DFd = 1, 7, p = 0.0348, One way ANOVA). In addition, expression of Gdnf transcripts show sig-

nificant reduction between the ages of P30 and P90 (Fig 8h; n = 6 mice/age-group, P30: 1.070

± 0.179 SEM, P90: 0.539 ± 0.129 SEM, p = 0.037, Student"s t-test). On the other hand, HGF

expression increases progressively between P10 and P90 ages (Fig 8i, j: n = 3 mice/age-group,

Protein levels: F = 8.820, DFn, DFd = 1, 7, p = 0.02; mRNA levels: F = 36.98, DFn, DFd = 1, 16,

p < 0.0001, One way ANOVA). The ratio of HGF to GDNF expression in ileal LM-MP tissue

shows significant increase from the age of P10 to P90 in our data (Suppl. Fig 7a, F = 48.60,

DFn, DFd = 1, 7, p = 0.0002, One way ANOVA), and HGF expression is consistently more than

GDNF expression in the full thickness small intestinal tissue from adult and aging mice in the

Tabula muris data[69] (Suppl. Fig 7b). We then queried the plasma proteome levels from the

LonGenity cohort of 1,025 aging individuals and found that HGF levels correlated positively,

while GDNF and RET levels correlated negatively with age (Suppl. Fig 7c)[70], suggesting paral-

lels between our data from murine intestine and human plasma proteome data.

Since GDNF tissue levels are correlated with NENs proportions, we hypothesized that GDNF

signaling regulates NENs proportions in maturing and adult ENS. On administration of GDNF or

saline to cohorts of P10 Wnt1-cre:*Rosa26*-tdTomato mice over 10 days[62,71], we found that

GDNF treatment promoted the juvenile phenotype by enhancing the proportions of tdTomato[+]

NENs (Fig 8k). GDNF supplementation reduced the proportion of tdTomato[-] MENs in P20 mice

to a level similar to that seen at the P10 age, while the MENs proportions in saline-treated con-

trol mice remained at a level expected of its age (Fig 8k; Controls: %MENs: 25.87 ± 4.37 SEM

of 1350 neurons from 3 mice; GDNF: %MENs: 3.86 ± 0.07 SEM of 1301 neurons from 3 mice; p

= 0.0072; Student"s t-test). Consistent with a previous report[62], the GDNF-driven expansion of

NENs and associated contraction of MENs conserved the total neuronal numbers (Fig 8k; Con-

trols: neurons/ganglia = 20.60 ± 4.00 SEM; GDNF: neurons/ganglia = 24.39 ± 6.96 SEM; p =

0.52; Student"s t-test).

Since HGF tissue levels increase with age, we hypothesized that increasing HGF signaling

drives the expansion of MENs in the maturing gut. HGF administration to cohorts of P10 Wnt1-

cre:*Rosa26*-tdTomato mice over 10 days promoted an increase in the tdTomato$^-$ MENs popula-

tion in P20 mice to levels previously observed in P60 mice, while tissues from saline-treated

control mice exhibited a MENs:NENs ratio that is expected at P20 (Fig 8l; Controls: %MENs:

27.40 ± 5.49 SEM of 1970 neurons from 5 mice; HGF: %MENs: 49.37 ± 5.52 SEM of 1704 neu-

rons from 5 mice; p = 0.02, Student"s t-test).  Similar to GDNF treatment, HGF treatment also

did not cause any significant change in total neuronal numbers (Fig 8l; Controls: neu-

rons/ganglia = 20.35 ± 2.22 SEM; HGF: neurons/ganglia = 18.21 ± 0.69 SEM; p = 0.38, Stu-

dent"s t-test).

**Reduced GDNF-RET signaling accelerates ENS aging to cause intestinal dysfunction.**

Since reduced GDNF or RET levels are associated with intestinal dysfunction in patients[61,72,73],

we hypothesized that alterations in GDNF-RET signaling unrelated to those seen with normal

aging, would cause dysfunction. To test this hypothesis, we studied lineage proportions and in-

testinal function in a mouse model of reduced RET signaling. Ret-null heterozygous mice, which

have been previously used to study the effect of reduced RET signaling in the adult ENS, have

normal ENS structure but altered gut physiology[62]. A similar mouse model with a Ret$^{CFP}$ allele

has a CFP reporter inserted at its *Ret* locus rendering it null[74]. Ret$^{+/CFP}$ (or Ret$^{+/-}$) mice carrying

a single copy of the Ret gene showed significant reduction in Ret transcript and RET protein

expression in the murine gut[74]. In these mice, using antibodies against CFP/GFP, HuC/D, and

MHCst, we confirmed that the NENs marker Ret-CFP, and the MENs marker MHCst were ex-

pressed by different neuronal subpopulations (Fig 9a). Using the adult Ret$^{+/-}$ mice, we tested

the effect of partial Ret loss on ENS lineages at two adult ages: 9 weeks (~P60) and 16 weeks

(~P110). Using antibody against CFP/GFP to detect CFP$^+$ RET-expressing neurons, we found

that Ret$^{+/-}$ mice show a significant reduction in the proportions of Ret-CFP$^+$ NENs (Fig 9b; at 9

weeks: %CFP$^+$ neurons: 24.91 ± 5.42 SEM of 837 neurons from 3 mice; at 16 weeks: %CFP$^+$

neurons: 13.13 ± 0.98 SEM of 1227 neurons from 5 mice; p = 0.03, Student"s t-test). We ob-

served a corresponding significant increase in the proportions of MHCst$^+$ MENs with age in

Ret$^{+/-}$ mice (Fig 9c; at 9 weeks: %MENs: 58.74 ± 7.33 SEM of 644 neurons from 3 mice; at 16

weeks: %MENs: 82.84 ± 3.58 SEM of 935 neurons from 5 mice, One-way ANOVA, p = 0.014),

while control Ret$^{+/+}$ mice showed no significant age-associated change in the proportions of

MENs (Fig 9c; 9 weeks: %MENs: 43.27 ± 3.24 SEM of 780 neurons from 3 mice; 16 weeks:

%MENs: 54.48 ± 4.07 SEM of 1276 neurons from 5 mice, One-way ANOVA, p = 0.36), which is

consistent with our previous results that MENs proportions in wildtype animals are relatively

stable in this time window (Fig 8c). The expedited loss of NENs in Ret$^{+/-}$ mice confirms that de-

pletion of endogenous RET signaling in the adult ENS accelerates the aging-associated loss of

NENs.

Having previously shown that aging mice have intestinal dysmotility[75], we tested whether the

increased loss of NENs in the Ret$^{+/-}$ ENS, concomitant with the expansion of MENs accelerated

ENS aging, causes an early onset of aging-associated intestinal dysmotility. We studied whole

gut transit time (WGTT) in a cohort (n = 10) of adult Ret$^{+/-}$ mice and their littermate control (n =

8) Ret$^{+/+}$ mice over 7 weeks, between 9 and 16 weeks of age. While 9-week adult Ret$^{+/-}$ mice

were similar to control Ret$^{+/+}$ mice, WGTT between the two genotypes diverged with age. Con-

sistent with a prior report[62], 16-week old Ret[+/-] mice displayed significantly delayed intestinal

transit compared to age-matched control Ret[+/+] mice (Fig 9d; WGTT (in min) Ret[+/+]: 121.4 ±

4.01 SEM; Ret[+/-]: 157.3 ± 14.62 SEM, p = 0.048; One way ANOVA).

**GDNF reverts aging in the ENS to normalize intestinal motility.** Along with others, we have

previously shown that aging is associated with slowing of intestinal motility[76,77]. We hypothe-

sized that this may be a consequence of the replacement of the juvenile NENs population by

MENs and therefore GDNF supplementation, by restoring a more balanced MENs:NENs ratio,

may prevent age related changes in motility. We studied 17-month-old male mice (at an age

where NENs constitute only ~5% of all myenteric neurons; Fig 8c) before and after they re-

ceived 10 days of intraperitoneal injection of GDNF or saline (n=5 for each cohort). While the

two cohorts showed no significant difference in their intestinal transit times at baseline (WGTT

(in min) Control: 192.8 ± 11.55 SEM; GDNF: 202.4 ± 7.60 SEM, p = 0.50, Student"s t-test),

GDNF treatment caused significant improvement in intestinal transit (Fig 10a, WGTT (in min)

Control: 175.0 ± 8.89 SEM; GDNF: 101.0 ± 8.91 SEM, p = 0.0004, Student"s t-test), reaching

levels previously observed in healthy mice (Fig 9d).

GDNF treatment significantly reduced the proportions of MHCst[+] MENs (Fig 10b; Suppl. Fig 8a,

b; Control: %MENs: 88.76 ± 1.48 SEM of 909 neurons from 5 mice; GDNF: %MENs: 74.12 ±

5.98 SEM of 799 neurons from 5 mice; p = 0.045; Student"s t-test) while increasing the numbers

of RET[+] NENs (Fig 10c; Suppl. Fig 8c, d; Control: RET[+] neurons: 1.35 ± 0.05 SEM in forty-two

40X fields from 4 mice; GDNF: RET[+] neurons: 3.19 ± 0.56 SEM in forty-one 40X fields from 4

mice, p = 0.017; Student"s t-test). Consistent with our earlier observation, GDNF treatment did

not change the average neuronal numbers in the ileal myenteric ganglia (Control: 17.56 ± 1.82

SEM neurons per ganglia, GDNF: 16.01 ± 1.22 SEM neurons per ganglia, p = 0.49, Student"s t-

test).

**The two neuronal lineages in human disease.** In addition to the detection of MENs in the

adult healthy gut by bioinformatic and immunofluorescence means (Fig 5, 6, 7), we next exam-

ined the relevance of the two neuronal lineages NENs and MENs in human disease. For this,

we tested whether gut dysfunction associated with pathological reductions in NENs-signaling

mechanisms are also associated with increased abundance of MENs. We mined previously

generated and publicly available transcriptomic data from intestinal tissues of humans with nor-

mal gut motility (controls) and of patients with obstructed defecation (OD) associated with enter-

ic neuropathy.[75] In this dataset, Kim et al have previously shown that OD patients have signifi-

cantly reduced intestinal expression of Gdnf and Ret[75]. Again, using our transfer learning tool

*ProjectR*[57,78], which allowed us to query bulk-RNAseq data across species to estimate relative

use of MENs-specific NMF-patterns generated earlier, we tested whether the usage of MENs-

specific NMF-patterns was significantly altered between transcriptomes of OD patients and

healthy humans (n = 3/group)[75]. This projection analysis indicated that 2 of the 4 MENs-specific

NMF-patterns showed significantly higher usage in OD samples compared to controls (Fig 11a,

Suppl. Fig 9; Student"s t-test, $p < 0.05$), providing evidence that the enteric neuropathy in these

patients is associated with a relative increase in MENs-specific transcriptional signatures. We

further queried the differential gene expression analysis performed by Kim et al and found that

in addition to Gdnf and Ret (as reported by them) other important NEN genes (Snap25 and

Nos1) were also significantly downregulated while the MEN marker genes identified in this study

(Clic3, Cdh3, Slc17a9) (Fig 11b) were significantly upregulated.

**Discussion**

Current dogma states that the adult ENS is exclusively derived from neural crest precursors that

populate the gut during fetal development[79]. The results of this study indicate a much more

complex system, one in which the fetal and early life "juvenile" ENS consisting of neural crest-

derived enteric neurons (NENs) is incrementally replaced during maturation by mesoderm-

derived enteric neurons (MENs). Eventually, the aging ENS consists almost exclusively of the

neurons of the MEN lineage. Using a combination of floxed transgenic reporters bred into multi-

ple lineage fate mapping transgenic mice, this study also provides the first definitive evidence of

a significant mesodermal contribution to any neural tissue. Previously, the mesoderm and espe-

cially the Lateral Plate Mesoderm (LPM) was known to give rise to diverse cell-types within the

viscera, including several in the gut[80], and our study shows that cells of this embryonic deriva-

tion also give rise to ENS neurons. A previous report on a dual origin of the ENS described adult

enteric neurons from Foxa2$^+$ or Sox17$^+$ precursors[14] and inferred that these were endodermal in

origin. However, Foxa2 and Sox17 are also expressed by mesoderm-derived cells[24-27]. By con-

trast, using two NC-lineage-traced mouse lines (Wnt1-cre and Pax3-cre), two lineage-traced

mouse lines marking mesodermal derivatives (Tek-cre, and Mesp1-cre), and robust adult mes-

oderm markers, we identify a population of Mesp1-derived MHCst- and MET-expressing adult

enteric neurons, which makes up the entire non-NC-derived population of myenteric neurons.

These results provide robust evidence that the second source of adult enteric neurons is the

mesoderm, and not the endoderm.

In our single cell transcriptomic analysis, we used the markers Calcb, Met, and Cdh3 that we

had validated to identify and annotate the MEN cell cluster. When compared with the Ret and

Sox10-expressing cells within the various clusters of NC-derived cell populations, an additional

set of marker genes were found to be expressed widely or selectively within MENs. These

markers, which include both novel as well as established enteric neuronal markers[2,55,81], help

establish the neuronal nature of MENs and at the same time help differentiate them from NENs.

Further, our bioinformatic analyses and immunofluorescence analyses of select MENs markers

helped identify MENs in publicly available murine dataset generated by May-Zhang et al[56], and

in transcriptomic data from the human gut cell atlas[59], thus extending the validity of our observa-

tions to other published data. While May-Zhang-et al. did not further interrogate the develop-

mental origins of the „mesenchymal" enteric neurons to establish the etiology of their distinct

transcriptomic signature, identification and correct annotation of a population of cells as MENs

in other murine and human datasets would not have been possible since these were performed

either exclusively on neural crest-derived ENS cells[46], or when done in a more agnostic manner,

only applied known canonical neural markers (such as Ret and Snap25, now known to be spe-

cific to NENs) to identify the ENS population[47,59,82]. Along with our data, the lack of SNAP-25

expression in a large population of hippocampal neurons[83] shows the limitations of using it as a

pan-neuronal marker. Importantly, we were also able to confirm the presence of cells express-

ing MENs and neuronal markers in human scRNAseq data, and the expression of MENs mark-

ers MHCst, MET and SLC17A9 in many enteric neurons in adult humans, suggesting that the

mesoderm-derived ENS may be a feature common to mice and humans alike. MENs in both

species can readily be identified by their expression of MENs-specific markers, thus providing a

convenient tool to further discriminate and study these neurons.

While the co-expression of pan-neuronal markers such as Elavl2, Pde10a, Tubb2b, Stmn2,

Vsnl1, and Stx3 establishes the neuronal nature of MENs, the differential expression of set of

key genes between the two neuronal populations (MENs versus NENs) suggests putative spe-

cialized functional roles of these subpopulations when they co-exist in adults. These include the

expression of genes associated with sensory (Calcb)[84], neuromodulatory (Ntf3)[85], secretory

(Cftr)[86], immunomodulatory (Il18)[2], and motility-regulating roles (Slc17a9 or VNUT)[87]. These

MEN-enriched or MEN-specific expression profiles suggest that MENs perform various roles in

the regulation of diverse gastrointestinal functions. While the current scRNAseq data is restrict-

ed in providing meaningful cell clusters that can inform the true functional diversity of MENs, fu-

ture work will provide detailed analyses and functional characterization of MENs.

The presence of small population of MENs in the gut at the post-natal age day 10 and their ex-

pansion during maturation suggests that key timepoints in their development correspond with

the post-natal development of the ENS. A recent study by Parathan et al.[88] highlighted important

neuronal and neurochemical changes during this time. Further studies on ENS maturation thus

need to be performed in the context of the two neuronal lineages to truly understand the nature

of the adult ENS. A crucial aspect of this work will require identifying the cellular population that

serves as the progenitor for MENs. Our earlier work showed that Nestin-expressing neural pro-

genitor cells are neural crest-derived cells that contribute to steady state neurogenesis in the

adult ENS[19], thus suggesting that MENs are derived from non-Nestin$^+$ precursor cells.

Since our scRNAseq data highlighted the lineage-specific nature of the expression of Ret and

Met, we studied whether these genes regulated the origin, expansion, and maintenance of the

neuronal populations that express them. While it is known that Gdnf expression in the mature

gut is significantly downregulated[43,62,71,89], the functional consequences of this loss have been

unclear[62]. We found that reduced GDNF-RET signaling drives the age-dependent loss of NENs

as this loss can be slowed or reversed by GDNF supplementation or accelerated by further re-

duction of Ret expression. In aging animals, GDNF-driven resuscitation of NENs was associat-

ed with a functional normalization of age-associated delay in intestinal transit. Our results identi-

fy a novel role for GDNF in maintaining or resuscitating the canonical NEN population, while

preserving overall enteric neuronal numbers. In the last few years, studies have focused on

identifying juvenile protective factors (JPFs), the loss of which correlates with or drives matura-

tion- and aging-associated phenotypes[90]. In this context, GDNF may therefore qualify as a JPF

or a *senolytic* as its presence maintains the dominance of NENs in maturing ENS, correspond-

ing to a juvenile phenotype; and its re-introduction promotes and resuscitates the genesis of

NENs in adult and aging gut to restore healthy gut function. The exact nature of the cells that

respond to GDNF re-introduction and generate NENs is yet unknown, but it can be hypothe-

sized that these may include Nestin$^+$ enteric neural stem cells and/or GDNF-responsive

Schwann cells[19,91]. In addition to its effect on resuscitating NENs, acute administration of GDNF

also augments peristaltic motility by increasing neuronal activity[22,92]. Since the aging gut has a

very small number of RET-expressing GDNF-responsive NENs, we assume that GDNF-driven

normalization of intestinal motility would first increase the proportions of RET-expressing NENs

and then augment their activation. This further emphasizes the importance of using GDNF or

similar RET agonists in the treatment of age-associated intestinal dysmotility.

The observed mutually exclusive expression of RET and MET by the two lineages of enteric

neurons in the mature adult ENS is consistent with an earlier study that reported that a deletion

of Met in the Wnt1-cre expressing NC-lineage did not alter the abundance of MET$^+$ neurons in

the adult ENS[43]. The immunofluorescence analyses on the basis of protein expression on the

mutually exclusive nature of RET and MET provided by us and by Avetisyan et al.[43] contrasts

with the co-expression of Ret and Met transcripts observed in the scRNAseq data on neural

crest-derived enteric neurons from Zeisel et al.[46], which we believe highlights the well-known

discrepancy between single cell/nuclear transcriptomic approaches and actual protein expres-

sion[93]. In this study, we also show that MENs are dependent on HGF-MET signaling in a man-

ner analogous to the requirement of GDNF-RET signaling by NENs. Further, Ret haploinsuffi-

ciency-mediated loss of the NEN lineage causes a proportional increase in the MEN lineage.

This suggests the presence of yet unknown quorum sensing signaling mechanisms between the

two lineages that reciprocally regulate their populations to maintain the structure of the post-

natal ENS. This again implies the existence of a yet undefined precursor cell responsible for the

expansion of the MEN population in a manner analogous to what we have previously described

for NENs[19].

While Avetisyan et al. showed that MET expression highly correlated with CGRP-expression in

the adult murine myenteric plexus[43], MET co-expression by all MHCst$^+$ human enteric neurons

in our data and by all PGP9.5$^+$ neurons in human enteric neurons in the earlier report by Avetis-

yan et al.[43] suggests that MET-expression in the human ENS may not be restricted to CGRP-

expressing IPANs. In addition, recent data shows that our understanding of the nature of mark-

ers that can correctly identify IPANs in the murine and human ENS is still evolving[56,94], suggest-

ing that ascribing function to neurons using a single marker may be incorrect.

The detection of MENs cluster in the post-natal and adult scRNAseq data in the gut cell atlas

generated by Elmentaite et al.[59], their increasing representation with age, and the subsequent

validation of the MEN-specific markers MHCst, MET and SLC17A9 in a subset of adult human

ENS neurons shows that our findings extend to human ENS and may be of clinical importance.

Many gastrointestinal motility functions are compromised with advancing age, resulting in clini-

cally significant disorders[95]. Although the exact mechanisms will need to be worked out, our re-

sults indicate that a MENs-predominant ENS is associated with significant differences in gut mo-

tility. The progressive nature of the change in enteric neuronal lineages with age may have

pathological implications for the elderly gut when the balance is overwhelmingly in favor of

MENs. Understanding the forces that regulate parity between these two different sources of

neurons and the associated processes of genesis of these neurons therefore holds the promise

of arresting or reverting progressive loss of gut motility with increasing age. Our results may al-

so have implications for the pathogenesis of disordered motility unrelated to aging, as downreg-

ulation of Gdnf and Ret expression has been associated with diverticular disease and obstruct-

ed defecation in adult patients[72,75,96]. GWAS analyses further showed that Gdnf gene was in the

eQTL associated with increased incidence of diverticular disease[97]. It is in this context that the

identification of SNAP-25 as a NENs marker gene and the transcriptomic pattern analyses of

patients with chronic obstructed defecation are significant. Expression of Snap25 is upregulated

by GDNF and is significantly downregulated in gut tissues of patients with diseases associated

with significant reduction in GDNF-RET signaling (diverticular disease and obstructed defeca-

tion)[61,72,75]. Thus, while earlier thought to be a pan-neuronal marker[47,98], establishing the identity

of SNAP-25 as a NEN lineage-restricted marker provides us not only with an important tool to

query proportions of NENs in murine and human tissue but also suggests the NENs-limited na-

ture of prior observations based on Snap25 transgenic animals[60,98]. Gut dysmotility disorders

that present with conserved ENS neuronal numbers[72] have puzzled investigators on the etiology

of the disease. Our results suggest an explanation based on NENs-to-MENs lineage shifts that

conserve overall enteric neuronal numbers with variable effects on function.

In conclusion, using multiple lines of evidence, we show that the ENS of the juvenile and the

aging mammalian gut is dominated by two different classes of neurons that have distinct on-

togenies and trophic factors, associated with functional differences. The shift in neuronal lineage

may represent a functional adaptation to changes in nutrient handling, microbiota or other phys-

iological factors associated with different ages.  Further research in the functional differences

between the neuronal lineages and factors that regulate the parity between these two nervous

systems in humans during the lifetime will be important to advance our understanding of the

adult ENS and the treatment of age-related and other pathological disorders of gut motility.

**Acknowledgements**

We would like to thank Dr. Chulan Kwon (JHU) for his kind gift of the Mesp1-cre mice, Dr. Jeremy Nathans (JHU) for his kind gift of the Tek-EGFP, Tek-cre and *Hprt*-tdTomato lineage-traced mice and his support, and Dr. Vanda Lennon (Mayo Clinic) for her kind gift of the ANNA1 anti-Hu antisera. We thank Dr. Akira Sawa (JHU) for his help with the *Rosa26*-EGFP mouse line. We thank Dr. Xinzhong Dong (JHU) and Dr. Mark Donowitz (JHU) for their support. Microscopy was performed on the Ross Imaging Core at the Hopkins Conte Digestive Disease Center at the Johns Hopkins University (P30DK089502) using the Olympus FV 3000rs (procured with the NIH-NIDDK S10 OD025244 grant) and we thank Dr. George McNamara for his help with confocal training and imaging. The 10X Genomics Chromium processing for scRNAseq was performed at the GRCF Core and the sequencing was performed at the CIDR core at the Johns Hopkins University. S.K. was funded through a grant from the Ludwig Foundation, a grant from the NIA (R01AG066768), a pilot award from the Hopkins Digestive Diseases Basic & Translational Research Core Center grant (P30DK089502). L.A.G was funded in part by a Johns Hopkins Catalyst Award and a grant from the NIA (R01AG066768). J.S. was funded through the Maryland Genetics, Epidemiology, and Medicine training program sponsored by the Burroughs Welcome Fund. P.J.P. was funded through the Hopkins Conte Digestive Disease Center at the Johns Hopkins University (P30DK089502), NIDDK (R01DK080920), the Maryland Stem Cell Research Foundation (MSCRF130005), and a grant from the AMOS family.

**Author contributions**

S.K., and P.J.P. conceived the study; S.K., M.A., M.L, L.A.G. and P.J.P. designed the research studies; S.K. L.B, C.Z, A.S., A.B., and M.S. conducted the experiments; S.K., Z.H., M.K., J.L., L.B., S.B., E.V., J.S., S.N., M.S., G.L., and L.A.G. acquired and analyzed the data; S.K., and P.J.P drafted the manuscript.

**Methods**

*Animals*:

Experimental protocols were approved by The Johns Hopkins University"s Animal Care and Use Committee in accordance with the guidelines provided by the National Institutes of Health. Presence of vaginal plug was ascertained as 0.5 days post-fertilization and this metric was used to calculate age of mice. Only male mice were used for the studies detailed in this report. The Wnt1-cre:*Rosa26*-tdTomato lineage-traced line was generated as detailed before by breeding the B6.Cg-Tg(*Wnt1-cre*) with the Ai14 transgenic mouse line (Jax #: 007914) containing the *Rosa26*- tdTomato transgene[19,99,100]. The Wnt1-cre line used in this study is the well validated line for studying neural crest derivatives previously used by us and others[18,19,22,46,101], and not the Wnt1-cre2 line used by Drokhlyansky et al.[102]. We again validated the fidelity of this line by observing whether any aberrant recombination occurred in our Wnt1-cre:*Rosa26*-tdTomato line by studying the intestinal mucosal tissue and found that the line behaved expectedly to label neural crest-derivatives and epithelial cells (Suppl. Fig 1i). Pax3-cre:*Rosa26*-tdTomato lineage-traced line was generated by breeding the Ai9 transgenic mouse line (Jax #: 007909) with the Pax3-cre transgenic mouse (Jax #: 005549). The Wnt1-cre:*Hprt*-tdTomato mouse was generated by breeding our aforementioned Wnt1-cre transgenic mouse line with the *Hprt*-tdTomato transgenic mouse line (Jax #: 021428, kind gift from Prof. Jeremy Nathans). Mesp1-cre:*Rosa26*-tdTomato mice were generated by breeding the Mesp1-cre transgenic mice[103] (gift from Dr. Chulan Kwon, JHU) with the Ai14 transgenic mice. Mesp1-cre:*Rosa26*-GFP mice were generated by breeding the Mesp1-cre transgenic mice with the *Rosa26*-EGFP transgenic mouse line (gift from Dr. Akira Sawa, JHU). Neither the *Rosa26*-tdTomato (Ai14) line nor the *Rosa26*-EGFP line showed any reporter expression in the absence of a cre driver (Suppl. Fig 1j, k). Ret[+/CFP] mice (MGI:3777556) were inter-bred to get a colony of adult Ret[+/+] and Ret[+/CFP] mice. Ret[CFP/CFP]

mice died at or before term. Tek-cre:*Hprt*-tdTomato mice were generated by breeding Tek-cre

transgenic mice (also known as Tie2-cre; Jax #: 004128) with the *Hprt*-tdTomato transgenic

mouse line. 17-month-old male C57BL/6 mice from the aging mouse colony of the National In-

stitute of Aging were procured for the GDNF-treatment experiment.

*Human Tissues:*

Human tissues were obtained under IRB protocol IRB00181108 that was approved by Institu-

tional Review Board at the Johns Hopkins University. Pathological normal specimens of human

duodenum and colon were obtained post-resection. Tissues were obtained from adult donors

and were de-identified such that the exact age, gender, and ethnicity of the donors was un-

known.

*Tissue preparation*:

Mice were anesthetized with isoflurane and sacrificed by cervical dislocation. A laparotomy was

performed, and the ileum was removed and lavaged with PBS containing penicillin-streptomycin

(PS; Invitrogen), then cut into 1-cm-long segments and placed over a sterile plastic rod. A su-

perficial longitudinal incision was made along the serosal surface and the LM-MP was peeled off

from the underlying tissue using a wet sterile cotton swab[19] and placed in Opti-MEM medium

(Invitrogen) containing Pen-Strep (Invitrogen). The tissue was then laid flat and fixed within 30

minutes of isolation with freshly prepared ice cold 4% paraformaldehyde (PFA) solution for 45

minutes in the dark to preserve fluorescence intensity and prevent photo-bleaching. All LM-MP

tissues post-isolation were fixed within 30 minutes of their isolation. After the fixation, the tissue

was removed and stored in ice cold sterile PBS with Pen-Strep for immunofluorescence staining

and subsequent microscopy.

For human tissues, duodenal tissue from adult human patients (n = 3 patients), who did not

have any prior history of chronic intestinal dysmotility, that was removed by Whipple procedure

was obtained. A colonic sample from a pathologically normal colonic resection from an adult

donor suffering from colon carcinoma who similarly did not have prior history of chronic intesti-

nal dysmotility was also obtained. The resected tissue was placed in ice cold Opti-MEM medium

(Invitrogen) containing Pen-Strep (Invitrogen). The mucosal and sub-mucosal tissue was dis-

sected out in the medium under light microscope and the muscularis layer containing myenteric

plexus tissue was obtained. The tissue was laid out between two glass slides and fixed over-

night in ice cold 4% PFA after which it was removed and stored in ice cold sterile PBS with Pen-

Strep for immunofluorescence staining, optical clarification, and subsequent microscopy.

*Immunohistochemistry*:

For murine tissue: The fixed LM-MP tissue was washed twice in ice-cold PBS in the dark at 16

°C. The tissue was then incubated in blocking-permeabilizing buffer (BPB; 5% normal goat se-

rum with 0.3% Triton-X) for 1 hour. While staining for antibodies that were mouse monoclonal,

5% normal mouse serum was added to the BPB. The tissue was then removed from the BPB

and was incubated with the appropriate primary antibody at the listed concentration (Suppl. Ta-

ble 1) for 48 h at 16 °C in the dark with shaking at 55 rpm. Following incubation with primary an-

tibody, the tissue was washed three times (15-min wash each) in PBS at room temperature in

the dark. The tissue was then incubated in the appropriate secondary antibody at room temper-

ature for 1 hour while on a rotary shaker (65 rpm). The tissue was again washed three times in

PBS at room temperature, counterstained with DAPI to stain the nuclei, overlaid with Prolong

Antifade Gold mounting medium, cover-slipped, and imaged.

Colchicine treatment: For CGRP immunostaining, mice were injected with Colchicine at a con-

centration of 5 mg/Kg body weight 16 hours (overnight) before they were sacrificed. The mice

were housed singly during this time and adequate gel packs were provided. Food and water

were provided *ad libitum*. On the following day, the mice were sacrificed, and their LM-MP tis-

sues were harvested as detailed above.

For human tissue: The fixed muscularis layer containing myenteric plexus tissue was removed

from ice cold PBS and incubated in blocking-permeabilizing buffer (BPB; 5% normal goat se-

rum, 5% normal mouse serum with 0.3% Triton-X) for 4 hours. The tissue was then removed

from the BPB and was incubated with the appropriate primary antibody at the listed concentra-

tion (Suppl. Table 1) for 5 days at 16°C in the dark with shaking at 55 rpm. Following incubation

with primary antibody, the tissue was washed five times (15-min wash each) in PBS at room

temperature in the dark. The tissue was then incubated in the appropriate secondary antibody at

16°C in the dark with shaking at 55 rpm for 2 days. The tissue was again washed in dark for five

times in PBS that contained DAPI at room temperature. After the final wash, the tissue was

suspended in tissue clarification buffer CUBIC[104] for 1 hour at 4°C in the dark after which it was

overlaid with Prolong Antifade Gold mounting medium, cover-slipped, and imaged. Briefly, the

CUBIC optical clarification buffer was made by mixing 2.5 g of urea  25   by wt , 2.5 g of N, N,

N ,N -tetrakis (2-hydroxy-propyl) ethylenediamine (25% by wt), 1.5 g of Triton X-100 (15% by

wt) in 35 ml of Distilled Water. The solution was shaken till the ingredients were dissolved and

yielded a clear viscous solution.

*Microscopy*:

Imaging was done by using the oil immersion 63X objective on the Leica SP8 confocal micro-

scope and by using the oil immersion 40X objective on the Olympus Fluoview 3000rs confocal

microscope with resonance scanning mode. For thick tissues, such as human tissues, the Gal-

vano mode of the Olympus Fluoview 3000rs microscope that enabled higher resolution imaging

and averaging was used. Images obtained were then analyzed using Fiji (https://fiji.sc/).

Live tissue imaging of the mouse gut tissue was performed by harvesting small intestinal tissue

from an adult Wnt1-cre:*Rosa26*-tdTomato mouse and immediately putting it in OptiMEM solu-

tion. The tissue was then immediately put on a chamber slide containing OptiMEM and imaged

in live tissue culture conditions of the EVOS M7000 microscope under the 20X objective.

*Enumeration of neurons*:

Enumeration of tdTomato$^+$ and tdTomato$^-$ neurons was performed on tissue with native tdToma-

to fluorescence. The native tdTomato fluorescence in fixed tissue was intense, thus there was

no need to increase fluorescent signal using anti-RFP/tdTomato antibodies.

Enumeration of enteric neurons to study alterations to ENS neuronal numbers was performed

by following the well-established protocol[105-108] of counting the numbers of neurons in myenteric

ganglia and comparing them between animals for statistically significant differences. Identifica-

tion of myenteric ganglia was performed according to our pre-determined method published ear-

lier[19]. Briefly, contiguous clusters of neurons were defined as a ganglia and the total numbers of

neurons within these clusters were enumerated as numbers of myenteric neurons per ganglion.

As a rule, clusters of 3 neurons or more were deemed to consist a ganglion and our enumera-

tion strategy did not count extra-ganglionic neurons. We imaged ~10 ganglia per tissue for our

enumeration and each group studied had n ≥ 3 mice. Identification and enumeration of neurons

and detection of co-localization was performed manually by trained laboratory personnel. Fig 1d

comprises of the data from 6 P60 mice, which includes data from the 3 mice that were used for

the development experiment presented in Fig 8c.

*Protein isolation and detection:*

After the LM-MP tissue was isolated, it was weighed and placed in a sterile 1.5 ml microfuge

tube. 1X RIPA buffer (Cell Signaling Technology) with Halt Protease Inhibitor Cocktail (Thermo

Scientific) at 5X concentration, Phosphatase Inhibitor Cocktails II and III (Sigma-Aldrich) at 2X

concentrations were added to the tissue lysate buffer. Tissue was disrupted using 1.0 mm silica

beads in Bullet Blender 24 (Next Advance) for 5 mins at highest setting. The lysate was incu-

bated at 4°C with shaking for 30 mins, centrifuged at 14,000rpm for 20 mins and the superna-

tant was taken and stored in -80°C in aliquots.  Protein concentration was estimated using Brad-

ford assay solution (Biorad) following the manufacturer"s protocol. For immunoblotting, 40 µg of

protein was loaded per well of 4%-20% gradient denaturing gel (Biorad). Protein marker used

was Precision Plus Dual Color standards (Biorad). After fractionating the proteins, they were

blotted onto ImmunBlot PVDF membrane (Biorad) overnight at 4°C at 17V for 12-16 hours. After

blotting, membrane was blocked with Odyssey TBS blocking buffer (Li-Cor) for 1 hour at room

temperature with shaking. Incubation with primary antibodies were carried out at 4°C with shak-

ing for 24 hours. Following binding, the blot was washed 4 times with TBS-T (Tris Buffered Sa-

line with 0.5% Tween) for 15 mins each with shaking at room temperature. Secondary antibody

incubation was carried out in dark at room temperature for 1.5 hours with shaking. The blot was

then washed 4 times for 15 mins each and imaged on Odyssey CLx system (Li-Cor).

Antibody used to detect RET was a well validated antibody used previously for detecting RET in

ENS[109] and was further validated by us using a western blot on total proteins isolated from mu-

rine small intestinal LM-MP (Suppl. Fig 8e). Antibodies used are detailed in the Supplementary

Table 1.

*RNA isolation and quantitative detection of specific transcripts:*

The isolated tissue was stored in RNALater Solution (Ambion). RNA was isolated using RNeasy

Mini Kit Qiagen following manufacturer"s protocol. RNA quantification was carried out using

Epoch Microplate Spectrophotometer (BioTek). cDNA synthesis was carried by SuperScript IV

VILO Master Mix (Invitrogen). Quantitative Real-time PCR was carried out using Taqman Gene

Expression Master Mix (Applied Biosystems) and Roto-Gene Q (Qiagen). The probes used are

listed in Supplementary Table 1.

*Single Cell RNA sequencing and analyses:*

Single cell preparation from adult murine ileal tissues: Ileal tissues from two 6-month old adult

male littermate C57/BL6 wildtype mice were isolated by peeling as previously described. The

tissues were then dissociated in Digestion Buffer containing 1 mg/ml Liberase (Sigma-Aldrich) in

OptiMEM. Tissues from mouse 1 were dissociated in the Digestion buffer containing Liberase

TH and tissues from mouse 2 were dissociated in the Digestion buffer containing Liberase TL.

Dissociation was performed at 37°C for 30 minutes on a rotary shaker, after which the cells

were centrifuged at 200g for 7 minutes, and the pellet was resuspended in ice cold sterile PBS.

The cell suspension was passed through a 40 μm cell sieve and the resulting filtered cell sus-

pension was again centrifuged at 200g for 7 minutes. This process of cell centrifugation and fil-

tration was repeated two more times, after which the cells were resuspended in 1 ml ice cold

sterile PBS. The repeated steps of serial cell washes and filtration removed clumps and debris

and the viability of the resulting cell suspension was estimated to be >90% using Trypan Blue

dye test. The cells were then processed through 10X Genomics Chromium V2.0 system accord-

ing to the manufacturer"s suggested workflow. The processing was done at the GRCF Core Fa-

cility at the Johns Hopkins University. The pooled libraries were sequenced on an Illumina HiS-

eq 2500 to an average depth of $3.125 \times 10^8$ reads per sample library. The sequencing was per-

formed at the CIDR core facility at the Johns Hopkins University.

Pre-processing of FASTQs to Expression Matrices: FASTQ sequence files were processed fol-

lowing a Kallisto Bustools workflow compatible with downstream RNA velocity[110]. References

required for pseudo-alignment of transcripts were obtained using the get_velocity_files (func-

tionality of BUSpaRSE (https://github.com/BUStools/BUSpaRse)), with "L=98" for 10X Ge-

nomics v2.0 sequencing chemistry. Reads were pseudo-aligned to an index built from Ensembl

97 transcriptome annotation (Gencode vM22; GRCm38). Across two samples processed, a total

of 578,529,125 reads were successfully pseudo-aligned. Barcodes within a Hamming distance

of one to known 10X Genomics v2.0 barcodes were corrected. Reads were classified as

"spliced" or "unspliced" by their complement to the target list of intronic sequences and exonic

sequences, respectively, and subsequently quantified separately into expression count matri-

ces. Spliced counts are used for all analyses.

Single cell gene expression analysis: scRNA-seq count matrices were analyzed using Mono-

cle3. 11,123 high-quality cells were identified as meeting a 200 UMI minimum threshold with a

mitochondrial read ratio of less than 20%; droplets that did not meet these criteria were exclud-

ed from the analysis. Mitochondrial counts were determined as the sum of reads mapping to 37

genes annotated to the mitochondrial genome. All genes with non-zero expression were includ-

ed for downstream analysis. Raw counts were first scaled by a cell-size scaling factor and sub-

sequently log10 transformed with a pseudo-count of 1. Normalized values are used in place of

raw counts unless otherwise noted.

Prior to UMAP dimensionality reduction, batch effects between the two biological replicates

were assessed and corrected via the mutual nearest neighbors (MNN) algorithm as implement-

ed by Batchelor in Monocle3[111]. 15 clusters of cells in the UMAP embedding were identified by

Leiden community detection. 30 marker genes for each cluster were identified based on great-

est pseudo $R^2$ values and used for supervised annotation of cell types by searching UniProt,

Allen Cell Atlas and through literature search with Pubmed.

NC-derived cell clusters were identified by expression of NC markers Ret and Sox10. MEN clus-

ter was identified by its expression of CGRP-coding Calcb, Met, and Cdh3. The pan-MENs pro-

tein marker MHCst was identified by labeling with an antibody S46 which labels all members of

the MHCst family[40]. Since the antibody does not identify a single gene product, MHCst im-

munostaining could not be used to identify a specific gene marker for use in the annotation of

the MEN cluster. For further analysis into the MENs population, the full LM-MP dataset was

subset to include only the 2,223 cells annotated as MENs. These cells were re-processed as

above, but with a reduced PCA dimensionality of k=20 as input for the UMAP embedding. 5

clusters of cells in the UMAP embedding were identified by Leiden community detection (k= 10,

resolution = 5e -4).

External datasets: Spliced and unspliced count matrices for mesenchymal subset of the Gut

Cell Atlas were obtained from https://www.gutcellatlas.org/, and existing annotations were used

when available. SnRNAseq counts matrices from May-Zhang et al.[56] were downloaded from

GEO GSE153192. Bulk RNAseq dataset from the Human Obstructed defecation study by Kim

et al.[75] was obtained from GEO GSE101968.

Pattern discovery and ProjectR analyses: Pattern discovery was utilized to identify sets of co-

expressed genes that define cell-type specific transcriptional signatures. The normalized ex-

pression matrix was decomposed via non-negative matrix factorization (NMF) as implemented

in the R package *NNLM*, with k=50 and default parameters. Cell weights for each pattern were

grouped by assigned cell-type and represented by heatmap. Pattern vectors were hierarchically

clustered by a Euclidian distance metric, implemented in ComplexHeatmap[112]. These patterns

were then tested on the bulk RNA-Seq expression matrix for the Human Obstructed Defecation

study[75]. The log2 expression (log2(rpkm + 1)) from this study was projected onto the NMF pat-

terns using projectR[57]. Students" t tests were performed on the projection weights from the Con-

trol and OD groups to test for differences between them. For the projection of datasets generat-

ed by May-Zhang et al. and Elmentaite et al., log10-transformed normalized counts were pro-

jected.

**Analysis of May-Zhang et al. dataset**

Raw UMI count matrices for the single-nucleus RNA sequencing of enteric neurons performed

by May-Zhang et al.[56] from the ileum of adult mice were obtained from GEO (GSE153192). The

DropletUtils package (v1.10.3)[113,114] was used to compute barcode ranks, knee, and inflection

point statistics for all the samples. All barcodes in the 10X genomics libraries with total UMI

greater than the inflection point were considered to be cell-containing droplets and retained for

further analysis. The inDrop libraries were filtered to retain all barcodes with UMI greater than

200. The filtered matrices were combined to form a SingleCellExperiment object.

Cell-specific scaling factors accounting for composition biases were computed using com-

puteSumFactors (scran v1.18.7)[115]. The raw UMI counts were then scaled using the scaling fac-

tors and log 2 transformed after addition of a pseudo count of 1 using logNormCounts (scater

v1.18.6)[116]. Batch effects between the samples were corrected using fastMNN (batchelor

v1.6.3)[111].

Unsupervised clustering was performed using a graph-based clustering technique. The graph

was obtained using buildSNNgraph (scran v1.18.7)[115] and the louvain algorithm implemented in

the igraph package (v1.2.7)[117] was used to identify communities. Differential expression testing

was performed between all pairs of clusters using findMarkers (scran v1.18.7)[115]. One-sided

Welch"s t-test was used to identify genes that were upregulated in a cluster as compared to any

others in the dataset. Benjamini-Hochberg method was applied (to combined p-values) to cor-

rect for multiple testing and the genes were ranked by significance. The top ranked genes were

used as marker genes for the clusters. The log fold change from the pairwise comparison with

the lowest p-value was used as the summary log fold change.

The log normalized counts from this dataset were then projected into the NMF patterns learned

on our scRNAseq data. This was performed using transfer learning implemented in projectR

(v1.6.0). The projection weights corresponding to the four MEN-specific patterns were then clus-

tered    using    graph-based    clustering    and    visualized    by    UMAP    (uwot    v0.1.10)(

https://github.com/jlmelville/uwot).

*Whole gut transit time analyses*:

Whole-gut transit time (WGTT) for every mouse was analyzed by the method using the carmine

red protocol[19]. Mice were placed in individual cages and deprived of food for 1 hour before re-

ceiving 0.3 mL 6% (wt/vol) carmine solution in 0.5% methylcellulose by oral gavage into the an-

imal"s stomach. The time taken for each mouse to produce a red fecal pellet after the admin-

istration of carmine dye was recorded in minutes. The experiment was terminated at 210

minutes post-gavage and the WGTT of any mice that did not expel the red dye at the termina-

tion was marked at the value of 210 min. The mean difference in whole gut transit time (in

minutes) between both the $Ret^{+/+}$ and $Ret^{+/-}$ mice cohorts, and the GDNF and Saline-treated

Control cohorts were analyzed statistically.

*In vivo Injections:*

GDNF injection: Similar to prior report that gave sub-cutaneous injections of GDNF to post-natal

mice[71], we took 6 littermate 10 day old (P10) male Wnt1-cre:*Rosa26*-tdTomato mice and divided

into two subgroups, GDNF and Control. Each mouse in the GDNF group was injected sub-

cutaneously with 50 µl of 2mg/ml of GDNF (Peprotech Catalogue #: 450-44) every other day,

while the Control group was injected with 50 µl of sterile saline. The mice were given 5 doses

and then sacrificed on P20, after which their LM-MP tissues were isolated as detailed above.

The tissues were then immunostained with antibodies against HuC/D and imaged. For studying

the effect of GDNF on aging mice, two cohorts of 17-month-old male C57BL/6 mice (n=5

mice/cohort) were obtained from the aging colony of the National Institute of Aging. Before the

start of dosing, the whole gut transit time was assayed. The animals were then injected daily

sub-cutaneously either with 100 µl of saline (Control) or 100 µl of GDNF (500 µg/ml) for 10 con-

secutive days, after which the mice were sacrificed and their LM-MP tissues were isolated as

detailed above. The tissues were then immunostained with appropriate antibodies and imaged.

HGF injection: Similar to prior report that gave sub-cutaneous injections of HGF to post-natal

mice, we took 5 littermate 10 day old (P10) male Wnt1-cre:*Rosa26*-tdTomato mice and divided

into two subgroups, HGF and Control. Each mouse in the HGF group was injected sub-

cutaneously with 100 µl of 2mg/ml of HGF (Peprotech Catalogue #:315-23) every other day,

while the saline group was injected with 100 µl of sterile saline. The mice were given 5 doses

and then sacrificed on P20, after which their LM-MP tissues were isolated as detailed above.

The tissues were then immunostained with antibodies against HuC/D and imaged.

*Statistics:*

Data was analyzed using Graphpad Prism 8.3.1 and R using Unpaired Students t-test, Simple

Linear Regression, and Ordinary One-Way ANOVA.

*Data:*

All raw data are provided in Supplementary Table 2 - 4. Raw single cell RNA sequencing data is

archived on the NCBI GEO server and can be accessed under the accession number

GSE156146.

*Code:*

Code generated for the scRNA/snRNA analyses is available at

https://github.com/jaredslosberg/6month_ENS/.

**Figures**

**Fig 1: Half of all adult small intestinal myenteric neurons are derived from a non-neural crest lineage.**

**(a)** Enteric glia, labeled with GFAP (green) in the myenteric plexus from an adult Wnt1-cre:*Rosa26*-tdTomato mouse. Inset is magnified in color segregated panels, showing GFAP$^+$ glia co-expressing tdTomato (red, red arrows). Nuclei are labeled with DAPI (blue). Scale bar = 10 μm.

**(b)** 2 dimensional representation of a 3 dimensional stack of images shows enteric neurons, labeled with HuC/D (green), in the myenteric plexus from an adult Wnt1-cre:*Rosa26*-tdTomato mouse. Enteric neurons either express tdTomato (red arrow) or not (white arrows). Nuclei are labeled with DAPI (blue). Scale bar = 10 μm.

**(c)** Orthogonal views of the image **(b)** shows the lack of tdTomato expression in a particular HuC/D-immunolabeled enteric neuron.

**(d)** Quantification of tdTomato$^+$ and tdTomato$^-$ neurons in the myenteric ganglia of 6 P60 Wnt1-cre:*Rosa26*-tdTomato mice. Data represents Mean ± SEM.

**Fig 2: Mesoderm-lineage-tracing and marker expression provide evidence of the meso-dermal-derivation of half of all adult small intestinal myenteric neurons.**

**(a)** Small intestinal LM-MP from adult male Tek-cre:*Hprt*-tdTomato shows the presence of Tek-derived and hence mesoderm-derived tdTomato$^+$ (red) HuC/D$^+$ (green) neurons (white arrows) and non-mesoderm-derived tdTomato$^-$ neurons (green arrows). Vascular cells depict higher flu-orescence intensity (yellow arrow) than cells in the ganglia. Nuclei are labeled with DAPI. Scale bar = 10 µm.

**(b)** Cardiac tissue from heart and **(c)** small intestinal LM-MP tissue of adult male Mesp1-cre:*Rosa26*-tdTomato mice shows the presence of variable tdTomato expression in cells of the-se tissues. While cardiomyocytes in heart and cells within the myenteric ganglia (white arrows) exhibit lower tdTomato fluorescence, similar to the tissues from Tek-cre:*Hprt*-tdTomato mice in **(a)**, the vascular cells exhibit higher fluorescence intensity (yellow arrows). Scale bar = 10 µm.

**(d)** Small intestinal LM-MP from adult male Mesp1-cre:*Rosa26*-tdTomato mouse shows the presence of mesoderm-derived tdTomato$^+$ (red) HuC/D$^+$ (green) neurons (white arrows) and non-mesoderm-derived tdTomato$^-$ neurons (green arrows). Nuclei are labeled with DAPI. Scale bar = 10 µm.

**(e)** Small intestinal LM-MP from adult male Mesp1-cre:*Rosa26*-EGFP mouse, when im-munostained with antibodies against GFP and against HuC/D, shows the presence of meso-derm-derived EGFP$^+$ (green) HuC/D$^+$ (red) neurons (white arrows) and non-mesoderm-derived EGFP$^-$ neurons (red arrows). Nuclei are labeled with DAPI. Scale bar = 10 µm.

**(f)** Quantification of tdTomato-expressing and non-expressing neurons in the myenteric ganglia of 3 P60 Mesp1-cre:*Rosa26*-tdTomato mice. Data represents Mean ± SEM.

MHCst immunolabeling exclusively marks all the **(g)** mesoderm-derived adult HuC/D$^+$ (green)

neurons in the small intestinal LM-MP of Mesp1-cre:*Rosa26*-tdTomato mice; and **(h)** all the non-

NC-derived neurons in the small intestinal LM-MP of Wnt1-cre:*Rosa26*-tdTomato mice (white

arrows). MHCst does not label tdTomato$^+$ NC-derived cells (yellow arrows). Nuclei are stained

with DAPI (blue). Scale bar = 10 μm.

**Fig 3: Cellular and molecular phenotyping of MENs and NENs.**

MET immunostaining (green) labels **(a)** Wnt1-cre:*Rosa26*-tdTomato⁻ MENs and **(b)** a population

of Mesp1-cre:*Rosa26*-tdTomato⁺ MENs (white arrow) while not labeling all MENs (red arrow).

**(c)** RET immunostaining (green) only labels Wnt1-cre:*Rosa26*-tdTomato⁺ NENs (green arrow)

and not tdTomato⁻ MENs (white arrows).

**(d)** NOS1 is expressed by both tdTomato⁺ NENs (red, white arrows) and tdTomato⁻ MENs (blue

arrows) in an adult Wnt1-cre:*Rosa26*-tdTomato mouse, but most MENs do not express NOS1

(marked by *).

**(e)** Both tdTomato⁺ (red) and tdTomato⁻ neurons in the myenteric plexus of an adult Wnt1-

cre:*Rosa26*-tdTomato mouse (HuC/D, green) express CGRP (blue) Inset showing a tdTomato⁻

CGRP⁺ neuron (white arrow) is magnified on the right.

**(f)** NEN lineage contains significantly higher proportions of NOS1⁺ neurons compared to MEN

lineage in Wnt1-cre:*Rosa26*-tdTomato mice. Data represent mean ± S.E.M. Student"s t-test ** p

< 0.01.

**(g)** MEN lineage contains significantly higher proportions of CGRP⁺ neurons compared to NEN

lineage in Wnt1-cre:*Rosa26*-tdTomato male mice. Data represent mean ± S.E.M. Student"s t-

test **** p < 0.0001.

**(h)** MENs express the protein P-Cadherin/CDH-3 (green; white arrows) while tdTomato⁺ (red)

NENs do not (blue arrows) in myenteric ganglia from adult Wnt1-cre:*Rosa26*-tdTomato mice.

Nuclei in **(b), (c), (h)** are labeled with DAPI (blue). Scale bar for all images denotes 10 μm.

**Fig 4: scRNAseq-analyses identifies the distinct transcriptomic profile of the MENs.**

**(a)** UMAP representation of 11,123 sequenced high-quality cells that were identified as meeting a 200 UMI minimum threshold with a mitochondrial read ratio of less than 20%. Clusters were annotated by markers that were found to be cell-type specific by searching UniProt, Allen Cell Atlas and Pubmed databases. Cells of the neural crest lineage were then identified as NENs by their expression of neural crest marker gene Ret and Neuroglia by their expression of Ncam1 and Sox10, or as MENs by co-expression of Calcb (CGRP), Cdh3, and Met genes.

**(b)** Visualization of expression of select MEN-specific and neuronal markers using quasi-violin plots. "Max value" represents the scale for the log-normalized expression of each gene.

**(c)** Validation of the MENs-specific marker genes discovered in the scRNAseq analyses by immunohistochemistry and confocal microscopy of small intestinal LM-MP from adult male Wnt1-cre:*Rosa26*-tdTomato mice. Expression of neuronal markers STX3, VSNL1, STMN2, HAND2, GPR88, PDE10A, ELAVL2, and TUBB2B (Gray, green arrows) was found in tdTomato$^-$ MENs. Immunostaining of the proteins AEBP1, CFTR, CLIC3, NT-3, SLPI, SMO, IL-18, SLC17A9, FMO2 and MYL7 (green; green arrows) was found to be localized to tdTomato$^-$ MENs. tdTomato$^+$ (red) NC-cells did not immunostain for these markers (red arrows). Panel also shows Wnt1-cre:*Rosa26*-tdTomato tissue with no primary controls (stained with AlexaFluor 488 and AlexaFluor 647 antibodies). Figures with * annotations are 2D representation of 3D stacks of images, whose orthogonal views are presented in Suppl Fig 3. Nuclei in some panels were labeled with DAPI (blue). Scale bar denotes 10 μm.

**Fig 5: Computational analyses using *Project R* identifies MENs in publicly available mu-**

**rine and human transcriptomic datasets.**

**(a)** Heatmap of cell weights for the patterns learned by NMF (k=50). Hierarchical clustering was

calculated using Euclidean distance. Multiple clusters annotated as the same cell type are

merged. The four most specific MENs patterns (16, 27, 32 and 41) are selected.

**(b)** In addition to using the four MEN-specific NMF patterns to label the MENs cluster in our

scRNAseq dataset (plots in top row), two other transcriptomic datasets: May-Zhang et al."s mu-

rine ileal ENS snRNAseq dataset[56] (plots in second row), and Elmentaite et al."s gut mesen-

chymal scRNAseq dataset[59] (plots in third row), were projected into the defined four MEN-

specific pattern space using *ProjectR*. Raw projected cell weights were visualized using previ-

ously learned UMAP embedding. The cell clusters that show high pattern usage are shown

bounded by the red square. Plots in the third row shows projection of healthy post-natal mesen-

chymal cells from Elmentaite et al"s data into the four MEN-specific NMF patterns, which again

show a population of cells showing high MEN-specific pattern utilization.

1017

**Fig 6. Detection of MENs in human single cell RNA sequencing data.**

**(a)** UMAP representation of scRNAseq data from Elmentaite et al."s[59] healthy post-natal mesen-

chymal cells from the human gut.

**(b)** Breakdown for previously published annotated features of the post-natal health subset of the

gut cell atlas as presented in Elmentaite et al.

**(c)** Expression of neuronal marker genes Slc17a9, Stx3, Uchl1, Tubb2b, Pde10a, and Hand2 in

the UMAP representation of healthy post-natal data from Elmentaite et al., which was annotated

as clusters of mesenchymal cells.

1026

**Fig 7: Validation of SNAP-25 as a NENs marker and detection of MENs in human tissues**

**(a)** SNAP-25 expression is restricted to the neural crest lineage in the adult myenteric ganglia. SNAP-25 expression (green) co-localizes with tdTomato (red) but not with the MENs marker MHCst (cyan) as observed in 2D and orthogonal views of a myenteric ganglia from a Wnt1-cre:*Rosa26*-tdTomato mouse that was immunolabeled with antibodies against MHCst and SNAP-25. Nuclei were labeled with DAPI (blue). Scale bar denotes 10 µm.

**(b)** Putative human MENs (red arrows) and NENs (green arrows) identified by presence or absence of the murine MENs markers, MET (blue) and MHCst (red) in subpopulations of HuC/D[+] (green) small intestinal myenteric neurons from normal human duodenal tissue. Scale bar denotes 10 µm.

**(c)** MENs marker SLC17A9 expression (green, green arrows) in a subset of MHCst-expressing (red) neurons. Subset of MHCst-expressing cells do not express SLC17A9 (red arrow). Nuclei are labeled with DAPI (blue) in the normal human duodenal tissue. Scale bar denotes 10 µm.

**Fig 8: GDNF and HGF signaling regulate age-dependent changes in NENs and MENs proportions.**

**(a, b)** Immunostaining myenteric plexus tissue from juvenile and aging Wnt1-cre:*Rosa26*-tdTomato mice with antibodies against the pan-neuronal marker HuC/D (green).

**(c)** Age-associated loss of NENs and gain of MENs in the small intestinal LM-MP of maturing and aging Wnt1-cre:*Rosa26*-tdTomato mice. Data represent mean ± S.E.M.

**(d)** Mean ± SEM of the four MEN-specific pattern weights in the human mesenchymal cell populations from Elmentaite et al.[59], wherein data from age ranges of 4 – 12 years was clubbed together as Juvenile (Juv), data from age ranges of 20 – 50 years was clubbed together as Adult (Ad), and data from age ranges of 60 – 75 years was clubbed together as Aged (Ag). Every datapoint refers to mean projected pattern weight for cells within a defined age or age range, where Juv data comprise of ages 4, 6, 9, 10, 12 years; Ad data comprise of age ranges 20 - 25, 25 – 30, and 45-50 years; and finally Ag data comprise of age ranges 60 – 65, 65 – 70, and 70 – 75 years. One-way ANOVA ** = $p < 0.01$).

**(e)** Western blot analyses of GDNF (green) and the house-keeping protein β-actin (red) expression in LM-MP tissues from mice of ages P10, P30, and P90. (n = 3 mice per group; each sample is a biological replicate). Fluorescent intensities of the two bands of GDNF (that correspond to ~25 kD bands of protein marker) were quantified together. The lower band of GDNF is present only in the P10 tissues and disappears in P30 and P90 adult murine tissues.

**(f)** Western blot analyses of HGF (green) and the house-keeping protein β-Actin (red) expression in LM-MP tissues from mice of ages P10, P30, and P90. (n = 3 mice per group; each sample is a biological replicate). Fluorescent intensities of the two bands of HGF (that are between 50 kD and 37 kD bands of the protein marker) were quantified together.

**(g)** The normalized fluorescent intensity of GDNF protein to house-keeping protein β-Actin compared between the three age groups. GDNF presence was highest in P10 group and was significantly reduced in P30 and P90 groups. Data represent mean ± S.E.M. One-way ANOVA **** $p < 0.001$.

**(h)** Age-dependent decrease in Gdnf mRNA transcript expression (normalized to the house-keeping gene Hprt) in the myenteric plexuses of P30 and P90 old mice. Data represent mean ± S.E.M. Student"s t-test *$p < 0.05$.

**(i)** The normalized fluorescent intensity of HGF protein to house-keeping protein β-Actin was compared between the three age groups. HGF expression significantly increased from P10 through P90. Data represent mean ± S.E.M. One-way ANOVA *$p < 0.05$.

**(j)** Age-dependent increase in Hgf mRNA transcript expression (normalized to the house-keeping gene Hprt) in the myenteric plexuses of P10, P30, and P90 old mice. Data represent mean ± S.E.M. One-way ANOVA * $p < 0.05$, *** $p < 0.01$, **** $p < 0.001$.

**(k)** Percent proportions of tdTomato$^-$ MENs and mean HuC/D$^+$ neurons/ganglia in LM-MP of cohorts of Wnt1-cre:*Rosa26*-tdTomato mice that were dosed with GDNF or Saline from P10 to P20 age. Data represent mean ± S.E.M. Student"s t-test ** $p < 0.01$.

**(l)** Percent proportions of tdTomato$^-$ MENs and mean HuC/D$^+$ neurons/ganglia in LM-MP of cohorts of Wnt1-cre:*Rosa26*-tdTomato mice that were dosed with HGF or Saline from P10 to P20 age. Data represent mean ± S.E.M. Student"s t-test * $p < 0.05$.

**Fig 9: Reduced RET signaling accelerates ENS aging to cause pathology**

**(a)** HuC/D immunostaining (green) LM-MP tissues from 16-week-old $Ret^{+/CFP}$ ($Ret^{+/-}$) mouse shows mutually exclusive expression of Ret-CFP (cyan, green arrow) and MHCst (red, red arrow) MENs. Nuclei are stained with DAPI (blue). Scale bar = 10 µm.

**(b)** Quantification of Ret-CFP$^+$ neurons from 9- and 16-week-old $Ret^{+/-}$ mice show age-associated loss of Ret-CFP$^+$ neurons. Data represent mean ± S.E.M. Student"s t-test * $p < 0.05$.

**(c)** Quantification of MHCst$^+$ MENs shows significant increase in their proportions in $Ret^{+/-}$ mice but not in $Ret^{+/+}$ mice with age. Data represent mean ± S.E.M. One way ANOVA * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

**(d)** Measures of whole gut transit time (WGTT) in cohorts of $Ret^{+/-}$ and $Ret^{+/+}$ mice MENs show significant slowing of whole gut transit of $Ret^{+/-}$ but not $Ret^{+/+}$ mice with age. Data represent mean ± S.E.M. One way ANOVA * = $p < 0.05$.

**Fig 10: GDNF normalizes altered intestinal motility by increasing NENs proportions in the aging gut.**

**(a)** Measures of whole gut transit time (WGTT) in GDNF (treated with GDNF) and Control (treated with Saline) cohorts of 17-month-old mice taken before the start of treatments and after the end of 10 consecutive days of treatment shows that the two groups are matched in their transit times before treatment, but GDNF treatment significant decreases average transit times when compared to the control cohort. Data represent mean ± S.E.M. Student"s t-test *** = $p <$ 0.001.

**(b)** Quantification of percent MHCst$^{+}$ MENs per HuC/D-labeled neurons in myenteric ganglia in the GDNF and Control cohorts shows significant decrease in their proportions in GDNF-treated cohort but not in saline-treated control cohort. Data represent mean ± S.E.M. Student"s t-test ** $p < 0.01$.

**(c)** Quantification of numbers of RET$^{+}$ NENs per 40X field views of myenteric ganglia shows significant increase in their numbers in GDNF cohort mice when compared with Control cohort mice. Data represent mean ± S.E.M. Student"s t-test * $p < 0.05$.

**Fig 11: Patients with chronic gut dysmotility show significant shifts in their normal pro-portions of the two neuronal lineages.**

**(a)** *ProjectR*-based projection of bulkRNAseq data from intestinal specimens of patients with normal motility (controls) and patients with obstructive defecation (OD), a chronic condition of intestinal dysmotility, into the 50 different NMF patterns learnt earlier, shows that the MEN-specific NMF patterns 32 and 41 were significantly upregulated in bulk-RNAseq of OD patients compared to controls. Data represent mean ± S.E.M. One way ANOVA. Data mined from raw data generated by Kim et al.[75]

**(b)** OD patients show a significant decrease in the expression of important NENs-associated genes such as Ret, Gdnf, Snap-25, Nos1, Klhl1, and Chat, while showing a significant increase in the expression of important MENs-specific genes such as Clic3, Upk3a, Cdh3, Slpi, and Slc17a9. Data taken from Kim et al.[75]

**Suppl. Figure 1: Presence and absence of tdTomato reporter expression in the myenteric**

**ganglia of adult neural crest lineage-traced mice**

**(a)** HuC/D immunostained Wnt1-cre:*Rosa26*-tdTomato LM-MP tissue, where tdTomato-

expression (red) marks neural crest-derived cells, and HuC/D (green) labels myenteric neurons,

we observed the presence of higher intensity fluorescence (red arrows) and lower intensity fluo-

rescence (orange arrows) in various tdTomato$^+$ neurons. The myenteric ganglia also contained

HuC/D$^+$ neurons that did not express tdTomato (green arrow) and these neurons alone were

deemed to be non-neural crest-derived neurons. Scale bar denotes 10 μm.

**(b)** Reporter non-expressing HuC/D-immunolabeled neurons (green, green arrows) can be ob-

served both through the expression of the tdTomato reporter (red), as well as on antibody label-

ing of the tdTomato reporter (gray).

**(c)** Freshly harvested small intestinal tissue from Wnt1-cre:*Rosa26*-tdTomato mice when im-

aged under 20X magnification of a live tissue fluorescence microscope shows the presence of

cells that express tdTomato red and those that don"t.

**(d)** Cells showing hyper-aggregation of the tdTomato reporter can be found in freshly harvested

full-thickness small intestinal myenteric plexus in tissues imaged under 20X magnification of a

live tissue fluorescence microscope.

**(e)** Myenteric ganglia from fixed LM-MP tissue of adult Wnt1-cre:*Rosa26*-tdTomato mice show

show hyper-aggregation of tdTomato (white arrow) when imaged using a confocal microscope.

Nuclei are stained with nuclear marker DAPI (blue). Scale bar denotes 10 μm.

**(f)** Cells showing hyper-aggregation of tdTomato (red, red arrow) show no detectable nuclear

labeling with DAPI (blue) or of pan-neuronal marker HuC/D (green, green arrow). Scale bar de-

notes 10 μm.

In addition, interrogation of two other neural crest-specific lineage-traced mouse lines confirms

our observations with Wnt1-cre:*Rosa26*-tdTomato mice. **(g)** In Wnt1-cre:*Hprt*-tdTomato adult

male mice, tdTomato expression (red) is present only in a subset of HuC/D-immunolabeled

(green) neurons (red arrows) while it is not detected in other HuC/D-expressing neurons

(green). Similarly, **(h)** only a subpopulation of HuC/D[+] adult enteric neurons (green) is derived

from the neural crest, as seen by its co-expression of tdTomato (red, red arrows) in the Pax3-

cre:*Rosa26*-tdTomato mice while other neurons that do not express tdTomato (white arrows)

are inferred to be not derived from the neural crest. Scale bar denotes 10 µm.

**(i)** Orthogonal views of small intestinal villus from a Wnt1-cre:*Rosa26*-tdTomato mouse that

shows presence of tdTomato-expressing (red) neural crest-derived fibers inside the villus but a

complete lack of tdTomato-expression by intestinal epithelial cells. Nuclei are labeled with DAPI

(blue). Scale bar denotes 10 µm.

**(j)** Representational image of the myenteric plexus from a *Rosa26*-tdTomato mouse without any

cre driver showing that in absence of cre recombinase to drive tdTomato expression, there is a

lack of tdTomato expression in this layer. Nuclei are labeled with DAPI (blue). Scale bar denotes

10 µm.

**(k)** Representational image of the myenteric plexus from a *Rosa26*-EGFP mouse without any

cre driver showing that in absence of cre recombinase to drive EGFP expression, there is a lack

of EGFP expression in this layer. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 µm.

1164

**Suppl. Figure 2: Tek-expressing and Mesp1-expressing mesodermal lineage contributes**

**to adult enteric neurons.**

**(a)** Adult small intestinal LM-MP layer from a male Tek-EGFP mouse, where EGFP expression

(green) labels Tek-expressing cells was immunostained with specific antibodies against the

pan-neuronal marker PGP9.5 (red) and imaged under a confocal microscope. The Tek-EGFP

expression did not label any cells within the myenteric plexus marked by PGP9.5 expression

showing that adult small intestinal myenteric neurons did not express Tek. Nuclei are labeled

with DAPI (blue). Scale bar denotes 10 µm.

**(b)** We observed the presence of higher intensity fluorescence (red arrows) and lower intensity

fluorescence (green arrow) in various tdTomato$^+$ cells in a section of heart tissue from a 6-

month-old Mesp1-cre:*Rosa26*-tdTomato adult male mouse, where tdTomato (red) labels

Mesp1-derived cells of the mesodermal lineage that includes cardiomyocytes and vasculature.

Nuclei are stained by DAPI (blue). Scale bar denotes 10 µm.

**(c)** Adult small intestinal LM-MP layer from a 6-month-old male Mesp1-cre:*Rosa26*-tdTomato

mouse, where tdTomato (red) cells are mesoderm-derived cells, was immunostained with spe-

cific antibodies against the pan-neuronal marker HuC/D (green) and imaged under a confocal

microscope. tdTomato-expressing cells outside the ganglia (red arrows) were brighter than

tdTomato$^+$ neurons inside the ganglia (green arrows). Nuclei are labeled with DAPI (blue).

Scale bar denotes 10 µm.

**(d)** Adult small intestinal LM-MP layer from a 2-month old male Mesp1-cre:*Rosa26*-EGFP

mouse, where EGFP cells are mesoderm-derived, was immunostained with specific antibodies

against the reporter EGFP (green) and the pan-neuronal marker HuC/D (red) and imaged under

a confocal microscope. A population of HuC/D-immunolabeled neurons were also labeled with

antibodies against EGFP (green arrows) while other neurons were not (red arrows), suggesting

the derivation of EGFP$^+$ neurons from the Mesp1-expressing mesoderm. **(e)** No primary control

for the Anti-GFP immunostaining showed very weak EGFP expression in the fixed LM-MP tis-

sue from these animals. Nuclei were labeled with DAPI (blue). Scale bar = 10 µm. Consistent

with the observations made with the tdTomato reporter in this transgenic line, the Mesp1-

cre:*Rosa26*-EGFP animals also show a brighter reporter signal in the vasculature and a dimmer

reporter signal in the myenteric neurons.

**(f)** MHCst$^+$ (cyan) labels all tdTomato$^+$ (red) HuC/D$^+$ (green) adult neurons (white arrows) but

not the tdTomato$^-$ neurons (yellow arrow) in the mesoderm-specific Tek-cre:*Hprt*-tdTomato line-

age-traced lines. Nuclei are stained with DAPI (blue). Scale bar = 10 µm.

**Suppl. Figure 3: Validation of MENs-specific markers by immunohistochemistry and confocal microscopy.**

Small intestinal LM-MP from several adult 5-month-old male Wnt1-cre:*Rosa26*-tdTomato mice, where tdTomato-expression (red) labels neural crest-derived cells, was used to ascertain and validate the expression of the novel MENs marker and of known neuronal marker genes.

**(a)** MHCst-specific antibody S46 (green) was found to label tdTomato non-expressing cells within the myenteric ganglia as well as cells outside of the myenteric ganglia. Neural crest-derived cells within the myenteric ganglia expressed tdTomato. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 μm.

The immunostaining of the MENs-markers elucidated through scRNAseq analyses: **(b)** CDH3, **(c)** AEBP1, **(d)** SLPI, **(e)** CFTR, **(f)** NT-3, **(g)** CLIC3, **(h)** FMO2, **(i)** IL-18, **(j)** MYL-7, and **(k)** SLC17A9 (all immunolabeled cells in these panels are green; green arrows) was found to be localized to or enriched in cells without tdTomato expression, while the tdTomato$^+$ cells were found to show not to express these MENs-specific markers (red arrow). In addition, we also immunolabeled the myenteric ganglia from this transgenic mouse line with known neuronal markers (gray) and with antibodies against the pan-neuronal marker HuC/D (green) to show that the neuronal markers Elavl2 or HuB **(l)**, GPR88 **(m)**, HAND2 **(n)**, PDE10A **(o)**, TUBB2B **(p)** and VSNL1 **(q)** were expressed by both tdTomato$^+$ (red arrow) and tdTomato$^-$ HuC/D$^+$ neurons. Orthogonal views of images where 3D image z-stacks were obtained are provided for better visualization of marker expression in the 3D space. Nuclei in these tissues were labeled with DAPI (blue). Scale bar denotes 10 μm.

**Suppl. Figure 4: scRNAseq metrics**

**(a)** MENs are significantly larger in size as compared to the NENs, as the mean Feret"s Diameter of the Wnt1-cre:*Rosa26*-tdTomato⁻ MENs is significantly more than that of the Wnt1-cre:tdTomato⁺ NENs. Data presented as mean ± SEM. Student"s t-test; ** p value < 0.01.

**(b, c)** UMAP representation of sequenced cells generated from 10X Genomics Chromium-derived libraries from the two mouse samples, taken from separate littermate adult male mice, shows similar representation of their cells across the various clusters.

**(d)** Quasi-violin plots of expressed genes provide a comparison of important MEN-specific and neuronal genes between the NENs and MENs clusters in the adult ileal May-Zhang et al[56]. dataset and the NENs, Neuroglial, and MENs clusters in our scRNAseq dataset.

**(e)** Top genes that are pattern drivers for the four MEN-specific NMF patterns along with their gene weights.

**(f)** The four MEN-specific patterns were together used to label a group of cells (True) that together showed increased usage of the MEN-specific patterns, when compared to other cell types in the adult ileal snRNAseq dataset from May-Zhang et al.

**(g)** Top-expressed genes in the group of cells **(f)** that showed higher utilization of the MEN-specific patterns, compared to other cells in the dataset.

**(h)** Representation of the four MEN-specific NMF patterns in the UMAP of our data and the projection of the May-Zhang et al murine ileal dataset and the entire gut mesenchymal single cell RNA seq dataset by Elmentaite et al[59] into the MEN-specific patterns highlights cell clusters that show higher usage of the four NMF patterns. Bottom row shows increasing usage of these patterns with age in the entire mesenchymal gut cell atlas data.

**Suppl. Figure 5: Putative mesoderm-derived enteric neurons (MENs) are present in adult human myenteric ganglia.**

**(a, b)** The mesodermal specific markers, MET and MHCst are expressed by a subpopulation of human enteric neurons. Myenteric ganglia from normal duodenal tissue of 3 different adult human subjects express the pan-neuronal marker HuC/D (green), with distinct subpopulations positive and negative for MET (blue) and MHCst (red). MET and MHCst-expressing neurons (red arrows) in the human ENS are assumed to be mesoderm-derived enteric neurons (MENs) and those not expressing these MENs markers (green arrows) are presumably NENs. Nuclei are labeled with DAPI (gray). Scale bar = 10 μm.

**(c)** Distribution of the various mesenchymal clusters of the post-natal healthy gut cell atlas data (generated by Elmentaite et al.[59]) by donor age. For the age of each donor, the proportion of cells annotated as belonging to each cluster is shown. Proportions of Cluster 1 are observed to expand with age.

1254    **Suppl. Figure 6: Sub-clustering MENs in our dataset.**

1255    **(a)** Sub-clustering the MENs cluster from our scRNAseq dataset into 5 distinct clusters **(b)** was

1256    found to be not based on the expression of cluster-specific marker genes.

1257

**Suppl. Figure 7: HGF and GDNF ratios in maturing and adult murine gut.**

**(a)** Ratio of HGF:GDNF values in fluorescence intensities of their western blots (normalized to their own house-keeping protein β-Actin) from the small intestinal LM-MP of post-natal mice at different ages during their maturation.

**(b)** Ratio of HGF:GDNF expression values from the full thickness small intestinal tissue of adult mice taken from the Tabula muris database[69].

**(c)** Analyses of human plasma proteome from 1,025 individuals was performed by Sathyan et al.[70] and the correlation of various plasma proteomes with aging was calculated. Of the genes that showed significant correlation with age (p < 0.05), we found that the HGF levels in human plasma correlated positively with age (above 0), while the levels of GDNF and RET correlated negatively with age (below 0).

**Suppl. Figure 8: RET-CFP mice and the effect of GDNF treatment on MHCst-expressing**

**MENs and RET-expressing NENs in aging mice**

Representative image of MHCst (red) and HuC/D (green)-immunostained myenteric ganglia of

**(a)** saline-treated control and **(b)**. GDNF-treated 17-month-old mice show MHCst$^+$ (yellow ar-

rows) and MHCst$^-$ neurons. Note the reduction in MHCst-immunostaining on inter-ganglionic

fibers in the 17-month-old mice when treated with GDNF, compared to control mice. Nuclei are

stained with DAPI (blue). Scale bar indicates 10 μm.

Representative image of RET-immunostained (green) myenteric ganglia of **(c)** saline-treated

control and **(d)** GDNF-treated 17-month-old mice. Nuclei are stained with DAPI (blue). Scale bar

indicates 10 μm.

**(e)** Western blot image of a 4-20% gel performed using the RET antibody on total protein lysate

from small intestinal LM-MP layer from post-natal mice shows the expected band between150

and 250 kDa of RET protein.

**Suppl. Figure 9: Projection of the bulk RNA sequencing data of intestinal tissue from Control and Patients with Obstructed Defecation into our murine scRNAseq-derived NMF patterns using *projectR*.**

Transcriptomic data from intestinal tissues of patients with normal intestinal motility and those with obstructed defecation (OD) was procured from GEO (GSE101968). Non-negative matrix factorization (NMF), as implemented in the R package *NNLM* (https://github.com/linxihui/NNLM), was performed on the murine scRNAseq data using k=50 and default parameters; and cell weights for each pattern were grouped. Using projectR, the log2 expression (log2(rpkm + 1)) from the human bulk RNA sequencing data from control and OD patients were projected into the murine scRNAseq-derived NMF patterns. The mean projection weights from Control and OD groups were tested for statistically significant differences using Students" t tests.

**References**

1294

1    Kulkarni, S. *et al.* Advances in Enteric Neurobiology: The "Brain" in the Gut in Health and
Disease. *J Neurosci* **38**, 9346-9354, doi:10.1523/JNEUROSCI.1663-18.2018 (2018).
2    Jarret, A. *et al.* Enteric Nervous System-Derived IL-18 Orchestrates Mucosal Barrier
Immunity. *Cell* **180**, 50-63 e12, doi:10.1016/j.cell.2019.12.016 (2020).
3    Uesaka, T., Young, H. M., Pachnis, V. & Enomoto, H. Development of the intrinsic and
extrinsic innervation of the gut. *Dev Biol* **417**, 158-167, doi:10.1016/j.ydbio.2016.04.016
(2016).
4    Obermayr, F., Hotta, R., Enomoto, H. & Young, H. M. Development and developmental
disorders of the enteric nervous system. *Nat Rev Gastroenterol Hepatol* **10**, 43-57,
doi:10.1038/nrgastro.2012.234 (2013).
5    Hao, M. M. & Young, H. M. Development of enteric neuron diversity. *J Cell Mol Med* **13**,
1193-1210, doi:10.1111/j.1582-4934.2009.00813.x (2009).
6    Anderson, R. B., Newgreen, D. F. & Young, H. M. Neural crest and the development of
the enteric nervous system. *Adv Exp Med Biol* **589**, 181-196, doi:10.1007/978-0-387-
46954-6_11 (2006).
7    Young, H. M. & Newgreen, D. Enteric neural crest-derived cells: origin, identification,
migration, and differentiation. *Anat Rec* **262**, 1-15, doi:10.1002/1097-
0185(20010101)262:1<1::AID-AR1006>3.0.CO;2-2 (2001).
8    Young, H. M., Hearn, C. J. & Newgreen, D. F. Embryology and development of the
enteric nervous system. *Gut* **47 Suppl 4**, iv12-14; discussion iv26,
doi:10.1136/gut.47.suppl_4.iv12 (2000).
9    Bergner, A. J. *et al.* Birthdating of myenteric neuron subtypes in the small intestine of the
mouse. *J Comp Neurol* **522**, 514-527, doi:10.1002/cne.23423 (2014).
10    Uesaka, T., Nagashimada, M. & Enomoto, H. Neuronal Differentiation in Schwann Cell
Lineage Underlies Postnatal Neurogenesis in the Enteric Nervous System. *J Neurosci*
**35**, 9879-9888, doi:10.1523/JNEUROSCI.1239-15.2015 (2015).
11    Espinosa-Medina, I. *et al.* Dual origin of enteric neurons in vagal Schwann cell
precursors and the sympathetic neural crest. *Proc Natl Acad Sci U S A* **114**, 11980-
11985, doi:10.1073/pnas.1710308114 (2017).
12    Burns, A. J. Migration of neural crest-derived enteric nervous system precursor cells to
and within the gastrointestinal tract. *Int J Dev Biol* **49**, 143-150,
doi:10.1387/ijdb.041935ab (2005).
13    Laranjeira, C. *et al.* Glial cells in the mouse enteric nervous system can undergo
neurogenesis in response to injury. *J Clin Invest* **121**, 3412-3424, doi:10.1172/JCI58200
(2011).
14    Brokhman, I. *et al.* Dual embryonic origin of the mammalian enteric nervous system. *Dev*
*Biol* **445**, 256-270, doi:10.1016/j.ydbio.2018.11.014 (2019).
15    Sohal, G. S., Ali, M. M. & Farooqui, F. A. A second source of precursor cells for the
developing enteric nervous system and interstitial cells of Cajal. *Int J Dev Neurosci* **20**,
619-626 (2002).
16    Habeck, M. VENT cells--a load of hot air? *Drug Discov Today* **8**, 332-333,
doi:10.1016/s1359-6446(03)02663-1 (2003).

17    Saffrey, M. J. Cellular changes in the enteric nervous system during ageing. *Dev Biol*
**382**, 344-355, doi:10.1016/j.ydbio.2013.03.015 (2013).

18    Becker, L., Kulkarni, S., Tiwari, G., Micci, M. A. & Pasricha, P. J. Divergent fate and
origin of neurosphere-like bodies from different layers of the gut. *Am J Physiol*
*Gastrointest Liver Physiol* **302**, G958-965, doi:10.1152/ajpgi.00511.2011 (2012).

19    Kulkarni, S. *et al.* Adult enteric nervous system in health is maintained by a dynamic
balance between neuronal apoptosis and neurogenesis. *Proc Natl Acad Sci U S A* **114**,
E3709-E3718, doi:10.1073/pnas.1619406114 (2017).

20    Antal, M., Beneduce, B. M. & Regehr, W. G. The substantia nigra conveys target-
dependent excitatory and inhibitory outputs from the basal ganglia to the thalamus. *J*
*Neurosci* **34**, 8032-8042, doi:10.1523/JNEUROSCI.0236-14.2014 (2014).

21    Freyer, L., Aggarwal, V. & Morrow, B. E. Dual embryonic origin of the mammalian otic
vesicle forming the inner ear. *Development* **138**, 5403-5414, doi:10.1242/dev.069849
(2011).

22    Wright, C. M. *et al.* scRNA-Seq Reveals New Enteric Nervous System Roles for GDNF,
NRTN, and TBX3. *Cell Mol Gastroenterol Hepatol* **11**, 1548-1592 e1541,
doi:10.1016/j.jcmgh.2020.12.014 (2021).

23    Honig, G., Liou, A., Berger, M., German, M. S. & Tecott, L. H. Precise pattern of
recombination in serotonergic and hypothalamic neurons in a Pdx1-cre transgenic
mouse line. *J Biomed Sci* **17**, 82, doi:10.1186/1423-0127-17-82 (2010).

24    Bardot, E. *et al.* Foxa2 identifies a cardiac progenitor population with ventricular
differentiation potential. *Nat Commun* **8**, 14428, doi:10.1038/ncomms14428 (2017).

25    Burtscher, I., Barkey, W., Schwarzfischer, M., Theis, F. J. & Lickert, H. The Sox17-
mCherry fusion mouse line allows visualization of endoderm and vascular endothelial
development. *genesis* **50**, 496-505, doi:10.1002/dvg.20829 (2012).

26    Wilm, B., Ipenberg, A., Hastie, N. D., Burch, J. B. & Bader, D. M. The serosal
mesothelium is a major source of smooth muscle cells of the gut vasculature.
*Development* **132**, 5317-5328, doi:10.1242/dev.02141 (2005).

27    Kawaguchi, M., Bader, D. M. & Wilm, B. Serosal mesothelium retains vasculogenic
potential. *Dev Dyn* **236**, 2973-2979, doi:10.1002/dvdy.21334 (2007).

28    Herrick, S. E. & Mutsaers, S. E. Mesothelial progenitor cells and their potential in tissue
engineering. *Int J Biochem Cell Biol* **36**, 621-642, doi:10.1016/j.biocel.2003.11.002
(2004).

29    Chan, Sunny S.-K. *et al.* Mesp1 Patterns Mesoderm into Cardiac, Hematopoietic, or
Skeletal Myogenic Progenitors in a Context-Dependent Manner. *Cell Stem Cell* **12**, 587-
601, doi:10.1016/j.stem.2013.03.004 (2013).

30    Devine, W. P., Wythe, J. D., George, M., Koshiba-Takeuchi, K. & Bruneau, B. G. Early
patterning and specification of cardiac progenitors in gastrulating mesoderm. *Elife* **3**,
doi:10.7554/eLife.03848 (2014).

31    Lescroart, F. *et al.* Early lineage restriction in temporally distinct populations of Mesp1
progenitors during mammalian heart development. *Nat Cell Biol* **16**, 829-840,
doi:10.1038/ncb3024 (2014).

32    Klotz, L. *et al.* Cardiac lymphatics are heterogeneous in origin and respond to injury.
*Nature* **522**, 62-67, doi:10.1038/nature14483 (2015).

33    Chan, S. S. *et al.* Mesp1 patterns mesoderm into cardiac, hematopoietic, or skeletal
myogenic progenitors in a context-dependent manner. *Cell Stem Cell* **12**, 587-601,
doi:10.1016/j.stem.2013.03.004 (2013).
34    Baup, D., Moser, M., Schurmans, S. & Leo, O. Developmental regulation of the
composite CAG promoter activity in the murine T lymphocyte cell lineage. *Genesis* **47**,
799-804, doi:10.1002/dvg.20569 (2009).
35    Alexopoulou, A. N., Couchman, J. R. & Whiteford, J. R. The CMV early
enhancer/chicken beta actin (CAG) promoter can be used to drive transgene expression
during the differentiation of murine embryonic stem cells into vascular progenitors. *BMC
Cell Biol* **9**, 2, doi:10.1186/1471-2121-9-2 (2008).
36    Hu, X. *et al.* Reprogramming progressive cells display low CAG promoter activity. *Stem
Cells* **39**, 43-54, doi:10.1002/stem.3295 (2021).
37    Akagi, Y. *et al.* Transcriptional activation of a hybrid promoter composed of
cytomegalovirus enhancer and beta-actin/beta-globin gene in glomerular epithelial cells
in vivo. *Kidney Int* **51**, 1265-1269, doi:10.1038/ki.1997.172 (1997).
38    Agah, R. *et al.* Gene recombination in postmitotic cells. Targeted expression of Cre
recombinase provokes cardiac-restricted, site-specific rearrangement in adult ventricular
muscle in vivo. *J Clin Invest* **100**, 169-179, doi:10.1172/JCI119509 (1997).
39    Sokoloff, A. J., Li, H. & Burkholder, T. J. Limited expression of slow tonic myosin heavy
chain in human cranial muscles. *Muscle Nerve* **36**, 183-189, doi:10.1002/mus.20797
(2007).
40    Stockdale, F. E. & Miller, J. B. The cellular basis of myosin heavy chain isoform
expression during development of avian skeletal muscles. *Dev Biol* **123**, 1-9,
doi:10.1016/0012-1606(87)90420-9 (1987).
41    Miller, J. B., Crow, M. T. & Stockdale, F. E. Slow and fast myosin heavy chain content
defines three types of myotubes in early muscle cell cultures. *J Cell Biol* **101**, 1643-1650,
doi:10.1083/jcb.101.5.1643 (1985).
42    Vaughan, S. K., Stanley, O. L. & Valdez, G. Impact of Aging on Proprioceptive Sensory
Neurons and Intrafusal Muscle Fibers in Mice. *J Gerontol A Biol Sci Med Sci* **72**, 771-
779, doi:10.1093/gerona/glw175 (2017).
43    Avetisyan, M. *et al.* Hepatocyte Growth Factor and MET Support Mouse Enteric Nervous
System Development, the Peristaltic Response, and Intestinal Epithelial Proliferation in
Response to Injury. *J Neurosci* **35**, 11543-11558, doi:10.1523/JNEUROSCI.5267-
14.2015 (2015).
44    Rappolee, D. A., Iyer, A. & Patel, Y. Hepatocyte growth factor and its receptor are
expressed in cardiac myocytes during early cardiogenesis. *Circ Res* **78**, 1028-1036,
doi:10.1161/01.res.78.6.1028 (1996).
45    Abraira, V. E. *et al.* The Cellular and Synaptic Architecture of the Mechanosensory
Dorsal Horn. *Cell* **168**, 295-310 e219, doi:10.1016/j.cell.2016.12.010 (2017).
46    Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-
1014 e1022, doi:10.1016/j.cell.2018.06.021 (2018).
47    Drokhlyansky, E. *et al.* The Human and Mouse Enteric Nervous System at Single-Cell
Resolution. *Cell* **182**, 1606-1622, doi:10.1016/j.cell.2020.08.00310.1101/746743 (2020).

48    Heanue, T. A. & Pachnis, V. Expression profiling the developing mammalian enteric
nervous system identifies marker and candidate Hirschsprung disease genes.
*Proceedings of the National Academy of Sciences* **103**, 6919,
doi:10.1073/pnas.0602152103 (2006).
49    D'Autreaux, F., Morikawa, Y., Cserjesi, P. & Gershon, M. D. Hand2 is necessary for
terminal differentiation of enteric neurons from crest-derived precursors but not for their
migration into the gut or for formation of glia. *Development* **134**, 2237-2249,
doi:10.1242/dev.003814 (2007).
50    Strick, C. A. *et al.* Alterations in gene regulation following inhibition of the striatum-
enriched phosphodiesterase, PDE10A. *Neuropharmacology* **58**, 444-451,
doi:10.1016/j.neuropharm.2009.09.008 (2010).
51    Bernstein, H. G. *et al.* Hippocampal expression of the calcium sensor protein visinin-like
protein-1 in schizophrenia. *Neuroreport* **13**, 393-396, doi:10.1097/00001756-200203250-
00006 (2002).
52    Massart, R., Guilloux, J. P., Mignon, V., Sokoloff, P. & Diaz, J. Striatal GPR88
expression is confined to the whole projection neuron population and is regulated by
dopaminergic and glutamatergic afferents. *Eur J Neurosci* **30**, 397-414,
doi:10.1111/j.1460-9568.2009.06842.x (2009).
53    Soo Hoo, L. *et al.* The SNARE Protein Syntaxin 3 Confers Specificity for Polarized
Axonal Trafficking in Neurons. *PLoS One* **11**, e0163671,
doi:10.1371/journal.pone.0163671 (2016).
54    Breuss, M. *et al.* The Expression of Tubb2b Undergoes a Developmental Transition in
Murine Cortical Neurons. *J Comp Neurol* **523**, 2161-2186, doi:10.1002/cne.23836
(2015).
55    De Giorgio, R., Arakawa, J., Wetmore, C. J. & Sternini, C. Neurotrophin-3 and
neurotrophin receptor immunoreactivity in peptidergic enteric neurons. *Peptides* **21**,
1421-1426, doi:10.1016/s0196-9781(00)00286-2 (2000).
56    May-Zhang, A. A. *et al.* Combinatorial Transcriptional Profiling of Mouse and Human
Enteric Neurons Identifies Shared and Disparate Subtypes In Situ. *Gastroenterology*,
doi:10.1053/j.gastro.2020.09.032 (2020).
57    Stein-O'Brien, G. L. *et al.* Decomposing Cell Identity for Transfer Learning across
Cellular Measurements, Platforms, Tissues, and Species. *Cell Syst* **8**, 395-411 e398,
doi:10.1016/j.cels.2019.04.004 (2019).
58    Stein-O'Brien, G. L. *et al.* Enter the Matrix: Factorization Uncovers Knowledge from
Omics. *Trends Genet* **34**, 790-805, doi:10.1016/j.tig.2018.07.003 (2018).
59    Elmentaite, R. *et al.* Cells of the human intestinal tract mapped across space and time.
*Nature* **597**, 250-255, doi:10.1038/s41586-021-03852-1 (2021).
60    Matheis, F. *et al.* Adrenergic Signaling in Muscularis Macrophages Limits Infection-
Induced Neuronal Loss. *Cell* **180**, 64-78 e16, doi:10.1016/j.cell.2019.12.002 (2020).
61    Barrenschee, M. *et al.* SNAP-25 is abundantly expressed in enteric neuronal networks
and upregulated by the neurotrophic factor GDNF. *Histochem Cell Biol* **143**, 611-623,
doi:10.1007/s00418-015-1310-x (2015).

62    Gianino, S., Grider, J. R., Cresswell, J., Enomoto, H. & Heuckeroth, R. O. GDNF
availability determines enteric neuron number by controlling precursor proliferation.
*Development* **130**, 2187-2198, doi:10.1242/dev.00433 (2003).

63    Natarajan, D., Marcos-Gutierrez, C., Pachnis, V. & de Graaff, E. Requirement of
signalling by receptor tyrosine kinase RET for the directed migration of enteric nervous
system progenitor cells during mammalian embryogenesis. *Development* **129**, 5151-
5160 (2002).

64    Taraviras, S. *et al.* Signalling by the RET receptor tyrosine kinase and its role in the
development of the mammalian enteric nervous system. *Development* **126**, 2785-2797
(1999).

65    Rodrigues, D. M., Li, A. Y., Nair, D. G. & Blennerhassett, M. G. Glial cell line-derived
neurotrophic factor is a key neurotrophin in the postnatal enteric nervous system.
*Neurogastroenterol Motil* **23**, e44-56, doi:10.1111/j.1365-2982.2010.01626.x (2011).

66    Amano, O. *et al.* Hepatocyte growth factor is essential for migration of myogenic cells
and promotes their proliferation during the early periods of tongue morphogenesis in
mouse embryos. *Dev Dyn* **223**, 169-179, doi:10.1002/dvdy.1228 (2002).

67    Brun, P. *et al.* Toll-like receptor 2 regulates intestinal inflammation by controlling integrity
of the enteric nervous system. *Gastroenterology* **145**, 1323-1333,
doi:10.1053/j.gastro.2013.08.047 (2013).

68    Jangphattananont, N. *et al.* Distinct Localization of Mature HGF from its Precursor Form
in Developing and Repairing the Stomach. *Int J Mol Sci* **20**, doi:10.3390/ijms20122955
(2019).

69    Almanzar, N. *et al.* A single-cell transcriptomic atlas characterizes ageing tissues in the
mouse. *Nature* **583**, 590-595, doi:10.1038/s41586-020-2496-1 (2020).

70    Sathyan, S. *et al.* Plasma proteomic profile of age, health span, and all-cause mortality
in older adults. *Aging Cell* **n/a**, e13250, doi:10.1111/acel.13250.

71    Wang, H. *et al.* The timing and location of glial cell line-derived neurotrophic factor
expression determine enteric nervous system structure and function. *J Neurosci* **30**,
1523-1538, doi:10.1523/JNEUROSCI.3861-09.2010 (2010).

72    Barrenschee, M. *et al.* No neuronal loss, but alterations of the GDNF system in
asymptomatic diverticulosis. *PLoS One* **12**, e0171416,
doi:10.1371/journal.pone.0171416 (2017).

73    Rossi, V. *et al.* Chronic intestinal pseudo-obstruction in a child harboring a founder
Hirschsprung RET mutation. *Am J Med Genet A* **170**, 2400-2403,
doi:10.1002/ajmg.a.37787 (2016).

74    Uesaka, T., Nagashimada, M., Yonemura, S. & Enomoto, H. Diminished Ret expression
compromises neuronal survival in the colon and causes intestinal aganglionosis in mice.
*J Clin Invest* **118**, 1890-1898, doi:10.1172/JCI34425 (2008).

75    Kim, M. *et al.* Obstructed defecation-an enteric neuropathy? An exploratory study of
patient samples. *Int J Colorectal Dis* **34**, 193-196, doi:10.1007/s00384-018-3160-1
(2019).

76    Sun, T. *et al.* Aging-dependent decrease in the numbers of enteric neurons, interstitial
cells of Cajal and expression of connexin43 in various regions of gastrointestinal tract.
*Aging (Albany NY)* **10**, 3851-3865, doi:10.18632/aging.101677 (2018).

77   Becker, L. *et al.* Age-dependent shift in macrophage polarisation causes inflammation-
mediated degeneration of enteric nervous system. *Gut* **67**, 827-836, doi:10.1136/gutjnl-
2016-312940 (2018).

78   Sharma, G., Colantuoni, C., Goff, L. A., Fertig, E. J. & Stein-O'Brien, G. projectR: an
R/Bioconductor package for transfer learning via PCA, NMF, correlation and clustering.
*Bioinformatics* **36**, 3592-3593, doi:10.1093/bioinformatics/btaa183 (2020).

79   Young, H. M. & Newgreen, D. Enteric neural crest-derived cells: Origin, identification,
migration, and differentiation. *The Anatomical Record* **262**, 1-15, doi:10.1002/1097-
0185(20010101)262:1<1::AID-AR1006>3.0.CO;2-2 (2001).

80   Prummel, K. D., Nieuwenhuize, S. & Mosimann, C. The lateral plate mesoderm.
*Development* **147**, dev175059, doi:10.1242/dev.175059 (2020).

81   Xue, R. *et al.* Expression of Cystic Fibrosis Transmembrane Conductance Regulator in
Ganglia of Human Gastrointestinal Tract. *Sci Rep* **6**, 30926, doi:10.1038/srep30926
(2016).

82   Morarach, K. *et al.* Diversification of molecularly defined myenteric neuron classes
revealed by single cell RNA-sequencing. *bioRxiv*, 2020.2003.2002.955757,
doi:10.1101/2020.03.02.955757 (2020).

83   Frassoni, C. *et al.* Analysis of SNAP-25 immunoreactivity in hippocampal inhibitory
neurons during development in culture and in situ. *Neuroscience* **131**, 813-823,
doi:10.1016/j.neuroscience.2004.11.042 (2005).

84   Wang, W. *et al.* Selective Expression of a SNARE-Cleaving Protease in Peripheral
Sensory Neurons Attenuates Pain-Related Gene Transcription and Neuropeptide
Release. *Int J Mol Sci* **22**, doi:10.3390/ijms22168826 (2021).

85   Tender, G. C., Kaye, A. D., Li, Y. Y. & Cui, J. G. Neurotrophin-3 and tyrosine kinase C
have modulatory effects on neuropathic pain in the rat dorsal root ganglia. *Neurosurgery*
**68**, 1048-1055; discussion 1055, doi:10.1227/NEU.0b013e318208f9c4 (2011).

86   de Jonge, H. R., Ardelean, M. C., Bijvelds, M. J. C. & Vergani, P. Strategies for cystic
fibrosis transmembrane conductance regulator inhibition: from molecular mechanisms to
treatment for secretory diarrhoeas. *FEBS Lett* **594**, 4085-4108, doi:10.1002/1873-
3468.13971 (2020).

87   Chaudhury, A., Dendi, V. S. & Mirza, W. Colligative Property of ATP: Implications for
Enteric Purinergic Neuromuscular Neurotransmission. *Front Physiol* **7**, 500,
doi:10.3389/fphys.2016.00500 (2016).

88   Parathan, P., Wang, Y., Leembruggen, A. J., Bornstein, J. C. & Foong, J. P. The enteric
nervous system undergoes significant chemical and synaptic maturation during
adolescence in mice. *Dev Biol* **458**, 75-87, doi:10.1016/j.ydbio.2019.10.011 (2020).

89   Golden, J. P., DeMaro, J. A., Osborne, P. A., Milbrandt, J. & Johnson, E. M., Jr.
Expression of neurturin, GDNF, and GDNF family-receptor mRNA in the developing and
mature mouse. *Exp Neurol* **158**, 504-528, doi:10.1006/exnr.1999.7127 (1999).

90   Papaconstantinou, J. *et al.* Attenuation of p38alpha MAPK stress response signaling
delays the in vivo aging of skeletal muscle myofibers and progenitor cells. *Aging (Albany
NY)* **7**, 718-733, doi:10.18632/aging.100802 (2015).

91 Soret, R. *et al.* Glial Cell-Derived Neurotrophic Factor Induces Enteric Neurogenesis and
Improves Colon Structure and Function in Mouse Models of Hirschsprung Disease.
*Gastroenterology* **159**, 1824-1838 e1817, doi:10.1053/j.gastro.2020.07.018 (2020).

92 Grider, J. R., Heuckeroth, R. O., Kuemmerle, J. F. & Murthy, K. S. Augmentation of the
ascending component of the peristaltic reflex and substance P release by glial cell line-
derived neurotrophic factor. *Neurogastroenterol Motil* **22**, 779-786, doi:10.1111/j.1365-
2982.2010.01489.x (2010).

93 Li, J., Zhang, Y., Yang, C. & Rong, R. Discrepant mRNA and Protein Expression in
Immune Cells. *Curr Genomics* **21**, 560-563, doi:10.2174/1389202921999200716103758
(2020).

94 Melo, C. G. S. *et al.* Identification of intrinsic primary afferent neurons in mouse jejunum.
*Neurogastroenterol Motil* **32**, e13989, doi:10.1111/nmo.13989 (2020).

95 Saffrey, M. J. Aging of the mammalian gastrointestinal tract: a complex organ system.
*Age (Dordr)* **36**, 9603, doi:10.1007/s11357-013-9603-2 (2014).

96 Cossais, F. *et al.* Altered enteric expression of the homeobox transcription factor Phox2b
in patients with diverticular disease. *United European Gastroenterology Journal* **7**, 349-
357, doi:10.1177/2050640618824913 (2019).

97 Schafmayer, C. *et al.* Genome-wide association analysis of diverticular disease points
towards neuromuscular, connective tissue and epithelial pathomechanisms. *Gut* **68**, 854,
doi:10.1136/gutjnl-2018-317619 (2019).

98 Muller, P. A. *et al.* Microbiota-modulated CART(+) enteric neurons autonomously
regulate blood glucose. *Science* **370**, 314-321, doi:10.1126/science.abd6176 (2020).

99 Becker, L., Kulkarni, S., Tiwari, G., Micci, M.-A. & Pasricha, P. J. Divergent fate and
origin of neurosphere-like bodies from different layers of the gut. *American Journal of
Physiology-Gastrointestinal and Liver Physiology* **302**, G958-G965,
doi:10.1152/ajpgi.00511.2011 (2012).

100 Becker, L., Peterson, J., Kulkarni, S. & Pasricha, P. J. Ex vivo neurogenesis within
enteric ganglia occurs in a PTEN dependent manner. *PLoS One* **8**, e59452 (2013).

101 Becker, L., Peterson, J., Kulkarni, S. & Pasricha, P. J. Ex vivo neurogenesis within
enteric ganglia occurs in a PTEN dependent manner. *PLoS One* **8**, e59452,
doi:10.1371/journal.pone.0059452 (2013).

102 Drokhlyansky, E. *et al.* The Human and Mouse Enteric Nervous System at Single-Cell
Resolution. *Cell* **182**, 1606-1622 e1623, doi:10.1016/j.cell.2020.08.003 (2020).

103 Shenje, L. T. *et al.* Precardiac deletion of Numb and Numblike reveals renewal of
cardiac progenitors. *Elife* **3**, e02164, doi:10.7554/eLife.02164 (2014).

104 Susaki, E. A. *et al.* Whole-brain imaging with single-cell resolution using chemical
cocktails and computational analysis. *Cell* **157**, 726-739, doi:10.1016/j.cell.2014.03.042
(2014).

105 Young, H. M., Furness, J. B., Sewell, P., Burcher, E. F. & Kandiah, C. J. Total numbers
of neurons in myenteric ganglia of the guinea-pig small intestine. *Cell Tissue Res* **272**,
197-200, doi:10.1007/BF00323587 (1993).

106 Zhou, Y. *et al.* Enteric nervous system abnormalities are present in human necrotizing
enterocolitis: potential neurotransplantation therapy. *Stem Cell Res Ther* **4**, 157,
doi:10.1186/scrt387 (2013).

107    Hosie, S. *et al.* Gastrointestinal dysfunction in patients and mice expressing the autism-
1597            associated R451C mutation in neuroligin-3. *Autism Res* **12**, 1043-1056,
1598            doi:10.1002/aur.2127 (2019).
1599    108    Lefevre, C. *et al.* Enteric Nervous System Remodeling in a Rat Model of Spinal Cord
1600            Injury: A Pilot Study. *Neurotrauma Rep* **1**, 125-136, doi:10.1089/neur.2020.0041 (2020).
1601    109    Qiu, X. *et al.* NEDL2 regulates enteric nervous system and kidney development in its
1602            Nedd8 ligase activity-dependent manner. *Oncotarget* **7**, 31440-31453,
1603            doi:10.18632/oncotarget.8951 (2016).
1604    110    Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq
1605            quantification. *Nat Biotechnol* **34**, 525-527, doi:10.1038/nbt.3519 (2016).
1606    111    Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in single-cell
1607            RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nat
1608            Biotechnol* **36**, 421-427, doi:10.1038/nbt.4091 (2018).
1609    112    Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
1610            multidimensional genomic data. *Bioinformatics* **32**, 2847-2849,
1611            doi:10.1093/bioinformatics/btw313 (2016).
1612    113    Lun, A. T. L. *et al.* EmptyDrops: distinguishing cells from empty droplets in droplet-based
1613            single-cell RNA sequencing data. *Genome Biol* **20**, 63, doi:10.1186/s13059-019-1662-y
1614            (2019).
1615    114    Griffiths, J. A., Richard, A. C., Bach, K., Lun, A. T. L. & Marioni, J. C. Detection and
1616            removal of barcode swapping in single-cell RNA-seq data. *Nat Commun* **9**, 2667,
1617            doi:10.1038/s41467-018-05083-x (2018).
1618    115    Lun, A. T., McCarthy, D. J. & Marioni, J. C. A step-by-step workflow for low-level
1619            analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122,
1620            doi:10.12688/f1000research.9501.2 (2016).
1621    116    McCarthy, D. J., Campbell, K. R., Lun, A. T. & Wills, Q. F. Scater: pre-processing, quality
1622            control, normalization and visualization of single-cell RNA-seq data in R. *Bioinformatics*
1623            **33**, 1179-1186, doi:10.1093/bioinformatics/btw777 (2017).
1624    117    Csárdi, G. & Nepusz, T.

1625

**a** Wnt1-cre:*Rosa26*-tdTomato  GFAP  DAPI

**b** Wnt1-cre:*Rosa26*-tdTomato  HuC/D  DAPI

**c** Wnt1-cre:*Rosa26*-tdTomato  HuC/D  DAPI

X-Y  Y-Z  X-Z

**d**

% of all HuC/D⁺ neurons

Wnt1-cre
tdTomato⁺

Wnt1-cre
tdTomato⁻

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

**Figure 10**

**a**

**Cell Pattern 32**
p = 0.05

**Cell Pattern 41**
p = 0.01

Control

OD

**b**

| HGNC Symbol | Direction | Gene type | Description | Coding | Mean RC | log FC | P-value | Adj. P-value |
|---|---|---|---|---|---|---|---|---|
| SNAP25 | Reduced in OD | protein_coding | synaptosomal-associated protein, 25kDa [Source:HGNC Symbol;Acc:11132] | yes | 419 | -3.56 | 2.68E-15 | 1.25E-12 |
| KLHL1 | Reduced in OD | protein_coding | kelch-like family member 1 [Source:HGNC Symbol;Acc:6352] | yes | 5 | -2.92 | 2.43E-06 | 7.07E-05 |
| NOS1 | Reduced in OD | protein_coding | nitric oxide synthase 1 (neuronal) [Source:HGNC Symbol;Acc:7872] | yes | 775 | -2.91 | 3.29E-06 | 9.06E-05 |
| CHAT | Reduced in OD | protein_coding | choline O-acetyltransferase [Source:HGNC Symbol;Acc:1912] | yes | 17 | -2.78 | 2.00E-07 | 9.34E-06 |
| GDNF | Reduced in OD | protein_coding | glial cell derived neurotrophic factor [Source:HGNC Symbol;Acc:4232] | yes | 649 | -2.75 | 2.27E-10 | 2.98E-08 |
| RET | Reduced in OD | protein_coding | ret proto-oncogene [Source:HGNC Symbol;Acc:9967] | yes | 474 | -2.59 | 6.38E-18 | 5.29E-15 |
| CLIC3 | Increased in OD | protein_coding | chloride intracellular channel 3 [Source:HGNC Symbol;Acc:2064] | yes | 98 | 2.95 | 1.74E-12 | 4.54E-10 |
| UPK3A | Increased in OD | protein_coding | uroplakin 3A [Source:HGNC Symbol;Acc:12580] | yes | 49 | 2.93 | 1.47E-06 | 4.68E-05 |
| CDH3 | Increased in OD | protein_coding | cadherin 3, type 1, P-cadherin (placental) [Source:HGNC Symbol;Acc:1762] | yes | 252 | 2.66 | 9.87E-06 | 2.20E-04 |
| SLPI | Increased in OD | protein_coding | secretory leukocyte peptidase inhibitor [Source:HGNC Symbol;Acc:11092] | yes | 1,924 | 2.56 | 9.84E-06 | 2.20E-04 |
| SLC17A9 | Increased in OD | protein_coding | solute carrier family 17 (Vesicular nucleotide transporter). [Source: HGNC] | yes | 1,997 | 2.15 | 9.67E-06 | 2.17E-07 |

Figure 11

**Table 1:** Top expressed genes by the Calcb-expressing clusters and the putative MENs cluster in data from May-Zhang et al, along with top-expressed genes in the MENs cluster in our data. Highlighted genes between MENs clusters from our data and data from May-Zhang et al show similar gene expression profiles between the two clusters.

| Calcb+ cluster corresponding to Ileum cluster in May-Zhang et al. which expresses Calcb, Calb2, Nmu, Pcdh10, Cysltr2 | | | Calcb+ cluster corresponding to Ileum cluster in May-Zhang et al. which expresses Calcb, Calb2, Sst | | | MENs-like cluster corresponding to Ileum cluster in May-Zhang et al. which expresses Upk3b and Igfbp6 | | | MENs Cluster (Our Data) | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Gene | FDR | summary.logFC | Gene | FDR | summary.logFC | Gene | FDR | summary.logFC | Gene | FDR | summary.logFC |
| Zfp804a | 0 | 5.314372906 | Nlgn1 | 1.56E-227 | 4.503778932 | Abi1 | 4.98E-36 | 3.316005072 | Bcam | 0 | 1.310889756 |
| Pcdh10 | 0 | 2.727131014 | Plxna4 | 5.40E-170 | 3.513261638 | Dcn | 2.08E-52 | 5.091738535 | Mgst1 | 0 | 1.466554292 |
| Gpr149 | 0 | 2.928565459 | Pcdh15 | 6.75E-269 | 4.552606891 | Col3a1 | 7.95E-31 | 3.039859427 | Krt18 | 1.02E-252 | 0.722876212 |
| Necab1 | 0 | 3.126115371 | Nrg3 | 9.23E-246 | 4.051012028 | C3 | 1.11E-38 | 4.827731701 | Igfbp6 | 0 | 4.277079428 |
| Grin3a | 0 | 2.477871618 | Eml6 | 1.02E-166 | 3.739252675 | Slpi | 2.75E-31 | 4.598725251 | Rspo1 | 0 | 1.356888212 |
| Fam19a1 | 0 | 4.658451729 | Ryr2 | 4.15E-122 | 2.724692337 | Lrrn4 | 7.45E-21 | 2.797813295 | Gpm6a | 0 | 2.007236002 |
| Ano2 | 0 | 3.443066822 | Nrxn3 | 1.20E-156 | 4.845162261 | S100a6 | 9.36E-29 | 3.230324987 | Upk3b | 0 | 2.136663175 |
| Ntrk3 | 0 | 4.380099903 | Rbfox1 | 0.00E+00 | 4.833341666 | Upk3b | 1.90E-26 | 3.101085936 | Aebp1 | 0 | 2.502295384 |
| Cntn5 | 0 | 6.308342607 | Sst | 6.45E-135 | 5.862426929 | Gpm6a | 1.13E-24 | 3.730157809 | Dcn | 0 | 4.082052904 |
| Ccbe1 | 0 | 3.076522552 | Chsy3 | 4.53E-122 | 3.602794203 | Sparc | 1.37E-28 | 3.316214748 | Slpi | 0 | 4.928306229 |
| Zbtb7c | 0 | 2.656109043 | Gria2 | 3.32E-126 | 3.083755369 | Meg3 | 3.89E-23 | 2.952545493 | Rarres2 | 0 | 4.202866323 |
| Astn2 | 0 | 3.366649926 | Ptprd | 3.17E-139 | 2.906173926 | Igfbp6 | 4.99E-26 | 3.61531349 | Efemp1 | 0 | 1.672848073 |
| Dapk2 | 0 | 1.703870376 | Pde4b | 3.69E-210 | 3.947198917 | Agap1 | 4.03E-21 | 2.507795342 | Aldh1a2 | 4.60E-277 | 0.796263425 |
| Nrxn3 | 0 | 5.160806272 | Fam19a2 | 8.41E-100 | 3.227261386 | Pbx1 | 4.47E-25 | 2.530783608 | Fmo2 | 0 | 1.582315372 |
| Dgkg | 0 | 4.15397884 | Mdga2 | 2.01E-132 | 3.667573403 | Efemp1 | 2.67E-22 | 2.915648257 | Upk1b | 0 | 1.081912196 |
| Galr1 | 0 | 1.673088209 | Snhg11 | 6.24E-112 | 4.344115315 | Gas6 | 2.54E-20 | 2.742900314 | Gas1 | 0 | 1.464482623 |
| Nmu | 0 | 3.521779106 | Ralyl | 1.59E-128 | 3.809356707 | Muc16 | 2.43E-21 | 3.168081514 | Krt7 | 0 | 1.315380677 |
| Hspb8 | 1.42E-236 | 1.244433448 | Bnc2 | 1.58E-137 | 3.131344621 | Gas1 | 2.13E-23 | 2.708510446 | Crip1 | 0 | 4.56883561 |
| Cpne4 | 0 | 4.163433255 | Robo1 | 8.43E-84 | 2.405353602 | Dpp4 | 4.03E-21 | 2.876742349 | Clu | 0 | 1.502811937 |
| Tmeff2 | 0 | 3.49539549 | Nrxn1 | 1.51E-150 | 4.662080188 | Cobll1 | 1.10E-20 | 2.231586143 | C3 | 0 | 3.494711523 |
| Myl1 | 0 | 3.653772641 | Pcsk2 | 5.79E-169 | 3.258681935 | Rarres2 | 1.64E-20 | 3.226437143 | Dpp4 | 1.12E-210 | 0.556370013 |
| Calcb | 0 | 3.200679451 | Syn2 | 1.17E-132 | 3.619085064 | Myo1d | 6.05E-21 | 2.158327438 | Lrrn4 | 1.58E-271 | 0.736486954 |
| Cysltr2 | 0 | 2.487957211 | Ntrk3 | 1.05E-108 | 2.657539356 | Tbce | 1.69E-12 | 3.139158995 | Nkain4 | 0 | 1.298360281 |
| Clstn2 | 0 | 2.655215539 | Scube1 | 1.03E-95 | 2.265338351 | Nfib | 7.62E-23 | 2.57202827 | S100a6 | 0 | 4.128650316 |
| Robo1 | 1.98E-297 | 1.9960685 | Man2a1 | 9.20E-139 | 3.375536511 | Crip1 | 7.17E-19 | 2.764477578 | Krt19 | 0 | 1.900927424 |
| Hunk | 0 | 1.918886092 | Kcnn2 | 7.39E-85 | 2.418320757 | Sema5a | 1.97E-19 | 2.852546212 | Cfb | 8.74E-203 | 0.608707666 |
| Snhg11 | 0 | 2.098574269 | Fam19a1 | 1.79E-147 | 3.622699285 | Fmo2 | 1.80E-18 | 3.253371192 | Gabarapl1 | 5.26E-250 | 0.709728291 |
| Aff2 | 0 | 2.087576143 | Nell1 | 3.21E-100 | 2.793128177 | Ezr | 2.84E-20 | 2.166036039 | Lgals3 | 0 | 1.26072921 |
| Otud7a | 3.38E-263 | 2.545208984 | Galntl6 | 1.72E-109 | 3.815091788 | Ahnak | 6.05E-21 | 2.714253195 | Krt8 | 1.39E-245 | 0.689105423 |
| Pcdh9 | 0 | 4.216229313 | Sgcd | 5.24E-120 | 2.88253651 | Bnc2 | 6.54E-20 | 3.061680372 | Tm4sf1 | 0 | 1.367716307 |
| Htr3b | 0 | 1.553388667 | Pde4d | 1.18E-97 | 2.285404689 | Krt19 | 6.15E-20 | 2.380492921 | Slc39a8 | 7.17E-197 | 0.489802511 |
| Scn11a | 0 | 1.908976823 | Slc8a1 | 1.08E-180 | 3.289543486 | Crim1 | 3.00E-20 | 2.434254471 | Sema3c | 2.63E-255 | 0.71554962 |
| Iqgap2 | 0 | 1.75599309 | Tcf4 | 1.52E-159 | 2.592805317 | Plxna4 | 2.07E-19 | 2.707560817 | Pcolce | 0 | 1.823133603 |
| Rbfox1 | 0 | 4.70241843 | Pak3 | 2.89E-91 | 2.580345128 | Sox6 | 3.93E-19 | 2.803585619 | Gsta4 | 1.09E-226 | 0.62733526 |
| Tcf7l2 | 0 | 2.84136652 | Kcnma1 | 1.27E-136 | 2.945051283 | Aebp1 | 1.57E-19 | 2.548178022 | Ezr | 0 | 1.019712533 |
| Tbx2 | 0 | 1.871954414 | Adgrb3 | 5.63E-143 | 4.396153392 | Cfh | 9.08E-18 | 2.422456138 | Cldn15 | 4.76E-291 | 0.798661223 |
| Robo2 | 0 | 4.088780154 | Unc5c | 2.09E-114 | 3.291001111 | Col1a1 | 2.72E-13 | 2.033502612 | Serpinb6b | 2.66E-212 | 0.562587863 |
| Kcnt2 | 0 | 3.926721004 | Cpne4 | 4.42E-81 | 2.730862875 | Wt1 | 2.03E-17 | 2.196826568 | Nbl1 | 0 | 1.239754358 |
| Srrm4 | 2.23E-261 | 2.303709088 | Meg3 | 7.04E-124 | 4.308953415 | Ccdc171 | 1.09E-14 | 2.022034827 | Slc16a1 | 2.28E-244 | 0.65901005 |
| Plxna4 | 0 | 3.31892195 | Snap25 | 4.22E-135 | 3.917101813 | Cd2ap | 2.30E-18 | 2.290803108 | Smim1 | 2.76E-245 | 0.659125894 |
| Sgcz | 0 | 4.739595343 | Syt1 | 7.05E-140 | 4.346683131 | Arhgap29 | 3.72E-18 | 2.409211927 | Snhg18 | 3.11E-181 | 0.475288967 |
| 9530059O14 | 0 | 3.600146025 | Grik4 | 8.53E-115 | 2.35016852 | Ano1 | 2.38E-18 | 3.039709984 | Cavin3 | 6.15E-293 | 0.855731534 |
| Sox4 | 7.89E-259 | 1.211156367 | Ptprn2 | 1.75E-115 | 3.282060606 | Col1a2 | 5.02E-13 | 2.132928671 | Gpc3 | 0 | 1.393181642 |
| Hcn1 | 8.400165280 | 1.861283774 | Arhgap26 | 1.46E-133 | 2.381300441 | Cav1 | 5.10E-18 | 2.235452113 | Serpinb1a | 3.54E-178 | 0.472277374 |
| Scube1 | 0 | 2.077291022 | Bcl2 | 1.53E-91 | 2.27434062 | Bicc1 | 2.27E-19 | 2.486459956 | Chst4 | 3.42E-232 | 0.599828027 |
| Boc | 5.82E-199 | 0.854637195 | Ppm1h | 6.46E-139 | 3.466729684 | Msln | 2.26E-14 | 1.829701191 | Ptgis | 5.59E-231 | 0.598750529 |
| Cbln2 | 0 | 2.229769856 | Fgf14 | 3.30E-126 | 4.330212767 | Dlc1 | 1.57E-18 | 2.561801087 | Slc9a3r1 | 7.03E-265 | 0.883988091 |
| Kcnn3 | 0 | 1.899022238 | Gm12216 | 9.54E-78 | 1.937528878 | Upk1b | 1.96E-17 | 1.970013811 | Ecm1 | 1.88E-172 | 0.514360839 |
| Rph3a | 0 | 1.741082515 | Camk2a | 1.73E-88 | 2.352765766 | Sulf1 | 3.39E-17 | 2.542517734 | Igfbp7 | 0 | 2.347561986 |
| Cpne8 | 0 | 2.360533782 | Fgf13 | 5.32E-123 | 3.932808506 | Vim | 1.68E-16 | 2.45290991 | Msln | 0 | 2.266588157 |
| Calcrl | 1.12695240 | 1.788279932 | Ppp3ca | 1.96E-101 | 2.03501536 | Sema3c | 1.17E-17 | 2.547300887 | Gpx3 | 0 | 2.209730215 |
| Lingo2 | 0 | 4.714175039 | Adamts9 | 1.24E-60 | 1.993937158 | Ifitm3 | 6.57E-15 | 2.071790767 | Col14a1 | 1.17E-163 | 0.423875564 |
| Slc4a4 | 0 | 2.575716085 | Calcb | 4.08E-76 | 2.288867891 | Rspo1 | 1.96E-17 | 1.964755911 | Anxa1 | 4.24E-246 | 0.691007825 |
| Cux2 | 0 | 2.614397251 | Pcbp3 | 1.17E-136 | 3.893280923 | Aldh1a2 | 4.16E-15 | 2.160694165 | Bace2 | 7.94E-223 | 0.603024779 |
| Chat | 1.62E-252 | 1.075493057 | Dlgap2 | 1.21E-96 | 2.470978745 | Serping1 | 4.91E-14 | 2.101997058 | Cfh | 0 | 1.190621234 |
| Htr3a | 0 | 1.82434275 | Nkain2 | 1.01E-107 | 2.761261609 | Timp2 | 8.53E-10 | 1.596556868 | Tmem176a | 1.85E-170 | 0.915766916 |
| Hs3st2 | 1.18369186 | 1.564650584 | Chat | 2.41E-66 | 1.619335139 | Sdc4 | 9.47E-13 | 1.640281568 | Tmsb10 | 0 | 2.299200297 |
| Cdh8 | 0.00E+00 | 2.544193879 | Gfra2 | 4.04E-106 | 2.379843878 | Cd34 | 2.04E-10 | 1.324668078 | Timp2 | 0 | 1.918063641 |
| Gm45881 | 0 | 2.045509731 | Pde10a | 7.37E-115 | 2.924056301 | Serpinh1 | 2.72E-11 | 1.617681506 | Pdpn | 2.36E-252 | 0.846386263 |
| Cdh6 | 0 | 2.996825969 | Ptprr | 1.01E-130 | 3.346248491 | Heg1 | 1.56E-17 | 1.795303412 | Prss23 | 0 | 0.980677037 |
| Gm21847 | 9.83E-301 | 2.668713923 | Plod2 | 7.61E-111 | 2.338515945 | Arhgap18 | 1.69E-15 | 2.494302228 | Cd151 | 2.85E-279 | 0.795276372 |
| Frmd4b | 0 | 2.260822381 | Hs6st3 | 2.71E-78 | 3.027568338 | Igfbp7 | 3.27E-15 | 2.01497489 | P3h4 | 8.94E-212 | 0.512747523 |
| Ndrg4 | 5.28E-172 | 1.033574626 | Rbms3 | 2.24E-117 | 1.826125797 | Anxa2 | 1.55E-16 | 1.783316817 | Cd200 | 0 | 0.975177086 |
| Efr3a | 0 | 2.922824682 | Ccbe1 | 2.81E-69 | 1.972703173 | Celf2 | 8.01E-17 | 2.016354207 | Ltbp4 | 1.80E-268 | 0.859534481 |
| Syt2 | 0 | 1.996197662 | Trpm3 | 3.63E-112 | 3.250658339 | Slc4a4 | 2.64E-14 | 2.058956296 | Gstt1 | 2.26E-142 | 0.347320974 |
| Pak7 | 2.65E-280 | 1.50563649 | Frmd4a | 4.22E-106 | 3.509014084 | Mast4 | 7.11E-16 | 2.591248418 | Ssbp3 | 1.429035064 | 0.888557965 |
| Lhfpl2 | 0 | 1.681014338 | Galnt17 | 5.29E-90 | 2.14675017 | Bcam | 3.45E-09 | 1.453472527 | Tmem98 | 3.90E-240 | 0.62900078 |
| Adgrg6 | 0 | 3.290333725 | Itpr1 | 1.09E-85 | 2.066072139 | Cobl | 3.19E-14 | 2.113252622 | Spint2 | 1.42E-186 | 0.478143896 |
| Spock3 | 0 | 3.61350554 | Srrm4 | 1.17E-82 | 2.302220849 | Airn | 1.00E-16 | 2.159009448 | Nupr1 | 3.42E-226 | 0.705142527 |
| Smad6 | 3.00E-194 | 0.829179685 | Slc7a14 | 2.27E-93 | 2.415887276 | Rbbp8 | 1.25E-07 | 1.460802144 | Basp1 | 6.06E-239 | 0.625848924 |
| Zeb2 | 1.58E-304 | 2.305509194 | Sema6d | 4.44E-99 | 2.246929266 | Piezo2 | 6.47E-15 | 2.155464659 | C1ra | 2.06E-263 | 0.752700756 |
| Hs3st5 | 0 | 2.356357004 | Pcdh7 | 8.46E-105 | 3.303407952 | Wls | 8.02E-16 | 1.68899014 | Serpinh1 | 4.94065645 | 1.065045464 |
| Kcnh1 | 1.71E-232 | 1.145753317 | Nwd2 | 6.48E-47 | 1.597139988 | Pcolce | 2.35E-09 | 1.299424275 | Adamts2 | 9.08E-129 | 0.322228993 |
| Scn9a | 0 | 1.985971133 | Slit3 | 2.09E-90 | 2.828190275 | Prkg1 | 6.92E-15 | 2.705845188 | Dmkn | 3.43E-187 | 0.652807485 |
| Ppp3ca | 0 | 2.635476275 | Asic2 | 3.26E-63 | 3.678490279 | Syne2 | 1.79E-15 | 2.085470282 | C1s1 | 2.25E-276 | 0.820741368 |
| Agrn | 0 | 2.149338255 | Stxbp5l | 1.22E-98 | 2.855884955 | Zfpm2 | 7.87E-15 | 2.689571135 | Sparc | 0 | 1.954167099 |
| Cntnap2 | 0 | 3.976062671 | Smarca2 | 2.65E-97 | 2.286413106 | Adamtsl1 | 1.19E-15 | 2.937092165 | Espn | 3.75E-166 | 0.392816171 |
| Ngfr | 0 | 2.251817207 | Ptprt | 3.14E-93 | 2.816471863 | Gm26809 | 0.0001518 | 1.800107543 | Anxa2 | 0 | 1.579365041 |
| Tnr | 0 | 2.255094218 | Dpyd | 1.72E-91 | 3.059193551 | Chst4 | 1.09E-13 | 1.534679673 | Krt14 | 6.29E-117 | 0.368125408 |
| Mcf2l | 6.65E-174 | 0.813444301 | Tln2 | 1.49E-97 | 2.183191767 | Ctdspl | 1.02E-13 | 1.587097481 | Sdc4 | 0 | 1.913582811 |
| Kcnb2 | 0 | 4.170860912 | Satb2 | 1.74E-60 | 2.016803391 | Ptgis | 1.96E-10 | 1.436656186 | Serping1 | 0 | 1.802367854 |
| Gpr85 | 0 | 1.49922453 | Tshz2 | 5.54E-95 | 2.764995869 | Kcnab1 | 4.33E-13 | 2.347466136 | Pmp22 | 5.43E-232 | 0.6401978 |
| Nell1 | 0 | 2.464621478 | Plcxd3 | 4.17E-89 | 2.590078205 | Nav2 | 2.26E-14 | 2.852022736 | Cplx2 | 1.77E-251 | 0.917909984 |
| Arhgef3 | 0 | 1.961289599 | Tcf7l2 | 1.76E-95 | 2.141154837 | C1ra | 6.88E-07 | 1.318506858 | Il18 | 6.70E-154 | 0.396176068 |
| Syt15 | 3.73E-275 | 1.216092892 | Caln1 | 9.30E-64 | 2.04008457 | Wnt5a | 4.94E-12 | 1.540387353 | Il6st | 1.73E-262 | 0.71225818 |
| Lsamp | 7.25E-253 | 2.548136091 | Dgki | 6.15E-121 | 3.238923425 | Dab2 | 2.15E-13 | 1.86057873 | Meg3 | 1.02E-173 | 0.558817278 |
| 5330434G04 | 4.84E-185 | 2.358912068 | Syt17 | 5.83E-76 | 1.809205079 | Ankrd12 | 0.0001106 | 1.841901593 | Cald1 | 2.88E-262 | 0.803200156 |
| Ptger3 | 1.23E-256 | 1.370213397 | 9530059O14 | 5.05E-74 | 2.430525524 | Cavin2 | 6.09E-12 | 1.976778556 | Mt2 | 1.37E-218 | 0.736087263 |
| 1700111E14 | 4.26E-258 | 1.561674061 | Unc79 | 7.38E-102 | 2.530692527 | Rnase4 | 1.39E-11 | 1.688218465 | Nenf | 7.12E-165 | 0.869917467 |
| Mgat4c | 0 | 3.086139006 | Pvt1 | 8.90E-90 | 2.163637608 | Meis1 | 1.26E-13 | 1.908870378 | Col6a1 | 2.61E-252 | 0.693856752 |
| Fgf14 | 0 | 3.456636428 | Mirg | 4.73E-63 | 1.711809423 | Anxa1 | 1.86E-12 | 1.531275304 | Rnase4 | 0.00E+00 | 1.162670662 |
| Mafg | 2.98E-74 | 1.352690002 | Ret | 1.04E-68 | 1.699484732 | Ildr2 | 9.47E-13 | 1.550620875 | Vim | 0 | 1.805718628 |
| Kif26b | 0 | 2.174461286 | Nrg1 | 5.72E-62 | 3.678288908 | Slc9a3r1 | 1.86E-10 | 1.325162783 | F11r | 6.92E-219 | 0.535570277 |
| Plppr5 | 1.60E-265 | 1.61715395 | Myt1l | 1.56E-120 | 3.008757293 | 9530026P05 | 2.17E-13 | 2.515612487 | Fam180a | 2.29E-131 | 0.294690768 |
| Srrm3 | 5.65E-165 | 1.711562875 | Mapk10 | 9.27E-117 | 2.871225578 | Man1a | 3.13E-12 | 1.539400492 | Npdc1 | 2.34E-164 | 0.415479342 |
| Hmbox1 | 7.78E-79 | 0.6148506 | Gm15398 | 4.78E-73 | 2.683512094 | Gpc3 | 1.20E-12 | 1.790707073 | Cd2ap | 2.16E-225 | 0.595177252 |
| Rbfox3 | 1.95E-259 | 1.230611205 | Arhgef28 | 5.64E-99 | 2.170007537 | Col14a1 | 3.60E-11 | 2.039467045 | Pamr1 | 3.15E-177 | 0.443000918 |
| Ncam2 | 0 | 2.906633451 | Zbtb20 | 6.81E-94 | 1.566902687 | | | | Stk26 | 4.71E-112 | 0.258746973 |
| Slc8a1 | 0 | 2.937485628 | Dscam | 2.70E-101 | 2.808126988 | | | | | | |