Age-associated changes in lineage composition of the enteric nervous system regulate gut health and disease

Subhash Kulkarni[1#], Monalee Saha[1], Jared Slosberg[2], Alpana Singh[1], Sushma Nagaraj[1], Laren Becker[3], Chengxiu Zhang[1], Alicia Bukowski[1], Zhuolun Wang[1], Guosheng Liu[1], Jenna Leser[1,] Mithra Kumar[1], Shriya Bakhshi[1], Matthew Anderson[4], Mark Lewandoski[4], Elizabeth Vincent[3], Loyal A. Goff [2, 5, 6], Pankaj Jay Pasricha[1]

[1]Center for Neurogastroenterology, Department of Medicine, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[2]Department of Genetic Medicine, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[3]Division of Gastroenterology, Stanford University – School of Medicine, Stanford CA 94301

[4]Center for Cancer Research, National Cancer Institute, Frederick, MD 21702

[5]Department of Neuroscience, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[6]Kavli Neurodiscovery Institute, Johns Hopkins University – School of Medicine, Baltimore, MD 21205

[#]Address all correspondence to:

Subhash Kulkarni, Ph.D.

Assistant Professor

Department of Medicine

skulkar9@jh.edu

Abstract

The enteric nervous system (ENS), a collection of neurons contained in the wall of the gut, is of fundamental importance to gastrointestinal and systemic health. According to the prevailing paradigm, the ENS arises from progenitor cells migrating from the neural crest and remains largely unchanged thereafter. Here, we show that the lineage composition of maturing ENS changes with time, with a decline in the canonical lineage of neural-crest derived neurons and their replacement by a newly identified lineage of mesoderm-derived neurons. Single cell transcriptomics and immunochemical approaches establish a distinct expression profile of mesoderm-derived neurons. The dynamic balance between the proportions of neurons from these two different lineages in the post-natal gut is dependent on the availability of their respective trophic signals, GDNF-RET and HGF-MET. With increasing age, the mesoderm-derived neurons become the dominant form of neurons in the ENS, a change associated with significant functional effects on intestinal motility which can be reversed by GDNF supplementation. Transcriptomic analyses of human gut tissues show reduced GDNF-RET signaling in patients with intestinal dysmotility which is associated with reduction in neural crest-derived neuronal markers and concomitant increase in transcriptional patterns specific to mesoderm-derived neurons. Normal intestinal function in the adult gastrointestinal tract therefore appears to require an optimal balance between these two distinct lineages within the ENS.

Introduction

The enteric nervous system (ENS) is a large collection of neurons, glial, and precursor cells that resides within the gastrointestinal wall and regulates gut motility and secretion along with modulating epithelial and immune cell function[1,2]. During fetal development, the mammalian ENS is populated by neurons and glia derived from neural crest (NC)-derived precursors[3-9]. These precursors follow diverse migratory routes to colonize and innervate various parts of the gut before birth[10-12]. It is not clear, however, that this lineage persists in its entirety in the adult gut, as indicated by the observed lack of expression of fluorescent reporter protein in a subpopulation of adult enteric neurons in NC-lineage-traced mice[13,14]. Alternative sources of enteric neurons that have been proposed in the literature include the ventral neural tube (VENT)[15], or the Pdx1-expressing pancreatic endoderm[14], but the interpretation of these studies has been limited by the lack of robust lineage markers for non-NC derived neurons[16]. In addition, while prior studies have documented cellular changes to the ageing ENS[17], the developmental mechanisms behind these changes are unknown. Thus, confirmation of a second, distinct lineage of enteric neurons in adults is important for our understanding of the healthy post-natal development and aging of the ENS, as well as for the pathogenesis of acquired disorders of the ENS.

In this study, we found that while the early post-natal ENS is derived from the canonical NC-lineage, this pattern changes rapidly as the ENS matures, due to the arrival and continual expansion of a novel population of <u>M</u>esoderm-derived <u>E</u>nteric <u>N</u>euron<u>s</u> (MENs) which represent an equal proportion of the ENS in young adulthood and with increasing age, eventually outnumber the <u>N</u>C-derived <u>E</u>nteric <u>N</u>euron<u>s</u> (NENs). We also found that, while the NEN population is regulated by glial derived neurotrophic factor (GDNF) signaling through its receptor RET, the MEN population is regulated by hepatocyte growth factor (HGF) signaling. Increasing HGF levels during maturation or by pharmacological dosing increase proportions of MENs. Similarly, decrease in

GDNF with age decrease NENs; and increasing GDNF levels by pharmacological dosing increase

NENs proportions in the adult ENS to impact intestinal motility.

These results indicate for the first time that the mesoderm is an important source of neurons in

the second largest nervous system of the body. The increasing proportion of neurons of meso-

dermal lineage is a natural consequence of maturation and aging; further, this lineage can be

expected to have vulnerabilities to disease that are distinct from those affecting the NEN popula-

tion. These findings therefore provide a new paradigm for understanding the structure and func-

tion of the adult and aging ENS in health, age-related gut dysfunction and other acquired disorders

of gastrointestinal motility.

Results

Only half of all mid-age adult enteric neurons are derived from the neural crest. We ana-

lyzed small intestinal longitudinal muscle-myenteric plexus (LM-MP) from adult (post-natal day

60; P60) Wnt1-cre:*Rosa26*-tdTomato mice, in which tdTomato is expressed by all derivatives of

Wnt1[+] NC-cells[18]. In these tissues, while GFAP, a glial marker, was always co-expressed with

tdTomato (Fig 1a), tdTomato-expression was absent in many myenteric neurons (Fig 1b, c). By

careful enumeration, almost half of all myenteric neurons expressing the pan-neuronal marker Hu

were found to not express the reporter (percent tdTomato[-] neurons: 44.27 ± 2.404 SEM; enumer-

ated from 2216 neurons from 6 mice, Fig 1d). In these lineage-traced mice, myenteric ganglia

were found to contain tdTomato*high* and tdTomato*low* neurons and due care was taken to image

subsets (Suppl. Fig 1a). Both tdTomato*high* and tdTomato*low* neurons were classified as tdTomato[+]

and only neurons that did not show any tdTomato-expression were classified as tdTomato[-] neu-

rons. Immunofluorescence staining of small intestinal LM-MP tissue from adult Wnt1-cre:*Rosa26*-

tdTomato mice using antibodies against tdTomato protein showed a consistent lack of reporter

expression in a population of myenteric neurons (Suppl. Fig 1b). This lack of reporter expression

is not a function of tissue peeling or fixation, as the tdTomato⁻ cells within the myenteric ganglia

were also observed in freshly harvested unpeeled gut tissue, when imaged using a live tissue

fluorescence microscope (Suppl. Fig 1c). In addition, we noticed that myenteric ganglia of the

Wnt1-cre:*Rosa26*-tdTomato transgenic mouse show similar tdTomato aggregation in cells of the

myenteric plexus from both freshly harvested and immediately fixed tissue (Suppl. Fig 1d, e).

Since we previously showed ongoing neuronal loss in healthy ENS[19], we reasoned that cells that

show hyper-aggregation of tdTomato are in advanced stages of cell death. Cells with hyper-ag-

gregation of tdTomato showed a lack of staining for the nuclear marker DAPI (Suppl. Fig 1e),

suggesting that these are indeed cells in the advanced stages of death, and given that they are

not labeled with antibodies against Hu, their presence does not alter our estimation of tdTomato-

expressing neurons (Suppl. Fig 1f). Thus, the absence of reporter expression in myenteric ganglia

of freshly harvested tissue and the presence of tdTomato aggregation in both freshly harvested

and fixed tissues shows that our observations are not caused due to any technical issues in tissue

isolation and preservation.

Additional confirmation was provided by an analogous Wnt1-cre:*Hprt*-tdTomato lineage-traced

mouse line, in which tdTomato was expressed from the *Hprt* locus in a Wnt1-cre-dependent man-

ner (*Hprt* locus is X-inactivated in females, hence only adult male mice were used)[20]; and by the

Pax3-cre:*Rosa26*-tdTomato lineage-traced mouse line, where *Rosa26*-tdTomato driven by the

Pax3-cre transgene labels the derivatives of the neural tube and pre-migratory NC (Suppl. Fig 1g,

h)[21]. Similar lack of reporter expression was previously observed in adult myenteric neurons from

Sox10-cre NC-lineage-traced mice, and more recently in significant numbers of myenteric neu-

rons from Wnt1-cre:*Rosa26*-H2B-mCherry mice, which expresses nuclear localized reporter

mCherry in NC-lineage cells, further confirming our observations of the presence of non-NC-de-

rived neurons in the adult ENS[13,22]. In addition to the ANNA-1 antisera, which is known to detect

all neuronally significant Hu proteins (HuB, HuC, and HuD)[23], we found that another often used

antibody thought to be specific to HuC and HuD also detects HuB protein (Suppl. Fig 1l) – showing

that these antibodies do not detect the expression of a specific neuronal Hu antigen.

Lineage-tracing confirms a mesodermal derivation for half of all adult myenteric neurons.

Alternative sources of enteric neurons proposed previously include the ventral neural tube[15], and

the pancreatic endoderm[14], but the interpretation of these studies was limited by the lack of robust

lineage markers[16]. Brokhman et al.[14] found evidence of labeled neurons in inducible Pdx1-cre,

Foxa2-cre, and Sox17-cre lineage-traced mouse lines and inferred their derivation from pancre-

atic endoderm. However, in a Pdx1-cre lineage-traced mouse line, many neuroectoderm-derived

neurons of the central nervous system have also been previously shown to be derived from Pdx1-

expressing cells[24], suggesting that Pdx1 is not an exclusive endodermal marker. Foxa2 is ex-

pressed by mesoderm-derived cells in the heart[25] and Sox17 also labels mesoderm-derived cells,

especially cells of the intestinal vasculature[26-29]. We therefore hypothesized that the embryonic

mesoderm may be the true developmental source of the non-NC enteric neurons.

Mesoderm-derived cells during embryogenesis express Tek[25] and analysis of LM-MP tissues from

adult male Tek-cre:*Hprt*-tdTomato lineage-traced mice revealed the presence of a population of

tdTomato$^+$ neurons (Fig 2a). Since Tek is not expressed in the adult ENS (Suppl. Fig 2a), the

presence of Tek-derived adult myenteric neurons suggested their mesodermal origin. We then

used *mesoderm posterior 1* or Mesp1 as a definitive developmental marker for the embryonic

mesoderm[30-34] to confirm the mesodermal derivation of the non-NC derived adult enteric neurons.

Reporter expression in this mouse line has been previously studied in cardiac development, which

we confirmed by observing its expression in the adult heart tissue (Fig 2b). Expression of

tdTomato in vascular cells of both Tek-cre and Mesp1-cre mice was more pronounced than in

myenteric neurons (Fig 2a, c, d) and in cardiac myocytes of Mesp1-cre mice (Fig 2b, Suppl. Fig

2b), which reflects the variable expression of the reporter gene in both the neural crest-lineage

and mesoderm-lineage specific transgenic mouse lines (Fig 2b, c; Suppl. Figs 1a, 2c). Using an

analogous Mesp1-cre:*Rosa26*-EGFP lineage-traced mouse line, in which EGFP was expressed

from the *Rosa26* locus in a Mesp1-cre-dependent manner, we next confirmed the derivation of a

population of myenteric neurons from Mesp1-expressing mesoderm (Fig 2e, Suppl. Fig 2d, e).

Similar to our observations with the tdTomato reporter (Fig 2a-d), we observed lower expression

of the reporter EGFP in myenteric neurons as compared to vascular cells (Fig 2e; Suppl. Fig 2d,

e). Variable expression of CAG and other pan-cellular promoters in various cell-types and during

processes of maturation have previously been reported[35-38]. In addition, an earlier study by Agah

et al. showed differing degrees of CAG-driven LacZ reporter activation in a cardiac-specific trans-

genic cre mouse line, which were unrelated to copy number, suggesting insertional and positional

effects or, potentially, differential methylation[39]. These reports are consistent with our observation

and could potentially help explain the observed variable expression of CAG and CMV promoter-

driven reporter genes in our study.

Analysis of small intestinal LM-MP from P60 Mesp1-cre:*Rosa26*-tdTomato lineage-traced mice

showed that tdTomato-expression was observed in about half of all myenteric neurons (Fig 2f;

percent tdTomato$^+$ neurons: 50.28 ± 2.89 SEM; enumerated from 484 neurons from 3 mice). Be-

cause of the variable expression of the reporter gene in the transgenic mice described above, we

also labeled Mesp1-cre:*Rosa26*-tdTomato$^+$ neurons with the S46 antibody, which was raised

against the slow tonic myosin heavy chain protein MHCst derived from avian embryonic upper

leg muscle, and is expressed exclusively by mesoderm-derived cell populations (Fig 2g)[40-43].

MHCst immunostaining was exclusively observed in all Tek-cre:*Rosa26*-tdTomato$^+$ MENs (Suppl.

Fig 2f). None of the 107 MHCst$^+$ intra-ganglionic cells observed (35.67 ± 6.17 SEM MHCst$^+$ cells

per mouse) across 3 Wnt1-cre:*Rosa26*-tdTomato mice expressed tdTomato, suggesting that

MHCst immunostaining was exclusive to non-NC lineage neurons (Fig 2h, Suppl. Fig 3a). MHCst

antibody was also found to label extra-ganglionic smooth muscle cells in the LM-MP tissue (Suppl.

Fig 3a). Thus, both lineage tracing and protein biomarker expression provides strong support for

their mesodermal origin (Fig 2g, Suppl. Fig 3a). Along with our observations on Wnt1-cre:*Rosa26*-

tdTomato mice (Fig 1c), our results indicate that these two distinct lineages together account for

all the adult small intestinal myenteric neurons.

The proteins RET, a receptor tyrosine kinase that transduces GDNF signaling in NC-derived en-

teric neuronal precursors, and MET, a receptor for hepatocyte growth factor (HGF), are expressed

by different subsets of adult myenteric neurons[44]. MET is classically expressed by mesoderm-

derived cells[45], and by using immunostaining of small intestinal LM-MP tissues from Wnt1-

cre:*Rosa26*-tdTomato and Mesp1-cre:*Rosa26*-tdTomato mice, we found that the expression of

MET was restricted to a sub-population of adult MENs (Fig 3a, b). By contrast, RET expression

was confined to NENs (Fig 3c).

We then studied whether MENs and NENs differed phenotypically. In the current paradigm re-

garding the neurochemical basis for the functional classification of adult enteric neurons, inhibitory

enteric motor neurons express the nitric oxide-producing enzyme nitric oxide synthase 1 (NOS1),

excitatory motor neurons express the acetylcholine-producing enzyme choline acetyl transferase

(ChAT), and enteric sensory neurons, called intrinsic primary afferent neurons (IPANs) express

the neuropeptide calcitonin gene related peptide (CGRP)[9,44,46]. Immunostaining using antibodies

against NOS1, ChAT, and CGRP in the LM-MP from P60 Wnt1-cre:*Rosa26*-tdTomato mice

showed that both tdTomato$^+$ NENs and tdTomato$^-$ MENs express these neuronal markers (Fig

3d, f, h). Quantification of proportions in these neuronal lineages shows that the majority of NOS1$^+$

inhibitory neurons and ChAT$^+$ excitatory neurons are NENs (Fig 3e; of the 616 NOS1$^+$ neurons

observed across 3 mice, NOS1$^+$ tdTomato$^-$: 27.06% ± 4.46 SEM; $p = 0.0019$; Student's t-test);

(Fig 3g; of the 912 ChAT+ neurons observed across 3 mice, ChAT$^+$ tdTomato$^-$: 26.34% ± 2.54

SEM; $p = 0.0002$; Student's t-test). By contrast, the majority (~75%) of CGRP$^+$ neurons were

found to be MENs (Fig 3i; of the 146 CGRP$^+$ neurons observed across 3 mice, CGRP$^+$ tdTomato$^-$

: 75.76% ± 0.63 SEM; p < 0.0001; Student's t-test). These results are in keeping with a previous

report by Avetisyan et al. that shows low expression of NOS1 (0% of MET$^+$ neurons were NOS$^+$)

and ChAT (<8% of MET$^+$ neurons were ChAT$^+$) and abundant expression of CGRP by MET$^+$

neurons in the adult ENS[44], previously not known to be derived from a different lineage. In addi-

tion, MENs also express Cadherin-3 (CDH3, Suppl. Fig 3b), which is known to also mark a sub-

population of mechanosensory spinal neurons[47]. Apart from their derivation from a distinct devel-

opmental lineage, the mean cell size of MENs was significantly larger than that of NENs (Suppl.

Fig 4a; n = 143 neurons/group observed across 3 mice; Feret diameter (µm) MENs: 17.47 ± 0.50

SEM, NENs: 13.03 ± 0.36 SEM, p = 0.002; Student's t-test).

An expanded molecular characterization of MENs using unbiased single cell RNA se-

quencing (scRNAseq)-based analyses. Recent studies have used single cell or single nuclei-

based transcriptomic analyses to query molecular diversity of enteric neurons. These studies

have used either used enrichment methods based on FACS sorting neural crest-derived cells

(using reporter-expression from neural crest-specific lineage fate mapping mice)[22,48,49], Phox2b-

expressing cells[50], Baf53-derived cells[51] or used classical NENs markers such as Snap25 to drive

identification and/or enrichment of enteric neurons[49]. Instead of using these strategies, we relied

upon another accepted use of scRNAseq, which allows for sequencing of diverse un-enriched

cell populations from a tissue, to discover any novel cell types in that tissue[52]. We performed truly

unbiased and agnostic clustering of the scRNAseq data from tissues of two 6-month-old adult

male C57BL/6 wildtype mice (Fig 4a; Suppl. Fig 4b, c). We identified clusters of NC-derived cells

by exclusive expression of canonical NC markers Ret and Sox10 (which include the clusters of

Ret-expressing NENs and Sox10-expressing Neuroglia); and the MENs cluster by its co-expres-

sion of the genes Calcb (CGRP), Met, and Cdh3 (Fig 4a, b). With cells from both samples pooled

together, we compared 1,737 NC-derived cells across the NENs (77 cells) and Neuroglia (1660

cells) clusters with 2,223 cells in the MENs cluster. A list of top marker genes for every cluster are

provided in Table S2.

The neuronal nature of MENs was first established by testing our scRNAseq dataset for expres-

sion of genes known to be expressed by enteric or other neurons. We found detectable expres-

sion of genes Elavl2 (HuB), Hand2, Pde10a, Vsnl1, Tubb2b, Stmn2, Stx3, and Gpr88 in the MENs

as well as in the NC-derived cell clusters (Fig 4b). While expression of Elavl2, Stmn2 and Hand2

has been previously observed in enteric neurons[53,54], the expression of neuronal marker genes

Pde10a (expressed by medium spiny neurons in the striatum[55]), Vsnl1 (expressed by hippocam-

pal neurons[56]), Gpr88 (expressed by striatal neurons[57]), Stx3 (expressed by hippocampal neu-

rons[58]), and Tubb2b (expressed by adult retinal neurons[59]) has not been studied. Using antibodies

against these markers along with co-staining with anti-Hu ANNA1 antisera to label neurons, we

visualized the expression of these neuronal markers in both tdTomato⁺ NENs and tdTomato⁻

MENs of the small intestinal myenteric plexus tissue from adult Wnt1-cre:*Rosa26*-tdTomato mice

(Fig 4c, Suppl. Fig 3l-q). In addition, the MENs cluster was also enriched in previously character-

ized ENS markers, such as Ntf3 and Il18[2,60] (Fig 4b). We validated the MEN-specificity of these

markers using immunochemical analyses (Fig 4c; Suppl. Fig 3b - k). In addition, we also detected

>40 neuronally significant genes whose expression was unique to or enriched in MENs scRNAseq

cluster, when compared with other larger cell clusters of macrophages, smooth muscle cells, neu-

roglia, and vascular endothelium (Suppl. Fig 5). These genes include neurotransmitter receptor

genes (such as Gabra1, Gabra3, Gria3, Grik5, Grind2d, Npy1r); ion channels genes (such as

Cacna1a, Cacna1g, Cacnb3, Clcn3); gap junction genes (Gjc1, Gjb5); transient receptor potential

channel genes (Trpv4, Trpc1, Trpc6); potassium channel genes (such as Kcnn1, Kcnj8, Kcnd3,

Kcnh3, Kcns3); hormone encoding genes (Ghrh, Gnrh1, Nucb2) along with other neuronal genes

such as, Prss12 (encoding for Neurotrypsin)[61], Uchl1 (encoding for pan-neuronal marker

PGP9.5)[62], Cplx2 (encoding for Complexin 2)[63], Gpm6a (encoding for neuronal membrane

glycoprotein M6-A)[64], and Vamp2 (encoding for Synaptobrevin 2)[65] (Suppl. Fig 5). Owing to the

small size of the NENs cluster, a comparative analysis between these two neuronal lineages for

these neuronally significant genes was not possible. However, the discovery of these neuronally

significant genes in the MENs cluster, and the subsequent validation of a set of them provides

evidence for the neuronal nature of MENs. Further examination of the MENs cluster yielded ad-

ditional MENs-specific marker genes Slpi, Aebp1, Clic3, Fmo2, Smo, Myl7, and Slc17a9, whose

expression by adult enteric neurons has not been previously described and which we also vali-

dated (Fig 4b).

Next, we examined the MENs cluster for expression of a mesenchymal gene that would under-

score the mesodermal nature of MENs. Decorin (Dcn) is a gene coding for a chondroitin-dermatan

sulphate proteoglycan, which is expressed by mesoderm-derived cells, including fibroblasts and

smooth muscle cells[66]. Decorin was found to be highly expressed in the MENs cluster in our data

(Fig 5a), and not by the neurons in the Zeisel et al. database (Fig 5b)[48]. We used a validated

antibody against DCN[67] and found that this protein, known to be specifically expressed by meso-

derm-derived cells, is expressed specifically by tdTomato-negative small intestinal myenteric neu-

rons of the Wnt1-cre:*Rosa26*-tdTomato lineage fate mapping mouse (Fig 5c).

These data show that the MENs scRNAseq cluster, which can be identified by its co-expression

of neuronal markers (Elavl2, Pde10a, Hand2, etc), validated cell surface marker (Cdh3), and the

mesodermal marker Dcn is the same as the population of Hu-expressing neurons that do not

express tdTomato in the Wnt1-cre:*Rosa26*-tdTomato neural crest lineage fate mapping mouse.

We thus establish the distinct nature of MENs based on the co-expression of many neuronally

significant genes and of mesodermal genes by this cell population of Wnt1-cre:*Rosa26*-tdTomato-

negative neuron. Our experimental strategy of performing scRNAseq on LM-MP cells without any

pre-enrichment helps detects this newly characterized cell type. However, it does not allow us to

compare and contrast the transcriptomic profiles of MENs and NENs – especially given that we

do not know the true size of NENs represented in our dataset. This is exemplified by the fact that

cells within the Neuroglia cluster showed expression of glial genes such as Sox10 as well as

neuronal genes such as Ncam1, Hand2 and Stmn2, suggesting that NC-derived neurons may be

present within both the smaller NENs cluster as well as the larger Neuroglia cluster (Fig 4b).

Detection of MENs in other murine and human datasets. Since Phox2b, the transcription fac-

tor expressed by all ENS cells, shows significant higher expression in some neurons than in all

ENS glial cells[68], May-Zhang et al.[50] reasoned that flow sorted nuclei of Phox2b-CFP*high* cells from

a Phox2b-CFP reporter mouse would provide for an enriched population of ENS neurons that can

be used for single nuclei RNA sequencing (snRNAseq). They first performed bulk-RNAseq on

Phox2b-CFP+ nuclei that were flow sorted based on CFP-expression level into CFP*low* and CFP*high*

fractions, of which the CFP*low* fraction was found enriched with glial-specific gene expression pro-

file. May-Zhang et al.[50] used this to provide a rationale for performing snRNAseq on Phox2b-

CFP*high* nuclei, which they presumed to consist of all enteric neurons. Since, MENs make up

roughly half of all adult ENS neurons, we hypothesized that they should also express Phox2b. By

using a PHOX2b-specific antibody, we validated the expression of this important transcription

factor in MENs (Fig 5d). Next, we tested whether we could detect some of the genes that show

enriched expression in MENs (namely, Smo, Aebp1, Cdh3, Fmo2, Il18, Slpi, Upk3b, Msln, Hand2)

in the Phox2b+ bulkRNAseq data from May-Zhang et al.[50], and found that not only were these

genes expressed by their Phox2b+ cells, but expression of these genes was also significantly

enriched in the Phox2b-CFP*low* fraction (Fig 5e) that was not used for subsequent snRNAseq

experiments. These data provide evidence that MENs express Phox2b, both transcriptomically,

as well as at the protein level. We next queried whether any MENs were represented in the May-

Zhang et al.'s snRNAseq data generated on Phox2b-CFP*high* nuclei, given that they reported on

the presence of a cluster of adult enteric neurons that expressed mesenchymal markers. We

again tested whether the MENs-expressed genes, such as Decorin, etc were also expressed by

the 'mesenchymal' neurons in the May-Zhang et al.[50] dataset. We found a high degree of similarity

between the top genes expressed by the MENs cluster in our scRNAseq dataset and the 'mes-

enchymal' neurons in the May-Zhang et al.'s snRNAseq dataset (Table 1, Suppl. Fig 4d). Further-

more, examination of the top genes shows that this Decorin-expressing neuronal sub-population

is significantly different from the other Calcb-expressing neuronal populations, which we presume

to be the CGRP-expressing NENs (Table 1).

To confirm that the 'mesenchymal' neurons identified by May-Zhang et al.[50] are MENs, we used

an agnostic bioinformatics projection-based approach which we have previously published[69]. This

approach allows us to learn latent space representations of gene expression patterns using single

cell transcriptomic data, generating patterns that correspond to cell-type-specific gene expression

signatures[69]. We can then project other single cell and bulk RNA transcriptomic datasets into

these learned patterns to accurately quantify the differential use of these transcriptional signatures

in target data across platforms, tissues, and species[69]. In this instance, gene expression patterns

were learned on our murine single cell transcriptomic data using non-negative matrix factorization

(NMF) dimensionality reduction and the data was decomposed into 50 distinct NMF-patterns[69,70]

(Fig 6a). Four NMF patterns were identified that were specific to MENs (Fig 6a, b; Suppl. Fig 4e).

We next projected the snRNAseq dataset from May-Zhang et al.[50] into the four MEN-specific

patterns and found that all these patterns specifically labeled the cluster of 'mesenchymal' neu-

rons in the May-Zhang et al. dataset of the adult ileal ENS (Fig 6b). Further, upon clustering the

nuclei based on their projection weights from the four MEN-specific patterns, we identified a clus-

ter of nuclei that showed higher usage of all four patterns, when compared to all the other nuclei

sequenced in the dataset (Suppl. Fig 4f). The genes that define this cluster of cells are similar to

the top-expressed genes that define our MENs cluster (Suppl. Fig 4g, Table 1). This approach,

together with the high similarities in the top-expressed genes, establishes that cluster of neurons

annotated as 'mesenchymal' neurons by May-Zhang et al. are indeed MENs. Further, one of the

markers used by May-Zhang et al. to describe their mesenchymal neuronal cluster was the gene

Myh11[50], which is known to be expressed by mesoderm-derived smooth muscle cells in the adult

gut[71]. To test whether Myh11 is expressed by MENs, we immunolabeled the small intestinal LM-

MP tissue from an adult Wnt1-cre:*Rosa26*-tdTomato mouse with a monoclonal antibody against

MYH11 (Invitrogen) and found that tdTomato-negative and Hu⁺ neurons exclusively immuno-

labeled for MYH11 (Suppl. Fig. 6a). We confirmed that a subpopulation of adult enteric neurons

expressed Myh11 by immunolabeling small intestinal LM-MP tissues from a tamoxifen treated

Myh11-cre^ERT2:*Rosa26*-YFP transgenic mouse line with anti-YFP/GFP and anti-Hu antibodies

(Suppl. Fig. 6b). Anti-MYH11 antibodies labeled circular smooth muscle in the human duodenum,

where it did not immunolabel nerve fibers and some submucosal neurons immunostained with

antibodies against the pan-neuronal marker PGP9.5 but showed immunolabeling against

PGP9.5-expressing myenteric neurons in the adult healthy small intestinal tissue (Supp. Fig 6c,

d). Thus, we confirm that the MENs identified by us correspond to the cluster of 'mesenchymal

neurons' observed by May-Zhang et al. As compared to our scRNAseq dataset, where MENs

comprise of a significantly large numbers of cells, the MENs cluster in the May-Zhang et al. da-

taset is significantly smaller, possibly due to the fact that a large number of MENs were left unse-

quenced in their Phox2b-CFP*low* fraction.

Similarly, Drokhlyansky et al.[49] using MIRACL-seq found a small cluster of cells that express the

MENs-signature genes (Cdh3, Dcn, Slpi, Aebp1, Wt1, Msln, Fmo2). However, the neuronal nature

of these cells was not correctly identified as the study used a gene signature profile specific for

neural crest-derived neurons (derived from their experiment with enriched Wnt1-cre2:*Rosa26*-

tdTomato-expressing neural crest-derived cells). As a result, this cluster of cells – which shows

similarity to the MENs transcriptomic profile, was instead annotated as *Mesothelial*. This 'meso-

thelial' cluster in the Drokhlyansky et al.[49] dataset shows similar gene expression profile as that

of the cluster of 'mesenchymal neurons' from the May-Zhang et al.[50] dataset (Table 1), especially

given that many of the marker genes for both clusters are similar (Lrrn4, Rspo1, Msln, Upk3b,

Upk1b, Gpm6a, Wnt5a, Gpc3, Sulf1, Muc16). The 'mesothelial' cluster also expresses neuronally

significant genes such as Gabra3 (GABA receptor A subunit 3[72]), Prss12 (Neurotrypsin[61,73]),

Synpr (Synaptoporin[74]), Trpv4 (Transient receptor potential cation channel subfamily V member

4[75]), and Nxph1 (Neurexophillin-1[76]). Unfortunately, a deeper interrogation of this dataset was not

possible due to the size and the manner in which the original data was processed[49]. In addition,

MENs were not represented in the snRNAseq carried out in Wright et al.[22], which was based

solely on neural crest-derived cells.

In contrast to these snRNAseq based studies, the Marklund Lab in two studies performed single

cell RNA-sequencing (scRNAseq) on juvenile ENS cells, first on tdTomato[+] cells from P21 Wnt1-

cre:*Rosa26*:tdTomato lineage fate mapping mice in Zeisel et al.[48] and then from the newly char-

acterized Baf53b-cre:*Rosa26*-tdTomato BAC-transgenic mouse line[77] in Morarach et al.[51]. Barring

2 small clusters (ENC5 and ENC11), the other neuronal clusters identified in Morarach et al.[51]

mapped to the neural crest-derived clusters identified earlier by them in Zeisel et al.[48]. The gene

expression signature of these two clusters (ENC5: Sst, Calb2; ENC11: Npy, Th, Dbh) did not

match the signature of our MENs cluster or that of the cluster of mesenchymal neurons in the

May-Zhang et al.[50] study (Table 1). We reasoned that the lack of MENs in the Morarach et al.[51]

data may be driven by the non-comprehensive nature of Baf53b-cre in this BAC transgenic mouse

line labeling all neurons. On immunolabeling LM-MP tissues from two P21 Baf53b-cre:*Rosa26*-

tdTomato mice with antibodies against Hu, we enumerated 1312 Hu[+] cells in the ileum (759 from

mouse 1 and 553 from mouse 2), 5758 Hu[+] cells in the proximal colon (3411 from mouse 1 and

2347 from mouse 2), and 3352 Hu[+] cells in the distal colon (1894 from mouse 1 and 1458 from

mouse 2), and found that a significant population of myenteric neurons did not express tdTomato

(Suppl. Fig. 6e, f). Given this non-comprehensive nature of the Baf53b-cre BAC transgenic line in

labeling all myenteric neurons, and the congruence between the transcriptomic profiles of cells

analyzed using the Baf53b-cre and the neural crest-specific Wnt1-cre lines, we infer that molec-

ular taxonomy performed on enteric neurons in the Morarach et al.[51] dataset is restricted to NENs.

In a recent report, Elmentaite et al.[78] performed scRNAseq on dissociated single cells from the

human gut at fetal, juvenile, and adult ages and catalogued multiple intestinal cell-types to gen-

erate a comprehensive gut cell atlas. Canonical enteric neurons, while adequately represented in

the fetal ages, were not found in the juvenile and adult ages which were dominated by mesen-

chymal and other cell types[78]. Given that our scRNAseq approach of sequencing diverse cell

types from the murine gut wall without any marker-based pre-selection was similar to this ap-

proach in human tissue, we reasoned that MENs would also be represented in their description

of mesenchymal cell clusters, especially in the juvenile and adult ages. To detect putative MENs

in this human gut cell atlas data, we projected the entire human mesenchymal cell scRNAseq

dataset into the four murine MEN-specific NMF patterns and detected putative human MENs

clusters (Suppl. Fig 4h). Since our identification of MENs and the generation of MEN-specific NMF

patterns was based on post-natal healthy murine tissue, we next projected data from only post-

natal healthy mesenchymal clusters into the four MEN-specific NMF patterns (Fig 6b). The four

patterns identified a population of cells within the post-natal intestinal mesenchymal cell clusters

suggesting that these cells are putative MENs in the human gut (Fig 6b). These putative MENs,

which map to Cluster 1 in our representation of the intestinal mesenchymal cells from the post-

natal gut cell atlas data (Fig 7a) were annotated previously by Elmentaite et al. as transitional

stromal cells (Fig 7b). We next tested and found that the MENs marker Dcn (Fig 7c) along with

pan-neuronal Uchl1 among other neuronal markers Hand2, Stx3, Slc17a9, Pde10a and Tubb2b

(Fig 7d) that we earlier showed to be expressed by adult murine MENs, were expressed by the

putative MENs in the adult human gut.

We hypothesized that the reason these cells were not annotated as enteric neurons was due to

the use of a NENs-restricted gene list. Apart from the NENs marker Ret, such gene lists often

contain the pre-synaptic gene Snap25, a component of the SNARE complex, which is widely

assumed to be a pan-neuronal marker in the adult ENS[48,49,79,80]. While we have previously ob-

served the NENs-specific nature of RET, our transcriptomic data suggests that MENs either do

not express Snap25 or do so in very small amounts (Suppl. Fig 4d). We tested the expression of

SNAP-25 by NENs and MENs in the myenteric ganglia of an adult Wnt1-cre:*Rosa26*-tdTomato

mouse and found that SNAP-25 expressing neurons expressed tdTomato, but not the MENs-

marker MHCst, suggesting the NEN-specific expression of this canonical marker for synaptic neu-

rons (Fig 8a). Next, we used a validated Snap25-Gcamp knock-in mouse line[81-83], where

GCamp/GFP is knocked in at the Snap25 locus and hence is expressed by Snap25-expressing

cells, to confirm that the expression of Snap25 is indeed restricted to a subset of myenteric neu-

rons in the adult LM-MP layer. By using antibodies against Hu and against GFP, we found that

the expression of Snap25-GFP is indeed restricted to a subset of adult myenteric neurons (Fig

8b). Finally, by immunolabeling adult small intestinal LM-MP of Snap25-Gcamp mice with anti-

bodies against MHCst and against GFP, we confirmed the lack of Snap25 expression by a pop-

ulation of MENs (Fig 8c). The low or lack of expression of Ret, Snap-25, Elavl3 and Elavl4 in

MENs as observed both by us and May-Zhang et al.[50] may help explain why these canonical gene

lists missed the correct identification and annotation of MENs' neuronal nature in human da-

tasets[78]. Next, we examined the expression of MENs markers MHCst, MET, SLC17A9, and

DECORIN in LM-MP tissues from adult humans with no known gut motility disorder and found

them to be expressed by a population of myenteric neurons in normal adult human ENS (Fig 9;

Suppl. Fig 7a, b, c). Thus, our bioinformatic approaches and immunofluorescence analyses of

MEN-specific markers in adult human gut together shows that the human gut similarly contains a

population of MENs.

We next tried sub-clustering the MENs cell cluster in our scRNAseq data to study whether indi-

vidual subclusters could be discriminated on the basis of MENs markers or other genes. However,

at the current sequencing depth, these clusters did not yield meaningful information on specific

marker genes that could be used to define them (Suppl. Fig 8a, b).

The proportion of mesoderm-derived neurons expands with age to become the dominant

population in the aging ENS. Since the ENS at birth consists solely of NENs[13], we next studied

the birth-date and eventual expansion of the MEN lineage in the post-natal gut. Using Wnt1-

cre:*Rosa26*-tdTomato mouse line, we enumerated the tdTomato⁻ MENs in LM-MP at different

ages and found a significant age-associated increase in MENs proportions (Fig 10a, b, c; One-

way ANOVA, F = 117.6, DFn, DFd = 5, 12; p < 0.0001). At P11, MENs were found only in few

isolated myenteric ganglia (Fig 10a) and together represented only ~5% of all myenteric neurons

(tdTomato⁻ neurons: 4.12% ± 2.98 SEM; n = 1327 Hu⁺ neurons counted from 3 mice), suggesting

that they originate shortly before this age. The proportion of MENs in myenteric ganglia rises

sharply thereafter: by P22, they account for ~30% of all neurons (tdTomato⁻ neurons: 29.63% ±

1.229 SEM; n = 742 neurons counted from 3 mice); and at P60 they represent ~45% of all neurons

(tdTomato⁻ neurons: 46.38% ± 4.62 SEM; n = 700 neurons counted from 3 mice). During the adult

ages, the proportions of MENs in the adult ENS remains relatively stable as at P180, MENs con-

tinue to represent roughly half of all myenteric neurons (tdTomato⁻ neurons: 57.29% ± 3.62 SEM;

n = 586 neurons counted from 3 mice). However, MENs dominate the proportions of the aging

ENS as at the very old age of 17 months (P510), the ENS is populated almost exclusively by

MENs (Fig 10b; tdTomato⁻ neurons: 95.99% ± 1.62 SEM; n = 996 neurons counted from 3 mice).

Thus, the proportions of NENs and MENs in the myenteric ganglia can be used as a biomarker

for deducing ENS age, as a healthy ENS dominant in NENs would be juvenile, one with roughly

equal proportions would be adult, and an aging ENS is dominated by MENs.

We next tested whether the representation of MEN-specific transcriptomic signatures show a sim-

ilar increase during the maturation and aging of the human gut. We tested whether the represen-

tation of the four MEN-specific NMF patterns (Fig 5a) increased with age in the human tran-

scriptomic data from Elmentaite et al.[78]. We grouped the transcriptomic data from the wide-range

of healthy post-natal specimens of the gut cell atlas into three groups: Juvenile (containing data

on 2910 cells represented by 10 samples within the age range of 4 – 12 years), Adult (containing

data on 3696 cells represented by 36 samples within the age range of 20 – 50 years), and Aging

(containing data on 2848 cells represented by 37 samples within age range of 60 – 75 years) and

tested whether the representation of the four MEN-specific patterns changed significantly be-

tween these age groups. Barring pattern 16 which showed a non-significant increase in projection

weights between the Juvenile and Aging samples (One-way ANOVA, F = 2.10, DFn, DFd = 2, 8,

p = 0.18), projection weights of MEN-specific pattern 27 (One-way ANOVA, F = 16.08, DFn, DFd

= 2, 8, p = 0.0016), pattern 32 (One-way ANOVA, F = 8.13, DFn, DFd = 2, 8, p = 0.01) and pattern

41 (One-way ANOVA, F = 9.818, DFn, DFd = 2, 8, p = 0.007) all show significant age-associated

increase in the human gut tissue (Fig 10d, every datapoint refers to mean projected pattern weight

for cells within a defined age or age range). This suggests that analogous to the age-associated

increase in MENs we observed in murine gut (Fig 10b), the proportions of MENs in the human

gut also might increase with age. We also found that the representation of MENs cluster (Cluster

1) in the post-natal human gut cell atlas data expanded with maturation and age (Suppl. Fig 7d)

GDNF and HGF levels regulate the populations of the neural crest-derived and the meso-

derm-derived neurons, respectively, in the maturing ENS. GDNF-RET signaling is responsi-

ble for proliferation and neurogenesis from NC-derived ENS precursor cells during development

as well as for the survival of Ret-expressing enteric neurons[84-87]. Similarly, HGF signaling has

been shown to be essential for the proliferation of mesoderm-derived cells[88]. Since the expression

of the HGF receptor MET and the GDNF receptor RET is exclusive to MENs and NENs

respectively, we studied the correlation between age and levels of HGF and GDNF in LM-MP (Fig

10e, f). Given that both GDNF, as well as HGF expression is found in the LM-MP layer[44,89,90], we

used LM-MP tissue to study how the levels of these two trophic factors change in and around the

ENS tissue during maturation. We found that in agreement with a previous report[88], GDNF protein

levels are markedly reduced between the P10 and the P30 ages and remains reduced thereafter

up to the P90 age (Fig 10g; n = 3 mice/age-group, F = 6.821, DFn, DFd = 1, 7, p = 0.0348, One

way ANOVA). In addition, expression of Gdnf transcripts show significant reduction between the

ages of P30 and P90 (Fig 10h; n = 6 mice/age-group, P30: 1.070 ± 0.179 SEM, P90: 0.539 ±

0.129 SEM, p = 0.037, Student's t-test). On the other hand, HGF expression increases progres-

sively between P10 and P90 ages (Fig 10i, j: n = 3 mice/age-group, Protein levels: F = 8.820,

DFn, DFd = 1, 7, p = 0.02; mRNA levels: F = 36.98, DFn, DFd = 1, 16, p < 0.0001, One way

ANOVA). The ratio of HGF to GDNF expression in ileal LM-MP tissue shows significant increase

from the age of P10 to P90 (Suppl. Fig 9a, F = 48.60, DFn, DFd = 1, 7, p = 0.0002, One way

ANOVA), and HGF expression is consistently higher than GDNF expression in the full thickness

small intestinal tissue from adult and aging mice in the Tabula muris data[91] (Suppl. Fig 9b). We

then queried the plasma proteome levels from the LonGenity cohort of 1,025 aging individuals

and found that HGF levels correlated positively, while GDNF and RET levels correlated negatively

with age (Suppl. Fig 9c)[92], suggesting parallels between our data from murine intestine and hu-

man plasma proteome data.

Since GDNF tissue levels are correlated with NENs proportions, we hypothesized that GDNF

signaling regulates NENs proportions in maturing and adult ENS. On administration of GDNF or

saline to cohorts of P10 Wnt1-cre:*Rosa26*-tdTomato mice over 10 days[84,93], we found that GDNF

treatment promoted the juvenile phenotype by enhancing the proportions of tdTomato[+] NENs and

correspondingly reduced the proportion of tdTomato[-] MENs in P20 mice to a level similar to that

seen at the P10 age, while retaining the MENs proportions in saline-treated control mice remained

at a level expected of its age (Fig 10k; Controls: %MENs: 25.87 ± 4.37 SEM of 1350 neurons

from 3 mice; GDNF: %MENs: 3.86 ± 0.07 SEM of 1301 neurons from 3 mice; p = 0.0072; Stu-

dent's t-test). Consistent with a previous report[84], the GDNF-driven expansion of NENs and as-

sociated contraction of MENs conserved the total neuronal numbers (Fig 10k; Controls: neu-

rons/ganglia = 20.60 ± 4.00 SEM; GDNF: neurons/ganglia = 24.39 ± 6.96 SEM; p = 0.52; Stu-

dent's t-test).

Since HGF tissue levels increase with age, we hypothesized that increasing HGF signaling drives

the expansion of MENs in the maturing gut. HGF administration to cohorts of P10 Wnt1-

cre:*Rosa26*-tdTomato mice over 10 days promoted an increase in the tdTomato⁻ MENs popula-

tion in P20 mice to levels previously observed in P60 mice, while tissues from saline-treated con-

trol mice exhibited a MENs:NENs ratio that is expected at P20 (Fig 10l; Controls: %MENs: 27.40

± 5.49 SEM of 1970 neurons from 5 mice; HGF: %MENs: 49.37 ± 5.52 SEM of 1704 neurons

from 5 mice; p = 0.02, Student's t-test). Similar to GDNF treatment, HGF treatment also did not

cause any significant change in total neuronal numbers (Fig 10l; Controls: neurons/ganglia =

20.35 ± 2.22 SEM; HGF: neurons/ganglia = 18.21 ± 0.69 SEM; p = 0.38, Student's t-test).

Transcriptomic evidence of MENs genesis. As the proportions of MENs in the myenteric plexus

rise significantly between the ages of P10 and P30, we reasoned that this phase was accompa-

nied by significant neurogenesis of MENs. To provide transcriptomic evidence that MENs popu-

lations indeed expand during this phase, using 10X Genomics v3.1, we again performed unbiased

scRNAseq and agnostic clustering, this time on unsorted cells from the myenteric plexus layer of

two male mice of P21 age, when the MENs population is still expanding (Suppl. Fig 10a). Using

NMF-generated MENs patterns to run projectR-based analyses (Suppl. Fig 10b) and by using the

NENs and Neuroglia-specific markers, we again annotated the neuroglia, the NENs, and the

MENs as before and found that at this age, we were able to sequence a similar number of MENs

and NENs (~500 cells) along with a large number (~800) of neuroglia cells (Fig 11a). The

expression of Met was again detected in the MENs cluster, as well as in a small subset of Ret-

expressing NENs in the P21 cluster (Suppl. Fig 10c). MENs showed significantly higher UMI per

cell, when compared to neural crest-derived cells (NENs and neuroglial cells) (Suppl. Table 4,

One-way ANOVA; $F = 187.4$, DFn, DFd = 2, 1877, $p < 0.0001$), which we expected given the

larger size of MENs[94] (Suppl. Fig 4a). By immunolabeling LM-MP tissue from 3 P21 Wnt1-

cre:*Rosa26*-tdTomato male mice with antibodies against MET, we found that proportions of MET-

expressing neurons at this age were significantly higher in the population of tdTomato-negative

MENs than in the population of tdTomato⁺ NENs (of 152 MET-immunolabeled neurons across 3

mice, 81.51 ± 1.55 S.E.M. were tdTomato-negative MENs). The Met and Ret co-expressing NENs

cluster in our data were described as populations of neural crest-derived Chat⁺ Calcb⁺ neurons

by Zeisel et al.[48] at the P21 age (Suppl. Fig 10d), where they were annotated as ENT6, 8, and 9

clusters (and which correspond to clusters ENC1, 3, 4, 7, 6, and 7 in the Morarach et al.[51] data-

base). These Met- and Ret-co-expressing neurons were also described by Avetisyan et al.[44] as

fetal-born neural crest-derived neurons that respond to both GDNF and HGF. By contrast, MENs

do not express Ret and hence are the only cell population that can respond to solely to HGF. This

data provides further evidence on the HGF-induced expansion of MENs during the juvenile phase.

We next performed *in silico* analyses on the MENs cluster using our recently published Tricycle

software[95], which is capable of inferring continuous cell-cycle position and can be applied to da-

tasets with multiple cell types, across species and experimental conditions, including sparse data

and shallow sequenced droplet-based dataset - thus allowing us to find evidence of cell cycling

in the MENs subset. P21 MENs scRNAseq data was projected into universal cell cycle principal

components (PCs) defined by the expression of 500 cell-cycle correlated genes and the continu-

ous cell cycle position (theta) was measured as the angle from the origin (Fig 11b). Based on

expression of 500 cell-cycle correlated genes, cells between 0.5π and 1.5π show hallmarks of

being cycling cells (Fig 11b). Our analyses showed that a significant number of cells in the MENs

cluster (213 out of 510) were present between 0.5 and 1.5π, which represents the mitotic position

in the continuous cell-cycle phase (Fig 11c). We confirmed our analyses by observing that the

expression of the cell cycle gene Top2a in the cells of the MENs cluster was highly correlated

with this phase (Fig 11d). Next, we found that a key cell cycle regulator gene Ect2 (epithelial cell

transforming 2), which encodes for the guanine nucleotide exchange factor protein ECT2, which

activates RhoA in a narrow zone at the cell equator in anaphase during cell division[96], was ex-

pressed in the MENs cluster and its expression was highly correlated with the cell's mitotic posi-

tion on the Tricycle plot (Fig 11e). We then used antibodies against ECT2 to immunolabel the LM-

MP from P21 mice and found expression of this key cell cycle regulator protein in DECORIN-

expressing and MHCst-expressing MENs at this age (Fig 11f, g). Thus, the Tricycle-based com-

putational analyses found evidence of cell cycling in the MENs cluster and identified a key cell

cycle gene Ect2 as a marker for putative MENs precursor cells.

Reduced GDNF-RET signaling accelerates ENS aging to cause intestinal dysfunction.

Since reduced GDNF or RET levels are associated with intestinal dysfunction in patients[80,97,98],

we hypothesized that alterations in GDNF-RET signaling unrelated to those seen with normal

aging, would cause dysfunction. To test this hypothesis, we studied lineage proportions and in-

testinal function in a mouse model of reduced RET signaling. Ret-null heterozygous mice, which

have been previously used to study the effect of reduced RET signaling in the adult ENS, have

normal ENS structure but altered gut physiology[84]. A similar mouse model with a Ret^CFP allele has

a CFP reporter inserted at its *Ret* locus rendering it null[99]. Ret$^{+/CFP}$ (or Ret$^{+/-}$) mice carrying a

single copy of the Ret gene showed significant reduction in Ret transcript and RET protein ex-

pression in the early post-natal murine gut[99]. Similarly, we found significantly reduced Ret tran-

script expression in the adult LM-MP of Ret$^{+/-}$ mice compared to age-matched wildtype Ret$^{+/+}$ mice

(Suppl. Fig 10e). In these mice, using antibodies against CFP/GFP, Hu, and MHCst, we confirmed

that the NENs marker Ret-CFP, and the MENs marker MHCst were expressed by different

neuronal subpopulations (Fig 12a). Using the adult Ret$^{+/-}$ mice, we tested the effect of partial Ret

loss on ENS lineages at two adult ages: 9 weeks (~P60) and 16 weeks (~P110). Using antibody

against CFP/GFP to detect CFP$^+$ RET-expressing neurons, we found that Ret$^{+/-}$ mice show a

significant reduction in the proportions of Ret-CFP$^+$ NENs (Fig 12b; 9 weeks: %CFP$^+$ neurons:

24.91 ± 5.42 SEM of 837 neurons from 3 mice; 16 weeks: %CFP$^+$ neurons: 13.13 ± 0.98 SEM of

1227 neurons from 5 mice; p = 0.03, Student's t-test). We observed a corresponding significant

increase in the proportions of MHCst$^+$ MENs with age in Ret$^{+/-}$ mice (Fig 12c; 9 weeks: %MENs:

58.74 ± 7.33 SEM of 644 neurons from 3 mice; 16 weeks: %MENs: 82.84 ± 3.58 SEM of 935

neurons from 5 mice, One-way ANOVA, p = 0.014), while control Ret$^{+/+}$ mice showed no signifi-

cant age-associated change in the proportions of MENs (Fig 12c; 9 weeks: %MENs: 43.27 ± 3.24

SEM of 780 neurons from 3 mice; 16 weeks: %MENs: 54.48 ± 4.07 SEM of 1276 neurons from 5

mice, One-way ANOVA, p = 0.36), which is consistent with our previous results that MENs pro-

portions in wildtype animals are relatively stable in this time window. The expedited loss of NENs

in Ret$^{+/-}$ mice confirms that depletion of endogenous RET signaling in the adult ENS accelerates

the aging-associated loss of NENs.

Having previously shown that aging mice have intestinal dysmotility[100], we tested whether the

increased loss of NENs in the Ret$^{+/-}$ ENS, concomitant with the expansion of MENs accelerated

ENS aging, causes an early onset of aging-associated intestinal dysmotility. We studied whole

gut transit time (WGTT) in a cohort (n = 8) of adult Ret$^{+/-}$ mice and their littermate control (n = 10)

Ret$^{+/+}$ mice over 7 weeks, between 9 and 16 weeks of age. While 9-week adult Ret$^{+/-}$ mice were

similar to control Ret$^{+/+}$ mice, WGTT between the two genotypes diverged with age. Consistent

with a prior report[84], 16-week old Ret$^{+/-}$ mice displayed significantly delayed intestinal transit

compared to age-matched control Ret[+/+] mice (Fig 12d; WGTT (in min) Ret[+/+]: 121.4 ± 4.01 SEM;

Ret[+/-]: 157.3 ± 14.62 SEM, p = 0.048; Student's t-test).

GDNF reverts aging in the ENS to normalize intestinal motility. Along with others, we have

previously shown that aging is associated with slowing of intestinal motility[101,102]. We hypothesized

that this may be a consequence of the replacement of the juvenile NENs population by MENs and

therefore GDNF supplementation, by restoring a more balanced MENs:NENs ratio, may prevent

age related changes in motility. We studied 17-month-old male mice (at an age where NENs

constitute only ~5% of all myenteric neurons; Fig 10c) before and after they received 10 days of

intraperitoneal injection of GDNF or saline (n=5 for each cohort). While the two cohorts showed

no significant difference in their intestinal transit times at baseline (WGTT (in min) Control: 192.8

± 11.55 SEM; GDNF: 202.4 ± 7.60 SEM, p = 0.50, Student's t-test), GDNF treatment caused

significant improvement in intestinal transit (Fig 13a, WGTT (in min) Control: 175.0 ± 8.89 SEM;

GDNF: 101.0 ± 8.91 SEM, p = 0.0004, Student's t-test), reaching levels previously observed in

healthy mice (Fig 12d).

GDNF treatment significantly reduced the proportions of MHCst[+] MENs (Fig 13b; Suppl. Fig 11a,

b; Control: %MENs: 88.76 ± 1.48 SEM of 909 neurons from 5 mice; GDNF: %MENs: 74.12 ± 5.98

SEM of 799 neurons from 5 mice; p = 0.045; Student's t-test) while increasing the numbers of

RET[+] NENs (Fig 13c; Suppl. Fig 11c, d; Control: RET[+] neurons: 1.35 ± 0.05 SEM in forty-two 40X

fields from 4 mice; GDNF: RET[+] neurons: 3.19 ± 0.56 SEM in forty-one 40X fields from 4 mice, p

= 0.017; Student's t-test). Consistent with our earlier observation, GDNF treatment did not change

the average neuronal numbers in the ileal myenteric ganglia (Control: 17.56 ± 1.82 SEM neurons

per ganglia, GDNF: 16.01 ± 1.22 SEM neurons per ganglia, p = 0.49, Student's t-test).

The two neuronal lineages in human disease. In addition to the detection of MENs in the adult

healthy gut by bioinformatic and immunofluorescence means (Fig 4, 5, 6, 7), we next examined

the relevance of the two neuronal lineages NENs and MENs in human disease. For this, we tested

whether gut dysfunction associated with pathological reductions in NENs-signaling mechanisms

are also associated with increased abundance of MENs. We mined previously generated and

publicly available transcriptomic data from intestinal tissues from humans with normal gut motility

(controls) and from patients with obstructed defecation (OD) associated with enteric neuropa-

thy.[100]. In this dataset, Kim et al. have previously shown that OD patients have significantly re-

duced intestinal expression of Gdnf and Ret[100]. Again, using our transfer learning tool *Pro-*

jectR[69,103], which allowed us to query bulk-RNAseq data across species to estimate relative use

of MENs-specific NMF-patterns generated earlier, we tested whether the usage of MENs-specific

NMF-patterns was significantly altered between transcriptomes of OD patients and healthy hu-

mans (n = 3/group)[100]. This projection analysis indicated that 2 of the 4 MENs-specific NMF-pat-

terns showed significantly higher usage in OD samples compared to controls (Fig 14a, Suppl. Fig

12; Student's t-test, $p < 0.05$), providing evidence that the enteric neuropathy in these patients is

associated with a relative increase in MENs-specific transcriptional signatures. We further queried

the differential gene expression analysis performed by Kim et al and found that in addition to Gdnf

and Ret (as reported by them) other important NEN genes (Snap25 and Nos1) were also signifi-

cantly downregulated while the MEN marker genes identified in this study (Clic3, Cdh3, Slc17a9)

(Fig 14b) were significantly upregulated.

Discussion

Current dogma states that the adult ENS is exclusively derived from neural crest precursors that

populate the gut during fetal development[104]. The results of this study indicate a much more com-

plex system, one in which the fetal and early life "juvenile" ENS consisting of neural crest-derived

enteric neurons (NENs) is incrementally replaced during maturation by mesoderm-derived enteric

neurons (MENs). Eventually, the aging ENS consists almost exclusively of the neurons of the

MEN lineage. Using a combination of floxed transgenic reporters bred into multiple lineage fate

mapping transgenic mice, this study also provides the first definitive evidence of a significant

mesodermal contribution to any neural tissue. Previously, the mesoderm and especially the Lat-

eral Plate Mesoderm (LPM) was known to give rise to diverse cell-types within the viscera, includ-

ing several in the gut[105], and our study shows that cells of this embryonic derivation also give rise

to ENS neurons. A previous report on a dual origin of the ENS described adult enteric neurons

from Foxa2[+] or Sox17[+] precursors[14] and inferred that these were endodermal in origin. However,

Foxa2 and Sox17 are also expressed by mesoderm-derived cells[25-28]. By contrast, using two NC-

lineage-traced mouse lines (Wnt1-cre and Pax3-cre), two lineage-traced mouse lines marking

mesodermal derivatives (Tek-cre, and Mesp1-cre), and robust adult mesoderm markers, we iden-

tify a population of Mesp1-derived MHCst- and MET-expressing adult enteric neurons, which

makes up the entire non-NC-derived population of myenteric neurons. These results provide ro-

bust evidence that the second source of adult enteric neurons is the mesoderm, and not the

endoderm.

Distinct neuronal nature of MENs. While the developmental origins of neurons have been

thought to be restricted to the neural tube and the neural crest, many neurons – such as placode-

derived neurons, have been known to be derived from non-neural tube, non-neural crest devel-

opmental lineages[106]. Similarly, while supposed pan-neuronal markers, such as Snap25, are ex-

pressed specifically by neurons, not all neurons express Snap25[107]. Indeed, release of neuro-

transmitters may occur independently of Snap25[108-113], suggesting that the expression of this pro-

tein is not central to neuronal functions. These present us with important context based on which,

one can study the data generated using multiple lines of evidence to infer the neuronal nature of

MENs. *Firstly*, we show that MENs are intra-ganglionic cells, which conditionally express report-

ers under mesoderm-specific lineage fate mapping models and are labeled with antibodies

against the neuronal marker Hu. *Secondly,* MENs show expression of important ENS neurotrans-

mitter-encoding or generating proteins CGRP, NOS1, and ChAT. *Thirdly,* with our single cell

transcriptomic analysis, where we used the markers Calcb, Met, and Cdh3 that we had validated

to identify and annotate the MEN cell cluster, we discovered an additional set of neuronal-specific

marker genes (Pde10a, Vsnl1, Stmn2, Stx3, Tubb2b, Slc17a9, Hand2, Gpr88, Elavl2, Ntf3) which

were found to be expressed widely or selectively within MENs. These markers include both novel

as well as established enteric neuronal markers[2,60,114]. *Fourthly*, further transcriptomic analyses

of the MENs scRNAseq cluster identifies the expression of many other neuronally significant

genes – including those encoding for ion channels, neurotransmitter receptors, hormones, cal-

cium channels, etc. – which provide a further glimpse into the diverse neuronal functions of MENs.

Finally, our analyses shows that MENs express the pan-ENS-expressed gene Phox2b, and that

our MENs scRNAseq cluster maps to a neuronal cluster in the May-Zhang et al.[50] database, which

is annotated as a 'mesenchymal neuron' cluster[50]. Our data, together with our analyses of publicly

available data, thus provides evidence on the neuronal nature of MENs. While the co-expression

of aforementioned pan-neuronal markers establishes the neuronal nature of MENs, the expres-

sion of set of key genes in MENs suggests putative specialized functional roles of these subpop-

ulations when they co-exist with NENs in adults. These include the expression of genes associ-

ated with putative sensory (Calcb)[115], neuromodulatory (Ntf3)[116], secretory (Cftr)[117], immunomod-

ulatory (Il18)[2], hormone regulating (Ghrh)[118], extracellular matrix (ECM)-modulating (Dcn), and

motility-regulating roles (Slc17a9 or VNUT)[119]. These MEN-enriched or MEN-specific expression

profiles suggest that MENs perform various roles in the regulation of diverse gastrointestinal func-

tions. While the current scRNAseq data is restricted in providing meaningful cell clusters that can

inform the true functional diversity of MENs, future work will provide detailed analyses and func-

tional characterization of MENs.

Beyond establishing the neuronal nature of MENs, our analyses also help establish the distinct

neuronal nature of MENs. Based on the lineage fate mapping experiments with multiple combi-

nations of neural crest and mesoderm lineage fate mapping mice models and based on the

expression of mesoderm-specific markers such as MHCst and Decorin, we identified that MENs

are a population of mesoderm-derived cells. MHCst is a subset of myosin heavy chain isoforms

that has been previously shown to be specifically expressed by a subset of mesoderm-derived

muscle cells[120]. While prior reports do not clarify the nature of the exact epitope and the gene

responsible for MHCst, the expression of the smooth muscle myosin heavy chain protein MYH11

specifically in MENs suggests that the epitope detected by the anti-MHCst antibody in MENs is

of the mesoderm specific protein MYH11 protein.

Further, our bioinformatic analyses of the human gut cell atlas data generated by Elmentaite et

al.[78] also shows the existence of a putative MENs-like cell cluster in the human gut, which shows

a similar co-expression of mesenchymal and neuronal markers as observed by us in our MENs

cluster and by May-Zhang et al.[50] in their 'mesenchymal neuronal' cluster. Indeed, Elmentaite et

al.[78] identified this supposed mesenchymal cluster by the expression of the mesoderm-specific

gene Decorin. Further, by observing the co-expression of neuronal and mesoderm-specific mark-

ers in human gut tissue, we validate the presence of neurons that express mesoderm-specific

markers or MENs in adult human tissue.

Detection and identification of MENs in other datasets. While May-Zhang-et al.[50] observed

the presence of a small neuronal cluster that expressed mesenchymal marker genes, their study

did not further interrogate the developmental origins of the 'mesenchymal' enteric neurons.

Drokhlyansky et al.[49] in their unbiased MIRACL-seq analyses and Elmentaite et al.[78] in their hu-

man gut cell atlas datasets detected the MENs cluster but could not correctly identify their neu-

ronal nature, possibly owing to the expression of mesenchymal genes in this cluster, and also

due to the absent or lower than expected expression of canonical neural markers such as Ret

and Snap25. In contrast to these studies that performed single cell or nuclear transcriptomic stud-

ies that were unbiased by lineage markers, other transcriptomic datasets from Drokhlyansky et

al.[49], Wright et al.[22], and Zeisel et al.[48] were generated using enriched neural crest-derived cells,

which contributed to the absence of MENs in these datasets. An important outlier to these studies
is Morarach et al.[51] from the Marklund group, which utilized the newly characterized BAC-trans-
genic line Baf53b-cre to label and flow sort enteric neuronal cells for subsequent single cell RNA
sequencing. Our analysis of this line shows that it does not comprehensively label all enteric
neurons. While this may seem at odds from the Morarach et al.[51] report, which claims a near-
comprehensive expression of tdTomato reporter in neurons, this apparent difference in the be-
havior of this transgenic line can be explained by the fact that in contrast to the established and
accepted practice of counting all Hu-immunolabeled neurons employed by us and others[22,50,121],
the Morarach et al. study only enumerated neurons that showed only cytoplasmic and not nuclear
expression of Hu. Neuronal Hu proteins (HuB, HuC, and HuD) show both nuclear and cytoplasmic
localization[122], and since the Morarach et al.[51] study only enumerated cytoplasmic Hu-expressing
neurons, their characterization of this transgenic mouse line was performed using only a subset
of all neurons. In addition, the Marklund group (in both Zeisel et al.[48] and Morarach et al.[51]) are
the only research group that performed flow sorting of late post-natal enteric neuronal cells (com-
pared to flow sorting neuronal nuclei as performed by Drokhlyansky et al.[49], Wright et al.[22], and
May-Zhang et al.[50]), which can cause significant stress on neuronal cells. These factors, together
with the BAC transgenic nature of the Baf53b-cre line, may have contributed to the lack of MENs
in the Morarach et al.[51] data. Further, all neuronal cells described in the Morarach et al.[51] study
express Snap25 (which we show is not expressed by all adult enteric neurons), and mostly map
to known neural crest-derived cell clusters identified in Zeisel et al.[48]. This suggests that the deep
transcriptomic profiling of enteric neurons carried out by Morarach et al.[51] is restricted only to the
NENs, which comprise of half of all adult enteric neurons. Thus, prior experimental data could not
establish the true developmental identity, correct annotation, and transcriptomic definition of a
population of mesoderm-derived neurons or MENs in murine and human tissue.

By contrast, we were able to correctly identify the MENs cluster in unbiased scRNAseq of adult

murine LM-MP cells and use it to both identify and validate the expression of MENs-specific mark-

ers in the adult murine LM-MP tissue, as well as use bioinformatic analyses to identify the MENs

in publicly available murine data from May-Zhang et al.[50] and human data from Elmentaite et al.[78].

This allowed us to show the expression of MENs markers MHCst, MET, DCN, and SLC17A9 in

many enteric neurons in adult humans, suggesting that the mesoderm-derived ENS may be a

feature common to mice and humans alike. MENs in both species can readily be identified by

their expression of MENs-specific markers, thus providing a convenient tool to further identify and

study these neurons.

Development of MENs and MENgenesis. The presence of small population of MENs in the gut

at the post-natal age day 10 (P10) and their expansion during maturation suggests that key

timepoints in their development correspond with the post-natal development of the ENS. Im-

portantly, since MENs populations are expanding at this age, performing scRNAseq-based bioin-

formatic analyses of the juvenile LM-MP cells not only provided us with the computational evi-

dence of significant cell cycling in the MENs cluster at this age, but also provided us with the

information on Ect2, a cell-cycle marker expressed by cycling putative MENs precursor cells. Ect2

transcripts were found expressed by cycling cells within the MENs cluster at the P21 age, which

also showed significant enrichment in the expression of the cell-cycle gene Top2a. Except for a

cluster of cells annotated as proliferating glial cells that show expression of the cell cycle gene

Top2a, Ect2 gene expression is not detected in other murine myenteric neural crest-derived cells

in the scRNAseq study performed by Zeisel et al.[48] at the P21 juvenile age. Our anatomical ob-

servation on the expansion of the MEN population in the adolescent gut along with the computa-

tional evidence of cell cycling in the MENs cluster at this age, and the detection and validation of

Ect2 gene expression in a subset of MENs provide evidence of significant MENgenesis during

the juvenile phase. ECT2 expression was found in MHCst- and DCN-expressing myenteric cells

at this age, suggesting that many of these are proliferating MEN-precursor cells, and may not be

terminally differentiated functional neurons.

The rapidly expanding MEN population is 10-11 days old in the P21 post-natal gut (as they origi-

nate in the gut just before P10 post-natal age) and thus, the age at which these cells partake in

regulation of diverse gastrointestinal functions is yet unknown. This important issue precludes us

from using the two MENs transcriptomic datasets to compare how MENs functions change with

age. Prior studies have reported on the differing birthdates of various neural crest-derived enteric

neuronal sub-populations in the developing fetal gut[9,51,123]. Similar studies need to be performed

in the future to understand the birthdates of the various subpopulations that comprise MENs, and

to ascertain when do they become functional. A recent study by Parathan et al.[121] highlighted

important neuronal and neurochemical changes in the adolescent gut. Further studies on how the

ENS matures and ages thus need to be performed in the context of NENs and MENs to truly

understand how shared and distinct functions of these two lineages change with maturation and

age. A crucial aspect of this work will require further identifying the molecular markers for the

Ect2-expressing proliferating MEN progenitor cells. Our earlier work showed that Nestin-express-

ing neural progenitor cells are neural crest-derived cells that contribute to steady state neurogen-

esis in the adult ENS[19], thus suggesting that MENs are derived from Ect2-expressing non-Nestin[+]

precursor cells.

Use of scRNAseq to identify MENs. The use of single cell and single nucleus transcriptomic

approaches provides an important tool that has been recently used to perform molecular taxon-

omy of fetal and post-natal enteric neurons and glial cells[22,48,50,51]. In addition to performing deep

sequencing that allows for cluster-based molecular taxonomy, these tools can also be utilized for

identification of novel cell types, where prior information on their presence and marker expression

is unknown[52,95,124-126]. Compared to the other studies that utilized this tool to study the ENS, we

used the single cell transcriptomics to both confirm the presence of MENs and identify more MEN-

specific markers; as well as to use the MEN-specific transcriptomic signature to query their pres-

ence in other murine and human datasets. In contrast to its use for performing molecular taxon-

omy, which requires deep sequencing of an enriched cell population, our strategy of opting for a

lower read depth (shallow sequencing) is efficient, practical, and economical for discovering and

identifying novel celltypes[52]. In addition, even if a gene is being expressed at a medium - low

level, there is a certain probability that it will not be detected by scRNAseq in both deep and

shallow sequencing approaches. This is important since the older v2.0 10X Genomics chemistry

(used for our 6-month dataset) captures only 10-14% of the cellular transcriptome, and that from

the newer v3.1 chemistry (for our P21 data) captures only a third of the transcriptome from a

suspension of homogenous single cells[127,128]. While a partial solution to this issue is to increase

sequencing depth, beyond a certain point, this strategy leads to diminishing returns as the fraction

of PCR duplicates increases with deeper sequencing. While getting insights into cellular hetero-

geneity of MENs is important for furthering our understanding of how they regulate diverse gut

functions, the transgenic mice that can specifically label only MENs or all enteric neurons to allow

for their enrichment are currently not available[49,48,22,51,68,50]. Without the availability of MENs-spe-

cific transgenic models, the scRNAseq strategy employed by us is the only currently available tool

for querying the transcriptome of MENs. These reflect significant bottlenecks that have disallowed

the comprehensive and an in-depth characterization of all enteric neurons in all the current studies

on the adult ENS.

Expression of MET and RET by MENs and NENs. Since our scRNAseq data highlighted the

lineage-specific nature of the expression of Ret and Met in the adult ENS, we studied whether

these genes regulated the origin, expansion, and maintenance of the neuronal populations that

express them. The expression of MET by MENs in the mature adult ENS is consistent with an

earlier study by Avetisyan et al. that reported that a deletion of Met in the Wnt1-cre expressing

NC-lineage did not alter the abundance of MET[+] neurons in the adult ENS[44]. If Met was expressed

solely by neural crest-derived cells, then the loss of MET by enteric neurons would have been

expected since the conditional deletion of Met gene in this floxed Met transgenic animal has been

observed to cause a significant or comprehensive loss of MET protein expression in other or-

gans[129,130,131,132]. While at first glance, observations of the mutually exclusive nature of RET and

MET provided by us and by Avetisyan et al.[44] in the adult gut might not agree with Zeisel et al.'s[48]

observed co-expression of Ret and Met transcripts in NENs at the P21 developmental stage, our

observation that a small proportion of NENs in the P21 gut express MET helps provide an expla-

nation. Our P21 data and the data from Zeisel et al.[48] are consistent with one of the observations

made by Avetisyan et al.[44], who showed the existence of a population of fetal neural crest-derived

enteric neurons that are co-dependent on GDNF and HGF and hence would express both RET

and MET. In contrast to these observations made in the fetal and juvenile ENS, the anatomical

observations made by Avetisyan et al.[44] in the adult ENS (where MET immunoreactivity in the

adult enteric neurons is exclusive of Ret expression) suggests that the population of fetal-derived

Ret and Met co-expressing neurons is lost after maturation. Compared to the P21 scRNAseq data

from Zeisel et al.[48] that shows that all Met-expressing neural crest-derived neurons co-express

Calcb (CGRP), Ret, and Chat, Avetisyan et al.'s anatomical data in the adult gut shows that de-

spite accounting for a third of all enteric neurons and expressing CGRP, MET-expressing adult

enteric neurons do not co-express Ret and only a small proportion of these MET$^+$ neurons co-

express ChAT. Thus, Avetisyan et al.'s data agrees with our data on the adult ENS which shows

exclusive expression of RET and MET by NENs and MENs, respectively, and that while MENs

consist of a large proportion of CGRP-expressing MET$^+$ neurons, they contain only a small pop-

ulation of ChAT-expressing neurons. Along with Avetisyan et al.'s data, our data suggests that

the adult small intestinal myenteric plexus contains a significant proportion of neurons that do not

immunolabel with antibodies against NOS1 or ChAT. This observation is consistent with earlier

data from Parathan et al.[121] who showed that ~35-40% of small intestinal myenteric neurons from

6-week-old mice did not immunolabel with antibodies against NOS1 or ChAT.

Since they do not express the neural crest-marker RET, MENs are the only neuronal population

that respond to HGF alone. In this study, we show that MENs are dependent on HGF-MET sig-

naling in a manner analogous to the requirement of GDNF-RET signaling by NENs. While our

study does not discount that the exogenous HGF supplementation in juvenile mice may alter

proportions of RET and MET-co-expressing neurons within the NENs subpopulation, our report

focuses on HGF's ability to drive the expansion of MENs population in the ENS. Similarly, Ret

haploinsufficiency-mediated loss of the NEN lineage causes a proportional increase in the MEN

lineage, while maintaining overall neuronal numbers. Conservation of neuronal number, despite

significant shifts in the proportions of the two neuronal lineages suggests the presence of yet

unknown quorum sensing signaling mechanisms between the two lineages that reciprocally reg-

ulate their populations to maintain the structure of the post-natal ENS. This again implies the

existence of a yet undefined precursor cell responsible for the expansion of the MEN population

in a manner analogous to what we have previously described for NENs[19].

Relevance to aging and human disease. While it is known that Gdnf expression in the mature

gut is significantly downregulated[44,84,93,133], the functional consequences of this loss have been

unclear[84]. We found that reduced GDNF-RET signaling drives the age-dependent loss of NENs

as this loss can be slowed or reversed by GDNF supplementation and accelerated by further

reduction of Ret expression. In aging animals, GDNF-driven resuscitation of NENs was associ-

ated with a functional normalization of age-associated delay in intestinal transit. Our results iden-

tify a novel role for GDNF in maintaining or resuscitating the canonical NEN population, while

preserving overall enteric neuronal numbers. In the last few years, studies have focused on iden-

tifying juvenile protective factors (JPFs), the loss of which correlates with or drives maturation-

and aging-associated phenotypes[134]. In this context, GDNF may therefore qualify as a JPF or a

senolytic as its presence maintains the dominance of NENs in maturing ENS, corresponding to a

juvenile phenotype; and its re-introduction promotes and resuscitates the genesis of NENs in

adult and aging gut to restore healthy gut function. The exact nature of the cells that respond to

GDNF re-introduction and generate NENs is yet unknown, but it can be hypothesized that these

may include Nestin+ enteric neural stem cells and/or GDNF-responsive Schwann cells[19,135]. In

addition to its effect on resuscitating NENs, acute administration of GDNF also augments peri-

staltic motility by increasing neuronal activity[22,136]. Since the aging gut has a very small number

of RET-expressing GDNF-responsive NENs, we assume that GDNF-driven normalization of in-

testinal motility would first increase the proportions of RET-expressing NENs and then augment

their activation. This further emphasizes the importance of using GDNF or similar RET agonists

in the treatment of age-associated intestinal dysmotility.

While Avetisyan et al. showed that MET expression highly correlated with CGRP-expression in

the adult murine myenteric plexus[44], MET co-expression by all MHCst+ human enteric neurons in

our data and by all PGP9.5+ neurons in human enteric neurons by Avetisyan et al.[44] suggests that

MET-expression in the human ENS may not be restricted to CGRP-expressing IPANs. In addition,

recent data shows that our understanding of the nature of markers that can correctly identify

IPANs in the murine and human ENS is still evolving[50,137], suggesting that ascribing function to

neurons using a single marker may be incorrect.

The detection of MENs cluster in the post-natal and adult scRNAseq data in the gut cell atlas

generated by Elmentaite et al.[78], their increasing representation with age, and the subsequent

validation of the MEN-specific markers MHCst, MET and SLC17A9 in a subset of adult human

ENS neurons shows that our findings may be of clinical importance. Many gastrointestinal motility

functions are compromised with advancing age, resulting in clinically significant disorders[138] Alt-

hough the exact mechanisms will need to be worked out, our results indicate that a MENs-pre-

dominant ENS is associated with significant differences in gut motility. The progressive nature of

the change in enteric neuronal lineages with age may have pathological implications for the el-

derly gut when the balance is overwhelmingly in favor of MENs. Understanding the forces that

regulate parity between these two different sources of neurogenesis therefore holds the promise
of arresting or reverting progressive loss of gut motility with increasing age. Our results may also
have implications for the pathogenesis of disordered motility unrelated to aging, as downregula-
tion of Gdnf and Ret expression has been associated with diverticular disease and obstructed
defecation in adult patients[97,100,139]. GWAS analyses further showed that Gdnf gene was in the
eQTL associated with increased incidence of diverticular disease[140]. It is in this context that the
identification of SNAP-25 as a NENs marker gene and the transcriptomic pattern analyses of
patients with chronic obstructed defecation are significant. Expression of Snap25 is upregulated
by GDNF and is significantly downregulated in gut tissues of patients with diseases associated
with significant reduction in GDNF-RET signaling (diverticular disease and obstructed defeca-
tion)[80,97,100]. Thus, while earlier thought to be a pan-neuronal[49,141], establishing the identity of
SNAP-25 as a NEN lineage-restricted marker provides us not only with an important tool to query
proportions of NENs in murine and human tissue but also suggests the NENs-limited nature of
prior observations based on Snap25 transgenic animals[79,141]. Gut dysmotility disorders that pre-
sent with conserved ENS neuronal numbers[97] have puzzled investigators on the etiology of the
disease. Our results suggest an explanation based on NENs-to-MENs lineage shifts that conserve
overall enteric neuronal numbers with variable effects on function.

In conclusion, using multiple lines of evidence, we show that the ENS of the juvenile and mature
mammalian gut is dominated by two different classes of neurons that have distinct ontogenies
and trophic factors, associated with functional differences. The shift in neuronal lineage may rep-
resent a functional adaptation to changes in nutrient handling, microbiota or other physiological
factors associated with different ages. Further research in the functional differences between the
neuronal lineages and factors that regulate the parity between these two nervous systems in
humans during the lifetime will be important to advance our understanding of the adult ENS and
the treatment of age-related and other pathological disorders of gut motility.

Acknowledgements

We would like to thank Dr. Chulan Kwon (JHU) for his kind gift of the Mesp1-cre mice, Dr. Jeremy Nathans (JHU) for his kind gift of the Tek-EGFP, Tek-cre and *Hprt*-tdTomato lineage-traced mice and his support, and Dr. Vanda Lennon (Mayo Clinic) for her kind gift of the ANNA1 anti-Hu anti-sera. We thank Dr. Akira Sawa (JHU) for his help with the *Rosa26*-EGFP mouse line, and Dr Meera Murgai (NCI) for her help with the Myh11-cre^ERT2:*Rosa26*-YFP mouse tissues. We thank Dr. Xinzhong Dong (JHU) and Dr. Mark Donowitz (JHU) for their support. The microscopy was performed on the Ross Imaging Core at the Hopkins Conte Digestive Disease Center at the Johns Hopkins University (P30DK089502) using the Olympus FV 3000rs (procured with the NIH-NIDDK S10 OD025244 grant) and we thank Dr. George McNamara for his help with confocal training and imaging. The 10X Genomics Chromium processing for scRNAseq was performed at the GRCF Core and the sequencing was performed at the CIDR core at the Johns Hopkins University. S.K. was funded through a grant from the Ludwig Foundation, a grant from the NIA (R01AG066768), a pilot award from the Hopkins Digestive Diseases Basic & Translational Research Core Center grant (P30DK089502), and a pilot award from the Diacomp initiative through Augusta University. L.A.G was funded in part by a Johns Hopkins Catalyst Award and a grant from the NIA (R01AG066768). J.S. was funded through the Maryland Genetics, Epidemiology, and Medicine training program sponsored by the Burroughs Welcome Fund. P.J.P. was funded through the Hopkins Conte Digestive Disease Center at the Johns Hopkins University (P30DK089502), NIDDK (R01DK080920), the Maryland Stem Cell Research Foundation (MSCRF130005), and a grant from the AMOS family.

Author contributions

S.K., and P.J.P. conceived the study; S.K., M.A., M.L, L.A.G. and P.J.P. designed the research studies; S.K. L.B, C.Z, A.S., A.B., and M.S. conducted the experiments; S.K., Z.H., M.K., J.L.,

929   L.B., S.B., E.V., J.S., S.N., M.S., G.L., and L.A.G. acquired and analyzed the data; S.K., and P.J.P

930   drafted the manuscript.

Methods

Animals:

Experimental protocols were approved by The Johns Hopkins University's Animal Care and Use

Committee in accordance with the guidelines provided by the National Institutes of Health. Pres-

ence of vaginal plug was ascertained as 0.5 days post-fertilization and this metric was used to

calculate age of mice. Only male mice were used for the studies detailed in this report. The Wnt1-

cre:*Rosa26*-tdTomato lineage-traced line was generated as detailed before by breeding the

B6.Cg-Tg(*Wnt1-cre*) with the Ai14 transgenic mouse line (Jax #: 007914) containing the *Rosa26*-

tdTomato transgene[19,142,143]. The Wnt1-cre line used in this study is the well validated line for

studying neural crest derivatives previously used by us and others[18,19,22,48,144], and not the Wnt1-

cre2 line used by Drokhlyansky et al.[49]. We again validated the fidelity of this line by observing

whether any aberrant recombination occurred in our Wnt1-cre:*Rosa26*-tdTomato line by studying

the intestinal mucosal tissue and found that the line behaved expectedly to label neural crest-

derivatives and epithelial cells (Suppl. Fig 1i). Pax3-cre:*Rosa26*-tdTomato lineage-traced line was

generated by breeding the Ai9 transgenic mouse line (Jax #: 007909) with the Pax3-cre trans-

genic mouse (Jax #: 005549). The Wnt1-cre:*Hprt*-tdTomato mouse was generated by breeding

our aforementioned Wnt1-cre transgenic mouse line with the *Hprt*-tdTomato transgenic mouse

line (Jax #: 021428, kind gift from Prof. Jeremy Nathans). Mesp1-cre:*Rosa26*-tdTomato mice

were generated by breeding the Mesp1-cre transgenic mice[145] (gift from Dr. Chulan Kwon, JHU)

with the Ai14 transgenic mice. Mesp1-cre:*Rosa26*-GFP mice were generated by breeding the

Mesp1-cre transgenic mice with the *Rosa26*-EGFP transgenic mouse line (gift from Dr. Akira

Sawa, JHU). Neither the *Rosa26*-tdTomato (Ai14) line nor the *Rosa26*-EGFP line showed any

reporter expression in the absence of a cre driver (Suppl. Fig 1j, k). Ret$^{+/CFP}$ mice (MGI:3777556)

were inter-bred to get a colony of adult Ret$^{+/+}$ and Ret$^{+/CFP}$ mice. Ret$^{CFP/CFP}$ mice died at or

before term. Tek-cre:*Hprt*-tdTomato mice were generated by breeding Tek-cre transgenic mice

(also known as Tie2-cre; Jax #: 004128) with the *Hprt*-tdTomato transgenic mouse line. 17-month-

old male C57BL/6 mice from the aging mouse colony of the National Institute of Aging were pro-

cured for the GDNF-treatment experiment.

Human Tissues:

Human tissues were obtained under IRB protocol IRB00181108 that was approved by Institutional

Review Board at the Johns Hopkins University. Pathological normal specimens of human duode-

num and colon were obtained post-resection. Tissues were obtained from adult donors and were

de-identified such that the exact age, gender, and ethnicity of the donors was unknown.

Tissue preparation:

Mice were anesthetized with isoflurane and sacrificed by cervical dislocation. A laparotomy was

performed, and the ileum was removed and lavaged with PBS containing penicillin-streptomycin

(PS; Invitrogen), then cut into 1-cm-long segments and placed over a sterile plastic rod. A super-

ficial longitudinal incision was made along the serosal surface and the LM-MP was peeled off

from the underlying tissue using a wet sterile cotton swab [19] and placed in Opti-MEM medium

(Invitrogen) containing Pen-Strep (Invitrogen). The tissue was then laid flat and fixed within 30

minutes of isolation with freshly prepared ice cold 4% paraformaldehyde (PFA) solution for 45

minutes in the dark to preserve fluorescence intensity and prevent photo-bleaching. All LM-MP

tissues post-isolation were fixed within 30 minutes of their isolation. After the fixation, the tissue

was removed and stored in ice cold sterile PBS with Pen-Strep for immunofluorescence staining

and subsequent microscopy.

For human tissues, duodenal tissue from adult human patients (n = 3 patients), who did not have

any prior history of chronic intestinal dysmotility, that was removed by Whipple procedure was

obtained. A colonic sample from a pathologically normal colonic resection from an adult donor

suffering from colon carcinoma who similarly did not have prior history of chronic intestinal dys-

motility was also obtained. The resected tissue was placed in ice cold Opti-MEM medium (Invi-

trogen) containing Pen-Strep (Invitrogen). The mucosal and sub-mucosal tissue was dissected

out in the medium under light microscope and the muscularis layer containing myenteric plexus

tissue was obtained. The tissue was laid out between two glass slides and fixed overnight in ice

cold 4% PFA after which it was removed and stored in ice cold sterile PBS with Pen-Strep for

immunofluorescence staining, optical clarification, and subsequent microscopy.

Immunohistochemistry:

For murine tissue: The fixed LM-MP tissue was washed twice in ice-cold PBS in the dark at 16°C.

The tissue was then incubated in blocking-permeabilizing buffer (BPB; 5% normal goat serum

with 0.3% Triton-X) for 1 hour. While staining for antibodies that were mouse monoclonal, 5%

normal mouse serum was added to the BPB. The tissue was then removed from the BPB and

was incubated with the appropriate primary antibody at the listed concentration (Table S1) for 48

h at 16°C in the dark with shaking at 55 rpm. Following incubation with primary antibody, the

tissue was washed three times (15-min wash each) in PBS at room temperature in the dark. The

tissue was then incubated in the appropriate secondary antibody at room temperature for 1 hour

while on a rotary shaker (65 rpm). The tissue was again washed three times in PBS at room

temperature, counterstained with DAPI to stain the nuclei, overlaid with Prolong Antifade Gold

mounting medium, cover-slipped, and imaged.

Immunostaining for ChAT was performed using fixed and unpeeled murine tissues from adult

male Wnt1-cre:*Rosa26*-tdTomato mice. The intestinal segments were obtained by perfuse fixing

the mouse with ice-cold PBS and then freshly prepared 4% PFA solution. Once the intestine was

harvested, the ileum was flushed with cold HBSS with 0.4 mol/L N-acetyl-l-cysteine and then PBS

to remove luminal contents. After, the tissue was fixed in 4% PFA overnight. The fixed tissue was

then immersed in 2% Triton-X 100 solution for 2 d at 15°C for permeabilization. The tissue was

immunostained with antibodies against Hu (ANNA1) and against ChAT (rabbit, Abnova). The pri-

mary antibody was then diluted (1:100) in the dilution buffer (0.25% Triton X-100, 1% normal goat

serum, and 0.02% sodium azide in PBS) and incubated for 1 day at 15°C. An Alexa Fluor 647-

conjugated goat anti-rabbit secondary antibody (1:200, Invitrogen) and an Alexa Fluor 488-con-

jugated goat anti-human secondary antibody (1:200, Invitrogen) were then used to reveal the

immunopositive structure. Finally, the labeled specimens were immersed in FocusClear solution

(CelExplorer) for optical clearing before being imaged by laser confocal microscopy using a Zeiss

LSM 510 META. The LSM 510 software and the Avizo 6.2 image reconstruction software (VSG)

were used for the 2D and 3D projection of the confocal micrographs. The Avizo software was

operated under a Dell T7500 workstation and the "Gaussian Filter" function of Avizo was used for

noise reduction of the micrographs.

Colchicine treatment: For CGRP immunostaining, mice were injected with Colchicine at a con-

centration of 5 mg/Kg body weight 16 hours (overnight) before they were sacrificed. The mice

were housed singly during this time and adequate gel packs were provided. Food and water were

provided *ad libitum*. On the following day, the mice were sacrificed, and their LM-MP tissues were

harvested as detailed above.

For human tissue: The fixed muscularis layer containing myenteric plexus tissue was removed

from ice cold PBS and incubated in blocking-permeabilizing buffer (BPB; 5% normal goat serum,

5% normal mouse serum with 0.3% Triton-X) for 4 hours. The tissue was then removed from the

BPB and was incubated with the appropriate primary antibody at the listed concentration (Table

S1) for 5 days at 16°C in the dark with shaking at 55 rpm. Following incubation with primary

antibody, the tissue was washed five times (15-min wash each) in PBS at room temperature in

the dark. The tissue was then incubated in the appropriate secondary antibody at 16°C in the dark

with shaking at 55 rpm for 2 days. The tissue was again washed in dark for five times in PBS that

contained DAPI at room temperature. After the final wash, the tissue was suspended in tissue

clarification buffer CUBIC[146] for 1 hour at 4°C in the dark after which it was overlaid with Prolong

Antifade Gold mounting medium, cover-slipped, and imaged. Briefly, the CUBIC optical clarifica-

tion buffer was made by mixing 2.5 g of urea (25% by wt), 2.5 g of N, N, N´, N´-tetrakis (2-hydroxy-

propyl) ethylenediamine (25% by wt), 1.5 g of Triton X-100 (15% by wt) in 35 ml of Distilled Water.

The solution was shaken till the ingredients were dissolved and yielded a clear viscous solution.

Microscopy:

Except for ChAT microscopy, all other imaging was done by using the oil immersion 63X objective

on the Leica SP8 confocal microscope and by using the oil immersion 40X objective on the Olym-

pus Fluoview 3000rs confocal microscope with resonance scanning mode. For thick tissues, such

as human tissues, the Galvano mode of the Olympus Fluoview 3000rs microscope that enabled

higher resolution imaging and averaging was used. Images obtained were then analyzed using

Fiji (https://fiji.sc/).

Live tissue imaging of the mouse gut tissue was performed by harvesting small intestinal tissue

from an adult Wnt1-cre:*Rosa26*-tdTomato mouse and immediately putting it in OptiMEM solution.

The tissue was then immediately put on a chamber slide containing OptiMEM and imaged in live

tissue culture conditions of the EVOS M7000 microscope under the 20X objective.

Enumeration of neurons:

Enumeration of tdTomato[+] and tdTomato[-] neurons was performed on tissue with native tdTomato

fluorescence. The native tdTomato fluorescence in fixed tissue was intense, thus there was no

need to increase fluorescent signal using anti-RFP/tdTomato antibodies.

Enumeration of enteric neurons to study alterations to ENS neuronal numbers was performed by

following the well-established protocol[147-150] of counting the numbers of neurons in myenteric gan-

glia and comparing them between animals for statistically significant differences. Identification of

myenteric ganglia was performed according to our pre-determined method published earlier[19].

Briefly, contiguous clusters of neurons were defined as a ganglion and the total numbers of neu-

rons within these clusters were enumerated as numbers of myenteric neurons per ganglion. As a

rule, clusters of 3 neurons or more were deemed to consist of a ganglion and our enumeration

strategy did not count extra-ganglionic neurons. We imaged ~10 ganglia per tissue for our enu-

meration and each group studied had n ≥ 3 mice. Identification and enumeration of neurons and

detection of co-localization was performed manually by trained laboratory personnel. Fig 1d com-

prises of the data from 6 P60 mice, which includes data from the 3 mice that were used for the

development experiment presented in Fig 8c.

Protein isolation and detection:

After the LM-MP tissue was isolated, it was weighed and placed in a sterile 1.5 ml microfuge tube.

1X RIPA buffer (Cell Signaling Technology) with Halt Protease Inhibitor Cocktail (Thermo Scien-

tific) at 5X concentration, Phosphatase Inhibitor Cocktails II and III (Sigma-Aldrich) at 2X concen-

trations were added to the tissue lysate buffer. Tissue was disrupted using 1.0 mm silica beads

in Bullet Blender 24 (Next Advance) for 5 mins at highest setting. The lysate was incubated at

4°C with shaking for 30 mins, centrifuged at 14,000rpm for 20 mins and the supernatant was

taken and stored in -80°C in aliquots. Protein concentration was estimated using Bradford assay

solution (Biorad) following the manufacturer's protocol. For immunoblotting, 40 µg of protein was

loaded per well of 4%-20% gradient denaturing gel (Biorad). Protein marker used was Precision

Plus Dual Color standards (Biorad). After fractionating the proteins, they were blotted onto Im-

munBlot PVDF membrane (Biorad) overnight at 4°C at 17V for 12-16 hours. After blotting, mem-

brane was blocked with Odyssey TBS blocking buffer (Li-Cor) for 1 hour at room temperature with

shaking. Incubation with primary antibodies were carried out at 4°C with shaking for 24 hours.

Following binding, the blot was washed 4 times with TBS-T (Tris Buffered Saline with 0.5%

Tween) for 15 mins each with shaking at room temperature. Secondary antibody incubation was

carried out in dark at room temperature for 1.5 hours with shaking. The blot was then washed 4

times for 15 mins each and imaged on Odyssey CLx system (Li-Cor).

Part of the aforementioned protocol was also followed for detecting HuB protein (recombinant

Human ELAVL2, 1-346aa, from Biosource; Catalogue number MBS205995). The protein was

loaded in two amounts (0.5 μg and 1.25 μg) in separate lanes of a 4-20% gradient denaturing gel

(Biorad) along with protein marker Precision Plus Dual Color standards (Biorad). After fractionat-

ing the proteins, they were blotted onto ImmunBlot PVDF membrane (Biorad) overnight at 4°C at

17V for 12-16 hours. After blotting, membrane was blocked with Odyssey TBS blocking buffer (Li-

Cor) for 1 hour at room temperature with shaking. Incubation with primary antibodies were carried

out at 4°C with shaking for 24 hours. Following binding, the blot was washed 4 times with TBS-T

(Tris Buffered Saline with 0.5% Tween) for 15 mins each with shak-ing at room temperature.

Secondary antibody incubation was carried out in dark at room tempera-ture for 1.5 hours with

shaking. The blot was then washed 4 times for 15 mins each and imaged on Odyssey CLx system

(Li-Cor).

Antibody used to detect RET was a well validated antibody used previously for detecting RET in

ENS[151] and was further validated by us using a western blot on total proteins isolated from murine

small intestinal LM-MP (Suppl Figure 8e). Antibodies used are detailed in the Supplementary

Table 1.

RNA isolation and quantitative detection of specific transcripts:

The isolated tissue was stored in RNALater Solution (Ambion). RNA was isolated using RNeasy

Mini Kit (Qiagen) following manufacturer's protocol. RNA quantification was carried out using

Epoch Microplate Spectrophotometer (BioTek). cDNA synthesis was carried by SuperScript IV

VILO Master Mix (Invitrogen). Quantitative Real-time PCR was carried out using Taqman Gene

Expression Master Mix (Applied Biosystems) and Roto-Gene Q (Qiagen). The probes used are

listed in Supplementary Table 1.

Single Cell RNA sequencing and analyses:

Single cell preparation from murine ileal tissues: Ileal tissues from male littermate C57/BL6

wildtype mice were isolated by peeling as previously described. The tissues were then dissociated

in Digestion Buffer containing 1 mg/ml Liberase (Sigma-Aldrich) in OptiMEM. Tissues from mouse

1 were dissociated in the Digestion buffer containing Liberase TH and tissues from mouse 2 were

dissociated in the Digestion buffer containing Liberase TL. Dissociation was performed at 37°C

for 30 minutes on a rotary shaker, after which the cells were centrifuged at 200g for 7 minutes,

and the pellet was resuspended in ice cold sterile PBS. The cell suspension was passed through

a 40 μm cell sieve and the resulting filtered cell suspension was again centrifuged at 200g for 7

minutes. This process of cell centrifugation and filtration was repeated two more times, after which

the cells were resuspended in 1 ml ice cold sterile PBS. The repeated steps of serial cell washes

and filtration removed clumps and debris and the viability of the resulting cell suspension was

estimated to be >90% using Trypan Blue dye test. The cells were then processed through 10X

Genomics Chromium V2.0 system (for the 6-month-old murine tissues) and 10X Genomics Chro-

mium V3.1 system (for the P21 murine tissues) according to the manufacturer's suggested work-

flow. The processing was done at the GRCF Core Facility at the Johns Hopkins University. The

pooled libraries were sequenced on an Illumina HiSeq 2500 (for 6-month-old tissues) and Illumina

Novaseq (for P21 tissues) to an average depth of 3.125×10^8 reads per sample library. The se-

quencing was performed at the CIDR core facility at the Johns Hopkins University.

For processing data from scRNAseq on 6-month-old murine LM-MP tissues: Pre-processing of

FASTQs to Expression Matrices: FASTQ sequence files were processed following a Kallisto

Bustools workflow compatible with downstream RNA velocity[152]. References required for pseudo-

alignment of transcripts were obtained using the get_velocity_files (functionality of BUSpaRSE

(https://github.com/BUStools/BUSpaRse), with "L=98" for 10X Genomics v2.0 sequencing chem-

istry. Reads were pseudo-aligned to an index built from Ensembl 97 transcriptome annotation

(Gencode vM22; GRCm38). Across two samples processed, a total of 578,529,125 reads were

successfully pseudo-aligned. Barcodes within a Hamming distance of one to known 10X Ge-

nomics v2.0 barcodes were corrected. Reads were classified as "spliced" or "unspliced" by their

complement to the target list of intronic sequences and exonic sequences, respectively, and sub-

sequently quantified separately into expression count matrices. Spliced counts are used for all

analyses.

Single cell gene expression analysis: scRNA-seq count matrices were analyzed using Monocle3.

11,123 high-quality cells were identified as meeting a 200 UMI minimum threshold with a mito-

chondrial read ratio of less than 20%; droplets that did not meet these criteria were excluded from

the analysis. Mitochondrial counts were determined as the sum of reads mapping to 37 genes

annotated to the mitochondrial genome. All genes with non-zero expression were included for

downstream analysis. Raw counts were first scaled by a cell-size scaling factor and subsequently

log10 transformed with a pseudo-count of 1. Normalized values are used in place of raw counts

unless otherwise noted.

Prior to UMAP dimensionality reduction, batch effects between the two biological replicates were

assessed and corrected via the mutual nearest neighbors (MNN) algorithm as implemented by

Batchelor in Monocle3[153]. 15 clusters of cells in the UMAP embedding were identified by Leiden

community detection. 30 marker genes for each cluster were identified based on greatest pseudo

R^2 values and used for supervised annotation of cell types by searching UniProt, Allen Cell Atlas

and through literature search with Pubmed.

NC-derived cell clusters were identified by expression of NC markers Ret and Sox10. MEN cluster

was identified by its expression of CGRP-coding Calcb, Met, and Cdh3. The pan-MENs protein

marker MHCst was identified by labeling with an antibody S46 which labels all members of the

MHCst family[41]. Since the antibody does not identify a single gene product, MHCst immunostain-

ing could not be used to identify a specific gene marker for use in the annotation of the MEN

cluster. For further analysis into the MENs population, the full LM-MP dataset was subset to in-

clude only the 2,223 cells annotated as MENs. These cells were re-processed as above, but with

a reduced PCA dimensionality of k=20 as input for the UMAP embedding. 5 clusters of cells in

the UMAP embedding were identified by Leiden community detection (k= 10, resolution = 5e -4).

P20 LM-MP scRNA processing and analysis: Pre-processing of FASTQs to Expression Matrices:

FASTQ sequence files were processed following a Kallisto Bustools workflow compatible with

downstream RNA velocity. References required for pseudo-alignment of transcripts were ob-

tained using the get_velocity_files (functionality of BUSpaRSE

(https://github.com/BUStools/BUSpaRse)), with "L=91" for 10X Genomics v3.1 sequencing chem-

istry. Reads were processed following the same steps as the six month LMMP dataset. Spliced

counts are used for all analyses.

Single cell gene expression analysis: scRNA-seq count matrices were analyzed using Monocle3.

11,264 high-quality cells were identified as meeting a 600 UMI minimum threshold with a mito-

chondrial read ratio of less than 20%; droplets that did not meet these criteria were excluded from

the analysis. Mitochondrial counts were determined as the sum of reads mapping to 37 genes

annotated to the mitochondrial genome. All genes with non-zero expression were included for

downstream analysis. Raw counts were first scaled by a cell-size scaling factor and subsequently

log10 transformed with a pseudo-count of 1. Normalized values are used in place of raw counts

unless otherwise noted.

NC-derived and MENs cell clusters were identified by expression of established marker genes.

Further, the identity of MENs was confirmed by a projection analysis using MENs-specific patterns

learned in the six month dataset. To investigate differences in division potential within the ENS,

the data was subset to NENs, neuroglia, and MENs for computational cell cycle analysis (NENs:

n = 526; neuroglia: n = 844; MENs: n = 510). Continuous scores for cell cycle position (0-2π) were

calculated via Tricycle by projecting each cell into the default reference space. The relevance of

these scores was confirmed by profiling the scores' correlation with expression of genes with

known variation over the cell cycle. Lastly, each cell was unambiguously characterized as "non-

cycling" (0-0.5π or 1.5-2π) or "cycling" (0.5-1.5π) by binarizing Tricycle scores.

External datasets: Spliced and unspliced count matrices for mesenchymal subset of the Gut Cell

Atlas were obtained from https://www.gutcellatlas.org/[78], and existing annotations were used

when available. SnRNAseq counts matrices from May-Zhang et al.[50] were downloaded from GEO

GSE153192. Bulk RNAseq dataset from the Human Obstructed defecation study by Kim et al.[100]

was obtained from GEO GSE101968.

Pattern discovery and ProjectR analyses: Pattern discovery was utilized to identify sets of co-

expressed genes that define cell-type specific transcriptional signatures. The normalized expres-

sion matrix was decomposed via non-negative matrix factorization (NMF) as implemented in the

R package *NNLM*, with k=50 and default parameters. Cell weights for each pattern were grouped

by assigned cell-type and represented by heatmap. Pattern vectors were hierarchically clustered

by a Euclidian distance metric, implemented in ComplexHeatmap[154]. These patterns were then

tested on the bulk RNA-Seq expression matrix for the Human Obstructed Defecation study[100].

The log2 expression (log2(rpkm + 1)) from this study was projected onto the NMF patterns using

projectR[69]. Students' t tests were performed on the projection weights from the Control and OD

groups to test for differences between them. For the projection of datasets generated by May-

Zhang et al.[50] and Elmentaite et al.[78], log10-transformed normalized counts were projected.

Analysis of May-Zhang et al. dataset

Raw UMI count matrices for the single-nucleus RNA sequencing of enteric neurons performed by

May-Zhang et al.[50] from the ileum of adult mice were obtained from GEO (GSE153192). The

DropletUtils package (v1.10.3)[155,156] was used to compute barcode ranks, knee, and inflection

point statistics for all the samples. All barcodes in the 10X genomics libraries with total UMI greater

than the inflection point were considered to be cell-containing droplets and retained for further

analysis. The inDrop libraries were filtered to retain all barcodes with UMI greater than 200. The

filtered matrices were combined to form a SingleCellExperiment object.

Cell-specific scaling factors accounting for composition biases were computed using com-

puteSumFactors (scran v1.18.7)[157]. The raw UMI counts were then scaled using the scaling fac-

tors and log 2 transformed after addition of a pseudo count of 1 using logNormCounts (scater

v1.18.6)[158]. Batch effects between the samples were corrected using fastMNN (batchelor

v1.6.3)[153].

Unsupervised clustering was performed using a graph-based clustering technique. The graph

was obtained using buildSNNgraph (scran v1.18.7)[157] and the louvain algorithm implemented in

the igraph package (v1.2.7)[159] was used to identify communities. Differential expression testing

was performed between all pairs of clusters using findMarkers (scran v1.18.7)[157]. One-sided

Welch's t-test was used to identify genes that were upregulated in a cluster as compared to any

others in the dataset. Benjamini-Hochberg method was applied (to combined p-values) to correct

for multiple testing and the genes were ranked by significance. The top ranked genes were used

as marker genes for the clusters. The log fold change from the pairwise comparison with the

lowest p-value was used as the summary log fold change.

The log normalized counts from this dataset were then projected into the NMF patterns learned

on our scRNAseq data. This was performed using transfer learning implemented in projectR

(v1.6.0). The projection weights corresponding to the four MEN-specific patterns were then clus-

tered using graph-based clustering and visualized by UMAP (uwot v0.1.10)(

https://github.com/jlmelville/uwot).

Whole gut transit time analyses:

Whole-gut transit time (WGTT) for every mouse was analyzed by the method using the carmine

red protocol[19]. Mice were placed in individual cages and deprived of food for 1 hour before receiv-

ing 0.3 mL 6% (wt/vol) carmine solution in 0.5% methylcellulose by oral gavage into the animal's

stomach. The time taken for each mouse to produce a red fecal pellet after the administration of

carmine dye was recorded in minutes. The experiment was terminated at 210 minutes post-ga-

vage and the WGTT of any mice that did not expel the red dye at the termination was marked at

the value of 210 min. The mean difference in whole gut transit time (in minutes) between both the

$Ret^{+/+}$ and $Ret^{+/-}$ mice cohorts, and the GDNF and Saline-treated Control cohorts were analyzed

statistically.

In vivo Injections:

GDNF injection: Similar to prior report that gave sub-cutaneous injections of GDNF to post-natal

mice[93], we took 6 littermate 10 day old (P10) male Wnt1-cre:*Rosa26*-tdTomato mice and divided

into two subgroups, GDNF and Control. Each mouse in the GDNF group was injected sub-cuta-

neously with 50 µl of 2mg/ml of GDNF (Peprotech Catalogue #: 450-44) every other day, while

the Control group was injected with 50 µl of sterile saline. The mice were given 5 doses and then

sacrificed on P20, after which their LM-MP tissues were isolated as detailed above. The tissues

were then immunostained with antibodies against Hu and imaged. In a separate experiment, adult

(P60) mice were also injected sub-cutaneously with GDNF (100 µl of 100 µg/ml of GDNF). The

mice were given 5 doses over a course of 10 days and then sacrificed on P70, after which their

LM-MP tissues were isolated as detailed above. For studying the effect of GDNF on aging mice,

two cohorts of 17-month-old male C57BL/6 mice (n=5 mice/cohort) were obtained from the aging

colony of the National Institute of Aging. Before the start of dosing, the whole gut transit time was

assayed. The animals were then injected daily sub-cutaneously either with 100 µl of saline

(Control) or 100 µl of GDNF (500 µg/ml) for 10 consecutive days, after which the mice were sac-

rificed, and their LM-MP tissues were isolated as detailed above. The tissues were then im-

munostained with antibodies against Hu and imaged.

HGF injection: Similar to prior report that gave sub-cutaneous injections of HGF to post-natal

mice, we took 6 littermate 10 day old (P10) male Wnt1-cre:*Rosa26*-tdTomato mice and divided

into two subgroups, HGF and Control. Each mouse in the HGF group was injected sub-cutane-

ously with 100 µl of 2mg/ml of HGF (Peprotech Catalogue #:315-23) every other day, while the

saline group was injected with 100 µl of sterile saline. The mice were given 5 doses and then

sacrificed on P20, after which their LM-MP tissues were isolated as detailed above. The tissues

were then immunostained with antibodies against Hu and imaged.

Statistics:

Data was analyzed using Graphpad Prism 8.3.1 and R using Unpaired Students t-test, Simple

Linear Regression, and Ordinary One-Way ANOVA.

Data:

All raw data are provided in Supplementary Table 2 and 3. We imaged atleast 10 ganglia per

tissue for our enumeration and each group studied had n ≥ 3 mice. Raw single cell RNA sequenc-

ing data is archived on the NCBI GEO server and can be accessed under the accession numbers

GSE156146 and GSE213604.

Code:

Code generated for the scRNA/snRNA analyses is available at https://github.com/jared-

slosberg/6month_ENS/.

1268

Figures

Fig 1: Half of all adult small intestinal myenteric neurons are derived from a non-neural
crest lineage.

(a) Enteric glia, labeled with GFAP (green) in the myenteric plexus from an adult Wnt1-
cre:*Rosa26*-tdTomato mouse. Inset is magnified in color segregated panels, showing GFAP$^+$ glia
co-expressing tdTomato (red, red arrows). Nuclei are labeled with DAPI (blue). Scale bar = 10
μm.

(b) 2 dimensional representation of a 3 dimensional stack of images shows enteric neurons, la-
beled with Hu (green), in the myenteric plexus from an adult Wnt1-cre:*Rosa26*-tdTomato mouse.
Enteric neurons either express tdTomato (red arrow) or not (white arrows). Nuclei are labeled with
DAPI (blue). Scale bar = 10 μm.

(c) Orthogonal views of the image **(b)** shows the lack of tdTomato expression in a particular Hu-
immunolabeled enteric neuron.

(d) Quantification of tdTomato$^+$ and tdTomato$^-$ neurons in the myenteric ganglia of 6 P60 Wnt1-
cre:*Rosa26*-tdTomato mice. Data represents Mean ± SEM.

Fig 2: Mesoderm-lineage-tracing and marker expression provide evidence of the mesoder-mal-derivation of half of all adult small intestinal myenteric neurons.

(a) Small intestinal LM-MP from adult male Tek-cre:*Hprt*-tdTomato shows the presence of Tek-derived and hence mesoderm-derived tdTomato$^+$ (red) Hu$^+$ (green) neurons (white arrows) and non-mesoderm-derived tdTomato$^-$ neurons (green arrows). Vascular cells depict higher fluores-cence intensity (yellow arrow) than cells in the ganglia. Nuclei are labeled with DAPI. Scale bar = 10 µm.

Cardiac and small intestinal tissues from **(b)** heart and **(c)** small intestinal LM-MP layer of adult male Mesp1-cre:*Rosa26*-tdTomato mice shows the presence of variable tdTomato expression in cells of these tissues. While cardiomyocytes in heart and cells within the myenteric ganglia (white arrows) exhibit lower tdTomato fluorescence, similar to the tissues from Tek-cre:*Hprt*-tdTomato mice in (a), the vascular cells exhibit higher fluorescence intensity (yellow arrows). Scale bar = 10 µm.

(d) Small intestinal LM-MP from adult male Mesp1-cre:*Rosa26*-tdTomato mouse shows the pres-ence of mesoderm-derived tdTomato$^+$ (red) Hu$^+$ (green) neurons (white arrows) and non-meso-derm-derived tdTomato$^-$ neurons (green arrows). Nuclei are labeled with DAPI. Scale bar = 10 µm.

(e) Small intestinal LM-MP from adult male Mesp1-cre:*Rosa26*-EGFP mouse, when im-munostained with antibodies against GFP and against Hu, shows the presence of mesoderm-derived EGFP$^+$ (green) Hu$^+$ (red) neurons (white arrows) and non-mesoderm-derived EGFP$^-$ neurons (red arrows). Nuclei are labeled with DAPI. Scale bar = 10 µm.

(f) Quantification of tdTomato-expressing and non-expressing neurons in the myenteric ganglia of 3 P60 Mesp1-cre:*Rosa26*-tdTomato mice. Data represents Mean ± SEM.

MHCst immunolabeling exclusively marks all the **(g)** mesoderm-derived adult Hu$^+$ (green) neu-

rons in the small intestinal LM-MP of Mesp1-cre:*Rosa26*-tdTomato mice; and **(h)** all the non-NC-

derived neurons in the small intestinal LM-MP of Wnt1-cre:*Rosa26*-tdTomato mice (white arrows).

MHCst does not label tdTomato$^+$ NC-derived cells (yellow arrows). Nuclei are stained with DAPI

(blue). Scale bar = 10 μm.

Fig 3: Cellular and molecular phenotyping of MENs and NENs.

MET immunostaining (green) labels **(a)** Wnt1-cre:*Rosa26*-tdTomato⁻ MENs and **(b)** a population

of Mesp1-cre:*Rosa26*-tdTomato⁺ MENs (white arrow) while not labeling all MENs (red arrow).

(c) RET immunostaining (green) only labels Wnt1-cre:*Rosa26*-tdTomato⁺ NENs (green arrow)

and not tdTomato⁻ MENs (white arrows).

(d) NOS1 is expressed by both tdTomato⁺ NENs (red, white arrows) and tdTomato⁻ MENs (blue

arrows) in an adult Wnt1-cre:*Rosa26*-tdTomato mouse, but most MENs do not express NOS1

(marked by *).

(e) NEN lineage contains significantly higher proportions of NOS1⁺ neurons compared to MEN

lineage in Wnt1-cre:*Rosa26*-tdTomato mice. Data represent mean ± S.E.M. Student's t-test ** p

< 0.01.

(f) ChAT is expressed by both tdTomato⁺ NENs (red, white arrows) and tdTomato⁻ MENs (blue

arrows) in an adult Wnt1-cre:*Rosa26*-tdTomato mouse.

(g) NEN lineage contains significantly higher proportions of ChAT⁺ neurons compared to MEN

lineage in Wnt1-cre:*Rosa26*-tdTomato mice. Data represent mean ± S.E.M. Student's t-test *** p

< 0.001.

(h) Both tdTomato⁺ (red) and tdTomato⁻ neurons in the myenteric plexus of an adult Wnt1-

cre:*Rosa26*-tdTomato mouse (Hu, green) express CGRP (blue) Inset showing a tdTomato⁻

CGRP⁺ neuron (white arrow) is magnified on the right.

(i) MEN lineage contains significantly higher proportions of CGRP$^+$ neurons compared to NEN

lineage in Wnt1-cre:*Rosa26*-tdTomato male mice. Data represent mean ± S.E.M. Student's t-test

**** $p < 0.0001$.

Nuclei in **(b) and (c)** are labeled with DAPI (blue). Scale bar for all images denotes 10 μm.

Fig 4: scRNAseq-analyses identifies the distinct transcriptomic profile of the MENs.

(a) UMAP representation of 11,123 sequenced high-quality cells that were identified as meeting a 200 UMI minimum threshold with a mitochondrial read ratio of less than 20%. Clusters were annotated by markers that were found to be cell-type specific by searching UniProt, Allen Cell Atlas and Pubmed databases. Cells of the neural crest lineage were then identified as NENs by their expression of neural crest marker gene Ret and Neuroglia by their expression of Ncam1 and Sox10, or as MENs by co-expression of Calcb (CGRP), Cdh3, and Met genes.

(b) Visualization of expression of select MEN-specific and neuronal markers using quasi-violin plots. "Max value" represents the scale for the log-normalized expression of each gene.

(c) Validation of the MENs-specific marker genes discovered in the scRNAseq analyses by im-munohistochemistry and confocal microscopy of small intestinal LM-MP from adult male Wnt1-cre:*Rosa26*-tdTomato mice. Expression of neuronal markers STX3, VSNL1, STMN2, HAND2, GPR88, PDE10A, ELAVL2, and TUBB2B (Gray, green arrows) was found in tdTomato⁻ MENs. Immunostaining of the proteins AEBP1, CFTR, CLIC3, NT-3, SLPI, SMO, IL-18, SLC17A9, FMO2 and MYL7 (green; green arrows) was found to be localized to tdTomato⁻ MENs. tdTomato⁺ (red) NC-cells did not immunostain for these markers (red arrows). Panel also shows Wnt1-cre:*Rosa26*-tdTomato tissue with no primary controls (stained with AlexaFluor 488 and AlexaFluor 647 antibodies). Figures with * annotations are 2D representation of 3D stacks of images. Nuclei in some panels were labeled with DAPI (blue). Scale bar denotes 10 μm.

Fig 5: MENs express the mesenchymal gene Decorin and the ENS-specific gene Phox2b.

(a) Sparkline representation plot of the top 90 percentile of expressed genes in the various

scRNAseq cell clusters from the adult murine LM-MP tissue of a 6-month-old mouse shows that

the MENs cluster, which expresses the genes Cdh3, Pde10a, and Hand2, also expresses signif-

icant amounts of the mesenchymal gene Decorin (Dcn). Darker colors of the sparkline plot repre-

sent higher expression.

(b) Sparkline representation plot of the neural crest-derived enteric neurons from the Zeisel et al.

dataset shows that the Decorin gene is not found expressed by most neural crest-derived enteric

neurons.

(c) 2 dimensional representation views and orthogonal view of a myenteric ganglion from the

small intestinal LM-MP of adult male Wnt-cre:*Rosa26*-EGFP mouse, where tdTomato (red) is

expressed by neural crest-derived cells, when immunostained with antibodies against DECORIN

(grey) shows that the DECORIN-expressing myenteric cells (green arrows) do not express

tdTomato and hence are non-neural crest-derived MENs. Nuclei are labeled with DAPI. Scale bar

= 10 µm.

(d) Representative image of a myenteric ganglion from the small intestinal LM-MP of adult male

Wnt-cre:*Rosa26*-EGFP mouse, where tdTomato (red) is expressed by neural crest-derived cells,

when immunostained with antibodies against PHOX2b (green) shows that the PHOX2b-expres-

sion is found in myenteric cells that do not express tdTomato (green arrows, and hence are not

neural crest-derived cells, or MENs) as well as in tdTomato-expressing neural crest-derived NENs

(red arrow). Nuclei are labeled with DAPI. Scale bar = 10 µm.

(e) Volcano plot of gene expression profiles of Phox2b-expressing cells, which were sorted based

on their CFP expression level, shows that the expression of MENs marker genes Smo, Aebp1,

Cdh3, Fmo2, Il18, Slpi, Upk3b, Msln, and Hand2 is significantly enriched in the Phox2b-CFP^{low}

fraction. Red dotted line shows the $p_{adjusted}$ value of 0.05.

1380

Fig 6: Computational analyses using *projectR* identifies MENs in publicly available murine and human transcriptomic datasets.

(a) Heatmap of cell weights for the patterns learned by NMF (k=50). Hierarchical clustering was calculated using Euclidean distance. Multiple clusters annotated as the same cell type are merged. The four most specific MENs patterns (16, 27, 32 and 41) are selected.

(b) In addition to using the four MEN-specific NMF patterns to label the MENs cluster in our scRNAseq dataset (plots in top row), two other transcriptomic datasets: May-Zhang et al.'s murine ileal ENS snRNAseq dataset[50] (plots in second row), and Elmentaite et al.'s gut mesenchymal scRNAseq dataset[78] (plots in third row), were projected into the defined four MEN-specific pattern space using *ProjectR*. Raw projected cell weights were visualized using previously learned UMAP embedding. The cell clusters that show high pattern usage are shown bounded by the red square. Plots in the final row shows projection of healthy post-natal mesenchymal cells from Elmentaite et al's data[78] into the four MEN-specific NMF patterns, which again show a population of cells showing high MEN-specific pattern utilization.

 Fig 7. Detection of MENs in human single cell RNA sequencing data.

(a) UMAP representation of scRNAseq data from Elmentaite et al.'s healthy post-natal mesen-

chymal cells from the human gut[78].

(b) Breakdown for previously published annotated features of the post-natal health subset of the

gut cell atlas as presented in Elmentaite et al.[78].

(c) Expression of the MENs marker gene Dcn in the UMAP representation of healthy post-natal

data from Elmentaite et al.[78]., which was annotated as clusters of mesenchymal cells.

(d) Expression of neuronal marker genes Slc17a9, Stx3, Uchl1, Tubb2b, Pde10a, and Hand2 in

the UMAP representation of healthy post-natal data from Elmentaite et al.[78], which was annotated

as clusters of mesenchymal cells.

Fig 8: Validation of SNAP-25 as a NENs marker

(a) SNAP-25 expression is restricted to the neural crest lineage in the adult myenteric ganglia. SNAP-25 expression (green) co-localizes with tdTomato (red) but not with the MENs marker MHCst (cyan) as observed in 2D and orthogonal views of a myenteric ganglia from a Wnt1-cre:*Rosa26*-tdTomato mouse that was immunolabeled with antibodies against MHCst and SNAP-25. Nuclei were labeled with DAPI (blue). Scale bar denotes 10 μm.

(b) In the adult male Snap25-Gcamp knock-in mice, the expression of Snap25-Gcamp/GFP (green) is restricted to a subset of Hu-expressing (red) myenteric neurons in the adult murine small intestinal myenteric ganglia (yellow arrows), while many neurons (red arrows) do not show any detectable expression of Snap25. Nuclei were labeled with DAPI (blue). Scale bar denotes 10 μm.

(c) Orthogonal views of z-stack of an image of the myenteric ganglion from the small intestinal LM-MP tissue from an adult male Snap25-Gcamp knock-in mice shows that the expression of Snap25-Gcamp/GFP (green) is exclusive of the expression of the MENs marker MHCst (red). Nuclei were labeled with DAPI (blue). Scale bar denotes 10 μm.

Fig 9: Observation of MENs marker expression in the adult human myenteric ganglia

(a) Hu-expressing small intestinal myenteric neurons (green) from the normal human duodenal tissue, when immunolabeled with antibodies against the MENs marker DECORIN (red) identifies putative human MENs (red arrows) and NENs (green arrows) by presence or absence of DECORIN immunolabeling. Scale bar denotes 10 μm.

(a) Hu-expressing small intestinal myenteric neurons (green) from the normal human duodenal tissue, when immunolabeled with antibodies against the MENs marker MHCst (red) and MET (blue) identifies putative human MENs (red arrows) and NENs (green arrows) by presence or absence of these MENs markers. Scale bar denotes 10 μm.

(c) Immunolabeling myenteric tissue with antibodies against the MENs markers SLC17A9 (green, green arrows) and MHCst (red) shows SLC17A9 expression in a subset of MHCst-expressing (red) neurons. Subset of MHCst-expressing cells do not express SLC17A9 (red arrow). Nuclei are labeled with DAPI (blue) in the normal human duodenal tissue. Scale bar denotes 10 μm.

Fig 10: GDNF and HGF signaling regulate age-dependent changes in NENs and MENs proportions.

(a, b) Immunostaining myenteric plexus tissue from juvenile and mature Wnt1-cre:*Rosa26*-tdTomato mice with antibodies against the pan-neuronal marker Hu (green).

(c) Age-associated loss of NENs and gain of MENs in the small intestinal LM-MP of maturing and aging Wnt1-cre:*Rosa26*-tdTomato mice. Data represent mean ± S.E.M.

(d) Mean ± SEM of the four MEN-specific pattern weights in the human mesenchymal cell populations from Elmentaite et al.[78], wherein data from age ranges of 4 – 12 years was clubbed together as Juvenile (Juv), data from age ranges of 20 – 50 years was clubbed together as Adult (Ad), and data from age ranges of 60 – 75 years was clubbed together as Aged (Ag). Every datapoint refers to mean projected pattern weight for cells within a defined age or age range, where Juv data comprise of ages 4, 6, 9, 10, 12 years; Ad data comprise of age ranges 20 - 25, 25 – 30, and 45-50 years; and finally Ag data comprise of age ranges 60 – 65, 65 – 70, and 70 – 75 years. One-way ANOVA ** = $p < 0.01$).

(e) Western blot analyses of GDNF (green) and the house-keeping protein β-actin (red) expression in LM-MP tissues from mice of ages P10, P30, and P90. (n = 3 mice per group; each sample is a biological replicate). Fluorescent intensities of the two bands of GDNF (that correspond to ~25 kD bands of protein marker) were quantified together. The lower band of GDNF is present only in the P10 tissues and disappears in P30 and P90 adult murine tissues.

(f) Western blot analyses of HGF (green) and the house-keeping protein β-Actin (red) expression in LM-MP tissues from mice of ages P10, P30, and P90. (n = 3 mice per group; each sample is a biological replicate). Fluorescent intensities of the two bands of HGF (that are between 50 kD and 37 kD bands of the protein marker) were quantified together.

(g) The normalized fluorescent intensity of GDNF protein to house-keeping protein β-Actin com-

pared between the three age groups. GDNF presence was highest in P10 group and was signifi-

cantly reduced in P30 and P90 groups. Data represent mean ± S.E.M. One-way ANOVA **** p <

0.001.

(h) Age-dependent decrease in Gdnf mRNA transcript expression (normalized to the house-keep-

ing gene Hprt) in the myenteric plexuses of P30 and P90 old mice. Data represent mean ± S.E.M.

Student's t-test *p < 0.05.

(i) The normalized fluorescent intensity of HGF protein to house-keeping protein β-Actin was

compared between the three age groups. HGF expression significantly increased from P10

through P90. Data represent mean ± S.E.M. One-way ANOVA *p < 0.05.

(j) Age-dependent increase in Hgf mRNA transcript expression (normalized to the house-keeping

gene Hprt) in the myenteric plexuses of P10, P30, and P90 old mice. Data represent mean ±

S.E.M. One-way ANOVA * p < 0.05, *** p < 0.01, **** p < 0.001.

(k) Percent proportions of tdTomato⁻ MENs and mean Hu⁺ neurons/ganglia in LM-MP of cohorts

of Wnt1-cre:*Rosa26*-tdTomato mice that were dosed with GDNF or Saline from P10 to P20 age.

Data represent mean ± S.E.M. Student's t-test ** p < 0.01.

(l) Percent proportions of tdTomato⁻ MENs and mean Hu⁺ neurons/ganglia in LM-MP of cohorts

of Wnt1-cre:*Rosa26*-tdTomato mice that were dosed with HGF or Saline from P10 to P20 age.

Data represent mean ± S.E.M. Student's t-test * p < 0.05.

Fig 11: Ect2-expression labels a population of cycling MENs

(a) UMAP representation of the cells of the MENs, NENs, and Neuroglia clusters from scRNAseq of cells from the small intestinal LM-MP tissue of two post-natal day 21 (P21) mice.

(b) Tricycle analyses of ENS cell-types with their cell cycle positions. The continuous cell cycle position (theta) is measured as the angle from the origin. Anti-clockwise representation of cells that are represented between 1.5π and 0.5π space on the embedding are increasingly present within the quiescent G1/G0 cell cycle phase. In contrast, cells that are represented between the 0.5π and 1π space on the embedding are present within the S phase, and those within the 1π and 1.5π space are present within the G2M phase.

(c) Cells within the P21 MENs cluster were projected into Tricycle software. 213 of the 504 P21 MENs were present between the 0.5π and 1.5π space on the embedding and hence were inferred to be undergoing cell cycling.

(d, e) Expression of cell-cycle correlated genes Top2a and Ect2 in MENs shows significant expression of these two genes in cells of the P21 MENs cluster that are present between the values of 0.5π and 1.5π in the Tricycle embedding. Loess curve fittings (black) represent the dynamics of gene expression over the cell cycle (theta).

Orthogonal views of small intestinal myenteric ganglia from a P21 wildtype mouse when immunostained with antibodies against ECT (green) and against MENs markers **(f)** DECORIN (red) and **(g)** MHCst (red), shows that ECT2 is expressed in a subset of MENs at this age. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 μm.

Fig 12: Reduced RET signaling accelerates ENS aging to cause pathology

(a) Hu immunostaining (green) LM-MP tissues from 16-week-old Ret$^{+/CFP}$ (Ret$^{+/-}$) mouse shows

mutually exclusive expression of Ret-CFP (cyan, green arrow) and MHCst (red, red arrow) MENs.

Nuclei are stained with DAPI (blue). Scale bar = 10 µm.

(b) Quantification of Ret-CFP$^+$ neurons from 9- and 16-week-old Ret$^{+/-}$ mice show age-associated

loss of Ret-CFP$^+$ neurons. Data represent mean ± S.E.M. Student's t-test * p < 0.05.

(c) Quantification of MHCst$^+$ MENs shows significant increase in their proportions in Ret$^{+/-}$ mice

but not in Ret$^{+/+}$ mice with age. Data represent mean ± S.E.M. One way ANOVA * p < 0.05, ** p

< 0.01, *** p < 0.001).

(d) Measures of whole gut transit time (WGTT) in cohorts of Ret$^{+/-}$ and Ret$^{+/+}$ mice MENs show

significant slowing of whole gut transit of Ret$^{+/-}$ but not Ret$^{+/+}$ mice with age. Data represent mean

± S.E.M. One way ANOVA * = p < 0.05.

Fig 13: GDNF normalizes altered intestinal motility by increasing NENs proportions in the aging gut.

(a) Measures of whole gut transit time (WGTT) in GDNF (treated with GDNF) and Control (treated with Saline) cohorts of 17-month-old mice taken before the start of treatments and after the end of 10 consecutive days of treatment shows that the two groups are matched in their transit times before treatment, but GDNF treatment significant decreases average transit times when compared to the control cohort. Data represent mean ± S.E.M. Student's t-test *** = $p < 0.001$.

(b) Quantification of percent MHCst$^+$ MENs per Hu-labeled neurons in myenteric ganglia in the GDNF and Control cohorts shows significant decrease in their proportions in GDNF-treated cohort but not in saline-treated control cohort. Data represent mean ± S.E.M. Student's t-test ** $p < 0.01$.

(c) Quantification of numbers of RET$^+$ NENs per 40X field views of myenteric ganglia shows significant increase in their numbers in GDNF cohort mice when compared with Control cohort mice. Data represent mean ± S.E.M. Student's t-test * $p < 0.05$.

Fig 14: Patients with chronic gut dysmotility show significant shifts in their normal proportions of the two neuronal lineages.

(a) ProjectR-based projection of bulkRNAseq data from intestinal specimens of patients with normal motility and patients with obstructive defecation (OD), a chronic condition of intestinal dysmotility, into the 50 different NMF patterns learnt earlier, shows that the MEN-specific NMF patterns 32 and 41 were significantly upregulated in bulk-RNAseq of OD patients compared to controls. Data represent mean ± S.E.M. One way ANOVA. Data mined from raw data generated by Kim et al.[100]

(b) OD patients show a significant decrease in the expression of important NENs-associated genes such as Ret, Gdnf, Snap-25, Nos1, Klhl1, and Chat, while showing a significant increase in the expression of important MENs-specific genes such as Clic3, Upk3a, Cdh3, Slpi, and Slc17a9. Data taken from Kim et al.[100]

Suppl Figure 1: Presence and absence of tdTomato reporter expression in the myenteric ganglia of adult neural crest lineage-traced mice

(a) Hu-immunostained Wnt1-cre:*Rosa26*-tdTomato LM-MP tissue, where tdTomato-expression (red) marks neural crest-derived cells, and Hu (green) labels myenteric neurons, we observed the presence of higher intensity fluorescence (red arrows) and lower intensity fluorescence (orange arrows) in various tdTomato$^+$ neurons. The myenteric ganglia also contained Hu$^+$ neurons that did not express tdTomato (green arrow) and these neurons alone were deemed to be non-neural crest-derived neurons. Scale bar denotes 10 μm.

(b) Reporter non-expressing Hu-immunolabeled neurons (green, green arrows) can be observed both through the expression of the tdTomato reporter (red), as well as on antibody labeling of the tdTomato reporter (gray).

(c) Freshly harvested small intestinal tissue from Wnt1-cre:*Rosa26*-tdTomato mice when imaged under 20X magnification of a live tissue fluorescence microscope shows the presence of cells that express tdTomato (red) and those that don't.

(d) Cells showing hyper-aggregation of the tdTomato reporter can be found in freshly harvested full-thickness small intestinal myenteric plexus in tissues imaged under 20X magnification of a live tissue fluorescence microscope.

(e) Myenteric ganglia from fixed LM-MP tissue of adult Wnt1-cre:*Rosa26*-tdTomato mice show show hyper-aggregation of tdTomato (white arrow) when imaged using a confocal microscope. Nuclei are stained with nuclear marker DAPI (blue). Scale bar denotes 10 μm.

(f) Cells showing hyper-aggregation of tdTomato (red, red arrow) show no detectable nuclear labeling with DAPI (blue) or of pan-neuronal marker Hu (green, green arrow). Scale bar denotes 10 μm.

In addition, interrogation of two other neural crest-specific lineage-traced mouse lines confirms our observations with Wnt1-cre:*Rosa26*-tdTomato mice. **(g)** In Wnt1-cre:*Hprt*-tdTomato adult male mice, tdTomato expression (red) is present only in a subset of Hu-immunolabeled (green) neurons (red arrows) while it is not detected in other Hu-expressing neurons (green). Similarly, **(h)** only a subpopulation of Hu⁺ adult enteric neurons (green) is derived from the neural crest, as seen by its co-expression of tdTomato (red, red arrows) in the Pax3-cre:*Rosa26*-tdTomato mice while other neurons that do not express tdTomato (white arrows) are inferred to be not derived from the neural crest. Scale bar denotes 10 μm.

(i) Orthogonal views of small intestinal villus from a Wnt1-cre:*Rosa26*-tdTomato mouse that shows presence of tdTomato-expressing (red) neural crest-derived fibers inside the villus but a complete lack of tdTomato-expression by intestinal epithelial cells. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 μm.

(j) Representational image of the myenteric plexus from a *Rosa26*-tdTomato mouse without any cre driver showing that in absence of cre recombinase to drive tdTomato expression, there is a lack of tdTomato expression in this layer. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 μm.

(k) Representational image of the myenteric plexus from a *Rosa26*-EGFP mouse without any cre driver showing that in absence of cre recombinase to drive EGFP expression, there is a lack of EGFP expression in this layer. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 μm.

(l) Western blot image showing detection of two different loading amounts of the recombinant HuB protein by the rabbit polyclonal antibody against HuC/D (abcam). Molecular Weight marker with defined band sizes indicated on the right.

Suppl Figure 2: Tek-expressing and Mesp1-expressing mesodermal lineage contributes to adult enteric neurons.

(a) Adult small intestinal LM-MP layer from a male Tek-EGFP mouse, where EGFP expression (green) labels Tek-expressing cells was immunostained with specific antibodies against the pan-neuronal marker PGP9.5 (red) and imaged under a confocal microscope. The Tek-EGFP expression did not label any cells within the myenteric plexus marked by PGP9.5 expression showing that adult small intestinal myenteric neurons did not express Tek. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 μm.

(b) We observed the presence of higher intensity fluorescence (red arrows) and lower intensity fluorescence (green arrow) in various tdTomato$^+$ cells in a section of heart tissue from a 6-month-old Mesp1-cre:*Rosa26*-tdTomato adult male mouse, where tdTomato (red) labels Mesp1-derived cells of the mesodermal lineage that includes cardiomyocytes and vasculature. Nuclei are stained by DAPI (blue). Scale bar denotes 10 μm.

(c) Adult small intestinal LM-MP layer from a 6-month-old male Mesp1-cre:*Rosa26*-tdTomato mouse, where tdTomato (red) cells are mesoderm-derived cells, was immunostained with specific antibodies against the pan-neuronal marker Hu (green) and imaged under a confocal microscope. tdTomato-expressing cells outside the ganglia (red arrows) were brighter than tdTomato+ neurons inside the ganglia (green arrows). Nuclei are labeled with DAPI (blue). Scale bar denotes 10 μm.

(d) Adult small intestinal LM-MP layer from a 2-month old male Mesp1-cre:Rosa26-EGFP mouse, where EGFP cells are mesoderm-derived, was immunostained with specific antibodies against the reporter EGFP (green) and the pan-neuronal marker Hu (red) and imaged under a confocal microscope. A population of Hu-immunolabeled neurons were also labeled with antibodies against EGFP (green arrows) while other neurons were not (red arrows), suggesting the

derivation of EGFP$^+$ neurons from the Mesp1-expressing mesoderm. **(e)** No primary control for

the Anti-GFP immunostaining showed very weak EGFP expression in the fixed LM-MP tissue

from these animals. Nuclei were labeled with DAPI (blue). Scale bar = 10 µm. Consistent with the

observations made with the tdTomato reporter in this transgenic line, the Mesp1-cre:*Rosa26*-

EGFP animals also show a brighter reporter signal in the vasculature and a dimmer reporter signal

in the myenteric neurons.

(f) MHCst$^+$ (cyan) labels all tdTomato$^+$ (red) Hu$^+$ (green) adult neurons (white arrows) but not the

tdTomato$^-$ neurons (yellow arrow) in the mesoderm-specific Tek-cre:*Hprt*-tdTomato lineage-

traced lines. Nuclei are stained with DAPI (blue). Scale bar = 10 µm.

Suppl Figure 3: Validation of MENs-specific markers by immunohistochemistry and con-

focal microscopy.

Small intestinal LM-MP from several adult 5-month-old male Wnt1-cre:*Rosa26*-tdTomato mice,

where tdTomato-expression (red) labels neural crest-derived cells, was used to ascertain and

validate the expression of the novel MENs marker and of known neuronal marker genes.

(a) MHCst-specific antibody S46 (green) was found to label tdTomato non-expressing cells within

the myenteric ganglia as well as cells outside of the myenteric ganglia. Neural crest-derived cells

within the myenteric ganglia expressed tdTomato. Nuclei are labeled with DAPI (blue). Scale bar

denotes 10 µm.

The immunostaining of the MENs-markers elucidated through scRNAseq analyses: **(b)** CDH3, **(c)**

AEBP1, **(d)** SLPI, **(e)** CFTR, **(f)** NT-3, **(g)** CLIC3, **(h)** FMO2, **(i)** IL-18, **(j)** MYL-7, and **(k)** SLC17A9

(all immunolabeled cells in these panels are green; green arrows) was found to be localized to or

enriched in cells without tdTomato expression, while the tdTomato$^+$ cells were found to show not

to express these MENs-specific markers (red arrow). In addition, we also immunolabeled the

myenteric ganglia from this transgenic mouse line with known neuronal markers (gray) and with

antibodies against the pan-neuronal marker Hu (green) to show that the neuronal markers Elavl2

or HuB **(l)**, GPR88 **(m)**, HAND2 **(n)**, PDE10A **(o)**, TUBB2B **(p)** and VSNL1 **(q)** were expressed

by both tdTomato$^+$ (red arrow) and tdTomato$^-$ Hu$^+$ neurons. Orthogonal views of images where

3D image z-stacks were obtained are provided for better visualization of marker expression in the

3D space. Nuclei in these tissues were labeled with DAPI (blue). Scale bar denotes 10 µm.

Suppl Figure 4: scRNAseq metrics

(a) MENs are significantly larger in size as compared to the NENs, as the mean Feret's Diameter of the Wnt1-cre:*Rosa26*-tdTomato⁻ MENs is significantly more than that of the Wnt1-cre:tdTomato⁺ NENs. Student's t-test; ** p value < 0.01.

(b, c) UMAP representation of sequenced cells generated from 10X Genomics Chromium-derived libraries from the two mouse samples, taken from separate littermate adult male mice, shows similar representation of their cells across the various clusters.

(d) Quasiviolin plot of expressed genes provide a comparison of important MEN-specific and neuronal genes between the NENs and MENs clusters in the adult ileal May-Zhang et al. dataset and the NENs, Neuroglial, and MENs clusters in our scRNAseq dataset.

(e) Top genes that are pattern drivers for the four MEN-specific NMF patterns along with their gene weights.

(f) The four MEN-specific patterns were together used to label a group of cells (True) that together showed increased usage of the MEN-specific patterns, when compared to other cell types (False) in the adult ileal snRNAseq dataset from May-Zhang et al.[50].

(g) Top-expressed genes in the group of cells **(f)** that showed higher utilization of the MEN-specific patterns, compared to other cells in the dataset.

(h) Representation of the four MEN-specific NMF patterns in the UMAP of our data and the projection of the May-Zhang et al.[50] murine ileal dataset and the entire gut mesenchymal single cell RNA seq dataset by Elmentaite et al.[78] into the MEN-specific patterns highlights cell clusters that show higher usage of the four NMF patterns. Bottom row shows increasing usage of these patterns with age in the entire mesenchymal gut cell atlas data.

 Suppl Figure 5: Expression of neuronally significant genes by MENs

Sparklines plot of gene expression in various cell clusters, as revealed by single cell transcriptom-

ics of diverse LM-MP cells from small intestinal tissue of two 6-month-old mice, show that expres-

sion of some genes is enriched in the MENs cluster, when compared to the other major clusters

of cells in the dataset. Lower to higher expression in the plot is represented by yellow to blue

gradient. Subsequent analyses shows that proteins coded by these genes are neuronally signifi-

cant.

1662

Suppl Figure 6: Expression of mesodermal marker MYH11, and reporter expression under the Baf53b-cre transgenic line in post-natal myenteric neurons

(a) Orthogonal views of a myenteric ganglion of an adult Wnt1-cre:*Rosa26*-tdTomato mouse, where tdTomato (red) is expressed by neural crest-derived cells, shows that immunolabeling with a MYH11-specific antibody (green) labels tdTomato non-expressing cells. Nuclei are labeled with DAPI (blue). Scale bar denotes 10 µm.

(b) Representative image of the myenteric plexus layer from a Myh11-cre[ERT2]:*Rosa26*-YFP adult male mouse, which was dosed with tamoxifen for two consecutive weeks, and which when immunostained with antibodies against YFP/GFP (green) and the pan-neuronal marker Hu (red), shows the presence of YFP-expressing neurons (green arrows) in the myenteric ganglia that also contains YFP negative neurons (red arrow). Nuclei are labeled with DAPI (blue). Scale bar denotes 10 µm.

(c) Immunostaining of healthy human adult duodenal tissue with antibodies against the pan-neuronal marker PGP9.5 (green) and MYH11 (red), shows that in the circular muscularis layer of the tissue, MYH11 labels the smooth muscle and not the PGP9.5 immunostained nerve fibers and cell bodies. Scale bar denotes 10 µm.

(d) Immunostaining of healthy human adult duodenal tissue with antibodies against the pan-neuronal marker PGP9.5 (green) and MYH11 (red), shows that in the myenteric ganglia, MYH11 immunolabels a subset of PGP9.5-expressing myenteric neurons (white arrows). Scale bar denotes 10 µm.

(e) Representative image of a myenteric ganglion from a Baf53b-cre:*Rosa26*-tdTomato adult male mouse, where tdTomato (red) is expressed by Baf53b expressing cells or those derived from Baf53-expressing cells, when immunostained with antibodies against the pan-neuronal marker Hu (green), shows the presence of neurons that express tdTomato (white arrows) along with

neurons that do not express tdTomato (green arrows). Nuclei are labeled with DAPI (blue). Scale

bar denotes 10 μm.

(f) Graphical representation of the proportions of tdTomato-expressing Hu$^+$ myenteric neurons in

the LM-MP tissue of two male P21 Baf53b-cre:*Rosa26*-tdTomato mice shows that myenteric gan-

glia from ileum, proximal colon, and distal colon contain significant proportions of neurons that do

not express tdTomato under the Baf53b-cre transgenic line.

1693

Suppl Figure 7: Putative mesoderm-derived enteric neurons (MENs) are present in adult human myenteric ganglia.

(a, b) The mesodermal specific markers, MET and MHCst are expressed by a subpopulation of human enteric neurons. Myenteric ganglia from normal duodenal tissue of 2 different adult human subjects express the pan-neuronal marker Hu (green), with distinct subpopulations positive and negative for MET (blue) and MHCst (red). MET and MHCst-expressing neurons (red arrows) in the human ENS are assumed to be mesoderm-derived enteric neurons (MENs) and those not expressing these MENs markers (green arrows) are presumably NENs. Nuclei are labeled with DAPI (gray). Scale bar = 10 µm.

(c) The mesodermal specific marker DECORIN is expressed by a subpopulation of human enteric neurons. Myenteric ganglia from normal human duodenal tissue when immunostained with anti-bodies against the pan-neuronal marker Hu (green) and DECORIN (red) shows the presence of DECORIN-expressing myenteric neurons, which are assumed to be MENs. Nuclei are labeled with DAPI (blue). Scale bar = 10 µm.

(d) Distribution of the various mesenchymal clusters of the post-natal healthy gut cell atlas data (generated by Elmentaite et al.[78]) by donor age. For the age of each donor, the proportion of cells annotated as belonging to each cluster is shown. Proportions of Cluster 1 are observed to expand with age.

1712 **Suppl Fig 8: Sub-clustering MENs in the scRNAseq dataset generated from 6-month-old**

1713 **mice.**

1714 **(a)** Sub-clustering the MENs cluster from our scRNAseq dataset into 5 distinct clusters **(b)** was

1715 found to be not based on the expression of cluster-specific marker genes.

1716

Suppl Fig 9: HGF and GDNF ratios in maturing and adult murine gut.

(a) Ratio of HGF:GDNF values in fluorescence intensities of their western blots (normalized to

their own house-keeping protein Beta Actin) from the small intestinal LM-MP of post-natal mice

at different ages during their maturation.

(b) Ratio of HGF:GDNF expression values from the full thickness small intestinal tissue of adult

mice taken from the Tabula muris database[91].

(c) Analyses of human plasma proteome from 1,025 individuals was performed by Sathyan et

al.[92] and the correlation of various plasma proteomes with aging was calculated. Of the genes

that showed significant correlation with age ($p < 0.05$), we found that the HGF levels in human

plasma correlated positively with age (above 0), while the levels of GDNF and RET correlated

negatively with age (below 0).

Suppl Fig 10: Identification of the MENs cluster in the P21 scRNAseq data and expression of Ret in NENs and in the Ret heterozygous mice.

(a) UMAP representation of all the cell clusters generated from scRNAseq performed on all cells from the P21 murine LM-MP.

(b) Projection of scRNAseq from the P21 murine LM-MP cells into the four MEN-specific NMF patterns generated using scRNAseq on 6-month-old murine LM-MP cells shows higher usage of the four NMF patterns in the putative MENs cluster in the P21 dataset (black arrow).

(c) Expression of the genes Ret and Met in the MENs and NENs clusters in the scRNAseq of P21 LM-MP cells shows that while Met-expression is detected in both MENs and a subset of NENs, the MET-expressing NENs subset shows significant expression of Ret, while the MENs do not express Ret.

(d) Sparklines plot of gene expression of various neuronally significant genes in the scRNAseq data from Zeisel et al., which was generated using flow sorted Wnt1-cre:*Rosa26*-tdTomato$^+$ cells from a P21 mouse, shows that Met expression is observed in the clusters ENT6, 8, and 9 which also show expression of CGRP-expressing gene Calcb, the neural crest-specific gene Ret, Choline acetyltransferase-expressing gene Chat and the pan-neuronal marker PGP9.5-expressing gene Uchl1.

(e) Fold change in transcript abundance of the gene Ret in the small intestinal LM-MP tissue from Ret wildtype (Ret WT) and Ret heterozygous (Ret Het) littermate adult mice shows that Ret transcripts are significantly less abundant in the tissue of Ret Het mice, when compared to those from Ret WT mice. Ret transcript expression in every sample was normalized to the expression of the house keeping gene Hprt. Student's t-test; * p value < 0.05.

Suppl Fig 11: Effect of GDNF treatment on MHCst-expressing MENs and RET-expressing NENs in aging mice

Representative image of MHCst (red) and Hu (green)-immunostained myenteric ganglia of **(a)** saline-treated control and **(b)**. GDNF-treated 17-month-old mice show MHCst$^+$ (yellow arrows) and MHCst$^-$ neurons. Note the reduction in MHCst-immunostaining on inter-ganglionic fibers in the 17-month-old mice when treated with GDNF, compared to control mice. Nuclei are stained with DAPI (blue). Scale bar indicates 10 µm.

Representative image of RET-immunostained (green) myenteric ganglia of **(c)**. saline-treated control and **(d)**. GDNF-treated 17-month-old mice. Nuclei are stained with DAPI (blue). Scale bar indicates 10 µm.

(e) Western blot image of a 4-20% gel performed using the RET antibody on total protein lysate from small intestinal LM-MP layer from post-natal mice shows the expected band between150 and 250 kDa of RET protein.

Suppl Fig 12: Projection of the bulk RNA sequencing data of intestinal tissue from Control and Patients with Obstructed Defecation into our murine scRNAseq-derived NMF patterns using *projectR*.

Transcriptomic data from intestinal tissues of patients with normal intestinal motility and those with obstructed defecation (OD) was procured from GEO (GSE101968). Non-negative matrix factorization (NMF), as implemented in the R package *NNLM* (https://github.com/linxihui/NNLM), was performed on the murine scRNAseq data using k=50 and default parameters; and cell weights for each pattern were grouped. Using projectR, the log2 expression (log2(rpkm + 1)) from the human bulk RNA sequencing data from control and OD patients were projected into the murine scRNAseq-derived NMF patterns. The mean projection weights from Control and OD groups were tested for statistically significant differences using Students' t tests.

References

References

1   Kulkarni, S. *et al.* Advances in Enteric Neurobiology: The "Brain" in the Gut in Health and Disease.
J Neurosci **38**, 9346-9354, doi:10.1523/JNEUROSCI.1663-18.2018 (2018).
2   Jarret, A. *et al.* Enteric Nervous System-Derived IL-18 Orchestrates Mucosal Barrier Immunity. *Cell*
180, 50-63 e12, doi:10.1016/j.cell.2019.12.016 (2020).
3   Uesaka, T., Young, H. M., Pachnis, V. & Enomoto, H. Development of the intrinsic and extrinsic
innervation of the gut. *Dev Biol* **417**, 158-167, doi:10.1016/j.ydbio.2016.04.016 (2016).
4   Obermayr, F., Hotta, R., Enomoto, H. & Young, H. M. Development and developmental disorders
of the enteric nervous system. *Nat Rev Gastroenterol Hepatol* **10**, 43-57,
doi:10.1038/nrgastro.2012.234 (2013).
5   Hao, M. M. & Young, H. M. Development of enteric neuron diversity. *J Cell Mol Med* **13**, 1193-
1210, doi:10.1111/j.1582-4934.2009.00813.x (2009).
6   Anderson, R. B., Newgreen, D. F. & Young, H. M. Neural crest and the development of the enteric
nervous system. *Adv Exp Med Biol* **589**, 181-196, doi:10.1007/978-0-387-46954-6_11 (2006).
7   Young, H. M. & Newgreen, D. Enteric neural crest-derived cells: origin, identification, migration,
and differentiation. *Anat Rec* **262**, 1-15, doi:10.1002/1097-0185(20010101)262:1<1::AID-
AR1006>3.0.CO;2-2 (2001).
8   Young, H. M., Hearn, C. J. & Newgreen, D. F. Embryology and development of the enteric nervous
system. *Gut* **47 Suppl 4**, iv12-14; discussion iv26, doi:10.1136/gut.47.suppl_4.iv12 (2000).
9   Bergner, A. J. *et al.* Birthdating of myenteric neuron subtypes in the small intestine of the mouse.
J Comp Neurol **522**, 514-527, doi:10.1002/cne.23423 (2014).
10 Uesaka, T., Nagashimada, M. & Enomoto, H. Neuronal Differentiation in Schwann Cell Lineage
Underlies Postnatal Neurogenesis in the Enteric Nervous System. *J Neurosci* **35**, 9879-9888,
doi:10.1523/JNEUROSCI.1239-15.2015 (2015).
11 Espinosa-Medina, I. *et al.* Dual origin of enteric neurons in vagal Schwann cell precursors and the
sympathetic neural crest. *Proc Natl Acad Sci U S A* **114**, 11980-11985,
doi:10.1073/pnas.1710308114 (2017).
12 Burns, A. J. Migration of neural crest-derived enteric nervous system precursor cells to and within
the gastrointestinal tract. *Int J Dev Biol* **49**, 143-150, doi:10.1387/ijdb.041935ab (2005).
13 Laranjeira, C. *et al.* Glial cells in the mouse enteric nervous system can undergo neurogenesis in
response to injury. *J Clin Invest* **121**, 3412-3424, doi:10.1172/JCI58200 (2011).
14 Brokhman, I. *et al.* Dual embryonic origin of the mammalian enteric nervous system. *Dev Biol* **445**,
256-270, doi:10.1016/j.ydbio.2018.11.014 (2019).
15 Sohal, G. S., Ali, M. M. & Farooqui, F. A. A second source of precursor cells for the developing
enteric nervous system and interstitial cells of Cajal. *Int J Dev Neurosci* **20**, 619-626 (2002).
16 Habeck, M. VENT cells--a load of hot air? *Drug Discov Today* **8**, 332-333 (2003).
17 Saffrey, M. J. Cellular changes in the enteric nervous system during ageing. *Dev Biol* **382**, 344-355,
doi:10.1016/j.ydbio.2013.03.015 (2013).
18 Becker, L., Kulkarni, S., Tiwari, G., Micci, M. A. & Pasricha, P. J. Divergent fate and origin of
neurosphere-like bodies from different layers of the gut. *Am J Physiol Gastrointest Liver Physiol*
302, G958-965, doi:10.1152/ajpgi.00511.2011 (2012).
19 Kulkarni, S. *et al.* Adult enteric nervous system in health is maintained by a dynamic balance
between neuronal apoptosis and neurogenesis. *Proc Natl Acad Sci U S A* **114**, E3709-E3718,
doi:10.1073/pnas.1619406114 (2017).

20   Antal, M., Beneduce, B. M. & Regehr, W. G. The substantia nigra conveys target-dependent
excitatory and inhibitory outputs from the basal ganglia to the thalamus. *J Neurosci* **34**, 8032-
8042, doi:10.1523/JNEUROSCI.0236-14.2014 (2014).
21   Freyer, L., Aggarwal, V. & Morrow, B. E. Dual embryonic origin of the mammalian otic vesicle
forming the inner ear. *Development* **138**, 5403-5414, doi:10.1242/dev.069849 (2011).
22   Wright, C. M. *et al.* scRNA-Seq Reveals New Enteric Nervous System Roles for GDNF, NRTN, and
TBX3. *Cell Mol Gastroenterol Hepatol* **11**, 1548-1592 e1541, doi:10.1016/j.jcmgh.2020.12.014
(2021).
23   King, P. H., Redden, D., Palmgren, J. S., Nabors, L. B. & Lennon, V. A. Hu Antigen Specificities of
ANNA-I Autoantibodies in Paraneoplastic Neurological Disease. *Journal of Autoimmunity* **13**, 435-
443, doi:https://doi.org/10.1006/jaut.1999.0337 (1999).
24   Honig, G., Liou, A., Berger, M., German, M. S. & Tecott, L. H. Precise pattern of recombination in
serotonergic and hypothalamic neurons in a Pdx1-cre transgenic mouse line. *J Biomed Sci* **17**, 82,
doi:10.1186/1423-0127-17-82 (2010).
25   Bardot, E. *et al.* Foxa2 identifies a cardiac progenitor population with ventricular differentiation
potential. *Nat Commun* **8**, 14428, doi:10.1038/ncomms14428 (2017).
26   Burtscher, I., Barkey, W., Schwarzfischer, M., Theis, F. J. & Lickert, H. The Sox17-mCherry fusion
mouse line allows visualization of endoderm and vascular endothelial development. *genesis* **50**,
496-505, doi:10.1002/dvg.20829 (2012).
27   Wilm, B., Ipenberg, A., Hastie, N. D., Burch, J. B. & Bader, D. M. The serosal mesothelium is a major
source of smooth muscle cells of the gut vasculature. *Development* **132**, 5317-5328,
doi:10.1242/dev.02141 (2005).
28   Kawaguchi, M., Bader, D. M. & Wilm, B. Serosal mesothelium retains vasculogenic potential. *Dev
Dyn* **236**, 2973-2979, doi:10.1002/dvdy.21334 (2007).
29   Herrick, S. E. & Mutsaers, S. E. Mesothelial progenitor cells and their potential in tissue
engineering. *Int J Biochem Cell Biol* **36**, 621-642, doi:10.1016/j.biocel.2003.11.002 (2004).
30   Chan, Sunny S.-K. *et al.* Mesp1 Patterns Mesoderm into Cardiac, Hematopoietic, or Skeletal
Myogenic Progenitors in a Context-Dependent Manner. *Cell Stem Cell* **12**, 587-601,
doi:10.1016/j.stem.2013.03.004 (2013).
31   Devine, W. P., Wythe, J. D., George, M., Koshiba-Takeuchi, K. & Bruneau, B. G. Early patterning
and specification of cardiac progenitors in gastrulating mesoderm. *Elife* **3**,
doi:10.7554/eLife.03848 (2014).
32   Lescroart, F. *et al.* Early lineage restriction in temporally distinct populations of Mesp1 progenitors
during mammalian heart development. *Nat Cell Biol* **16**, 829-840, doi:10.1038/ncb3024 (2014).
33   Klotz, L. *et al.* Cardiac lymphatics are heterogeneous in origin and respond to injury. *Nature* **522**,
62-67, doi:10.1038/nature14483 (2015).
34   Chan, S. S. *et al.* Mesp1 patterns mesoderm into cardiac, hematopoietic, or skeletal myogenic
progenitors in a context-dependent manner. *Cell Stem Cell* **12**, 587-601,
doi:10.1016/j.stem.2013.03.004 (2013).
35   Baup, D., Moser, M., Schurmans, S. & Leo, O. Developmental regulation of the composite CAG
promoter activity in the murine T lymphocyte cell lineage. *Genesis* **47**, 799-804,
doi:10.1002/dvg.20569 (2009).
36   Alexopoulou, A. N., Couchman, J. R. & Whiteford, J. R. The CMV early enhancer/chicken beta actin
(CAG) promoter can be used to drive transgene expression during the differentiation of murine
embryonic stem cells into vascular progenitors. *BMC Cell Biol* **9**, 2, doi:10.1186/1471-2121-9-2
(2008).
37   Hu, X. *et al.* Reprogramming progressive cells display low CAG promoter activity. *Stem Cells* **39**,
43-54, doi:10.1002/stem.3295 (2021).

38   Akagi, Y. *et al.* Transcriptional activation of a hybrid promoter composed of cytomegalovirus
enhancer and beta-actin/beta-globin gene in glomerular epithelial cells in vivo. *Kidney Int* **51**,
1265-1269, doi:10.1038/ki.1997.172 (1997).

39   Agah, R. *et al.* Gene recombination in postmitotic cells. Targeted expression of Cre recombinase
provokes cardiac-restricted, site-specific rearrangement in adult ventricular muscle in vivo. *J Clin*
Invest **100**, 169-179, doi:10.1172/JCI119509 (1997).

40   Sokoloff, A. J., Li, H. & Burkholder, T. J. Limited expression of slow tonic myosin heavy chain in
human cranial muscles. *Muscle Nerve* **36**, 183-189, doi:10.1002/mus.20797 (2007).

41   Stockdale, F. E. & Miller, J. B. The cellular basis of myosin heavy chain isoform expression during
development of avian skeletal muscles. *Dev Biol* **123**, 1-9, doi:10.1016/0012-1606(87)90420-9
(1987).

42   Miller, J. B., Crow, M. T. & Stockdale, F. E. Slow and fast myosin heavy chain content defines three
types of myotubes in early muscle cell cultures. *J Cell Biol* **101**, 1643-1650,
doi:10.1083/jcb.101.5.1643 (1985).

43   Vaughan, S. K., Stanley, O. L. & Valdez, G. Impact of Aging on Proprioceptive Sensory Neurons and
Intrafusal Muscle Fibers in Mice. *J Gerontol A Biol Sci Med Sci* **72**, 771-779,
doi:10.1093/gerona/glw175 (2017).

44   Avetisyan, M. *et al.* Hepatocyte Growth Factor and MET Support Mouse Enteric Nervous System
Development, the Peristaltic Response, and Intestinal Epithelial Proliferation in Response to
Injury. *J Neurosci* **35**, 11543-11558, doi:10.1523/JNEUROSCI.5267-14.2015 (2015).

45   Rappolee, D. A., Iyer, A. & Patel, Y. Hepatocyte growth factor and its receptor are expressed in
cardiac myocytes during early cardiogenesis. *Circ Res* **78**, 1028-1036,
doi:10.1161/01.res.78.6.1028 (1996).

46   Hao, M. M., Bornstein, J. C. & Young, H. M. Development of myenteric cholinergic neurons in
ChAT-Cre;R26R-YFP mice. *J Comp Neurol* **521**, 3358-3370, doi:10.1002/cne.23354 (2013).

47   Abraira, V. E. *et al.* The Cellular and Synaptic Architecture of the Mechanosensory Dorsal Horn.
Cell **168**, 295-310 e219, doi:10.1016/j.cell.2016.12.010 (2017).

48   Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-1014 e1022,
doi:10.1016/j.cell.2018.06.021 (2018).

49   Drokhlyansky, E. *et al.* The Human and Mouse Enteric Nervous System at Single-Cell Resolution.
Cell **182**, 1606-1622.e1623, doi:https://doi.org/10.1016/j.cell.2020.08.003 (2020).

50   May-Zhang, A. A. *et al.* Combinatorial Transcriptional Profiling of Mouse and Human Enteric
Neurons Identifies Shared and Disparate Subtypes In Situ. *Gastroenterology* **160**, 755-770 e726,
doi:10.1053/j.gastro.2020.09.032 (2021).

51   Morarach, K. *et al.* Diversification of molecularly defined myenteric neuron classes revealed by
single-cell RNA sequencing. *Nat Neurosci* **24**, 34-46, doi:10.1038/s41593-020-00736-x (2021).

52   Pollen, A. A. *et al.* Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity and
activated signaling pathways in developing cerebral cortex. *Nat Biotechnol* **32**, 1053-1058,
doi:10.1038/nbt.2967 (2014).

53   Heanue, T. A. & Pachnis, V. Expression profiling the developing mammalian enteric nervous
system identifies marker and candidate Hirschsprung disease genes. *Proceedings of the National*
Academy of Sciences **103**, 6919, doi:10.1073/pnas.0602152103 (2006).

54   D'Autreaux, F., Morikawa, Y., Cserjesi, P. & Gershon, M. D. Hand2 is necessary for terminal
differentiation of enteric neurons from crest-derived precursors but not for their migration into
the gut or for formation of glia. *Development* **134**, 2237-2249, doi:10.1242/dev.003814 (2007).

55   Strick, C. A. *et al.* Alterations in gene regulation following inhibition of the striatum-enriched
phosphodiesterase, PDE10A. *Neuropharmacology* **58**, 444-451,
doi:10.1016/j.neuropharm.2009.09.008 (2010).

56 Bernstein, H. G. *et al.* Hippocampal expression of the calcium sensor protein visinin-like protein-
1 in schizophrenia. *Neuroreport* **13**, 393-396, doi:10.1097/00001756-200203250-00006 (2002).

57 Massart, R., Guilloux, J. P., Mignon, V., Sokoloff, P. & Diaz, J. Striatal GPR88 expression is confined
to the whole projection neuron population and is regulated by dopaminergic and glutamatergic
afferents. *Eur J Neurosci* **30**, 397-414, doi:10.1111/j.1460-9568.2009.06842.x (2009).

58 Soo Hoo, L. *et al.* The SNARE Protein Syntaxin 3 Confers Specificity for Polarized Axonal Trafficking
in Neurons. *PLoS One* **11**, e0163671, doi:10.1371/journal.pone.0163671 (2016).

59 Breuss, M. *et al.* The Expression of Tubb2b Undergoes a Developmental Transition in Murine
Cortical Neurons. *J Comp Neurol* **523**, 2161-2186, doi:10.1002/cne.23836 (2015).

60 De Giorgio, R., Arakawa, J., Wetmore, C. J. & Sternini, C. Neurotrophin-3 and neurotrophin
receptor immunoreactivity in peptidergic enteric neurons. *Peptides* **21**, 1421-1426,
doi:10.1016/s0196-9781(00)00286-2 (2000).

61 Mitsui, S., Yamaguchi, N., Osako, Y. & Yuri, K. Enzymatic properties and localization of motopsin
(PRSS12), a protease whose absence causes mental retardation. *Brain Res* **1136**, 1-12,
doi:10.1016/j.brainres.2006.11.094 (2007).

62 Krammer, H. J., Karahan, S. T., Rumpel, E., Klinger, M. & Kuhnel, W. Immunohistochemical
visualization of the enteric nervous system using antibodies against protein gene product (PGP)
9.5. *Ann Anat* **175**, 321-325, doi:10.1016/s0940-9602(11)80029-4 (1993).

63 Ono, S. *et al.* Regulatory roles of complexins in neurotransmitter release from mature presynaptic
nerve terminals. *Eur J Neurosci* **10**, 2143-2152, doi:10.1046/j.1460-9568.1998.00225.x (1998).

64 Leon, A., Aparicio, G. I. & Scorticati, C. Neuronal Glycoprotein M6a: An Emerging Molecule in
Chemical Synapse Formation and Dysfunction. *Front Synaptic Neurosci* **13**, 661681,
doi:10.3389/fnsyn.2021.661681 (2021).

65 Hoogstraaten, R. I., van Keimpema, L., Toonen, R. F. & Verhage, M. Tetanus insensitive VAMP2
differentially restores synaptic and dense core vesicle fusion in tetanus neurotoxin treated
neurons. *Scientific Reports* **10**, 10913, doi:10.1038/s41598-020-67988-2 (2020).

66 Neill, T., Schaefer, L. & Iozzo, R. V. Decorin: A Guardian from the Matrix. *The American Journal of*
Pathology **181**, 380-387, doi:https://doi.org/10.1016/j.ajpath.2012.04.029 (2012).

67 Scott, P. G., Dodd, C. M. & Pringle, G. A. Mapping the locations of the epitopes of five monoclonal
antibodies to the core protein of dermatan sulfate proteoglycan II (decorin). *Journal of Biological*
Chemistry **268**, 11558-11564, doi:https://doi.org/10.1016/S0021-9258(19)50237-X (1993).

68 Corpening, J. C., Cantrell, V. A., Deal, K. K. & Southard-Smith, E. M. A Histone2BCerulean BAC
transgene identifies differential expression of Phox2b in migrating enteric neural crest derivatives
and enteric glia. *Dev Dyn* **237**, 1119-1132, doi:10.1002/dvdy.21498 (2008).

69 Stein-O'Brien, G. L. *et al.* Decomposing Cell Identity for Transfer Learning across Cellular
Measurements, Platforms, Tissues, and Species. *Cell Syst* **8**, 395-411 e398,
doi:10.1016/j.cels.2019.04.004 (2019).

70 Stein-O'Brien, G. L. *et al.* Enter the Matrix: Factorization Uncovers Knowledge from Omics. *Trends*
Genet **34**, 790-805, doi:10.1016/j.tig.2018.07.003 (2018).

71 Lee, M. Y. *et al.* Smooth Muscle Cell Genome Browser: Enabling the Identification of Novel Serum
Response Factor Target Genes. *PLOS ONE* **10**, e0133751, doi:10.1371/journal.pone.0133751
(2015).

72 Purves-Tyson, T. D., Brown, A. M., Weissleder, C., Rothmond, D. A. & Shannon Weickert, C.
Reductions in midbrain GABAergic and dopamine neuron markers are linked in schizophrenia. *Mol*
Brain **14**, 96, doi:10.1186/s13041-021-00805-7 (2021).

73 Mitsui, S. *et al.* A mental retardation gene, motopsin/neurotrypsin/prss12, modulates
hippocampal function and social interaction. *Eur J Neurosci* **30**, 2368-2378, doi:10.1111/j.1460-
9568.2009.07029.x (2009).

Singec, I. *et al.* Synaptic vesicle protein synaptoporin is differently expressed by subpopulations of mouse hippocampal neurons. *J Comp Neurol* **452**, 139-153, doi:10.1002/cne.10371 (2002).

Fichna, J. *et al.* Transient receptor potential vanilloid 4 inhibits mouse colonic motility by activating NO-dependent enteric neurotransmission. *J Mol Med (Berl)* **93**, 1297-1309, doi:10.1007/s00109-015-1336-5 (2015).

Born, G. *et al.* Modulation of synaptic function through the alpha-neurexin-specific ligand neurexophilin-1. *Proc Natl Acad Sci U S A* **111**, E1274-1283, doi:10.1073/pnas.1312112111 (2014).

Zhan, X. *et al.* Generation of BAF53b-Cre transgenic mice with pan-neuronal Cre activities. *genesis* **53**, 440-448, doi:https://doi.org/10.1002/dvg.22866 (2015).

Elmentaite, R. *et al.* Cells of the human intestinal tract mapped across space and time. *Nature* **597**, 250-255, doi:10.1038/s41586-021-03852-1 (2021).

Matheis, F. *et al.* Adrenergic Signaling in Muscularis Macrophages Limits Infection-Induced Neuronal Loss. *Cell* **180**, 64-78 e16, doi:10.1016/j.cell.2019.12.002 (2020).

Barrenschee, M. *et al.* SNAP-25 is abundantly expressed in enteric neuronal networks and upregulated by the neurotrophic factor GDNF. *Histochem Cell Biol* **143**, 611-623, doi:10.1007/s00418-015-1310-x (2015).

Shore, A. N. *et al.* Reduced GABAergic Neuron Excitability, Altered Synaptic Connectivity, and Seizures in a KCNT1 Gain-of-Function Mouse Model of Childhood Epilepsy. *Cell Reports* **33**, 108303, doi:https://doi.org/10.1016/j.celrep.2020.108303 (2020).

Steinmetz, N. A. *et al.* Aberrant Cortical Activity in Multiple GCaMP6-Expressing Transgenic Mouse Lines. *eNeuro* **4**, doi:10.1523/ENEURO.0207-17.2017 (2017).

Zatka-Haas, P., Steinmetz, N. A., Carandini, M. & Harris, K. D. Sensory coding and the causal impact of mouse cortex in a visual decision. *eLife* **10**, e63163, doi:10.7554/eLife.63163 (2021).

Gianino, S., Grider, J. R., Cresswell, J., Enomoto, H. & Heuckeroth, R. O. GDNF availability determines enteric neuron number by controlling precursor proliferation. *Development* **130**, 2187-2198, doi:10.1242/dev.00433 (2003).

Natarajan, D., Marcos-Gutierrez, C., Pachnis, V. & de Graaff, E. Requirement of signalling by receptor tyrosine kinase RET for the directed migration of enteric nervous system progenitor cells during mammalian embryogenesis. *Development* **129**, 5151-5160 (2002).

Taraviras, S. *et al.* Signalling by the RET receptor tyrosine kinase and its role in the development of the mammalian enteric nervous system. *Development* **126**, 2785-2797 (1999).

Rodrigues, D. M., Li, A. Y., Nair, D. G. & Blennerhassett, M. G. Glial cell line-derived neurotrophic factor is a key neurotrophin in the postnatal enteric nervous system. *Neurogastroenterol Motil* **23**, e44-56, doi:10.1111/j.1365-2982.2010.01626.x (2011).

Amano, O. *et al.* Hepatocyte growth factor is essential for migration of myogenic cells and promotes their proliferation during the early periods of tongue morphogenesis in mouse embryos. *Dev Dyn* **223**, 169-179, doi:10.1002/dvdy.1228 (2002).

Brun, P. *et al.* Toll-like receptor 2 regulates intestinal inflammation by controlling integrity of the enteric nervous system. *Gastroenterology* **145**, 1323-1333, doi:10.1053/j.gastro.2013.08.047 (2013).

Jangphattananont, N. *et al.* Distinct Localization of Mature HGF from its Precursor Form in Developing and Repairing the Stomach. *Int J Mol Sci* **20**, doi:10.3390/ijms20122955 (2019).

Almanzar, N. *et al.* A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. *Nature* **583**, 590-595, doi:10.1038/s41586-020-2496-1 (2020).

Sathyan, S. *et al.* Plasma proteomic profile of age, health span, and all-cause mortality in older adults. *Aging Cell* **n/a**, e13250, doi:10.1111/acel.13250.

Wang, H. *et al.* The timing and location of glial cell line-derived neurotrophic factor expression determine enteric nervous system structure and function. *J Neurosci* **30**, 1523-1538, doi:10.1523/JNEUROSCI.3861-09.2010 (2010).

Nadal-Ribelles, M. *et al.* Sensitive high-throughput single-cell RNA-seq reveals within-clonal transcript correlations in yeast populations. *Nat Microbiol* **4**, 683-692, doi:10.1038/s41564-018-0346-9 (2019).

Zheng, S. C. *et al.* Universal prediction of cell-cycle position using transfer learning. *Genome Biol* **23**, 41, doi:10.1186/s13059-021-02581-y (2022).

Schneid, S. *et al.* The BRCT domains of ECT2 have distinct functions during cytokinesis. *Cell Rep* **34**, 108805, doi:10.1016/j.celrep.2021.108805 (2021).

Barrenschee, M. *et al.* No neuronal loss, but alterations of the GDNF system in asymptomatic diverticulosis. *PLoS One* **12**, e0171416, doi:10.1371/journal.pone.0171416 (2017).

Rossi, V. *et al.* Chronic intestinal pseudo-obstruction in a child harboring a founder Hirschsprung RET mutation. *Am J Med Genet A* **170**, 2400-2403, doi:10.1002/ajmg.a.37787 (2016).

Uesaka, T., Nagashimada, M., Yonemura, S. & Enomoto, H. Diminished Ret expression compromises neuronal survival in the colon and causes intestinal aganglionosis in mice. *J Clin Invest* **118**, 1890-1898, doi:10.1172/JCI34425 (2008).

Kim, M. *et al.* Obstructed defecation-an enteric neuropathy? An exploratory study of patient samples. *Int J Colorectal Dis* **34**, 193-196, doi:10.1007/s00384-018-3160-1 (2019).

Sun, T. *et al.* Aging-dependent decrease in the numbers of enteric neurons, interstitial cells of Cajal and expression of connexin43 in various regions of gastrointestinal tract. *Aging (Albany NY)* **10**, 3851-3865, doi:10.18632/aging.101677 (2018).

Becker, L. *et al.* Age-dependent shift in macrophage polarisation causes inflammation-mediated degeneration of enteric nervous system. *Gut* **67**, 827-836, doi:10.1136/gutjnl-2016-312940 (2018).

Sharma, G., Colantuoni, C., Goff, L. A., Fertig, E. J. & Stein-O'Brien, G. projectR: an R/Bioconductor package for transfer learning via PCA, NMF, correlation and clustering. *Bioinformatics* **36**, 3592-3593, doi:10.1093/bioinformatics/btaa183 (2020).

Young, H. M. & Newgreen, D. Enteric neural crest-derived cells: Origin, identification, migration, and differentiation. *The Anatomical Record* **262**, 1-15, doi:10.1002/1097-0185(20010101)262:1<1::AID-AR1006>3.0.CO;2-2 (2001).

Prummel, K. D., Nieuwenhuize, S. & Mosimann, C. The lateral plate mesoderm. *Development* **147**, dev175059, doi:10.1242/dev.175059 (2020).

Carlson, B. M. in *Human Embryology and Developmental Biology (Fifth Edition)* (ed Bruce M. Carlson) 269-293 (W.B. Saunders, 2014).

Verderio, C. *et al.* SNAP-25 Modulation of Calcium Dynamics Underlies Differences in GABAergic and Glutamatergic Responsiveness to Depolarization. *Neuron* **41**, 599-610, doi:https://doi.org/10.1016/S0896-6273(04)00077-7 (2004).

Nouvian, R. *et al.* Exocytosis at the hair cell ribbon synapse apparently operates without neuronal SNARE proteins. *Nature Neuroscience* **14**, 411-413, doi:10.1038/nn.2774 (2011).

Grabs, D., Bergmann, M., Urban, M., Post, A. & Gratzl, M. Rab3 proteins and SNAP-25, essential components of the exocytosis machinery in conventional synapses, are absent from ribbon synapses of the mouse retina. *Eur J Neurosci* **8**, 162-168, doi:10.1111/j.1460-9568.1996.tb01177.x (1996).

Hellström, J., Arvidsson, U., Elde, R., Cullheim, S. & Meister, B. Differential expression of nerve terminal protein isoforms in VAChT-containing varicosities of the spinal cord ventral horn. *Journal of Comparative Neurology* **411**, 578-590, doi:https://doi.org/10.1002/(SICI)1096-9861(19990906)411:4<578::AID-CNE4>3.0.CO;2-L (1999).

111 Morris, J. L., Lindberg, C. E. Y. & Gibbins, I. L. Different levels of immunoreactivity for synaptosomal-associated protein of 25 kDa in vasoconstrictor and vasodilator axons of guinea-pigs. *Neuroscience Letters* **294**, 167-170, doi:https://doi.org/10.1016/S0304-3940(00)01568-8 (2000).

112 Greenlee, M. H., Roosevelt, C. B. & Sakaguchi, D. S. Differential localization of SNARE complex proteins SNAP-25, syntaxin, and VAMP during development of the mammalian retina. *J Comp Neurol* **430**, 306-320, doi:10.1002/1096-9861(20010212)430:3<306::aid-cne1032>3.0.co;2-b (2001).

113 Gibbins, I. L., Jobling, P., Teo, E. H., Matthew, S. E. & Morris, J. L. Heterogeneous expression of SNAP-25 and synaptic vesicle proteins by central and peripheral inputs to sympathetic neurons. *J Comp Neurol* **459**, 25-43, doi:10.1002/cne.10527 (2003).

114 Xue, R. *et al.* Expression of Cystic Fibrosis Transmembrane Conductance Regulator in Ganglia of Human Gastrointestinal Tract. *Sci Rep* **6**, 30926, doi:10.1038/srep30926 (2016).

115 Wang, W. *et al.* Selective Expression of a SNARE-Cleaving Protease in Peripheral Sensory Neurons Attenuates Pain-Related Gene Transcription and Neuropeptide Release. *Int J Mol Sci* **22**, doi:10.3390/ijms22168826 (2021).

116 Tender, G. C., Kaye, A. D., Li, Y. Y. & Cui, J. G. Neurotrophin-3 and tyrosine kinase C have modulatory effects on neuropathic pain in the rat dorsal root ganglia. *Neurosurgery* **68**, 1048-1055; discussion 1055, doi:10.1227/NEU.0b013e318208f9c4 (2011).

117 de Jonge, H. R., Ardelean, M. C., Bijvelds, M. J. C. & Vergani, P. Strategies for cystic fibrosis transmembrane conductance regulator inhibition: from molecular mechanisms to treatment for secretory diarrhoeas. *FEBS Lett* **594**, 4085-4108, doi:10.1002/1873-3468.13971 (2020).

118 de Lima, J. B. M. *et al.* ARC(GHR) Neurons Regulate Muscle Glucose Uptake. *Cells* **10**, doi:10.3390/cells10051093 (2021).

119 Chaudhury, A., Dendi, V. S. & Mirza, W. Colligative Property of ATP: Implications for Enteric Purinergic Neuromuscular Neurotransmission. *Front Physiol* **7**, 500, doi:10.3389/fphys.2016.00500 (2016).

120 Daugherty, M., Luo, Q. & Sokoloff, A. J. Myosin heavy chain composition of the human genioglossus muscle. *J Speech Lang Hear Res* **55**, 609-625, doi:10.1044/1092-4388(2011/10-0287) (2012).

121 Parathan, P., Wang, Y., Leembruggen, A. J., Bornstein, J. C. & Foong, J. P. The enteric nervous system undergoes significant chemical and synaptic maturation during adolescence in mice. *Dev Biol* **458**, 75-87, doi:10.1016/j.ydbio.2019.10.011 (2020).

122 Zhu, H., Hasman, R. A., Barron, V. A., Luo, G. & Lou, H. A nuclear function of Hu proteins as neuron-specific alternative RNA processing regulators. *Mol Biol Cell* **17**, 5105-5114, doi:10.1091/mbc.e06-02-0099 (2006).

123 Epstein, M. L., Saffrey, M. J. & Poulsen, K. T. Development and birthdates of vasoactive intestinal peptide immunoreactive neurons in the chick proventriculus. *Journal of Comparative Neurology* **321**, 83-92, doi:https://doi.org/10.1002/cne.903210108 (1992).

124 Zanini, F. *et al.* Northstar enables automatic classification of known and novel cell types from tumor samples. *Scientific Reports* **10**, 15251, doi:10.1038/s41598-020-71805-1 (2020).

125 Wang, Z. *et al.* Single-cell RNA sequencing reveals a novel cell type and immunotherapeutic targets in papillary thyroid cancer. *medRxiv*, 2021.2002.2024.21251881, doi:10.1101/2021.02.24.21251881 (2021).

126 Zhang, Y., Aevermann, B., Gala, R. & Scheuermann, R. H. Cell type matching in single-cell RNA-sequencing data using FR-Match. *Scientific Reports* **12**, 9996, doi:10.1038/s41598-022-14192-z (2022).

127   Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nature*
2104   *Communications* **8**, 14049, doi:10.1038/ncomms14049 (2017).
2105 128   Genomics, X. *What fraction of mRNA transcripts are captured per cell?*,
2106   <https://kb.10xgenomics.com/hc/en-us/articles/360001539051-What-fraction-of-mRNA-
2107   transcripts-are-captured-per-cell-> (
2108 129   Song, Y., Su, M., Panchatsharam, P., Rood, D. & Lai, L. c-Met signalling is required for efficient
2109   postnatal thymic regeneration and repair. *Immunology* **144**, 245-253,
2110   doi:https://doi.org/10.1111/imm.12365 (2015).
2111 130   Dai, C., Huh, C. G., Thorgeirsson, S. S. & Liu, Y. Beta-cell-specific ablation of the hepatocyte growth
2112   factor receptor results in reduced islet size, impaired insulin secretion, and glucose intolerance.
2113   *Am J Pathol* **167**, 429-436, doi:10.1016/s0002-9440(10)62987-2 (2005).
2114 131   Ishibe, S. *et al.* Met and the epidermal growth factor receptor act cooperatively to regulate final
2115   nephron number and maintain collecting duct morphology. *Development* **136**, 337-345,
2116   doi:10.1242/dev.024463 (2009).
2117 132   Arechederra, M. *et al.* Met signaling in cardiomyocytes is required for normal cardiac function in
2118   adult mice. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease* **1832**, 2204-2215,
2119   doi:https://doi.org/10.1016/j.bbadis.2013.08.008 (2013).
2120 133   Golden, J. P., DeMaro, J. A., Osborne, P. A., Milbrandt, J. & Johnson, E. M., Jr. Expression of
2121   neurturin, GDNF, and GDNF family-receptor mRNA in the developing and mature mouse. *Exp
2122   Neurol* **158**, 504-528, doi:10.1006/exnr.1999.7127 (1999).
2123 134   Papaconstantinou, J. *et al.* Attenuation of p38alpha MAPK stress response signaling delays the in
2124   vivo aging of skeletal muscle myofibers and progenitor cells. *Aging (Albany NY)* **7**, 718-733,
2125   doi:10.18632/aging.100802 (2015).
2126 135   Soret, R. *et al.* Glial Cell-Derived Neurotrophic Factor Induces Enteric Neurogenesis and Improves
2127   Colon Structure and Function in Mouse Models of Hirschsprung Disease. *Gastroenterology* **159**,
2128   1824-1838 e1817, doi:10.1053/j.gastro.2020.07.018 (2020).
2129 136   Grider, J. R., Heuckeroth, R. O., Kuemmerle, J. F. & Murthy, K. S. Augmentation of the ascending
2130   component of the peristaltic reflex and substance P release by glial cell line-derived neurotrophic
2131   factor. *Neurogastroenterol Motil* **22**, 779-786, doi:10.1111/j.1365-2982.2010.01489.x (2010).
2132 137   Melo, C. G. S. *et al.* Identification of intrinsic primary afferent neurons in mouse jejunum.
2133   *Neurogastroenterol Motil* **32**, e13989, doi:10.1111/nmo.13989 (2020).
2134 138   Saffrey, M. J. Aging of the mammalian gastrointestinal tract: a complex organ system. *Age (Dordr)*
2135   **36**, 9603, doi:10.1007/s11357-013-9603-2 (2014).
2136 139   Cossais, F. *et al.* Altered enteric expression of the homeobox transcription factor Phox2b in
2137   patients with diverticular disease. *United European Gastroenterology Journal* **7**, 349-357,
2138   doi:10.1177/2050640618824913 (2019).
2139 140   Schafmayer, C. *et al.* Genome-wide association analysis of diverticular disease points towards
2140   neuromuscular, connective tissue and epithelial pathomechanisms. *Gut* **68**, 854,
2141   doi:10.1136/gutjnl-2018-317619 (2019).
2142 141   Muller, P. A. *et al.* Microbiota-modulated CART(+) enteric neurons autonomously regulate blood
2143   glucose. *Science* **370**, 314-321, doi:10.1126/science.abd6176 (2020).
2144 142   Becker, L., Kulkarni, S., Tiwari, G., Micci, M.-A. & Pasricha, P. J. Divergent fate and origin of
2145   neurosphere-like bodies from different layers of the gut. *American Journal of Physiology-
2146   Gastrointestinal and Liver Physiology* **302**, G958-G965, doi:10.1152/ajpgi.00511.2011 (2012).
2147 143   Becker, L., Peterson, J., Kulkarni, S. & Pasricha, P. J. Ex vivo neurogenesis within enteric ganglia
2148   occurs in a PTEN dependent manner. *PLoS One* **8**, e59452 (2013).

144   Becker, L., Peterson, J., Kulkarni, S. & Pasricha, P. J. Ex vivo neurogenesis within enteric ganglia occurs in a PTEN dependent manner. *PLoS One* **8**, e59452, doi:10.1371/journal.pone.0059452 (2013).

145   Shenje, L. T. *et al.* Precardiac deletion of Numb and Numblike reveals renewal of cardiac progenitors. *Elife* **3**, e02164, doi:10.7554/eLife.02164 (2014).

146   Susaki, E. A. *et al.* Whole-brain imaging with single-cell resolution using chemical cocktails and computational analysis. *Cell* **157**, 726-739, doi:10.1016/j.cell.2014.03.042 (2014).

147   Young, H. M., Furness, J. B., Sewell, P., Burcher, E. F. & Kandiah, C. J. Total numbers of neurons in myenteric ganglia of the guinea-pig small intestine. *Cell Tissue Res* **272**, 197-200, doi:10.1007/BF00323587 (1993).

148   Zhou, Y. *et al.* Enteric nervous system abnormalities are present in human necrotizing enterocolitis: potential neurotransplantation therapy. *Stem Cell Res Ther* **4**, 157, doi:10.1186/scrt387 (2013).

149   Hosie, S. *et al.* Gastrointestinal dysfunction in patients and mice expressing the autism-associated R451C mutation in neuroligin-3. *Autism Res* **12**, 1043-1056, doi:10.1002/aur.2127 (2019).

150   Lefevre, C. *et al.* Enteric Nervous System Remodeling in a Rat Model of Spinal Cord Injury: A Pilot Study. *Neurotrauma Rep* **1**, 125-136, doi:10.1089/neur.2020.0041 (2020).

151   Qiu, X. *et al.* NEDL2 regulates enteric nervous system and kidney development in its Nedd8 ligase activity-dependent manner. *Oncotarget* **7**, 31440-31453, doi:10.18632/oncotarget.8951 (2016).

152   Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nat Biotechnol* **34**, 525-527, doi:10.1038/nbt.3519 (2016).

153   Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in single-cell RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nat Biotechnol* **36**, 421-427, doi:10.1038/nbt.4091 (2018).

154   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847-2849, doi:10.1093/bioinformatics/btw313 (2016).

155   Lun, A. T. L. *et al.* EmptyDrops: distinguishing cells from empty droplets in droplet-based single-cell RNA sequencing data. *Genome Biol* **20**, 63, doi:10.1186/s13059-019-1662-y (2019).

156   Griffiths, J. A., Richard, A. C., Bach, K., Lun, A. T. L. & Marioni, J. C. Detection and removal of barcode swapping in single-cell RNA-seq data. *Nat Commun* **9**, 2667, doi:10.1038/s41467-018-05083-x (2018).

157   Lun, A. T., McCarthy, D. J. & Marioni, J. C. A step-by-step workflow for low-level analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122, doi:10.12688/f1000research.9501.2 (2016).

158   McCarthy, D. J., Campbell, K. R., Lun, A. T. & Wills, Q. F. Scater: pre-processing, quality control, normalization and visualization of single-cell RNA-seq data in R. *Bioinformatics* **33**, 1179-1186, doi:10.1093/bioinformatics/btw777 (2017).

159   Csárdi, G. & Nepusz, T.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

a

DECORIN

Hu

Merged

b

MHCst

MET

Hu

Merged

c

MHCst

SLC17A9

DAPI

Merged

Figure 9

Figure 10

Figure 11

Figure 12

Figure 13

a

b

HGNC Symbol	Direction	Gene type	Description	Coding	Mean RC	log FC	P-value	Adj. P-value
SNAP25	Reduced in OD	protein_coding	synaptosomal-associated protein, 25kDa [Source:HGNC Symbol;Acc:11132]	yes	419	-3.56	2.68E-15	1.25E-12
KLHL1	Reduced in OD	protein_coding	kelch-like family member 1 [Source:HGNC Symbol;Acc:6352]	yes	5	-2.92	2.43E-06	7.07E-05
NOS1	Reduced in OD	protein_coding	nitric oxide synthase 1 (neuronal) [Source:HGNC Symbol;Acc:7872]	yes	775	-2.91	3.29E-06	9.06E-05
CHAT	Reduced in OD	protein_coding	choline O-acetyltransferase [Source:HGNC Symbol;Acc:1912]	yes	17	-2.78	2.00E-07	9.34E-06
GDNF	Reduced in OD	protein_coding	glial cell derived neurotrophic factor [Source:HGNC Symbol;Acc:4232]	yes	649	-2.75	2.27E-10	2.98E-08
RET	Reduced in OD	protein_coding	ret proto-oncogene [Source:HGNC Symbol;Acc:9967]	yes	474	-2.59	6.38E-18	5.29E-15
CLIC3	Increased in OD	protein_coding	chloride intracellular channel 3 [Source:HGNC Symbol;Acc:2064]	yes	98	2.95	1.74E-12	4.54E-10
UPK3A	Increased in OD	protein_coding	uroplakin 3A [Source:HGNC Symbol;Acc:12580]	yes	49	2.93	1.47E-06	4.68E-05
CDH3	Increased in OD	protein_coding	cadherin 3, type 1, P-cadherin (placental) [Source:HGNC Symbol;Acc:1762]	yes	252	2.66	9.87E-06	2.20E-04
SLPI	Increased in OD	protein_coding	secretory leukocyte peptidase inhibitor [Source:HGNC Symbol;Acc:11092]	yes	1,924	2.56	9.84E-06	2.20E-04
SLC17A9	Increased in OD	protein_coding	solute carrier family 17 (Vesicular nucleotide transporter). [Source: HGNC]	yes	1,997	2.15	9.67E-06	2.17E-07

Figure 14