Transcutaneous vagus nerve stimulation in humans induces pupil dilation and attenuates alpha oscillations

Omer Sharon[1], Firas Fahoum[2,3], Yuval Nir[1,4,5]

1. Department of Physiology and Pharmacology, Sackler Faculty of Medicine, Tel Aviv University, Israel, Tel Aviv 69978, Israel; 2. Sackler Faculty of Medicine, Tel-Aviv University, Tel Aviv 69978, Israel; 3. EEG and Epilepsy Unit, Department of Neurology, Tel Aviv Sourasky Medical Center,Tel, Aviv, Israel; 4. Sagol School of Neuroscience, Sagol School of Neuroscience, Tel Aviv 69978, Israel; 5. Functional Neurophysiology and Sleep Research Lab, Tel-Aviv Sourasky Medical Center, Tel Aviv, Israel

Corresponding author email address: yuvalnir.tau@gmail.com

Conflict of interest: none

Abstract

Vagus nerve stimulation (VNS) is widely used to treat drug-resistant epilepsy and depression. While the precise mechanisms mediating its long-term therapeutic effects are not fully resolved, they likely involve Locus Coeruleus (LC) stimulation via the nucleus of the solitary tract (NTS) that receives afferent vagal inputs. In rats, VNS elevates LC firing and forebrain noradrenaline (NE) levels, whereas LC lesions suppress VNS therapeutic efficacy. Non-invasive transcutaneous VNS (tVNS) employs electrical stimulation targeting the auricular branch of the vagus nerve at the Cymba Conchae of the ear, but it remains unclear to what extent tVNS mimics VNS. Here, we investigated the short-term effects of tVNS in healthy human male volunteers (n=24) using high-density EEG and pupillometry during visual fixation at rest, comparing short (3.4s) trials of tVNS to sham electrical stimulation at the earlobe (far from the vagus nerve branch) to control for somatosensory stimulation. Although tVNS and sham stimulation did not differ in subjective intensity ratings, tVNS led to robust pupil dilation (peaking 4-5s after trial onset) that was significantly higher than following

sham stimulation. We further quantified how tVNS modulates idle occipital alpha (8-

13Hz) activity, identified in each participant using parallel factor analysis. We found

that tVNS attenuates alpha oscillations to a greater extent than does sham stimulation.

Thus, tVNS reliably induces pupillary and EEG markers of arousal beyond the effects

of somatosensory stimulation, supporting the hypothesis that it elevates noradrenaline

and other arousal-promoting neuromodulatory signaling, and mimics invasive VNS.

Significance statement

Current non-invasive brain stimulation techniques are mostly confined to modulating cortical

activity, as is typically the case with transcranial magnetic or transcranial direct/alternating-

current electrical stimulation. tVNS has been proposed to stimulate subcortical arousal-

promoting nuclei, but previous studies yielded inconsistent results. Here we show that short

(3.4s) tVNS pulses in naïve healthy male participants induce transient pupil dilation and

attenuation of occipital alpha oscillations. These markers of brain arousal are in line with the

established effects of invasive VNS on LC-NE signaling, and support the notion that tVNS

mimics VNS. Therefore, tVNS can be used as a tool for studying how endogenous subcortical

neuromodulatory signaling affects human cognition including perception, attention, memory,

and decision-making, as well as for developing novel clinical applications.

1 Introduction

Vagus nerve stimulation (VNS) has been successfully used to reduce epileptic seizures in patients with drug-resistant epilepsy since 1988 (Krahl and Clark, 2012) and is clinically effective for many patients treated with invasive VNS (Boon et al., 2018; Kwon et al., 2018). In addition, it is applied as a treatment for drug resistant major depression (e.g Nemeroff et al., 2006).

VNS modulates vagal afferent inputs to the brainstem Nucleus Tractus Solitaris that then activate the LC-NE system. Indeed, in rats VNS increases LC neuronal discharges (Takigawa and Mogenson, 1977; Groves et al., 2005; Hulsey et al., 2017) and elevates NE levels in hippocampus and cortex (Dorr and Debonnel, 2006; Roosevelt et al., 2006). The effects of VNS on LC-NE are considered key to reducing seizures, since a strong positive correlation is found between the noradrenergic and anticonvulsive effects of VNS (Raedt et al., 2011), and chemical lesions of the LC eliminate these anticonvulsive effects (Krahl et al., 1998). VNS also modulates signaling in other neuromodulatory pathways such as the serotonergic, dopaminergic, and cholinergic systems (Dorr and Debonnel, 2006; Manta et al., 2009; Mridha et al., 2019) . However, it is likely that some of these effects are secondary, i.e. occur later and mediated through the LC-NE system (Dorr and Debonnel, 2006).

In human patients, invasive VNS induces markers of brain arousal that are consistent with LC-NE activity such as pupil dilation (Desbeaumes Jodoin et al., 2015) that is tightly linked with LC-NE activity (Joshi et al., 2016; Reimer et al., 2016; Gelbard-Sagiv et al., 2018; Hayat et al., 2020). VNS may also lead to EEG desynchronization, but effects are subtler than pupil dilation, at least with the clinical parameters that typically employ long (30-60s) stimulation epochs. Accordingly, early studies with <10 patients each, did not find VNS effects on spontaneous intracranial EEG (Hammond et al., 1992) or scalp EEG (Salinsky and Burchiel, 1993), whereas a recent study with 19 participants that separated VNS 'responders' from 'non-responders' observed EEG desynchronization in the alpha and delta bands (Bodin et al., 2015).

Non-invasive transcutaneous vagal nerve stimulation (tVNS) applies electrical current at a high

frequency (typically 25Hz) through the left ear, targeting the auricular branch of the vagus nerve

at the Cymba Conchae (Figure1) (for anatomic evidence see Van Bockstaele et al., 1999;

Bermejo et al., 2017). tVNS has been shown to mimic the anticonvulsive and antidepressant

effects of invasive VNS (Stefan et al., 2012; He et al., 2013; Hein et al., 2013; Bauer et al.,

2016; Rong et al., 2016; Trevizol et al., 2016) and proved to be safe and tolerable (Redgrave et

al., 2018). Beyond its clinical efficacy, there is growing interest in using tVNS in healthy

individuals for basic neuroscience research (Van Leusden et al., 2015), but previous studies are

inconsistent in determining to what extent tVNS mimics invasive VNS with respect to the

effects on EEG or pupil dilation that could suggest LC-NE involvement (Ventura-Bort et al.,

2018; Warren et al., 2018; Keute et al., 2019). We suspected that such inconsistencies stem

from employing long (e.g. 30sec) stimulation epochs as used clinically, and due to the indirect

focus on the P300 component where LC-NE activity is assumed to play a key role.

Here, we set out to test if short-term tVNS induces EEG and pupillary markers of arousal, as is

the case for VNS-induced activation. We used short (3.4s) stimulation pulses during task-free

rest conditions in healthy naïve volunteers (to avoid long-term changes associated with

therapeutic effects (Follesa et al., 2007; Manta et al., 2013)). We hypothesized that if indeed

tVNS increases LC and neuromodulatory activities, it should lead to pupil dilation as observed

across multiple species (Joshi et al., 2016; Reimer et al., 2016; Hayat et al., 2020). In addition,

we hypothesized that tVNS will attenuate alpha oscillations that are anti-correlated with arousal

during rest (Torsvall and Akerstedt, 1987a; Drapeau and Carrier, 2004a; Amzica and Lopes da

Silva, 2017), and are attenuated by invasive VNS (Bodin et al., 2015). In line with these

predictions, we found that tVNS induces pupil dilation and alpha desynchronization above and

beyond the effects of sham (somatosensory) stimulation.

Materials and methods

Participants

High-density (256-channel) EEG and pupillometry recording were performed in 25 healthy young male adults (mean age: 28.08 ±5.84 years, 2 left-handed). Written informed consent was obtained from each participant. The study was approved by the Medical Institutional Review Board (IRB) at the Tel Aviv Sourasky Medical Center. Female subjects of child bearing age were not included per guidelines of the approved IRB. Participants reported to be healthy, without history of neuropsychiatric disorders, and indicated their dominant eye for pupillometry. One participant was excluded due to excessive blinking, after which 24 participants remained (mean age: 28.3 ± 1.2). Data from additional three subjects were excluded from EEG analysis due to lack of alpha activity, after which 21 subjects remained for EEG analysis (mean age 28.01±1.3).

Experimental design

Main experiment. After EEG setup (see below), participants performed a short 'method of limits' procedure to select tVNS/sham stimulation intensities while sitting. This procedure systematically finds the maximal comfortable stimulation levels for each individual, as in (Kraus et al., 2013; Yakunina et al., 2017; Ventura-Bort et al., 2018). We applied 5s-long stimulation trials, starting at 0.1mA, and increasing in each trial by 0.2mA. After each trial, participants rated the subjective intensity on a scale of 0-9 ([0] = no sensation; [3] = light tingling; [6] = strong tingling; [9] = painful) and we kept increasing the current until either reaching a level rated as [9] or a maximal level of 5mA. This procedure was carried out twice for each stimulation location (real tVNS at cymba cochnea vs. sham stimulation at ear lobe), and the selected currents were the average currents corresponding to a subjective rating of [8] (just below painful), for each stimulation location separately. Thus, tVNS intensity was adjusted for each subject and location separately to be above the detection threshold and below the pain threshold, as in clinical settings (Ellrich, 2011). Participants were then instructed to position

their heads in a chin-rest apparatus for adjusting and calibrating the eye-tracker (see below).

Then, participants were instructed to fixate on a white cross on a background of a gray computer

screen (HP model 2311x, positioned 80cm from participants' eyes) throughout experimental

"blocks" lasting 5 minutes, each including 11 trials of 3.4s stimulation epochs (in each trial,

tVNS intensity ramped up gradually up to the level defined above), separated by inter-stimulus-

intervals of 26s (±1s jitter). We performed two blocks of either tVNS or sham, and then

switched to position the stimulating electrode in the alternate location (order counterbalanced),

reaching a total of eight blocks altogether per session (Figure 1C). Before changing the

electrode location, participants answered questions regarding their subjective experience of

stimulation (Table 1). Participants were free to rest between the blocks ad lib. We refer to these

rest periods as "breaks", and data acquired during these times were used to characterize alpha

activity in each individual in a manner unbiased with respect to experimental conditions (see

below).

Pilot experiment. A similar experiment using the same device with a separate group of 29 male

participants (mean age: 26.82 ±1.1 years, 4 left-handed) used the default clinical stimulation

mode (30s on, 30s off) during fixation at rest while recording high-density EEG (n=15) and

pupillometry (n=29).

Transcutaneous Vagus Nerve Stimulation (tVNS)

tVNS was delivered using NEMOS®, (Cerbomed, Germany, now tVNS technologies; Figure

1B). In the tVNS condition, the electrodes were placed at the left Cymba Conchae, which is

heavily innervated by the auricular branch of the vagus nerve (Peuker and Filler, 2002; Safi et

al., 2016; Badran et al., 2018) (Figure 1A). In the sham condition, the electrodes were placed

at the left earlobe (Figure 1A), which is not expected to induce brainstem or cortical activation

(Kraus et al., 2007; Sellaro et al., 2015; Steenbergen et al., 2015). Pulses (200-300µs width)

were delivered at a rate of 25Hz (duty cycle of ~7% ON time) for 3.4s that included a ramping

up of intensity (as set by the device) to a level experienced as just-below painful, adjusted for

each participant and condition separately ('method of limits' procedure above), as often set

clinically in patients (Vonck et al., 2014). To achieve 3.4s stimulation trials, we controlled the

NEMOS stimulation device using linear actuators (Actounix, Canada) that pressed the ON/OFF

button automatically according to programmable times. These actuators were controlled by

Arduino mega (Arduino, Italy) directed by Psychopy python package (Peirce, 2007). Two

additional measures verified good electrode contact throughout, and consistent effectiveness of

stimulation: (i) the NEMOS device stops stimulation automatically whenever good physical

contact with the participant's skin is disrupted, and (ii) the experimenter verified in each

participant the presence of a visually-evident 25Hz stimulation artifact in EEG electrodes close

to the left ear.

Figure 1: Experimental design

Legend: (A) Schematic illustration of the rationale of tVNS (B) Stimulation electrode
placement – (1) location of tVNS effective location on the Cymba conchae of the left ear (2)
location of the sham ineffective location (3) Photo of the commercial stimulation electrode. (C)
Experimental design, each experiment started with a 'method of limits' procedure in order to
adjust stimulation current according to individual subjective pain report (Rating) and increase
in 0,2 mA every step until a current matched to a rating of 8 was selected. Then there 8 blocks
each of 5min including 11 stimulation trials of 3.4s and stimulation interval of 25-27s.

Pupillometry

Data acquisition. Eye movements/gaze and pupil size were recorded monocularly from the

dominant eye using infrared video-oculographic system with a chin-rest (Eyelink 1000 Plus,

SR Research). Gaze and pupil data were sampled at 500Hz, positions were converted into

degrees of visual angle based on a 9-point calibration performed at the beginning of the

experiment (on mid-gray background). The experiment was carried out in a room with constant

ambient light.

Data analysis.

Pupil data were low-pass filtered using a 10Hz 4th-order Butterworth filter with zero phase shift,

and periods of blinks were detected using the manufacturer's standard algorithms with default

settings. The remaining data analyses were performed using custom-made Matlab scripts (The

MathWorks). Blinks were removed by linear interpolation of values measured 100ms before

and after each identified blink (de Gee et al., 2014). Segmentation of peri-trial data was

performed by extracting pupil data [-10s +13.4s] around each stimulation onset. Trials where

interpolated data accounted for > 50% of data points were excluded (van Steenbergen and

Band, 2013). After excluding one participant that had no trials remaining, the process yielded

an average of 42.12±1.79 trials in the tVNS condition and 42.16±1.79 in the sham condition

(out of 45, range 42-45 for both) . To allow averaging across subjects with different pupil sizes

while avoiding arbitrary units, we converted pupil data to 'percent change' values relative to

10s baseline prior to stimulation: [(x-baseline/baseline)*100] (As in Reimer et al., 2016; Liu et

al., 2017).

The resulting pupil time-courses were the mean value across trials for each participant and

condition separately, as depicted in Figure 2A (for visualization only, single subject traces were

band-pass filtered again between 0.01-10Hz in Figure 2C). In order to present the individual

subject data, we reduced the pupil data for each participant and condition to a scalar value

(Figure 2B) by averaging the time-course across trials in the interval between the two points of half maximum (FDHM, 3.2-10.4s) following stimulation onset (see dashed bar in Figure 2A).

Gaze data and blink rate were also inspected and compared between conditions. Gaze was extracted, interpolated and averaged using the same procedure described above for pupil size, and data points marked as blinks were summed across participants to produce a blink rate time-locked to stimulation onset.

EEG

Data acquisition. High-density EEG was recorded continuously using a 256-channel hydrocel geodesic sensor net (Electrical Geodesics, Inc. [EGI], Eugene OR, USA). Each carbon-fiber electrode consists of a silver chloride carbon fiber pellet, a lead wire, a gold-plated pin, and was injected with conductive gel (Electro-Cap International). Signals were referenced to Cz, amplified via an AC-coupled high-input impedance amplifier with antialiasing analog filter (NetAmps 300, EGI) and digitized at 1000 Hz. Electrode impedance in all sensors was verified to be <50 kΩ before starting the recording.

EEG data analysis. EEG preprocessing was performed in MATLAB (The MathWorks) using custom-written code as well as the FieldTrip toolbox (Oostenveld et al., 2011). First, we used a subsample of 192 electrodes placed directly on the skull (avoiding cheek electrodes with higher muscle artifacts). Continuous data from these electrodes were segmented to 33s epochs, [-15s +18s] around each stimulation onset. First, in order to allow effective visual inspection, data epochs were linearly de-trended, notch-filtered (at 50Hz), and high-pass (>0.1Hz) filtered using a 2nd order Butterworth filter to remove DC shifts. We then visually confirmed that all sham and tVNS trials show 25Hz stimulation artifact around the left ear. Trials without the artifact were excluded (on average, 14.75%±3.08 of trials were excluded). To focus on alpha oscillations, data were further band-passed filtered between 5-15Hz using a 3rd-order two-pass Butterworth filter as in previous studies applying parallel factor analysis (Barzegaran et al., 2017) as well as an additional notch filter in 25Hz (stimulation frequency) and its harmonics

up until 475Hz to remove any residual artifact stemming from stimulation that were not

removed by previous filters. Then, we removed the minimal number of channels or trials whose

data crossed an absolute amplitude threshold of 100µV in an automatic iterative process – that

is, each 3s epoch in each channel had a Boolean value [max(abs(x))>100?] – and then in each

iteration, either a channel or a trial was excluded such that a minimal number of channel×trial

3s   data   epochs   were   discarded   (code   is   available   at:

https://github.com/sharomer/eeg_2d_minimal_rejection).   This   process   removed   large

movement artifacts, but not all blink artifacts, which were separated later using the parallel

factor analysis.

This preprocessing resulted in identifying an average of 18.76±2.86% of bad channels per

participant (out of 192, data were interpolated using a linear distance weighted interpolation),

and 22±3.42% bad trials per participant (discarded from subsequent analysis). Only then, trials

were divided to tVNS or sham trials to avoid any bias in preprocessing. The average number of

valid trials in the tVNS condition was 35.61±1.09, and in the sham condition 35.61±1.09 (out

of 44, number of trials was not significantly different between conditions). Next, data of each

trial were transformed to the time-frequency domain using the Fast Fourier Transform (FFT)

after multiplying by a moving hamming window of 3s, yielding a frequency resolution of

0.33Hz and temporal resolution of 0.33s.

Parallel factor (PARAFAC) decomposition. We first extracted data from "break" periods (in

between stimulation blocks) to identify each participant's alpha topography and frequency in

an unbiased manner with respect to study objectives. These data were segmented to 5s epochs

with 1s overlap with its preceding epoch and 1s with the following epoch, band-passed filtered

(as for stimulation data, above) and reduced to 3s trials (discarding the overlap) to avoid

filtering artifacts at edges. Then, "break" data epochs were cleaned as described for stimulation

data, using the same procedure described above (resulting in 117.84±7.15 trials, on average,

per subject, with 175.28±2.71 clean channels on average). These 3s time-frequency epochs

were used to identify each participant's alpha topography and precise frequency range using

the parallel factor analysis as implemented in the N-way toolbox (Andersson and Bro, 2000),

as presented in Figure 3. The type of constraint for each dimension was set to non-negativity.

The proper number of components was determined by using the Core Consistency Diagnostic

(CCD) in which number of components is highest when the minimal value of CCD was 55%

and 90.60±3.18% on average (Bro and Kiers, 2003).

Next, in order to assess the changes in alpha oscillations during stimulation, the individual

weights for alpha component topography and frequency (Figure 4A) were derived from the

break data, and multiplied by the spectrum of all channels such that a single channel

representing the weighted activity was achieved. We then subtracted the mean baseline activity

in [-1:0]s relative to stimulation onset for each subject, for each trial, and calculated the mean

activity across subjects (N=21) to receive the results depicted in Figure 4D.

To assess more carefully the brain activity following stimulation beyond the a-priori electrode-

and frequency-band of interest, (i) we rigidly set the alpha topography (to investigate time

frequency changes in the entire spectrogram). To this end, we used the topography of interest

derived from the PARAFAC decomposition of the break data (Figure 4A, lower panel), ignored

the frequency of interest, plotted the entire spectrogram in 5-15Hz in % change relative to the

same baseline ([-1:0]s prior to stimulation onset, Figure 4G,4H,4E), and used a cluster

permutation test (see below). We also confirmed the difference between tVNS and sham

conditions using post-hoc direct comparison using Wilcoxon sign rank tests (Results).

Alternatively, (ii) we rigidly set the frequency-band of interest (to investigate changes in all

electrodes). To this end, we used the frequency of interest derived from the PARAFAC

decomposition of the break data (Figure 4A upper panel) and ignored the topography of interest,

and plotted the entire topographical changes in voltage around times of stimulation, while

subtracting the activity [-1:0s] prior to stimulation (Figure 4F).

Statistical Analyses

Unless stated otherwise all statistical test were carried using a Wilcoxon sign rank test

(Wilcoxon, 1945), including the significance of the pupil time-course that was corrected to

multiple comparison using FDR correction (Benjamini and Yekutieli, 2011). The significance

for alpha attenuation in the EEG spectrogram was assessed using a cluster permutation test with

the Monte Carlo method and a dependent samples T-statistic with 10,000 permutations as

implemented in the fieldtrip toolbox (Oostenveld et al., 2011). An alpha of 0.05 was considered

significant after FDR correction for clusters (Benjamini and Yekutieli, 2011). In Figure 4, we

plotted alpha attenuation at the individual subject level, so the reader could assess effect size

across subjects; these per-subject values were tested also using Wilcoxon sign rank test (Figure

4C). Data are expressed as mean ± standard error of the mean throughout.

Results

To investigate the short-term effects of tVNS in naïve humans, we compared pupil dynamics

and EEG alpha oscillations in healthy young male volunteers (n=24) induced by multiple trials

of short (3.4s) electrical stimulation at the Cymba Conche (tVNS) or at the earlobe (sham)

(Figure 1). Stimulation was applied at a frequency of 25Hz with intensity ramping up during

the trial up to a maximal value selected per participant and location separately (Methods).

First, we verified that sham and tVNS did not differ in subjective averseness. There was no

significant difference between tVNS and sham in any subjective reports such as pain or

irritation (Table 1, P>0.05 for all comparisons after FDR correction). In terms of objective

current intensity, the average currents applied were 2.20±0.24mA in the tVNS condition and

2.79±0.27mA in the sham condition. The higher current intensity in the sham condition

represented a statistically significant difference (p=0.0125 Wilcoxon signed rank test), likely

due to lower sensitivity at the earlobe, and "works against" our a-priori hypothesis (larger

effects in tVNS *despite* higher current intensity in sham, see Discussion). Therefore, earlobe

stimulation provides a good somatosensory control to distill the changes related specifically to

tVNS.

Table 1: Subjective ratings of tVNS/sham stimulation

	tVNS	sham
I feel headache	1.82±0.20	1.89±0.22
I feel nausea	1.67±0.19	1.63±0.19
I feel dizziness	1.82±0.22	1.97±0.25
I feel neck pain	1.87±0.16	1.97±0.19
I feel muscle contraction in the neck	2.10±0.20	2.00±0.21
I feel stinging under the electrode	2.22±0.22	2.18±0.23
I feel skin irritation in the ear	2.17±0.28	2.15±0.27
I feel fluctuation in concentration	4.02±0.39	3.73±0.36
I feel unpleasant feeling	2.52±0.29	2.71±0.31
I am in good mood	4.82±0.32	5.23±0.27
I am alert	4.07±0.32	4.15±0.31

Transcutaneous vagus nerve stimulation induces pupil dilation

tVNS led to robust pupil dilation that increased gradually (in line with the ramping up of the

stimulation intensity), reaching half maximum at 2.53s after stimulation onset, peaking at 4.25s

after stimulation onset, decreasing back to half maximum at 8.17s and returning to baseline

levels 10s after stimulation. During peak pupil dilation, pupil size (in pixels) was 4.05±0.92%

above baseline (Figure 2A).

By contrast, sham stimulation only led to modest 1.67%+0.63% pupil dilation, peaking around

the same time, and this dilation level was significantly weaker than following tVNS ($p < 0.05$

between 2.88-5.96s via repeated Wilcoxon sign rank test across all time points, and FDR

correction for multiple comparisons, red bar, Figure 2A). These results were largely consistent

across individual subjects (Figure 2B) and evident in most (21 out of 24) participants (for

examples see Figure 2C).

There was no significant difference in baseline pupil values between tVNS and sham conditions (p = 0.5). In both conditions, the smaller was the pupil before a specific trial, the higher was the stimulation-evoked response (Pearson's correlation R=-0.27, $p<10^{-20}$). We found no significant differences between conditions when comparing blink rate or gaze position (p>0.6 for all comparison, using the same statistical procedure). In addition, the higher pupil dilation upon tVNS remained significant and robust across individual subjects when avoiding blink interpolation and simply discarding blink data. In order to verify that the pupil dilation effect was not mediated by the difference in objective currents, we calculated the correlation between differences in pupil dilation (across tVNS and sham), and the difference in current (across tVNS and sham) and did not find a significant correlation (Spearman correlation R=-0.12, p=0.56).

In a pilot experiment employing 30s ON / 30s OFF 'clinical-like' stimulation we only observed a modest trend for greater pupil dilation for tVNS than sham stimulation (p = 0.053, n=23), and there was no significant difference in pupil size between ON and OFF periods. Thus, short tVNS pulses lead to robust pupil dilation that is significantly greater than following sham stimulation, indicating that it promotes arousal above and beyond somatosensory stimulation at the ear.

Figure 2: tVNS leads to greater pupil dilation than sham stimulation

Legend: **(A)** grand average pupil dilation in response to tVNS (red trace) and sham stimulation
(black trace). Shaded areas around the trace indicate SEM. Grey transparent rectangle indicate
active current is on. Upper red line indicates FDR-corrected statistical significance using
Wilcoxon rank sum test. Dashed black bar indicates time interval used to compute individual
subject dilation values in B. **(B)** single subject values in both tVNS and sham condition between
the 2 points of half maximum (FDHM, 3.2-10.4s dashed black bar in A). Solid black lines
denotes tVNS>sham, while dashed grey lines denotes sham>tVNS. I,II refer to the single
subject traces shown in C. **(C)** Representative single-subject pupil time-courses as indicated in
B, with identical graphic representation as in A.

Transcutaneous vagus nerve stimulation attenuates alpha oscillations

Alpha oscillations exhibit considerable inter-individual variability in frequency and scalp

topography (Haegens et al., 2014). To facilitate a sensitive examination of how tVNS or sham

stimulation affects alpha activity, we first identified the frequency and topography of alpha

oscillations in each participant separately using parallel factor analysis (PARAFAC; Harshman,

1970). PARAFAC provides a unique solution to decompose the EEG signal to three factors

(time, frequency, channel; Figure 3A) and may enhance sensitivity. It was previously applied

to electrophysiological recordings (Miwakeichi et al., 2004; Yanagawa et al., 2013; Meij et al.,

2016), and specifically for assessing individual alpha oscillations (Barzegaran et al., 2017;

Knyazeva et al., 2018); for a detailed review of its EEG applications, see Cong et al (2015). We

identified the regions and frequencies of interest for alpha oscillations in each participant

separately using unbiased "break" data between stimulation blocks (Figure 3B). Figure 3C,D

presents the result of this process in representative subjects, and Figure 4A shows the median

region and frequency profile of alpha oscillations across all participants. PARAFAC

successfully identified alpha activity (see examples in Figure 3C,D) capturing the subject's

specific alpha frequency around 7-13Hz with the typical occipital topography.

Figure 3: Parallel Factor Analysis (PARAFAC) to identify individual alpha activity

Legend: Graphic illustration of the Parallel Factor Analysis (PARAFAC) method we used to
decompose the stimulus-free (break) data and create subject-specific topographical and
frequency bands of interest. (A) Illustration of the PARAFAC model with two components,
where f1 and f2 refer to the frequency features, t1 and t2 indicate temporal features, and c1 and
c2 represent the spatial features of the components in the channel space. (B) Spectrogram of 5
single 3s "trials" derived from the break, same subject as top left in C. (C,D) Representative
examples of the decomposition result for 2 different subjects. Each panel includes two
components:1 (pink), and 2 (blue), along with their associated frequency (f) and trial (t) profiles.
The spatial (channel) dimension is presented as scalp topographies on the right side.

After having identified alpha activity in each individual, we quantified the extent to which alpha

activity may be decreased by tVNS or sham stimulation in each subject separately. We found

that tVNS attenuated alpha activity (mean: 94.35±2.2% of baseline) to a greater extent

(p=0.0027, Wilcoxon signed rank test) than sham stimulation, which was not associated with

significant alpha attenuation (103.55 ±2.4% of baseline). Baseline alpha was not significantly

different between conditions (p = 0.3 via sign rank test). Greater alpha attenuation following

tVNS was evident in most (19/21) individual subjects (Figure 4B).

We correlated the differences in alpha attenuation (between tVNS and sham conditions) in each

individual with the differences in applied current (between tVNS and sham conditions) and

found a significant negative correlation (Spearman correlation R=-0.49. p=0.02). Thus, subjects

with stronger sham stimulation current showed less difference in alpha attenuation. Along this

line, repeating the analysis for alpha attenuation while removing one third of participants with

highest difference in current between conditions, revealed that for the remaining n=14 subjects

the difference in alpha attenuation was even more significant despite a smaller number of

subjects (p=0.0001, Wilcoxon signed rank test). This indicates that the alpha attenuation we

observed upon tVNS constitutes a lower bound (if objective current intensity was equal in the

two conditions, a stronger difference in alpha attenuation between tVNS and sham would

emerge, see also Discussion).

Figure 4: tVNS attenuates EEG alpha activity more than sham stimulation

Legend: (A) Median alpha component topography. Pink marks the median weights across
participants. Blue points mark electrodes with highest alpha activity (selected using a threshold
applied to the median weights) to facilitate visualization in subsequent panel E and for the direct
statistical test.(B) Alpha attenuation relative to baseline in individual subject data between
0:4sec, using the weighted topography in A and using the spectral profile in C. Black solid lines
mark subjects showing higher alpha decrease in tVNS condition (19/21), whereas dashed gray
lines mark subjects with higher alpha decrease in sham condition (2/21). (C) Alpha component
spectral profile (median across subjects). (D) Average alpha component time-course (using
spectral profile depicted in C and topographical profile depicted in A). (E) Difference in
induced power between tVNS and sham conditions (shown separately in G and H). White
contours marks statistically significant time-frequency cluster (after correction for multiple
comparisons). Note that tVNS causes alpha attenuation lasting several seconds (F)
Topographical dynamics following stimulation in 1s resolution reveals occipital alpha
attenuation upon tVNS (upper panel) but not in sham (lower panel), yellow points mark
electrodes comprising the statistically significant time-space cluster exhibiting tVNS
attenuation > sham attenuation (after correction for multiple comparisons). Note tha tVNS
cause attenuation in occipital electrodes (G) Average induced spectrogram upon tVNS; white
contour identical to that shown in E. (H) Average induced spectrogram upon sham stimulation;
white contour identical to that shown in E.

tVNS-induced alpha attenuation was not observed in our pilot experiment employing 30s ON /

30s OFF 'clinical-like' stimulation (p > 0.05, n=15). In both experiments, we did not find

significant correlation between alpha attenuation and subjective (or objective) individual scores

of stimulation intensities, or significant correlation between alpha attenuation and pupil dilation at the single subject level (all $p \geq 0.1$).

To complement the PARAFAC-based analysis and better understand the precise time-frequency dynamics and topographical changes of alpha attenuation, we used the weighted alpha topography from the break data as a 'weighted region of interest' to reduce the data to two dimensions (time and frequency). This approach ignored the frequency-of-interest and inspected the induced power changes in the 5-15Hz frequency range for the (weighted) occipital region derived from the PARAFAC decomposition (Figure 4A). In line with the previous results, we found that tVNS significantly attenuated activity in the alpha band (8-12Hz, Figure 4G). Along this line, examining the effects of stimulation on EEG dynamics using cluster based permutation (Maris and Oostenveld, 2007) revealed a significant ($p=0.0063$, white contour in Figure 4G-H) cluster around 8-12Hz in the seconds following stimulation onset where mean alpha power was $90.84 \pm 2.77\%$ in the tVNS condition, significantly lower than the $106.66\% \pm 2.70\%$ observed following sham condition ($p<0.0001$ in a direct comparison). We also compared the two conditions using the classical alpha frequency range (8-12Hz), during stimulation (0-4s) (Means: $94.41 \pm 2.15\%$ in tVNS condition and $105.25 \pm 2.41\%$ in the sham condition, $p=0.0012$ (both in signed rank test)).

Finally, we tested to what extent the observed alpha attenuation was specific for occipital electrodes. We inspected the topographical changes in voltage around stimulation relative to baseline (Figure 4F). This analysis was carried out by focusing on the a-priori frequency-band of interest derived from the PARAFAC decomposition (Figure 4C) while ignoring the topography-of-interest derived from the break. Examining topographical effects of stimulation on EEG dynamics using topographical cluster-based permutation (Maris and Oostenveld, 2007) revealed a significant ($p<0.05$) cluster over occipital electrodes exhibiting tVNS attenuation > sham attenuation (yellow points in Figure 4F), indicating that alpha attenuation is specific to occipital areas. We also compared the two conditions directly using the occipital electrodes marked in Figure 4A during 0:4sec (Mean 3.75µV in tVNS and -1.78µV, $p=0.007$

via sign rank test). Importantly, the regions showing tVNS-induced alpha attenuation overlapped with electrodes showing alpha activity in the independent "break" intervals between stimulation blocks (compare blue dots in Figure 4A with yellow dots in Figure 4F). Altogether, the EEG data establish that short tVNS pulses, but not sham stimulation, attenuate occipital alpha activity.

Discussion

We examined the effects of short tVNS pulses (and sham stimulation at the ear lobe) on pupil dynamics and EEG alpha activity in naïve healthy subjects. While subjective stimulation intensities were not significantly different in the two conditions (Table 1), we found that short tVNS pulses induce pupil dilation (Figure 2) and EEG alpha attenuation (Figure 4) to a greater extent than sham stimulation. These effects support the hypothesis that tVNS activates endogenous arousal-promoting neuromodulatory signaling such as LC-NE activity, as is known for invasive VNS (Hulsey et al., 2017; Mridha et al., 2019), and therefore suggest that tVNS mimics VNS.

Validity and limitations. Our results were obtained during fixation at rest. Although they may generalize well to other conditions, future studies need to determine the effects of short-pulse tVNS during other states such as specific cognitive tasks or during drowsiness and sleep. For example, high arousal at baseline could create a ceiling effect for pupil dilation and alpha attenuation; conversely, during decreased vigilance the EEG effects may attenuate idle activity at different frequency bands (e.g. changing theta/alpha ratio during drowsiness, or suppressing slow wave activity in sleep). Another limitation is that we could only study tVNS in male volunteers. Since there may be sex-specific differences in LC-NE and neuromodulatory activity (Bangasser et al., 2016), future studies with females are warranted. Lastly, our experimental design equated the subjective intensity of tVNS and sham stimulation. Our data revealed that a significantly higher current at the earlobe (sham condition) was necessary to achieve equal

subjective intensity. We could not find a significant correlation between current differences (tVNS vs. sham) and pupil dilation differences (tVNS vs. sham), but current differences were negatively correlated with differences in alpha attenuation (tVNS vs. sham). Accordingly, participants for which current intensity was higher in sham than tVNS showed a smaller difference in alpha attenuation between conditions. Indeed, restricting the analysis to a subset of 14 subjects in a way that eliminated the significant difference in current intensity revealed a stronger effect of alpha attenuation in tVNS vs. sham. Thus, our results represent a conservative lower-bound on the actual difference between alpha attenuation in tVNS and sham, that would be even higher in the event of comparable currents in tVNS and sham. More generally, this touches upon an inherent limitation of using earlobe sham stimulation as a control condition. Despite its extensive use (Yap et al., 2020) and advantages, the lower sensitivity at the earlobe requires higher currents to produce a comparable subjective intensity. Future studies could apply additional control conditions (e.g. stimulating at other frequencies) to go beyond this limitation.

Previous tVNS studies. Our finding that tVNS attenuates alpha oscillations is compatible with some recent studies (Bodin et al., 2015; Lewine et al., 2019) while other studies reported mixed results or could not reveal EEG effects (Hammond et al., 1992; Salinsky and Burchiel, 1993). Our use of short tVNS pulses likely contributed to our ability to reveal alpha attenuation, but it should also be acknowledged that revealing alpha effects required sensitive analysis that used PARAFAC to identify alpha in each individual, and was observed in many, but not all subjects.

Most other previous tVNS studies did not focus on ongoing EEG or pupillometry as done here, but tried to demonstrate the effectiveness of tVNS by focusing on the EEG P300 or on salivary alpha amylase as readouts. The P300 is a positive deflection with maximal amplitude over centro-parietal midline electrodes, 300-500ms after stimulus onset, whose amplitude is modulated by the probability of stimulus appearance regardless of sensory modality (Desmedt et al., 1965; Sutton et al., 1965). It is hypothesized that the P300 is a marker of LC-NE activity (Nieuwenhuis et al., 2005), because LC neurons are likewise activated by infrequent stimuli

independent of sensory modality (Aston-Jones et al., 1991), and deviant stimuli elicit greater

pupil dilation than standard stimuli (Murphy et al., 2011). However, the P300 may not

constitute a straightforward test of tVNS efficacy since the link between P300 and LC-NE

activity is still debated (Nieuwenhuis et al., 2011), and there could be significant contributions

to P300 from dopaminergic (Glover et al., 1988) and glutamatergic (Hall et al., 2015) systems.

Ventura-Bort et al (2018) demonstrated that tVNS amplifies the parietal component of the P300

effect (P3b), selectively for easy targets in their task. However, this effect was modest and could

not be replicated using weaker fixed currents (0.5mA) and simpler classical P300 task (Warren

et al., 2018). Another study by Keute et al (2019) capitulated on the difference in pupil dilation

between deviant and standard stimuli using a classical auditory oddball task. They used a

constant 3mA tVNS in all subjects and found no effect of stimulation on event-related or

baseline pupil size. One possibility is that 30s tVNS modulates tonic NE levels but does not

affect phasic stimulus-evoked NE changes associated with the P300. In agreement with this

possibility, using clonidine (an α2 adrenergic receptor agonist that reduces NE signaling),

provided similar mixed results (Pineda and Swick, 1992; Halliday et al., 1994; Pineda et al.,

1997; Brown et al., 2015). Future studies could use short tVNS pulses as used here to better

assess its effects on the P300.

Both Warren et al (Warren et al., 2018) and Ventura-Bort (2018) showed that tVNS increases

levels of salivary alpha amylase, a peripheral measure of sympathetic activity associated with

LC-NE signaling (Rohleder and Nater, 2009). However, this measure has poor temporal

resolution and can only reveal differences between time intervals before vs. after stimulation

blocks lasting many minutes. This approach does not leverage the superior temporal resolution

of tVNS compared to pharmacological NE manipulations in humans, which are highly effective

in studying the effects of slower NE dynamics (Gelbard-Sagiv et al., 2018). By contrast, the

transient (within seconds) tVNS-mediated effects revealed here offer significant advantages

over slow modulations brought about by NE drugs.

Pupil dilation and alpha attenuation as indices of arousal and LC-NE activity. Pupil size was suggested as a proxy for noradrenergic signaling since Aston Jones and Cohen first provided an example of correlated dynamics in simultaneous pupil and LC single-unit activities in a monkey (Aston-Jones and Cohen, 2005; and see a recent review by Joshi and Gold, 2020) This relation was recently established in monkeys (Varazzani et al., 2015; Joshi et al., 2016), rats (Liu et al., 2017; Hayat et al., 2020), and mice (Reimer et al., 2016; Breton-Provencher and Sur, 2019), as well as in human BOLD fMRI (Murphy et al., 2014). The tVNS-induced pupil dilation time-course we observed (Figure 3) resembles the pupil dynamics in response to LC electrical stimulation in monkeys (Joshi et al., 2016) and optogenetic stimulation in rats (Hayat et al., 2020) supporting the hypothesis that tVNS activates the LC, as is established for invasive VNS.

Alpha oscillations are abundant during detachment from the sensory environment in wakefulness and considered an index of low arousal (Torsvall and Akerstedt, 1987b; Drapeau and Carrier, 2004b). Alpha oscillations are believed to represent an "idling" state of cortical activity (Steriade, 2001; Palva and Palva, 2007) that is expected to be anti-correlated with arousal-promoting activity, such as that of the LC-NE system, and bias sensory perception (Waschke et al., 2019). A recent study that used long 2min tVNS on the neck also found that tVNS attenuates alpha and theta oscillation (Lewine et al., 2019).

LC-NE vs. other neuromodulatory systems. While pupil dilation and EEG alpha attenuation are both *compatible* with noradrenergic signaling, LC-NE involvement is unlikely to be the only modulatory system at play, given the overlap and redundancy among neuromodulatory systems. Other elements such as the cholinergic system also contribute to brain arousal and are associated with both pupil dilation (Reimer et al., 2016) and with EEG activation (Szerb, 1967). However, cholinergic activation alone is unlikely to drive the effects observed here since during Rapid Eye Movement (REM) sleep, when cholinergic activation occurs without LC-NE activity (Nir and Tononi, 2010), the EEG is activated but pupils remain constricted (Siegel, 2005). Moreover, given that VNS robustly activates the LC and no such relation has been reported for

cholinergic nuclei, the most parsimonious interpretation is that the primary neuromodulatory effects of tVNS are noradrenergic, while cholinergic modulation (Mridha et al., 2019) are likely secondary. tVNS may engage additional subcortical neuromodulatory systems such as the dorsal raphe and ventral tegmental area, as observed with tVNS-induced BOLD fMRI (Frangos et al., 2015). Thus, the possible relation of tVNS to other neuromodulatory systems beyond LC-NE is an important topic for further investigation.

tVNS as a novel tool for transient neuromodulation. There is great interest in investigating the contribution of the LC-NE system to human cognition including perception, learning and memory, decision-making, and in aging and neurodegeneration. In this context, tVNS entails important advantages over existing tools. While important contributions have been made by relying on correlation with pupil dynamics (e.g de Gee et al., 2017), correlated LC-NE activity and pupil dynamics could still reflect hidden factors (such as fluctuations in arousal and attention) that constitute the causal root of the observed correlations (Clewett et al., 2018; Dragone et al., 2018) . Previous human studies also employed *causal* perturbations using NE drugs to study effects on perception (Gelbard-Sagiv et al., 2018), memory (see van Stegeren, 2008 for review), and decision making (Warren et al., 2017) . However, systemic delivery of NE drugs is inherently limited to affecting tonic LC-NE activity and has poor temporal resolution, where tVNS has a clear added value.

Using invasive VNS in patients, previous studies reported effects on memory (Clark et al., 1999; Jacobs et al., 2015; for review see Hansen, 2017; Sun et al., 2017). However, such studies are only possible in patients suffering from severe epilepsy or depression. In addition, ongoing daily VNS induces complex long-term plastic changes that make interpretation difficult.

Due to limitations of existing techniques, the potential of tVNS has been recognized and it is being increasingly used to study cognition (Van Leusden et al., 2015). However, to date, the evidence supporting the notion that it is effective in mimicking invasive VNS was mixed. By using short stimulation pulses and maximal possible currents per participant, and by focusing on simple pupillary and ongoing EEG readouts, we show that such stimulation transiently elicits

markers of brain arousal that are compatible with arousal-promoting neuromodulatory signaling such as NE/Ach (supporting the hypothesis that it does mimic invasive VNS) thereby extending the experimental toolkit for non-pharmacological neuromodulation in humans with high temporal resolution. Therefore, it could be used to further investigate how transient neuromodulation contributes to human cognition Future studies could compare several levels of both sham stimulation and tVNS to conduct a parametric investigation. Of note, stronger stimulation may not necessarily produce stronger effects on neuronal activity and behavior since some effects could turn out to elicit "U-shape" profiles (for example see Clark et al., 1999).

Finally, tVNS could be used to further understand the processes mediating the clinical improvements brought about by VNS in epilepsy and depression, for example the role of arousal-promoting neuromodulatory signaling in improving mood in depressed patients (Grimonprez et al., 2015; Liu et al., 2016; Fang et al., 2017; Tu et al., 2018). In particular, tVNS-induced pupillary and EEG effects in individual patients could possibly predict the future clinical efficacy of invasive VNS and help guide triaging patients into either conservative therapy or surgical implantation of VNS stimulation devices.

Software Accessibility code is available upon request.

Acknowledgements: Supported by ISF 51/11 (I-CORE cognitive sciences) and the Adelis Foundation (YN), The Herczeg Institute on Aging, the TAU global research fund, and the Naomi Foundation (OS). The authors thank Nir lab members for discussions, Leon Deouell and Amit Marmelshtein for their comments on earlier drafts of the manuscript.

References

Amzica F, Lopes da Silva FH (2017) Niedermeyer's Electroencephalography (Schomer DL, Lopes da Silva FH, eds). Oxford University Press. Available at: http://www.oxfordmedicine.com/view/10.1093/med/9780190228484.001.0001/med-9780190228484.

Andersson CA, Bro R (2000) The N -way Toolbox for MATLAB. :3–6.

Aston-Jones G, Chiang C, Alexinsky T (1991) Discharge of noradrenergic locus coeruleus neurons in behaving rats and monkeys suggests a role in vigilance. Prog Brain Res 88:501–520 Available at: http://www.ncbi.nlm.nih.gov/pubmed/1813931.

Aston-Jones G, Cohen JD (2005) An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. Annu Rev Neurosci 28:403–450 Available at: http://www.ncbi.nlm.nih.gov/pubmed/16022602.

Badran BW, Brown JC, Dowdle LT, Mithoefer OJ, LaBate NT, Coatsworth J, DeVries WH, Austelle CW, McTeague LM, Yu A, Bikson M, Jenkins DD, George MS (2018) Tragus or cymba conchae? Investigating the anatomical foundation of transcutaneous auricular vagus nerve stimulation (taVNS). Brain Stimul 11:947–948 Available at: https://doi.org/10.1016/j.brs.2018.06.003.

Bangasser DA, Wiersielis KR, Khantsis S (2016) Sex differences in the locus coeruleus-norepinephrine system and its regulation by stress. Brain Res 1641:177–188 Available at: http://www.ncbi.nlm.nih.gov/pubmed/26607253.

Barzegaran E, Vildavski VY, Knyazeva MG (2017) Fine Structure of Posterior Alpha Rhythm in Human EEG : Frequency Components , Their Cortical Sources , and Temporal Behavior. :1–12.

Bauer S, Baier H, Baumgartner C, Bohlmann K, Fauser S, Graf W, Hillenbrand B, Hirsch M, Last C, Lerche H, Mayer T, Schulze-Bonhage A, Steinhoff BJ, Weber Y, Hartlep A, Rosenow F, Hamer HM (2016) Transcutaneous Vagus Nerve Stimulation (tVNS) for

Treatment of Drug-Resistant Epilepsy: A Randomized, Double-Blind Clinical Trial (cMPsE02). Brain Stimul 9:356–363 Available at: http://dx.doi.org/10.1016/j.brs.2015.11.003.

Benjamini Y, Yekutieli D (2011) The Control of the False Discovery Rate in Multiple Testing under Dependency Source. Statistics (Ber) 29:1165–1188.

Bermejo P, López M, Larraya I, Chamorro J, Cobo JL, Ordóñez S, Vega JA (2017) Innervation of the Human Cavum Conchae and Auditory Canal : Anatomical Basis for Transcutaneous Auricular Nerve Stimulation. 2017.

Bodin C, Aubert S, Daquin G, Carron R, Scavarda D, McGonigal A, Bartolomei F (2015) Responders to vagus nerve stimulation (VNS) in refractory epilepsy have reduced interictal cortical synchronicity on scalp EEG. Epilepsy Res 113:98–103 Available at: http://dx.doi.org/10.1016/j.eplepsyres.2015.03.018.

Boon P, De Cock E, Mertens A, Trinka E (2018) Neurostimulation for drug-resistant epilepsy. Curr Opin Neurol 31:198–210 Available at: http://insights.ovid.com/crossref?an=00019052-201804000-00015.

Breton-Provencher V, Sur M (2019) Active control of arousal by a locus coeruleus GABAergic circuit. Nat Neurosci 22:218–228 Available at: http://www.ncbi.nlm.nih.gov/pubmed/30643295.

Bro R, Kiers HAL (2003) A new efficient method for determining the number of components in PARAFAC models. J Chemom 17:274–286 Available at: http://doi.wiley.com/10.1002/cem.801.

Brown SBRE, van der Wee NJA, van Noorden MS, Giltay EJ, Nieuwenhuis S (2015) Noradrenergic and cholinergic modulation of late ERP responses to deviant stimuli. Psychophysiology 52:1620–1631 Available at: http://doi.wiley.com/10.1111/psyp.12544.

Clark KB, Naritoku DK, Smith DC, Browning RA, Jensen RA (1999) Enhanced recognition

memory following vagus nerve stimulation in human subjects. Nat Neurosci 2:94–98

Available at: http://www.nature.com/doifinder/10.1038/4600.

Clewett D V., Huang R, Velasco R, Lee T-H, Mather M (2018) Locus Coeruleus Activity
Strengthens Prioritized Memories Under Arousal. J Neurosci 38:1558–1574 Available
at: http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.2097-17.2017.

Cong F, Lin Q, Kuang L, Gong X, Astikainen P (2015) Tensor decomposition of EEG
signals : A brief review. J Neurosci Methods 248:59–69 Available at:
http://dx.doi.org/10.1016/j.jneumeth.2015.03.018.

de Gee JW, Colizoli O, Kloosterman NA, Knapen T, Nieuwenhuis S, Donner TH (2017)
Dynamic modulation of decision biases by brainstem arousal systems. Elife 6 Available
at: http://www.ncbi.nlm.nih.gov/pubmed/28383284.

de Gee JW, Knapen T, Donner TH (2014) Decision-related pupil dilation reflects upcoming
choice and individual bias. Proc Natl Acad Sci 111:E618–E625 Available at:
http://www.pnas.org/lookup/doi/10.1073/pnas.1317557111.

Desbeaumes Jodoin V, Lespérance P, Nguyen DK, Fournier-Gosselin M-PP, Richer F (2015)
Effects of vagus nerve stimulation on pupillary function. Int J Psychophysiol 98:455–
459 Available at: http://dx.doi.org/10.1016/j.ijpsycho.2015.10.001.

Desmedt JE, Debecker J, Manil J (1965) [Demonstration of a cerebral electric sign associated
with the detection by the subject of a tactile sensorial stimulus. The analysis of cerebral
evoked potentials derived from the scalp with the aid of numerical ordinates]. Bull Acad
R Med Belg 5:887–936 Available at: http://www.ncbi.nlm.nih.gov/pubmed/5864251.

Dorr AE, Debonnel G (2006) Effect of Vagus Nerve Stimulation on Serotonergic and
Noradrenergic Transmission. 318:890–898.

Dragone A, Lasaponara S, Pinto M, Rotondaro F, De Luca M, Doricchi F (2018) Expectancy
modulates pupil size during endogenous orienting of spatial attention. Cortex 102:57–66
Available at: https://linkinghub.elsevier.com/retrieve/pii/S0010945217303179.

Drapeau C, Carrier J (2004a) Fluctuation of waking electroencephalogram and subjective

alertness during a 25-hour sleep-deprivation episode in young and middle-aged subjects.

Sleep 27:55–60 Available at: http://www.ncbi.nlm.nih.gov/pubmed/14998238.

Drapeau C, Carrier J (2004b) Fluctuation of waking electroencephalogram and subjective

alertness during a 25-hour sleep-deprivation episode in young and middle-aged subjects.

Sleep 27:55–60.

Ellrich J (2011) Transcutaneous Vagus Nerve Stimulation. Eur Neurol Rev 6:254 Available

at: http://www.touchneurology.com/articles/transcutaneous-vagus-nerve-stimulation.

Fang J, Egorova N, Rong P, Liu J, Hong Y, Fan Y, Wang X, Wang H, Yu Y, Ma Y, Xu C, Li

S, Zhao J, Luo M, Zhu B, Kong J (2017) Early cortical biomarkers of longitudinal

transcutaneous vagus nerve stimulation treatment success in depression. NeuroImage

Clin 14:105–111 Available at: http://dx.doi.org/10.1016/j.nicl.2016.12.016.

Follesa P, Biggio F, Gorini G, Caria S, Talani G, Dazzi L, Puligheddu M, Marrosu F, Biggio

G (2007) Vagus nerve stimulation increases norepinephrine concentration and the gene

expression of BDNF and bFGF in the rat brain. 79.

Frangos E, Ellrich J, Komisaruk BR (2015) Non-invasive access to the vagus nerve central

projections via electrical stimulation of the external ear: FMRI evidence in humans.

Brain Stimul 8:624–636 Available at: http://dx.doi.org/10.1016/j.brs.2014.11.018.

Gelbard-Sagiv H, Magidov E, Sharon H, Hendler T, Nir Y (2018) Noradrenaline Modulates

Visual Perception and Late Visually Evoked Activity. Curr Biol 28:2239-2249.e6

Available at: http://www.ncbi.nlm.nih.gov/pubmed/29983318.

Glover A, Ghilardi MF, Bodis-Wollner I, Onofrj M (1988) Alterations in event-related

potentials (ERPs) of MPTP-treated monkeys. Electroencephalogr Clin Neurophysiol

Potentials Sect 71:461–468 Available at:

https://linkinghub.elsevier.com/retrieve/pii/0168559788900500.

Grimonprez A et al. (2015) The antidepressant-like effect of vagus nerve stimulation is

mediated through the locus coeruleus. J Psychiatr Res 68:1–7 Available at: http://dx.doi.org/10.1016/j.jpsychires.2015.05.002.

Groves DA, Bowman EM, Brown VJ (2005) Recordings from the rat locus coeruleus during acute vagal nerve stimulation in the anaesthetised rat. Neurosci Lett 379:174–179.

Haegens S, Cousijn H, Wallis G, Harrison PJ, Nobre AC (2014) Inter- and intra-individual variability in alpha peak frequency. Neuroimage 92:46–55 Available at: http://dx.doi.org/10.1016/j.neuroimage.2014.01.049.

Hall M-H, Jensen JE, Du F, Smoller JW, O'Connor L, Spencer KM, Öngür D (2015) Frontal P3 event-related potential is related to brain glutamine/glutamate ratio measured in vivo. Neuroimage 111:186–191 Available at: https://linkinghub.elsevier.com/retrieve/pii/S1053811915001159.

Halliday R, Naylor H, Brandeis D, Callaway E, Yano L, Herzig K (1994) The effect of D-amphetamine, clonidine, and yohimbine on human information processing. Psychophysiology 31:331–337.

Hammond EJ, Uthman BM, Reid SA, Wilder BJ (1992) Electrophysiological Studies of Cervical Vagus Nerve Stimulation in Humans: I. EEG Effects. Epilepsia 33:1013–1020.

Hansen N (2017) The Longevity of Hippocampus-Dependent Memory Is Orchestrated by the Locus Coeruleus-Noradrenergic System. Neural Plast 2017.

Harshman RA (1970) Foundations of the PARAFAC procedure: Models and conditions for an" explanatory" multimodal factor analysis.

Hayat H, Regev N, Matosevich N, Sales A, Paredes-Rodriguez E, Krom AJ, Bergman L, Li Y, Lavigne M, Kremer EJ, Yizhar O, Pickering AE, Nir Y (2020) Locus coeruleus norepinephrine activity mediates sensory-evoked awakenings from sleep. Sci Adv 6:eaaz4232 Available at: http://www.ncbi.nlm.nih.gov/pubmed/32285002.

He W, Jing X-H, Zhu B, Zhu X-L, Li L, Bai W-Z, Ben H (2013) The auriculo-vagal afferent

pathway and its role in seizure suppression in rats. BMC Neurosci 14:85 Available at: http://bmcneurosci.biomedcentral.com/articles/10.1186/1471-2202-14-85.

Hein E, Nowak M, Kiess O, Biermann T, Bayerlein K, Kornhuber J, Kraus T (2013) Auricular transcutaneous electrical nerve stimulation in depressed patients: A randomized controlled pilot study. J Neural Transm 120:821–827.

Hulsey DR, Riley JR, Loerwald KW, Ii RLR, Kilgard MP, Hays SA (2017) Parametric characterization of neural activity in the locus coeruleus in response to vagus nerve stimulation. Exp Neurol 289:21–30 Available at: http://dx.doi.org/10.1016/j.expneurol.2016.12.005.

Jacobs HIL, Riphagen JM, Razat CM, Wiese S, Sack AT (2015) Transcutaneous vagus nerve stimulation boosts associative memory in older individuals. Neurobiol Aging 36:1860–1867.

Joshi S, Gold JI (2020) Pupil Size as a Window on Neural Substrates of Cognition. Trends Cogn Sci 24:466–480 Available at: https://linkinghub.elsevier.com/retrieve/pii/S1364661320300802.

Joshi S, Li Y, Kalwani RM, Gold JI (2016) Relationships between Pupil Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. Neuron 89:221–234 Available at: https://linkinghub.elsevier.com/retrieve/pii/S089662731501034X.

Keute M, Demirezen M, Graf A, Mueller NG, Zaehle T (2019) No modulation of pupil size and event-related pupil response by transcutaneous auricular vagus nerve stimulation (taVNS). Sci Rep 9:11452 Available at: http://www.nature.com/articles/s41598-019-47961-4.

Knyazeva MG, Barzegaran E, Vildavski VY, Demonet J (2018) Neurobiology of Aging of human alpha rhythm. Neurobiol Aging 69:261–273 Available at: https://doi.org/10.1016/j.neurobiolaging.2018.05.018.

Krahl SE, Clark KB (2012) Vagus nerve stimulation for epilepsy: A review of central mechanisms. Surg Neurol Int 3:S255-9 Available at: http://www.ncbi.nlm.nih.gov/pubmed/23230530.

Krahl SE, Clark KB, Smith DC, Browning RA (1998) Locus coeruleus lesions suppress the seizure-attenuating effects of vagus nerve stimulation. Epilepsia 39:709–714.

Kraus T, Hösl K, Kiess O, Schanze A, Kornhuber J, Forster C (2007) BOLD fMRI deactivation of limbic and temporal brain structures and mood enhancing effect by transcutaneous vagus nerve stimulation. J Neural Transm 114:1485–1493.

Kraus T, Kiess O, Hösl K, Terekhin P, Kornhuber J, Forster C (2013) CNS BOLD fMRI effects of sham-controlled transcutaneous electrical nerve stimulation in the left outer auditory canal - A pilot study. Brain Stimul 6:798–804 Available at: http://dx.doi.org/10.1016/j.brs.2013.01.011.

Kwon C-S, Ripa V, Al-Awar O, Panov F, Ghatan S, Jetté N (2018) Epilepsy and Neuromodulation—Randomized Controlled Trials. Brain Sci 8:69 Available at: http://www.mdpi.com/2076-3425/8/4/69.

Lewine JD, Paulson K, Bangera N, Simon BJ (2019) Exploration of the Impact of Brief Noninvasive Vagal Nerve Stimulation on EEG and Event-Related Potentials. Neuromodulation Technol Neural Interface 22:564–572 Available at: https://onlinelibrary.wiley.com/doi/abs/10.1111/ner.12864.

Liu J, Fang J, Wang Z, Rong P, Hong Y, Fan Y, Wang X, Park J, Jin Y, Liu C, Zhu B, Kong J (2016) Transcutaneous vagus nerve stimulation modulates amygdala functional connectivity in patients with depression. J Affect Disord 205:319–326 Available at: http://dx.doi.org/10.1016/j.jad.2016.08.003.

Liu Y, Rodenkirch C, Moskowitz N, Schriver B, Wang Q (2017) Dynamic Lateralization of Pupil Dilation Evoked by Locus Coeruleus Activation Results from Sympathetic, Not Parasympathetic, Contributions. Cell Rep 20:3099–3112 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/28954227.

Manta S, Dong J, Debonnel G, Blier P (2009) Enhancement of the function of rat serotonin and norepinephrine neurons by sustained vagus nerve stimulation. J Psychiatry Neurosci 34:272–280.

Manta S, Mansari M El, Debonnel G, Blier P (2013) Electrophysiological and neurochemical effects of long-term vagus nerve stimulation on the rat monoaminergic systems. :459–470.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. J Neurosci Methods 164:177–190.

Meij R Van Der, Ede F Van, Maris E (2016) Rhythmic Components in Extracranial Brain Signals Reveal Multifaceted Task Modulation of Overlapping Neuronal Activity. :1–28.

Miwakeichi F, Martínez-Montes E, Valdés-Sosa PA, Nishiyama N, Mizuhara H, Yamaguchi Y (2004) Decomposing EEG data into space-time-frequency components using Parallel Factor Analysis. Neuroimage 22:1035–1045 Available at: http://www.ncbi.nlm.nih.gov/pubmed/15219576.

Mridha Z, Gee JW de, Shi Y, Alkashgari R, Williams J, Suminski A, Ward MP, Zhang W, McGinley MJ (2019) Graded recruitment of pupil-linked neuromodulation by parametric stimulation of the vagus nerve. bioRxiv:2019.12.28.890111 Available at: http://biorxiv.org/content/early/2019/12/30/2019.12.28.890111.abstract.

Murphy PR, O'Connell RG, O'Sullivan M, Robertson IH, Balsters JH (2014) Pupil diameter covaries with BOLD activity in human locus coeruleus. Hum Brain Mapp 35:4140–4154 Available at: http://www.ncbi.nlm.nih.gov/pubmed/24510607.

Murphy PR, Robertson IH, Balsters JH, O'connell RG (2011) Pupillometry and P3 index the locus coeruleus-noradrenergic arousal function in humans. Psychophysiology 48:1532–1543.

Nemeroff CB, Mayberg HS, Krahl SE, McNamara J, Frazer A, Henry TR, George MS, Charney DS, Brannan SK (2006) VNS therapy in treatment-resistant depression: clinical evidence and putative neurobiological mechanisms. Neuropsychopharmacology 31:1345–1355 Available at: http://www.ncbi.nlm.nih.gov/pubmed/16641939.

Nieuwenhuis S, Aston-Jones G, Cohen JD (2005) Decision making, the P3, and the locus coeruleus-norepinephrine system. Psychol Bull 131:510–532 Available at: http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-2909.131.4.510.

Nieuwenhuis S, De Geus EJ, Aston-Jones G (2011) The anatomical and functional relationship between the P3 and autonomic components of the orienting response. Psychophysiology 48:162–175 Available at: http://www.ncbi.nlm.nih.gov/pubmed/20557480.

Nir Y, Tononi G (2010) Dreaming and the brain: from phenomenology to neurophysiology. Trends Cogn Sci 14:88–100 Available at: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2814941&tool=pmcentrez&rendertype=abstract [Accessed May 28, 2014].

Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011) FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. Comput Intell Neurosci 2011:1–9.

Palva S, Palva JM (2007) New vistas for α-frequency band oscillations. Trends Neurosci 30:150–158.

Peirce JW (2007) PsychoPy--Psychophysics software in Python. J Neurosci Methods 162:8–13 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17254636.

Peuker ET, Filler TJ (2002) The nerve supply of the human auricle. Clin Anat 15:35–37.

Pineda JA, Swick D (1992) Visual P3-like potentials in squirrel monkey: Effects of a noradrenergic agonist. Brain Res Bull 28:485–491 Available at: https://linkinghub.elsevier.com/retrieve/pii/036192309290051X.

Pineda JA, Westerfield M, Kronenberg BM, Kubrin J (1997) Human and monkey P3-like

responses in a mixed modality paradigm: effects of context and context-dependent

noradrenergic influences. Int J Psychophysiol 27:223–240 Available at:

https://linkinghub.elsevier.com/retrieve/pii/S0167876097000615.

Raedt R, Clinckers R, Mollet L, Vonck K, El Tahry R, Wyckhuys T, De Herdt V, Carrette E,

Wadman W, Michotte Y, Smolders I, Boon P, Meurs A (2011) Increased hippocampal

noradrenaline is a biomarker for efficacy of vagus nerve stimulation in a limbic seizure

model. J Neurochem 117:461–469 Available at: http://doi.wiley.com/10.1111/j.1471-

4159.2011.07214.x.

Redgrave J, Day D, Leung H, Laud PJ, Ali A, Lindert R, Majid A (2018) Safety and

tolerability of Transcutaneous Vagus Nerve stimulation in humans; a systematic review.

Brain Stimul 11:1225–1238 Available at: https://doi.org/10.1016/j.brs.2018.08.010.

Reimer J, McGinley MJ, Liu Y, Rodenkirch C, Wang Q, McCormick DA, Tolias AS (2016)

Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex.

Nat Commun 7:13289 Available at: http://www.nature.com/articles/ncomms13289.

Rohleder N, Nater UM (2009) Determinants of salivary alpha-amylase in humans and

methodological considerations. Psychoneuroendocrinology 34:469–485 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/19155141.

Rong P, Liu J, Wang L, Liu R, Fang J, Zhao J, Zhao Y, Wang H, Vangel M, Sun S, Ben H,

Park J, Li S, Meng H, Zhu B, Kong J (2016) Effect of transcutaneous auricular vagus

nerve stimulation on major depressive disorder: A nonrandomized controlled pilot study.

J Affect Disord 195:172–179 Available at: http://dx.doi.org/10.1016/j.jad.2016.02.031.

Roosevelt RW, Smith DC, Clough RW, Jensen RA, Browning RA (2006) Increased

extracellular concentrations of norepinephrine in cortex and hippocampus following

vagus nerve stimulation in the rat. Brain Res 1119:124–132.

Safi S, Ellrich J, Neuhuber W (2016) Myelinated Axons in the Auricular Branch of the

Human Vagus Nerve. Anat Rec 299:1184–1191.

Salinsky MC, Burchiel KJ (1993) Vagus Nerve Stimulation Has No Effect on Awake EEG

Rhythms in Humans. Epilepsia 34:299–304.

Sellaro R, van Leusden JWR, Tona K-D, Verkuil B, Nieuwenhuis S, Colzato LS (2015)

Transcutaneous Vagus Nerve Stimulation Enhances Post-error Slowing. J Cogn

Neurosci 27:2126–2132 Available at:

http://www.mitpressjournals.org/doi/10.1162/jocn_a_00851.

Siegel JM (2005) Chapter 10 - REM Sleep. In: Principles and Practice of Sleep Medicine

(Fourth Edition), Fourth Edi. (Kryger MH, Roth T, Dement WC, eds), pp 120–135.

Philadelphia: W.B. Saunders. Available at:

http://www.sciencedirect.com/science/article/pii/B0721607977500173.

Steenbergen L, Sellaro R, Stock AK, Verkuil B, Beste C, Colzato LS (2015) Transcutaneous

vagus nerve stimulation (tVNS) enhances response selection during action cascading

processes. Eur Neuropsychopharmacol 25:773–778 Available at:

http://dx.doi.org/10.1016/j.euroneuro.2015.03.015.

Stefan H, Kreiselmeyer G, Kerling F, Kurzbuch K, Rauch C, Heers M, Kasper BS, Hammen

T, Rzonsa M, Pauli E, Ellrich J, Graf W, Hopfengärtner R (2012) Transcutaneous vagus

nerve stimulation (t-VNS) in pharmacoresistant epilepsies: a proof of concept trial.

Epilepsia 53:e115-8 Available at: http://www.ncbi.nlm.nih.gov/pubmed/22554199.

Steriade M (2001) Impact of network activities on neuronal properties in corticothalamic

systems. J Neurophysiol 86:1–39 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/11431485.

Sun L, Peräkylä J, Holm K, Haapasalo J, Lehtimäki K, Ogawa KH, Peltola J, Hartikainen KM

(2017) Vagus nerve stimulation improves working memory performance. J Clin Exp

Neuropsychol 00:1–11 Available at:

https://www.tandfonline.com/doi/full/10.1080/13803395.2017.1285869.

Sutton S, Braren M, Zubin J, John ER (1965) Evoked-potential correlates of stimulus uncertainty. Science 150:1187–1188 Available at: http://www.ncbi.nlm.nih.gov/pubmed/5852977.

Szerb JC (1967) Cortical acetylcholine release and electroencephalographic arousal. J Physiol 192:329–343 Available at: http://www.ncbi.nlm.nih.gov/pubmed/6050151.

Takigawa M, Mogenson GJ (1977) A study of inputs to antidromically identified neurons of the locus coeruleus. Brain Res 135:217–230 Available at: http://www.ncbi.nlm.nih.gov/pubmed/922473.

Torsvall L, Akerstedt T (1987a) Sleepiness on the job: continuously measured EEG changes in train drivers. Electroencephalogr Clin Neurophysiol 66:502–511 Available at: http://www.ncbi.nlm.nih.gov/pubmed/2438115.

Torsvall L, Akerstedt T (1987b) Sleepiness on the job: continuously measured EEG changes in train drivers. Electroencephalogr Clin Neurophysiol 66:502–511.

Trevizol AP, Shiozawa P, Taiar I, Soares A, Gomes JS, Barros MD, Liquidato BM, Cordeiro Q (2016) Transcutaneous Vagus Nerve Stimulation (taVNS) for Major Depressive Disorder: An Open Label Proof-of-Concept Trial. Brain Stimul 9:453–454 Available at: http://dx.doi.org/10.1016/j.brs.2016.02.001.

Tu Y, Fang J, Cao J, Wang Z, Park J, Jorgenson K, Lang C, Liu J, Zhang G, Zhao Y, Zhu B, Rong P, Kong J (2018) A distinct biomarker of continuous transcutaneous vagus nerve stimulation treatment in major depressive disorder. Brain Stimul 11:501–508 Available at: https://doi.org/10.1016/j.brs.2018.01.006.

Van Bockstaele EJ, Peoples J, Telegan P (1999) Efferent projections of the nucleus of the solitary tract to peri-Locus coeruleus dendrites in rat brain: Evidence for a monosynaptic pathway. J Comp Neurol 412:410–428.

Van Leusden JWR, Sellaro R, Colzato LS (2015) Transcutaneous Vagal Nerve Stimulation (tVNS): A new neuromodulation tool in healthy humans? Front Psychol 6:2013–2016.

van Steenbergen H, Band GPH (2013) Pupil dilation in the Simon task as a marker of conflict processing. Front Hum Neurosci 7 Available at: http://journal.frontiersin.org/article/10.3389/fnhum.2013.00215/abstract.

van Stegeren AH (2008) The role of the noradrenergic system in emotional memory. Acta Psychol (Amst) 127:532–541 Available at: https://linkinghub.elsevier.com/retrieve/pii/S0001691807001278.

Varazzani C, San-Galli A, Gilardeau S, Bouret S (2015) Noradrenaline and dopamine neurons in the reward/effort trade-off: a direct electrophysiological comparison in behaving monkeys. J Neurosci 35:7866–7877 Available at: http://www.ncbi.nlm.nih.gov/pubmed/25995472.

Ventura-Bort C, Wirkner J, Genheimer H, Wendt J, Hamm AO, Weymar M (2018) Effects of Transcutaneous Vagus Nerve Stimulation (tVNS) on the P300 and Alpha-Amylase Level: A Pilot Study. Front Hum Neurosci 12:202 Available at: http://www.ncbi.nlm.nih.gov/pubmed/29977196.

Vonck K, Raedt R, Naulaerts J, De Vogelaere F, Thiery E, Van Roost D, Aldenkamp B, Miatton M, Boon P (2014) Vagus nerve stimulation. . .25 years later! What do we know about the effects on cognition? Neurosci Biobehav Rev 45:63–71 Available at: http://dx.doi.org/10.1016/j.neubiorev.2014.05.005.

Warren CM, Tona KD, Ouwerkerk L, van Paridon J, Poletiek F, van Steenbergen H, Bosch JA, Nieuwenhuis S (2018) The neuromodulatory and hormonal effects of transcutaneous vagus nerve stimulation as evidenced by salivary alpha amylase, salivary cortisol, pupil diameter, and the P3 event-related potential. Brain Stimul 12:1–8 Available at: https://doi.org/10.1016/j.brs.2018.12.224.

Warren CM, Wilson RC, van der Wee NJ, Giltay EJ, van Noorden MS, Cohen JD, Nieuwenhuis S (2017) The effect of atomoxetine on random and directed exploration in humans. PLoS One 12:e0176034 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/28445519.

Waschke L, Tune S, Obleser J (2019) Local cortical desynchronization and pupil-linked

arousal differentially shape brain states for optimal sensory performance. Elife 8

Available at: http://www.ncbi.nlm.nih.gov/pubmed/31820732.

Wilcoxon F (1945) Individual Comparisons by Ranking Methods. Biometrics Bull 1:80

Available at: https://www.jstor.org/stable/10.2307/3001968?origin=crossref.

Yakunina N, Kim SS, Nam E-C (2017) Optimization of Transcutaneous Vagus Nerve

Stimulation Using Functional MRI. Neuromodulation 20:290–300 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/27898202.

Yanagawa T, Chao ZC, Hasegawa N, Fujii N (2013) Large-scale information flow in

conscious and unconscious states: An ECoG study in monkeys. PLoS One 8:1–13.

Yap JYY, Keatch C, Lambert E, Woods W, Stoddart PR, Kameneva T (2020) Critical Review

of Transcutaneous Vagus Nerve Stimulation: Challenges for Translation to Clinical

Practice. Front Neurosci 14:284 Available at:

http://www.ncbi.nlm.nih.gov/pubmed/32410932.

16