Using Multilayer Heterogeneous Networks to Infer Functions of Phosphorylated Sites

*Joanne Watson[1,2], Jean-Marc Schwartz[1] and Chiara Francavilla[2]**

* Corresponding Author: <u>chiara.francavilla@manchester.ac.uk</u>

[1]Division of Evolution & Genomic Sciences, School of Biological Sciences, Faculty of Biology, Medicine & Health, University of Manchester, Manchester, M13 9PT

[2]Division of Molecular and Cellular Function, School of Biological Sciences, Faculty of Biology, Medicine & Health, University of Manchester, Manchester, M13 9PT.

1. Abstract

Mass spectrometry-based quantitative phosphoproteomics has become an essential approach in the study of cellular processes such as signaling. Commonly used methods to analyze phospho-proteomics datasets depend on generic, gene-centric annotations such as Gene Ontology terms which do not account for the function of a protein in a particular phosphorylation state. Analysis of phosphoproteomics data is hampered by a lack of phosphorylated site-specific annotations. We propose a method that combines shotgun phosphoproteomics data, protein-protein interactions, and functional annotations into a heterogeneous multilayer network. Phosphorylation sites are associated to potential functions using a random walk on heterogeneous network (RWHN) algorithm. We validated our approach against a model of the MAPK/ERK pathway and functional annotations from PhosphoSite Plus and were able to associate differentially regulated sites on the same proteins to their previously described specific functions. We further tested the algorithm on three previously published datasets and were able to reproduce their experimentally validated conclusions and to associate phosphorylation sites with known functions based on their regulatory patterns. Our approach provides a refinement of commonly used analysis methods and accurately predicts context-specific functions for sites with similar phosphorylation profiles.

Keywords: phosphoproteomics, functional analysis, bioinformatics, multilayer networks, random walk, gene ontology, pathways

2. Introduction

Phosphorylation is the most studied post-translational modification (PTM) due to its central role in cellular regulation. It is thought to be the principal PTM in the human proteome, as well as an essential mediator of protein-protein interaction (PPI) and protein function [1]. Transient changes occur at specifically regulated phosphorylation sites, of which there may be multiple on each protein. Regulation of phosphorylation is often dependent on perturbations such as the activity of extracellular ligands, drug treatment or physical stimuli in the extracellular environment [2]. By comparing changes in the phosphoproteome of cells in different experimental conditions through mass spectrometry-based phosphoproteomics, phosphorylated sites that are key players in cellular processes and functions can be uncovered in an unbiased, high-throughput manner [3].

Functional analysis of phosphoproteomics datasets is typically based on gene-centric enrichment of Gene Ontology (GO) terms or involvement in known pathways [4]. However, this approach disregards information captured by phosphoproteomics data on changes at specific phosphorylated sites, by limiting the analysis to the protein level. The modification state of a protein is inherently coupled to its function; PTMs alter protein activity, as well as the ability to interact with different sets of proteins. Furthermore, if a protein is phosphorylated on multiple sites, each with a different function and regulatory pattern, this information is not revealed by gene-centric analysis [5]. For instance, the well-studied signalling protein MAPK1 has 18 known phosphorylated sites recorded in the database PhosphositePlus® (PSP), however only 6 have been annotated to a downstream function[6]. In the most recent release of PSP (v6.5.9.3), only 4271 of the more than 230,000 human phosphorylated sites recorded in the database are associated to 19 generic biological functions (e.g., "cell cycle", "transcription"), which are qualified with one of: "induced", "inhibited", "regulated" or "altered". Enrichment analyses that rely on generalizations based on protein-level or gene-centric descriptions exclude the details that are encoded in the phosphorylation

signature. Analyses are thus hampered by the lack of phosphorylation site-specific functional an-

notations.

Several network-based methods have been proposed to move towards phosphorylation site-spe-

cific analyses of phosphoproteomics data. A significant focus has been on the inference of kinase-

substrate networks, such as NetworKIN [7], KSEA [8], and IKAP [9], among others. These methods are

useful for reconstructing the architecture of signalling intracellular networks, which can be in-

formative for identifying modules of modified proteins involved in cellular processes, but again

remain hampered by the lack of site-specific functional annotation [10]. They may also be biased

towards the most studied kinases and the exclusion of non-kinase proteins. Rudolph *et al*. [11] pro-

posed a method to address this issue named PHOTON which identified differentially regulated

proteins based on the level of phosphorylation of their binding partners in a high-confidence PPI

network, and then used logistic regression to identify the involvement of the phosphorylated pro-

teins in known signalling pathways. PHOTON is not truly site-centric, however, as it relies on

summarized quantitative values of phosphorylated sites. An alternative approach described by

Krug *et al*. [12] uses a modification to the gene-centric method gene set enrichment analysis (GSEA),

which relies on a curated resource containing literature-derived phosphorylated site-specific sig-

natures to assign functions to sites. However, the use of such a resource is limited by the lack of

phosphorylated-site annotation to these signatures or specific cellular processes; though useful for

identifying well-studied sites, predictions of functions of under-studied or sites in alternative con-

texts would not be captured. These do not fulfil the same role as popular gene-centric methods

such as over-representation analysis (ORA)[13] in aiding prediction and hypothesis generation when

performing exploratory analysis of shotgun phosphoproteomics data.

In recent years, heterogeneous or multilayer networks have been used to represent many types

of 'omics datasets [14]. These specialized networks are used to describe multiple types of associa-

tions with nodes representing different entities. To identify relationships between the different

biological layers, random walk algorithms have been applied to these networks. The random walk

on heterogeneous network (RWHN) with restart (RWR) method [15] has been particularly popular.

Jiang [16] used RWR to prioritize disease candidate genes in a PPI-phenome network; similarly,

Soul *et al*. [17] applied it to a PPI-phenome network to identify disease mechanisms. Recent work

has extended the method to multiple layers of biological information, for example to infer disease

associated m6A RNA methylation via known gene-disease associations [18]. Similar methodology

was used to associate phosphorylation sites recorded in the PSP database to diseases via kinase-

substrate interactions [19].

Here, we propose an algorithm that uses RWHN to associate phosphorylated sites to context-

specific function via a heterogeneous multilayer network using shotgun phosphoproteomics data.

The network combines three layers of information: phosphorylated sites, protein interactions and

GO terms. Clustering of phosphoproteomics data is used to find common features within datasets

and is generally followed by enrichment analyses. This is based on the assumption that common

patterns of phosphorylation, based on temporal changes or in response to a particular stimulus,

treatment or environmental context, are a likely indicator of involvement in common functions or

processes [20]. We utilize this concept in our algorithm by connecting phosphorylated sites that

have been clustered together, and therefore share regulatory patterns, within the multilayer net-

work. The algorithm is intended for use on large phosphoproteomics datasets, assessing perturba-

tions or processes as opposed to an interpretation of the full phosphoproteomics network in the

cell.

We first apply our algorithm to a small-scale, manually curated validation network. We also

assess the ability of our method and the most commonly used alternative, ORA, to capture the

functional descriptions recorded in PSP, which are based on experimental analysis. We then

demonstrate the utility of our algorithm using three previously published shotgun phosphoprote-

omics datasets, describing early signalling events in HeLa cells upon EGF and TGF-α stimulation

[21], phosphorylation-mediated changes in breast cancer cells resistant to lapatinib treatment [22] and

subcellular location dependent signalling events downstream of HRAS [23], respectively. We

demonstrate that phosphorylated sites can be differentially assigned to functional annotations and

this is driven by changes in the context-dependent modification of these sites. Our method is suit-

able for use with any phosphoproteomics dataset and could be generalized for data describing

other PTMs.

101   **3.   Experimental Procedures**

A multilayer heterogeneous network was constructed to relate biological function to phosphor-

ylation sites via a protein-protein interaction network (Figure 1). Three types of nodes are con-

tained within the network: phosphorylation sites (called 'sites' here on in for brevity), proteins

and functional annotations (either Gene Ontology Biological Process (GOBP) terms or KEGG

pathways). The edges describe five possible associations which are either bipartite (i.e., site to

protein, protein to function) or between the same type of node.

Figure 1. Overview of multilayer heterogeneous network construction from phosphoproteomics

data.

Dataset Accession and Processing

Sites annotated to have a role in biological processes were extracted from the `Regulatory sites`

dataset, available for download from PhosphoSitePlus® (version 6.5.9.3). The dataset was filtered

on the "ON_PROCESS" column to only include sites annotated to biological roles, rather than

molecular or interaction roles.

The results of the phosphoproteomics experiment described in Francavilla *et al*. [21] were taken

from Supplementary Table S1 of their manuscript. The authors had processed the raw mass spec-

trometry data using MaxQuant [24], filtered identified phosphorylated sites based on a localization

probability of greater than 0.75 and normalized remaining data to the ratio of EGFR at 1 min

after stimulation with EGF or TGF-α. Prior to multilayer heterogeneous network construc-

tion, the data were further filtered to remove sites with two or more missing ratios in their

time series and containing at least one SILAC ratio higher than 2 or lower than 0.5 (as de-

scribed in the methods of Francavilla *et al.* [21]). Missing data were imputed with random draws

from a truncated distribution, as previously described [25], using the *impute.QRLIC* function from

the *imputeLCMD* R package.

Data from the Ruprecht *et al*. [22] study were taken from the Supplementary Data file named

'Filtered and normalized phosphoproteome dataset'. The raw mass-spectrometry data had been

processed using MaxQuant, filtered to include only identified phosphorylated sites with a locali-

zation probability of greater than 0.75, and normalized. Sites that did not show a significant change

(FDR <1%) either between the untreated parental and lapatinib-treated resistant (SILAC ratios

H/L) or between the lapatinib treated parental and lapatinib-treated resistant conditions (SILAC

ratios H/M) experimental conditions were filtered from the data, as described in the methods of

Ruprecht *et al*. [22]. Missing data were imputed with the same method used for the Francavilla *et*

al. dataset.

Data from the Santra *et al*. study [23] were taken from Supplemental table S2. The raw mass-

spectrometry data had been processed using MaxQuant, filtered to include only identified phos-

phorylated sites with a localization probability of greater than 0.75; significant sites were identi-

fied using a two-sample t-test with FDR. The downloaded dataset was normalized using the *limma*

R package function *normalizeBetweenArrays* using the quantile method [26]. Missing data were

imputed with the same method used for the Francavilla *et al*. dataset. The data were then filtered

to exclude sites that had not been identified by significant in the original analysis by Santra *et al*.

Before applying the algorithm, all datasets were clustered based on what was reported in the

original publication, where available. The model dataset was clustered using the Fuzzy C-Means

(FCM) method [27], with the number of clusters selected based on the temporal trends we could

visually identify in the data; generally, silhouette analysis is recommended for selecting the num-

ber of clusters using FCM for real experimental datasets [28]. The sites extracted from PSP were

clustered based on what process they were annotated to (as indicated in the "ON_PROCESS"

column in the Regulatory_sites file available for download on the PSP website). The regulated

phosphorylated sites from Francavilla et al. were clustered using FCM (Figure 3); since the orig-

inal publication used a cluster number of 6, the same was done here for better comparison. The

regulated phosphorylated sites from Ruprecht et al. and Santra et al. were both clustered using the

k-means method, with the cluster number selected using the elbow method (Figure S4 and S5,

respectively) [29].

Construction of Multilayer Network
Phosphorylation Site – Phosphorylation Site Subnetwork
Edges were drawn between sites in the same cluster that had a Pearson correlation (R2) between

all the data points greater than or equal to 0.99 (Figure 1).

Protein - Protein Subnetwork
The protein-protein interaction network was extracted from STRING [30]. All interactors of the

proteins included in the phosphoproteomics datasets with an experimental confidence score of

greater than 0.4 were included (Figure 1).

Function – Function Subnetwork
We used either GOBP terms or KEGG pathways as functional annotators in this work. If GOBP

terms were included in the multilayer network, the *GOSemSim* package from Bioconductor [31] was

used to calculate the semantic similarity of enriched GOBP terms. An edge was drawn between

terms with semantic similarity greater than 0.7, as calculated using the Wang method included in

the *goSim* function of *GOSemSim* [31,32]. In the case of KEGG pathways, edges were drawn between

pathways that had greater than 70% pairwise similarity in their functional annotation profiles,

following the method described in Stoney *et al*. [33] (Figure 1).

 Phosphorylation Site – Protein Bipartite Edges
Sites and proteins had an edge between them if the residue was found on that protein. Therefore,

sites will only have one edge, but proteins will have edges to all the sites found on that protein

that were differentially regulated in the data set (Figure 1).

Protein – Function Bipartite Edge
We assumed that closely connected nodes in the protein-protein subnetwork would more likely

be involved in similar biological processes. Therefore, we computed modules of the protein-pro-

tein subnetwork using the Louvain module-detection method [34]. Proteins from each module were

analyzed for enrichment (hypergeometric test with Benjamini-Hochberg correction, FDR <

0.05%) of functional annotations (from either the "GO_Biological_Process_2018" or

"KEGG_2019_Human" libraries, included in enrichR and listed on the EnrichR website) using

the *enrichR* [35] R interface. This increased the specificity of terms to be included in the network.

When GO terms were included, high frequency (annotated to more than 5% of genes) and seman-

tically redundant terms (similarity > 0.9) were filtered using the Bioconductor *GOSemSim* [31] and

GO.db packages [36] (Figure 1).

Random Walk on Heterogeneous Network
188   The heterogeneous network can be represented as an adjacency matrix as follows:

$$\mathrm{A} = \begin{bmatrix} A_{RR} & A_{RP} & 0 \\ A_{PR} & A_{PP} & A_{PF} \\ 0 & A_{FP} & A_{FF} \end{bmatrix} \tag{1}$$

189   where: A_{RR} are site-site associations, A_{PP} are protein-protein associations, A_{FF} are function-func-

190   tion associations, A_{RP} are site-protein associations with A_{PR} as the transpose and A_{PF} are protein-

191   function associations with A_{FP} as the transpose.

192   As described in previous work [15], a transition matrix (M) was calculated for use in the first stage

193   of the algorithm:

$$M = \begin{bmatrix} M_{RR} & M_{RP} & 0 \\ M_{RP^T} & M_{PP} & M_{PF} \\ 0 & M_{PF^T} & M_{FF} \end{bmatrix} \qquad (2)$$

194   The bipartite inter-subgraph transition matrices (M_{RP} and M_{PF}) were calculated as:

$$(M_{RP})_{i,j} = \begin{cases} \lambda\,(A_{RP})_{i,j} \Big/ \sum_j (A_{RP})_{i,j}, & if \sum_j (A_{RP})_{i,j} \neq 0 \\ 0, & otherwise \end{cases} \qquad (3)$$

$$(M_{PF})_{i,j} = \begin{cases} \lambda\,(A_{PF})_{i,j} \Big/ \sum_j (A_{PF})_{i,j}, & if \sum_j (A_{PF})_{i,j} \neq 0 \\ 0, & otherwise \end{cases} \qquad (4)$$

195   where λ is the transition probability (i.e., the likelihood of the walker moving between two lay-

196   ers of the network).

197   The intra-subgraph transition matrices (M_{RR}, M_{PP} and M_{FF}) were calculated as:

$$(M_{RR})_{i,j} = \begin{cases} (A_{RR})_{i,j} \Big/ \sum_j (A_{RR})_{i,j}, & if \sum_j (A_{RP})_{i,j} = 0 \\ (1 - \lambda)\,(A_{RR})_{i,j} \Big/ \sum_j (A_{RR})_{i,j}, & otherwise \end{cases} \qquad (5)$$

$$(M_{PP})_{i,j} = \begin{cases} (A_{PP})_{i,j} \Big/ \sum_j (A_{PP})_{i,j}, & if \sum_j (A_{PR})_{i,j}, (A_{PF})_{i,j} = 0 \\ (1 - \lambda)\,(A_{PP})_{i,j} \Big/ \sum_j (A_{PP})_{i,j}, & otherwise \end{cases} \qquad (6)$$

$$(M_{FF})_{i,j} = \begin{cases} (A_{FF})_{i,j} \Big/ \sum_j (A_{FF})_{i,j}, & if \sum_j (A_{FP})_{i,j} = 0 \\ (1 - \lambda)\,(A_{FF})_{i,j} \Big/ \sum_j (A_{FF})_{i,j}, & otherwise \end{cases} \qquad (7)$$

198   RWHN is a ranking algorithm; nodes are ranked based on the probabilities of finding the ran-

199   dom walker at a given node in the steady state, having started at a given seed node or set of seed

200   nodes. In this work we set the seed nodes to be those sites belonging to a particular cluster. The

201   probability of finding the random walker at each node for each step is calculated based on the

202   iterative equation:

$$\begin{cases} P_{s+1} = (1-r) \times M^T \times P_s + r \times P_0 & \quad\quad\quad (8) \\ \quad\quad P_{s+1} - P_s \leq 10^{-6} \end{cases}$$

where r is the restart probability (set to 0.7 as described in Kohler *et al.* [37]), P_0 is the initial

probability vector and P_s is the probability vector at step s. P0 was calculated such that all seed

nodes were given equal probabilities with their sum equal to 1. All other nodes in the site-site

subnetwork were assigned an initial probability of 0. Nodes in other subnetworks were assigned

equal probabilities with their sum equal to 1 and weighted with the tunable parameters ηP and ηF,

that are used to weight the influence of each layer.

The output of the algorithm is a ranked list of all the nodes, based on the probability of finding

the random walker at each node in the steady state. This list is filtered by retaining functional

annotations and removing proteins and sites. We remove annotations that are ranked in the same

position regardless of seed node, in order to focus on specific functions related to each cluster.

The top 5% of functions in the probability distribution are retained.

We implemented the algorithm in R using packages available from Bioconductor and CRAN.

The source code is available at www.github.com/jowatson2011/RWHN_phosphoproteomics and

an R package for general use of the tool can be found at github.com/JoWatson2011/phosphoR-

[WHN](WHN). RWHN on the validation network took less than 30 seconds to run however the larger

experimental data sets took several hours on a moderately powerful computer (32GB RAM, Intel

i7-4770 Processor).

Overrepresentation Analysis (ORA)

The names of modified proteins in each cluster were used as input for the *enrichR* R interface

[35]. The libraries "GO_Biological_Process_2018" or "KEGG_2019_Human" (included in enrichR

and listed on the EnrichR website) were used depending on which was used for the RWHN

analysis of the same data. The list of overrepresented terms was filtered to only include those with

an FDR < 0.05.

4. Results

Overview of Algorithm

To associate phosphorylated sites of unknown function to potential cellular functions we devel-

oped an algorithm to apply to shotgun phosphoproteomics data. First, in order to help experimental

biologists to identify the context- or perturbation-specific roles of modified proteins changing in

each given experiment, a multilayer heterogeneous network is constructed based on the regulation

of phosphorylated sites. Regulated phosphorylated sites may be defined differently by the user

depending on the quantitation method used in their phosphoproteomics experiment (e.g., label-

free quantification, SILAC) or their experimental question. The network represents three layers

of biological entities and information: regulated phosphorylated sites, protein-protein interaction,

and biological function (Figure 1). We then apply a ranking algorithm, RWHN, which ranks nodes

of each layer based on: (i) distance from the phosphorylated sites of interest, which are assigned

as 'seed' nodes and (ii) the topology of the multilayer heterogeneous network [15]. Functions that

are highly ranked can be considered more correlated with a set of seed nodes.

In the multilayer heterogeneous network, edges are drawn between sites based on similarity in

the regulatory pattern of their phosphorylation which we determine using k-means or fuzzy C-

means clustering. For the protein layer, a PPI network is constructed of the phosphorylated pro-

teins and their interactors using the STRING database [30]. Interactors are included to account for

non-phosphorylated proteins, or those below the limit of detection of the experiment [38]. To get a

comprehensive and specific list of functional annotations, we identified closely connected groups

of proteins in the PPI network using module detection and performed functional enrichment

analysis on these modules. Edges between functional annotations are drawn based on functional

similarity and overlap [31,33].

The RWHN algorithm simulates a walker moving from a starting node(s) (called a seed) and

then node to node through the multilayer network. Each step is influenced by the probability of

transition to another layer (λ), the weighting of the protein and function layers (η_P and η_F) and the

probability of restart (that is, teleportation back to the seed node(s), r). The output of RWHN

consists in a list of ranks for all the nodes in the network based on the likelihood of the walker

reaching that node.

To optimize the algorithm, we ran RWHN with each of these parameters (λ, r, η_P and η_F) tested

over a range of values (0.1 – 0.9), changing one whilst setting all others to 0.5 (Figure S1A).

Performance was decidedly stable over the range of η_P and η_F however altering λ and r resulted

in differential ranking depending on the parameter value. Previous work has suggested an ideal λ

and r of 0.7 [15,37,39], whilst η_P and η_F were set to 0.7 and 0.3 respectively to prioritize movement in

the protein layer and reduce the number of terms having the same rank regardless of seed.

The parameters used in the construction of the PPI network were also assessed. Proteins inter-

acting with the phosphorylated proteins found in a given sample were extracted from STRING;

those proteins whose interaction had a STRING experimental confidence score above a given

threshold were included in the multilayer heterogeneous network. To investigate the impact of

choosing different STRING confidence scores on the results, we tested a confidence score range

of 0.1 – 0.9. First, we looked at whether connectivity of the graph was altered by the score. As

expected, the greater the confidence score, the more components the PPI were split into; above a

score of 0.7, the PPI had two or more components (Figure S1B). We next investigated whether

the RWHN results were affected by changing the STRING confidence score increasing

components within the protein layer. Using weighted-tau correlation[40] to compare the rankings

from each set of results, we found that the correlation was generally in the range of 0.8 – 1.0, with

the lowest value being 0.54 between networks constructed with 0.1 or 0.9 confidence threshold,

which were the extremes of the range tested.

Validation 1: Model of MAPK/ERK pathway

MAPK/ERK signalling has been well studied and the temporal phosphorylation status of com-

ponent proteins in response to growth factor stimulus are relatively established [41,42](Figure 2A).

Based on this general understanding, we compiled a simple validation dataset of phosphorylation

dynamics at particular sites of the main signalling proteins within the pathway (Figure 2B). Sites

were chosen based on whether their phosphorylation is known to activate or inactivate protein

activity, as recorded in the PSP database [6]. We clustered the data into 5 clusters, referred to here

as clusters 1-5, using the fuzzy c-means algorithm (Figure 2B). The features of the multilayer

network that was constructed are summarized in Table 1. Enrichment of GOBP terms was used

to form the functional annotation layer of the network.

We ran the RWHN algorithm over the multilayer heterogeneous network with seed nodes set to

all the sites belonging to one of the clusters; this was then repeated for each cluster (Figure 2C).

The highest ranked GOBP term was the same ("protein autophosphorylation") regardless of the

seed nodes. However, it was possible to differentiate between the five clusters based on the rank-

ing of terms below the first one.

Table 1. Multilayer heterogeneous network, constructed from MAPK/ERK validation data. GOBP

terms were included as functional annotations.

Subnetwork	Edges	Nodes

Site-site	28	19
Protein-protein	932	313
Function-function	148	88
Site-protein	19	19 sites, 8 proteins
Protein-function	1796	261 proteins, 298 functions

Figure 2. Site- and gene-centric analysis of validation data. A) Construction of validation data was based on a traditional representation of the MAPK/ERK pathway. B) A validation dataset was constructed based on the MAPK/ERK pathway, simulating phosphorylation dynamics over

time after pathway activation. The sites were clustered using fuzzy c-means clusters. These data

were used to construct a heterogeneous multilayer network. C) RWHN results with sites annotated

to functional roles in PSP; seed nodes are equivalent to the clusters. D) ORA results with modified

proteins contained in toy data. E) GO hierarchy, which has been subset to include only the parents

of terms that occurred in either the RHWN or ORA analysis. Terms that occur in either set of

results are colored in purple or green, respectively, with terms that occur in both colored in light

blue.

Where available, we could use the biological process annotations in PSP (v6.5.9.3) as a 'bench-

mark' for the highly ranked terms associated to each cluster. RPS6KA3_Y529, annotated to

"apoptosis, altered" in PSP is found in cluster 2, along with DUSP6 sites; this cluster highly ranks

"positive regulation of programmed cell death". Of the four RAF1 sites included in this model

dataset, two were annotated to "cell cycle regulation" in PSP: RAF1_S289, in cluster 2, and

RAF1_S338, in cluster 5. These two clusters both rank "regulation of mitotic cell cycle" in the

top 5% of terms.

Largely due to the inherent step in our method in which the redundant terms are reduced in the

multilayer heterogeneous network, fewer terms were found in the results of RWHN than ORA.

We next wanted to assess whether the terms in the RWHN result were drawn from a particular

level of the GO hierarchy; if they were, this would suggest a bias towards more specific or more

broad terms compared to ORA. From the GO hierarchy, we extracted the parent and ancestor

terms of the terms found in each set of the results (Figure 2E). We found that RWHN terms were

drawn from the same range of the hierarchy as the ORA terms, with slightly more coming from

lower in the hierarchy, indicating the results have more specificity than the ORA results. We note

that there is also less redundancy in the RWHN results, despite the ORA results also being reduced

for semantically similar / redundant terms. For example, the terms "regulation of ERK1 and ERK2

cascade" and "ERK1 and ERK2 cascade" both come up in the ORA results and are associated to

different clusters.

To assess the robustness of the algorithm, we performed a random permutation control. RWHN

was run one hundred times with the seeds set as before, random permutations of the subnetworks

and bipartite edges maintained. Kernel Density Estimation (KDE) was calculated to assess how

often each GO term occurred at each rank in the 100 random permutations for each set of seed

nodes. In each case, there were a subset of terms that were more likely to have a rank greater than

50, however none strongly correlated with the actual rankings for each set of seed nodes (Figure

S2A). This confirms that the rankings are not random, but primarily determined by the network

topology. We next investigated the rate of potential false positives in the results. Previous imple-

mentations of RWHN, primarily using genomics data, have utilized an arbitrary threshold of the

top x ranked terms, or the top $x\%$ of ranked terms (based on the probability vector as calculated

in equation 8). A threshold that is too stringent would not remove lower-ranking terms from the

results, whilst a too-low threshold would increase the amount of noise in the data. Using the results

of RWHN run on the randomly permuted networks we calculated the probability that the "true"

ranking (i.e. from RWHN run on the non-permuted network) was different from the random rank-

ings ($p < 0.05$, Mann-Whitney U test with Benjamini-Hochberg adjustment [43,44]). We found that

most of the rankings differed from random (Figure S2B). The terms that had $p > 0.05$ were

checked against the RWHN results with different thresholds selected (top 1%, 5%, 10% and 15%

of ranked terms); none were found above the 1% thresholds for any of the clusters, and only one

term ("peptidyl-serine modification" in cluster 2) was found above the 5% and 10% thresholds

(Figure S2B). We chose to set the default threshold to the top 5% of terms to be more conservative

whilst retaining useful annotations.

For a given set of seed nodes, the algorithm is capable of giving high ranks to functions with

known associations to those sites. It is also capable of predicting non-random, reasonable func-

tions for sites of unknown function in this context.

Validation 2: Classification of PhosphoSitePlus annotated sites
A comprehensive benchmark of dynamically regulated phosphorylated sites is not currently

available, due to the lack of phosphorylated sites with a regulatory pattern universally defined

independent of the experimental conditions. Therefore, we used the static classifications of phos-

phorylated site function recorded in PSP to benchmark the predictive power of our algorithm. The

PSP database has annotations to biological processes for ~4000 human phosphorylated sites. As

these annotations are non-standard, each one was mapped to the most closely related GOBP term,

where available (Table S1). The majority of sites were annotated to signal transduction, gene ex-

pression or cell differentiation (Figure 3A). Of these, 1496 sites were annotated to more than one

function; as these annotations are manually assigned based on literature searches, the multiple

functions may be due to the differences in experimental questions in the source datasets. The sites

were clustered based on which function they were annotated to and analyzed with RWHN and

ORA. The GOBP terms in the results of both methods were associated to the child terms of each

mapped PSP GO term, per the GO hierarchy. The terms in the results of both methods showed a

similar distribution to the terms in PSP as a whole, the main exception being in the cell differen-

tiation category (Figure 3B). The two methods showed a considerable difference in the number of

terms in the output, with ORA resulting in several hundred more terms in the output than RWHN.

To test whether the two sets of results were drawn from the same parts of the GO hierarchy (with

those from the top of the hierarchy being broader and those at the bottom being more specific),

we constructed a tree based on the GO hierarchy. The tree was then divided into levels, as in

Figure 2E, to capture terms that were of similar descriptiveness. Finally, we counted the number

of terms that came up in the RWHN or ORA results (Figure 3C). Both sets of results draw from

the same part of the hierarchy, with a skew towards the broader terms. We therefore concluded

that RWHN resulted in as descriptive but more restricted and focused annotations than ORA. This

was notable given that RWHN did not associate any terms to the smaller categories in PSP, such

as exocytosis, endocytosis, cell growth and autophagy (Figure 3B).

Figure 3. Distribution of annotations and classifications of sites extracted from PhosphoSitePlus. A) Distributions of functionally annotated sites (to mapped GO term). B) Distribution of terms in ORA results and RWHN results when analyzing functionally annotated sites from PSP. C) Number of terms from ORA or RWHN that are from different levels of the GO hierarchy

Experimental Case Study 1: Dissecting EGF- and TGF-α-Induced Dynamic Phosphory-lation

Given that PSP contains static annotations of sites from various sources, we wanted to test our

algorithm on experimental datasets with controlled perturbations. We reasoned that this would

allow us to assess whether our algorithm could differentially associate sites to their context-de-

pendent function. The phosphoproteomics data retrieved from Francavilla *et al*. [21] describes the

effect in HeLa cells of stimulation with EGF or TGF-α over a period of 90 minutes (with time

points at 1, 8, 40 and 90 minutes). In this study, one of the findings was that EGF or TGF-α

stimulation of EGFR induce receptor degradation or recycling, respectively. TGF-α was associ-

ated with a more potent mitogenic and migratory response than EGF over time [21]. Further details

on the experimental design and results from the publication can be found in Supplementary Table

S2. Phosphorylated sites from Supplementary Table S1 of the original publication were filtered

based on regulation by EGF or by TGF-α and divided based on which stimuli the regulation was

dependent on. The separated data were clustered into 6 clusters using the fuzzy C-means method,

as per the original publication, with each cluster representing a distinct dynamic profile of phos-

phorylation (Figure S3). A multilayer heterogeneous network was constructed for both sets of

data, described in Table 2 and Table 3, respectively.

Table 2. Multilayer heterogeneous network, constructed from Francavilla *et al*. [21] EGF-regulated

phosphorylated sites. GOBP terms were included as functional annotations.

Subnetwork	Edges	Nodes
Site-site	4347	733
Protein-protein	16654	4072
Function-function	532	544
Site-protein	733	733 sites, 433 proteins
Protein-function	1223	115 proteins, 532 functions

398   Table 3. Multilayer heterogeneous network, constructed from Francavilla *et al*. [21] TGFα-regulated

399   phosphorylated sites. GOBP terms were included as functional annotations.

Subnetwork	Edges	Nodes
Site-site	3085	675
Protein-protein	16556	3960
Function-function	332	496
Site-protein	675	675 sites, 403 proteins
Protein-function	1015	109 proteins, 486 functions

Several biologically relevant terms were differentially ranked in the EGF and TGF-α networks

(Figure 4A,C). Consistent with the original publication, the term "regulation of ERK1 and ERK2

cascade" was highly ranked using RWHN when seeds were set to sites in cluster 2 (representing

EGF late responders) for the EGF network and in cluster 1 (representing TGF-α early responders)

for the TGF-α network. Despite their central role to TGF-α signalling, when we performed

standard ORA enrichment terms relating to MAPK/ERK were not found in any of the clusters

(Figure 4D). However, both ORA and our method highlighted the role of the proteins belonging

to TGF-α cluster 1 in regulating EGFR/ERBB signaling, with terms referring to regulation of

these pathways ranked highly (Figure 4C,D). We used cluster 2 in the EGF condition and cluster

1 in the TGF-α condition to verify the robustness and accuracy of our approach. As the

MAPK/ERK cascade is well studied in respect to EGF/TGF-α signaling, many of its associated

sites have documented functions that we could use to benchmark our results. EGF cluster 2 con-

tains sites of several known players of the MAPK/ERK cascades, including RSP6KA3_S375,

SOS1 (T1119, S1137), RAF1 (S289, S96, S301), JUN_S243 and JUNB_259. All of these sites

are predicted MAPK1/MAPK3 targets, with the exception of SOS1_T1119, which is a MAP2K1

target [7]. As this cluster represents those sites phosphorylated late in the time course, we would

expect ERK1/2 feedback regulation to emerge. Besides confirming the term "regulation of ERK1

and ERK2 cascade" in EGF cluster 2, results from the RHWN algorithm add detail to this picture,

with "establishment of protein localization to organelle" ranked highly in this cluster (Figure 4A).

This finding confirms the conclusion of Francavilla *et al*. of EGFR trafficking to the lysosome,

and subsequent degradation, when stimulated by EGF [21]. EGF cluster 2 is the only one which did

not highly rank "regulation of clathrin-dependent endocytosis" compared to the 5 other clusters;

as this cluster represents sites that are highly phosphorylated late in the time-course, this coincides

with the understanding of receptor endocytosis as a process regulated in the early stage. This clus-

ter is one of the five that highly ranks "regulation of cell migration" (Figure 4A); PSP also anno-

tates several of the sites in this cluster to cell motility (PTPN12_S571, SLC9A1_S703,

FLNA_S2152).

Figure 4. Site-centric and gene-centric analysis of Francavilla *et al*.[21] datasets. (A) RWHN results related to EGF regulated sites. (B) ORA results related to proteins modified upon EGF stimulation. (C) RWHN results related to TGF-α regulated sites. (D) ORA results related to proteins modified upon TGF-α stimulation.

TGF-α cluster 1, on the other hand, contains sites that are phosphorylated within the first eight minutes upon TGF-α stimulation. Accordingly, this cluster contains different members of the MAPK/ERK cascade, such as the canonical activating sites of MAPK1, MAPK3 and MAPK14 (also known as p38) and several sites on EGFR (EGFR_S991, EGFR_S995, EGFR_T693). EGFR_S991. A few of these sites, such as EGFR_T693, are annotated to "receptor internalization" in PSP. This cluster also contains several phosphorylated sites on known players in receptor trafficking, including RAB7A_Y183 and RAB11B_Y8 which are highlighted in the original publication [21], along with CBL_Y674 and SH3BP4_T118. Our algorithm ranked terms such as "endosomal transport,", "establishment of protein localization to organelle" and "rab protein signal transduction" high when seed nodes were set to the sites found in this cluster (Figure 4C). This was only partially captured by the ORA (Figure 4D), which showed enrichment of the term "regulation of vesicle mediated transport" but also "protein localization to the lysosome", the opposite of the experimentally derived conclusion of Francavilla *et al*.

EGF cluster 1 representing phosphorylated cycling sites ranks both "regulation of cell migration" and "positive regulation of cell migration" in the top 5% of terms but did not have a similar association with trafficking terms as TGF-α cluster 1 in the RWHN output. Consistent with Francavilla *et al*., we concluded that there was not an association between trafficking and regulation of migration upon EGF stimulation [21]. Upon further investigation of this cluster, we found several

sites that were shared with TGF-α cluster 1, such as the MAPK1/3/14 sites, EGFR_S991 and

EGFR_T693. We also found the inhibitory RAF1 site S259 and SCRIB_1448. SCRIB has a pu-

tative role in cell migration [45,46].

TGF-α clusters 4 and 5 also highly rank terms related to intracellular trafficking. In the ORA

analysis (Figure 4D) cluster 4 was also enriched for "regulation of clathrin-dependent endocyto-

sis", corroborating this association. Of particular interest in cluster 4 are the EGFR (Y1110,

Y1125 and Y1197), CBL (Y700) and CBLB (Y665. Y763. Y889) sites. None of the CBL/CBLB

sites have roles described in PSP, but these proteins are known regulators of receptor in trafficking

via clathrin [47,48]. Moreover, EGFR_Y1110 and EGFR_Y1197 are both annotated to receptor inter-

nalization in PSP, as well as inducing cell motility. This is in line with "regulation of epithelial

cell migration" highly ranked in this cluster (Figure 4C). Taken together, this data indicates that

TGF cluster 4 sites may have an initial role in regulating receptor internalization and migration,

whilst the cluster 1 sites are regulating later parts of the process.

As a positive control for assignment of functions to clustered sites, sites annotated in PSP to the

function "cell growth, inhibited" were added as a "spike-in" cluster to the EGF and TGF-α data,

with sites already found in the data excluded from this new cluster. RWHN and ORA were then

applied as previously. We found that both RWHN and ORA could assign these sites to the function

"negative regulation of cell growth" (Figure 5A). However, ORA assigned 134 and 122 other

terms to the sites added to the EGF and TGF data respectively, whilst RWHN assigned a refined

list of 36 and 39 respectively (Figure 5B). This result suggests that RWHN assigns more specific

functions to each phosphorylated site. Indeed, for each of the investigated clusters, RWHN was

able to assign sites belonging to those clusters to functions that closely resemble those that have

been experimentally proven by Francavilla *et al*. or described in PSP. The ORA results, whilst not

inaccurate, assigned more generic biological functions and did not assign specific terms. These

results suggest the RWHN algorithm can distinguish between sites of known function and can

assign functional terms to differentially regulated phosphorylation sites that could be further in-

vestigated experimentally.

Figure 5. Cluster containing sites annotated to "Cell growth, inhibited" in Phosphosite Plus, spiked into Francavilla et al. networks. A) Terms in common between ORA and RWHN results with both the EGF and TGF-α networks. B) Total number of terms in the output of both types of analyses on both networks.

Data	Analysis	No. terms
EGF	ORA	134
TGF	ORA	122
EGF	RWHN	36
TGF	RWHN	29

Experimental Case study 2: Phosphorylation-dependent response to Lapatinib treatment and resistance in breast cancer

To verify the applicability of the algorithm with non-temporal shotgun phosphoproteomics data,

we applied it to the experimental dataset from Ruprecht *et al*. [22] describing changes in the phos-

phoproteome upon treatment with the breast cancer drug lapatinib in sensitive ('parental' BT-474)

or lapatinib-resistant (BT-474-J4) cells. The paper uncovered and experimentally validated the

role of several metabolic enzymes and signaling pathways that were driving lapatinib resistance

in breast cancer. In particular, proteins that form the spliceosome, those involved in glycolysis

and glycogen catabolism and PI3K/AKT/mTOR pathway members were differentially phosphor-

ylated in lapatinib-resistant cells compared to parental lapatinib-treated cells. Further details on

the experimental design and results from the publication can be found in Supplementary Table

S2.

We first constructed a multilayer heterogeneous network, as described in Experimental Proce-

dures, which included 1603 phosphorylated sites that showed significant changes in response to

lapatinib treatment in either the parental or resistant cell lines (Table 4). Data were grouped into

5 clusters (referred to as clusters 1 – 5) using the k-means method. The number of clusters to use

was determined using the elbow plot method [29](Figure S4A), confirming that 5 clusters were suf-

ficient to capture all features found in the data. Indeed, sites sharing similar regulation profiles

between the resistant and parental cells were clustered together (Figure S4B). As the original pub-

lication used KEGG pathways rather than GO terms, we incorporated these in the functional-

annotation layer of our network to investigate the flexibility of our approach.

Table 4. Multilayer heterogeneous network, constructed from Ruprecht *et al*. [22] lapatinib-regu-

lated phosphorylated sites in parental and lapatinib-resistant cell lines. KEGG pathways were in-

cluded as functional annotations.

Subnetwork	Edges	Nodes
Site-site	75688	1599
Protein-protein	12752	3129
Function-function	15	152
Site-protein	1599	1599 sites, 930 proteins
Protein-function	692	103 proteins, 152 pathways

The results of RWHN applied to the multilayer heterogeneous network (Figure 6A) show that

when the seed nodes are set to sites belonging to cluster 4, representing sites that are less phos-

phorylated in lapatinib-resistant cells, the spliceosome pathway was highly ranked. This cluster

interestingly contained signalling sites such as ERBB2_Y1233, ERBB2_T1227, ERBB2_Y1233,

ERBB3_S627 and MAPK1_Y187, alongside sites on known spliceosome factors such as SRRM2

(S454, S2449), SRSF6 (S301) CWC25 (S170). Figure 6B shows that there was no enrichment for

any pathways in this cluster using standard ORA. However, ORA did show enrichment for meta-

bolic terms in several clusters, unlike RWHN, despite this being one of the clear conclusions of

the original paper. Metabolic pathways were enriched in clusters 3 ("glycolysis / gluconeogene-

sis") and 5 ("central carbon metabolism in cancer") using ORA. Since much of the investigation

and experimental validation by Ruprecht *et al*. [22] was done on the sites found to be regulated only

in lapatinib-treated resistant cells, we investigated whether our approach could uncover more nu-

anced roles of sites within these datasets. We constructed a second network using only those sites

that were regulated in lapatinib-treated resistant cells (Table 5). These were grouped into four

clusters (referred to as clusters 1 – 4) by K-means clustering as before (Figure S4A,C) and KEGG

pathways were used again in the functional annotation layer.

Table 5. Multilayer heterogeneous network, constructed from Ruprecht *et al.* [22] lapatinib-regu-

lated phosphorylated sites in lapatinib-resistant cell line. KEGG pathways were included as func-

tional annotations.

Subnetwork	Edges	Nodes
Site-site	668491	1910
Protein-protein	14770	3880
Function-function	15	187
Site-protein	1910	1910 sites, 1087 proteins
Protein-function	874	96 proteins, 187 pathways

As described in Figure 6C, two sets of seed nodes (clusters 2 and 3, representing sites substan-

tially upregulated and downregulated, Figure S4C) tended to rank metabolic pathways more

highly ("starch and sucrose metabolism", "galactose metabolism", "pentose phosphate pathway",

"purine metabolism", "amino sugar and nucleotide sugar metabolism"). Cluster 3, along with

cluster 1 (representing moderately downregulated sites), ranked "spliceosome" highly. These ini-

tial interpretations indicate that using a specific subset of the regulated sites in the data yielded

more specific and functionally relevant terms being ranked highly by the RWHN algorithm. More-

over, only cluster 2 showed enrichment for any metabolic pathways using ORA (Figure 6D). As

this is a large cluster containing all the sites shown to be significantly upregulated in the resistant

cell line, there is a considerable amount of 'noise' in the ORA results. The enrichment of two

metabolic pathways ("glycolysis / gluconeogenesis" and "central carbon metabolism in cancer")

potentially corroborate the role of these sites as metabolism regulators, as indicated by RWHN.

This cluster contained several of the sites classified as metabolic drivers of the resistance

phenotype, such as ENO1_Y44, PKM_Y175, PDHA_S293/S300. Intriguingly, S17 on the diverse

signalling regulator SRC is also found in this cluster; this site is annotated to induction of enzy-

matic activity in PSP.

Figure 6. Site-centric and gene-centric analysis of Ruprecht *et al.* [22] datasets. (A) RWHN results

related to sites regulated by lapatinib in lapatinib-resistant and parental cell lines. (B) ORA results

related to proteins modified by lapatinib in lapatinib-resistant and parental cell lines. (C) RWHN

results related to sites regulated by lapatinib in lapatinib-resistant cell lines only. (D) ORA results

related to proteins modified by lapatinib in lapatinib-resistant cell lines only.

A closer look at cluster 3 uncovered sites on spliceosome factors (RBMX_S58, CDK13_S525,

CWC25_S17) and known signaling proteins (ERBB2_Y1233, FGFR4_S505). A site on the adap-

tor protein IRS2 (S1176) was also found in this cluster; as a known player in glucose and lipid

metabolism [49]. This points towards the link between rewired signalling and metabolism investi-

gated by Ruprecht *et al*. [22]. It is also annotated to "metastatic potential" in PSP, indicating its

therapeutic potential. ORA also showed enrichment for "spliceosome" in cluster 1. Indeed, this

cluster contained several of the sites highlighted in the paper. For example, the known lapatinib

responsive sites SF3B2_S309 and SF3B1_T261 could be found in this cluster, alongside 17 of the

30 SRRM2 sites that were significantly regulated by lapatinib.

From the comparison of ORA and the RWHN method we have extracted meaningful associa-

tions between specific phosphorylated sites and their functions in driving lapatinib resistance. Our

algorithm also highlighted potential further sites, such as IRS2_S1176, compared to the ORA

method, that may serve as hubs or intersections of crosstalk between key signalling pathways and

the rewired metabolic processes. It can be noted that the RWHN output was substantially more

refined than the ORA output, with the conclusions of the original paper being captured in 14 terms

in our analysis of the sites regulated in resistance, as opposed to 51 using ORA.

Experimental Case Study 3: Spatially resolved activity of HRAS.
The final dataset we analyzed described immunoprecipitation (IP) samples, in order for us to

see if our algorithm would be useful for analyzing local as well as global phosphoproteomes. The

subcellular localization of signaling proteins has been established as crucial in regulating cascades

and cellular processes. There are now numerous methods that allow experimental biologists to

detect spatial effects of proteins and phosphorylation, for example through proximity labeling and

targeted proteomics[50,51]. Recent work by Santra *et al*.[23] took a similar approach to investigate the

differential roles of a mutant, constitutively active form of HRAS, HRASV12, at different subcel-

lular locations. HRASV12 constructs tagged with different signal peptides (targeting them to dif-

ferent subcellular locations) were stably transfected into HeLa cells; this allowed the authors to

collect multi-omic samples with endogenous HRAS, unlocalized HRASV12 and HRASV12 lo-

calized to either the plasma membrane in disordered membrane regions (DM) or lipid rafts (LR),

Golgi apparatus (GA) and endoplasmic reticulum (ER). Using this approach, it was confirmed

that the majority of HRASV12 activity is mediated from the plasma membrane, with ER- and GA-

localized HRASV12 involved in regulating some of HRASV12's role in cell survival. Further

details on the experimental design and results from the publication can be found in Supplementary

Table S2.

The phosphoproteomics data from this study were clustered using k-means clustering, with an

optimal cluster number of 6 determined using an elbow plot (Figure S5A,B). Table 6 describes

the properties of the multilayer network constructed with these clusters. The first three clusters

represented dynamics discussed in the original paper: cluster 1 represented phosphorylated events

of mutant HRAS regardless of localization (location-independent); cluster 4 represented HRAS

in the secretory pathway, at the ER and GA (ER/GA); cluster 6 represented HRAS at the plasma

membrane, in the DM or LR (DM/LR). As these experimental conditions were best characterized

in the original paper, downstream analysis focused on these three clusters.

Table 6. Multilayer heterogeneous network, constructed using regulated sites from the Santra *et*

al. [23] dataset. GOBP terms were included as functional annotations.

Subnetwork	Edges	Nodes
Site-site	766	621
Protein-protein	7262	2163
Function-function	370	452
Site-protein	621	621 sites, 474 proteins
Protein-function	900	122 proteins, 442 functions

Figure 7. Site-centric and gene-centric analysis of Santra *et al*. [23] datasets; results from RWHN

and ORA are in green and purple, respectively. Clusters shown in the figure refer to phosphory-

lated sites that are location-independent or found at the endoplasmic reticulum (ER)/Golgi

apparatus (GA) or disordered membrane (DM)/ lipid rafts (LR) (marked in the figure as clusters

1, 2, and 3, respectively).

Using RWHN (Figure 7, in purple), we found that the ER/GA and DM/LR clusters both highly

ranked key signaling pathways such as "ERK1 and ERK2 cascade" and "TOR signaling", along-

side signaling regulation such as "Receptor mediated endocytosis". This is consistent with the

finding from the original paper, that the majority of HRAS mediated effects derive from the lo-

calized population. Specifically, the DM/LR cluster ranks mitogenic processes highly, such as

"regulation of epithelial cell proliferation" and "positive regulation of metaphase/anaphase tran-

sition of cell cycle". The original paper also concluded that the majority of HRAS' mitogenic

effects were mediated from the plasma membrane. The ER/GA cluster independently ranks "pos-

itive regulation of growth" and "Notch signaling pathway". This potentially implicates ER/GA

localized HRAS, in the known developmental roles of HRAS in development; this was touched

upon but not investigated in the original paper.

Conversely, where a term is ranked highly in all three clusters, this could indicate that this is

part of the localization-independent regulation of HRAS. Twenty of the highly regulated terms fit

this definition; of these, several are related to immunological processes, such as T-help cell lineage

commitment" and "cellular response to interleukin-25". The location-independent cluster also in-

dependently ranks "interleukin-27-mediated signaling pathway" and "STAT cascade". This sug-

gests that HRAS' role in regulating cytokine and immune response is not location dependent.

An obvious outcome of the results using RWHN and ORA on this data is the difference in the

number terms associated to each cluster. There were 132 terms describing these clusters with ORA

(Figure 7, in green), compared to 86 with RWHN, with many of the enriched terms related to the

overrepresented categories depicted in Figure 3, such as "gene expression". Several comparisons

can be drawn between the results of the two analyses to add credence to the accuracy of results

from RWHN. For example, cluster 1 shows enrichment for the term "Interleukin-6-mediated sig-

naling pathway", and cluster 3 is enriched for the term "regulation of mitotic cell cycle". These

results from ORA are consistent with the results from RWHN and the original paper, but with a

significant amount of noise and redundancy in the ORA output relative to RWHN.

5. Discussion

Phosphorylation has an important impact on protein function and thus cellular behavior. Within

a network of kinase-substrate interactions, phosphorylation modulates the flow of information and

regulation of disparate processes throughout the cell. Understanding the impact of phosphoryla-

tion on protein function traditionally required the experimental investigation and manipulation of

individual sites, however with the advent of high throughput phosphoproteomics thousands of

novel phosphorylation sites of unknown function have been discovered. As there is limited ex-

perimentally validated information available on how regulation at individual phosphorylated sites

impacts the downstream cellular output, current methods for analyzing high throughput phospho-

proteomics data rely on general descriptors of phosphorylated protein function such as GO terms

or involvement in known pathways. Their use in proteomics and transcriptomics analysis, though

widespread, has come under recent criticism. Maertens *et al* [52] demonstrated that annotations are

lacking for a substantial number of genes known to be important in cancer, whilst Haynes *et al*. [53]

demonstrate a "rich-getting-richer" effect, with well-studied genes having higher annotation rates

despite having less -omic or molecular evidence for the association with processes or diseases.

These problems are exacerbated when used in analyzing phosphoproteomics data, as enrichment

analyses also disregard site-specific or multiple-site regulation, masking the roles of proteins in a

particular modification state [10,12].

Here, we developed and tested a method that associates phosphorylated sites to potential func-

tion using RWHN. By incorporating the pattern of phosphorylation upon perturbation, we con-

sider more of the information available in phosphoproteomics datasets and establish the phosphor-

ylated sites as the key drivers in the functional analysis. The algorithm is capable of recapturing

experimentally proven functions of phosphorylated sites in a non-gene centric manner, refining

analysis and extracting more information from phosphoproteomics data.

To prove the utility of our algorithm we applied it to a validation network, which represented a

simple model of phosphorylation dynamics in the MAPK/ERK pathway, and to two previously

published phosphoproteomics dataset describing temporal cellular signalling events and the im-

pact of resistance to drug treatment in breast cancer, respectively. Our algorithm successfully dis-

tinguished between differentially regulated phosphorylated sites from the same protein and asso-

ciated them to both known and previously uninvestigated functions. For example, when applying

this approach to the data from Francavilla *et al*. [21], we can recapture the roles of EGFR sites with

functions described in PSP. Both EGFR_Y1110 and EGFR_Y1197 are annotated to receptor in-

ternalization and cell motility in PSP; in our analysis, we found that these sites are both found in

clusters that associate with "regulation of clathrin dependent endocytosis", however only the TGF-

α cluster also ranked "regulation of cell migration" highly, reproducing the experimental evidence

from Francavilla *et al*. Whilst the authors of the original manuscript associated these sites with

their functions using a combination of gene-centric approaches and experimental validation, here

we use a single site-centric algorithm to extract these associations directly from the phosphopro-

teomics dataset, demonstrating the power of our approach in quickly and more specifically nar-

rowing down candidates for further functional studies. By associating ligand-dependent regulation

of phosphorylated sites, it is possible to disentangle their multifaceted role in regulating cellular

signalling networks and downstream cellular behaviors.

From the Ruprecht *et al*. data, we can use the example of SRRM2 to illustrate the validity of

our approach. This component of the spliceosome is known to be highly phosphorylated, with 675

phosphorylation sites recorded in PSP. At the time of writing none of these sites was ascribed to

a function in the database, despite the importance of phosphorylation and dephosphorylation

events in orchestrating splicing events [54]. It is also regularly mutated (>5% of cases in TGCA) in

lung, stomach, bladder, endometrial and colorectal cancers [6]. Ruprecht *et al*. uncovered a new role

for the spliceosome in modulating lapatinib-resistance, highlighting several sites on spliceosome

proteins (including SRRM2_S1132, S1987 and S970) that were differentially modified in resistant

and parental cell populations. Previous work has described how SRRM2-depletion in HER2-pos-

itive breast and ovarian cancer cells reduced the rate of migration [55]; the spliceosome plays myriad

roles in the breast cancer environment, but site-specific analysis is lacking. Using our algorithm,

we can begin to uncover the nuanced modification-specific roles of proteins such as SRRM2 in

metabolic rewiring. We suggest that eleven of the SRRM2 sites found in cluster 2 of the lapatinib-

resistant sites (Figure 6C) could be at the intersection between the spliceosome and metabolic

processes; the conjunction between these processes was central to the work of Ruprecht *et al*. The

sites in this cluster could be investigated as points of crosstalk between these different cellular

processes. This example highlights the value of our approach in generating hypotheses of modi-

fied protein function and signaling pathway crosstalk.

Although the associations predicted here between sites and functions cannot be interpreted as

causative, phosphorylation sites can act as indirect, and context-dependent, regulators in cellular

processes. This is demonstrated by the clustering of SRC_S17, a protein that is typically associated

with signalling downstream of cell-surface receptors, with a cluster that ranked metabolic path-

ways higher than cellular processing or signalling pathways. Critical to this is a robust method of

defining clusters, to capture meaningful regulatory patterns at different phosphorylated sites. Here

we define clusters using established techniques (e.g., elbow method for K-means) or visually iden-

tifying those with differing time-resolved behaviors. By using clusters to define edges between

phosphorylated sites in the multilayer heterogeneous network, the pattern of experimentally per-

turbed modules in the cellular system can be investigated. Although having fixed edges between

sites, as with the PPI layer of the multilayer network, would likely allow for a closer representation

of the whole cellular system, this would not be computationally viable nor in the interest of ex-

perimental biologists. This algorithm is targeted towards datasets that describe the effect of spe-

cific treatments or perturbations, allowing for a focused view on the impacted processes to identify

candidates to investigate further. Quantitatively defining the edges between phosphorylated sites

based on the dataset collected also prevents the issues plaguing standard enrichments analyses as

discussed before; fixed edges, defined on information in databases or previously published data,

would almost certainly skew the analysis towards well studied phosphorylated sites and biological

processes.

Multilayer heterogeneous networks are increasingly being used to integrate 'omic data; here,

their use allows phosphoproteomics data to be incorporated with PPI networks and functional

annotations, overcoming the issue of considering phosphoproteomics data primarily on the protein

level. A potential drawback of our approach is the reliance on large semi-curated resources such

GO or STRING. For instance, different clusters may have many terms or pathways in common

given the involvement of many proteins in the same biological functions and the high proportion

of frequently used GO terms [56]. We theorize that this may be rectified if the data were clustered

into more groups, in order to capture more nuanced phosphorylation patterns. However, enrich-

ment of non-specific terms or false positives remains an issue when analyzing high-throughput

'omics data by other commonly used methods too. By only including edges in the protein layer of

the network with a STRING experimental confidence score of greater than 0.4 and only including

non-redundant terms in the function layer we have reduced the noise in the resulting list of asso-

ciated terms. This could be taken further by filtering for only high confidence functional annota-

tions or incorporating annotations from multiple sources into the network. Moreover, RWHN

does not result in a different set of terms in the output compared to the accepted standard method

ORA, but rather provides a refinement that will allow for biological insights to be reached more

clearly.

Our method is flexible enough to be used with any discovery phosphoproteomics data that de-

scribes a change between conditions. This is an improvement on the previously published use of

RWHN using multiple sources of phosphoproteomics data to uncover disease-dependent regula-

tion [19]. Moreover, it could easily be generalized to any post-translational modification proteomics

dataset, as it incorporates readily available PPI and functional annotation data, as demonstrated

here. The fundamental aspect would be maintained with any of these expansions: specific patterns

of regulation at modified sites dictate movement through the multilayer heterogeneous network.

6. Conclusion

We have proposed a site-centric approach to analyze phosphoproteomics data, that provides a

robust alternative to gene-centric methods of analysis. We integrated clustered quantitative phos-

phoproteomics data, a context-specific PPI network and functional annotations into a multilayer,

heterogeneous network and used the RWHN method to predict the functions of phosphorylation

sites with similar regulatory patterns. Using our algorithm, we extracted experimentally validated

associations between phosphorylated sites and their role in cellular processes, which could not be

captured using the typical gene-centric methods. Moreover, our algorithm has the potential to be

used by researchers in predicting novel site-function associations and generating hypotheses to be

experimentally validated.

746   **7. Supporting Information**
747

Figure S1. Optimization of RWHN parameters using the validation dataset.

Figure S2. Validation using randomly permuted networks.

Figure S3. Clustering of Francavilla et al. data.

Figure S4. Clustering of Ruprecht et al. data.

Figure S5. Clustering of Santra et al. data.

Table S1. PSP ON_PROCESS annotations, that describe biological function of sites, mapped to

most closely related GOBP term.

Table S2. Summary of the experimental methods and results from each of the Case Study da-

tasets.

Data and source code for the analysis described in the manuscript can be found at

github.com/JoWatson2011/RWHN_Phosphoproteomics

An R package for use of the tool can be found at github.com/JoWatson2011/phosphoRWHN

8. Author Contributions

760
761 The manuscript was written through contributions of all authors. All authors have given ap-

762 proval to the final version of the manuscript.

9. Funding Sources

764
765 JW is supported by a Biotechnology and Biological Sciences Research Council (BBSRC) Doc-

766 toral Training Programme grant (BB/M011208/1). CF is supported by a Wellcome Trust Sir Henry

767 Dale Fellowship (107636/Z/15/Z), the Biotechnology and Biological Sciences Research Council

768 (BB/R015864/1), and the Medical Research Council (MR/T016043/1).

10. Acknowledgements

770 We would like to acknowledge the assistance given by Research IT and the use of the Compu-

771 tational Shared Facility at The University of Manchester. We would like to thank current and

772 former members of the Schwartz and Francavilla labs for their thoughtful and insightful discussion

773 on this work and Dr Joe Swift (University of Manchester) for useful comments on our manuscript.

11. Abbreviations

774
775 RWHN, Random walk on heterogeneous network; PTM, Post-translational modification; PPI,

776 Protein-protein interaction; GO, Gene Ontology; PSP, Phosphosite Plus; EGF, Epidermal Growth

777 Factor; TGF-α, Tumor growth factor α; RWR, Random walk with restart; GOBP, Gene Ontology

778 Biological Processes; FDR, False discovery rate; SILAC, Stable Isotope Labelling of Amino Ac-

779 ids in Cell culture; KDE, Kernel Density Estimation

12. References

(1) Sugiyama, N.; Imamura, H.; Ishihama, Y. Large-Scale Discovery of Substrates of the Human Kinome. *Scientific Reports* **2019**, *9* (1), 10503. https://doi.org/10.1038/s41598-019-46385-4.

(2) Day, E. K.; Sosale, N. G.; Lazzara, M. J. Cell Signaling Regulation by Protein Phosphorylation: A Multivariate, Heterogeneous, and Context-Dependent Process. *Current Opinion in Biotechnology* **2016**, *40*, 185–192. https://doi.org/10.1016/j.copbio.2016.06.005.

(3) Vyse, S.; Desmond, H.; Huang, P. H. Advances in Mass Spectrometry Based Strategies to Study Receptor Tyrosine Kinases. *IUCrJ* **2017**, *4* (Pt 2), 119–130. https://doi.org/10.1107/S2052252516020546.

(4) Munk, S.; Refsgaard, J. C.; Olsen, J. V. Systems Analysis for Interpretation of Phosphoproteomics Data. In *Phospho-Proteomics: Methods and Protocols*; von Stechow, L., Ed.; Springer New York: New York, NY, 2016; pp 341–360. https://doi.org/10.1007/978-1-4939-3049-4_23.

(5) Sacco, F.; Perfetto, L.; Cesareni, G. Combining Phosphoproteomics Datasets and Literature Information to Reveal the Functional Connections in a Cell Phosphorylation Network. *PROTEOMICS* **2018**, *18* (5–6), 1700311. https://doi.org/10.1002/pmic.201700311.

(6) Hornbeck, P. V.; Zhang, B.; Murray, B.; Kornhauser, J. M.; Latham, V.; Skrzypek, E. PhosphoSitePlus, 2014: Mutations, PTMs and Recalibrations. *Nucleic Acids Research* **2015**, *43* (D1), D512–D520. https://doi.org/10.1093/nar/gku1267.

(7) Linding, R.; Jensen, L. J.; Pasculescu, A.; Olhovsky, M.; Colwill, K.; Bork, P.; Yaffe, M. B.; Pawson, T. NetworKIN: A Resource for Exploring Cellular Phosphorylation Networks. *Nucleic Acids Research* **2008**. https://doi.org/10.1093/nar/gkm902.

(8) Casado, P.; Rodriguez-Prados, J.-C.; Cosulich, S. C.; Guichard, S.; Vanhaesebroeck, B.; Joel, S.; Cutillas, P. R. Kinase-Substrate Enrichment Analysis Provides Insights into the Heterogeneity of Signaling Pathway Activation in Leukemia Cells. *Science signaling* **2013**, *6* (268), rs6. https://doi.org/10.1126/scisignal.2003573.

(9) Mischnik, M.; Sacco, F.; Cox, J.; Schneider, H.-C.; Schäfer, M.; Hendlich, M.; Crowther, D.; Mann, M.; Klabunde, T. IKAP: A Heuristic Framework for Inference of Kinase Activities from Phosphoproteomics Data. *Bioinformatics* **2016**, *32* (3), 424–431. https://doi.org/10.1093/bioinformatics/btv699.

(10) Needham, E. J.; Parker, B. L.; Burykin, T.; James, D. E.; Humphrey, S. J. Illuminating the Dark Phosphoproteome. *Science Signaling* **2019**, *12* (565), eaau8645. https://doi.org/10.1126/scisignal.aau8645.

(11) Rudolph, J. D.; de Graauw, M.; van de Water, B.; Geiger, T.; Sharan, R. Elucidation of Signaling Pathways from Large-Scale Phosphoproteomic Data Using Protein Interaction Networks. *Cell systems* **2016**, *3* (6), 585-593.e3. https://doi.org/10.1016/j.cels.2016.11.005.

(12) Krug, K.; Mertins, P.; Zhang, B.; Hornbeck, P.; Raju, R.; Ahmad, R.; Szucs, M.; Mundt, F.; Forestier, D.; Jane-Valbuena, J.; Keshishian, H.; Gillette, M. A.; Tamayo, P.; Mesirov, J. P.; Jaffe, J. D.; Carr, S. A.; Mani, D. R. A Curated Resource for Phosphosite-Specific Signature Analysis. *Molecular & cellular proteomics : MCP* **2019**, *18* (3), 576–593. https://doi.org/10.1074/mcp.TIR118.000943.

(13) Boyle, E. I.; Weng, S.; Gollub, J.; Jin, H.; Botstein, D.; Cherry, J. M.; Sherlock, G.
GO::TermFinder—Open Source Software for Accessing Gene Ontology Information and
Finding Significantly Enriched Gene Ontology Terms Associated with a List of Genes. *Bi-*
oinformatics **2004**, *20* (18), 3710–3715. https://doi.org/10.1093/bioinformatics/bth456.

(14) Battiston, F.; Nicosia, V.; Chavez, M.; Latora, V. Multilayer Motif Analysis of Brain Net-
works. *Chaos: An Interdisciplinary Journal of Nonlinear Science* **2017**, *27* (4), 047404.
https://doi.org/10.1063/1.4979282.

(15) Li, Y.; Patra, J. C. Genome-Wide Inferring Gene–Phenotype Relationship by Walking on
the Heterogeneous Network. *Bioinformatics* **2010**, *26* (9), 1219–1224.
https://doi.org/10.1093/bioinformatics/btq108.

(16) Jiang, R. Walking on Multiple Disease-Gene Networks to Prioritize Candidate Genes.
JOURNAL OF MOLECULAR CELL BIOLOGY **2015**, *7* (3, SI), 214–230.
https://doi.org/10.1093/jmcb/mjv008.

(17) Soul, J.; Hardingham, T. E.; Boot-Handford, R. P.; Schwartz, J.-M. PhenomeExpress: A
Refined Network Analysis of Expression Datasets by Inclusion of Known Disease Pheno-
types. *Scientific Reports* **2015**, *5* (1), 8117. https://doi.org/10.1038/srep08117.

(18) Tang, Y.; Chen, K.; Wu, X.; Wei, Z.; Zhang, S.-Y.; Song, B.; Zhang, S.-W.; Huang, Y.;
Meng, J. DRUM: Inference of Disease-Associated M6A RNA Methylation Sites From a
Multi-Layer Heterogeneous Network. *Frontiers in Genetics* **2019**, *10*, 266.
https://doi.org/10.3389/fgene.2019.00266.

(19) Xu, X.; Wang, M. Inferring Disease Associated Phosphorylation Sites via Random Walk
on Multi-Layer Heterogeneous Network. *IEEE/ACM Transactions on Computational Biol-*
ogy and Bioinformatics **2016**, *13* (5), 836–844.
https://doi.org/10.1109/TCBB.2015.2498548.

(20) Behar, M.; Hoffmann, A. Understanding the Temporal Codes of Intra-Cellular Signals.
Current Opinion in Genetics & Development **2010**, *20* (6), 684–693.
https://doi.org/10.1016/J.GDE.2010.09.007.

(21) Francavilla, C.; Papetti, M.; Rigbolt, K. T. G.; Pedersen, A.-K.; Sigurdsson, J. O.; Cazza-
mali, G.; Karemore, G.; Blagoev, B.; Olsen, J. V. Multilayered Proteomics Reveals Molec-
ular Switches Dictating Ligand-Dependent EGFR Trafficking. *Nature Structural & Molec-*
ular Biology **2016**, *23* (6), 608–618. https://doi.org/10.1038/nsmb.3218.

(22) Ruprecht, B.; Zaal, E. A.; Zecha, J.; Wu, W.; Berkers, C. R.; Kuster, B.; Lemeer, S. Lapa-
tinib Resistance in Breast Cancer Cells Is Accompanied by Phosphorylation-Mediated Re-
programming of Glycolysis. *Cancer Research* **2017**, *77* (8), 1842–1853.
https://doi.org/10.1158/0008-5472.CAN-16-2976.

(23) Santra, T.; Herrero, A.; Rodriguez, J.; von Kriegsheim, A.; Iglesias-Martinez, L. F.;
Schwarzl, T.; Higgins, D.; Aye, T.-T.; Heck, A. J. R.; Calvo, F.; Agudo-Ibáñez, L.; Crespo,
P.; Matallanas, D.; Kolch, W. An Integrated Global Analysis of Compartmentalized HRAS
Signaling. *Cell reports* **2019**, *26* (11), 3100-3115.e7.
https://doi.org/10.1016/j.celrep.2019.02.038.

(24) Cox, J.; Mann, M. MaxQuant Enables High Peptide Identification Rates, Individualized
p.p.b.-Range Mass Accuracies and Proteome-Wide Protein Quantification. *Nature Biotech-*
nology **2008**, *26* (12), 1367–1372. https://doi.org/10.1038/nbt.1511.

(25) Tyanova, S.; Temu, T.; Sinitcyn, P.; Carlson, A.; Hein, M. Y.; Geiger, T.; Mann, M.; Cox,
J. The Perseus Computational Platform for Comprehensive Analysis of (Prote)Omics Data.
Nature Methods **2016**, *13* (9), 731–740. https://doi.org/10.1038/nmeth.3901.

(26) Ritchie, M. E.; Phipson, B.; Wu, D.; Hu, Y.; Law, C. W.; Shi, W.; Smyth, G. K. Limma
Powers Differential Expression Analyses for RNA-Sequencing and Microarray Studies.
Nucleic Acids Research **2015**, *43* (7), e47–e47. https://doi.org/10.1093/nar/gkv007.

(27) Schwämmle, V.; Jensen, O. N. A Simple and Fast Method to Determine the Parameters for
Fuzzy c–Means Cluster Analysis. *Bioinformatics* **2010**, *26* (22), 2841–2848.
https://doi.org/10.1093/bioinformatics/btq534.

(28) Campello, R. J. G. B.; Hruschka, E. R. A Fuzzy Extension of the Silhouette Width Criterion
for Cluster Analysis. *Fuzzy Sets and Systems* **2006**, *157* (21), 2858–2875.
https://doi.org/10.1016/J.FSS.2006.07.006.

(29) Zambelli, A. E. A Data-Driven Approach to Estimating the Number of Clusters in Hierar-
chical Clustering. *F1000Research* **2016**, *5*, 2809. https://doi.org/10.12688/f1000re-
search.10103.1.

(30) Szklarczyk, D.; Gable, A. L.; Lyon, D.; Junge, A.; Wyder, S.; Huerta-Cepas, J.; Simonovic,
M.; Doncheva, N. T.; Morris, J. H.; Bork, P.; Jensen, L. J.; Mering, C. von. STRING V11:
Protein–Protein Association Networks with Increased Coverage, Supporting Functional
Discovery in Genome-Wide Experimental Datasets. *Nucleic Acids Research* **2018**, *47* (D1),
D607–D613. https://doi.org/10.1093/nar/gky1131.

(31) Yu, G.; Li, F.; Qin, Y.; Bo, X.; Wu, Y.; Wang, S. GOSemSim: An R Package for Measuring
Semantic Similarity among GO Terms and Gene Products. *Bioinformatics* **2010**, *26* (7),
976–978. https://doi.org/10.1093/bioinformatics/btq064.

(32) Wang, J. Z.; Du, Z.; Payattakool, R.; Yu, P. S.; Chen, C.-F. A New Method to Measure the
Semantic Similarity of GO Terms. *Bioinformatics* **2007**, *23* (10), 1274–1281.
https://doi.org/10.1093/bioinformatics/btm087.

(33) Stoney, R. A.; Ames, R. M.; Nenadic, G.; Robertson, D. L.; Schwartz, J.-M. Disentangling
the Multigenic and Pleiotropic Nature of Molecular Function. *BMC systems biology* **2015**,
9 Suppl 6 (Suppl 6), S3. https://doi.org/10.1186/1752-0509-9-S6-S3.

(34) Blondel, V. D.; Guillaume, J.-L.; Lambiotte, R.; Lefebvre, E. Fast Unfolding of Communi-
ties in Large Networks. *J. Stat. Mech.* **2008**, *2008* (10), P10008.
https://doi.org/10.1088/1742-5468/2008/10/P10008.

(35) Kuleshov, M. V; Jones, M. R.; Rouillard, A. D.; Fernandez, N. F.; Duan, Q.; Wang, Z.;
Koplev, S.; Jenkins, S. L.; Jagodnik, K. M.; Lachmann, A.; McDermott, M. G.; Monteiro,
C. D.; Gundersen, G. W.; Ma'ayan, A. Enrichr: A Comprehensive Gene Set Enrichment
Analysis Web Server 2016 Update. *Nucleic Acids Research* **2016**, *44* (W1), W90–W97.
https://doi.org/10.1093/nar/gkw377.

(36) Carlson, M. GO.Db: A Set of Annotation Maps Describing the Entire Gene Ontology. R
Package Version 3.8.2. *Bioconductor 3.10*. 2019.

(37) Köhler, S.; Bauer, S.; Horn, D.; Robinson, P. N. Walking the Interactome for Prioritization
of Candidate Disease Genes. *The American Journal of Human Genetics* **2008**, *82* (4), 949–
958. https://doi.org/10.1016/J.AJHG.2008.02.013.

(38) Riley, N. M.; Coon, J. J. Phosphoproteomics in the Age of Rapid and Deep Proteome Pro-
filing. *Analytical Chemistry* **2016**, *88* (1), 74–94. https://doi.org/10.1021/acs.anal-
chem.5b04123.

(39) Blatti, C.; Sinha, S. Characterizing Gene Sets Using Discriminative Random Walks with
Restart on Heterogeneous Biological Networks. *Bioinformatics* **2016**, *32* (14), 2167–2175.
https://doi.org/10.1093/bioinformatics/btw151.

(40) Shieh, G. S. A Weighted Kendall's Tau Statistic. *Statistics & Probability Letters* **1998**, *39* (1), 17–24. https://doi.org/10.1016/S0167-7152(98)00006-6.

(41) Fey, D.; Croucher, D. R.; Kolch, W.; Kholodenko, B. N. Crosstalk and Signaling Switches in Mitogen-Activated Protein Kinase Cascades. *Frontiers in Physiology* **2012**, *3*. https://doi.org/10.3389/fphys.2012.00355.

(42) Arkun, Y.; Yasemi, M. Dynamics and Control of the ERK Signaling Pathway: Sensitivity, Bistability, and Oscillations. *PLOS ONE* **2018**, *13* (4), e0195513. https://doi.org/10.1371/journal.pone.0195513.

(43) H. B. Mann; D. R. Whitney. On a Test of Whether One of Two Random Variables Is Stochastically Larger than the Other. *The Annals of Mathematical Statistics* **1947**, *18* (1), 50–60. https://doi.org/10.1214/aoms/1177730491.

(44) Benjamini, Y.; Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **1995**, *57* (1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x.

(45) Dow, L. E.; Elsum, I. A.; King, C. L.; Kinross, K. M.; Richardson, H. E.; Humbert, P. O. Loss of Human Scribble Cooperates with H-Ras to Promote Cell Invasion through Deregulation of MAPK Signalling. *Oncogene* **2008**, *27* (46), 5988–6001. https://doi.org/10.1038/onc.2008.219.

(46) Nola, S.; Sebbagh, M.; Marchetto, S.; Osmani, N.; Nourry, C.; Audebert, S.; Navarro, C.; Rachel, R.; Montcouquiol, M.; Sans, N.; Etienne-Manneville, S.; Borg, J. P.; Santoni, M. J. Scrib Regulates PAK Activity during the Cell Migration Process. *Human Molecular Genetics* **2008**, *17* (22), 3552–3565. https://doi.org/10.1093/hmg/ddn248.

(47) Pennock, S.; Wang, Z. A Tale of Two Cbls: Interplay of c-Cbl and Cbl-b in Epidermal Growth Factor Receptor Downregulation. *Molecular and Cellular Biology* **2008**, *28* (9), 3020–3037. https://doi.org/10.1128/mcb.01809-07.

(48) Roepstorff, K.; Grandal, M. V.; Henriksen, L.; Knudsen, S. L. J.; Lerdrup, M.; Grøvdal, L.; Willumsen, B. M.; Van Deurs, B. Differential Effects of EGFR Ligands on Endocytic Sorting of the Receptor. *Traffic* **2009**, *10* (8), 1115–1127. https://doi.org/10.1111/j.1600-0854.2009.00943.x.

(49) Neukamm, S. S.; Toth, R.; Morrice, N.; Campbell, D. G.; MacKintosh, C.; Lehmann, R.; Haering, H. U.; Schleicher, E. D.; Weigert, C. Identification of the Amino Acids 300-600 of IRS-2 as 14-3-3 Binding Region with the Importance of IGF-1/Insulin-Regulated Phosphorylation of Ser-573. *PLoS ONE* **2012**, *7* (8). https://doi.org/10.1371/journal.pone.0043296.

(50) Gingras, A.-C.; Abe, K. T.; Raught, B. Getting to Know the Neighborhood: Using Proximity-Dependent Biotinylation to Characterize Protein Complexes and Map Organelles. *Current Opinion in Chemical Biology* **2019**, *48*, 44–54. https://doi.org/10.1016/j.cbpa.2018.10.017.

(51) Orre, L. M.; Vesterlund, M.; Pan, Y.; Arslan, T.; Zhu, Y.; Fernandez Woodbridge, A.; Frings, O.; Fredlund, E.; Lehtiö, J. SubCellBarCode: Proteome-Wide Mapping of Protein Localization and Relocalization. *Molecular Cell* **2019**, *73* (1), 166-182.e7. https://doi.org/10.1016/j.molcel.2018.11.035.

(52) Maertens, A.; Tran, V. P.; Maertens, M.; Kleensang, A.; Luechtefeld, T. H.; Hartung, T.; Paller, C. J. Functionally Enigmatic Genes in Cancer: Using TCGA Data to Map the Limitations of Annotations. *Scientific reports* **2020**, *10* (1), 4106. https://doi.org/10.1038/s41598-020-60456-x.

(53) Haynes, W. A.; Tomczak, A.; Khatri, P. Gene Annotation Bias Impedes Biomedical Research. *Scientific Reports* **2018**, *8* (1), 1362. https://doi.org/10.1038/s41598-018-19333-x.

(54) Naro, C.; Sette, C. Phosphorylation-Mediated Regulation of Alternative Splicing in Cancer. *International Journal of Cell Biology*. 2013. https://doi.org/10.1155/2013/151839.

(55) Mukherji, M.; Brill, L. M.; Ficarro, S. B.; Hampton, G. M.; Schultz, P. G. A Phosphoproteomic Analysis of the ErbB2 Receptor Tyrosine Kinase Signaling Pathways. *Biochemistry* **2006**, *45* (51), 15529–15540. https://doi.org/10.1021/bi060971c.

(56) Gaudet, P.; Dessimoz, C. Gene Ontology: Pitfalls, Biases, and Remedies. In *The Gene Ontology Handbook*; Dessimoz, C., Škunca, N., Eds.; Springer New York: New York, NY, 2017; pp 189–205. https://doi.org/10.1007/978-1-4939-3743-1_14.

For table of contents only

We confirm that the eTOC figure contains original material drawn by the authors.