Facilitation and biodiversity jointly drive mutualistic networks

Gianalberto Losapio[1], Elizabeth Norton Hasday[2], Xavier Espadaler[3], Christoph

Germann[4], Francisco Javier Ortiz-Sánchez[5], Adrian Pont[6], Daniele Sommaggio[7], Christian

Schöb[9]

[1] Department of Biology, Stanford University, 94305 California, USA.

[2] Department of Evolutionary Biology and Environmental Studies, University of Zurich,

8057 Zurich, Switzerland.

[3] CREAF, Autonomous University of Barcelona, 08193 Bellaterra, Spain.

[4] Naturhistorisches Museum Basel, Augustinergasse 2, 4001 Basel, Switzerland.

[5] Grupo de Investigación "Transferencia de I+D en el Área de Recursos Naturales",

Universidad de Almería, E-04120 La Cañada Almería, Spain.

[6] Oxford University Museum of Natural History, Parks Road, Oxford OX1 3PW, UK.

[7] Department of Agricultural and Food Sciences, Alma Mater Studiorum-Università di

Bologna, 40127 Bologna, Italy.

[9] Institute of Agricultural Sciences, ETH Zurich, 8092 Zurich, Switzerland.

Correspondence author: Gianalberto Losapio, losapiog@stanford.edu

ABSTRACT

1. Facilitation by nurse plants increases understorey diversity and supports ecological

communities. In turn, biodiversity shapes ecological networks and enhances ecosystem

functioning. However, whether and how facilitation and increased biodiversity jointly

influence community structure and ecosystem functioning remains unclear.

2. We performed a field experiment disentangling the relative contribution of nurse

plants and increasing understorey plant diversity in driving pollination interactions.

Both the presence of nurse shrubs as well as increased understorey plant diversity

increased pollinator diversity and visitation rates. While nurse and understorey

diversity effects on pollinator visitation rates did not interact, the effects of increasing

understorey plant diversity on pollinator diversity were stronger in the absence than in

the presence of shrubs, meaning that nurse shrubs attenuated the effects of high

understorey diversity and buffered the effects of low understorey diversity.

3. We also found positive complementarity effects among understorey species as well as

complementarity between nurse plants and understorey species at high diversity.

Results also indicate negative selection effects, suggesting that species with generally

few pollinators benefit the most in the polyculture, while a species (possibly the nurse

plant) with generally lots of pollinators does not. The corresponding changes in

pollination networks with the experimental treatments were due to both changes in the

frequency of visits and turnover in pollinator community composition.

4. *Synthesis* Plant–plant facilitative systems, where a nurse plant increases understorey

plant diversity, are common in stressful environments. Here, we show that these

facilitative systems positively influence mutualistic interactions with pollinators via

both direct nurse effects and indirect positive effects of increasing plant diversity.

Conserving and supporting nurse plant systems is crucial not only for maintaining plant

diversity but also for supporting ecosystem functions and services.

<u>Keywords</u>

Cooperation, Ecosystem functioning and services, Leguminous shrubs, Mutualism,

Nurse plants, Plant–Plant–Insect interactions, Pollination, *Retama sphaerocarpa*

1 INTRODUCTION

Positive interactions among plants often increase community diversity (Bruno et al.,

2003; Callaway, 2007; Levin, 2009; Keddy, 2017; Ellison, 2019). In turn, community

diversity increases the functioning and stability of ecological communities (Chapin et al.,

2000; Hooper et al., 2005; Tilman et al., 2014; IPBES, 2019) and facilitation contributes

to these diversity–ecosystem function relationships (Wright et al. 2017). However,

whether and how facilitation and biodiversity interact to influence community

structure and ecosystem functioning is not known.

Nurse plants (e.g., leguminous shrubs and trees, cushion plants, kelp) can ameliorate

local environmental conditions, increase biodiversity in their understorey and in the

entire ecosystem, and organize entire communities (Callaway, 2007; Brooker et al.,

2009; Pugnaire, 2010; Butterfield et al., 2013; Cavieres et al., 2014; McIntire & Fajardo,

2014; Rodríguez-Echeverría et al., 2016; Ellison, 2019). By increasing both resources

and biodiversity, nurse plants can shape networks of ecological interactions (Losapio et

al., 2018) and ultimately influence ecosystem functioning (Losapio et al., 2021).

Facilitation has been shown for example to increase plant primary productivity (Wright

& Jones, 2004; Michalet & Touzard, 2010; Wright et al. 2017, Schöb et al. 2019) or rates

of flower visitation by pollinators (Losapio et al., 2021). In fact, numerous studies have

demonstrated the positive effects of diverse flower assemblages on pollination (Laverty

& Plowright, 1988; Laverty 1992; O'Connell & Johnston 1998; Feldman et al., 2004;

Gazhoul, 2006; Braun & Lortie 2019; Losapio & Schöb 2020). Understanding the

influence of nurse species and their positive effects for understorey plant diversity on

pollination interactions is crucial for predicting how facilitation can influence

ecosystem functioning.

Wright *and colleagues* (2017) proposed three main classes of facilitation

mechanisms that positively contribute to ecosystem functioning: (i) indirect biotic

facilitation, via reducing species-specific natural enemies (pathogens, herbivores); (ii)

abiotic facilitation via nutrient enrichment, such as the legume–rhizobia symbiosis that

directly increases nitrogen availability for neighbouring plants; (iii) abiotic amelioration

via improving microclimatic conditions. These mechanisms are deduced from literature

on biodiversity experiments carried out mainly on plant-species-richness gradients and

biomass productivity in temperate meadows (Jiang, Pu & Nemergut, 2008). Thus, the

potential of such biodiversity experiments to gather generalizable knowledge is limited

to a small and limited range of latitudes, systems, and ecological functions.

Nevertheless, there is an increasing number of studies showing that diversity has

similar effects in other types of ecosystems (Albrecht et al. 2012; Fründ et al. 2013;

Schleuning et al., 2015; Rohr et al., 2020; Losapio et al., 2021). For example, increasing

plant species richness increases pollinator diversity and supports mutualistic network

structure (Ebeling et al., 2008; Scherber et al., 2010; Blüthgen & Klein 2011).

Analogously, pollinator diversity increases plant fitness and ecosystem services

(Albrecht et al., 2012; Fründ et al., 2013; Winfree et al., 2018). As nurse plants increase

biodiversity and plant diversity increases pollinator diversity, we hypothesize that the

presence of a nurse plant and increased understorey plant diversity, i.e., the two

components of a facilitative nurse plant system (e.g., Cavieres et al., 2014;

Rodríguez-Echeverría et al., 2016), have joint positive effects on pollinator richness and

their flower visitation rates (Fig. 1). Possible mechanisms include microhabitat

amelioration by nurse species and polycultures, niche creation and complementarity,

cluster and magnet effect, resource sharing, redundancy and diversification of functions.

Here, we address this hypothesis and disentangle the relative contribution of nurse

plants and understorey plant diversity in driving pollination interactions. We asked the

following questions: (i) What are the joint effects of nurse plants and understorey plant

diversity on the pollinator community? (ii) What are the costs and benefits for the nurse

plant in terms of pollination? (iii) How does a facilitative nurse plant system shape

pollination interactions and mutualistic networks?

2 MATERIALS AND METHODS

2.1 Study system

We used a well-known nurse plant system for the study of plant–plant interactions and

particularly facilitation: *Retama sphaerocarpa* (L.) BOISS. (Fabaceae) and its understorey.

This leguminous shrub creates "islands of fertility" (Pugnaire et al., 1996; Schlesinger et

al., 1996). Mutualistic interactions with *Rhizobia* bacteria in their roots contribute to the

improvement of soil resources and amelioration of microhabitat conditions (Moro et al.,

1997; Armas, Rodríguez-Echeverría & Pugnaire, 2011). This nurse plant facilitates a

wide diversity of understorey species, supports and structures plant communities

(Rodríguez-Echeverría et al., 2013, 2016; Lozano et al., 2017; Losapio et al., 2018).

Furthermore, *Retama* produces copious yellow blooms pollinated by diverse insects,

including small-, medium-, and large-sized Hymenoptera as well as several species of

ants (Rodríguez-Riaño et al., 1999).

To experimentally disentangle the direct effect of the nurse plant presence on

pollination and its indirect effect through understorey plant diversity, we artificially

created gradients of plant diversity in the understorey of *Retama* and in the absence of

Retama. For this purpose, we selected three annual herbaceous species that commonly

grow in *Retama* shrublands at our study site: *Matricaria chamomilla* L. (Asteraceae),

Echium plantagineum L. (Boraginaceae), and *Carduus bourgeanus* Boiss. & Reut.

(Asteraceae). *M. chamomilla* has white and yellow flowers in an open capitulum, *E.*

plantagineum has purple tubular flowers along a raceme, and *C. bourgeanus* has blue

flowers in a dense capitulum. These three species therefore represent a broad set of

flower morphology and pollination niches.

The study was carried out in an oak (*Quercus ilex* L.) savannah in a Mediterranean-

type ecosystem at the Aprisco de Las Corchuelas research station in Torrejón el Rubio,

Spain (39.81337 N -6.00022 W, 350 m a.s.l., mean rainfall of 637 mm year^{-1} and mean

annual temperature of 18 °C).

2.2 Experimental design

In order to disentangle the role of direct nurse effects from the indirect effects of the

nurse plant on pollinators through increased understorey plant diversity, and to further

examine their joint effects on the mutualistic interaction between plants and

pollinators, a fully-factorial experimental design was established with two factors (Fig.

1): (i) absence and presence of the legume shrub (open vs nurse), and (ii) understorey

plant diversity with zero (0), one (1) and three (3) plant species. This design results in

five treatment combinations of open–1, open–3, nurse–0, nurse–1 and nurse–3sp. The

treatment combination open–0 was excluded due to the inherent lack of flowers for

visitation.

A randomised block design was adopted by grouping together the five treatments

and replicating them three times in each block over four blocks, for a total of $n = 60$

plots. Distance between plots within the same block was approx. 1 m. Blocks were

distributed randomly over an area of about 4,800 m². Understorey plant species

richness was manipulated with plants of the three species transplanted into pots and

then the plots placed beneath *Retama* or in the open. The density of understorey plants

and flowers was kept constant over all understorey diversity levels. Plants were taken

from the neighbouring field and replaced every second day to assure fresh flowers were

available. Pots were kept aggregated or separated (*c.* 30 cm apart) below the nurse or in

the open. This additional factor (hereafter, aggregation) was replicated per block. In the

nurse alone treatment with no understorey species, three empty pots with only soil

were placed under the shrub to control for any potential effect of the pots. For the nurse

plant treatments, *Retama* shrubs were chosen of approx. the same size (height 127–178

cm and width 125–220 cm). An area of 1 m² at 1 m height was used as the pollinator

observation area in each shrub. Flowers of the surrounding vegetation within at least 1

m around each shrub and open area were cleared with a sickle.

Flower visitation was considered as a proxy for the ecosystem function of

pollination (Schleuning et al., 2015; Winfree et al., 2018; IPBES, 2019) as it correlates

well with seed production (Losapio & Schöb, 2020; but see Lázaro et al., 2013). Flower

visits were documented by sampling, identifying, and recording all insects visiting the

flowers of each plant in each plot. Observations were conducted between 9 AM and 7

PM over eight days between 1 May and 14 May 2017, covering the blooming phase of

the four plant species. Each plot was observed during three surveys of 20 min randomly

allocated over the day. Nurse plants and corresponding open plots were observed

simultaneously, thereby reducing the potential confounding effects of changing weather

conditions within blocks. Each block was sampled within two days. Pollinators were

identified at the species level whenever possible, otherwise to the genus. Specimens are

conserved in 90% alcohol at our institution collections for inspection and further

investigations.

2.3 Data analysis

To answer the first question, we calculated the abundance and richness of pollinators in

each plot, i.e., at the community level. We assessed the individual and combined effects

of facilitation, i.e., nurse presence and understorey plant diversity, on pollinator

abundance and richness (two separate models) by means of Zero-inflated Generalized

Linear Mixed Modelling (Zi-GLMM) with a negative binomial distribution (Brooks et al.,

2017). Understorey aggregation was included as additional factor. Plant species

composition and plot nested within block were considered as random effects.

To answer the second question, we calculated flower visits for each single plant

species in each plot and conducted cost–benefit analysis. For understorey plants, we

assessed the individual and combined effects of nurse shrubs and understorey species

richness on pollinator abundance by means of Zi-GLMM with a negative binomial

distribution. Understorey aggregation was included as additional factor. Plant species

identity and plot nested within block were considered as random effects. For nurse

shrubs, we assessed the effects of understorey species richness (nurse alone, 1 species

and 3 species; second-degree polynomial) on pollinator abundance by means of Zi-

GLMM with a negative binomial distribution. Understorey aggregation was included as

additional factor. Understorey composition and plot nested within block were

considered as random effects.

To answer the third question, we used a framework based on the variance

partitioning of biodiversity effects (Loreau and Hector 2001) for the pollinator

community (Losapio et al., 2021). This framework allows comparing the net impact of a

diverse plant community on flower visits, distinguishing between complementarity and

selection effects (Loreau and Hector 2001). First, we calculated complementarity effects

(CE) and selection effects (SE) among understorey species as:

$\text{CE} = 3 \overline{\frac{Y_{mc}}{M_{mc}} - 1, \frac{Y_{ep}}{M_{ep}} - 1, \frac{Y_{cb}}{M_{cb}} - 1} \; \overline{M_{mc}, M_{ep}, M_{cb}}$, and

$\text{SE} = 3 \, cov \, (\overline{\frac{Y_{mc}}{M_{mc}} - 1, \frac{Y_{ep}}{M_{ep}} - 1, \frac{Y_{cb}}{M_{cb}} - 1}, \; \overline{M_{mc}, M_{ep}, M_{cb}})$, where Y and M indicate flower

visits in polyculture (three understorey species) and monoculture (one of the three

understorey species *M. chamomilla* [mc], *E. plantagineum* [ep], *C. bourgeanus* [cb]),

respectively. These effects were calculated both in the absence and presence of nurse

shrubs. This way, we tested the impact of plant facilitation on CE and SE. Then, the

effects of nurse presence, type (CE vs SE) and their interaction on the diversity effect

size was tested using a linear model.

Second, we calculated CE and SE between nurse shrubs and understorey species.

This way, nurse and understorey were considered as two distinct functional groups.

These functional diversity effects were calculated as: $\text{CE} = 2 \overline{\frac{Y_{ns}}{M_{ns}} - 1, \frac{Y_{us}}{M_{us}} - 1} \; \overline{M_{ns}, M_{us}}$,

and $\text{SE} = 2 \, cov \, (\overline{\frac{Y_{ns}}{ns} - 1, \frac{Y_{us}}{M_{us}} - 1}, \; \overline{M_{ns}, M_{us}})$, where Y and M indicate flower visits in

polyculture (two functional diversity groups of nurse *ns* and understorey *us*) and

monoculture (nurse and understorey alone), respectively. These effects were calculated

with both one and three understorey species (nurse–1sp and open–1sp; nurse–3sp and

open–3sp). Then, the effect of understorey plant diversity (categorical), effect type (CE

vs SE), and their interaction on the diversity effect sizes was tested using a linear model.

Finally, we built mutualistic networks between the four plant species and each of

their pollinator species (or genus) according to the additive matrix framework (Losapio

et al., 2021). This approach consists of building and comparing observed networks

(hereafter, 'observed') with 'additive' networks. 'Observed' networks are built using the

plant–pollinator interactions data collected from the empirical plant community, here

composed by the nurse shrub and the three understorey species. Instead, 'additive'

networks are built using data collected from the four treatments of nurse shrub and

understorey species monocultures and by pooling plant–pollinator interactions into a

single 'additive' matrix.

To quantify network structure, we measured network eigenvector centrality

(Bonacich, 1987; Csardi & Nepusz, 2006). This metric quantifies the extent to which

plant species that are visited by many pollinators are central to the network, or poorly

visited plants are strongly connected to few central pollinators. Then, to understand the

drivers of differences in mutualistic networks, we measured the dissimilarity between

the 'observed' and 'additive' networks using the framework of beta-diversity of species

interactions (Poisot, 2016). In particular, we considered the dissimilarity in species

composition and pairwise plant–pollinator interactions. In this case, networks were

considered at the block level. Network dissimilarity was calculated within 'additive'

networks, within 'observed' networks, and between 'additive' and 'observed' networks.

Differences among networks were tested in response to the dissimilarity index (species

or interactions), dissimilarity within networks nested within dissimilarity between

networks, and their interaction using a linear model.

Statistical results are reported in terms of variances explained, using type-II ANOVA

(Fox & Weisberg, 2019), and parameter estimates with 95% Confidence Interval. In case

of significant statistical interactions, contrasts among factor combinations were

computed using estimated marginal means (Lenth, 2020).

3 RESULTS

3.1 Pollinator community

We found that nurse plants and understorey plant diversity increased the number of

pollinators (Fig. 2a) as differences in visitor abundance at the community level were

explained by both nurse presence ($P < 0.001$) and understorey plant diversity ($P =$

0.002). Instead, neither aggregation nor the interaction between nurse presence and

understorey diversity were significant. *Retama* increased flower visitor abundance on

average fivefold compared to open areas (β, 95% CI = 1.72, 0.99–2.45). Furthermore,

increasing understorey plant diversity from one to three species increased flower

visitor abundance by 82% (0.29, 0.02–0.56).

Similarly, *Retama* increased the richness of the pollinator community by 74% (1.6,

1.07–2.13, $P < 0.001$), while understorey plant diversity increased pollinator species

richness by 24% (0.34, 0.15–0.54, $P = 0.002$). Nurse presence and understorey diversity

jointly influenced pollinator species richness ($P = 0.040$, Fig. 2b). The effect of

understorey plant diversity was stronger in the open ($c = 0.69 \pm 0.20$ SE, $P = 0.005$) than

underneath the nurse canopy ($c = 0.19 \pm 0.14$ SE, $P = 0.495$), while the effect of nurse

presence was stronger at low understorey diversity ($c = 1.35 \pm 0.17$, $P < 0.001$)

compared with high understorey diversity ($c = 0.86 \pm 0.17$ SE, $P < 0.001$). Understorey

aggregation had no significant effects.

3.2 Benefits and costs

We then explored the effects (i.e., costs and benefits) of nurse plant presence and

understorey diversity on flower visitation rate for each species. For understorey

species, flowering with nurse shrubs did not have a cost nor a benefit in terms of

attracting pollinators (Fig. 3). Instead, understorey diversity had positive effects on

visitation rate for understorey plants ($P < 0.001$). These effects were independent of

aggregation, nurse presence or their interaction (Fig. 3a). In particular, increasing

understorey diversity increased the number of flower visits on each understorey

species by 34% on average (0.40, 0.14–0.54). Variance among species was low (0.013).

Both aggregation and understorey diversity influenced visitation rate to *Retama*

flowers (P = 0.045 and P = 0.002, respectively). In particular, aggregating understorey

plants increased visitor abundance for the nurse shrub by 10% (0.39, 0.01–0.76).

Understorey diversity had non-linear effects on flower visitation rate to the nurse plant

(Fig. 3b), being positive only at high understorey diversity (quadratic term 2.28, 0.98–

3.59).

3.3 Complementarity and selection effects

We then explored diversity effects (i.e., complementarity and selection effects) among

understorey species (Fig. 4a) as well as between nurse and understorey plants (Fig. 4b).

We found that diversity effects of understorey species were independent of nurse plant

presence but varied between complementarity and selection effects (P < 0.001).

Selection effects were more negative than complementarity effects were positive (Fig.

4a): complementarity effects were marginally positive in the absence and presence of

nurse shrubs (499, -33.1–1031; 517, -15.5–1049), respectively, while selection effects

were negative in both cases (-754, -1286.3– -222; -771, -1302.9– -238).

Furthermore, diversity effects of combining nurse and understorey plants changed

with understorey plant diversity (P = 0.033) depending on effect type (P = 0.009). While

complementarity effects increased with increasing understorey diversity (c = 160.0 ±

41.1 SE, P = 0.002), selection effects were the same (c = 20.1 ± 41.4 SE, P = 0.495).

Complementarity effects were negative and positive at low and high understorey

diversity (-124.6, -188.3– -60.89; 36.2, -27.5–99.93), respectively, while selection effects

were on average always negative (-38.5, -102.2–25.26; -58.6, -122.3–5.14).

3.4 Network structure

Finally, we explored centrality of the mutualistic network involving plants and

pollinators as well as the dissimilarity between additive and observed networks

(Appendix S1). We found that observed networks were less centralized than additive

networks (β = -0.35, P < 0.001, Fig. 5). Looking at network dissimilarity components, we

found that species turnover contributed twice as much as interaction rewiring (c = 0.20

$\pm$ 0.02 SE, P < 0.002, Fig. S1).

4 DISCUSSION

Facilitation and species richness play crucial roles in shaping ecological networks and

supporting ecosystem functioning, yet their joint effects remain poorly understood.

Here, we provide new evidence for the mingled consequences of the presence of a nurse

plant and plant diversity for mutualistic networks of pollination, a key ecosystem

function. The results of our field experiment confirm the hypothesis that the presence of

a leguminous nurse shrub and high understorey plant diversity jointly increased

pollinator diversity and visitation rates. Our findings reflect that complementarity and

cooperation among different plants resulted in no costs nor benefits in terms of

pollination for *Retama* but benefits for pollination of the understorey plant community

and the ecosystem overall.

4.1 Community-scale benefits

Our results support the hypothesis that nurse plant presence and understorey plant

diversity jointly influence pollination, as they both increase pollinator abundance and

richness.

Plant facilitation can positively influence ecological networks that include taxa other

than plants (Valiente-Banuet et al., 2015), such as pollinators and herbivores (Losapio

et al., 2021), ground-dwelling arthropods (van der Zee et al., 2016), mammals (Lortie et

al., 2016), and soil microorganisms (Rodríguez-Echeverría et al., 2013; Montesinos-

Navarro et al., 2019). Benefits of facilitation involved not only the plant community

itself but influenced flower visitors, whose diversity increased in the presence of

Retama and with increasing understorey richness. Nurse shrubs therefore favour

understorey plant diversity, most likely through amelioration of the biophysical

environment (Pugnaire 2010), and together with the more diverse understorey plant

community also enhanced flower visits via supporting species-rich pollinator

communities. Underlying mechanisms for a more diverse pollinator community may be

the enhanced floral display via increasing community-level attractiveness to generalist

pollinators (Losapio et al. 2021), i.e., 'cluster effect' (Krugman, 1991; Laverty, 1992),

and resource and service sharing (McIntire & Fajardo, 2014). Moreover, enhanced floral

resources via soil symbionts of the leguminous nurse shrub (Harris, 2009;

Rodríguez-Echeverría et al., 2016) may also be responsible for improving floral

attractiveness in natural conditions.

The fact that increasing understorey plant diversity increases flower visitation rate

is consistent with previous biodiversity experiments showing that plant species

richness, along with the co-varying factors including blossom cover and presence of

particularly attractive flowering species, enhances both the frequency and the temporal

stability of pollinator visits (Ebeling et al., 2008; Albrecht et al., 2012). In addition,

flower visits increase with plant diversity when diverse flower displays increase the

duration of flower provision (Fornoff et al., 2017). Since plant and flower density were

kept constant in our experiment, the positive effects of understorey diversity on

ecosystem functions are mainly driven by niche complementarity of pollinators and loss

of poorly-attractive species, as discussed in the paragraph below (4.2). We used three

plant species that have different colours, shapes, volatile cues, nectar and degree of

specialization. Such a diversified set of flowers may have contributed to attracting more

and diverse flower visitors and have compensated for conditions where the nurse shrub

was absent.

Our results show that plant–plant facilitation is an important driver of mutualistic

networks. *Retama* increased biodiversity directly and indirectly through facilitation of

other plant species, as the increase in plant diversity also increased pollinator diversity,

flower visitation rates and ecosystem functioning. The combination of nurse plant

presence and associated high understorey plant diversity produced even greater

benefits for mutualistic networks than expected by these two factors alone. In fact, the

highest levels of pollination were observed in diverse plant communities in the

understorey of *Retama.* The presence of nurse shrubs was more important at low

diversity, while increasing understorey diversity was more relevant in the absence of

Retama. Adding more understorey species (at constant plant and flower density)

produced stronger effects on pollinator diversity in the absence of nurse shrubs as

compared to their presence.

The increased flower visitation rate of communities consisting of *Retama* nurse

shrubs and a diverse understorey community compared with communities consisting

only of *Retama* or only of herbaceous understorey plants came at no costs for the

individual plant species. This shows that *Retama* was not deterring visitors to

understorey species. Furthermore, the presence of understorey plants was harmful for

Retama only at low diversity (one species), while it was neutral at high diversity (three

species). These results indicate that diversity decreased the competitive impact of

understorey plants on *Retama* visitors, shifting it to non-competitive, neutral effects. A

likely process is associated with increasing pollinator attractiveness of a more diverse

understorey community following the increase in floral display and flower trait

diversity. These results align with studies indicating that the presence of various

facilitated plant species can be positive for nurse plant pollination (Laverty, 1992;

Braun & Lortie, 2019; Losapio & Schöb, 2020). Increasing understorey diversity to

naturally-occurring levels, which are higher than three species, might even shift the

effects of understorey plants on *Retama* pollination to positive.

Expanding the understorey diversity gradient would provide further insights into

the compensation effect of understorey diversity on pollination of the nurse plant.

Beyond that, future studies might want to improve measurements of the ecosystem

service by addressing the contribution of flower visits to pollen transfer, visitor

effectiveness and seed production. Furthermore, as we controlled for plant diversity

and density by using potted plants, our experiment lacks the 'direct' facilitation effect of

the nurse plant on the understorey plants but allowed us to control for differences in

understorey density and floral display in natural communities. We refer here to

previous research demonstrating the positive direct effect of *Retama* on understorey

plants (e.g., Pugnaire et al., 1996; Rodríguez-Echeverría et al., 2013; Lozano et al., 2017).

4.2 Complementarity and selection effects

Our current understanding of the relationships between biodiversity and ecosystem

functioning comes primarily from studies focusing on the effects of plant species

richness on biomass productivity in temperate meadows (Jiang et al., 2008; Tilman et

al., 2014). While competition is often claimed to play a central role in these systems,

facilitation is overlooked or lumped within several less explicitly defined processes as

complementarity effects (Blüthgen & Klein 2011; Wright et al., 2017). The experimental

framework we adopted here (Losapio et al., 2021) allowed us manipulating both

taxonomic and functional diversity in combination with facilitation, then measuring

complementarity among understorey species as well as between nurse shrubs and

understorey plants.

Our results indicate positive complementarity effects among understorey species.

Furthermore, complementarity between nurse plants and understorey species

increased with diversity. Results also indicate negative selection effects, suggesting that

species with generally few pollinators benefit the most in the polyculture (understorey

species), while a species (possibly the nurse plant) with generally lots of pollinators

does not. These results provide new evidence for a novel facilitation process based on

community-scale facilitation (Callaway, 2007; Liancourt & Dolezal 2020) and on the

'cluster effect' (Krugman, 1991; Laverty 1992; Losapio et al., 2021): diverse flower

assemblages including nurse shrubs are more attractive than monocultures thanks to

increased visibility of the community as a whole for attracting a wider spectrum of

visitors. This way, being part of the polyculture cluster (nurse shrub with diverse

understorey) would increase the chances of being visited, and eventually pollinated.

The positive effects of increased biodiversity for pollination at the community level

also translates into increased pollination for each individual species. This is not always

the case for biodiversity experiments, where an increase in community productivity

does not necessarily follow an increase in species-specific biomass (Tilman et al., 2014).

4.3 Network change

By means of the additive matrix framework (Losapio et al., 2021), we compared

observed networks that emerge from the plant community as a whole with additive

networks that result from pooling plant species as separate components. These two

networks appear to be very different, highlighting the non-additivity of nurse and

understorey plants. This shows that interactions among plants influence interactions

between plants and pollinators. In particular, facilitative interactions among nurse and

understorey plants influence pollination networks by changing the identity and

frequency of flower visitors. Such nonadditive interactions change network structure,

making the network more de-centralised than expected by additive effects, which may

ultimately improve the overall resistance and stability of pollinator communities

(Blüthgen & Klein 2011). Furthermore, our results show that observed networks were

more similar among each other in terms of species interactions as compared to the

higher dissimilarity of species interactions observed among additive networks. Drivers

of network rewiring are associated with increasing frequency of visits and potentially

pollinator population density, thus affecting interaction strength, as well as changes in

floral attractiveness and pollination niches, which ultimately promote species turnover

and interaction rewiring.

In conclusion, our study shows that plant facilitation systems composed of nurse

plants and associated high understorey diversity have joint positive impacts on

pollination networks and the functioning of the ecosystem.

Acknowledgements

GL and CS are grateful for financial support by the Swiss National Science Foundation

(P2ZHP3_187938, PZ00P3_148261 and PP00P3_170645).

Authors' contributions

GL and CS designed the study; ENH conducted the experiment with the help of GL; XE,

CG, FJOS, AP, DS identified the specimens; GL analysed the data and wrote the

manuscript with inputs from CS. All authors commented and approved the final

publication.

The authors have declared no competing interest.

Data availability

Data and R code available from the Dryad Digital Repository

https://doi.org/10.5061/dryad.bzkh1897m

Supporting Information

Additional supporting information may be found online in the Supporting Information

section.

References

Albrecht, M., Schmid, B., Hautier, Y. & Müller, C.B. (2012) Diverse pollinator

communities enhance plant reproductive success. *Proceedings of the Royal Society B:*

Biological Sciences, **279**, 1748.

Armas, C., Rodríguez-Echeverría, S. & Pugnaire, F.I. (2011) A field test of the stress-

gradient hypothesis along an aridity gradient. *Journal of Vegetation Science*, **22**, 818-

827.

Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting Linear Mixed-Effects

Models Using lme4. *Journal of Statistical Software*, **67**, 1-48.

Bertness, M.D. & Callaway, R. (1994) Positive interactions in communities. *Trends in*

Ecology and Evolution, **9**, 191-193.

Blüthgen, N. & Klein, A. (2011) Functional complementarity and specialisation: the role

of biodiversity in plant–pollinator interactions. *Basic and Applied Ecology*, **12**, 282-

291.

Bonacich, P. (1987) Power and Centrality: A Family of Measures. *American Journal of*

Sociology, **92**, 1170-1182.

Braun, J. & Lortie, C.J. (2019) Finding the bees knees: A conceptual framework and

systematic review of the mechanisms of pollinator-mediated facilitation.

Perspectives in Plant Ecology, Evolution and Systematics, **36**, 33-40.

Brooker, R.W., Maestre, F.T., Callaway, R.M., Lortie, C.J., Cavieres, L.A., Kunstler, G....

Michalet, R. (2008) Facilitation in plant communities: the past, the present, and the

future. *Journal of Ecology*, **96**, 18-34.

Brooker, R.W., Callaway, R.M., Cavieres, L.A., Kikvidze, Z., Lortie, C.J., Michalet, R.,

Pugnaire, F.I., Valiente-Banuet, A., Whitham, T.G. (2009) Don't Diss Integration: A

Comment on Ricklefs's Disintegrating Communities. *The American Naturalist*, **174**,

919-927.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A.,

Skaug, H.J., Maechler, M. & Bolker, B.M. (2017). glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling.

The R Journal, **9**, 378-400.

Bruno, J.F., Stachowicz, J.J. & Bertness, M.D. (2003) Inclusion of Facilitation into

Ecological Theory. *Trends in Ecology and Evolution*, **18**, 119-125.

Butterfield, B.J., Cavieres, L.A., Callaway, R.M., Cook, B.J., Kikvidze, Z., Lortie, C.J....

Brooker, R.W. (2013) Alpine cushion plants inhibit the loss of phylogenetic diversity

in severe environments. *Ecology Letters,* **16**, 478-486.

Callaway, R.M. (2007) *Positive interactions and interdependence in plant communities.*

Springer, The Netherlands.

Cavieres, L.A., Broojer, R.W., Butterfield, B.J., Cook, B.J., Kikvidze, Z., Lortie, C.J....

Callaway, R.M. (2014) Facilitative plant interactions and climate simultaneously

drive alpine plant diversity. *Ecology Letters*, **17**, 193-202.

Chapin, F.S., Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L.,

Hooper, D.U., Lavorel, S., Sala, O.E., Hobbie, S.E., Mack, M.C. & Díaz, S. (2000)

Consequences of changing biodiversity. *Nature*, **405**, 234-242.

Csardi, G. & Nepusz, T. (2006) The igraph software package for complex network

research. *International Journal Complex Systems,* **1695**. 2006. http://igraph.org

Connell, J. H. & Slatyer, R. O. (1977) Mechanisms of succession in natural communities

and their role in community stability and organization. *The American Naturalist*,

111, 1119–1144.

DeAngelis, D.L., Post, W.M. & Travis, C.C. (1986) *Positive Feedback in Natural Systems*.

Springer, Berlin, Germany.

Ebeling, A., Klein, A.M., Schumacher, J., Weisser, W.W. & Tscharntke, T. (2008) How does

plant richness affect pollinator richness and temporal stability of flower visits?

Oikos, **117**, 1808-1815.

Ellison, A.M. (2019) Foundation species, Non-trophic interactions, and the value of

being common. *iScience*, **13**, 254-268.

Feldman, T. S., Morris, W. F. & Wilson, W. G. (2004). When can two plant species

facilitate each other's pollination? *Oikos*, **105**, 197–207.

Fox, J. & Weisberg, S. (2019). An {R} Companion to Applied Regression, Third Edition.

Thousand Oaks CA: Sage. URL:

https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Fründ, J., Dormann, C.F., Holzschuh, A. & Tscharntke, T. (2013) Bee diversity effects on

pollination depend on functional complementarity and niche shifts. *Ecology*, **94**,

2042-2054.

Gazhoul, J. (2006) Floral diversity and the facilitation of pollination. *Journal of Ecology,*

94, 295-304.

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J.H.,

Lodge, D.M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A.J., Vandermeer, J.

& Wardle, D.A. (2005) Effects of biodiversity on ecosystem functioning: a consensus

of current knowledge. *Ecological Monographs*, **75**, 3-35.

Hunter, A.F. & Aarssen, L.W. (1988) Plants helping plants. *Bioscience*, **38**, 34–40.

Keddy, P. (2017) *Plant Ecology: Origins, Processes, Consequences*. Cambridge University

Press, UK.

IPBES (2019) *Global assessment report on biodiversity and ecosystem services of the*

Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. E.

S. Brondizio, J. Settele, S. Díaz, and H. T. Ngo (editors). IPBES secretariat, Bonn,

Germany.

Krugman, P. (1991). Geography and Trade. MIT Press.

Jiang, L., Pu, Z. & Nemergut, D.R. (2008) On the importance of the negative selection

effect for the relationship between biodiversity and ecosystem functioning. *Oikos*,

117, 488–493.

Laverty, T.M. & Plowright, R.C. (1988) Fruit and seed set in Mayapple (*Podophyllum*

peltatum): influence of intraspecific factors and local enhancement near *Pedicularis*

canadensis. Canadian Journal of Botany, **66***,* 173-178.

Laverty, T.M. (1992) Plant interactions for pollinator visits: a test of the magnet species

effect. *Oecologia*, **89**, 502–508.

Lázaro, A., Jakobsson, A. & Totland, O. (2013) How do pollinator visitation rate and seed

set relate to species' floral traits and community context? *Oecologia*, **173**, 881-893.

Lenth, R. (2020). emmeans: Estimated Marginal Means, aka Least-Squares Means. R

package version 1.4.8. https://CRAN.R-project.org/package=emmeans

Levin, S.A. (2009) *The Princeton Guide to Ecology*. Princeton University Press, USA.

Liancourt, P. & Dolezal, J. (2020) Community-scale effects and strain: Facilitation

beyond conspicuous patterns. *Journal of Ecology*, in press.

Loreau, M. & Hector, A. (2001) Partitioning selection and complementarity in

biodiversity experiments. *Nature*, **412**, 72-76.

Lortie, C.J., Filazzola, A. & Sotomayor, D.A. (2016) Functional assessment of animal

interactions with shrub-facilitation complexes: a formal synthesis and conceptual

framework. *Functional Ecology*, **30**, 41-51.

Losapio, G., Pugnaire, F.I., O'Brien, M.J. & Schöb, C. (2018) Plant life history stage and

nurse age change the development of ecological networks in an arid ecosystem.

Oikos, **128**, 1390–1397.

Losapio, G. & Schöb, C. (2020) Pollination interactions reveal direct costs and indirect

benefits of plant-plant facilitation for ecosystem engineers. *Journal of Plant Ecology*,

13, 107–113.

Losapio, G., Schmid, B., Bascompte, J., Michalet, R., Cerretti, P., Germann, C., Haenni, J.P.,

Neumeyer, R., Ortiz-Sánchez, F.J., Pont, A.C., Rousse, P., Schmid, J., Sommaggio, D. &

Schöb, C. (2021) An experimental approach to assessing the impact of ecosystem

engineers on biodiversity and ecosystem functions. *Ecology,* **e03243.**

Lozano, Y.M., Armas, C., Hortal, S., Casanoves, F. & Pugnaire, F.I. (2017) Disentangling

above- and below-ground facilitation drivers in arid environments: the role of soil

microorganisms, soil properties and microhabitat. *New Phytologist,* **216**, 1236–

1246.

McIntire, E.J. & Fajardo, A. (2014) Facilitation as a ubiquitous driver of biodiversity. *New*

Phytologist, **201**, 403-416.

Michalet, R. &Touzard, B. (2010) *Biotic interactions, biodiversity, and community*

productivity. In (eds. F.I. Pugnaire) Positive plant interactions and community

dynamics. Taylor & Francis, USA.

Montesinos-Navarro, A., Valiente-Banuet, A. & Verdú, M. (2019) Mycorrhizal symbiosis

increases the benefits of plant facilitative interactions. *Ecography,* **42**, 447-455.

Moro, M. J., Pugnaire, F. I., Haase, P. & Puigdefabregas, J. (1997). Effect of the Canopy of

Retama sphaerocarpa on Its Understorey in a Semiarid Environment. *Functional*

Ecology, **11**, 425–431.

O'Connell, L.M. & Johnston, M.O. (1998) Male and female pollination success in a

deceptive orchid, a selection study. *Ecology,* **79**, 1246-1260.

Poisot, T. (2016). betalink: Beta-Diversity of Species Interactions. R package version

2.2.1. https://CRAN.R-project.org/package=betalink

Pugnaire, F.I., Haase, P. & Puigdefabregas, J. (1996) Facilitation between higher plant

species in a semiarid environment. *Ecology*, **77**, 1420-1426.

Pugnaire, F.I. (2010) *Positive Plant Interactions and Community Dynamics*. CRC Press,

Cleveland, Ohio, United States.

Rodríguez-Echeverría S, Armas C, Pistón N, Hortal S & Pugnaire FI. (2013) A role for

below ground biota in plant–plant facilitation. Journal of Ecology 101: 1420–1428.

Rodríguez-Echeverría S, Lozano YM &Bardgett RD (2016) Influence of soil microbiota in

nurse plant systems. Functional Ecology 30: 30–40.

Rodríguez-Riaño, T., Ortega-Olivencia, A. & Devesa, J.A. (1999) Reproductive phenology

in three Genisteae (Fabaceae) shrub species of the W Mediterranean Region. *Nordic*

J. Bot. 19: 345-3

Rohr, J.R., Civitello, D.J., Hlliday, F.W., Hudson, P.J., Lafferti, K., Wood, C.L. &Mordecai, E.A.

(2020) Towards common ground in the biodiversity–disease debate. *Nature Ecology*

& Evolution, **4**, 24–33

Scherber, C., Eisenhauer, N., Weisser, W., Schmid, B., Voigt, W., Fischer, M.... Tscharntke,

T. (2010) Bottom-up effects of plant diversity on multitrophic interactions in a

biodiversity experiment. *Nature*, **468**, 553-556.

Schlesinger, W.H, Raikes, J.A., Hartley A.E. & Cross, A.F. (1996) On the Spatial Pattern of

Soil Nutrients in Desert Ecosystems. *Ecology*, **77**, 364-374.

Schleuning, M., Fründ, J. & García, D. (2015) Predicting ecosystem functions from

biodiversity and mutualistic networks: an extension of trait-based concepts to

plant–animal interactions. *Ecography*, **38**, 380-392.

Schöb, C., Brooker, R.W. & Zuppinger-Dingley, D. (2018) Evolution of facilitation

requires diverse communities. *Nature Ecology & Evolution*, **2**, 1381-1385.

Tilman, D., Isbell, F. & Cowles, J.M. (2014) Biodiversity and Ecosystem Functioning.

Annual Review of Ecology, Evolution, and Systematics, **45**, 471-493.

Valiente-Banuet, A. et al. (2015) Beyond species loss: the extinction of ecological

interactions in a changing world. *Functional Ecology*, **29**, 299-307.

Winfree, R., Reilly, J.R., Bartomeus, I., Cariveau, D.P., Williams, N.W. & Gibbs, J. (2018)

Species turnover promotes the importance of bee diversity for crop pollination at

regional scales. *Science,* **359**, 791-793.

Wright, A.J., Wardle, D.A., Callaway, R. & Gaxiola, A. (2017) The overlooked role of

facilitation in biodiversity experiments. *Trends in Ecology and Evolution*, **32**, 383-

390.

Wright, J.P. & Jones, C.G.P. (2004) Predicting effects of ecosystem engineers on patch-

scale species richness from primary productivity. *Ecology*, **85**, 2071-2081.

Figures and tables

Fig. 1 Conceptual framework and experimental design. (A) A legume nurse shrub

(*Retama sphaerocarpa*) facilitates diverse herbaceous plant species in the understorey.

(B) To examine the joint effects of the nurse plant facilitation and biodiversity change

on mutualistic networks, the experimental design includes the treatments of nurse

presence/absence and understorey plant diversity. Plant and flower density of

understorey species were kept constant. Furthermore, a nurse alone treatment (i.e.,

shrub without understorey plants) was included. (C) We then examined the response of

pollination interactions, such as those between *Apis mellifera* and *Retama sphaerocarpa*.

(D) Process pathways among facilitation, diversity, mutualistic networks and ecosystem

functioning.

624

Fig. 2 Effects of the nurse plant presence (open vs nurse) and understorey plant

biodiversity (1 vs 3 species) on abundance (A) and richness (B) of the pollinator

community (i.e., flower visitors) per 20 min per m^2. Bars indicate standard errors.

Fig. 3 Cost and benefit analysis of nurse plant presence (open vs nurse) and

understorey plant diversity (1 vs 3 understorey species) for understorey plants (A) and

nurse shrubs (B) in terms of flower visits per 20 min per m^2. Bars indicate standard

errors.

Fig. 4 Consequences of nurse plant presence (open vs nurse) for complementarity and

selection effects among understorey species (A). Complementarity and selection effects

between nurse plants and understorey species (B). Bars indicate standard errors.

Fig. 5 Network centrality of additive networks (red) and observed networks (blue).