On the joint effect of spatial attention and defocus blur on acuity: attentional limit to the resolving power of the eye

De Lestrange-Anginieur E[1*], Leung TW[1], and Kee CS[1,2]

1. School of Optometry, Hong Kong Polytechnic University, Hong Kong
2. Interdisciplinary Division of Biomedical Engineering, Hong Kong Polytechnic University, Hong Kong SAR, China

Correspondence and requests for materials should be addressed to E.D.L. (email: elie.delestrange@hotmail.com).

Abstract

Defocus blur and spatial attention both act on our ability to see clearly over time. However, it is currently unknown how these two factors interact because studies on acuity resolution only focused on the separate effects of attention and defocus blurs. In this study, resolution acuity was measured along the diagonal 135°/315° with horizontal, at 8° eccentricity for clear and blur Landolt C images under various manipulations of covert endogenous attention. We observe that attention not just improves the resolution of clear stimuli, but also modulates the resolution of defocused stimuli for compensating the loss of resolution caused by retinal blur. Our results show, however, that as the degree of attention decreases, the differences between clear and blurred images largely diminish, thus limiting the benefit of an image quality enhancement. It also appeared that attention tends to enhance the resolution of clear images more than blurred targets, suggesting potential variations in the gain of vision correction with the level of attention. This demonstrates that the interaction between spatial attention and focus plays a role in the way we see things. In view of these findings, the development of adaptive (neuro-optical) interventions, which adjust the eye's focus to attention, may hold promise.

Significance statement Visual technologies are now attaining a degree of extreme sophistication and diversity, which allows more comprehensive, but often complex manipulations of the optical image formed onto the back of the eye. It is therefore an enigma how those fine and immersive manipulations of the sensory environment are integrated in the brain. In this study, we show that the resolving power of the eye can depend complexly on the interaction between spatial attention and focus. This discovery suggests that perception might be advantageously guided by technologies tailoring optical focus to individual attentional patterns.

Our ability to perceive the constituents of a visual stimulation strongly depends on the optical focus of the eye that defines the state for which the smallest spatial feature of the environment can be resolved. The control of optical focus has been one the most significant challenges of optics, and continues to drive a large number of visual technologies, such as ophthalmic visual aids (such as spectacles, contact lenses), but also display technologies (Virtual reality, vision-correcting light field display) aimed at improving and correcting the human sensory experience. Often controlling more than one visual target, those technologies have become both adaptive [**1-**

2] and more complex [**3**], capable of manipulating image quality over time and space, which may

broaden our experience and interaction with the environment. However, even with such

heightened optical manipulation, predicting the impacts of an optical correction on visual function

remains challenging due to an uncertainty as to how and to what extent optical modification

affects the way retinal images are processed by the brain. Considering a capacity-limited visual

system [**4**] and the infinite sum of information present in the visual world, augmentation of optical

information could be traded off by increase amount and time for neural computation to access a

neural representation, veridical with respect to the retinal image. It is known that usually not all

information entering the eye is effectively processed over our visual field, with some spatial and

temporal information never reaching consciousness or receiving attention [**5**]. Incompleteness of

processing or absence of attention could limit the gain of correcting or augmenting optical signals.

A vivid illustration of this phenomenon, the inattentional blindness paradigm [**6**], is the failure to

report a simple, suprathreshold stimulus or a stimulus attribute of the visual field in the absence

of attention, when the eye is engaged in another attention demanding primary task. Despite this

deficit, it has been shown that some amount of processing still occurs in absence of attention,

when the stimulus fails to be detected [**7-8**]. It has been demonstrated that the effect of attention

strongly depends on the type of spatial information [**9**], and so our visual experience may vary

with the characteristics of the environment. However, the effect of movement of attention on the

processing of optical signals (i.e., signals having a complex distribution of spatial frequency-

dependent contrast) has received little attention. Thus, to date, it is unknown to what extent

attention contributes to the perception of the fine optical signals unveiled by ocular correction. A

recent examination of the relationship between blur and attention proposed that blur detection

might be pre-attentively processed by the visual system [**10**]. Interestingly, this suggests that, even

in the absence of attention, when a stimulus attribute failed to be reported, visual blur might be

partly processed. Nevertheless, the degree to which blur analysis remains independent from

attention deployments is still unclear. Comparing the effect of blurred and clear images under

variation of attention could help determine the potential benefit of correcting visual blur under

real life contexts, when attention varies. To comprehensively examine this relationship, several

forms of attention and type of blur may be of interest. This study focused on the interaction

between endogenous covert attention and visual blur.

In covert attention [11], sensory enhancement occurs, without eye movements, by granting

priority of processing to a certain location of the visual field, which usually takes place at the

expense of processes in other regions stimulating the retina. This prioritization can occur via two

modes of control: The first mode, exogenous attention, is a rapid, reflex, automatic deployment

of attention in response to external visual stimulation, such as an abrupt, peripheral visual event,

requiring an immediate response. The second mode, endogenous attention, is under conscious,

voluntary control and can be directed according to the observer's visual goal provided there is

sufficient time of activation. In real life, both endogenous and exogenous factors [12] affect the

deployment of attention, which implies a complex interaction between environmental

characteristics, neural architecture, and cognitive behaviors. A major interest of endogenous

attention is that, unlike exogenous attention, it can be flexibly sustained at a position of the visual

field suitable for the task requirements [13-15]. In laboratory conditions, the strength of

endogenous attention can be manipulated via a central symbolic cue (e.g., an arrow pointing

towards the region of the cued location), and takes about 300ms from cue onset to deploy.

Neurophysiological studies have shown that as attention deploys, sensitivity (contrast gain [16])

or/and firing rate (response gain [17]) of neurons are increased, altering the relationship between

stimulus contrast and neurons response. This can manifest in an enhancement of several

perceptual tasks and stimulus properties, including resolution acuity [18-27], resolvable and

apparent contrast [28-29]. Failure to adequately direct attention to environmental stimulation has

been shown to drastically impair sensory performance. How does this attentional variability affect

an optical ocular correction? Recent behavioral studies demonstrate the existence of a preferential

enhancement of neurons tuned to high spatial frequency **[30]**, with endogenous attentional

modulation at both low and high spatial frequencies **[31]**. This differential effect suggests that

variations of neural filtering by the focus of attention could potentially alter the *complex*

appearance of broadband stimuli, having more than one spatial frequency. Such alteration could

take place under various forms depending on the spatial extent of attention modulation in

broadband images containing more than one spatial frequency. It is however unknown how

attention varies with the complex distribution of spatial frequency pattern set by retinal blur. An

account of both optical transfer function and neural transfer function could predict the

neurooptical transformation of individual retinal images [**32**]. Unfortunately, because previous

measurements of the contrast sensitivity function [**31, 33**] did not control the effect of blur on

attention, it remains unknown how attention reshapes the neural transfer function, which is the

neural part determining perception independently of the optics of the eye. One of the questions

that ensues is whether the modulation of attended and unattended stimulus is affected by the

differentiation of ocular optical filtering across spatial frequencies and the variation of ocular blur

across individuals. For example, the attenuation of high spatial frequencies in uncorrected stimuli

could increase or decrease focus of attention with neural adaptation to the image the visual system

decodes.

In this study, we were interested to examine whether, or not, there is a possible interaction

between blur and attention on acuity, and its potential implication when correcting the eye via

optical or attentional manipulations. We noted that previous studies assessing the impact of blur

in acuity [**34-40**] systematically overlooked the effect of attention on acuity, and vice versa. To

the best of our knowledge, this is the first studies that examined the joint effect of blur and spatial

attention on resolution acuity. Since attention can affect differentially spatial frequencies [9,30-

32], the variation of the spatial frequency distribution with retinal blurs could vary the effect of

attention, and thus have practical implications when correcting retinal blurs with an ophthalmic

correction. Then, to which extent, and how? Here, we hypothesized that blurred stimuli,

exhibiting greater attenuation of contrast-dependent spatial frequency [41], may benefit less from

attentional enhancement, as compared to clear stimuli. To test the effect of retinal defocus blur

on attention, we used the "source method" proposed by Haig et al [42], which is a widely used

method for assessing the visual impact of ocular aberrations [34-40], and the most optimum

approach for simulating spatiotemporally-varying blurs stimuli, which cannot be simulated with

adaptive optics. This study shows evidence that the deployments of attention in non-foveal

locations act on the visual effect of correction by altering the acuity distinctions between clear

and blur stimuli. These findings demonstrate the existence of an attentional modulation of the

effect of retinal blurs, and may motivate the development of novel interventions based on the

adjustment of focus.

Results

Adapting a Posner's cueing paradigm introduced by other researchers [18], resolution acuity

was tested under the manipulation of blur and covert endogenous spatial attention. The visual

stimuli consisted of a pair of Landolt C letters (Fig. 1, step 3; "target" and "non-target", with

different gap orientations in each) briefly displayed at two locations in the near-periphery of

the visual field at 8° eccentricity. Spatial attention was manipulated using three cueing

conditions (Fig. 1, step 2), whereby a cue preceded the Landolt C stimuli: (1) cued condition,

in which a cue was displayed and pointed at the upcoming target Landolt C location; (2) uncued

condition, in which the cue pointed at the non-target upcoming Landolt C location; and (3)

neutral condition, in which two cues were simultaneously displayed and pointed at both target

and non-target locations. In cued and uncued trials, observers were required to attend to a cued

location (either north-west [NW] or south-east [SE]) while fixating on a small central cross. In

neutral trials, observers were required to spread their attention and focus on both locations (i.e.,

NW and SE). At the end of the stimulus presentation sequence, the location of the target Landolt

C (Fig. 1, step 4) was indicated by a response cue, which consists of a central line symbol

pointing towards one of the two stimuli.

The visual task was to identify the orientation of the gap in the "target" Landolt C (6AFC; see

Methods for details), as indicated by the response cue. Visual acuity performance was measured

under both clear and blurred conditions (Fig. 1, step 3). The gap size was controlled using an

adaptive 1-down-1-up staircase procedure. A computerized image processing technique was

employed to simulate the effects of a moderate amount of optical defocus on the retinal image

of the Landolt C stimulus [**43**] (defocus blur, two waves of RMS wavefront error, about 1.25

diopters; see Methods for details). We chose to use a *clear* attentional cue and fixation stimuli

in all the conditions so that the magnitude of the attention deployment was identical when

comparing clear and blurred target conditions. There was a total of 12 interleaved staircases,

for a total of 12 conditions (i.e., two retinal locations x three cue conditions x two blur levels).

In addition to visual acuity (VA), response time (RT) was also measured.

Eleven young adults with corrected-to-normal vision participated. The visual acuity and

response time data were processed using a three-way repeated measures analysis of variance

(RANOVA) test (retinal location: NW and SW; visual blur: clear and blurred; spatial cueing:

cued, uncued, and neutral), and post-hoc pairwise comparisons were performed with the

Bonferroni correction using SPSS.

Spatial location. Spatial location of the target had a statistically significant effect on the

overall performance for RT (three-way RANOVA: $F_{(1,10)}= 5.473$ p=.041, $\eta^2=0.364$), but not

for VA (three-way RANOVA: $F_{(1,10)}=2.313$ p=.159, $\eta^2=0.188$). There was no significant

interaction between spatial cueing effect and spatial location for either RT (three-way

RANOVA RT: $F(1.184,20)=3.227$, p= .094, $\eta^2=0.244$) or VA (three-way RANOVA VA:

$F(2,20)= .081$ p=.922, $\eta^2=0.008$), suggesting that spatial cueing effect was little affected by

the position of the target along the diagonal (135°/315° with horizontal).

Spatial cueing. Spatial cueing was found to significantly influence VA (Fig. 2a, three-way

RANOVA: $F(1.109,11.091)=21.962$, p=0.001, $\eta^2=0.687$), as reported previously [18-27], and

response time (Fig. 2c, three-way RANOVA: $F(2,20)=28.919$, p<0.001, $\eta^2=0.743$). The

strength of spatial cueing tended to vary with the participant's acuity level (see Fig.1 in the

Supplementary Materials). Visual acuity performance was superior for cued stimuli when

compared to that for neutral stimuli (Bonferroni post-hoc test VA: mean difference=1.981,

arcmin, p=0.012; RT: mean difference=-0.052 ms, p=0.019) or uncued stimuli (Bonferroni

post-hoc test VA: mean difference=5.963, arcmin, p=0.002; RT: mean difference=0.160 ms,

p=0.001). Using the neutral condition as a reference, the data sets were replotted as changes in

visual acuity in Fig. 2b and changes in response time in Fig. 2d. Note that positive values

represent an increase in visual acuity performance, and vice versa. The overall cost of attending

to an incorrect location was greater than the benefit of attending to a correct location by two-

fold for both RT and VA.

Effect of spatial cueing on clarity We replotted in Fig. 3 the spatial cueing data for both clear

and blurred targets showing that, under both conditions, VA and RT increased under the cued

condition (Bonferroni post-hoc test VA: blurred target: mean difference = 1.528, p=0.02; clear

target: mean difference = 2.433, p=0.017) and decreased under the uncued condition

(Bonferroni post-hoc test VA: blurred target: mean difference = 3.855, p=0.002; clear target:

mean difference = 4.109, p=0.003), when compared with the neutral cue condition.

Nevertheless, visual blur significantly influenced VA (three-way RANOVA: $F(1,10)=54.203$,

p<0.001, $\eta^2=0.844$). As for spatial cueing, the strength of blurring tended to vary with the

participant's acuity level (see Fig.2 in the Supplementary materials). Interestingly, there were

also a tendency from subjects suffering the most from blurring to show less effect of attention

effect (see Fig.3 in the Supplementary materials). This could suggest a relationship between

perceived blur and attentional deployment. Notably, a significant interaction was found

between visual blur and spatial cueing for VA (three-way RANOVA VA: $F(2,20)=3.586$,

$p=0.047$, $\eta^2=0.264$), but not for RT (3-way RANOVA: $F(2,20)=0.142$, $p=0.869$, $\eta^2=0.014$). A

significant impact of blur (Fig. 3e, VA_{blur}-VA_{clear}) was observed for both cued and neutral

conditions, but not for the uncued condition (Bonferroni post-hoc test VA: cued: mean

difference = 1.857 arcmin, $p<0.001$; neutral: mean difference= 0.952 arcmin, $p=0.015$; uncued:

mean difference= 0.698, arcmin, $p=0.074$). Most remarkably, the suboptimal attentional

conditions (i.e., neutral and uncued) reduced the difference in resolution between blurred and

clear stimuli, suggesting that as attention is diverted from the resolution task at hand, visual

differences between stimuli having distinct focus are no longer processed.

Existence of a neuro-optical balance. This equalization of in and out of focus stimulation

reveals that the alteration and augmentation of acuity involve an interaction between optical

quality of the eye and the movement of attention over the visual field. As depicted in Fig. 4, a

paired sampled t-test shows that a defocus system under full attention (attended blur image:

6.72 ± 1.42) exhibits superior performance than a perfectly focused system with reduced

attention (unattended clear image: 11.40 ± 5.14); $t(21) = 4.58$, $p<0.001$. This indicates that,

by increasing a person's attention, it is possible to compensate for a large drop in optical

resolution. Similarly, a defocus system under full attention (attended blur image: $6.716 \pm$

1.417), has similar performance than a focused system with reduced attention (neutral clear

image: 7.292 ± 3.347); as a paired sampled t-test shows no statistical difference $t(21) = 1.04$,

$p=0.312$. This highlights that for a given acuity level, it is possible to have varying combinations

of attention and focus. This highlights that natural modulation of attention (e.g., occurring under

prolonged sustained attention [44-46]) can be counterbalanced by an increase in optical

resolution.

Effect of clarity on spatial cueing Interestingly, it appears that the global cueing effect (Fig.

5b, VA_{cued}/VA_{uncued}) was stronger for clear targets than blurred targets (Paired-samples t-test:

mean difference: 0.591, paired t(10)=3.37, p=0.007), which clearly revealed that spatial

attention enhanced acuity more with clear than blurred vision. Albeit speculative, a plausible

explanation is that the effective width of the attentional filter is broader for clear vision, because

of the augmentation of the suprathreshold components of the image (e.g., towards the higher

spatial frequencies) on which attention can operate, as illustrated in Fig. 6. While the cueing

gain (Fig. 5a, $VA_{cued}/VA_{neutral}$) was significantly augmented in clear targets compared to blurred

targets (Paired-samples t-test: mean difference: 0.290, paired t(10)=2.37, p=0.039), the cueing

cost (Fig. 5a, $VA_{uncued}/VA_{neutral}$) did not appear to be influenced by the blurring condition

(Paired-samples t-test: mean difference: -0.02, paired t(10)=-0.061, p=0.557). The absence of

difference in acuity reduction (from the neutral baseline) for blurred and clear targets is

compatible with the idea that, below a certain level of attention (here, referring to the neutral

conditions), the information modulated by attention in clear targets becomes similar to that for

blurred targets. Those findings indicate that the efficiency of attention deployment is somewhat

dependent on the image quality of the human eye at a given location. If attention do depend on

the retinal blurred image, the particular distribution of blur across the visual field, which is

known to vary with refractive populations, could matter in the distribution of spatial attention.

This asks whether the retinal blurs resulting from the eye growth are merely an outcome of

biological constraints, or could involve some neural feedback mechanisms involved in the

regulation of the attentional resources?

Simulated refractive gain. The data were replotted in terms of visual acuity gain

(VA_{blur}/VA_{clear}) as shown in Fig. 7a. A significant decrease (one-way RANOVA, F(2,20)=7.697,

p=0.003, η^2=0.435) in refractive gain by a factor of approximately 1.26 and 1.19 was found

under neutral and uncued conditions, respectively, compared to the cued condition, indicating

that a certain degree of attention is required for attaining a benefit of sufficient value from blur

correction. Fig. 7b shows the simulated refractive gain variations associated with modulation

in attention from the neutral condition. It is worth noting that, under suboptimal attentional

conditions, the correction of retinal blur may not increase acuity although stimuli are still

detected. This finding identifies attention as a prerequisite of vision enhancement in the

perifovea.

Discussion

In this study, it was demonstrated that acuity depends on both attentional and optical factors,

throwing light, for the first time, on the way resolution acuity is modulated by variations in

endogenous covert attention and the eye's focus (Fig. 2-4, 6-7). This complements a large

number of studies in the literature that have till now discounted the interaction between

attention and optics, focusing only on the separate effects of attention [**18-32**] and ocular optics

[**34-40**] per se in acuity resolution.

Tolerance to retinal blurs We showed that processing of blur is strongly influenced by the

level of attention, and so not simply automatically processed [**10**]. Specifically, it was

determined that attention not only improved images that are sharp [**18-27**], but can also

drastically improved the resolution of degraded retinal images (Fig. 2a-c) to provide more

tolerance to blur. This tolerance suggests the possibility of an adaptive compensation by

attention of the retinal blurs degrading vision, which, might enable the eye to rely less on the

optical ocular quality. This may be particularly useful to adapt the unwanted spatiotemporal

blurs of the visual field that are caused by the motion of the eye and visual targets. Besides, it

could allow relaxation of the constraint on optical quality required to achieve a certain

performance when recomposing these blurs (as is the trend with progressive addition lenses).

On the other hand, under diminished attention (Fig. 3e), an optical enhancement could present

the sure advantage to providing greater acuity compared to a blurred image. A riveting question

remains how optical modulations affect our attentional responses and resources.

Attentional limit to supervision Our results reveal that blurring does modulate the impact of

attention. We show that attention boosts more acuity resolution in clear stimuli than blurred

stimuli (Fig. 5). While this seems to accord with the idea that attention enhance more retinal

stimuli having the highest level of details [**30**], it is also plausible that the increased contrast

across spatial frequencies in clear stimuli favor attentionnal modulation. Supposing that clear

retinal stimuli are more attentionally demanding, ocular blurs brought by the eye growth could

be a way for the visual system to regulate the deployment of attentional resources across the

visual field, and adjust the cues of the environment. A practical, and potentially interesting,

consequence of a differential attentional modulation is that the decrease of attention can alter

the resolution gain of optical correction (Fig. 7): the conditionality of beneficial effects of

correction in daily activities can have certain implications for visual technologies aimed at

augmenting visual resolution. For instance, a super-resolved optical system [**40**] may only be

effective if subjects allocate sufficient attentional resources to a given location. Such

contingency of performance requires an individual's attentional pattern to be considered when

determining the level of correction. Indeed, the observed effect of attention and blur and their

interaction could vary between individuals because of several factors, including ocular

aberration, neural sampling, neural adaptation but also maybe individual attentionnal patterns.

Dynamic focus correction. An expected, but seminal, finding of this study is that a given

level of acuity can involve different combinations of attention and focus (Fig. 4) -that is, a

focused system with reduced attention can perform closely to a defocused system under full

attention. This provides evidence that changes in attention might be balanced by external or/and

changes in optical focus, and vice-versa. Given the incessant modulation of attention with time

[15], environmental settings [12], but also among individuals [47], the use of adaptive optical

technology could constitute a unique opportunity to adjust the modulation of attention at the

optical level. While such optical compensation is not feasible using standard static corrections,

given their fixed focus, several emerging technologies, such as spectacles-free display [1] and

adaptive optics spectacles [2], show the potential to dynamically adjust the level of acuity to

the movement of attention via an adjustable focus, which could thus timely control the

resolution of the neural images. While a dynamic optical correction might be simply based on

the person's feedback, in the case of an automatic optical focus, it would require the ability -

not without challenges- to continuously sense and decode the eye responses to access the

temporal variations of attention. Combining real-time eye-tracking and control of the optical

focus of neural images may open up unique and exciting horizons to respond to individual

visual needs, which hold promises for the development of visual aids capable of dynamically

linking optical inputs and neural outputs.

Future development. This study has some limitations. First, the control of ocular aberrations

in the extrafoveal regions of the retina was not possible, as, to date, there is no visual simulator

capable to correct aberration over a wide viewing angle. Visual simulator correcting the

aberrations of the eye are restricted by the isoplanatic patch of the eye to a small visual angle

(of about 1-2degrees) that allows testing only one peripheral location at once [48]. This severely

limits the manipulation of spatial attention across distant locations of the visual field. By using

an adaptive optical system incorporating several deformable mirrors and wavefront sensors, it

may be possible in the future to enlarge the angular extent of adaptive ocular correction of the

eye [49], and so the limitation of conventional visual simulators, which incorporate only a

single deformable mirror. The use of a wide field AO would help, not just to simulate different

retinal blurs, but also compensate for the aberrations of the eye. Such compensation is essential

to investigate a possible effect of neural adaptation to natural peripheral blurs in spatial attention.

For example, sensitivity to the simulated blurred images could be influenced by individual

differences in natural peripheral blurs the eye may be adapted to [50]. Nevertheless, little is

known about the degree of adaptation to *natural* peripheral aberrations, though recent studies

in Yoon Lab suggests that adaptation to defocus could differ with individual refractive errors

in myopes and emmetropes [51]. Further works will be required to elucidate this. A second

limitation of the study is that the interleaved test was restricted to a limited number of retinal

conditions. It is plausible that the joint effect of attention and blur may vary as the characteristic

of ocular blur changes across the visual field, but a comprehensive understanding of these

parameters could involve controlling for the variation of ocular blur across eccentricities, as

recently performed in our Lab via a multiscale visual simulator [52]. To avoid the confounding

effects of the individual ocular aberrations on the simulated retinal blurred images, a

sufficiently large blur, producible on the display, was considered, excluding blur conditions

with very small and large amounts of retinal blurs. Other limitations could involve the

complexity of the stimulus and task. For instance, we show in a recent study [53] that, when

individuals perform a simple detection task, the effect of exogenous spatial attention on

simulated blurred images is small, suggesting a plausible decrease of interaction between blur

and attention as the attentional demand required by the task diminishes. Given that exogenous

and endogenous attentional filters involve different neuronal pathways, the dynamic of

attention might influence an interaction between blur and attention. Further works will be

needed to elucidate how these factors affect the processing of blur processing by attention.

Conclusion

In sum, our results proved, for the first time, a joint effect of optical and attentional factors on acuity resolution, showing that both attention and the eye's optics matters in the way we perceive things: both the degree of attention and optical correction vary the acuity resolution of the visual system, which potentially allows distinct (neuro-optical) combinations for achieving a given visual resolution. The interaction between those two visual factors is, however, more complex than thought, as acuity enhancement is not equal for focused and defocused images, the movements of attention could modulate an optical correction of the eye. This suggests an important avenue of exploration for adaptive optical technologies. Indeed, the neuro-optical interaction investigated in the present study for localized, defocused images is one only tiny facet of the iceberg, with a more important question perhaps, that is: how, and to which extent, a person's attention interact with the patterns of blurring on the retina? We believe that further research utilizing new optical advances to control ocular blurs and attention over the entire visual field could contribute to unfolding this mystery. Future works shall examine how the interaction that may link the brain and the optics of the eye spreads across the visual field in real-world contexts.

Methods

Experimental design. Subjects were asked to fixate on a small cross displayed at the center of the monitor screen (Fig. 1, "+"; size, $0.5° \times 0.5°$). The endogenous covert attention of the subject was manipulated using a central cue preceding the visual stimuli. On cued and uncued trials, a central cue was displayed (size, 18×0.6 arcmin; exposure time, 293 ms), pointing at either one of the two upcoming Landolt C locations, to which the observer was required to allocate attention. In neutral trials, two central cues were displayed, pointing at both upcoming Landolt C locations. Two Landolt Cs ("target" and "non-target"), with different gap orientations, were then presented simultaneously after a 300 ms inter-stimulus-interval. At the end of the

presentation sequence, a central line symbol (size, 18 x 0.6 arcmin) was displayed to point the

location of the target Landolt C.

Three cue conditions were tested: (1) in the cued condition, the cue pointed towards the target

Landolt C location; (2) in the uncued condition, the cue pointed towards the non-target Landolt

C location; and (3) in the neutral condition, two cues were displayed, which pointed towards

both target and non-target locations. Each cue condition was displayed for one-third of the

response trials in each session (cued: uncued: neutral = 1:1:1). It should be noted that the

participants were required to direct their attention as instructed by the cue(s), and that they were

not informed of the proportions of the three cue conditions.

The experimental procedures were approved by the University Committee for the Protection

of Human Subjects (HSEARS20170103003), and the research was conducted according to the

principles expressed in the Declaration of Helsinki. Informed consent was obtained from each

participant. All those involved, except two of the observers, the authors J.T.L and D.L.E, were

inexperienced with psychophysical procedures and not informed of the purpose of the

experiments.

Simulated defocus blur. Visual performance was assessed under both clear (zero blur) and

blurred (defocus blur, 2 waves of RMS wavefront error, about 1.25D) conditions. Note that the

blurred Landolt C stimuli were graphically generated by convolution of the two-dimensional

luminance profile of the Landolt C and a point spread function, $h(x, y)$, of a 5 mm pupil ($2r$):

$$h(x, y) = \|F[g(u, v)]\|^2$$

$$g(u, v) = A(u, v)exp\left[\frac{i2\pi}{\lambda}W(u, v)\right]$$

where F denotes Fourier transform, $A(u, v)$ denotes pupil function, $W(u, v)$ denotes

wavefront aberration described by a set of Zernike polynomials [54]. In this study, the visual

stimuli (i.e. Landolt C letters) were presented in black against a green background, and therefore

λ was set to 550 nm. The wavefront aberration was calculated from the Zernike defocus

polynomials Z_2^0, as:

$$W_{def}(u, v) = c_2^0 Z_2^0(u, v)$$

403   The amount of defocus used in these experiments is given by [55]:

$$c_2^0 = \frac{M\, r^2}{4\sqrt{3}}$$

where M denotes the spherical equivalent in diopters, r the radius of the simulated pupil.

Visual acuity measurement. Eleven young adults with corrected-to-normal vision (age 24

- 39; VA 6/6 or better in each eye) performed a Landolt C acuity task at two locations in the

near-peripheral visual field (Fig. 1, 8° eccentricity; quadrant, NW and SE; along 135°/315° with

horizontal). The placement of the pair of stimuli at the intercardinal locations aimed to minimize

field performance and attentional asymmetries. The visual task was to identify the orientation

of a gap in the "target" Landolt C (6AFC: 30°, 90°, 150°, 210°, 270°, 330°). The stroke width

was one-fifth of the Landolt C size. The stimulus exposure duration was 33 ms. Viewing was

binocular.

An interleaved 1-down-1-up staircase procedure that converged on the 50 % correct level

(adjusted for 6AFC, guess rate=1/6) was used to control the gap size and measure the visual

acuity threshold. There was a total of 12 interleaved staircases for various stimulus conditions

(i.e., three cue conditions x two locations x two blur levels). The acuity test was repeated a total

of 15 times for each stimulus condition over five sessions (approximately 5400 response trials

in total), and the average value of the threshold measurements was taken as visual acuity. The

number of trials to determine acuity threshold varied between 15 to 30 trials depending on the

sequence of correct and incorrect answer that we defined via a stopping rule. Observer

responded using a keyboard and audio feedback was provided after each response. The time

taken for an observer to respond after the offset of the Landolt C stimuli was measured as

response time. Only response taking place after the response cue offset were considered,

response time longer than 1 second were excluded.

A 32-inch Dell LCD monitor (screen resolution, 3840 x 2160; background luminance, 25

cd/m^2; contrast, ~100 %) was used to display visual stimuli. Viewing distance was 1m.

Throughout the experiment, an Eye tracker (Tobbi TX 300) monitored eye fixation, from the

onset of the cue to the offset of the target. Trials with unstable fixation (eye movements > 2˚)

were discarded.

References

1. Huang, F. C., Wetzstein, G., Barsky, B. A. & Raskar, R. Eyeglasses-free Display: Towards Correcting

Visual Aberrations with Computational Light Field Displays. Acm T Graphic 33 (2014). doi:Artn

5910.1145/2601097.2601122

2. Hasan, N., Banerjee, A., Kim, H. & Mastrangelo, C. H. Tunable-focus lens for adaptive eyeglasses.

Opt Express 25, 1221-1233 (2017). doi:10.1364/OE.25.001221

3. Krueger, R.R., Applegate, R.A., MacRae, S. Wavefront Customized Visual Correction: The Quest for

Super Vision II. (Thorofare, NJ: Slack Inc. , 2004).

4. Lennie, P. The cost of cortical computation. Curr Biol 13, 493-497 (2003).

doi:10.1016/S0960-9822(03)00135-0

5. Desimone, R. & Duncan, J. Neural mechanisms of selective visual attention. Annu Rev Neurosci 18,

193-222 (1995). doi:10.1146/annurev.ne.18.030195.001205

6. Rock, I., Linnett, C. M., Grant, P. & Mack, A. Perception without Attention - Results of a New

Method. Cognitive Psychol 24, 502-534 (1992). doi:Doi 10.1016/0010-0285(92)90017-V

7. Moore, C. M. & Egeth, H. Perception without attention: Evidence of grouping under conditions of

inattention. J Exp Psychol Human 23, 339-352 (1997). doi:Doi 10.1037//0096-1523.23.2.339

8. Scholte, H. S., Witteveen, S. C., Spekreijse, H. & Lamme, V. A. F. The influence of inattention on

the neural correlates of scene segmentation. Brain Res 1076, 106-115 (2005).

doi:10.1016/j.brainres.

9. Lee, D. K., Koch, C. & Braun, J. Spatial vision thresholds in the near absence of attention. Vision

Research 37, 2409-2418, doi:Doi 10.1016/S0042-6989(97)00055-2 (1997).

10. Loschky, L. C. et al. Blur detection is unaffected by cognitive load. Vis Cogn 22, 522-547,

(2014). doi:10.1080/13506285.2014.884203

11. Posner, M. I., Snyder, C. R. R. & Davidson, B. J. Attention and the Detection of Signals. J Exp
Psychol Gen 109, 160-174(1980). doi:Doi 10.1037//0096-3445.109.2.160

12. Corbetta, M., Patel, G. & Shulman, G. L. The reorienting system of the human brain: from
environment to theory of mind. Neuron 58, 306-324 (2008). doi:10.1016/j.neuron.2008.04.017

13. Muller, H. J. & Rabbitt, P. M. A. Reflexive and Voluntary Orienting of Visual-Attention - Time
Course of Activation and Resistance to Interruption. J Exp Psychol Human 15, 315-330, doi:Doi
10.1037/0096-1523.15.2.315 (1989).

14. Yantis, S. & Jonides, J. Abrupt Visual Onsets and Selective Attention - Voluntary Versus
Automatic Allocation. J Exp Psychol Human 16, 121-134, doi:Doi 10.1037//0096-1523.16.1.121
(1990).

15. Nakayama, K. & Mackeben, M. Sustained and transient components of focal visual attention.
Vision Res 29, 1631-1647 (1989).

16. Reynolds, J. H., Pasternak, T. & Desimone, R. Attention increases sensitivity of V4 neurons.
Neuron 26, 703-714 (2000). doi:Doi 10.1016/S0896-6273(00)81206-4.

17. Lee, J. & Maunsell, J. H. The effect of attention on neuronal responses to high and low contrast
stimuli. J Neurophysiol 104, 960-971 (2010). doi:10.1152/jn.01019.2009.

18. Montagna, B., Pestilli, F. & Carrasco, M. Attention trades off spatial acuity. Vision Research 49,
735-745, doi:10.1016/j.visres.2009.02.001 (2009).

19. Chan, C., Smith, G. & Jacobs, R. J. Simulating refractive errors: source and observer methods. Am
J Optom Physiol Opt 62, 207-216 (1985).

20. Shiu, L.P., Pashler, H. Neglible effect of spatial precuing on identification of single digits. J. Exp.
Psychol. Hum. Percept. Perform. 20, 1037–1054 (1994).

21. Shiu, L.P., Pashler, H. Spatial attention and vernier acuity. Vision Res. 35, 337–343 (1995).

22. Yeshurun, Y., Carrasco, M. Spatial attention improves performance in spatial resolution
tasks. Vision Res. 39, 293–306 (1999).

23. Yeshurun, Y., Carrasco, M. Spatial attention improves performance in spatial resolution
tasks. Vision Res.39, 293-306 (1999). doi:10.1016/s0042-6989(98)00114-x

24. Shalev, L., Tsal, Y. Detecting gaps with and without attention: Further evidence for attentional
receptive fields. European Journal of Cognitive Psychology. 14, 3–26 (2002).

25. Carrasco, M., Williams, P.E., Yeshurun, Y. Covert attention increases spatial resolution with or
without masks: support for signal enhancement. J Vis. 2, 467-479 (2002). doi:10.1167/2.6.4

26. Golla, H., Ignashchenkova, A., Haarmeier, T., Their, P. Improvement of visual acuity by spatial
cueing: a comparative study in human and non-human primates. Vision Res. 44, 1589-1600 (2004).
doi:10.1016/j.visres.2004.01.009

27. Yeshurun, Y., Carrasco, M. The effects of transient attention on spatial resolution and the size of the
attentional cue. Percept. & Psychophys. 70, 104–113 (2008).

28. Carrasco, M., Ling, S. & Read, S. Attention alters appearance. Nat Neurosci 7, 308-313,
doi:10.1038/nn1194 (2004).

29. Liu T, Abrams J, Carrasco M. Voluntary attention enhances contrast appearance. *Psychol*
Sci. ;20(3):354-362 (2009). doi:10.1111/j.1467-9280.2009.02300.x

30. Anton-Erxleben, K. & Carrasco, M. Attentional enhancement of spatial resolution: linking
behavioural and neurophysiological evidence. Nat Rev Neurosci 14, 188-200 (2013).
doi:10.1038/nrn3443.

31. Michael Jigo, Marisa Carrasco; Differential impact of exogenous and endogenous attention on
the contrast sensitivity function across eccentricity. *Journal of Vision* 20(6):11 (2020).
doi: https://doi.org/10.1167/jov.20.6.11.

32. Watson, A. B. & Ahumada, A. J. Modeling acuity for optotypes varying in complexity. J Vis 12,
doi:10.1167/12.10.19 (2012).

33. Carrasco, M., Penpeci-Talgar, C. & Eckstein, M. Spatial covert attention increases contrast
sensitivity across the CSF: support for signal enhancement. Vision Research 40, 1203-1215, doi:
10.1016/S0042-6989(00)00024-9 (2000).

34. Applegate, R. A., Ballentine, C., Gross, H., Sarver, E. J., Sarver, C. A. Visual acuity as a
function of zernike mode and level of root mean square error. Optom. & Vis. Sc. 80, 97–105
(2003).

35. Thibos, L.N., Hong, X., Bradley, A., Applegate, R.A. Accuracy and precision of objective
refraction from wavefront aberrations. J. Vis., 4, 329-351 (2004).

36. Marsack, J., Thibos, L.N., Applegate, R.A. Metrics of optical quality derived from wave
aberrations predict visual performance. J. Vis., 4, 322-328 (2004).

37. Villegas, E.A., Alcón, E., Artal, P. Optical Quality of the Eye in Subjects with Normal and
Excellent Visual Acuity. Invest. Ophthalmol. Vis. Sci. 49:4688-4696 (2008). doi:
10.1167/iovs.08-2316.

38. Ravikumar A., Sarver E. J., Applegate R. A. Change in visual acuity is highly correlated with
change in six image quality metrics independent of wavefront error and/or pupil diameter. J. Vis.
12, 1–13 (2012).

39. Ravikumar, A., Marsack, J. D., Bedell, H. E., Shi, Y., & Applegate, R. A. Change in visual
acuity is well correlated with change in image-quality metrics for both normal and keratoconic
wavefront errors. J. Vis. 13, 1–16 (2013).

40. Liang, J. Z., Williams, D. R. & Miller, D. T. Supernormal vision and high-resolution retinal imaging
through adaptive optics. J Opt Soc Am A 14, 2884-2892, doi:Doi 10.1364/Josaa.14.002884 (1997).

41. Artal, P. (2014). Optics of the eye and its impact in vision: A tutorial. Advances in Optics and
Photonics, 6, 340–367.

42. Burton GJ, Haig ND. Effects of the Seidel aberrations on visual target discrimination. *J Opt Soc*
Am A. 1984;1(4):373-385. doi:10.1364/josaa.1.000373

43. Chan, C., Smith, G. & Jacobs, R. J. Simulating refractive errors: source and observer methods.
Am J Optom Physiol Opt 62, 207-216 (1985).

44. Grier, R. A. et al. The vigilance decrement reflects limitations in effortful attention, not

mindlessness. Hum Factors 45, 349-359(2003). doi:10.1518/hfes.45.3.349.27253.

45. Warm, J. S., Parasuraman, R. & Matthews, G. Vigilance requires hard mental work and is stressful.

Hum Factors 50, 433-441   (2008). doi:10.1518/001872008X312152.

46. Szalma, J. L. et al. Effects of sensory modality and task duration on performance, workload, and

stress in sustained attention. Hum Factors 46, 219-233 (2004). doi:10.1518/hfes.46.2.219.37334

47. Brefczynski-Lewis, J. A., Datta, R., Lewis, J. W. & DeYoe, E. A. The topography of visuospatial

attention as revealed by a novel visual field mapping technique. J Cogn Neurosci 21, 1447-1460,

(2009). doi:10.1162/jocn.2009.21005

48. Nowakowski M, Sheehan M, Neal D, Goncharov AV. Investigation of the isoplanatic patch and

wavefront aberration along the pupillary axis compared to the line of sight in the eye. *Biomed Opt*

Express. 3(2):240-258 (2012) . doi:10.1364/BOE.3.000240

49. J. Thaung, P. Knutsson, Z. Popovic, M. Owner-Petersen, Dual-conjugate adaptive optics for wide-

field high-resolution retinal imaging, Opt. Express 17 (6) 4454-4467 (2009).

50. Zheleznyak L, Barbot A, Ghosh A, Yoon G. Optical and neural anisotropy in peripheral vision. *J*

Vis;16(5):1 (2016) . doi:10.1167/16.5.1

51. Ghosh A, Zheleznyak L, Barbot A, Jung H, Yoon G. Neural adaptation to peripheral blur in

myopes and emmetropes. *Vision Res*.132:69-77 (2017) . doi:10.1016/j.visres.2016.09.017

52. De Lestrange-Anginieur E, Kee C-s Testing impacts of global blur profiles using a multiscale

simulator. Heliyon 6 e04153 (2020).

53. De Lestrange-Anginieur E, Kee C-s Investigation of the impact of blur under mobile attentional

orientation using a vision simulator. PLoS ONE 15(6): e0234380 (2020).

https://doi.org/10.1371/journal.pone.0234380

54. Thibos, L. N., Applegate, R. A., Schwiegerling, J. T., Webb, R. & Members, V. S. T. Standards for

reporting the optical aberrations of eyes. J Refract Surg 18, S652-S660 (2002).

55. Thibos, L. N., Hong, X., Bradley, A., Cheng, X. "Statistical variation of aberration structure and

image quality in a normal population of healthy eyes," J Opt Soc Am A 19, 2329-2348 (2002).

56.

559
Figure 1

a   b

c   d

Figure 2

a   b

c   d

e

Figure 3

Figure 4

a   b

Figure 5

Figure 6

a   b

Figure 7

Figure Legends

Figure 1. Visual acuity in the near-peripheral retina. In this example, the subject was required

to respond to the Landolt C located in the SE quadrant. Visual acuity was tested at two locations

(NW and SE) under various blur levels (blurred and clear) and cueing conditions (cued, uncued,

and neutral). NW, north-west; SE, south-east; RT, response time.

Figure 2. Spatial cueing. (**a**) Mean visual acuity (VA) for the three cue conditions. (**b**) Changes

in VA at the attended and unattended locations using the neutral condition as a baseline

($VA_{neutral}$–VA_{cued} and $VA_{neutral}$–VA_{uncued}, respectively). (**c**) Mean response time (RT) for the three

cue conditions. (**d**) Changes in RT at the attended and unattended locations using the neutral

condition as a baseline ($RT_{neutral}$–RT_{cued} and $RT_{neutral}$–RT_{uncued}, respectively). In this and

following figures, error bars represent one standard error of the mean.

Figure 3. Spatial cueing and visual blur. (**a**) visual acuity (VA) for the three cue conditions and

the two blurred conditions. (**b**) Changes in VA at the attended and unattended locations for the

two blurred conditions using the neutral condition as a baseline ($VA_{neutral}$–VA_{cued} and $VA_{neutral}$–

VA_{uncued}, respectively). (**c**) Response time (RT) for the three cue conditions and the two blurred

conditions. (**d**) Changes in RT at the attended and unattended locations for the two blurred

conditions using the neutral condition as a baseline ($RT_{neutral}$–RT_{cued} and $RT_{neutral}$–RT_{uncued},

respectively). (**e**) Effect of visual blur on visual acuity under the three cue conditions. Acuity

difference between blurred and clear targets. When attention becomes diverted from the target

location, the difference of resolution between blurred and clear images is mitigated.

Figure 4. Interaction between attention and blur. Mean VA for different combinations of

attention and blur levels. The deployment of attention resulted in a defocused system (blur

image) that performs better than, or the same as, a focused system (clear image). This indicates

that different combinations of the degree of a person's attention and his/her optical correction

are possible for the visual system to achieve a given acuity resolution.

Figure 5. Impact of visual blur on the beneficial effect of attention. (**a**) Effect of attentional

modulation on VA. Ratio in VA between neutral and focused conditions ($VA_{neutral}/VA_{focused}$) at

cued and uncued locations (i.e., cued and uncued conditions, respectively). Visual blur

decreased both beneficial and cost effects of cued attentional orienting, which resulted in a

narrower range of resolvable stimulus by the focus of attention. (**b**) Acuity dynamic range

(VA_{cued}/VA_{uncued}) associated with the modulation occurred when attention shifted from uncued

to cued conditions.

Figure 6. Changes of optical and neural filter. Schematic diagram showing how the gain of a

spatially localized attentional filter (ΔNTF) could vary in response to the variation of the

modulation transfer function (*MTF*) of the eye for various level of defocus. The rate of the dash

line indicates the level of blur associated with the optical and attentional filter. As the magnitude

of blur increases, the maximum effective width of the attentional filter is reduced in the highest

spatial frequency of the image due to the shrinkage of the area under the modulation transfer

function.

Figure 7. Impact of attention on refractive gain. (**a**) Ratio in VA between blur and clear images

(R_x=VA$_{\text{blur}}$/VA$_{\text{clear}}$) as a function of cueing conditions. As attention diminished, the beneficial

effect of blur correction decreased. (**b**) Refractive gain change ($R_x$cueing/R_x neutral) from the

baseline condition (i.e., neutral attention) associated with attention modulation. Attention

focusing resulted in a large increase in the expected gain of blur correction with respect to

neutral condition, but only slight reduction accompanied attentional diversion.