Transport Mechanism of Class-3 P4 ATPase Lipid Flippases

Lin Bai[1,#,*], Qinglong You[1,#], Bhawik K. Jain[2,#], Amanda Kovach[1], Todd R. Graham[2], and Huilin Li[1,*]

[1] Structural Biology Program, Van Andel Institute, Grand Rapids, Michigan, USA

[2] Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[#] These authors contributed equally to the work.

Correspondence should be addressed to L.B. (Lin.Bai@vai.org) or H.L. (Huilin.Li@vai.org).

ABSTRACT

The P4 ATPases are a large family of membrane-embedded enzymes that use ATP hydrolysis to transport large lipid substrates across lipid bilayers. P4 ATPases differ in their cellular membrane location and their substrates. The structures of the endosome- and Golgi-localized class-1 phosphatidylserine flippases—such as the yeast Drs2 and human ATP8A1—have recently been reported, revealing a substrate binding site on the lumenal side and several transport states. However, a substrate binding site on the cytosolic side has not been found, and the transport mechanisms of P4 ATPases in other classes are still unknown. Here we report a systematic structural and functional study on two plasma-membrane localized, class-3 P4 ATPases that have broader substrate specificity, the *S. cerevisiae* Dnf1–Lem3 and Dnf2–Lem3 complexes. We have captured substrate lipids on both the exoplasmic and cytosolic sides, and we found that these two enzymes have very similar structures, consistent with their high sequence identity and redundant function. Unexpectedly, Lem3 contributes to substrate binding near the cytosolic surface. We found that the conformational transitions through the substrate transport cycle of these two class-3 enzymes match those of the class-1 enzymes, suggesting a conserved lipid-flipping mechanism among all classes of the P4 ATPases. Our study also revealed a helix-turn-helix insertion in the cytosolic P domain

34   that is unique to the class-3 enzymes and plays a crucial role in their function.

35   Therefore, the P4 ATPases may have retained an overall transport mechanism while

36   evolving distinct features for cellular membrane localization, regulatory mechanisms,

37   and transporting different lipid substrates.

INTRODUCTION

The large group of integral membrane proteins that mediate the transport of substrates across biological membranes by the formation of ATP-dependent phosphorylated intermediates are called P-type ATPases (Axelsen and Palmgren, 1998). They are widely expressed in prokaryotes, archaea, and eukaryotes, and they play essential roles in many cellular processes. P-type ATPases are phylogenetically divided into five subclasses (P1-P5), each have unique sequence motifs and transport different substrates (Bublitz et al., 2011; Palmgren and Axelsen, 1998). Most P4 ATPases function as a heterodimeric complex: a catalytic α-subunit containing 10 transmembrane α-helices (TMHs) and a regulatory β-subunit of the Cdc50 (cell division cycle 50) protein family containing two TMHs. The α-subunit of all P-type ATPases has a conserved architecture consisting of a transmembrane domain (TMD), an actuator domain (A-domain), a nucleotide-binding domain (N-domain), and a phosphorylation domain (P-domain) (Andersen et al., 2016; van der Mark et al., 2013). The "Post–Albers" model proposed that the substrate transport of P-type ATPases goes through a cyclic transition of E1–E1P–E2P–E2 states (Post et al., 1972). This model is supported by the structure and function studies of the most well-understood P2 ATPases (Dyla et al., 2019; Toyoshima and Cornelius, 2013).

P4 ATPases are involved in phospholipid translocation: they flip lipids from the extracellular side of the plasma membrane or from the lumenal side of internal organelles to the cytosolic side (Tang et al., 1996; Zhou and Graham, 2009). They are important for creating and maintaining lipid compositional asymmetry in cell membranes. Because a phospholipid substrate is large, the transport mechanism of P4 ATPases is substantially different from that of the cation-transporting P-type ATPases, such as the sarcoplasmic reticulum Ca^{2+}-ATPase (SERCA), the Na^+/K^+-ATPase, and the H^+-ATPase. There are five P4-ATPases belonging to four classes in *Saccharomyces cerevisiae*: Drs2 in class 1, Neo1 in class 2, Dnf1 and Dnf2 in class 3, and Dnf3 in class 4 (van der Mark et al., 2013). Humans have 14 P4-ATPases (Andersen et al., 2016; Best et al., 2019; van der Mark et al., 2013). The P4 ATPases

have different tissue-specific expression, subcellular localization, and substrate

specificity. For example, yeast Drs2 transports phosphatidylserine (PS) and

phosphatidylethanolamine (PE) in the trans-Golgi network (TGN) and endosomes

(Natarajan et al., 2004; Zhou and Graham, 2009), while the yeast Dnf1 and Dnf2 flip

PE, PC, and glucosylceramide (GlcCer) in the plasma membrane (Best et al., 2019;

Pomorski et al., 2003; Roland et al., 2020; Roland et al., 2019; Stevens et al., 2008).

The first structural and mechanistic insights into the P4 ATPases came from recent

cryo-EM structures of the *S. cerevisiae* class-1 flippase Drs2–Cdc50 in its auto-inhibited

and active E2P states (Bai et al., 2019a; Timcenko et al., 2019), and of its human

homolog, ATP8A1–CDC50a, in six transport states (E1, E1-ATP, E1P-ADP, E1P, E2P,

and E2Pi-PL) (Hiraizumi et al., 2019). Drs2 and ATP8A1 belong to the same subfamily

of P4 ATPases, and both transport the same substrates, PS and PE (Paterson et al.,

2006; van der Mark et al., 2013; Zhou and Graham, 2009). The structural studies

revealed an allosteric domain movement during lipid translocation and identified a lipid

binding site at the exoplasmic side of membrane of this subfamily. Prior mutational

studies had identified amino acids in this exoplasmic lipid binding site that are important

for substrate recognition and had implicated a second substrate binding site on the

cytosolic side of the membrane (Baldridge and Graham, 2012, 2013; Baldridge et al.,

2013; Roland et al., 2019).

However, many important questions remain regarding the lipid-flipping mechanism(s) of

the P4 ATPases. Chief among them are how substrate specificity is achieved by

different P4 ATPases and where the lipid binds on the cytoplasmic side of the

membrane. Addressing these questions requires a detailed structural and functional

comparison among P4 ATPases that transport different substrates. Toward this goal,

we have now studied the two *S. cerevisiae* class-3 P4 ATPases, the Dnf1–Lem3 and

Dnf2–Lem3 complexes. Dnf1 and Dnf2 share the same β-subunit, Lem3, and both

transport PE, PC, and glucosylceramide. Their human homologues, ATP10A and

ATP10D, are implicated in diabetes, obesity, myocardial infarction, atherosclerosis, and

Parkinson's disease (Folmer et al., 2009; Martin et al., 2020; van der Mark et al., 2013).

Here we report the cryo-EM structures of Dnf1–Lem3 in three different states (E1, E1P-

ADP, and E2P) and Dnf2–Lem3 in five different states (E1, E1-ATP, E1P-ADP, E2P,

and E2P-transition). We further performed structure-guided mutagenesis and *in vivo*

substrate transport activity assays. By comparing these class-3 flippase structures with

the class-1 structures previously reported, we found new phospholipid binding sites at

the cytoplasmic side and unique structural features during the ATP-dependent transport

cycle in Dnf1–Lem3 and Dnf2–Lem3. Our comprehensive structural and functional

study improves the mechanistic understanding of the essential P4 ATPases.

RESULTS AND DISCUSSION

1. Overall Dnf1 structure

Endogenous Dnf1–Lem3 and Dnf2–Lem3 complexes were purified from baker's yeast

using an anti-FLAG affinity column followed by size-exclusion chromatography

(**Supplemental Figure 1, online Methods**). We used the detergent dodecyl maltoside

(DDM) to solubilize these flippases and lauryl maltose neopentyl glycol (LMNG) and

cholesteryl hydrogen succinate (CHS) to stabilize the membrane protein complexes. We

performed single-particle cryo-EM on purified samples in apo form (E1 state) and on

samples incubated with AMPPCP (E1-ATP state), ALF4$^-$-ADP (E1P-ADP and E2P-

transition states), or BeF3$^-$ (E2P state). The substrate transport states were assigned

based on the established nomenclature in the P-type ATPase field (Bai et al., 2019a;

Hiraizumi et al., 2019; Timcenko et al., 2019). We obtained cryo-EM 3D maps of the *S.*

cerevisiae Dnf1–Lem3 in the E1, E1P-ADP, and E2P states and Dnf2–Lem3 in the E1,

E1-ATP, E1P-ADP, E2P-transition, and E2P states, at resolutions of 2.8 Å to 4.0 Å

(**Supplemental Figures 2-7, Supplemental Table 1**). The high-resolution 3D map at

2.8 Å of the Dnf1–Lem3 in the E2P state enabled us to build the first atomic model of

Dnf1–Lem3 (**Fig. 1, Supplemental Figure 2**). Except for a few short and disordered

loops, almost the entire complex is ordered and well resolved in this model.

Interestingly, many extra densities were observed in the transmembrane region of Dnf1

in the E2P state (**Fig. 1A**). Some clearly belonged to the detergent CHS that was added

during purification, while others located in the substrate transport path resembled

phospholipid, and they were so assigned, as will be described below.

We then used the 2.8-Å structure to help build atomic models for the seven remaining

3D maps of Dnf1p–Lem3p and Dnf2p–Lem3p, which are at slightly lower resolutions. All

these models were refined to good statistics (**Supplemental Table 1**) and fit well with

the 3D maps (**Supplemental Figure 3**). The modeled nucleotides and lipids all had

clear densities. By comparing the 3D maps of Dnf1–Lem3 and Dnf2–Lem3 in the E1,

E1P-ADP, and E2P states, which were the three states observed for both flippases, we

found that the structures were superimposable in all three states (**Supplemental Figure**

8). This comparison suggests that Dnf1 and Dnf2 have highly similar structures and go

through virtually the same functional states during their respective substrate transport

cycles. This finding was not surprising, because these two class-3 flippases share a

high sequence identity of about 69%, localize to the same membrane, transport the

same substrates, and appear to have redundant functions in yeast (Pomorski et al.,

2003). Because of their apparently interchangeable structural feature, we will discuss

these two flippases together in the following sections.

We first examined the Dnf1 structure in the E2P state because of the high resolution

and completeness of the structure. As expected of any P4 ATPase, Dnf1 has a 10-TMH

TMD (TMH1-10); a cytosolic A domain inserted between TMH2 and TMH3; and a

cytosolic P domain and a cytosolic N domain, both inserted between TMH4 and TMH5

(**Fig. 1**). The amino terminal peptide preceding TMH1 (N-tail) of Dnf1 is 204 residues

long and is largely disordered, except for the last 39-residue segment (Glu-166 to Leu-

204), which is stabilized by binding to the regulatory A domain. The carboxyl terminal

peptide (C-tail) of Dnf1 following the 10th TMH is 157 residues long (Pro-1405 to Asn-

1571) and is also largely disordered except for the TMD-proximal 35-residue segment

(Pro-1405 to Lys-1439).

Interestingly, this ordered Dnf1 C-tail segment contains a short α-helix, the equivalent of

which was reported to function as a switch that converts the Drs2 flippase from its auto-

inhibitory state to the active state in a unique PI4P-binding-dependent manner (Bai et

al., 2019a). However, the Dnf1 helix is stabilized by the C-tail of Lem3 in a conformation

corresponding to the active state of Drs2, suggesting that Dnf1 and Dnf2 do not require

binding to a phosphoinositide at this site for activation. The Dnf1 C-tail interacts

extensively with the cytoplasmic domains of Dnf1 in regulating the substrate transport

cycle, as described below. Dnf1 and Dnf2 activity is regulated by two flippase kinases

Fpk1 and Fpk2. Mutations in Fpk1 and Fpk2 do not affect the subcellular location of

Dnf1 and Dnf2, but these enzymes have almost no activity in a *fpk1,2Δ* strain (Nakano

et al., 2008; Roelants et al., 2010). We found that all 17 known phosphorylation sites

(P32660 in the UniProt Database (UniProt, 2019)) are located in disordered loop

regions in the Dnf1 structure: the N-tail, a loop between Thr-290 and Ala-378, and the

C-tail. These observations suggest that the class-3 flippases are autoinhibited and

phosphorylation, not PI4P binding as in the class-1 flippases, relieves the inhibition.

2. Lem3 has unique and ordered N- and C-terminal regions

Similar to the yeast Cdc50 and human CDC50a, Lem3 contains two TMHs flanking a

large lumenal domain. These TMHs are packed against the last TMH of Dnf1, TMH10

(**Fig. 1**). The Lem3 lumenal domain contains two disulfide bonds (Cys-110 with Cys-159

and Cys-216 with Cys-231) and four N-glycans at Asn-240, Asn-256, Asn-298, and Asn-

332. To our surprise, both the N-terminal and the C-terminal regions of Lem3 are stable,

in contrast to the yeast Cdc50 and human CDC50a, in which these regions are

disordered (**Fig. 1A-B**). The N-terminal peptide preceding Asn-50 folds into a two-turn

α-helix and binds the Dnf1 A domain (**Supplemental Figure 9A-B**). Due to a partial

flexibility, the exact sequence of the Lem3 N-tail α-helix was not resolved. However, we

found its binding interface in the A domain is highly positively charged, featuring Lys-

392, Lys-395, Arg-408, Arg-457, Lys-460, and Lys-516 (**Supplemental Figure 9C**), and

the Lem3 N-tail segment between Glu-28 to Glu-42 is highly negative, containing 11

Asp and Glu residues (**Supplemental Figure 9D**). Therefore, the short N-tail α-helix of

Lem3p is most likely from this negatively charged region. Because the A domain moves

significantly during the substrate transport cycle of P4 ATPase, this unique, A domain-

interacting, and highly negative Lem3 N-tail α-helix likely regulates lipid transport activity

of Dnf1 and Dnf2, as described below.

3. Phospholipids bind at the exoplasmic and cytoplasmic sites of Dnf1

Although the exact substrates a P4 ATPase transport differ, the substrate translocation

path appears to have been conserved in all P4 ATPase studied so far (Andersen et al.,

2016; Huang et al., 2020). Like the Drs2 and ATP8A1 flippases, we found that the Dnf1

substrate translocation path is composed of TMH1-4 and TMH6 (**Fig. 2a-d,**

Supplemental Figure 10). This substrate path is consistent with biochemical studies

demonstrating that TMH1-6 are responsible for phospholipid transport, with TM7-10

having a supporting role (Baldridge and Graham, 2013; Vestergaard et al., 2014). In the

E2P-state Dnf1 3D map, we observed three phospholipid-like densities in the substrate

transport path, and accordingly, we modeled three phospholipid molecules (PL1-PL3) in

that map (**Figs. 1A, 2A**). These phospholipids stack against Dnf1 TMH2, TMH4, and

TMH6. Because no lipids were added during purification, these phospholipids are

potentially endogenous substrates that were co-purified with the Dnf1–Lem3 complex.

This is not unusual, because previous studies have shown that substrate lipids bound in

the substrate pockets can be co-purified with the enzyme complexes (Bai et al., 2019b;

Bloch et al., 2020; Hiraizumi et al., 2019). Interestingly, the three phospholipids are out

of register with respect to the lipid bilayer by approximately 10 Å: PL1 is moved down

about 10 Å into the middle of the bilayer relative to lipids in the outer leaflet, and PL2

and PL3 are moved 10 Å into the cytosolic space relative to lipids in the inner leaflet of

the membrane.

4. Functional studies reveal that PL1 and PL2 are in substrate binding sites

Based on sequence conservation of the three phospholipid binding pockets, we expect

that PL1 and PL2—but not PL3—are located in the substrate path and are likely

physiologically relevant. To test this hypothesis, we carried out systematic, structure-

based mutagenesis and functional assays (**Fig. 3A-E, Supplemental Table 3**). PL1 is

bound in an open cleft of Dnf1 on the exoplasmic side of the plasma membrane, and

the phosphate group of PL1 interacts with several conserved residues, including Gln-

610, Ser-611, and Asn-1226 (**Fig. 2A-B**). We expressed the Q610A, S611A,

609YQS611-FSN, and N1226A Dnf1 mutants on a *dnf1Δ dnf2Δ* cell background and

measured their PC, PE, and GlcCer uptake abilities. We found that Q610A and the triple

mutation 609YQS611-FSN eliminated the GlcCer and PE, but not the PC, transport

activity (**Fig. 3A**). This observation was consistent with a previous finding in which

mutations of the conserved tripeptide motif 654YQS656 of Dnf2—the equivalent of

609YQS611 in Dnf1—caused substantial and specific loss of GlcCer transport (Roland

et al., 2020; Roland et al., 2019).

It is unclear why the S611A mutation led to increased transport activity for all three

substrates (**Fig. 3A**). In contrast, the N1226A mutation completely halted transport of all

three substrates, even though the mutant protein localized to the plasma membrane

comparably to WT Dnf1 (**Fig. 3D**). The ability of Dnf1-N1226A to exit the ER indicated it

was folded properly and was able to interact with Lem3, but consistent with the loss of

transport activity, Dnf1-N1226A failed to support the viability of a flippase-deficient yeast

strain, whereas Dnf1 variants that retained transport activity fully supported viability

(**Supplemental Figure 11**). The GA motif (230-231) in TMH1B is at the membrane-

exoplasmic interface and is next to the PL1 binding site (**Fig. 2D**). A recent study has

shown that the GA motif is crucial for GlcCer transport by Dnf2 but has little influence on

PC and PE transport (Roland et al., 2019). Furthermore, we found that the PL1 binding

site superimposed with the substrate binding site of ATP8A1, based on structural

alignment (**Supplemental Figure 11**). Taken together, we conclude that the PL1

binding site is the entrance of the substrate transporting path in Dnf1 and Dnf2.

Unlike PL1, both PL2 and PL3 bind Dnf1 on the cytosolic side of the lipid bilayer. The

head group of PL2 is in a positively charged groove and is stabilized by several

conserved residues, including Arg-264, Tyr-633, Thr-648, and Trp-652 of Dnf1, and Arg-

51 of Lem3. Because PL2 is in the putative substrate transport path, we suggest that

PL2 is at the substrate exit site of Dnf1. We generated a series of point mutations in

Dnf1, including R264A, Y633A, T648A, W652A, and W652S. We expressed these

mutations in *dnf1Δ dnf2Δ*, *lem3Δ*, or *lem3Δ dnf2Δ* cells and found that W652A and

W652S significantly disrupted transport of all substrates; that R264A enhanced PC, PE,

and GlcCer transport; and that Y633A and T648A had no significant effects on the

transport activities (**Fig. 3A**). The elimination of Dnf1 transport activity by the W652

mutations was consistent with this residue's strategic location between the PL2-binding

substrate exit site and the PL1-binding substrate entry site. However, this mutation

caused the retention of Dnf1 in the ER, possibly because it disrupts folding of the

protein (**Fig. 3E**). The increased activity of Dnf1 R264A suggests that this charged

residue may trap the substrates and slow their release into the cytosolic leaflet. This

observation supports our hypothesis that PL2 is a substrate lipid. Because the PL3 site

is located right next to the PL2 site, and the PL3 site is composed of nonconserved

residues (except for F631), we suggest that the PL3 site plays an accessory role,

perhaps by facilitating substrate release from the PL2 site.

5. Lem3 unexpectedly contributes to substrate binding on the cytosolic side

Because the Lem3 Arg-51 directly interacts with PL2, we generated a Lem3 R51A

mutant and found that the substitution changed the substrate specificity of both Dnf1

and Dnf2, with a 1.5- to 2-fold increase in GlcCer transport activity of Dnf2–Lem3 and

Dnf1–Lem3, but no change in PC or PE activity for either complex (**Fig. 3B-C**). Because

there is no phosphate in GlcCer, it is possible that R51A increases ability of GlcCer to

compete with endogenous lipids for this site. As expected, Lem3-R51A supported

trafficking of the complex out of the ER (**Fig. 3E**), and this result supports the role of

Lem3 R51 in controlling substrate specificity and further supports the assignment of PL2

as a substrate lipid.

To examine the functional importance of the N-terminal region of Lem3, we generated a

Lem3 variant missing the N-terminal peptide (Lem3[Δ2-49]) and performed a lipid

uptake assay. We found that with Lem3(Δ2-49), both Dnf1 and Dnf2 had significantly

reduced PC, PE, and GlcCer uptake ability (**Fig. 3B-C**). We next examined the

functional importance of the ordered C-terminal region of Lem3 (Gly-400 to Lys-414)

that stabilizes Dnf1 in an active state. In the lipid uptake assays with Lem3(Δ400-414),

both Dnf1 and Dnf2 had substantially decreased activity (**Fig. 3B-C**). We then labeled

Dnf1 with GFP and found that Lem3(Δ2-49) and Lem3(Δ400-414) abolished Lem3's

ability to escort Dnf1 out of the ER (**Fig. 3E**). Thus, neither Dnf1 or Dnf2 was able to

reach to its functional destination in the plasma membrane with a N- or C-terminus

truncated Lem3. Thus, these results support an unexpected role of Lem3 in substrate

binding and specificity. No such role was found in the β-subunit (Cdc50 or CDC50) of

the class-1 flippases Drs2 and ATP8A1 (Bai et al., 2019a; Hiraizumi et al., 2019;

Timcenko et al., 2019).

During preparation of this manuscript, the cryo-EM structure of Dnf1–Cdc50 of a

thermophilic fungus, *Chaetomium thermophilum*, was reported in the E1-ATP and E2P

states at 3.4-3.5 Å resolution (He et al., 2020). That Dnf1 structure contained two lipids,

but no evidence was provided that they were substrate lipids. Many membrane proteins,

including SERCA and Na^+/K^+ ATPases (Kanai et al., 2013 ; Toyoshima et al., 2013),

have specifically bound lipid molecules in their structures that are presumably not

substrate lipids. Because we have now functionally defined that the PL1 and PL2 in our

S. cerevisiae Dnf1–Lem3 and Dnf2–Lem3 structures are located at the physiological

substrate entry and exit, the observation that the two lipids in *C. thermophilum* Dnf1 are

equivalent to PL1 and PL2 in our structure indicates that in both species, the class-3

flippases share a similar substrate path.

We conclude that the substrate transport pathway is primarily located between TMH2

and TMH4 (**Fig. 2E**). This pathway is nearly perpendicular to the membrane plane and

is open to the lipid bilayer. Interestingly, this pathway is hydrophilic at both ends, the

entry and the exit sites, but is highly hydrophobic in the middle gating region (**Fig. 2B-**

E). In the Dnf1–Lem3 structure, the exoplasmic substrate entry site is lined by Gln-610,

Ser-611, and Asn-1226; the cytosolic substrate exit site is surrounded by Dnf1 residues

Gln-214, Asn-217, Asn-220, Thr-254, Lys-257, Asp-258, Glu-261, Arg-264, Arg-265,

Glu-622, and Thr-626, along with Lem3 Arg-51, and the middle hydrophobic region is

composed of the conserved Phe-223, Leu-247, Ile-250, Ile-615, and Ile-619 (**Fig. 2B-E**).

The physiochemical property of the transport pathway is compatible with the so-called

"credit card" mechanism, in which the hydrophilic or charged headgroups of the lipid

substrates simply slide through the groove while allowing the substrate acyl chains to

flip in the favorable membrane environment. It is likely that during transport, the lipid

head group inserts into and make direct interactions with the hydrophilic residues

between TMH2 and TMH4.

6. ATP dependent transport cycle

By cryo-EM analyses of samples incubated with or without the ATP analogs AMPPCP

and ADP-AlF$_4^-$ or the phosphate analogs AlF$_4^-$ and BeF$_3^-$, we obtained a total of eight

structures of the class-3 flippases. There were Dnf1–Lem3 in the E1 (apo), E1P-ADP

(ALF$_4^-$-ADP), and E2P (BeF$_3^-$) states; and Dnf2–Lem3 in the E1 (apo), E1-ATP

(AMPPCP), E1P-ADP (ADP-AlF$_4^-$), E2P-transition (AlF$_4^-$), and E2P (BeF$_3^-$) states.

Comparing these structures revealed how ATP binding and hydrolysis, ADP release,

and enzyme phosphorylation produce conformational changes in Dnf1 and Dnf2 to drive

the Post–Albers substrate transport cycle (**Fig. 4A**).

We found that a major part of the TMD region (that is, TMH3-10) and the P domain of

Dnf1 or Dnf2, and the most of Lem3 molecule except for its N-tail, were well aligned,

suggesting that these regions are largely stationary during the substrate transport cycle

(**Fig. 4A, Supplemental Figure 12A-B**). However, TMH1-2 and the A and N domains of

Dnf1 and Dnf2 moved significantly; these movements must be coupled with substrate

transport. In the E1 state, the A domain moves towards the membrane bilayer and

becomes separated from the N domain (**Fig. 4A**). Due to the lack of interaction, the A

and N domains are flexible and are visible in the 3D maps only at a low display

threshold (**Supplemental Figure 13**).

The binding of ATP in the E1-ATP state triggers the N domain to move towards and

interact with the A domain, thereby stabilizing both domains. However, the A domain

must be only partially stabilized and retain some flexibility, because its density is still

weak. With the ensuing phosphoryl transfer, the flippases transition from the E1-ATP

state to the E1P-ADP state. This transition does not trigger large-scale conformational

changes, because the structures of Dnf2–Lem3 in the E1-ATP and E1P-ADP are almost

the same except for their respective bound nucleotide (**Supplemental Figure 12**). In

the E1, E1-ATP, and E1P-ADP states, TMH1-2 have a small degree of flexibility, but

they do not move much (**Fig. 4B**). Transitioning from the E1P-ADP state to the E2P

state, the release of ADP triggers the N domain to rotate 40° away from the A domain

by pivoting on the N domain and P domain interface. The conserved DGET motif of the

A domain now occupies the phosphorylation site and stabilizes the A domain. During

this transition, TMH1 and TMH2 tilt to enlarge the exoplasmic substrate binding site for

the admission of PL1 and at the same time to narrow the cytosolic substrate site,

thereby stabilizing the bound PL2 (**Fig. 4B**).

During the subsequent phosphate (Pi) transition, when the enzyme changes from the

E2P state to the E2P-transition state, the enzyme structure largely holds still; the 3D

maps of Dnf2–Lem3 in these two states are nearly identical (**Supplemental Figure

12B**). Finally, the Dnf1 and Dnf2 switch back to the E1 state by auto-dephosphorylation.

During this switch, the A and N domains become flexible again, and the TMH1-2 tilt

back to narrow and close the exoplasmic substrate binding site and open the cytosolic

substrate binding site. It is during this last transition that PL1 in the exoplasmic site is

forced to flip and enter the cytosolic substrate site, replacing the resident PL2 there and

leading to the release of PL2 from the flippases.

7. Transport mechanisms of the class-3 and class-1 P4 ATPases are similar

As described above, the major domain rearrangements in Dnf1 and Dnf2 are triggered

by nucleotide binding (E1 to E1-ATP); nucleotide hydrolysis, enzyme phosphorylation,

nucleotide release (E1P-ADP to E2P); and enzyme dephosphorylation (E2P to E1). The

energy gain from ATP during these transitions is transduced from the phosphorylation

site in the P domain to the A domain, which in turn causes the TMH1-2 to tilt and drive

the flipping of the phospholipids. TMH3-10 of Dnf1 or Dnf2 and TMH1-2 of Lem3 largely

remain stationary throughout the transport cycle. Combining the finding of two substrate

binding sites and the transport cycle, we propose that the lipid head group transports

through the groove between TMH2 and TMH4, while lipid tails move accordingly on the

hydrophobic surface of the TM2 and TM4. This is essentially the same as the previously

described "credit card model" (Andersen et al., 2016; Roland and Graham, 2016).

The previous cryo-EM study of the human ATP8A1–CDC50a flippase captured multiple

transport states (Hiraizumi et al., 2019). Aligning our structures in this paper with those

of ATP8A1–CDC50a reveal that the corresponding E1, E1-ATP, E1P-ADP, E1P, and

E2P states are remarkably similar in both domain architecture and substrate transport

path (**Fig. 5**). This result was unexpected, because ATP8A1 is a class-1 flippase and

Dnf1 and Dnf2 are class-3. Our observation raises the possibility that all P4 ATPases

may employ a similar substrate-transport mechanism.

Despite their overall similarity in substrate transport cycle, notable structural differences

exist among the flippases characterized so far. In class-1 flippases such as the yeast

Drs2 and human ATP8A1, the C-terminal loop is auto-inhibitory; this loop, which

contains the conserved GF/YAFS motif, extends over and wraps around the

cytoplasmic catalytic domains to stabilize the class-1 enzymes in the E2P state. The C-

terminal loop of Dnf1 and Dnf2 doesn't contain the GF/YAFS motif and is disordered in

the E2P state. However, there is a unique helix-turn-helix motif (E692-S737) in the N

domain of Dnf1 and Dnf2 that is absent in Drs2 and ATP8A1 (**Fig. 5**). In the E1 state,

the helix-turn-helix interacts with the P domain, specifically, its Lys-702 and Arg-703

interacting with the P domain Glu-994, Asn-998, and Asn-1008. Upon ATP hydrolysis

and enzyme phosphorylation in the E1P-ADP state, this interaction is weakened

because only Arg-703 of the motif now interacts with the P domain Asn-998 and Asn-

1008. The release of ADP in the E2P state appears to shift the interaction of the Dnf1

and Dnf2 insertion from the P domain to the C-tail, with the helix-turn-helix Glu-709 and

Arg-713 forming electrostatic interactions with the C-tail Arg-1434 and Lys-1436. These

systematic changes suggest that the Dnf1 and Dnf2 insertion may play a regulatory role

in the substrate transport cycle. To examine this possibility, we deleted the motif in Dnf1

and performed an *in vivo* substrate uptake assay. The deletion strongly decreased

transport of all substrates (**Fig. 3A**). These results confirm a crucial role of the helix-

turn-helix motif and reveal a unique aspect of the class-3 flippases over the otherwise

conserved substrate transport mechanism of the P4 ATPases.

In summary, our systematic structural and biochemical studies of the class-3 Dnf1–

Lem3 and Dnf2–Lem3 complexes in five transport states provide the novel insight that

the overall substrate transport mechanism is similar to that of class-1 flippases and that

it may be conserved among all classes of the P4 ATPases. Our studies also show a

helix-turn-helix motif in the P domain that is unique to the class-3 enzymes and plays a

crucial regulatory function during the substrate transport cycle. Also unique to class-3

enzymes is the role of the beta-subunit Lem3 in substrate binding and specificity. Thus,

this work advances our mechanistic understanding of the highly conserved and

essential family of the P4 ATPases.

MATERIALS AND METHODS

Purification of the endogenous Dnf1p–Lem3p and Dnf2p–Lem3p complexes.

Using a PCR-based genomic epitope-tagging method (Funakoshi and Hochstrasser,

2009), we tagged Dnf1p or Dnf2p with a C-terminal triple-FLAG in the yeast strain

BY4741 (Horizon Discovery). Cells were grown in YPD medium for about 20 h before

harvest. Cells were resuspended in lysis buffer (20 mM Tris-HCl, pH 7.4, 0.2 M sorbitol,

50 mM potassium acetate, 2 mM EDTA, and 1 mM phenylmethylsulfonyl fluoride

[PMSF]) and then lysed using a French press at 15,000 psi. The lysate was centrifuged

at 10,000 × g for 30 min at 4 °C. The supernatant was collected and centrifuged at

100,000 × g for 60 min at 4 °C. The membrane pellet was collected and then

resuspended in buffer A containing 10% glycerol, 20 mM Tris-HCl (pH 7.4), 1% DDM,

0.1% CHS, 0.5 M NaCl, 1 mM MgCl$_2$, 1 mM MnCl$_2$, 1 mM EDTA, and 1 mM PMSF.

After incubation for 30 min at 4 °C, the mixture was centrifuged for 30 min at 120,000 ×

g to remove insoluble membrane. The supernatant was mixed with pre-washed anti-

FLAG (M2) affinity gel at 4 °C overnight with shaking. The affinity gel was then collected

and washed three times in buffer B containing 20 mM HEPES, pH 7.4, 150 mM NaCl,

0.01% LMNG, 0.001% CHS, and 1 mM MgCl$_2$. The proteins were eluted with buffer B

containing 0.15 mg/ml 3×FLAG peptide and was further purified in a Superose 6 10/300

gel filtration column in buffer C containing 20 mM HEPES, pH 7.4, 150 mM NaCl,

0.003% LMNG, 0.0003% CHS and 1 mM MgCl$_2$. Finally, the purified proteins were

assessed by SDS-PAGE gel and concentrated for cryo-EM analysis. Approximate 20 µg

Dnf1p–Lem3p and 40 µg Dnf2p–Lem3p can be purified from 18 L of cells.

Cryo-electron microscopy. To capture different states, the purified Dnf1p–Lem3p or

Dnf2p–Lem3p was mixed with different solutions for 1 h on ice: E1-ATP, 5 mM MgCl$_2$

and 2 mM AMPPCP; E1P-ADP and E2P-transition, 5 mM MgCl$_2$, 5 mM NaF, 1 mM

AlCl$_3$, and 5 mM ADP; E2P, 5 mM MgCl$_2$, 10 mM NaF, and 2 mM BeSO$_4$. After an

incubation, 2.5 µL aliquots of purified Dnf1p–Lem3p or Dnf2p–Lem3p at a concentration

of about 1.5 mg/mL were placed on glow-discharged holey carbon grids (Quantifoil Au

R2/2, 300 mesh) and were flash-frozen in liquid ethane using a FEI Vitrobot Mark IV.

Cryo-EM data was collected automatically with SerialEM in a 300-kV FEI Titan Krios

electron microscope with defocus values from −1.0 to −2.0 µm. The microscope was

operated with a K3 direct detector at a nominal magnification of 130,000× and a pixel

size of 0.413 Å per pixel. The dose rate was 8 electrons per Å^2 per s, and the total

exposure time was 8 s.

Cryo-EM image processing. Program MotionCorr 2.0 (Zheng et al., 2017) was used for motion correction, and CTFFIND 4.1 was used for calculating contrast transfer function parameters (Mindell and Grigorieff, 2003). All the remaining steps were performed using RELION 3 (Zivanov et al., 2018). The resolution of the map was estimated by the gold-standard Fourier shell correlation at a correlation cutoff value of 0.143.

For the Dnf1–Lem3 structure in the E2P state, we collected 4764 raw movie micrographs. A total of 2,094,437 particles were picked automatically. After 2D classification, a total of 1,789,831 particles were selected and used for 3D classification. Based on the quality of the four 3D classes, 590,043 particles were retained for further 3D reconstruction, refinement, and postprocessing, resulting in a 2.8-Å average resolution 3D map.

For the Dnf1–Lem3 structure in the E1 state, we collected 4074 raw movie micrographs. A total of 1,358,323 particles were picked automatically. After 2D classification, a total of 1,089,904 particles were selected and used for 3D classification. Based on the quality of the four 3D classes, 473,761 particles were selected for further 3D reconstruction, refinement, and postprocessing, resulting in a 3.1-Å average resolution 3D map.

For the Dnf1–Lem3 structure in the E1P-ADP state, we collected 6038 raw movie micrographs. A total of 2,257,199 particles were picked automatically. After 2D classification, a total of 1,789,831 particles were selected and used for 3D classification. Based on the quality of the four 3D classes, 716,510 particles were selected for further 3D reconstruction, refinement, and postprocessing, resulting in a 3.2-Å average resolution 3D map.

For the Dnf2–Lem3 structure in the E1 state, we collected 1438 raw movie micrographs. A total of 509,555 particles were picked automatically. After 2D classification, a total of 291,438 particles were selected and used for 3D classification. Based on the quality of

the four 3D classes, 190,753 particles were retained for further 3D reconstruction,

refinement, and postprocessing, resulting in a 3.1-Å average resolution 3D map.

For the Dnf2–Lem3 structure in the E1-AMPPCP state, we collected 3665 raw movie

micrographs. A total of 844,464 particles were picked automatically. After 2D

classification, a total of 519,537 particles were selected and used for 3D classification.

Based on the quality of the four 3D classes, 315,846 particles were selected for further

3D reconstruction, refinement, and postprocessing, resulting in a 3.8-Å average

resolution 3D map.

For the Dnf2–Lem3 structure in the E1P-ADP and E2P-transition states, we collected

5683 raw movie micrographs. A total of 1,010,708 particles were picked automatically.

After 2D classification, a total of 593,683 particles were retained and used for 3D

classification. Based on the conformation of the six 3D classes, the E1P-ADP and E2P-

transition states were separated. For the E1P-ADP state, 266,218 particles were

selected for further 3D reconstruction, refinement, and postprocessing, resulting in a

4.0-Å average resolution 3D map. The Dnf1 A domain was flexible and was not

modeled. The N domain density was weak but clearly present. This domain was

modeled as a rigid body and was excluded from final atomic model refinement. For the

E2P-transition state, 127,442 particles were selected for further 3D reconstruction,

refinement, and postprocessing, resulting in a 4.0-Å average resolution 3D map.

For the Dnf2–Lem3 in the E2P state, we collected 3260 raw movie micrographs. A total

of 1,095,601 particles were picked automatically. After 2D classification, a total of

1,065,397 particles were retained and used for 3D classification. Based on the quality of

the four 3D classes, 763,661 particles were selected for further 3D reconstruction,

refinement, and postprocessing, resulting in a 3.5-Å average resolution 3D density map.

Structural modeling, refinement, and validation. We first built the model of Dnf1–

Lem3 in the E2P state at 2.8-Å resolution. We generated the initial model based on the

structure of Drs2–Cdc50 (PDB ID 6PSY) by SWISSMODEL

(https://swissmodel.expasy.org), and then manually corrected in COOT and Chimera

(Emsley et al., 2010; Pettersen et al., 2004). The complete model of Dnf1–Lem3 was

refined by real-space refinement in the PHENIX program and subsequently adjusted

manually in COOT. Using the model of Dnf1–Lem3 in E2P as reference, models of

Dnf1–Lem3 in the E1 and E1P-ADP states and of Dnf2–Lem3 in the E1, E1-ATP, E1P-

ADP, E2P transition, and E2P states were built and refined using COOT, Chimera, and

PHENIX. Finally, all models were validated using MolProbity in PHENIX (Adams et al.,

2010; Chen et al., 2010). Structural figures were prepared in Chimera (Pettersen et al.,

2004) and PyMOL (https://pymol.org/2/).

Yeast strains and plasmid construction. Strains and plasmid used in the study are

listed in Supplemental Tables I and II. All yeast culture reagents were purchased from

Sigma-Aldrich and BD Scientific. Yeast strains were grown in YPD or minimal selective

media. Yeast transformation was performed using the standard LiAC-PEG method. For

plasmid shuffling assays, 10-fold serial dilutions starting from a cell suspension with cell

density of OD_{600} = 1 were spotted on synthetic complete media (SD) and SD-5-FOA and

incubated at 30 °C for 2 d before imaging. DNA constructs and mutations were created

by Gibson assembly and quick-change mutagenesis.

Lipid uptake assay. All the lipids used in this study were purchased from Avanti Polar

Lipids. NBD-lipid uptake was performed as previously described (Roland et.,2019).

Cells were grown to the mid-log phase with a cell density of approximately OD_{600} = 1.

Cells were collected for each strain. NBD-PC, NBD-PE, and NBD-GlcCer were dried

and resuspended in 100% ethanol, then added to the designated ice-cold growth

medium at a final concentration of 2 µg/mL. Cells were incubated on ice with the pre-

chilled NBD-lipid+medium for 30 min. After the incubation, the cells were washed twice

with pre-chilled SA medium (SD medium + 2% [w/v] sorbitol + 20 mM NaN_3)

supplemented with 4% (w/v) fatty acid-free BSA. Cells were resuspended in SA with 5

µM propidium iodide (PI) and NBD-lipid uptake was immediately measured by flow

cytometry. The order in which the samples were prepared and processed was varied in

each experiment to reduce potential experimental error. Each experiment consisted of

three independent biological replicates for each strain, repeated three times (n = 9).

Fluorescence microscopy. Cells were grown in minimal synthetic dropout medium to

log phase. Cells were washed with fresh medium 3 times and resuspended in fresh SD

medium. Cells were then mounted on glass slides and observed immediately at room

temperature. Images were acquired using a DeltaVision Elite Imaging System (GE

Healthcare Life Sciences, Pittsburgh, PA) equipped with a 100× objective lens followed

by deconvolution using softWoRx software (GE Healthcare Life Science).

Supplementary Information is available in the online version of the paper.

ACKNOWLEDGEMENTS. Cryo-EM images were collected in the David Van Andel

Advanced Cryo-Electron Microscopy Suite at Van Andel Institute. We thank Gongpu

Zhao and Xing Meng for facilitating data collection and David Nadziejka for proofreading

the manuscript. This work was supported by the U.S. National Institutes of Health

(CA231466 to H.L. and GM107978 to T.R.G.) and the Van Andel Institute (to H.L.).

CONTRIBUTIONS. L.B., H.L., B.K.J. and T.R.G. conceived and designed the

experiments. L.B., Q.Y., B.K.J., and A.K. performed the experiments. L.B., T.R.G., and

H.L. analyzed the data. L.B., T.R.G, and H.L. wrote the manuscript with input from all

authors.

DATA AVAILABILITY. The cryo-EM 3D maps and the corresponding atomic models of

the Dnf1–Lem3 complex have been deposited at the EMDB database and the RCSB

PDB with the respective accession codes of EMD-xxxx and yyyy (E1), EMD-xxxx and

yyyy (E1P-ADP), EMD-xxxx and yyyy (E2P). The cryo-EM 3D maps and the

corresponding atomic models of the Dnf2–Lem3 complex have been deposited at the

EMDB database and the RCSB PDB with the respective accession codes of EMD-xxxx

and yyyy (E1), EMD-xxxx and yyyy (E1-ATP), EMD-xxxx and yyyy (E1P-ADP), EMD-

xxxx and yyyy (E2P), EMD-xxxx and yyyy (E2P-transition).

REFERENCES

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd,
J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W.*, et al.* (2010). PHENIX: a
comprehensive Python-based system for macromolecular structure solution. Acta
Crystallogr D Biol Crystallogr *66*, 213-221.

Andersen, J.P., Vestergaard, A.L., Mikkelsen, S.A., Mogensen, L.S., Chalat, M., and
Molday, R.S. (2016). P4-ATPases as Phospholipid Flippases-Structure, Function,
and Enigmas. Front Physiol *7*, 275.

Axelsen, K.B., and Palmgren, M.G. (1998). Evolution of substrate specificities in the P-
type ATPase superfamily. J Mol Evol *46*, 84-101.

Bai, L., Kovach, A., You, Q., Hsu, H.C., Zhao, G., and Li, H. (2019a). Autoinhibition and
activation mechanisms of the eukaryotic lipid flippase Drs2p-Cdc50p. Nat Commun
10, 4142.

Bai, L., Kovach, A., You, Q., Kenny, A., and Li, H. (2019b). Structure of the eukaryotic
protein O-mannosyltransferase Pmt1-Pmt2 complex. Nat Struct Mol Biol *26*, 704-
711.

Baldridge, R.D., and Graham, T.R. (2012). Identification of residues defining
phospholipid flippase substrate specificity of type IV P-type ATPases. Proc Natl
Acad Sci U S A *109*, E290-298.

Baldridge, R.D., and Graham, T.R. (2013). Two-gate mechanism for phospholipid
selection and transport by type IV P-type ATPases. Proc Natl Acad Sci U S A *110*,
E358-367.

Baldridge, R.D., Xu, P., and Graham, T.R. (2013). Type IV P-type ATPases distinguish
mono- versus diacyl phosphatidylserine using a cytofacial exit gate in the membrane
domain. J Biol Chem *288*, 19516-19527.

Best, J.T., Xu, P., and Graham, T.R. (2019). Phospholipid flippases in membrane
remodeling and transport carrier biogenesis. Curr Opin Cell Biol *59*, 8-15.

Bloch, J.S., Pesciullesi, G., Boilevin, J., Nosol, K., Irobalieva, R.N., Darbre, T., Aebi, M.,
Kossiakoff, A.A., Reymond, J.L., and Locher, K.P. (2020). Structure and mechanism
of the ER-based glucosyltransferase ALG6. Nature *579*, 443-447.

Bublitz, M., Morth, J.P., and Nissen, P. (2011). P-type ATPases at a glance. J Cell Sci
608   *124*, 2515-2519.

609 Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral,
610   G.J., Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-
611   atom structure validation for macromolecular crystallography. Acta Crystallogr D Biol
612   Crystallogr *66*, 12-21.

613 Dyla, M., Basse Hansen, S., Nissen, P., and Kjaergaard, M. (2019). Structural dynamics
614   of P-type ATPase ion pumps. Biochem Soc Trans *47*, 1247-1257.

615 Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and
616   development of Coot. Acta Crystallogr D Biol Crystallogr *66*, 486-501.

617 Folmer, D.E., Elferink, R.P., and Paulusma, C.C. (2009). P4 ATPases - lipid flippases
618   and their role in disease. Biochim Biophys Acta *1791*, 628-635.

619 Funakoshi, M., and Hochstrasser, M. (2009). Small epitope-linker modules for PCR-
620   based C-terminal tagging in Saccharomyces cerevisiae. Yeast *26*, 185-192.

621 He, Y., Xu, J., Wu, X., and Li, L. (2020). Structures of a P4-ATPase lipid flippase in lipid
622   bilayers. Protein Cell *11*, 458-463.

623 Hiraizumi, M., Yamashita, K., Nishizawa, T., and Nureki, O. (2019). Cryo-EM structures
624   capture the transport cycle of the P4-ATPase flippase. Science *365*, 1149-1155.

625 Hua, Z., Fatheddin, P., and Graham, T.R. (2002). An essential subfamily of Drs2p-
626   related P-type ATPases is required for protein trafficking between Golgi complex and
627   endosomal/vacuolar system. Mol Biol Cell *13*, 3162-3177.

628 Hua, Z., and Graham, T.R. (2003). Requirement for neo1p in retrograde transport from
629   the Golgi complex to the endoplasmic reticulum. Mol Biol Cell *14*, 4971-4983.

630 Huang, Y., Takar, M., Best, J.T., and Graham, T.R. (2020). Conserved mechanism of
631   phospholipid substrate recognition by the P4-ATPase Neo1 from Saccharomyces
632   cerevisiae. Biochim Biophys Acta Mol Cell Biol Lipids *1865*, 158581.

633 Kanai, R., Ogawa, H., Vilsen, B., Cornelius, F., and Toyoshima, C. (2013). Crystal
634   structure of a Na+-bound Na+,K+-ATPase preceding the E1P state. Nature *502*,
635   201-206.

636 Martin, S., Smolders, S., Van den Haute, C., Heeman, B., van Veen, S., Crosiers, D.,
637   Beletchi, I., Verstraeten, A., Gossye, H., Gelders, G.*, et al.* (2020). Mutated ATP10B

increases Parkinson's disease risk by compromising lysosomal glucosylceramide

export. Acta Neuropathol *139*, 1001-1024.

Mindell, J.A., and Grigorieff, N. (2003). Accurate determination of local defocus and

specimen tilt in electron microscopy. J Struct Biol *142*, 334-347.

Nakano, K., Yamamoto, T., Kishimoto, T., Noji, T., and Tanaka, K. (2008). Protein

kinases Fpk1p and Fpk2p are novel regulators of phospholipid asymmetry. Mol Biol

Cell *19*, 1783-1797.

Natarajan, P., Wang, J., Hua, Z., and Graham, T.R. (2004). Drs2p-coupled

aminophospholipid translocase activity in yeast Golgi membranes and relationship to

in vivo function. Proc Natl Acad Sci U S A *101*, 10614-10619.

Palmgren, M.G., and Axelsen, K.B. (1998). Evolution of P-type ATPases. Biochim

Biophys Acta *1365*, 37-45.

Paterson, J.K., Renkema, K., Burden, L., Halleck, M.S., Schlegel, R.A., Williamson, P.,

and Daleke, D.L. (2006). Lipid specific activation of the murine P4-ATPase Atp8a1

(ATPase II). Biochemistry *45*, 5367-5376.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng,

E.C., and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory

research and analysis. J Comput Chem *25*, 1605-1612.

Pomorski, T., Lombardi, R., Riezman, H., Devaux, P.F., van Meer, G., and Holthuis,

J.C. (2003). Drs2p-related P-type ATPases Dnf1p and Dnf2p are required for

phospholipid translocation across the yeast plasma membrane and serve a role in

endocytosis. Mol Biol Cell *14*, 1240-1254.

Post, R.L., Hegyvary, C., and Kume, S. (1972). Activation by adenosine triphosphate in

the phosphorylation kinetics of sodium and potassium ion transport adenosine

triphosphatase. J Biol Chem *247*, 6530-6540.

Roelants, F.M., Baltz, A.G., Trott, A.E., Fereres, S., and Thorner, J. (2010). A protein

kinase network regulates the function of aminophospholipid flippases. Proc Natl

Acad Sci U S A *107*, 34-39.

Roland, B.P., and Graham, T.R. (2016). Decoding P4-ATPase substrate interactions.

Crit Rev Biochem Mol Biol *51*, 513-527.

Roland, B.P., Jain, B.K., and Graham, T.R. (2020). Exofacial membrane composition
and lipid metabolism regulates plasma membrane P4-ATPase substrate specificity.
bioRxiv.

Roland, B.P., Naito, T., Best, J.T., Arnaiz-Yepez, C., Takatsu, H., Yu, R.J., Shin, H.W.,
and Graham, T.R. (2019). Yeast and human P4-ATPases transport
glycosphingolipids using conserved structural motifs. J Biol Chem *294*, 1794-1806.

Sikorski, R.S., and Hieter, P. (1989). A system of shuttle vectors and yeast host strains
designed for efficient manipulation of DNA in Saccharomyces cerevisiae. Genetics
122, 19-27.

Stevens, H.C., Malone, L., and Nichols, J.W. (2008). The putative aminophospholipid
translocases, DNF1 and DNF2, are not required for 7-nitrobenz-2-oxa-1,3-diazol-4-
yl-phosphatidylserine flip across the plasma membrane of Saccharomyces
cerevisiae. J Biol Chem *283*, 35060-35069.

Tang, X., Halleck, M.S., Schlegel, R.A., and Williamson, P. (1996). A subfamily of P-
type ATPases with aminophospholipid transporting activity. Science *272*, 1495-1497.

Timcenko, M., Lyons, J.A., Januliene, D., Ulstrup, J.J., Dieudonne, T., Montigny, C.,
Ash, M.R., Karlsen, J.L., Boesen, T., Kuhlbrandt, W.*, et al.* (2019). Structure and
autoregulation of a P4-ATPase lipid flippase. Nature *571*, 366-370.

Toyoshima, C., and Cornelius, F. (2013). New crystal structures of PII-type ATPases:
excitement continues. Curr Opin Struct Biol *23*, 507-514.

Toyoshima, C., Iwasawa, S., Ogawa, H., Hirata, A., Tsueda, J., and Inesi, G. (2013).
Crystal structures of the calcium pump and sarcolipin in the Mg2+-bound E1 state.
Nature *495*, 260-264.

UniProt, C. (2019). UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res
47, D506-D515.

van der Mark, V.A., Elferink, R.P., and Paulusma, C.C. (2013). P4 ATPases: flippases
in health and disease. Int J Mol Sci *14*, 7897-7922.

Vestergaard, A.L., Coleman, J.A., Lemmin, T., Mikkelsen, S.A., Molday, L.L., Vilsen, B.,
Molday, R.S., Dal Peraro, M., and Andersen, J.P. (2014). Critical roles of isoleucine-
364 and adjacent residues in a hydrophobic gate control of phospholipid transport by
the mammalian P4-ATPase ATP8A2. Proc Natl Acad Sci U S A *111*, E1334-1343.

Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A.

(2017). MotionCor2: anisotropic correction of beam-induced motion for improved

cryo-electron microscopy. Nat Methods *14*, 331-332.

Zhou, X., and Graham, T.R. (2009). Reconstitution of phospholipid translocase activity

with purified Drs2p, a type-IV P-type ATPase from budding yeast. Proc Natl Acad

Sci U S A *106*, 16586-16591.

Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and

Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure

determination in RELION-3. Elife *7*.

Fig. 1. Structure of the *S. cerevisiae* class-3 lipid flippase Dnf1–Lem3 in the E2P

state. A) Cryo-EM 3D map of Dnf1–Lem3 in the E2P, in front (left) and back (right)

views. The major domains and motifs are labeled in different colors. **B**) Atomic model of

Dnf1–Lem3 in the E2P state in cartoon and colored as in panel A. The detergent and

phospholipid molecules and the covalently linked glycans on Lem3 are shown as sticks.

Fig. 2. The lipid binding sites and the putative substrate transport path of Dnf1–

Lem3. A) Cryo-EM 3D of the region around the putative substrate transporting path.

The transmembrane region is marked by the pink band. Three lipids densities are

shown in grey. PL1 in the exoplasmic side and PL2 in the cytosolic side of the

membrane are highlighted by a purple and a cyan box, respectively. **B**) Cartoon view of

the lipid binding region in the Dnf1–Lem3 structure. The purple box in panel A and the

cyan box in panel B are enlarged in panels C and D, respectively. Note that PL1 is

about 10 Å below the lipid bilayer at the exoplasmic side, and PL2 and PL3 ingress into

the cytosolic region by 10 Å. **C**) Close-up and top view of PL1 binding site. **D**) Close-up

view of PL2 binding site. **E**) Putative substrate transport path of Dnf1p, with the residues

lining the path and bound PL1 and PL2 shown as sticks. The hydrophobic gate in the

lipid transport path is highlighted by the yellow oval. PL3 is removed in the left panel for

clarity.

Figure 3. Influence of lipid binding site mutations on Dnf1 transport activity and

localization. A) NBD-lipid uptake catalyzed by Dnf1 variants expressed on a *dnf1,2Δ*

strain background. **B**) NBD-lipid uptake catalyzed by Lem3 variants expressed on a

lem3Δ background, in which lipid transport is mediated primarily by Dnf2–Lem3. **C**)

NBD-lipid uptake catalyzed by Lem3 variants expressed on a *lem3Δ dnf2Δ* background

to assay the influence of Lem3 mutations solely on Dnf1 activity. In (A-C), cells were

incubated with NBD-lipids for 30 min and uptake was measured by flow cytometry. The

value represents percentage uptake of NBD-lipid in comparison to WT Dnf1 (A) or WT

Lem3 (B-C) for each substrate lipid. The variance was assessed with one-way

ANOVAs, and comparison to WT was calculated with Tukey's post hoc analysis. *n* ≥ 9,

± SD, * indicates $p < 0.05$, **$p < 0.01$, ***$p < 0.001$. **D**) Localization of GFP-tagged Dnf1

variants expressed in *dnf1,2Δ* cells. **E**) WT Dnf1-GFP co-expressed with Lem3 variants.

Arrowheads indicate nuclear envelope/endoplasmic reticulum location. Scale bar = 2

μm (*n* = 40).

Fig. 4. The ATP-dependent transport cycle of Dnf1–Lem3 and Dnf2–Lem3. A) The

five structures of Dnf1 and Dnf2 in the E1 (apo), E1-ATP (AMPPCP), E1P-ADP (AlF$_4^-$-

ADP), E2P-transition (AlF$_4^-$), and E2P (BeF$_3^-$) states. The A domain in the E1, E1-ATP,

and E1P-ADP states is flexible and is shown here as a ribbon in the yellow rectangle to

indicate its position. The structures of the E1-ATP and E1P-ADP states within the black

dashed box are almost identical. The structures of the E2P and E2P-transition states

are similarly grouped for the same reason. Magenta arrows indicate the domain

movement during the Post–Albers cycle in the order E1 to E1-ATP to E1P-ADP to

E2P/E2P-transition. **B**) Structural comparison of the substrate binding sites in the E1,

E1P-ADP, and E2P states. TMH1-2 in the E1 and E1P-ADP states tilt with respect to

that in the E2P state; this movement closes the PL1 site and opens the PL2 site.

763
764

Fig. 5. Structural comparison between Dnf1 (color) and the ATP8A1 (grey) in the
E1 (A), E1P-ADP (B), and E2P (C) states. Bottom panels are close-up views of the
boxed regions in the top panels. These panels highlight the unique helix-turn-helix motif
in the N domain of Dnf1, which interacts with different regions in different states.

769
770

771 **Supplemental Table 1. Cryo-EM data collection, refinement, and validations.**

	Dnf1–Lem3			Dnf2–Lem3				
	Apo	E1-ADP	E2P	Apo	E1-ATP	E1-ADP	E2P-transition	E2P
Data collection and processing								
Magnification (k)	130	130	130	130	130	130	130	130
Voltage (kV)	300	300	300	300	300	300	300	300
Electron exposure (e⁻/Å²)	64	64	64	60	60	60	60	60
Defocus range (-μm)	1.0-2.0	1.0-2.0	1.0-2.0	1.0-2.0	1.5-3.0	1.5-3.0	1.5-3.0	1.5-3.0
Pixel size (Å)	0.826	0.826	0.826	1.029	0.826	0.826	0.826	0.826
Symmetry imposed	C1	C1	C1	C1	C1	C1	C1	C1
Micrograph #	4,074	6,038	4,764	1,438	3,665	5683	5683	3260
Initial particle #	1,358,323	2,257,199	2,094,437	509,555	844,464	1,010,708	1,010,708	1,095,601
Final particle #	473,761	716,510	590,043	190,753	358,687	266,218	127,442	763,661
Map resolution (Å)	3.00	3.25	2.82	3.08	3.85	4.05	3.98	3.50
FSC threshold	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143
Map resolution (Å)	3.12	3.25	2.82	3.08	3.85	4.05	3.98	3.50
Refinement								
Map sharpening B factor (Å²)	116.5	178.9	116.2	102.5	195.8	243.3	197.8	180.0
Model composition								
Non-hydrogen atoms	11150	11235	13121	10369	10684	11960	12521	12879
Protein residues	1332	1322	1552	1247	1274	1471	1534	1545
Ligands & glycans	22	26	26	18	24	15	16	24
B factors (Å²)								
Protein	81	39	47	86	48	175	166	58
Ligand	67	45	34	56	47	174	158	50
R.m.s. deviations								
Bond lengths (Å)	0.01	0.01	0.01	0.004	0.003	0.002	0.002	0.003
Bond angles (°)	0.8	0.9	1.1	0.6	0.6	0.6	0.6	0.6
Validation								
MolProbity score	2.4	2.5	2.0	2.1	2.1	2.1	2.0	2.1
Clashscore	13	11	11	14	12	12	13	13
Poor rotamers (%)	1	1	1	1	0	0	0	1
Ramachandran plot								
Favored (%)	93.0	91.9	95.6	93.0	90.7	93.1	93.9	92.9
Allowed (%)	7.0	8.1	4.4	6.9	9.2	6.9	5.9	7.1
Disallowed (%)	0	0	0	0	0	0	0	0

Supplemental Table 2. List of plasmids used in the study.

Plasmid	Source
pRS313	(Sikorski and Hieter, 1989)
pRS425	(Sikorski and Hieter, 1989)
pRS313-Dnf1	(Baldridge and Graham, 2012)
pRS313-Dnf1 R264A	
pRS313-Dnf1 Y633A	
pRS313-Dnf1 Y633G	
pRS313-Dnf1 T648A	
pRS313-Dnf1 T648V	
pRS313-Dnf1 W652A	
pRS313-Dnf1 W652S	
pRS313-Dnf1 Q610A	
pRS313-Dnf1 S611A	
pRS313-Dnf1 YQS-FSN	
pRS313-Dnf1 N1226A	
pRS313-Dnf1 (692-737)Δ	
pRS416-GFP-Dnf1	(Baldridge and Graham, 2012)
pRS416-GFP-Dnf1 R264A	
pRS416-GFP-Dnf1 W652A	
pRS416-GFP-Dnf1 W652S	
pRS416-GFP-Dnf1 Q610A	
pRS416-GFP-Dnf1 S611A	
pRS416-GFP-Dnf1 YQS-FSN	
pRS416-GFP-Dnf1 N1226A	
pRS425 Lem3	(Baldridge and Graham, 2012)
pRS425 Lem3 (2-49)Δ	
pRS425 Lem3 (400-414)Δ	
pRS425 Lem3 R51A	

775
776

Supplemental Table 3. List of strains used in the study (knockouts are *KanMX* replacements unless otherwise indicated).

Strain	Genotype	Plasmid	Source
BY4741	*MATa his3Δ1 leu2Δ0 ura3Δ0 met15Δ0*		Invitrogen & Horizon Discovery
PFY3275F	MAT**a** *his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 dnf1Δ dnf2Δ*		(Hua and Graham, 2003)
RBY9701	PFY3275F *lem3Δ*::natNT2		(Baldridge and Graham, 2012)
ZHY704	*MATα his3Δ1 leu2Δ0 ura3Δ0 lys2Δ0 dnf1Δ dnf2Δ dnf3Δ drs2Δ::LEU2* pRS416-*DRS2*	pRS416-*DRS2*	(Hua et al., 2002)
SCY119	*MATα his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 lem3Δ::KanMX*		(Baldridge and Graham, 2012)
BJ201	PFY3275F	pRS313	
BJ204	PFY3275F	pRS313-Dnf1	
BJ209	PFY3275F	pRS313-Dnf1R264A	
BJ212	PFY3275F	pRS313-Dnf1Y633A	
BJ215	PFY3275F	pRS313-Dnf1Y633G	
BJ218	PFY3275F	pRS313-Dnf1T648A	
BJ219	PFY3275F	pRS313-Dnf1T648V	
BJ222	PFY3275F	pRS313-Dnf1W652A	
BJ225	PFY3275F	pRS313-Dnf1W652S	
BJ230	PFY3275F	pRS313-Dnf1Q610A	
BJ231	PFY3275F	pRS313-Dnf1S611A	
BJ234	PFY3275F	pRS313-Dnf1YQS-FSN	
BJ237	PFY3275F	pRS313-Dnf1N1226A	
BJ240	PFY3275F	pRS313-Dnf1(692-737)Δ	
BJ241	PFY3275F	pRS313-Dnf1(692-737)Δ	
BJ242	PFY3275F	pRS313-Dnf1(692-737)Δ	
BJ245	PFY3275F	pRS416-GFP-Dnf1	
BJ248	PFY3275F	pRS416-GFP-Dnf1 R264A	
BJ251	PFY3275F	pRS416-GFP-Dnf1 W652A	
BJ254	PFY3275F	pRS416-GFP-Dnf1 W652S	
BJ255	PFY3275F	pRS416-GFP-Dnf1 Q610A	

BJ260	PFY3275F	pRS416-GFP-Dnf1 S611A	
BJ263	PFY3275F	pRS416-GFP-Dnf1 YQS-FSN	
BJ266	PFY3275F	pRS416-GFP-Dnf1 N1226A	
BJ269	ZHY704	pRS313	
BJ272	ZHY704	pRS313-Dnf1	
BJ275	ZHY704	pRS313-Dnf1R264A	
BJ276	ZHY704	pRS313-Dnf1Y633A	
BJ279	ZHY704	pRS313-Dnf1Y633G	
BJ282	ZHY704	pRS313-Dnf1T648A	
BJ285	ZHY704	pRS313-Dnf1T648V	
BJ288	ZHY704	pRS313-Dnf1W652A	
BJ291	ZHY704	pRS313-Dnf1W652S	
BJ294	ZHY704	pRS313-Dnf1Q610A	
BJ297	ZHY704	pRS313-Dnf1S611A	
BJ300	ZHY704	pRS313-Dnf1YQS-FSN	
BJ303	ZHY704	pRS313-Dnf1N1226A	
BJ308	ZHY704	pRS313-Dnf1(692-737)Δ	
BJ309	RBY9701	pRS313, pRS425	
BJ310	RBY9701	pRS313-Dnf1, pRS425-Lem3	
BJ311	RBY9701	pRS313-Dnf1, pRS425-Lem3 (2-49)Δ	
BJ312	RBY9701	pRS313-Dnf1, pRS425-Lem3 (400-414)Δ	
BJ313	RBY9701	pRS313-Dnf1, pRS425-Lem3 R51A	
BJ314	SCY119	pRS416-GFP-Dnf1, pRS425	
BJ315	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3	
BJ316	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3 (2-49)Δ	
BJ317	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3 (400-414)Δ	
BJ318	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3 R51A	

 Supplemental figures

Supplemental Figure 1. Purification of the Dnf1–Lem3 and Dnf2–Lem3 complexes.

A) Gel filtration profile of the Dnf1–Lem3 complex. **B**) The peak fraction in (A) was checked using the Coomassie blue-stained SDS-PAGE gel. **C**) Gel filtration profile of the Dnf2–Lem3 complex. **D**) Peak fraction in (C) was checked using the Coomassie blue-stained SDS-PAGE gel.

Supplemental Figure 2. Cryo-EM data processing and resolution estimation of Dnf1–Lem3 in the E2P state. A) A representative electron micrograph. **B**) Selected reference-free 2D class averages. **C**) Cryo-EM data processing procedure. **D**) Gold-standard Fourier shell correlation of two independent half 3D maps. **E**) Local resolution

map of the final 3D map. **F**) Angular distribution of raw particles used in the final

reconstruction of the 2.8-Å 3D map.

Supplemental Figure 3. Selected regions in the 3D map of Dnf1–Lem3 in the E2P

state, superimposed on the atomic model.

Supplemental Figure 4. **Cryo-EM data processing and validation of the Dnf1–**

Lem3 in the E1 and E1P-ADP states. A) Cryo-EM data processing procedure of Dnf1

in the E1 state. **B**) Gold-standard Fourier shell correlation of two independent half 3D

maps in the E1 state. **C**) Cryo-EM data processing procedure of Dnf1 in the E1P-ADP

state. **D**) Gold-standard Fourier shell correlations of two independent half 3D maps of

Dnf1p in the E1P-ADP state. In (**A**) and (**C**), the bottom panel shows the final map

colored according to the local resolution.

Supplemental Figure 5. **Cryo-EM data processing and resolution estimation of the**

Dnf2–Lem3 in the E1 and E1-ATP states. A) Cryo-EM data processing procedure of

Dnf2 in the E1 state. **B**) Gold-standard Fourier shell correlations of two independent half

3D maps in the E1 state. **C**) Cryo-EM data processing procedure of Dnf2in the E1-ATP

state. **D**) Gold-standard Fourier shell correlations of two independent half maps of the

Dnf2p-E1-ATP. In (**A**) and (**C**), the bottom panel shows the final map colored according

to the local resolution.

Supplemental Figure 6. Cryo-EM data processing and validation of the Dnf2–

Lem3 in the E2P-transition state and in the E1P-ADP state. A) Cryo-EM data

processing procedure of the Dnf2–Lem3 datasets in the E2P-transition and the Dnf2p-

E1P-ADP states. The two bottom panels show the two 3D maps colored according to

their respective local resolutions. **B**) Gold-standard Fourier shell correlations of two

independent half 3D maps of Dnf2–Lem3 in the E2P-transition state and in the E1P-

ADP state.

Supplemental Figure 7. Cryo-EM data processing and validation of the Dnf2–

Lem3 in the E2P state. A) Cryo-EM data processing procedure. The final 3D map

shown at the bottom is colored coded by the local resolution. **B**) Gold-standard Fourier

shell correlation of two independent half 3D maps.

Supplemental Figure 8. Structural alignments between the Dnf1 and Dnf2 in their

corresponding E1 (apo) (**A**), the E1P-ADP (**B**), and the E2P states (**C**) show a high

similarity.

Supplemental Figure 9. The N-terminal peptide of Lem3p is partially ordered in the 3D map of Dnf1–Lem3 in the E2P state. A) Cryo-EM 3D map of the Dnf1–Lem3 in the E2P state. The ordered Lem3 N-terminal peptide is highlighted by the black and red boxes. **B**) An enlargement of the black box in panel A, shown at a lower display threshold to show the weaker density. **C**) An enlarged view of the red box in panel A. The Lem3 N-terminal peptide interacts with the Dnf1 A domain. The interface is highly positive. **D**. Sequence alignment of yeast Lem3 (ScLem3) and other CDC50 family proteins (yeast Crf1, yeast Cdc50, human CDC50A, and human CDC50B) shows the Lem3-specific, N-terminal, negatively charged motif.

857 **Supplemental Figure 10. Structural comparison between Dnf1 (color) and the**

858 **ATP8A1 (grey) in the E2P state.** Lipids in the Dnf1 structure are shown in red, while

859 the only lipid in ATP8A1 is in blue. The PL binding sites in the exoplasmic side of

860 membrane of the two flippases are well aligned. The right panel is an enlargement of

861 the substrate transport path of the left panel.

865
866

Supplemental Figure 11. Ability of Dnf1 variants to support viability of flippase-

deficient cells. *HIS3*-marked plasmids (pRS313) carrying the indicated Dnf1 variants

were used to transform a *dnf1,2,3∆drs2∆* pRS416-*DRS2* strain. Strains were spotted

onto minimal media plates (SD) selecting for both plasmids, and pRS416-*DRS2* was

counter-selected on medium containing 5-fluoroorotic acid (SD-FOA). Strains were

incubated at 30 °C for 2 d and imaged. The growth phenotype on SD-FOA is an

indicator of how well the Dnf1 variant functions *in vivo*.

Supplemental Figure 12. High structural similarity between Dnf1 and Dnf2 in the corresponding substrate transport states. A) Structural alignments 3D maps of Dnf2–Lem3 in the E1-ATP and E1P-ADP states. **B**) Structural alignments of 3D maps of Dnf2 in the E2P and E2P-transition states.

Supplemental Figure 13. The N and A domain densities are weak in the 3D maps

of Dnf1–Lem3 in the E1 and E1P-ADP states, indicating their high flexibility. A)

Cryo-EM 3D map of the Dnf1-Lem3 in the E1 state displayed at a high (top panel) and a

low threshold (bottom panel). Red arrows indicate the domain movement from the E1

state to the E1-ATP state. **B**) Cryo-EM 3D map of the Dnf1–Lem3 in the E1-ATP state

displayed at a high (top panel) and a low threshold (bottom panel). Red arrows indicate

the domain movement from the E1P-ADP state to the E2P state. **C**) Cryo-EM 3D map of

the Dnf1–Lem3 in the E2P state. **D-F**, Selected reference-free 2D class averages of the

Dnf1–Lem3 in the E1 state (**D**), in the E1P-ADP state (**E**), and in the E2P state (**F**). The

averages in red or green boxes belong in the same views. The A domain locations are

labeled by red arrowheads. The weak and blurry A domain density indicates its flexibility

in the E1 and E1P-ADP states.