Transport Mechanism of Class-3 P4 ATPase Lipid Flippases

Lin Bai[1,#,*], Qinglong You[1,#], Bhawik K. Jain[2,#], Amanda Kovach[1], Todd R. Graham[2], and Huilin Li[1,*]

[1] Structural Biology Program, Van Andel Institute, Grand Rapids, Michigan, USA

[2] Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

[#] These authors contributed equally to the work.

Correspondence should be addressed to L.B. (Lin.Bai@vai.org) or H.L. (Huilin.Li@vai.org).

ABSTRACT

The P4 ATPases are a large family of membrane-embedded enzymes that use ATP hydrolysis to transport large lipid substrates across lipid bilayers. P4 ATPases differ in their cellular membrane location and their substrates. The structures of the endosome- and Golgi-localized class-1 phosphatidylserine flippases—such as the yeast Drs2 and human ATP8A1—have recently been reported, revealing a substrate binding site on the lumenal side and several transport states. However, a substrate binding site on the cytosolic side has not been found, and the transport mechanisms of P4 ATPases in other classes are still unknown. Here we report a systematic structural and functional study on two plasma-membrane localized, class-3 P4 ATPases that have broader substrate specificity, the *S. cerevisiae* Dnf1–Lem3 and Dnf2–Lem3 complexes. We have captured substrate lipids on both the exoplasmic and cytosolic sides, and we found that these two enzymes have very similar structures, consistent with their high sequence identity and redundant function. Unexpectedly, Lem3 contributes to substrate binding near the cytosolic surface. We found that the conformational transitions through the substrate

transport cycle of these two class-3 enzymes match those of the class-1 enzymes, suggesting a

conserved lipid-flipping mechanism among all classes of the P4 ATPases. Our study also

revealed a helix-turn-helix insertion in the cytosolic P domain that is unique to the class-3

enzymes and plays a crucial role in their function. Therefore, the P4 ATPases may have

retained an overall transport mechanism while evolving distinct features for cellular membrane

localization, regulatory mechanisms, and transporting different lipid substrates.

INTRODUCTION

The large group of integral membrane proteins that mediate the transport of substrates across biological membranes by the formation of ATP-dependent phosphorylated intermediates are called P-type ATPases (Axelsen and Palmgren, 1998). They are widely expressed in prokaryotes, archaea, and eukaryotes, and they play essential roles in many cellular processes. P-type ATPases are phylogenetically divided into five subclasses (P1-P5), each have unique sequence motifs and transport different substrates (Bublitz et al., 2011; Palmgren and Axelsen, 1998). Most P4 ATPases function as a heterodimeric complex: a catalytic α-subunit containing 10 transmembrane α-helices (TMHs) and a regulatory β-subunit of the Cdc50 (cell division cycle 50) protein family containing two TMHs. The α-subunit of all P-type ATPases has a conserved architecture consisting of a transmembrane domain (TMD), an actuator domain (A-domain), a nucleotide-binding domain (N-domain), and a phosphorylation domain (P-domain) (Andersen et al., 2016; van der Mark et al., 2013). The "Post–Albers" model proposed that the substrate transport of P-type ATPases goes through a cyclic transition of E1–E1P–E2P–E2 states (Post et al., 1972). This model is supported by the structure and function studies of the most well-understood P2 ATPases (Dyla et al., 2019; Toyoshima and Cornelius, 2013).

P4 ATPases are involved in phospholipid translocation: they flip lipids from the extracellular side of the plasma membrane or from the lumenal side of internal organelles to the cytosolic side (Tang et al., 1996; Zhou and Graham, 2009). They are important for creating and maintaining lipid compositional asymmetry in cell membranes. Because a phospholipid substrate is large, the transport mechanism of P4 ATPases is substantially different from that of the cation-transporting P-type ATPases, such as the sarcoplasmic reticulum Ca^{2+}-ATPase (SERCA), the Na^+/K^+-ATPase, and the H^+-ATPase. There are five P4-ATPases belonging to four classes in *Saccharomyces cerevisiae*: Drs2 in class 1, Neo1 in class 2, Dnf1 and Dnf2 in class 3, and Dnf3 in class 4 (van der Mark et al., 2013). Humans have 14 P4-ATPases (Andersen et al., 2016; Best et al., 2019; van der Mark et al., 2013). The P4 ATPases have different tissue-specific expression, subcellular localization, and substrate specificity. For example, yeast Drs2 transports phosphatidylserine (PS) and phosphatidylethanolamine (PE) in the trans-Golgi network (TGN) and endosomes (Natarajan et al., 2004; Zhou and Graham, 2009), while the yeast Dnf1 and Dnf2 flip PE, PC, and glucosylceramide (GlcCer) in the plasma membrane (Best

et al., 2019; Pomorski et al., 2003; Roland et al., 2020; Roland et al., 2019; Stevens et al.,

2008).

The first structural and mechanistic insights into the P4 ATPases came from recent cryo-EM

structures of the *S. cerevisiae* class-1 flippase Drs2–Cdc50 in its auto-inhibited and active E2P

states (Bai et al., 2019a; Timcenko et al., 2019), and of its human homolog, ATP8A1–CDC50a,

in six transport states (E1, E1-ATP, E1P-ADP, E1P, E2P, and E2Pi-PL) (Hiraizumi et al., 2019).

Drs2 and ATP8A1 belong to the same subfamily of P4 ATPases, and both transport the same

substrates, PS and PE (Paterson et al., 2006; van der Mark et al., 2013; Zhou and Graham,

2009). The structural studies revealed an allosteric domain movement during lipid translocation

and identified a lipid binding site at the exoplasmic side of membrane of this subfamily. Prior

mutational studies had identified amino acids in this exoplasmic lipid binding site that are

important for substrate recognition and had implicated a second substrate binding site on the

cytosolic side of the membrane (Baldridge and Graham, 2012, 2013; Baldridge et al., 2013;

Roland et al., 2019).

However, many important questions remain regarding the lipid-flipping mechanism(s) of the P4

ATPases. Chief among them are how substrate specificity is achieved by different P4 ATPases

and where the lipid binds on the cytoplasmic side of the membrane. Addressing these questions

requires a detailed structural and functional comparison among P4 ATPases that transport

different substrates. Toward this goal, we have now studied the two *S. cerevisiae* class-3 P4

ATPases, the Dnf1–Lem3 and Dnf2–Lem3 complexes. Dnf1 and Dnf2 share the same β-

subunit, Lem3, and both transport PE, PC, and glucosylceramide. Their human homologues,

ATP10A and ATP10D, are implicated in diabetes, obesity, myocardial infarction,

atherosclerosis, and Parkinson's disease (Folmer et al., 2009; Martin et al., 2020; van der Mark

et al., 2013). Here we report the cryo-EM structures of Dnf1–Lem3 in three different states (E1,

E1P-ADP, and E2P) and Dnf2–Lem3 in five different states (E1, E1-ATP, E1P-ADP, E2P, and

E2P-transition). We further performed structure-guided mutagenesis and *in vivo* substrate

transport activity assays. By comparing these class-3 flippase structures with the class-1

structures previously reported, we found new phospholipid binding sites at the cytoplasmic side

and unique structural features during the ATP-dependent transport cycle in Dnf1–Lem3 and

Dnf2–Lem3. Our comprehensive structural and functional study improves the mechanistic

understanding of the essential P4 ATPases.

RESULTS AND DISCUSSION

1. Overall Dnf1 structure

Endogenous Dnf1–Lem3 and Dnf2–Lem3 complexes were purified from baker's yeast using an

anti-FLAG affinity column followed by size-exclusion chromatography (**Supplemental Figure 1,**

online Methods). We used the detergent dodecyl maltoside (DDM) to solubilize these flippases

and lauryl maltose neopentyl glycol (LMNG) and cholesteryl hydrogen succinate (CHS) to

stabilize the membrane protein complexes. We performed single-particle cryo-EM on purified

samples in apo form (E1 state) and on samples incubated with AMPPCP (E1-ATP state),

$ALF4^-$-ADP (E1P-ADP and E2P-transition states), or $BeF3^-$ (E2P state). The substrate transport

states were assigned based on the established nomenclature in the P-type ATPase field (Bai et

al., 2019a; Hiraizumi et al., 2019; Timcenko et al., 2019). We obtained cryo-EM 3D maps of the

S. cerevisiae Dnf1–Lem3 in the E1, E1P-ADP, and E2P states and Dnf2–Lem3 in the E1, E1-

ATP, E1P-ADP, E2P-transition, and E2P states, at resolutions of 2.8 Å to 4.0 Å (**Supplemental**

Figures 2-7, Supplemental Table 1). The high-resolution 3D map at 2.8 Å of the Dnf1–Lem3 in

the E2P state enabled us to build the first atomic model of Dnf1–Lem3 (**Fig. 1, Supplemental**

Figure 2). Except for a few short and disordered loops, almost the entire complex is ordered

and well resolved in this model. Interestingly, many extra densities were observed in the

transmembrane region of Dnf1 in the E2P state (**Fig. 1A**). Some clearly belonged to the

detergent CHS that was added during purification, while others located in the substrate

transport path resembled phospholipid, and they were so assigned, as will be described below.

We then used the 2.8-Å structure to help build atomic models for the seven remaining 3D maps

of Dnf1p–Lem3p and Dnf2p–Lem3p, which are at slightly lower resolutions. All these models

were refined to good statistics (**Supplemental Table 1**) and fit well with the 3D maps

(**Supplemental Figure 3**). The modeled nucleotides and lipids all had clear densities. By

comparing the 3D maps of Dnf1–Lem3 and Dnf2–Lem3 in the E1, E1P-ADP, and E2P states,

which were the three states observed for both flippases, we found that the structures were

superimposable in all three states (**Supplemental Figure 8**). This comparison suggests that

Dnf1 and Dnf2 have highly similar structures and go through virtually the same functional states

during their respective substrate transport cycles. This finding was not surprising, because

these two class-3 flippases share a high sequence identity of about 69%, localize to the same

membrane, transport the same substrates, and appear to have redundant functions in yeast

(Pomorski et al., 2003). Because of their apparently interchangeable structural feature, we will

discuss these two flippases together in the following sections.

We first examined the Dnf1 structure in the E2P state because of the high resolution and

completeness of the structure. As expected of any P4 ATPase, Dnf1 has a 10-TMH TMD

(TMH1-10); a cytosolic A domain inserted between TMH2 and TMH3; and a cytosolic P domain

and a cytosolic N domain, both inserted between TMH4 and TMH5 (**Fig. 1**). The amino terminal

peptide preceding TMH1 (N-tail) of Dnf1 is 204 residues long and is largely disordered, except

for the last 39-residue segment (Glu-166 to Leu-204), which is stabilized by binding to the

regulatory A domain. The carboxyl terminal peptide (C-tail) of Dnf1 following the 10th TMH is

157 residues long (Pro-1405 to Asn-1571) and is also largely disordered except for the TMD-

proximal 35-residue segment (Pro-1405 to Lys-1439).

Interestingly, this ordered Dnf1 C-tail segment contains a short α-helix, the equivalent of which

was reported to function as a switch that converts the Drs2 flippase from its auto-inhibitory state

to the active state in a unique PI4P-binding-dependent manner (Bai et al., 2019a). However, the

Dnf1 helix is stabilized by the C-tail of Lem3 in a conformation corresponding to the active state

of Drs2, suggesting that Dnf1 and Dnf2 do not require binding to a phosphoinositide at this site

for activation. The Dnf1 C-tail interacts extensively with the cytoplasmic domains of Dnf1 in

regulating the substrate transport cycle, as described below. Dnf1 and Dnf2 activity is regulated

by two flippase kinases Fpk1 and Fpk2. Mutations in Fpk1 and Fpk2 do not affect the

subcellular location of Dnf1 and Dnf2, but these enzymes have almost no activity in a *fpk1,2*Δ

strain (Nakano et al., 2008; Roelants et al., 2010). We found that all 17 known phosphorylation

sites (P32660 in the UniProt Database (UniProt, 2019)) are located in disordered loop regions in

the Dnf1 structure: the N-tail, a loop between Thr-290 and Ala-378, and the C-tail. These

observations suggest that the class-3 flippases are autoinhibited and phosphorylation, not PI4P

binding as in the class-1 flippases, relieves the inhibition.

2. Lem3 has unique and ordered N- and C-terminal regions

Similar to the yeast Cdc50 and human CDC50a, Lem3 contains two TMHs flanking a large

lumenal domain. These TMHs are packed against the last TMH of Dnf1, TMH10 (**Fig. 1**). The

Lem3 lumenal domain contains two disulfide bonds (Cys-110 with Cys-159 and Cys-216 with

Cys-231) and four N-glycans at Asn-240, Asn-256, Asn-298, and Asn-332. To our surprise, both

the N-terminal and the C-terminal regions of Lem3 are stable, in contrast to the yeast Cdc50

and human CDC50a, in which these regions are disordered (**Fig. 1A-B**). The N-terminal peptide

preceding Asn-50 folds into a two-turn α-helix and binds the Dnf1 A domain (**Supplemental**

Figure 9A-B). Due to a partial flexibility, the exact sequence of the Lem3 N-tail α-helix was not

resolved. However, we found its binding interface in the A domain is highly positively charged,

featuring Lys-392, Lys-395, Arg-408, Arg-457, Lys-460, and Lys-516 (**Supplemental Figure**

9C), and the Lem3 N-tail segment between Glu-28 to Glu-42 is highly negative, containing 11

Asp and Glu residues (**Supplemental Figure 9D**). Therefore, the short N-tail α-helix of Lem3p

is most likely from this negatively charged region. Because the A domain moves significantly

during the substrate transport cycle of P4 ATPase, this unique, A domain-interacting, and highly

negative Lem3 N-tail α-helix likely regulates lipid transport activity of Dnf1 and Dnf2, as

described below.

3. Phospholipids bind at the exoplasmic and cytoplasmic sites of Dnf1

Although the exact substrates a P4 ATPase transport differ, the substrate translocation path

appears to have been conserved in all P4 ATPase studied so far (Andersen et al., 2016; Huang

et al., 2020). Like the Drs2 and ATP8A1 flippases, we found that the Dnf1 substrate

translocation path is composed of TMH1-4 and TMH6 (**Fig. 2a-d, Supplemental Figure 10**).

This substrate path is consistent with biochemical studies demonstrating that TMH1-6 are

responsible for phospholipid transport, with TM7-10 having a supporting role (Baldridge and

Graham, 2013; Vestergaard et al., 2014). In the E2P-state Dnf1 3D map, we observed three

phospholipid-like densities in the substrate transport path, and accordingly, we modeled three

phospholipid molecules (PL1-PL3) in that map (**Figs. 1A, 2A**). These phospholipids stack

against Dnf1 TMH2, TMH4, and TMH6. Because no lipids were added during purification, these

phospholipids are potentially endogenous substrates that were co-purified with the Dnf1–Lem3

complex. This is not unusual, because previous studies have shown that substrate lipids bound

in the substrate pockets can be co-purified with the enzyme complexes (Bai et al., 2019b; Bloch

et al., 2020; Hiraizumi et al., 2019). Interestingly, the three phospholipids are out of register with

respect to the lipid bilayer by approximately 10 Å: PL1 is moved down about 10 Å into the

middle of the bilayer relative to lipids in the outer leaflet, and PL2 and PL3 are moved 10 Å into

the cytosolic space relative to lipids in the inner leaflet of the membrane.

4. Functional studies reveal that PL1 and PL2 are in substrate binding sites

Based on sequence conservation of the three phospholipid binding pockets, we expect that PL1

and PL2—but not PL3—are located in the substrate path and are likely physiologically relevant.

To test this hypothesis, we carried out systematic, structure-based mutagenesis and functional

assays (**Fig. 3A-E, Supplemental Table 3**). PL1 is bound in an open cleft of Dnf1 on the

exoplasmic side of the plasma membrane, and the phosphate group of PL1 interacts with

several conserved residues, including Gln-610, Ser-611, and Asn-1226 (**Fig. 2A-B**). We

expressed the Q610A, S611A, 609YQS611-FSN, and N1226A Dnf1 mutants on a *dnf1Δ dnf2Δ*

cell background and measured their PC, PE, and GlcCer uptake abilities. We found that Q610A

and the triple mutation 609YQS611-FSN eliminated the GlcCer and PE, but not the PC,

transport activity (**Fig. 3A**). This observation was consistent with a previous finding in which

mutations of the conserved tripeptide motif 654YQS656 of Dnf2—the equivalent of 609YQS611

in Dnf1—caused substantial and specific loss of GlcCer transport (Roland et al., 2020; Roland

et al., 2019).

It is unclear why the S611A mutation led to increased transport activity for all three substrates

(**Fig. 3A**). In contrast, the N1226A mutation completely halted transport of all three substrates,

even though the mutant protein localized to the plasma membrane comparably to WT Dnf1 (**Fig.**

3D). The ability of Dnf1-N1226A to exit the ER indicated it was folded properly and was able to

interact with Lem3, but consistent with the loss of transport activity, Dnf1-N1226A failed to

support the viability of a flippase-deficient yeast strain, whereas Dnf1 variants that retained

transport activity fully supported viability (**Supplemental Figure 11**). The GA motif (230-231) in

TMH1B is at the membrane-exoplasmic interface and is next to the PL1 binding site (**Fig. 2D**). A

recent study has shown that the GA motif is crucial for GlcCer transport by Dnf2 but has little

influence on PC and PE transport (Roland et al., 2019). Furthermore, we found that the PL1

binding site superimposed with the substrate binding site of ATP8A1, based on structural

alignment (**Supplemental Figure 11**). Taken together, we conclude that the PL1 binding site is

the entrance of the substrate transporting path in Dnf1 and Dnf2.

Unlike PL1, both PL2 and PL3 bind Dnf1 on the cytosolic side of the lipid bilayer. The head

group of PL2 is in a positively charged groove and is stabilized by several conserved residues,

including Arg-264, Tyr-633, Thr-648, and Trp-652 of Dnf1, and Arg-51 of Lem3. Because PL2 is

in the putative substrate transport path, we suggest that PL2 is at the substrate exit site of Dnf1.

We generated a series of point mutations in Dnf1, including R264A, Y633A, T648A, W652A,

and W652S. We expressed these mutations in *dnf1Δ dnf2Δ*, *lem3Δ*, or *lem3Δ dnf2Δ* cells and

found that W652A and W652S significantly disrupted transport of all substrates; that R264A

enhanced PC, PE, and GlcCer transport; and that Y633A and T648A had no significant effects

on the transport activities (**Fig. 3A**). The elimination of Dnf1 transport activity by the W652

mutations was consistent with this residue's strategic location between the PL2-binding

substrate exit site and the PL1-binding substrate entry site. However, this mutation caused the

retention of Dnf1 in the ER, possibly because it disrupts folding of the protein (**Fig. 3E**). The

increased activity of Dnf1 R264A suggests that this charged residue may trap the substrates

and slow their release into the cytosolic leaflet. This observation supports our hypothesis that

PL2 is a substrate lipid. Because the PL3 site is located right next to the PL2 site, and the PL3

site is composed of nonconserved residues (except for F631), we suggest that the PL3 site

plays an accessory role, perhaps by facilitating substrate release from the PL2 site.

5. Lem3 unexpectedly contributes to substrate binding on the cytosolic side

Because the Lem3 Arg-51 directly interacts with PL2, we generated a Lem3 R51A mutant and

found that the substitution changed the substrate specificity of both Dnf1 and Dnf2, with a 1.5-

to 2-fold increase in GlcCer transport activity of Dnf2–Lem3 and Dnf1–Lem3, but no change in

PC or PE activity for either complex (**Fig. 3B-C**). Because there is no phosphate in GlcCer, it is

possible that R51A increases ability of GlcCer to compete with endogenous lipids for this site.

As expected, Lem3-R51A supported trafficking of the complex out of the ER (**Fig. 3E**), and this

result supports the role of Lem3 R51 in controlling substrate specificity and further supports the

assignment of PL2 as a substrate lipid.

To examine the functional importance of the N-terminal region of Lem3, we generated a Lem3

variant missing the N-terminal peptide (Lem3[Δ2-49]) and performed a lipid uptake assay. We

found that with Lem3(Δ2-49), both Dnf1 and Dnf2 had significantly reduced PC, PE, and GlcCer

uptake ability (**Fig. 3B-C**). We next examined the functional importance of the ordered C-

terminal region of Lem3 (Gly-400 to Lys-414) that stabilizes Dnf1 in an active state. In the lipid

uptake assays with Lem3(Δ400-414), both Dnf1 and Dnf2 had substantially decreased activity

(**Fig. 3B-C**). We then labeled Dnf1 with GFP and found that Lem3(Δ2-49) and Lem3(Δ400-414)

abolished Lem3's ability to escort Dnf1 out of the ER (**Fig. 3E**). Thus, neither Dnf1 or Dnf2 was

able to reach to its functional destination in the plasma membrane with a N- or C-terminus

truncated Lem3. Thus, these results support an unexpected role of Lem3 in substrate binding

and specificity. No such role was found in the β-subunit (Cdc50 or CDC50) of the class-1

flippases Drs2 and ATP8A1 (Bai et al., 2019a; Hiraizumi et al., 2019; Timcenko et al., 2019).

During preparation of this manuscript, the cryo-EM structure of Dnf1–Cdc50 of a thermophilic

fungus, *Chaetomium thermophilum*, was reported in the E1-ATP and E2P states at 3.4-3.5 Å

resolution (He et al., 2020). That Dnf1 structure contained two lipids, but no evidence was

provided that they were substrate lipids. Many membrane proteins, including SERCA and

Na^+/K^+ ATPases (Kanai et al., 2013 ; Toyoshima et al., 2013), have specifically bound lipid

molecules in their structures that are presumably not substrate lipids. Because we have now

functionally defined that the PL1 and PL2 in our *S. cerevisiae* Dnf1–Lem3 and Dnf2–Lem3

structures are located at the physiological substrate entry and exit, the observation that the two

lipids in *C. thermophilum* Dnf1 are equivalent to PL1 and PL2 in our structure indicates that in

both species, the class-3 flippases share a similar substrate path.

We conclude that the substrate transport pathway is primarily located between TMH2 and

TMH4 (**Fig. 2E**). This pathway is nearly perpendicular to the membrane plane and is open to the

lipid bilayer. Interestingly, this pathway is hydrophilic at both ends, the entry and the exit sites,

but is highly hydrophobic in the middle gating region (**Fig. 2B-E**). In the Dnf1–Lem3 structure, the exoplasmic substrate entry site is lined by Gln-610, Ser-611, and Asn-1226; the cytosolic substrate exit site is surrounded by Dnf1 residues Gln-214, Asn-217, Asn-220, Thr-254, Lys-257, Asp-258, Glu-261, Arg-264, Arg-265, Glu-622, and Thr-626, along with Lem3 Arg-51, and the middle hydrophobic region is composed of the conserved Phe-223, Leu-247, Ile-250, Ile-615, and Ile-619 (**Fig. 2B-E**). The physiochemical property of the transport pathway is compatible with the so-called "credit card" mechanism, in which the hydrophilic or charged headgroups of the lipid substrates simply slide through the groove while allowing the substrate acyl chains to flip in the favorable membrane environment. It is likely that during transport, the lipid head group inserts into and make direct interactions with the hydrophilic residues between TMH2 and TMH4.

6. ATP dependent transport cycle

By cryo-EM analyses of samples incubated with or without the ATP analogs AMPPCP and ADP-AlF$_4^-$ or the phosphate analogs AlF$_4^-$ and BeF$_3^-$, we obtained a total of eight structures of the class-3 flippases. There were Dnf1–Lem3 in the E1 (apo), E1P-ADP (ALF$_4^-$-ADP), and E2P (BeF$_3^-$) states; and Dnf2–Lem3 in the E1 (apo), E1-ATP (AMPPCP), E1P-ADP (ADP-AlF$_4^-$), E2P-transition (AlF$_4^-$), and E2P (BeF$_3^-$) states. Comparing these structures revealed how ATP binding and hydrolysis, ADP release, and enzyme phosphorylation produce conformational changes in Dnf1 and Dnf2 to drive the Post–Albers substrate transport cycle (**Fig. 4A**).

We found that a major part of the TMD region (that is, TMH3-10) and the P domain of Dnf1 or Dnf2, and the most of Lem3 molecule except for its N-tail, were well aligned, suggesting that these regions are largely stationary during the substrate transport cycle (**Fig. 4A, Supplemental Figure 12A-B**). However, TMH1-2 and the A and N domains of Dnf1 and Dnf2 moved significantly; these movements must be coupled with substrate transport. In the E1 state, the A domain moves towards the membrane bilayer and becomes separated from the N domain (**Fig. 4A**). Due to the lack of interaction, the A and N domains are flexible and are visible in the 3D maps only at a low display threshold (**Supplemental Figure 13**).

The binding of ATP in the E1-ATP state triggers the N domain to move towards and interact with

the A domain, thereby stabilizing both domains. However, the A domain must be only partially

stabilized and retain some flexibility, because its density is still weak. With the ensuing

phosphoryl transfer, the flippases transition from the E1-ATP state to the E1P-ADP state. This

transition does not trigger large-scale conformational changes, because the structures of Dnf2–

Lem3 in the E1-ATP and E1P-ADP are almost the same except for their respective bound

nucleotide (**Supplemental Figure 12**). In the E1, E1-ATP, and E1P-ADP states, TMH1-2 have

a small degree of flexibility, but they do not move much (**Fig. 4B**). Transitioning from the E1P-

ADP state to the E2P state, the release of ADP triggers the N domain to rotate 40° away from

the A domain by pivoting on the N domain and P domain interface. The conserved DGET motif

of the A domain now occupies the phosphorylation site and stabilizes the A domain. During this

transition, TMH1 and TMH2 tilt to enlarge the exoplasmic substrate binding site for the

admission of PL1 and at the same time to narrow the cytosolic substrate site, thereby stabilizing

the bound PL2 (**Fig. 4B**).

During the subsequent phosphate (Pi) transition, when the enzyme changes from the E2P state

to the E2P-transition state, the enzyme structure largely holds still; the 3D maps of Dnf2–Lem3

in these two states are nearly identical (**Supplemental Figure 12B**). Finally, the Dnf1 and Dnf2

switch back to the E1 state by auto-dephosphorylation. During this switch, the A and N domains

become flexible again, and the TMH1-2 tilt back to narrow and close the exoplasmic substrate

binding site and open the cytosolic substrate binding site. It is during this last transition that PL1

in the exoplasmic site is forced to flip and enter the cytosolic substrate site, replacing the

resident PL2 there and leading to the release of PL2 from the flippases.

7. Transport mechanisms of the class-3 and class-1 P4 ATPases are similar

As described above, the major domain rearrangements in Dnf1 and Dnf2 are triggered by

nucleotide binding (E1 to E1-ATP); nucleotide hydrolysis, enzyme phosphorylation, nucleotide

release (E1P-ADP to E2P); and enzyme dephosphorylation (E2P to E1). The energy gain from

ATP during these transitions is transduced from the phosphorylation site in the P domain to the

A domain, which in turn causes the TMH1-2 to tilt and drive the flipping of the phospholipids.

TMH3-10 of Dnf1 or Dnf2 and TMH1-2 of Lem3 largely remain stationary throughout the

transport cycle. Combining the finding of two substrate binding sites and the transport cycle, we

propose that the lipid head group transports through the groove between TMH2 and TMH4,

while lipid tails move accordingly on the hydrophobic surface of the TM2 and TM4. This is

essentially the same as the previously described "credit card model" (Andersen et al., 2016;

Roland and Graham, 2016).

The previous cryo-EM study of the human ATP8A1–CDC50a flippase captured multiple

transport states (Hiraizumi et al., 2019). Aligning our structures in this paper with those of

ATP8A1–CDC50a reveal that the corresponding E1, E1-ATP, E1P-ADP, E1P, and E2P states

are remarkably similar in both domain architecture and substrate transport path (**Fig. 5**). This

result was unexpected, because ATP8A1 is a class-1 flippase and Dnf1 and Dnf2 are class-3.

Our observation raises the possibility that all P4 ATPases may employ a similar substrate-

transport mechanism.

Despite their overall similarity in substrate transport cycle, notable structural differences exist

among the flippases characterized so far. In class-1 flippases such as the yeast Drs2 and

human ATP8A1, the C-terminal loop is auto-inhibitory; this loop, which contains the conserved

GF/YAFS motif, extends over and wraps around the cytoplasmic catalytic domains to stabilize

the class-1 enzymes in the E2P state. The C-terminal loop of Dnf1 and Dnf2 doesn't contain the

GF/YAFS motif and is disordered in the E2P state. However, there is a unique helix-turn-helix

motif (E692-S737) in the N domain of Dnf1 and Dnf2 that is absent in Drs2 and ATP8A1 (**Fig.**

5). In the E1 state, the helix-turn-helix interacts with the P domain, specifically, its Lys-702 and

Arg-703 interacting with the P domain Glu-994, Asn-998, and Asn-1008. Upon ATP hydrolysis

and enzyme phosphorylation in the E1P-ADP state, this interaction is weakened because only

Arg-703 of the motif now interacts with the P domain Asn-998 and Asn-1008. The release of

ADP in the E2P state appears to shift the interaction of the Dnf1 and Dnf2 insertion from the P

domain to the C-tail, with the helix-turn-helix Glu-709 and Arg-713 forming electrostatic

interactions with the C-tail Arg-1434 and Lys-1436. These systematic changes suggest that the

Dnf1 and Dnf2 insertion may play a regulatory role in the substrate transport cycle. To examine

this possibility, we deleted the motif in Dnf1 and performed an *in vivo* substrate uptake assay.

The deletion strongly decreased transport of all substrates (**Fig. 3A**). These results confirm a

crucial role of the helix-turn-helix motif and reveal a unique aspect of the class-3 flippases over

the otherwise conserved substrate transport mechanism of the P4 ATPases.

In summary, our systematic structural and biochemical studies of the class-3 Dnf1–Lem3 and

Dnf2–Lem3 complexes in five transport states provide the novel insight that the overall

substrate transport mechanism is similar to that of class-1 flippases and that it may be

conserved among all classes of the P4 ATPases. Our studies also show a helix-turn-helix motif

in the P domain that is unique to the class-3 enzymes and plays a crucial regulatory function

during the substrate transport cycle. Also unique to class-3 enzymes is the role of the beta-

subunit Lem3 in substrate binding and specificity. Thus, this work advances our mechanistic

understanding of the highly conserved and essential family of the P4 ATPases.

MATERIALS AND METHODS

Purification of the endogenous Dnf1p–Lem3p and Dnf2p–Lem3p complexes. Using a

PCR-based genomic epitope-tagging method (Funakoshi and Hochstrasser, 2009), we tagged

Dnf1p or Dnf2p with a C-terminal triple-FLAG in the yeast strain BY4741 (Horizon Discovery).

Cells were grown in YPD medium for about 20 h before harvest. Cells were resuspended in

lysis buffer (20 mM Tris-HCl, pH 7.4, 0.2 M sorbitol, 50 mM potassium acetate, 2 mM EDTA,

and 1 mM phenylmethylsulfonyl fluoride [PMSF]) and then lysed using a French press at 15,000

psi. The lysate was centrifuged at $10,000 \times g$ for 30 min at 4 °C. The supernatant was collected

and centrifuged at $100,000 \times g$ for 60 min at 4 °C. The membrane pellet was collected and then

resuspended in buffer A containing 10% glycerol, 20 mM Tris-HCl (pH 7.4), 1% DDM, 0.1%

CHS, 0.5 M NaCl, 1 mM $MgCl_2$, 1 mM $MnCl_2$, 1 mM EDTA, and 1 mM PMSF. After incubation

for 30 min at 4 °C, the mixture was centrifuged for 30 min at $120,000 \times g$ to remove insoluble

membrane. The supernatant was mixed with pre-washed anti-FLAG (M2) affinity gel at 4 °C

overnight with shaking. The affinity gel was then collected and washed three times in buffer B

containing 20 mM HEPES, pH 7.4, 150 mM NaCl, 0.01% LMNG, 0.001% CHS, and 1 mM

$MgCl_2$. The proteins were eluted with buffer B containing 0.15 mg/ml 3×FLAG peptide and was

further purified in a Superose 6 10/300 gel filtration column in buffer C containing 20 mM

HEPES, pH 7.4, 150 mM NaCl, 0.003% LMNG, 0.0003% CHS and 1 mM $MgCl_2$. Finally, the

purified proteins were assessed by SDS-PAGE gel and concentrated for cryo-EM analysis.

Approximate 20 μg Dnf1p–Lem3p and 40 μg Dnf2p–Lem3p can be purified from 18 L of cells.

Cryo-electron microscopy. To capture different states, the purified Dnf1p–Lem3p or Dnf2p–

Lem3p was mixed with different solutions for 1 h on ice: E1-ATP, 5 mM $MgCl_2$ and 2 mM

AMPPCP; E1P-ADP and E2P-transition, 5 mM $MgCl_2$, 5 mM NaF, 1 mM $AlCl_3$, and 5 mM ADP;

E2P, 5 mM $MgCl_2$, 10 mM NaF, and 2 mM $BeSO_4$. After an incubation, 2.5 μL aliquots of

purified Dnf1p–Lem3p or Dnf2p–Lem3p at a concentration of about 1.5 mg/mL were placed on

glow-discharged holey carbon grids (Quantifoil Au R2/2, 300 mesh) and were flash-frozen in

liquid ethane using a FEI Vitrobot Mark IV. Cryo-EM data was collected automatically with

SerialEM in a 300-kV FEI Titan Krios electron microscope with defocus values from −1.0 to −2.0

μm. The microscope was operated with a K3 direct detector at a nominal magnification of

130,000× and a pixel size of 0.413 Å per pixel. The dose rate was 8 electrons per $Å^2$ per s, and

the total exposure time was 8 s.

Cryo-EM image processing. Program MotionCorr 2.0 (Zheng et al., 2017) was used for motion

correction, and CTFFIND 4.1 was used for calculating contrast transfer function parameters

(Mindell and Grigorieff, 2003). All the remaining steps were performed using RELION 3 (Zivanov

et al., 2018). The resolution of the map was estimated by the gold-standard Fourier shell

correlation at a correlation cutoff value of 0.143.

For the Dnf1–Lem3 structure in the E2P state, we collected 4764 raw movie micrographs. A

total of 2,094,437 particles were picked automatically. After 2D classification, a total of

1,789,831 particles were selected and used for 3D classification. Based on the quality of the

four 3D classes, 590,043 particles were retained for further 3D reconstruction, refinement, and

postprocessing, resulting in a 2.8-Å average resolution 3D map.

For the Dnf1–Lem3 structure in the E1 state, we collected 4074 raw movie micrographs. A total

of 1,358,323 particles were picked automatically. After 2D classification, a total of 1,089,904

particles were selected and used for 3D classification. Based on the quality of the four 3D

classes, 473,761 particles were selected for further 3D reconstruction, refinement, and

postprocessing, resulting in a 3.1-Å average resolution 3D map.

For the Dnf1–Lem3 structure in the E1P-ADP state, we collected 6038 raw movie micrographs.

A total of 2,257,199 particles were picked automatically. After 2D classification, a total of

1,789,831 particles were selected and used for 3D classification. Based on the quality of the

four 3D classes, 716,510 particles were selected for further 3D reconstruction, refinement, and

postprocessing, resulting in a 3.2-Å average resolution 3D map.

For the Dnf2–Lem3 structure in the E1 state, we collected 1438 raw movie micrographs. A total

of 509,555 particles were picked automatically. After 2D classification, a total of 291,438

particles were selected and used for 3D classification. Based on the quality of the four 3D

classes, 190,753 particles were retained for further 3D reconstruction, refinement, and

postprocessing, resulting in a 3.1-Å average resolution 3D map.

For the Dnf2–Lem3 structure in the E1-AMPPCP state, we collected 3665 raw movie

micrographs. A total of 844,464 particles were picked automatically. After 2D classification, a

total of 519,537 particles were selected and used for 3D classification. Based on the quality of

the four 3D classes, 315,846 particles were selected for further 3D reconstruction, refinement,

and postprocessing, resulting in a 3.8-Å average resolution 3D map.

For the Dnf2–Lem3 structure in the E1P-ADP and E2P-transition states, we collected 5683 raw

movie micrographs. A total of 1,010,708 particles were picked automatically. After 2D

classification, a total of 593,683 particles were retained and used for 3D classification. Based on

the conformation of the six 3D classes, the E1P-ADP and E2P-transition states were separated.

For the E1P-ADP state, 266,218 particles were selected for further 3D reconstruction,

refinement, and postprocessing, resulting in a 4.0-Å average resolution 3D map. The Dnf1 A

domain was flexible and was not modeled. The N domain density was weak but clearly present.

This domain was modeled as a rigid body and was excluded from final atomic model

refinement. For the E2P-transition state, 127,442 particles were selected for further 3D

reconstruction, refinement, and postprocessing, resulting in a 4.0-Å average resolution 3D map.

For the Dnf2–Lem3 in the E2P state, we collected 3260 raw movie micrographs. A total of

1,095,601 particles were picked automatically. After 2D classification, a total of 1,065,397

particles were retained and used for 3D classification. Based on the quality of the four 3D

classes, 763,661 particles were selected for further 3D reconstruction, refinement, and

postprocessing, resulting in a 3.5-Å average resolution 3D density map.

Structural modeling, refinement, and validation. We first built the model of Dnf1–Lem3 in the

E2P state at 2.8-Å resolution. We generated the initial model based on the structure of Drs2–

Cdc50 (PDB ID 6PSY) by SWISSMODEL (https://swissmodel.expasy.org), and then manually

corrected in COOT and Chimera (Emsley et al., 2010; Pettersen et al., 2004). The complete

model of Dnf1–Lem3 was refined by real-space refinement in the PHENIX program and

subsequently adjusted manually in COOT. Using the model of Dnf1–Lem3 in E2P as reference,

models of Dnf1–Lem3 in the E1 and E1P-ADP states and of Dnf2–Lem3 in the E1, E1-ATP,

E1P-ADP, E2P transition, and E2P states were built and refined using COOT, Chimera, and

PHENIX. Finally, all models were validated using MolProbity in PHENIX (Adams et al., 2010;

Chen et al., 2010). Structural figures were prepared in Chimera (Pettersen et al., 2004) and

PyMOL (https://pymol.org/2/).

Yeast strains and plasmid construction. Strains and plasmid used in the study are listed in

Supplemental Tables I and II. All yeast culture reagents were purchased from Sigma-Aldrich

and BD Scientific. Yeast strains were grown in YPD or minimal selective media. Yeast

transformation was performed using the standard LiAC-PEG method. For plasmid shuffling

assays, 10-fold serial dilutions starting from a cell suspension with cell density of $OD_{600} = 1$ were

spotted on synthetic complete media (SD) and SD-5-FOA and incubated at 30 °C for 2 d before

imaging. DNA constructs and mutations were created by Gibson assembly and quick-change

mutagenesis.

Lipid uptake assay. All the lipids used in this study were purchased from Avanti Polar Lipids. NBD-lipid uptake was performed as previously described (Roland et.,2019). Cells were grown to the mid-log phase with a cell density of approximately OD_{600} = 1. Cells were collected for each strain. NBD-PC, NBD-PE, and NBD-GlcCer were dried and resuspended in 100% ethanol, then added to the designated ice-cold growth medium at a final concentration of 2 µg/mL. Cells were incubated on ice with the pre-chilled NBD-lipid+medium for 30 min. After the incubation, the cells were washed twice with pre-chilled SA medium (SD medium + 2% [w/v] sorbitol + 20 mM NaN_3) supplemented with 4% (w/v) fatty acid-free BSA. Cells were resuspended in SA with 5 µM propidium iodide (PI) and NBD-lipid uptake was immediately measured by flow cytometry. The order in which the samples were prepared and processed was varied in each experiment to reduce potential experimental error. Each experiment consisted of three independent biological replicates for each strain, repeated three times (n = 9).

Fluorescence microscopy. Cells were grown in minimal synthetic dropout medium to log phase. Cells were washed with fresh medium 3 times and resuspended in fresh SD medium. Cells were then mounted on glass slides and observed immediately at room temperature. Images were acquired using a DeltaVision Elite Imaging System (GE Healthcare Life Sciences, Pittsburgh, PA) equipped with a 100× objective lens followed by deconvolution using softWoRx software (GE Healthcare Life Science).

Supplementary Information is available in the online version of the paper.

ACKNOWLEDGEMENTS. Cryo-EM images were collected in the David Van Andel Advanced Cryo-Electron Microscopy Suite at Van Andel Institute. We thank Gongpu Zhao and Xing Meng for facilitating data collection and David Nadziejka for proofreading the manuscript. This work was supported by the U.S. National Institutes of Health (CA231466 to H.L. and GM107978 to T.R.G.) and the Van Andel Institute (to H.L.).

CONTRIBUTIONS. L.B., H.L., B.K.J. and T.R.G. conceived and designed the experiments. L.B., Q.Y., B.K.J., and A.K. performed the experiments. L.B., T.R.G., and H.L. analyzed the data. L.B., T.R.G, and H.L. wrote the manuscript with input from all authors.

DATA AVAILABILITY. The cryo-EM 3D maps and the corresponding atomic models of the

Dnf1–Lem3 complex have been deposited at the EMDB database and the RCSB PDB with the

respective accession codes of EMD-xxxx and yyyy (E1), EMD-xxxx and yyyy (E1P-ADP), EMD-

xxxx and yyyy (E2P). The cryo-EM 3D maps and the corresponding atomic models of the Dnf2–

Lem3 complex have been deposited at the EMDB database and the RCSB PDB with the

respective accession codes of EMD-xxxx and yyyy (E1), EMD-xxxx and yyyy (E1-ATP), EMD-

xxxx and yyyy (E1P-ADP), EMD-xxxx and yyyy (E2P), EMD-xxxx and yyyy (E2P-transition).

REFERENCES

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J.,

Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., *et al.* (2010). PHENIX: a comprehensive

Python-based system for macromolecular structure solution. Acta Crystallogr D Biol

Crystallogr *66*, 213-221.

Andersen, J.P., Vestergaard, A.L., Mikkelsen, S.A., Mogensen, L.S., Chalat, M., and Molday,

R.S. (2016). P4-ATPases as Phospholipid Flippases-Structure, Function, and Enigmas.

Front Physiol *7*, 275.

Axelsen, K.B., and Palmgren, M.G. (1998). Evolution of substrate specificities in the P-type

ATPase superfamily. J Mol Evol *46*, 84-101.

Bai, L., Kovach, A., You, Q., Hsu, H.C., Zhao, G., and Li, H. (2019a). Autoinhibition and

activation mechanisms of the eukaryotic lipid flippase Drs2p-Cdc50p. Nat Commun *10*,

4142.

Bai, L., Kovach, A., You, Q., Kenny, A., and Li, H. (2019b). Structure of the eukaryotic protein

O-mannosyltransferase Pmt1-Pmt2 complex. Nat Struct Mol Biol *26*, 704-711.

Baldridge, R.D., and Graham, T.R. (2012). Identification of residues defining phospholipid

flippase substrate specificity of type IV P-type ATPases. Proc Natl Acad Sci U S A *109*,

E290-298.

Baldridge, R.D., and Graham, T.R. (2013). Two-gate mechanism for phospholipid selection and

transport by type IV P-type ATPases. Proc Natl Acad Sci U S A *110*, E358-367.

Baldridge, R.D., Xu, P., and Graham, T.R. (2013). Type IV P-type ATPases distinguish mono-
versus diacyl phosphatidylserine using a cytofacial exit gate in the membrane domain. J Biol
Chem *288*, 19516-19527.

Best, J.T., Xu, P., and Graham, T.R. (2019). Phospholipid flippases in membrane remodeling
and transport carrier biogenesis. Curr Opin Cell Biol *59*, 8-15.

Bloch, J.S., Pesciullesi, G., Boilevin, J., Nosol, K., Irobalieva, R.N., Darbre, T., Aebi, M.,
Kossiakoff, A.A., Reymond, J.L., and Locher, K.P. (2020). Structure and mechanism of the
ER-based glucosyltransferase ALG6. Nature *579*, 443-447.

Bublitz, M., Morth, J.P., and Nissen, P. (2011). P-type ATPases at a glance. J Cell Sci *124*,
2515-2519.

Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J.,
Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom structure
validation for macromolecular crystallography. Acta Crystallogr D Biol Crystallogr *66*, 12-21.

Dyla, M., Basse Hansen, S., Nissen, P., and Kjaergaard, M. (2019). Structural dynamics of P-
type ATPase ion pumps. Biochem Soc Trans *47*, 1247-1257.

Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of
Coot. Acta Crystallogr D Biol Crystallogr *66*, 486-501.

Folmer, D.E., Elferink, R.P., and Paulusma, C.C. (2009). P4 ATPases - lipid flippases and their
role in disease. Biochim Biophys Acta *1791*, 628-635.

Funakoshi, M., and Hochstrasser, M. (2009). Small epitope-linker modules for PCR-based C-
terminal tagging in Saccharomyces cerevisiae. Yeast *26*, 185-192.

He, Y., Xu, J., Wu, X., and Li, L. (2020). Structures of a P4-ATPase lipid flippase in lipid
bilayers. Protein Cell *11*, 458-463.

Hiraizumi, M., Yamashita, K., Nishizawa, T., and Nureki, O. (2019). Cryo-EM structures capture
the transport cycle of the P4-ATPase flippase. Science *365*, 1149-1155.

Hua, Z., Fatheddin, P., and Graham, T.R. (2002). An essential subfamily of Drs2p-related P-
type ATPases is required for protein trafficking between Golgi complex and
endosomal/vacuolar system. Mol Biol Cell *13*, 3162-3177.

Hua, Z., and Graham, T.R. (2003). Requirement for neo1p in retrograde transport from the Golgi
complex to the endoplasmic reticulum. Mol Biol Cell *14*, 4971-4983.

Huang, Y., Takar, M., Best, J.T., and Graham, T.R. (2020). Conserved mechanism of
phospholipid substrate recognition by the P4-ATPase Neo1 from Saccharomyces
cerevisiae. Biochim Biophys Acta Mol Cell Biol Lipids *1865*, 158581.

Kanai, R., Ogawa, H., Vilsen, B., Cornelius, F., and Toyoshima, C. (2013). Crystal structure of a
Na+-bound Na+,K+-ATPase preceding the E1P state. Nature *502*, 201-206.

Martin, S., Smolders, S., Van den Haute, C., Heeman, B., van Veen, S., Crosiers, D., Beletchi,
I., Verstraeten, A., Gossye, H., Gelders, G.*, et al.* (2020). Mutated ATP10B increases
Parkinson's disease risk by compromising lysosomal glucosylceramide export. Acta
Neuropathol *139*, 1001-1024.

Mindell, J.A., and Grigorieff, N. (2003). Accurate determination of local defocus and specimen
tilt in electron microscopy. J Struct Biol *142*, 334-347.

Nakano, K., Yamamoto, T., Kishimoto, T., Noji, T., and Tanaka, K. (2008). Protein kinases
Fpk1p and Fpk2p are novel regulators of phospholipid asymmetry. Mol Biol Cell *19*, 1783-
1797.

Natarajan, P., Wang, J., Hua, Z., and Graham, T.R. (2004). Drs2p-coupled aminophospholipid
translocase activity in yeast Golgi membranes and relationship to in vivo function. Proc Natl
Acad Sci U S A *101*, 10614-10619.

Palmgren, M.G., and Axelsen, K.B. (1998). Evolution of P-type ATPases. Biochim Biophys Acta
1365, 37-45.

Paterson, J.K., Renkema, K., Burden, L., Halleck, M.S., Schlegel, R.A., Williamson, P., and
Daleke, D.L. (2006). Lipid specific activation of the murine P4-ATPase Atp8a1 (ATPase II).
Biochemistry *45*, 5367-5376.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and
Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research and
analysis. J Comput Chem *25*, 1605-1612.

Pomorski, T., Lombardi, R., Riezman, H., Devaux, P.F., van Meer, G., and Holthuis, J.C. (2003).
Drs2p-related P-type ATPases Dnf1p and Dnf2p are required for phospholipid translocation

across the yeast plasma membrane and serve a role in endocytosis. Mol Biol Cell *14*, 1240-
1254.

Post, R.L., Hegyvary, C., and Kume, S. (1972). Activation by adenosine triphosphate in the
phosphorylation kinetics of sodium and potassium ion transport adenosine triphosphatase. J
Biol Chem *247*, 6530-6540.

Roelants, F.M., Baltz, A.G., Trott, A.E., Fereres, S., and Thorner, J. (2010). A protein kinase
network regulates the function of aminophospholipid flippases. Proc Natl Acad Sci U S A
107, 34-39.

Roland, B.P., and Graham, T.R. (2016). Decoding P4-ATPase substrate interactions. Crit Rev
Biochem Mol Biol *51*, 513-527.

Roland, B.P., Jain, B.K., and Graham, T.R. (2020). Exofacial membrane composition and lipid
metabolism regulates plasma membrane P4-ATPase substrate specificity. bioRxiv.

Roland, B.P., Naito, T., Best, J.T., Arnaiz-Yepez, C., Takatsu, H., Yu, R.J., Shin, H.W., and
Graham, T.R. (2019). Yeast and human P4-ATPases transport glycosphingolipids using
conserved structural motifs. J Biol Chem *294*, 1794-1806.

Sikorski, R.S., and Hieter, P. (1989). A system of shuttle vectors and yeast host strains
designed for efficient manipulation of DNA in Saccharomyces cerevisiae. Genetics *122*, 19-
27.

Stevens, H.C., Malone, L., and Nichols, J.W. (2008). The putative aminophospholipid
translocases, DNF1 and DNF2, are not required for 7-nitrobenz-2-oxa-1,3-diazol-4-yl-
phosphatidylserine flip across the plasma membrane of Saccharomyces cerevisiae. J Biol
Chem *283*, 35060-35069.

Tang, X., Halleck, M.S., Schlegel, R.A., and Williamson, P. (1996). A subfamily of P-type
ATPases with aminophospholipid transporting activity. Science *272*, 1495-1497.

Timcenko, M., Lyons, J.A., Januliene, D., Ulstrup, J.J., Dieudonne, T., Montigny, C., Ash, M.R.,
Karlsen, J.L., Boesen, T., Kuhlbrandt, W.*, et al.* (2019). Structure and autoregulation of a
P4-ATPase lipid flippase. Nature *571*, 366-370.

Toyoshima, C., and Cornelius, F. (2013). New crystal structures of PII-type ATPases:
excitement continues. Curr Opin Struct Biol *23*, 507-514.

Toyoshima, C., Iwasawa, S., Ogawa, H., Hirata, A., Tsueda, J., and Inesi, G. (2013). Crystal
structures of the calcium pump and sarcolipin in the Mg2+-bound E1 state. Nature *495*, 260-
264.

UniProt, C. (2019). UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res *47*, D506-
D515.

van der Mark, V.A., Elferink, R.P., and Paulusma, C.C. (2013). P4 ATPases: flippases in health
and disease. Int J Mol Sci *14*, 7897-7922.

Vestergaard, A.L., Coleman, J.A., Lemmin, T., Mikkelsen, S.A., Molday, L.L., Vilsen, B., Molday,
R.S., Dal Peraro, M., and Andersen, J.P. (2014). Critical roles of isoleucine-364 and
adjacent residues in a hydrophobic gate control of phospholipid transport by the mammalian
P4-ATPase ATP8A2. Proc Natl Acad Sci U S A *111*, E1334-1343.

Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A. (2017).
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
microscopy. Nat Methods *14*, 331-332.

Zhou, X., and Graham, T.R. (2009). Reconstitution of phospholipid translocase activity with
purified Drs2p, a type-IV P-type ATPase from budding yeast. Proc Natl Acad Sci U S A *106*,
16586-16591.

Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and Scheres,
S.H. (2018). New tools for automated high-resolution cryo-EM structure determination in
RELION-3. Elife *7*.

Fig. 1. Structure of the *S. cerevisiae* class-3 lipid flippase Dnf1–Lem3 in the E2P state. A)
Cryo-EM 3D map of Dnf1–Lem3 in the E2P, in front (left) and back (right) views. The major
domains and motifs are labeled in different colors. **B**) Atomic model of Dnf1–Lem3 in the E2P
state in cartoon and colored as in panel A. The detergent and phospholipid molecules and the
covalently linked glycans on Lem3 are shown as sticks.

Fig. 2. The lipid binding sites and the putative substrate transport path of Dnf1–Lem3. A)
Cryo-EM 3D of the region around the putative substrate transporting path. The transmembrane
region is marked by the pink band. Three lipids densities are shown in grey. PL1 in the
exoplasmic side and PL2 in the cytosolic side of the membrane are highlighted by a purple and
a cyan box, respectively. **B**) Cartoon view of the lipid binding region in the Dnf1–Lem3 structure.
The purple box in panel A and the cyan box in panel B are enlarged in panels C and D,
respectively. Note that PL1 is about 10 Å below the lipid bilayer at the exoplasmic side, and PL2
and PL3 ingress into the cytosolic region by 10 Å. **C**) Close-up and top view of PL1 binding site.
D) Close-up view of PL2 binding site. **E**) Putative substrate transport path of Dnf1p, with the
residues lining the path and bound PL1 and PL2 shown as sticks. The hydrophobic gate in the
lipid transport path is highlighted by the yellow oval. PL3 is removed in the left panel for clarity.

Figure 3. Influence of lipid binding site mutations on Dnf1 transport activity and

localization. A) NBD-lipid uptake catalyzed by Dnf1 variants expressed on a *dnf1,2Δ* strain

background. **B**) NBD-lipid uptake catalyzed by Lem3 variants expressed on a *lem3Δ*

background, in which lipid transport is mediated primarily by Dnf2–Lem3. **C**) NBD-lipid uptake

catalyzed by Lem3 variants expressed on a *lem3Δ dnf2Δ* background to assay the influence of

Lem3 mutations solely on Dnf1 activity. In (A-C), cells were incubated with NBD-lipids for 30 min

and uptake was measured by flow cytometry. The value represents percentage uptake of NBD-

lipid in comparison to WT Dnf1 (A) or WT Lem3 (B-C) for each substrate lipid. The variance was

assessed with one-way ANOVAs, and comparison to WT was calculated with Tukey's post hoc

analysis. $n \geq 9$, ± SD, * indicates $p < 0.05$, **$p < 0.01$, ***$p < 0.001$. **D**) Localization of GFP-

tagged Dnf1 variants expressed in *dnf1,2Δ* cells. **E**) WT Dnf1-GFP co-expressed with Lem3

variants. Arrowheads indicate nuclear envelope/endoplasmic reticulum location. Scale bar = 2

μm (n = 40).

Fig. 4. The ATP-dependent transport cycle of Dnf1–Lem3 and Dnf2–Lem3. A) The five

structures of Dnf1 and Dnf2 in the E1 (apo), E1-ATP (AMPPCP), E1P-ADP (AlF$_4^-$-ADP), E2P-

transition (AlF_4^-), and E2P (BeF_3^-) states. The A domain in the E1, E1-ATP, and E1P-ADP

states is flexible and is shown here as a ribbon in the yellow rectangle to indicate its position.

The structures of the E1-ATP and E1P-ADP states within the black dashed box are almost

identical. The structures of the E2P and E2P-transition states are similarly grouped for the same

reason. Magenta arrows indicate the domain movement during the Post–Albers cycle in the

order E1 to E1-ATP to E1P-ADP to E2P/E2P-transition. **B**) Structural comparison of the

substrate binding sites in the E1, E1P-ADP, and E2P states. TMH1-2 in the E1 and E1P-ADP

states tilt with respect to that in the E2P state; this movement closes the PL1 site and opens the

PL2 site.

Fig. 5. Structural comparison between Dnf1 (color) and the ATP8A1 (grey) in the E1 (A), E1P-ADP (B), and E2P (C) states. Bottom panels are close-up views of the boxed regions in the top panels. These panels highlight the unique helix-turn-helix motif in the N domain of Dnf1, which interacts with different regions in different states.

719 **Supplemental Table 1. Cryo-EM data collection, refinement, and validations.**

	Dnf1–Lem3			Dnf2–Lem3				
	Apo	E1-ADP	E2P	Apo	E1-ATP	E1-ADP	E2P-transition	E2P
Data collection and processing								
Magnification (k)	130	130	130	130	130	130	130	130
Voltage (kV)	300	300	300	300	300	300	300	300
Electron exposure (e$^-$/Å^2)	64	64	64	60	60	60	60	60
Defocus range (-μm)	1.0-2.0	1.0-2.0	1.0-2.0	1.0-2.0	1.5-3.0	1.5-3.0	1.5-3.0	1.5-3.0
Pixel size (Å)	0.826	0.826	0.826	1.029	0.826	0.826	0.826	0.826
Symmetry imposed	C1	C1	C1	C1	C1	C1	C1	C1
Micrograph #	4,074	6,038	4,764	1,438	3,665	5683	5683	3260
Initial particle #	1,358,323	2,257,199	2,094,437	509,555	844,464	1,010,708	1,010,708	1,095,601
Final particle #	473,761	716,510	590,043	190,753	358,687	266,218	127,442	763,661
Map resolution (Å)	3.00	3.25	2.82	3.08	3.85	4.05	3.98	3.50
FSC threshold	0.143	0.143	0.143	0.143	0.143	0.143	0.143	0.143
Map resolution (Å)	3.12	3.25	2.82	3.08	3.85	4.05	3.98	3.50
Refinement								
Map sharpening B factor (Å^2)	116.5	178.9	116.2	102.5	195.8	243.3	197.8	180.0
Model composition								
Non-hydrogen atoms	11150	11235	13121	10369	10684	11960	12521	12879
Protein residues								
Ligands & glycans	1332	1322	1552	1247	1274	1471	1534	1545

	22	26	26	18	24	15	16	24
B factors (Å^2)								
Protein	81	39	47	86	48	175	166	58
Ligand	67	45	34	56	47	174	158	50
R.m.s. deviations								
Bond lengths (Å)	0.01	0.01	0.01	0.004	0.003	0.002	0.002	0.003
Bond angles (°)	0.8	0.9	1.1	0.6	0.6	0.6	0.6	0.6
Validation								
MolProbity score	2.4	2.5	2.0	2.1	2.1	2.1	2.0	2.1
Clashscore	13	11	11	14	12	12	13	13
Poor rotamers (%)	1	1	1	1	0	0	0	1
Ramachandran plot								
Favored (%)	93.0	91.9	95.6	93.0	90.7	93.1	93.9	92.9
Allowed (%)	7.0	8.1	4.4	6.9	9.2	6.9	5.9	7.1
Disallowed (%)	0	0	0	0	0	0	0	0

721 **Supplemental Table 2. List of plasmids used in the study.**

Plasmid	Source
pRS313	(Sikorski and Hieter, 1989)
pRS425	(Sikorski and Hieter, 1989)
pRS313-Dnf1	(Baldridge and Graham, 2012)
pRS313-Dnf1 R264A	
pRS313-Dnf1 Y633A	

pRS313-Dnf1 Y633G	
pRS313-Dnf1 T648A	
pRS313-Dnf1 T648V	
pRS313-Dnf1 W652A	
pRS313-Dnf1 W652S	
pRS313-Dnf1 Q610A	
pRS313-Dnf1 S611A	
pRS313-Dnf1 YQS-FSN	
pRS313-Dnf1 N1226A	
pRS313-Dnf1 (692-737)Δ	
pRS416-GFP-Dnf1	(Baldridge and Graham, 2012)
pRS416-GFP-Dnf1 R264A	
pRS416-GFP-Dnf1 W652A	
pRS416-GFP-Dnf1 W652S	
pRS416-GFP-Dnf1 Q610A	
pRS416-GFP-Dnf1 S611A	
pRS416-GFP-Dnf1 YQS-FSN	
pRS416-GFP-Dnf1 N1226A	
pRS425 Lem3	(Baldridge and Graham, 2012)
pRS425 Lem3 (2-49)Δ	
pRS425 Lem3 (400-414)Δ	
pRS425 Lem3 R51A	

Strain	Genotype	Plasmid	Source
BY4741	*MATa his3Δ1 leu2Δ0 ura3Δ0 met15Δ0*		Invitrogen & Horizon Discovery
PFY3275F	MAT**a** *his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 dnf1Δ dnf2Δ*		(Hua and Graham, 2003)
RBY9701	PFY3275F *lem3*Δ::natNT2		(Baldridge and Graham, 2012)
ZHY704	*MATα his3Δ1 leu2Δ0 ura3Δ0 lys2Δ0 dnf1Δ dnf2Δ dnf3Δ drs2Δ::LEU2* pRS416-*DRS2*	pRS416-*DRS2*	(Hua et al., 2002)
SCY119	*MATα his3Δ1 leu2Δ0 ura3Δ0 met15Δ0 lem3*Δ::KanMX		(Baldridge and Graham, 2012)
BJ201	PFY3275F	pRS313	
BJ204	PFY3275F	pRS313-Dnf1	
BJ209	PFY3275F	pRS313-Dnf1R264A	
BJ212	PFY3275F	pRS313-Dnf1Y633A	
BJ215	PFY3275F	pRS313-Dnf1Y633G	
BJ218	PFY3275F	pRS313-Dnf1T648A	
BJ219	PFY3275F	pRS313-Dnf1T648V	
BJ222	PFY3275F	pRS313-Dnf1W652A	
BJ225	PFY3275F	pRS313-Dnf1W652S	
BJ230	PFY3275F	pRS313-Dnf1Q610A	
BJ231	PFY3275F	pRS313-Dnf1S611A	

BJ234	PFY3275F	pRS313-Dnf1YQS-FSN	
BJ237	PFY3275F	pRS313-Dnf1N1226A	
BJ240	PFY3275F	pRS313-Dnf1(692-737)Δ	
BJ241	PFY3275F	pRS313-Dnf1(692-737)Δ	
BJ242	PFY3275F	pRS313-Dnf1(692-737)Δ	
BJ245	PFY3275F	pRS416-GFP-Dnf1	
BJ248	PFY3275F	pRS416-GFP-Dnf1 R264A	
BJ251	PFY3275F	pRS416-GFP-Dnf1 W652A	
BJ254	PFY3275F	pRS416-GFP-Dnf1 W652S	
BJ255	PFY3275F	pRS416-GFP-Dnf1 Q610A	
BJ260	PFY3275F	pRS416-GFP-Dnf1 S611A	
BJ263	PFY3275F	pRS416-GFP-Dnf1 YQS-FSN	
BJ266	PFY3275F	pRS416-GFP-Dnf1 N1226A	
BJ269	ZHY704	pRS313	
BJ272	ZHY704	pRS313-Dnf1	
BJ275	ZHY704	pRS313-Dnf1R264A	
BJ276	ZHY704	pRS313-Dnf1Y633A	
BJ279	ZHY704	pRS313-Dnf1Y633G	
BJ282	ZHY704	pRS313-Dnf1T648A	
BJ285	ZHY704	pRS313-Dnf1T648V	

BJ288	ZHY704	pRS313-Dnf1W652A	
BJ291	ZHY704	pRS313-Dnf1W652S	
BJ294	ZHY704	pRS313-Dnf1Q610A	
BJ297	ZHY704	pRS313-Dnf1S611A	
BJ300	ZHY704	pRS313-Dnf1YQS-FSN	
BJ303	ZHY704	pRS313-Dnf1N1226A	
BJ308	ZHY704	pRS313-Dnf1(692-737)Δ	
BJ309	RBY9701	pRS313, pRS425	
BJ310	RBY9701	pRS313-Dnf1, pRS425-Lem3	
BJ311	RBY9701	pRS313-Dnf1, pRS425-Lem3 (2-49)Δ	
BJ312	RBY9701	pRS313-Dnf1, pRS425-Lem3 (400-414)Δ	
BJ313	RBY9701	pRS313-Dnf1, pRS425-Lem3 R51A	
BJ314	SCY119	pRS416-GFP-Dnf1, pRS425	
BJ315	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3	
BJ316	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3 (2-49)Δ	
BJ317	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3 (400-414)Δ	
BJ318	SCY119	pRS416-GFP-Dnf1, pRS425-Lem3 R51A	

 Supplemental figures

Supplemental Figure 1. Purification of the Dnf1–Lem3 and Dnf2–Lem3 complexes. A) Gel
filtration profile of the Dnf1–Lem3 complex. **B**) The peak fraction in (A) was checked using the
Coomassie blue-stained SDS-PAGE gel. **C**) Gel filtration profile of the Dnf2–Lem3 complex. **D**)
Peak fraction in (C) was checked using the Coomassie blue-stained SDS-PAGE gel.

Supplemental Figure 2. Cryo-EM data processing and resolution estimation of Dnf1–Lem3 in the E2P state. A) A representative electron micrograph. **B**) Selected reference-free 2D class averages. **C**) Cryo-EM data processing procedure. **D**) Gold-standard Fourier shell correlation of two independent half 3D maps. **E**) Local resolution map of the final 3D map. **F**) Angular distribution of raw particles used in the final reconstruction of the 2.8-Å 3D map.

Supplemental Figure 3. Selected regions in the 3D map of Dnf1–Lem3 in the E2P state, superimposed on the atomic model.

Supplemental Figure 4. **Cryo-EM data processing and validation of the Dnf1–Lem3 in the E1 and E1P-ADP states. A**) Cryo-EM data processing procedure of Dnf1 in the E1 state. **B**) Gold-standard Fourier shell correlation of two independent half 3D maps in the E1 state. **C**) Cryo-EM data processing procedure of Dnf1 in the E1P-ADP state. **D**) Gold-standard Fourier shell correlations of two independent half 3D maps of Dnf1p in the E1P-ADP state. In (**A**) and (**C**), the bottom panel shows the final map colored according to the local resolution.

Supplemental Figure 5. **Cryo-EM data processing and resolution estimation of the Dnf2–Lem3 in the E1 and E1-ATP states. A**) Cryo-EM data processing procedure of Dnf2 in the E1 state. **B**) Gold-standard Fourier shell correlations of two independent half 3D maps in the E1 state. **C**) Cryo-EM data processing procedure of Dnf2in the E1-ATP state. **D**) Gold-standard

Fourier shell correlations of two independent half maps of the Dnf2p-E1-ATP. In (**A**) and (**C**), the
bottom panel shows the final map colored according to the local resolution.

Supplemental Figure 6. Cryo-EM data processing and validation of the Dnf2–Lem3 in the
E2P-transition state and in the E1P-ADP state. A) Cryo-EM data processing procedure of the
Dnf2–Lem3 datasets in the E2P-transition and the Dnf2p-E1P-ADP states. The two bottom
panels show the two 3D maps colored according to their respective local resolutions. **B**) Gold-
standard Fourier shell correlations of two independent half 3D maps of Dnf2–Lem3 in the E2P-
transition state and in the E1P-ADP state.

Supplemental Figure 7. Cryo-EM data processing and validation of the Dnf2–Lem3 in the E2P state. A) Cryo-EM data processing procedure. The final 3D map shown at the bottom is colored coded by the local resolution. **B**) Gold-standard Fourier shell correlation of two independent half 3D maps.

Supplemental Figure 8. Structural alignments between the Dnf1 and Dnf2 in their

corresponding E1 (apo) (**A**), the E1P-ADP (**B**), and the E2P states (**C**) show a high similarity.

Supplemental Figure 9. The N-terminal peptide of Lem3p is partially ordered in the 3D

map of Dnf1–Lem3 in the E2P state. A) Cryo-EM 3D map of the Dnf1–Lem3 in the E2P state.

The ordered Lem3 N-terminal peptide is highlighted by the black and red boxes. **B**) An

enlargement of the black box in panel A, shown at a lower display threshold to show the weaker

density. **C**) An enlarged view of the red box in panel A. The Lem3 N-terminal peptide interacts

with the Dnf1 A domain. The interface is highly positive. **D**. Sequence alignment of yeast Lem3

(ScLem3) and other CDC50 family proteins (yeast Crf1, yeast Cdc50, human CDC50A, and

human CDC50B) shows the Lem3-specific, N-terminal, negatively charged motif.

Supplemental Figure 10. Structural comparison between Dnf1 (color) and the ATP8A1

(grey) in the E2P state. Lipids in the Dnf1 structure are shown in red, while the only lipid in

ATP8A1 is in blue. The PL binding sites in the exoplasmic side of membrane of the two

flippases are well aligned. The right panel is an enlargement of the substrate transport path of

the left panel.

Supplemental Figure 11. Ability of Dnf1 variants to support viability of flippase-deficient

cells. *HIS3*-marked plasmids (pRS313) carrying the indicated Dnf1 variants were used to

transform a *dnf1,2,3Δdrs2Δ* pRS416-*DRS2* strain. Strains were spotted onto minimal media

plates (SD) selecting for both plasmids, and pRS416-*DRS2* was counter-selected on medium

containing 5-fluoroorotic acid (SD-FOA). Strains were incubated at 30 °C for 2 d and imaged.

The growth phenotype on SD-FOA is an indicator of how well the Dnf1 variant functions *in vivo*.

Supplemental Figure 12. High structural similarity between Dnf1 and Dnf2 in the corresponding substrate transport states. A) Structural alignments 3D maps of Dnf2–Lem3 in the E1-ATP and E1P-ADP states. **B**) Structural alignments of 3D maps of Dnf2 in the E2P and E2P-transition states.

Supplemental Figure 13. The N and A domain densities are weak in the 3D maps of Dnf1–Lem3 in the E1 and E1P-ADP states, indicating their high flexibility. A) Cryo-EM 3D map of the Dnf1-Lem3 in the E1 state displayed at a high (top panel) and a low threshold (bottom panel). Red arrows indicate the domain movement from the E1 state to the E1-ATP state. **B**) Cryo-EM 3D map of the Dnf1–Lem3 in the E1-ATP state displayed at a high (top panel) and a low threshold (bottom panel). Red arrows indicate the domain movement from the E1P-ADP state to the E2P state. **C**) Cryo-EM 3D map of the Dnf1–Lem3 in the E2P state. **D-F**, Selected reference-free 2D class averages of the Dnf1–Lem3 in the E1 state (**D**), in the E1P-ADP state (**E**), and in the E2P state (**F**). The averages in red or green boxes belong in the same views. The A domain locations are labeled by red arrowheads. The weak and blurry A domain density indicates its flexibility in the E1 and E1P-ADP states.