1 **Effect of relative social rank within a social hierarchy on neural activation in response to**
2 **familiar or unfamiliar social signals**
3

4 Won Lee[1,3], Hollie N. Dowd[2], Cyrus Nikain[1], Madeleine F. Dwortz[3], Eilene D. Yang[1], James P.
5 Curley [3] [*]
6
7 1 – Department of Psychology, Columbia University, New York, New York USA
8 2 – Department of Psychology, Barnard College, New York, New York USA
9 3 – Department of Psychology, University of Texas, Austin, Texas USA
10
* Corresponding author – curley@utexas.edu

Abstract
Competent social functioning of group-living species relies on the ability of individuals to detect
and utilize conspecific social cues to guide behavior. In many species, cues about an
individual's identity, including social status, are conveyed through urinary proteins. Previous
studies have identified numerous brain regions involved in processing these external cues,
collectively referred to as the Social Decision-Making Network. However, how the brain encodes
social information with respect to an individual's internal social status has not been thoroughly
examined. Here we show that neuronal activity in male mice, as indicated by cFos
immunoreactivity, is modulated by the social status and familiarity of the cue source and the
subject's own social status. The posteroventral medial amygdala was the only brain region that
responded exclusively to the social status of the cue source. Activity in other brain regions was
modulated by social status. Specifically, within the brain of dominant subjects there was robust
differential activity dependent on the status or familiarity of the cue source. Our results suggest
that individuals perceive social cues differently depending on their social experience. These
data inform further the investigation of neurobiological mechanisms underlying social-status
related brain differences and how social status-associated differences influence physiology and
behavior.

Introduction

        For social species living in groups, the ability to competently interact with others is vital for an individual's survival and reproductive fitness[1]. Appropriate behavioral responses require accurate social decision making, whereby animals evaluate their external environment and integrate salient social information with their current internal physiological status as well as contextual and social memories[2–5]. Social decision making should be thought of as a reciprocal process between two or more individuals, with one individual's behavioral output serving as input to another individual, that invokes rapid updating of information[3]. Brain circuits involved in social decision making that are conserved across social vertebrates constitute the Social Decision Making Network (SDMN)[5–8]. Studies in rodents have highlighted the role that subregions of the SDMN play in regulating social decisions in various social contexts including reproduction, parental care, aggression[3,9–13]. However, the functional role of the SDMN in the context of ongoing reciprocal group-level interactions has yet to be extensively explored.

        A social dominance hierarchy is a common social context that necessitates constant social decision making. The establishment and maintenance of social hierarchies requires individuals to learn and remember their experience with each social partner in the group[14] and dynamically express dominant or subordinate behavior towards animals of relatively lower or higher rank, respectively. Individuals must also remain vigilant to general social context changes, such as the presence or absence of high ranking individuals or alterations to group composition[15–19]. Despite these seemingly difficult cognitive requirements, numerous social species including fish, rats, mole rats, and mice are capable of rapidly organizing themselves into a hierarchy. Studies in primates, fish and rodents have recapitulated social hierarchies in the laboratory[17–19]. Such paradigms provide researchers with the unique opportunity to study the biological correlates of social behavior within an ethologically relevant context. Our lab has developed a paradigm that consistently generates autonomous hierarchies of 12 outbred CD-1 male mice[20,21]. We have shown that these mice exhibit flexible and context-appropriate social behaviors and adjust their behavioral response rapidly[15,22–24]. For example, individuals inhibit aggressive behavior in the presence of the alpha male[15], but are able to quickly ascend in social rank when the alpha male is removed from the group[12,17,19]. Furthermore, we have shown that altering the general social context (e.g. by removing the alpha male), is associated with immediate early gene activation throughout SDMN nodes.

        In mice, information about social context is often relayed through chemosensory cues present in urine. These cues can convey information relating to reproductive condition, dominance status, familiarity, and individual identity with exposure to these cues being shown to guide neural activity and behavior[25–27]. For instance, female mice show differential patterns of immediate early gene activation when exposed to dominant versus subordinate male urinary odors even when they have no previous experience with the urine donors[28]. Male mice scent mark over unfamiliar urinary odors more than familiar ones and show different patterns of immediate early gene expression when presented with familiar versus unfamiliar subjects[29,30]. There is also evidence across species that individuals respond differently to social cues depending upon individual factors such as their sex, hormonal state, past social experiences and social status[3,31]. For example, the rate of counter scent-marking over unfamiliar male urine is higher in dominant male mice than subordinates[32,33]. In cichlids, dominant male fish show a higher percentage of activated neurons when presented with food or female cues while subordinate males respond more sensitively towards dominant male odors[31]. These differential responses to social cues according to their own social status appear to be plastic adaptive responses. For instance, alpha males are required to distinguish between familiar and unfamiliar cues as they defend their territories from unknown conspecifics[34,35]. Conversely, mid-ranked or subordinate individuals should be able to recognize dominant or unfamiliar individuals and display attention or show fear-related responses to these individuals, while they should show less aversion towards social cues from familiar subordinate individuals. Indeed, there is strong

evidence that subordinate animals pay more attention to the behavior and cues of dominant
individuals than dominant individuals do to the behavior and cues of subordinates[15,36,37].
In the present study, we investigate how an individual's own internal state (social status)
modulates the neural activity of their SDMN in response to olfactory cues of conspecifics of
differing social status and familiarity. Specifically, we examine how differences in internal social
state (dominant versus subordinate social status) are associated with activation of the
immediate early gene (IEG) cFos in the brains of male mice in response to exposure to urinary
odors. Subjects were exposed to one of four different cue types: familiar dominant, familiar
subordinate, unfamiliar dominant or unfamiliar subordinate. Familiar urine donors originated
from the same social hierarchies as the subject, whereas unfamiliar urine donors originated
from different social hierarchies in separate housing systems. While we mainly focused on
SDMN brain regions, we also examined immunoreactivity in several other brain regions related
to stress responsivity and those regions previously described to have high sensitivity to different
social cues[12,28,30].
94
95 **Results**
96
Social hierarchy characteristics
All 8 social groups of 12 male mice formed significantly linear social hierarchies (Supplemental
Fig. S1). By the end of the group housing period, the median modified Landau's h-value and
interquartile range of 8 social groups (IQR) was 0.69 [0.63 – 0.71] (all p<0.01; see
Supplemental **Table S1** for values for each cohort). Triangle transitivity was 0.88 [0.84 – 0.93]
(all p<0.001) and directional consistency 0.89 [0.87 – 0.93] (all p<0.001). As noted in the
methods (see **Fig. 1.** for experimental design), urine samples were collected from group
housing between days 7-10, after the establishment of social hierarchies. All social groups in
this study established linear and stable social hierarchy by the 6th day of group housing (**Fig. 2**;
median 4 days [2.8 – 5.3]).
107
Figure 7.1: Schematic of the social stimulus exposure experimental design. 8 groups of 12
male CD-1 mice were housed in an ethologically relevant group housing system. All 8 groups
formed linear social hierarchies by the start of urine collection from the donors (alpha (Rank 1)
or the most subordinate (Rank 12) individual). On the last day of group housing (Day 10-12), all
mice in social groups excluding the urine donor mouse were exposed to either familiar or
unfamiliar urine from either alpha or the most subordinate males.

Figure 7.2: The triangle transitivity and Landau's Modified h-value of each social group across the group housing period. All 8 cohorts formed linear and stable social hierarchies by day 6 of group housing. Each line represents a different social hierarchy. Red lines indicate significant values.

120
121

Fos-immunoreactivity in response to social cues in dominant and subordinate male mice

For each brain region, we used Bayesian hierarchical modeling[38] to estimate posteriors of 12 planned comparisons (**Table 1**, **Fig. 3**, Supplemental **Fig. S2**) to examine the effects of a) internal status – dominant vs. subordinate social status in the hierarchies; b) familiarity of social cues – familiar vs. unfamiliar social cues; c) the source of social cue – from the alpha or the most subordinate males in the hierarchies.

Among 23 analyzed regions, the MEApv was the only region for which all individuals responded with significantly higher cFos immunoactivity to alpha male urine compared to subordinate urine regardless of the subject's own social status or the familiarity of the cue. In the VMHdm/c and VMHvl, alpha male urine induced significantly higher cFos immunoreactivity than subordinate urine for subordinate subjects exposed to familiar or unfamiliar urine, but only for dominant subjects when exposed to unfamiliar urine. Additionally, both dominant and subordinate animals showed a significant increase in cFos immunoreactivity in the PMv when exposed to familiar alpha urine compared to familiar subordinate urine but did not show any differences when exposed to unfamiliar alpha versus subordinate urine. Other brain regions showed context-dependent differences in cFos immunoreactivity when comparing exposure to alpha versus subordinate urine. For instance, both dominant and subordinate mice showed an increase in cFos immunoreactivity in the vlPAG when exposed to alpha versus subordinate urine. Conversely dominant mice showed an increase in vlPAG and dlPAG immunoreactivity when exposed to subordinate urine compared to alpha urine. Two brain regions, the mPFC and dmPAG showed increases in cFos immunoreactivity only in dominant animals that were exposed to unfamiliar subordinate urine compared to unfamiliar dominant urine. In the AH, dominant mice showed increase cFos responses to familiar alpha versus familiar subordinate urine, whereas subordinates showed an increased neural response to unfamiliar alpha versus unfamiliar subordinate urine. When considering the familiarity of the urine cue, the vlPAG showed the biggest effect. Both dominant and subordinate mice showed significantly increased responses to unfamiliar alpha urine compared to familiar alpha urine. Dominant mice also show increased vlPAG cFos immunoreactivity in response to familiar dominant urine compared to familiar subordinate urine. Further, dominant, but not subordinate, mice exhibited increase in

cFos immunoreactivity when exposed to familiar alpha compared to unfamiliar alpha urine in the PMv, dmPAG and AH. Dominant mice also showed increase responses to unfamiliar subordinate urine compared to familiar subordinate urine in the mPFC and AH, whereas the VMHvl had the opposite pattern with dominant mice responding significantly more to familiar subordinate urine compared to unfamiliar subordinate urine. It is also worth noting that subordinate males responded with significantly increased cFos immunoreactivity to unfamiliar alpha urine compared to familiar alpha urine in the mPFC.

When comparing differences in the responses between dominant and subordinate animals to urine cues, effects were highly dependent upon the context of the social cue as already outlined above. Further, the strongest effects observed were that subordinate animals exhibited a more robust response to familiar alpha urine than dominant animals in the LH, vlPAG, and dlPAG. Subordinates also responded more robustly than dominants to unfamiliar alpha urine in the PMv, dmPAG, dlPAG, AH and mPFC. When exposed to subordinate urine, subordinates responded more than dominants in the mPFC and AH, but dominants responded more than subordinates in the VMHvl and vlPAG. Only one region, the PMv showed a differential response between dominant and subordinate mice exposed to unfamiliar subordinate urine, with subordinates showing a significantly greater response compared to dominants.

We next applied Flexible discriminant analysis (FDA) using data from all brain regions from all experimental conditions to classify similar patterns of brain responses into eight groups, and to identify which brain regions drove the strongest differences in responses between groups. FDA classified the eight experimental groups correctly with seven discriminant functions. The first two functions explained 43.67% and 21.72% of the classification accuracy respectively (the corresponding spread of each data point is plotted in **Fig.4**). Function 1 was mostly driven by mPFC (factor loading coefficient for Function 1: -1.53), MEApv (0.97), and AH (0.95). Function 1 appears to be primarily driven by being exposed to alpha urine cues compared to subordinate urine cues of any familiarity. Function 2 was loaded heavily with vlPAG (-1.52), AH (1.04), BNSTpr (0.73), VMHdm/c (-0.72). Function 2 was heavily driven by the difference between being exposed to familiar alpha or familiar subordinate urine in dominant individuals. The coefficient values for each brain regions for all seven functions are listed in Supplemental **Table S2**. The same analysis only including SBN nodes (AH, BNSTpr, dlPAG, dmPAG, LSv, mPOA, MEApd, MEApd, vlPAG, VMHvl, VMHdm/c) did not yield as accurate a prediction as when we included data from all brain regions (Supplemental **Fig. S3**).

Table 7.1: Estimates and 95% highest density intervals (HDI) of 12 planned pairwise comparisons for cFos immunoreactivity in each brain regions. Bolded cells represent comparisons of which HDI did not overlap with 0.

Region	Alpha urine		Sub urine		Alpha urine		Sub urine		Fam		Unfam	
	Fam	Unfam	Fam	Unfam	Dom	Sub	Dom	Sub	Status: Dom	Sub	Dom	Sub
	Internal state: Dominant status vs. subordinate status				Familiarity: Familiar cue vs. Unfamiliar cue				Source: Alpha urine cue vs. Sub urine cue			
mPFC	-37 [-89, 21]	**-110 [-161, -53]***	**-114 [-174, -51]***	34 [-30, 97]	-2 [-51, 50]	**-74 [-132, -17]***	**-142 [-211, -80]***	4 [-59, 63]	32 [-26, 86]	-50 [-109, 8]	**-117 [-175, -58]***	22 [-34, 84]
NAcc	0 [-14, 12]	-1 [-16, 10]	0 [-27, 11]	0 [-12, 25]	0 [-10, 14]	0 [-13, 14]	-1 [-28, 10]	0 [-13, 2 4]	-1 [-22, 9]	-1 [-31, 6]	-14 [-35, 3]	-1 [-25, 8]
LSv	0 [-69, 46]	-1 [-72, 38]	0 [-55, 56]	0 [-57, 56]	0 [-51, 65]	0 [-53, 53]	0 [-51, 64]	0 [-49, 61]	-2 [-105, 24]	-1 [-83, 28]	-2 [-99, 25]	-2 [-80, 32]
BNSTpr	13 [-4, 35]	**7 [0, 12]***	-15 [-28, 2]	15 [-3, 29]	11 [-6, 32]	3 [-3, 9]	-14 [-30, 3]	14 [-2, 27]	13 [-5, 35]	-17 [-29, 1]	-13 [-27, 4]	-5 [-9, 1]
mPOA	-14 [-33, 5]	1 [-11, 29]	0 [-19, 27]	1 [-27, 59]	-6 [-30, 10]	10 [-6, 32]	0 [-48, 31]	0 [-29, 30]	-12 [-42, 7]	0 [-19, 19]	0 [-59, 22]	-1 [-41, 16]
AH	-20 [-57, 15]	**-52 [-95, -12]***	**-57 [-103, -8]***	36 [-3, 76]	**48 [8, 87]***	16 [-23, 54]	**-48 [-93, -1]***	**47 [5, 85]***	**65 [17, 108]***	27 [-12, 64]	-35 [-72, 8]	**57 [15, 97]***
MEApd	0 [-14, 19]	0 [-23, 14]	0 [-32, 10]	0 [-22, 20]	0 [-17, 17]	0 [-24, 12]	0 [-28, 13]	0 [-16, 24]	0 [-12, 26]	0 [-23, 12]	0 [-19, 19]	0 [-16, 24]
MEApv	-15 [-86, 59]	-50 [-128, 33]	-3 [-60, 58]	6 [-45, 60]	-27 [-99, 52]	-60 [-134, 22]	17 [-44, 73]	25 [-28, 77]	**73 [5, 139]***	**86 [16, 150]***	**107 [39, 179]***	**165 [92, 235]***
BLA	0 [-16, 18]	0 [-25, 11]	0 [-10, 28]	0 [-10, 30]	0 [-17, 19]	0 [-24, 10]	0 [-16, 22]	0 [-12, 21]	0 [-31, 10]	0 [-17, 15]	0 [-27, 11]	0 [-9, 28]
CEA	4 [-4, 14]	-4 [-11, 5]	0 [-13, 16]	6 [-10, 22]	5 [-6, 14]	-4 [-11, 3]	5 [-11, 24]	10 [-3, 23]	-13 [-29, 2]	**-18 [-28, -5]***	-11 [-28, 2]	-2 [-13, 7]
VMH dm/c	**-18 [-33, -2]***	-4 [-21, 13]	7 [-3, 17]	-2 [-9, 6]	-13 [-27, 2]	1 [-16, 19]	8 [-2, 18]	-1 [-8, 7]	4 [-7, 15]	**29 [14, 44]***	**25 [11, 39]***	**28 [13, 41]***
VMH vl	-14 [-33, 4]	-8 [-29, 15]	**8 [0, 16]***	-8 [-17, 1]	-17 [-36, 3]	-8 [-31, 12]	**12 [5, 19]***	-4 [-13, 5]	-1 [-10, 10]	**22 [5, 40]***	**27 [9, 47]***	**29 [9, 46]***
CA1	0 [-25, 26]	-9 [-42, 11]	-1 [-31, 16]	0 [-30, 24]	0 [-25, 28]	-9 [-40, 10]	-12 [-42, 11]	-6 [-35, 16]	17 [-7, 51]	11 [-8, 37]	1 [-20, 32]	12 [-10, 45]
CA3	0 [-11, 8]	0 [-7, 12]	0 [-11, 8]	0 [-11, 9]	0 [-14, 6]	0 [-9, 8]	0 [-11, 10]	0 [-10, 9]	0 [-9, 11]	0 [-7, 10]	0 [-6, 15]	0 [-8, 11]
DG	0 [-8, 9]	0 [-6, 12]	0 [-13, 6]	0 [-9, 10]	0 [-12, 7]	0 [-8, 8]	0 [-14, 6]	0 [-8, 9]	0 [-6, 15]	0 [-8, 9]	0 [-6, 14]	0 [-7, 10]
dlPAG	**-13 [-22, -4]***	**-17 [-31, -3]***	-2 [-15, 9]	-11 [-24, 1]	2 [-7, 12]	-1 [-15, 11]	7 [-4, 17]	-2 [-15, 12]	**-11 [-20, -1]***	0 [-12, 11]	-6 [-17, 3]	-1 [-16, 14]
dmPAG	1 [-8, 12]	**-9 [-14, -1]***	-2 [-10, 7]	-3 [-12, 4]	**13 [3, 23]***	4 [-4, 11]	2 [-6, 9]	0 [-9, 9]	3 [-6, 13]	0 [-8, 10]	**-8 [-15, 0]***	-4 [-11, 4]
vlPAG	**-106 [-171, -43]***	13 [-62, 95]	**64 [1, 126]***	-50 [-155, 52]	**-217 [-250,-174]***	**-90 [-188, -2]***	**128 [20, 217]***	2 [-70, 77]	**-149 [-207, -89]***	20 [-46, 88]	**185 [94, 272]***	**122 [17, 215]***
PMd	-10 [-37, 14]	15 [-7, 43]	5 [-11, 27]	13 [-6, 32]	-14 [-47, 10]	9 [-9, 34]	10 [-11, 32]	17 [-2, 33]	-11 [-40, 11]	1 [-14, 25]	11 [-9, 40]	10 [-8, 28]
PMv	12 [-34, 61]	**-56 [-91, -16]***	3 [-30, 30]	**-40 [-75, -7]***	**75 [30, 112]***	2 [-43, 48]	16 [-19, 49]	-27 [-56, 3]	**78 [35, 117]***	**64 [21, 107]***	22 [-13, 57]	35 [-1, 71]
PVN	0 [-11, 25]	0 [-17, 19]	0 [-20, 17]	0 [-15, 23]	0 [-22, 15]	0 [-26, 11]	0 [-18, 20]	0 [-12, 26]	0 [-23, 15]	0 [-30, 8]	0 [-18, 19]	0 [-15, 22]
LH	**-47 [-82, -7]***	-27 [-57, 6]	-43 [-89, 11]	17 [-23, 55]	-7 [-45, 27]	11 [-19, 40]	-5 [-49, 31]	47 [-4, 97]	-40 [-81, 4]	-31 [-78, 14]	-37 [-74, 0]	1 [-31, 39]
DMH	0 [-28, 13]	0 [-27, 14]	0 [-17, 25]	0 [-13, 32]	0 [-17, 23]	0 [-17, 24]	0 [-23, 20]	0 [-16, 24]	0 [-29, 15]	0 [-16, 23]	0 [-34, 10]	0 [-17, 24]

187 **Figure 7.3: Fos-immunoreactivity in selected nine brain regions in response to urine from familiar alpha, unfamiliar alpha, familiar**
188 **subordinate, unfamiliar subordinate males in dominant and subordinate male mice.** Boxplots show median (horizontal bars), IQR
189 (boxes) and 95% CI (whiskers). Raw data points of each group are also shown on the right side of each box plot. See Supplemental **Fig. S2**
190 for the data of all 23 brain regions.

191
192

Figure 7.4: Flexible discriminant analysis of the eight experimental groups (status x
familiarity x cue origin) as a function of cFos cell counts in all brain regions. Rhombi
represent group centroids of each experimental group. Grey lines and represent the degree of
separation (factor loading coefficient) each brain region contribute to Function 1 and Function 2.
The longer the line is the larger contribution that brain has to for each function. For legibility of
the entire plot, the grey lines were plotted with three times multiplication of the original (x, y)
coordinates and only those brain regions that are most contributing to each function are labeled.

Discussion

In this study we demonstrate that numerous brain regions respond to contextual features

of social odor cues, depending on the internal state, or social status, of the subject processing

the odor. We replicated our lab's behavioral paradigm that generates linear social hierarchies
within an ethologically relevant housing system. Groups of 12 outbred CD-1 mice were co-
housed and allowed to establish stable linear social hierarchies and do so within a median of 4
days. Then with cFos immunohistochemistry, we identify activity patterns across 23 brain
regions following exposure to a urinary odor cue. In total we examined eight different cue
conditions. In each condition, we varied three key features of the cue presentation: 1) the social
status of urine donors; 2) the familiarity of the urine source; 3) the subject's own internal state
(social status). This results in eight different treatment groups (four social cue types for
dominant and subordinate subjects). For each of the 23 brain regions we examined, we used
pair-wise comparisons for each unique combination of these key features.
FDA analysis categorized subjects based on collective neural activation in all 23 brain
regions. We were able to classify subjects into each experimental group with 100% accuracy,
indicating that collective brain activity from these brain regions is sufficient to determine which
group each subject originated from. In other words, we are able to predict the social status of an
individual and the type of odor cue they were exposed to based on their brain activity pattern.
The first two functions described by the FDA provide the strongest classification. Function 1
primarily discriminates between subjects that were exposed to urine from an alpha versus
subordinate mouse, and mostly ignored familiarity and social status. Function 2 most efficiently
discriminates between dominant subjects that were exposed to urine from a familiar alpha
versus subordinate mouse. It should be noted that Function 2 does not separate subordinate
subjects as efficiently. **Fig. 4**, which plots scores for Function 1 against Function 2 demonstrates
that there is no overlap of dominant subjects across the four social cue conditions, whereas
there is overlap between subordinate subjects exposed to each of the four social cue conditions.
This suggests that brain-wide neural activation patterns induced by various social odors are
more subtle in subordinate individuals and that discrimination of social cues relies on collective
changes across the multiple brain regions. We hypothesize that Function 2 may represent level
of saliency or the level of social vigilance towards given cues.
Notably, the FDA analysis is most successful when using data from all 23 brain regions
rather than just brain regions associated with the SDNM. This demonstrates the importance of
brain regions beyond this network for the differential processing of social cues from dominant
and subordinate animals. However, five out of the six brain regions (BNSTpr, AH, MEApv, VMH
dm/c, and vlPAG) that contribute most to the first two discrimination functions are also five of the
six nodes of the social behavior network (SBN)[6,7]. The SBN in conjunction with the mesolimbic
dopamine system comprises the SDMN and is considered to be an evolutionarily conserved set
of interconnected brain regions whose coordinated activity drives social decision making and
social behavior[5,7,39,40]. Consistent with our findings, other studies have also demonstrated that
differential patterns of whole brain activity can be used to classify subjects during social
discrimination tasks. For example, immediate early gene studies in African cichlid fish have also
used FDA to identify differential patterns of neural activity throughout the SDMN in response to
specific social contexts (e.g. exposure to dominant male, non-receptive female, aggressive
gravid female, juveniles, etc.)[41,42]. Similarly, differential patterns of neural activity in the SDMN,
olfactory bulb and mPFC can accurately predict whether male mice were exposed to novel or
familiar juvenile mice or an empty cage[30].
247
Processing of Alpha versus Subordinate Olfactory Cues
Many animals, especially mice, use urinary odors to convey information about their identity,
including social dominance[25,26,29,41,43]. The urine of male mice contains various chemosensory
cues such as ESP1 (a pheromone shown to enhance female sexual receptivity) and major
urinary proteins (MUPs)[44,45]. Dominant males show higher levels of MUPs, especially MUP20
(darcin)[27,46]. Our lab has previously shown that alpha CD-1 males living in social hierarchies of
12 mice produce higher levels of total MUPs and MUP20 (Lee et al., 2017). MUPs and MUP20

are used by dominant males to mark their territory and to signal dominance to other males, as
well as to attract females[23,27,34,45]. Thus, the ability to distinguish between urine from a dominant
versus a subordinate mouse is of high adaptive value in the context of territory marking, finding
a mate and evaluating the social environment. Function 1 of the FDA analysis distinguishes
whether the urine cue was from an alpha or subordinate mouse based on brain-wide activity
patterns. This is predominantly driven by responses in the MEApv, mPFC, and AH. That is,
these regions appear to be tuned to the dominance information conveyed by the urine cue. In
our pairwise analysis, the MEApv is the only region to show differential activation based on the
social status of the urine source alone, regardless of the familiarity with the source or the
subject's own social status. Specifically, this region shows greater activation in response to
urine from alpha mice. The MEA receives olfactory information directly from both MOB and AON
pathways[47]. Considering this region's sensitivity to purely the dominance of the cue source, it is
reasonable to hypothesize that it represents an early stage in an odor processing stream and is
responsible for parsing out the most critical information first. In line with this idea, we also
observe more refined responses to features of the cue presentation in downstream targets of
the MEA, including the VMHvl and PMV. Prior studies have shown increased neural activity in
MEApv neurons projecting to VMHdm in mice exposed to predator cat odor[48,49]. Indeed, it has
been generally suggested that MEApv is involved in predator defense and MEApd is activated
in the context of conspecific defense[50–52]. However, there is also evidence that the MEApv is
activated when male mice are exposed to conspecific male odors[53] or in social contexts with
conspecifics such as mating and aggressive same-sex encounters[11]. It has been shown that the
MEApv of female mice responds more robustly to dominant as opposed to subordinate urine
from male conspecifics[28], which is consistent with our results. Although we observed a robust
overall response to alpha urine, regardless of its familiarity or the social status of the subject
animal, further investigation into neuronal type might reveal different response characteristics
within this nucleus associated with these different social contexts further supporting a role for
the MEApd in decision making in competitive social contexts[11,54,55].
282
While the mPFC and AH also strongly contribute to Function 1, they also appear to have
surprisingly refined responses to additional features of social cues, as revealed by the pairwise
comparisons (**Table 1**). Importantly, the internal state of the subject (i.e. their own social status),
highly influenced activation patterns. For instance, the mPFC of dominant subjects, but not
subordinate subjects, exhibits increased activation in response to unfamiliar subordinate urine
compared to unfamiliar alpha urine. Interestingly, this differentiation of subordinate versus alpha
urine is not observed when the urine source is familiar. Furthermore, dominant subjects respond
significantly more to unfamiliar subordinate urine than familiar subordinate urine. Previous
studies corroborate our observation that social status is positively associated with activity in the
mPFC[56,57]. The mPFC is also implicated in the processing of reward-based motivation, decision-
making, cognition and fear and stress responses, all of which may be differentially regulated in
dominant versus subordinate mice. Indeed, it has been shown that dominant mice exhibit
greater motivation in social novelty exploration tasks[58]. In conjunction with our findings, it is
possible that the mPFC is engaged in social motivation-related processes when presented with
a novel social cue from a relatively subordinate source.
298
Processing of Familiar versus Unfamiliar Olfactory Cues
As with distinguishing alpha versus subordinate urine, no one brain region appears to be
exclusively tuned to familiar versus unfamiliar cues. The AH**,** which contributes heavily to
Functions 1 and 2, exhibits differential activation across all three features of the cue
presentation, with some activation patterns overlapping with those of the mPFC. While the AH
serves important non-social functions (e.g. thermoregulation), it has also been implicated in
social aggression, with elevated cFos immunoreactivity observed in mice selected for

aggression[59] as well as during offensive aggressive interactions in golden hamsters[60].
Interestingly, we find that the AH of dominant subjects responds more to a familiar alpha than
unfamiliar alpha. It is possible that AH activation is indicative of a response to a group-member
with which they share a history of agonistic interactions. We must stress, however, that the AH
exhibits differential activation in many pairwise comparisons, and therefore it's response to
social cues is likely to be influenced by many contextual factors.
Like the AH, the PMv of dominant subjects also exhibits increased activation in response
to familiar alpha urine but not unfamiliar alpha urine. Subordinate subjects also exhibited
increased PMv activation to unfamiliar urine, regardless of the cue source's social status.
Stimulating PMv neurons in male mice has been shown to trigger attack behavior towards
conspecifics, whereas silencing them interrupts attacks[61]. Activation of the PMv can also induce
general reward-related behaviors, including conditioned-place preference, and thus it has been
suggested that it may play a role in organizing goal-oriented aggression[61]. That different
olfactory cues activate the PMv of dominant and subordinate animals in our study may reflect
differences in such goal-oriented aggression.
The PMv is innervated by monosynaptic reciprocal excitatory connections from the
VMHvl[61,62]. Interestingly, the differential responses of VMH regions appear to compliment those
of the PMv, such that dominant subjects exhibit increased activation in response to unfamiliar
alpha urine compared to familiar alpha urine and subordinate subjects exhibit increased
activation to alpha urine in general, regardless of familiarity. VMHvl also contributes heavily to
Function 2, which we hypothesize represents level of saliency or level of social vigilance
towards given cues. Neurons in VMHvl, especially those expressing estrogen receptor 1, show
significant neural activation in aggressive, mating and defensive contexts[11,63–66]. These social
contexts are highly salient, either aversive or appetitive, thus the sensitivity of VMHvl to these
social contexts suggests that social information is gathered and the saliency of the contexts is
estimated in VMH. Furthermore, optogenetic activation of VMHvl induces male mice to increase
aggression[11], whereas the activation of VMHdm/c induces defensive behavior[66]. These results
suggest that VMH is significantly involved in decision-making of social behavior.
It has also been shown that the level of activation observed in the VMHvl is associated
with differences in internal state (e.g. hunger or thirst), and thus activation could represent
arousal-related information that guides social decision making[63]. The results from the current
study also suggest that social status, a component of internal state, influences responses to
social cue. While VMHvl activity in subordinate subjects increased in response to both familiar
and unfamiliar alpha urine, activity in dominant subjects only increased in response to unfamiliar
alpha urine. This data suggests that the familiar alpha cue is not as salient to dominant subjects
as subordinate subjects. While we've only discussed connections between the VMH, MEA and
PMv, the VMH shares reciprocal connections with many brain regions including those in the
SDMN[66–68]. The VMH's abundant connections further supports the idea that it integrates various
types of external social information with internal state.
The VMH, AH and PMd are sources of inputs to the PAG[50], another region that responds
significantly to urinary cues. Function 2 is most strongly driven by differences in the vlPAG
which, like the VMH, exhibits increased activation in response to unfamiliar alpha urine in both
dominant and subordinate subjects. Moreover, the vlPAG shows increased activation in
dominant subjects when exposed to familiar subordinate urine compared to familiar alpha urine.
In general, the PAG regulates the acquisition and expression of defensive behaviors in
response to threatening stimuli and plays a role in fear learning[69]. For instance, optogenetic
stimulation of the vlPAG leads to increased defensive behavior[70] and micturition[71], as well as
facilitating Pavlovian fear conditioning[69]. In the current study, it is possible that increased
activation of the vlPAG in response to an unfamiliar alpha reflects threat evaluation. In
subordinate males, threat evaluation may lead to a defensive fear response, whereas in
dominant subjects, it may elicit social vigilance and countermarking behavior. This would also

explain dominant subjects' increased activation in response to urine from a familiar subordinate
source, as subjects would typically display vigilant dominant behavior towards them in a
hierarchy context.
360
Conclusion
Virtually all brain regions that respond to social cues in this study are modulated by
social status. Dominant and subordinate subjects only share similar response patterns across
all conditions in the MEApv, where activity strongly predicted whether the cue source was an
alpha or subordinate. Considering the MEApv's early position in the olfactory processing
stream, we hypothesize that this region encodes cue information that is immediately important
for survival. Subcortical targets of the MEA and cortical regions also respond significantly to
urinary cues, including but not limited to the mPFC, AH, PMv, VMH, and vlPAG. All of these
regions exhibit differential activation across all three features of the cue presentation, meaning
they are receptive to the effects of the cue source's social status, familiarity with the source, and
the subject's internal state.
We believe these differences in neural activity across the brain could reflect the cognitive
processes that enable these animals to behave in a socially plastic and contextually appropriate
manner. The ability to extract relative social status information from social cues and to use that
information to guide behavior is crucial for the maintenance of social hierarchies[1]. In addition to
neural mechanisms that drive rapid behavioral flexibility from one context to the next, other
types of plasticity occur on longer timescales and with varying reversibility[72]. In the current
study, we show that brain-wide activity of dominant subjects is highly differentiated between
types of social cues, whereas activation changes in subordinate subjects is more subtle across
cues. This suggests that there are inherent differences in the processing of olfactory information
related to dominant status. This hypothesis is supported by existing evidence that synaptic
changes emerge in dominant individuals as they form dominance relationships[36,73]. These
status-associated changes in neural plasticity can have implications across cognitive domains,
such as learning and memory[74]. Further investigation into how reversible these status-related
changes are will yield valuable insight to how social experience shapes the brain and life-history
trajectories.
In conclusion, social information conveyed in odor cues is processed by a distributed
network of regions, most of which respond to multiple contextual features. Rather than
identifying distinct nodes where these features are encoded, this study demonstrates that the
multifaceted nature of social cues is reflected in the distributed and nuanced way the brain
processes them.
392
393
394 **Methods**
395
Subjects and housing
A total of 96 seven-week old male outbred Crl:CD-1 mice were obtained from Charles River
Laboratories (Wilmington, MA, USA) and habituated to the facility in pairs in standard cages (27
x 17 x 12 cm^3) with pine shaving bedding. Throughout the study, mice were housed under a
12:12 light:dark cycle (light cycle starting at 2400 hours and dark cycle starting at 1200hr) with
constant room temperature (22-23°C) and humidity (30-50%) and provided with standard chow
and water ad libitum. All mice were marked with a unique ID with non-toxic animal markers
(Stoelting Co., Wood Dale, IL, USA). During pair housing, we determined dyadic dominant-
subordinate relationships in each pair by focal observation during the dark phase of the light
cycle for 2 hours per day for 3-4 days prior to group housing. Animals that consistently initiated
and maintained aggressive behaviors were considered dominant and those that yielded to
attacks and showed persistent subordinate behavior were considered to be subordinate. After

11-15 days of pair housing, each batch of 12 pairs (four batches total) were assigned to two
separate social groups consisting of 12 mice each and placed into custom built vivaria as
previously described in So et al.[20]. Briefly, the vivarium (150 × 80 cm^2 and 220 cm high; Mid-
Atlantic, Hagerstown, MD, USA) consists of an upper level with three floor shelves and a lower
level with five nest-boxes connected by tubes and all surfaces were covered with pine shavings
(Supplemental **Fig. S4.**). The total surface of the vivarium is approximately 62,295 cm^2,
providing approximately 5191.25 cm^2 per mouse. Standard chow (LabDiet 5001, PMI, St. Louis,
MO, USA) and water were provided ad libitum at the top left and right side of the vivarium. Each
social group consisted of six dominant males and six subordinate males based on pair housing
observations. Live behavioral observations started on the first day of group housing and
continued for 10-13 days per social group with an average of 22 hours of total observation
evenly distributed across the housing period. During these observations, trained observers
recorded all interactions of fighting, chasing, mounting, subordinate posture and induce-flee
behaviors that occurred between any two individuals as further described in Williamson et al.[21].
All experiments were conducted in accordance with National Health Institute (NIH; Bethesda,
MD, USA) and the Columbia University Institutional Animal Care and Use Committee (IACUC
Protocol No. AC-AAAP5405) animal care guidelines.
425
Urine collection
On group housing Day 6-10, we collected urine daily from either alpha males or the most
subordinate males based on observation data (see **Fig.1**). Each day, mice were quickly
removed from the group housing environment and scruffed over Eppendorf tubes as described
in Lee et al.[23]. Collected urine samples were immediately stored at -20°C until the day of social
stimulus exposure. Urine samples were only defrosted once and pooled for each individual.
432
Social stimulus exposure
We habituated mice to scruffing and urine exposure for the last 5 days of pair housing by
applying 20ul of nuclease free water directly on their oronasal groove, following the procedure
previously described in Martel and Baum[75]. On the last day of group housing (group housing
day 9-12), mice were removed from the vivaria and isolated in a standard cage filled with pine
shaving and provided with ad libitum standard chow and water at zeitgeber hour 12 (start of the
dark cycle) for 60-140 minutes. Mice were randomly assigned to two experimental groups: a)
familiar urine exposure (the urine samples were collected from either alpha or the most
subordinate in the same group as subjects), b) unfamiliar urine exposure (the urine samples
were collected from either alpha or most subordinate individuals in different social group). After
the habituation, 7-9 mice in each social group (excluding the urine donor mouse and those with
midranks) were gently scruffed and 20ul of urine was applied onto the oronasal grove of each
mouse. A total of 64 mice from 8 social groups were exposed to urine stimuli (familiar alpha
male urine: 6 dominant mice and 10 subordinate mice; unfamiliar alpha male urine: 8 dominant
mice and 8 subordinate mice; familiar subordinate urine: 7 dominant mice and 9 subordinate
mice; unfamiliar subordinate urine: 7 dominant mice and 9 subordinate mice).
449
Fos protein immunohistochemistry
Ninety minutes after the exposure to urine stimulus, mice were deeply anesthetized with mixture
of ketamine and xylazine and perfused transcardially with pH7.4 0.1M phosphate-buffered
saline (PBS) followed by 4% paraformaldehyde (PFA) in 0.1M PBS. Brains were removed and
immersed overnight in 4% PFA then 30% sucrose in 0.1M PBS for at least 24 hours (4°C).
Whole brains were sectioned coronally at 40um thickness in a cryostat and stored in 0.1 M PB
azide. Every other series was used for processing according to free-floating avidin-biotin
procedure, using the Vectastain ABC Elite peroxidase rabbit IgG kit (Vector Laboratories,
Burlingame, CA, USA). Sections were first washed in 0.1 M PB for five minutes three times then

incubated with 3% H2O2 for 5 minutes followed by three times of 5-minute washes in 0.1M
PB/0.1% Triton-X (PBT). The sections were incubated with 1% normal goat serum (NGS) in
PBT for an hour followed by 4°C overnight incubation in primary Fos rabbit polyclonal IgG (sc-
52; Santa Cruz Biotechnology, Inc. Dallas, Texas, USA) at a concentration of 1:5000 in 1 %
NGS/PBT. Slices were washed three times in PBT for 5 minutes then incubated in biotinylated
anti-rabbit IgG (Vectastain, Vector Laboratories) at a concentration of 1:200 in PBS. The
sections were rinsed again three times with PBT for 5 minutes then incubated in avidin-biotin-
peroxidase complex at a concentration of 1:250 in PBS for an hour. Sections were washed
again with 0.1 M PBS three time for 5 minutes. H2O2 (final concentration as 0.02 %) was added
to 0.02 % 3,3'-diaminobenzidine (DAB) solution just prior to incubation. Sections were incubated
in DAB solution to visualize immunoreactivity for 2-4 minutes and the reaction was immediately
stopped by washing sections in PBS for 1 minute for the first wash and 5 minutes for the
remaining three washes. All sections were mounted on FisherBrand Plus slides within 24 hours
then processed for serial dehydration followed by coverslipping with Krystalon mounting
medium (Millipore Billerica, MA, USA).
474
Quantification of Fos-immunoreactive cells
Images were obtained with a Leica DMi8 microscope (Leica Microsystems, Wetzlar, Germany)
under a 10x objective at a magnification x100. Regions of interest were located using the mouse
brain atlas of Paxinos and Franklin[76]. A total of 23 brain regions were analyzed and their
bregmas are described in Supplemental Table. S1: anterior hypothalamus (AH), basolateral
amygdaloid nucleus (BLA), principal nucleus of the bed nucleus of the stria terminalis (BNSTpr),
field CA1 of hippocampus (CA1), field CA3 of hippocampus (CA3), central amygdaloid nucleus
(CEA), dentate gyrus (DG), dorsolateral periaqueductal gray (dlPAG), dorsomedial
hypothalamus (DMH), dorsomedial periaqueductal gray (dmPAG), lateral hypothalamus (LH),
ventral lateral septal nucleus (LSv), medial amygdaloid nucleus, posterodorsal part (MEApd),
medial amygdaloid nucleus, posteroventral part (MEApv), infralimbic/prelimbic regions of the
prefrontal cortex (mPFC), medial preoptic area (mPOA), nucleus accumbens core (NAcc),
dorsal premammillary nucleus (PMd), ventral premammillary nucleus (PMv), paraventricular
nucleus (PVN), ventrolateral periaqueductal gray (vlPAG), ventromedial hypothalamic nucleus,
dorsomedial central part (VMHdm/c) and ventromedial hypothalamic nucleus, ventrolateral part
(VMHvl) (see Supplemental **Fig. S5.** for example images). For each brain regions, 3 brain
sections per mouse were imaged and cell counts were summed. We overlaid images from The
Mouse Brain in Steretaxic Coordinates[76] to include only the exact portion of the region. We used
Fiji[77] to count cFos positive cells.
494
495   **Statistical analysis**
496   All statistical analyses were undertaken in R version 3.6.4.[78]
497
Group dominance structure, emergence of hierarchies and individual social status
For each cohort, we aggregated the total number of wins and losses by each mouse from all
recorded aggressive interaction over the entire group housing period into win/loss frequency
sociomatrices. To confirm the presence of a linear social hierarchy in each social cohort, we
calculated Landau's h-value, triangle transitivity (ttri) directional consistency (DC) and
associated p-values from 10,000 Monte-Carlo randomizations[79,80] using compete R package[81]
as previously described in Williamson et al.[21]. Further, we calculated ttri and Landau's h-value
and their significance for aggregated win/loss sociomatrices each day of group housing to
determine how quickly each cohort established a linear social hierarchy. Despotism of each
alpha was derived by calculating the proportion of wins by the alpha to all aggressive
interactions observed during group housing period. Emergence of linear and stable social
hierarchies were additionally confirmed by calculating individual Glicko ratings and visualizing

them for each cohort using PlayerRatings R package[82]. David's score, a win proportion measure
adjusted for opponent's relative dominance in the hierarchy[83], was calculated for each individual
to assign individual social rank in each hierarchy using the compete R package[81]. Based on our
observation of social hierarchy dynamics, we further categorized male mice living in social
hierarchies into three distinct social status groups: alpha (rank 1, the highest David's score),
subdominant (David's score >=0), and subordinate (David's score <0; see Supplemental **Fig.**
S6. for the David's score of each individual in the study) as previously described in Lee et
al.[23,84,85].
518
cFos immunoreactivity count data analysis
We used Bayesian hierarchical generalized linear modeling[38] fitted in STAN[86] with the brms R
package[87]. We took advantage of this Bayesian approach as it can handle outliers in the data by
implementing heavy-tailed distributions while accommodating heterogeneity of variances across
groups[38]. Similar to traditional frequentist analysis of variance (ANOVA), but free from
frequentist ANOVA's strict assumptions, we conducted 12 planned pairwise comparisons
among experimental groups (main effects; familiarity of the given social cue and individual
social rank) for each brain region. We used flexible discriminant analysis (FDA) to examine
whether cFos immunoreactivity across 23 brain regions can predict which of the eight
experimental groups (status x familiarity x cue source) each individual corresponded to. We
used the mda R package to conduct FDA[88]. As an extension of linear discriminant analysis
(LDA), FDA is capable of handling non-normality or non-linearity among variables, thus FDA
results in more robust classification of groups given the data compared to LDA when data
violates the assumptions of LDA[89].
533
534   **References**
535

1.   Taborsky, B. & Oliveira, R. F. Social competence: an evolutionary approach. *Trends Ecol.*

Evol. **27**, 679–688 (2012).

2.   Anderson, D. J. Circuit modules linking internal states and social behaviour in flies and mice.

Nat. Rev. Neurosci. **17**, 692–704 (2016).

3.   Chen, P. & Hong, W. Neural Circuit Mechanisms of Social Behavior. *Neuron* **98**, 16–30

(2018).

4.   Insel, T. R. & Fernald, R. D. How the brain processes social information: Searching for the

social brain. *Annu. Rev. Neurosci.* **27**, 697–722 (2004).

5.   O'Connell, L. A. & Hofmann, H. A. Genes, hormones, and circuits: An integrative approach

to study the evolution of social behavior. *Front. Neuroendocrinol.* **32**, 320–335 (2011).

6.   Goodson, J. L. The vertebrate social behavior network: Evolutionary themes and variations.

Horm. Behav. **48**, 11–22 (2005).

7.   Newman, S. W. The Medial Extended Amygdala in Male Reproductive Behavior A Node in

the Mammalian Social Behavior Network. *Ann. N. Y. Acad. Sci.* **877**, 242–257 (1999).

8.   O'Connell, L. A. & Hofmann, H. A. The Vertebrate mesolimbic reward system and social

behavior network: A comparative synthesis. *J. Comp. Neurol.* **519**, 3599–3639 (2011).

9.   Golden, S. A. *et al.* Compulsive addiction-like aggressive behavior in mice. *Biol. Psychiatry*

82, 239–248 (2017).

10. Hong, W., Kim, D.-W. & Anderson, D. J. Antagonistic control of social versus repetitive self-

grooming behaviors by separable amygdala neuronal subsets. *Cell* **158**, 1348–1361 (2014).

11. Lin, D. *et al.* Functional identification of an aggression locus in the mouse hypothalamus.

Nature **470**, 221–226 (2011).

12. Williamson, C. M., Klein, I. S., Lee, W. & Curley, J. P. Immediate early gene activation

throughout the brain is associated with dynamic changes in social context. *Soc. Neurosci.*

14, 253–265 (2019).

13. Wu, Z., Autry, A. E., Bergan, J. F., Watabe-Uchida, M. & Dulac, C. G. Galanin neurons in the

medial preoptic area govern parental behaviour. *Nature* **509**, 325–330 (2014).

14. Chase, I. D. Behavioral sequences during dominance hierarchy formation in chickens.

Science **216**, 439–440 (1982).

15. Curley, J. P. Temporal pairwise-correlation analysis provides empirical support for attention

hierarchies in mice. *Biol. Lett.* **12**, 20160192 (2016).

16. Desjardins, J. K., Hofmann, H. A. & Fernald, R. D. Social Context Influences Aggressive and

Courtship Behavior in a Cichlid Fish. *PLOS ONE* **7**, e32781 (2012).

17. Maruska, K. P. & Fernald, R. D. Behavioral and physiological plasticity: Rapid changes during

social ascent in an African cichlid fish. *Horm. Behav.* **58**, 230–240 (2010).

18. Snyder-Mackler, N. *et al.* Social status alters immune regulation and response to infection in

macaques. *Science* **354**, 1041–1045 (2016).

19. Williamson, C. M., Romeo, R. D. & Curley, J. P. Dynamic changes in social dominance and

mPOA GnRH expression in male mice following social opportunity. *Horm. Behav.* **87**, 80–88

(2017).

20. So, N., Franks, B., Lim, S. & Curley, J. P. A social network approach reveals associations

between mouse social dominance and brain gene expression. *PloS One* **10**, e0134509

(2015).

21. Williamson, C. M., Lee, W. & Curley, J. P. Temporal dynamics of social hierarchy formation

and maintenance in male mice. *Anim. Behav.* **115**, 259–272 (2016).

22. Lee, W., Fu, J., Bouwman, N., Farago, P. & Curley, J. P. Temporal microstructure of dyadic

social behavior during relationship formation in mice. *PLOS ONE* **14**, e0220596 (2019).

23. Lee, W., Khan, A. & Curley, J. P. Major urinary protein levels are associated with social

status and context in mouse social hierarchies. *Proc. R. Soc. B Biol. Sci.* **284**, 20171570

(2017).

24. Williamson, C. M., Franks, B. & Curley, J. P. Mouse Social Network Dynamics and

Community Structure are Associated with Plasticity-Related Brain Gene Expression. *Front.*

Behav. Neurosci. 152 (2016) doi:10.3389/fnbeh.2016.00152.

25. Hoffman, E., Pickavance, L., Thippeswamy, T., Beynon, R. J. & Hurst, J. L. The male sex

pheromone darcin stimulates hippocampal neurogenesis and cell proliferation in the

subventricular zone in female mice. *Front. Behav. Neurosci.* **9**, (2015).

26. Hurst, J. L., Thom, M. D., Nevison, C. M., Humphries, R. E. & Beynon, R. J. MHC odours are

not required or sufficient for recognition of individual scent owners. *Proc. R. Soc. B Biol. Sci.*

272, 715–724 (2005).

27. Nelson, A. C., Cunningham, C. B., Ruff, J. S. & Potts, W. K. Protein pheromone expression

levels predict and respond to the formation of social dominance networks. *J. Evol. Biol.* **28**,

1213–1224 (2015).

28. Veyrac, A., Wang, G., Baum, M. J. & Bakker, J. The main and accessory olfactory systems of

female mice are activated differentially by dominant versus subordinate male urinary

odors. *Brain Res.* **1402**, 20–29 (2011).

29. Borelli, K. G. *et al.* Neural correlates of scent marking behavior in C57BL/6J mice: detection

and recognition of a social stimulus. *Neuroscience* **162**, 914–923 (2009).

30. Lüscher Dias, T., Fernandes Golino, H., Oliveira, V. E. M. de, Dutra Moraes, M. F. &

Schenatto Pereira, G. c-Fos expression predicts long-term social memory retrieval in mice.

Behav. Brain Res. **313**, 260–271 (2016).

31. Nikonov, A. A. & Maruska, K. P. Male dominance status regulates odor-evoked processing in

the forebrain of a cichlid fish. *Sci. Rep.* **9**, 1–17 (2019).

32. Hurst, J. L. The priming effects of urine substrate marks on interactions between male

house mice, Mus musculus domesticus Schwarz & Schwarz. *Anim. Behav.* **45**, 55–81 (1993).

33. Rich, T. J. & Hurst, J. L. Scent marks as reliable signals of the competitive ability of mates.

Anim. Behav. **56**, 727–735 (1998).

34. Hurst, J. L. & Beynon, R. J. Scent wars: the chemobiology of competitive signalling in mice.

Bioessays **26**, 1288–1298 (2004).

35. Maruska, K. P. & Fernald, R. D. Contextual chemosensory urine signaling in an African cichlid

fish. *J. Exp. Biol.* **215**, 68–74 (2012).

36. Kingsbury, L. *et al.* Correlated Neural Activity and Encoding of Behavior across Brains of

Socially Interacting Animals. *Cell* **178**, 429-446.e16 (2019).

37. Pannozzo, P. L., Phillips, K. A., Haas, M. E. & Mintz, E. M. Social monitoring reflects

dominance relationships in a small captive group of brown capuchin monkeys (cebus

apella). *Ethology* **113**, 881–888 (2007).

38. Kruschke, J. *Doing bayesian data analysis: A tutorial with r, JAGS, and stan*. (Academic

Press, 2014).

39. O'Connell, L. A. & Hofmann, H. A. The Vertebrate mesolimbic reward system and social

behavior network: A comparative synthesis. *J. Comp. Neurol.* **519**, 3599–3639 (2011).

40. Prounis, G. S. & Ophir, A. G. One cranium, two brains not yet introduced: Distinct but

complementary views of the social brain. *Neurosci. Biobehav. Rev.* **108**, 231–245 (2020).

41. Field, K. E. & Maruska, K. P. Context-dependent chemosensory signaling, aggression and

neural activation patterns in gravid female African cichlid fish. *J. Exp. Biol.* **220**, 4689–4702

(2017).

42. Field, K. E., McVicker, C. T. & Maruska, K. P. Sexually-relevant visual and chemosensory

signals induce distinct behaviors and neural activation patterns in the social african cichlid,

astatotilapia burtoni. *Front. Behav. Neurosci.* **12**, (2018).

43. Arakawa, H., Blanchard, D. C. & Blanchard, R. J. Central oxytocin regulates social familiarity

and scent marking behavior that involves amicable odor signals between male mice.

Physiol. Behav. **146**, 36–46 (2015).

44. Haga, S. *et al.* The male mouse pheromone ESP1 enhances female sexual receptive

behaviour through a specific vomeronasal receptor. *Nature* **466**, 118–122 (2010).

45. Roberts, S. A. *et al.* Darcin: a male pheromone that stimulates female memory and sexual

attraction to an individual male's odour. *BMC Biol.* **8**, 75 (2010).

46. Guo, H., Fang, Q., Huo, Y., Zhang, Y. & Zhang, J. Social dominance-related major urinary

proteins and the regulatory mechanism in mice. *Integr. Zool.* (2015) doi:10.1111/1749-

4877.12165.

47. Brennan, P. A. & Zufall, F. Pheromonal communication in vertebrates. *Nature* **444**, 308–315

(2006).

48. Choi, G. B. *et al.* Lhx6 Delineates a Pathway Mediating Innate Reproductive Behaviors from

the Amygdala to the Hypothalamus. *Neuron* **46**, 647–660 (2005).

49. Miller, S. M., Marcotulli, D., Shen, A. & Zweifel, L. S. Divergent medial amygdala projections

regulate approach–avoidance conflict behavior. *Nat. Neurosci.* **22**, 565–575 (2019).

50. Canteras, N. S. The medial hypothalamic defensive system: Hodological organization and

functional implications. *Pharmacol. Biochem. Behav.* **71**, 481–491 (2002).

51. Motta, S. C. *et al.* Dissecting the brain's fear system reveals the hypothalamus is critical for

responding in subordinate conspecific intruders. *Proc. Natl. Acad. Sci.* **106**, 4870–4875

(2009).

52. Sokolowski, K. & Corbin, J. G. Wired for behaviors: from development to function of innate

limbic system circuitry. *Front. Mol. Neurosci.* **5**, (2012).

53. Samuelsen, C. L. & Meredith, M. Categorization of biologically relevant chemical signals in

the medial amygdala. *Brain Res.* **1263**, 33–42 (2009).

54. Hashikawa, K., Hashikawa, Y., Falkner, A. & Lin, D. The neural circuits of mating and fighting

in male mice. *Curr. Opin. Neurobiol.* **38**, 27–37 (2016).

55. Li, Y. *et al.* Neuronal Representation of Social Information in the Medial Amygdala of Awake

Behaving Mice. *Cell* **171**, 1176-1190.e17 (2017).

56. Wang, F. *et al.* Bidirectional control of social hierarchy by synaptic efficacy in medial

prefrontal cortex. *Science* **334**, 693–697 (2011).

57. Zink, C. F. *et al.* Know Your Place: Neural Processing of Social Hierarchy in Humans. *Neuron*

58, 273–283 (2008).

58. Kunkel, T. & Wang, H. Socially dominant mice in C57BL6 background show increased social

motivation. *Behav. Brain Res.* **336**, 173–176 (2018).

59. Haller, J., Tóth, M., Halasz, J. & De Boer, S. F. Patterns of violent aggression-induced brain c-

fos expression in male mice selected for aggressiveness. *Physiol. Behav.* **88**, 173–182

(2006).

60. Delville, Y., Vries, G. J. D. & Ferris, C. F. Neural Connections of the Anterior Hypothalamus

and Agonistic Behavior in Golden Hamsters. *Brain. Behav. Evol.* **55**, 53–76 (2000).

61. Stagkourakis, S. *et al.* A neural network for intermale aggression to establish social

hierarchy. *Nat. Neurosci.* **21**, 834–842 (2018).

62. Ziegler, D. R., Cullinan, W. E. & Herman, J. P. Distribution of vesicular glutamate transporter

mRNA in rat hypothalamus. *J. Comp. Neurol.* **448**, 217–229 (2002).

63. Falkner, A. L., Dollar, P., Perona, P., Anderson, D. J. & Lin, D. Decoding ventromedial

hypothalamic neural activity during male mouse aggression. *J. Neurosci.* **34**, 5971–5984

(2014).

64. Hashikawa, Y., Hashikawa, K., Falkner, A. L. & Lin, D. Ventromedial hypothalamus and the

generation of aggression. *Front. Syst. Neurosci.* **11**, (2017).

65. Lee, H. *et al.* Scalable control of mounting and attack by Esr1 + neurons in the ventromedial

hypothalamus. *Nature* **509**, 627–632 (2014).

66. Wang, L., Chen, I. Z. & Lin, D. Collateral pathways from the ventromedial hypothalamus

mediate defensive behaviors. *Neuron* **85**, 1344–1358 (2015).

67. Lo, L. *et al.* Connectional architecture of a mouse hypothalamic circuit node controlling

social behavior. *Proc. Natl. Acad. Sci.* **116**, 7503–7512 (2019).

68. Wang, L. *et al.* Hypothalamic control of conspecific self-defense. *Cell Rep.* **26**, 1747-1758.e5

(2019).

69. Assareh, N., Bagley, E. E., Carrive, P. & McNally, G. P. Brief optogenetic inhibition of rat

lateral or ventrolateral periaqueductal gray augments the acquisition of Pavlovian fear

conditioning. *Behav. Neurosci.* **131**, 454–459 (2017).

70. Assareh, N., Sarrami, M., Carrive, P. & McNally, G. P. The organization of defensive behavior

elicited by optogenetic excitation of rat lateral or ventrolateral periaqueductal gray. *Behav.*

Neurosci. **130**, 406–414 (2016).

71. Verstegen, A. M. J. *et al.* Non-crh glutamatergic neurons in barrington's nucleus control

micturition via glutamatergic afferents from the midbrain and hypothalamus. *Curr. Biol.* **29**,

2775-2789.e7 (2019).

72. Cardoso, S. D., Teles, M. C. & Oliveira, R. F. Neurogenomic mechanisms of social plasticity. *J.*

Exp. Biol. **218**, 140–149 (2015).

73. Nelson, A. C. *et al.* Molecular and Circuit Architecture of Social Hierarchy. *bioRxiv* 838664

(2019) doi:10.1101/838664.

74. Barnard, C. J. & Luo, N. Acquisition of dominance status affects maze learning in mice.

Behav. Processes **60**, 53–59 (2002).

75. Martel, K. L. & Baum, M. J. Sexually dimorphic activation of the accessory, but not the main,

olfactory bulb in mice by urinary volatiles. *Eur. J. Neurosci.* **26**, 463–475 (2007).

76. Paxinos, G. & Franklin, K. B. *Paxinos and Franklin's the mouse brain in stereotaxic*

coordinates. (Academic press, 2019).

77. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods*

9, 676–682 (2012).

78. R Core Team. *R: A language and environment for statistical computing*. (R Foundation for

Statistical Computing, 2019).

79. de Vries, H. An improved test of linearity in dominance hierarchies containing unknown or

tied relationships. *Anim. Behav.* **50**, 1375–1389 (1995).

80. Shizuka, D. & McDonald, D. B. A social network perspective on measurements of dominance

hierarchies. *Anim. Behav.* **83**, 925–934 (2012).

81. Curley, J. P. *compete: Organizing and analyzing social dominance hierarchy data.* (2016).

82. Stephenson, A. & Sonas, J. *PlayerRatings: Dynamic updating methods for player ratings*

estimation. (2012).

83. Gammell, M. P., Vries, de H., Jennings, D. J., Carlin, C. M. & Hayden, T. J. David's score: a

more appropriate dominance ranking method than Clutton-Brock et al.'s index. *Anim.*

Behav. **66**, 601–605 (2003).

84. Lee, W., Yang, E. & Curley, J. P. Foraging dynamics are associated with social status and

context in mouse social hierarchies. *PeerJ* **6**, e5617 (2018).

85. Lee, W. *et al.* Social status in mouse social hierarchies is associated with variation in

oxytocin and vasopressin 1a receptor densities. *Horm. Behav.* **114**, 104551 (2019).

86. Carpenter, B. *et al.* Stan: A probabilistic programming language. *J. Stat. Softw.* **76**, 1–32

(2017).

87. Bürkner, P.-C. *brms: Bayesian regression models using 'stan'*. (2020).

88. Leisch, S. original by T. H. & R. T. O. R. port by F., Ripley, K. H. & D., B. *mda: Mixture and*

flexible discriminant analysis. (2017).

89. Hastie, T., Tibshirani, R. & Buja, A. Flexible discriminant analysis by optimal scoring. *J. Am.*

Stat. Assoc. **89**, 1255–1270 (1994).