The Bromodomains of the mammalian SWI/SNF (mSWI/SNF) ATPases Brahma (BRM) and

Brahma Related Gene 1 (BRG1) promote chromatin interaction and are critical for skeletal

muscle differentiation

Tapan Sharma[a], Hanna Witwicka[a]*, Anthony N. Imbalzano[a,#]

[a] Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical

School, Worcester, MA 01605 USA

Running Title: BRG1/BRM bromodomains are critical for myogenesis

Address correspondence to Anthony N. Imbalzano, Anthony.Imbalzano@umassmed.edu.

*Present Address: Hanna Witwicka, Charles River Laboratories, Inc., Shrewsbury, MA 01545,

USA

Word count for Abstract: 200

Word count for Introduction, Results and Discussion: 4901 (combined upper limit is 8000 words)

Word count for Materials and Methods: 1565 (no limit)

ABSTRACT: Skeletal muscle differentiation induces changes in the epigenome of myoblasts as

they proceed towards a myogenic phenotype. mSWI/SNF chromatin remodeling enzymes

coordinate with lineage-determining transcription factors and are key regulators of differentiation.

Three mSWI/SNF proteins, the mutually exclusive ATPases, BRG1 and BRM, and the BAF180

protein (Polybromo1, PBRM1) contain bromodomains belonging to the same structural subfamily.

Bromodomains bind to acetylated lysines on histone N-terminal tails and on other proteins.

Pharmacological inhibition of mSWI/SNF bromodomain function using the selective inhibitor

PFI-3 reduced differentiation, decreased expression of myogenic genes, and increased the

expression of cell cycle-related genes, and the number of cells that remained in the cell cycle.

Knockdown of BAF180 had no effect on differentiation, suggesting that only the BRG1 and BRM

bromodomains contributed to differentiation. Comparison with existing gene expression data from

myoblasts subjected to knockdown of BRG1 or BRM showed that bromodomain function was

required for a subset of BRG1- and BRM-dependent gene expression. ChIP analysis revealed

decreased BRG1 and BRM binding to target gene promoters, indicating that the BRG1 and BRM

bromodomains promote chromatin binding. Thus mSWI/SNF ATPase bromodomains contribute

to cell cycle exit, to skeletal muscle-specific gene expression, and to stable promoter binding by

the mSWI/SNF ATPases.

INTRODUCTION

Regulation of gene expression is a tightly coordinated process that is dependent on transcription

factors, coactivators and chromatin remodelers. Some of these regulators are tissue-specific and

act on target genes in a context-dependent manner. Tissue-specific regulation is absolutely crucial

for proper development of multi-cellular life forms in which all cells contain the same genetic

information. Portions of the genome that are irrelevant to a particular tissue type are often

condensed into repressive heterochromatin as development and differentiation occur (1, 2). In

contrast, coordinated activity of lineage-determining transcription factors and chromatin

remodelers, in particular the mSWI/SNF family of chromatin remodeling enzymes, drives many

differentiation events, including skeletal muscle differentiation (3–7). The mSWI/SNF enzymes

remodel chromatin in an ATP-dependent manner (8–10) and form a family of enzymes assembled

into different configurations from a potential pool of more than twenty subunit proteins (11–13).

The BRG1 and BRM ATPases act as mutually exclusive catalytic subunits (10).

Skeletal muscle originates from the paraxial mesoderm during embryogenesis. Fetal skeletal

myogenesis is characterized by an abundance of myogenic progenitor cells that divide actively and

fuse to form multinucleated muscle fibers (14, 15). As the embryo develops into an adult, these

progenitor cells become relatively sparse and quiescent. These adult stem cells are known as

satellite cells and can be activated to proliferate and regenerate new myofibers in case of an injury

to adult skeletal muscle (16–18). Upon activation, expression of myogenic regulatory factors

(MRFs) – MYOD, MRF4, MYF5 and Myogenin - is initiated in a coordinated manner. MRFs are

basic helix-loop-helix (bHLH) proteins that are evolutionarily conserved from worms to humans

(19–21). They bind to consensus sequences called E-boxes at target muscle promoters and activate

muscle-specific gene expression (22). Another family of transcription factors called the myocyte

enhancer factor 2 (MEF2) family acts with the MRFs to promote expression of the myogenic

genes (23, 24).

During skeletal myogenesis, the mSWI/SNF complex is recruited to the myogenic loci by MRFs

(25–30), in some cases, in conjunction with PBX1 (25). Mechanistically, upon induction of

differentiation in myocytes, the p38 kinase responds to extracellular cues by phosphorylating the

BAF60c subunit of mSWI/SNF chromatin remodeling enzymes, which is associated with MYOD

on myogenic genes in the absence of other mSWI/SNF subunits in proliferating myoblasts (27,

29). The phospho-BAF60c-MYOD complex then recruits the rest of the mSWI/SNF complex to

myogenic loci, which promotes chromatin accessibility and activates gene expression (29). Once

recruited to myogenic loci, the ATPase activity of BRG1 or BRM in the complex is known to be

indispensable for expression of the differentiation-specific gene program (26, 28, 31).

The BRG1 and BRM ATPases possess bromodomains in the C-terminal part of the protein (32–

34). Bromodomains are well-characterized motifs known to interact with acetylated lysine residues

on the N-terminal tails of histones H3 and H4 (35, 36) and on other non-histone proteins (37). The

interaction of bromodomains with acetylated histones has been determined to be crucial for

regulation of some gene expression events (37). Based on structural homology, bromodomain-

containing proteins can be classified into eight families (35, 38). BRG1 and BRM belong to family

VIII of bromodomains along with a third mSWI/SNF protein called BAF180 (Polybromo1,

PBRM1, PB1) that contains six tandem bromodomains (38, 39)

In this study, we characterized the role of mSWI/SNF bromodomains in the context of skeletal

myogenesis. We showed that inhibiting bromodomain function using PFI-3, a specific

pharmacological inhibitor that binds to the BRG1, BRM and BAF180 bromodomains (40–42),

reduced the ability of mouse myoblasts to differentiate into myotubes. Using RNA-sequencing,

we identified the genes whose expression is dependent on mSWI/SNF bromodomains. Broadly,

proliferation-related genes were found to be upregulated by bromodomain inhibition while

myogenic genes were downregulated. We also demonstrated that bromodomain function is

essential for timely exit of myoblasts from the cell cycle upon induction of differentiation. We

determined that BAF180 is not required for myogenesis in mouse myoblasts and demonstrated

that the BRG1 and BRM bromodomains play a crucial role in skeletal muscle differentiation by

promoting the stable binding of BRG1 and BRM to target gene promoters. Thus, this study

mechanistically demonstrates the specific importance of mSWI/SNF bromodomains in context of

skeletal muscle differentiation.

RESULTS

Inhibition of bromodomain function results in aberrant myotube fusion

PFI-3 is a pharmacological inhibitor specific for the BRG1, BRM and BAF180 bromodomains,

members of bromodomain family VIII (40, 41).   Prior work showed that PFI-3 impaired

differentiation of immortalized pre-adipocytes and myoblasts (42). The mechanisms responsible

for the observed effects on differentiation were not defined, so we sought to investigate the roles

played by mSWI/SNF bromodomains during myogenesis.

C2C12 immortalized myoblasts and primary myoblasts isolated from the tibialis anterior muscles

of 1-week old C57BL/6 mice were assayed for their ability to differentiate in the presence of PFI-

3 or the vehicle (DMSO). DMSO-treated C2C12 myoblasts immunostained for myosin heavy

chain (MHC) showed formation of longer and thicker myotubes at 48h and 72h post induction of

differentiation than did C2C12 myoblasts treated with PFI-3 (Fig 1a). The efficiency of myogenic

differentiation can be scored by calculating fusion index, which is the ratio of the number of nuclei

in MHC-stained cells to the total number of nuclei (43). C2C12 cells treated with PFI-3 showed a

>50% decrease in fusion index at 24h, 48h and 72h as compared to control samples (Fig 1b).

Similar results were observed when primary myoblasts were exposed to PFI-3. While DMSO-

treated primary myoblasts showed elongated myotubes upon induction of differentiation, the PFI-

3-treated cells showed fewer and less elongated myotubes at corresponding timepoints (Supp fig

1a). Quantitative analysis of differentiated primary myoblasts immunostained for MHC showed

about a 25-30% decrease in fusion index (Supp fig 1b).

We further quantitatively analyzed the extent of differentiation by counting the number of nuclei

in MHC-positive myotubes and classifying them into groups at each timepoint. DMSO-treated

C2C12 cells shifted from the majority of 48h myotubes having <5 nuclei to the majority of

myotubes having >5 nuclei by 72h (Fig 1c). PFI-3-treated cells failed to make this switch; the

majority of 72h myotubes had <5 nuclei (Fig 1c). Similarly, in PFI-3 treated primary cells, the

number of myotubes with >5 nuclei at 36h was about one-third of those in DMSO-treated control

cells. In PFI-3 treated samples, cells with a single nucleus positively immunostained for MHC

were abundant, showing a failure of differentiating myoblasts to fuse (Supp fig 1c). These results

suggest that an initial myogenic stimulus is present but is not fully implemented due to inhibition

of bromodomain function.

Myogenic genes are downregulated upon PFI-3 induced inhibition of mSWI/SNF

bromodomains

The results show that PFI-3 treatment causes defects in myogenic differentiation, including an

inability of the differentiating myoblasts and/or nascent myotubes to fuse. Myomaker and

myomixer have been identified as master regulators of myoblast fusion (44–47). We therefore

determined whether the expression of these two regulators was altered upon PFI-3 induced

bromodomain inhibition. The results show that expression of these two genes was significantly

lower in PFI-3 treated C2C12 cells (Fig 2a). The expression of other myogenic genes like

myogenin, creatine kinase and myosin light chain 1 was also significantly decreased in PFI-3

treated samples, as was the expression of caveolin 3 and integrin 7A, two muscle differentiation-

related genes (Fig 2b). Western blot analysis confirmed the decreased expression of myosin

heavy chain in PFI-3 treated C2C12 cells (Fig 2c). Similar results were obtained for PFI-3

treated primary myoblasts (Supp fig 2). The gene expression signatures from both C2C12 cells

and primary myoblasts provides a molecular explanation for the differentiation phenotype caused

by bromodomain inhibition.

RNA-seq analysis of PFI-3 treated C2C12 cells shows upregulation of cell cycle genes and

downregulation of myogenic genes

To gain better insight into effect of the molecular mechanism of bromodomain inhibition on

skeletal muscle differentiation, we performed RNA-sequencing of C2C12 cells treated with

DMSO or PFI-3. Cells were harvested from proliferative stage (GM) and two differentiated stages

(DM 24h and DM 48h post-induction). Libraries generated from the samples had ~45M unique

reads. Transcripts were mapped to the mouse genome (mm10) and gene expression levels were

calculated. Genes that were identified to be differentially expressed in both replicates for each

condition and timepoint were considered for further analysis.

We first examined whether PFI-3 treatment affected gene expression of the subunits of mSWI/SNF

complexes. A recent characterization of sub-families of mSWI/SNF complexes identified 29

subunit proteins (13). Assessment of expression of the genes encoding each of these proteins at

each time point found only two instances of statistically significant differences (Supp. Table 1).

Arid1a expression was reduced ~7% at 24 h post-differentiation and *Actl6a* expression was

increased ~27% at 48 h post-differentiation. We conclude that PFI-3 treatment had essentially no

effect on the expression of the genes encoding mSWI/SNF subunits.

Inhibition of bromodomain function affected the expression of about 50% of the total genes

identified as expressed over the time course of the experiment (Fig 3a). The number of DEGs due

to bromodomain inhibition increased as a function of differentiation (Fig 3b). The total number of

DEGs for proliferating cells (GM) and differentiating cells at 24h or 48h post-differentiation (DM

24h and DM 48h) were 3144 (up 2216; down 928), 4675 (up 2878; down 1797) and 5261 (up

3105; down 2156), respectively (Supp. Table 2). Gene expression at DM 24h and DM 48h was

strongly correlated with 2359 common DEGs as compared to about 1634 common DEGs between

GM and DM 24h. There were 899 genes that were differentially expressed at all timepoints. Gene

ontology (GO) analysis was performed on DEGs to cluster genes into function-based categories

(48, 49) and the complete results are listed in Supp. Table 3. GO analysis of genes downregulated

at DM 48h showed that the top 10 biological process categories were related to skeletal muscle

contraction and skeletal muscle tissue development (Fig 3c; Supp. Table 3). This is in agreement

with our experimental results, thus identifying the importance of bromodomain function in

myogenesis. The top 10 categories from GO analysis of genes upregulated at DM 48h were related

to cell proliferation (Fig 3d; Supp. Table 3), which indicated altered proliferation due to PFI-3-

induced bromodomain inhibition. The promoters of the differentially expressed genes were also

analyzed using the HOMER motif enrichment software (49). Sequences 1kb upstream of the

transcription start sites were searched for presence of known consensus motifs (Supp. Table 4).

The analysis revealed that promoters of genes downregulated due to PFI-3 treatment were

significantly enriched in motifs corresponding to muscle specific transcription factors from the

MEF and MRF families. (Fig 3c; Supp. Table 4). In the case of upregulated genes, HOMER

analysis identified enrichment of motifs known to be bound by E2F family, NFY, KLF5 and Sp1

transcription factors (Fig 3d; Supp. Table 4). E2F and KLF5 TF families are known to play key

role in regulation of cell proliferation and differentiation (50–52). Thus, PFI-3 induced

bromodomain inhibition affects expression of genes which are involved in regulation of cell

proliferation and skeletal muscle differentiation.

PFI-3 treatment blocked cell-cycle exit of C2C12 cells induced for differentiation

Cell cycle exit is prerequisite for cellular differentiation to proceed in a number of cell types (53–

55). Results from the GO analysis of upregulated genes at DM 48h indicated that inhibition of

bromodomain function may interfere with cell cycle exit. To experimentally address the

requirement of mSWI/SNF bromodomain function in cell cycle exit, a BrdU (5-bromo-

deoxyuridine) incorporation assay was performed with samples treated with or without PFI-3 (Fig

4a). Confocal microscopy analysis showed that cells treated with PFI-3 continued to incorporate

BrdU even after the control cells showed no further incorporation, indicating a partial inability to

exit cell cycle. Quantification of these images is shown in Fig 4b. Increased mRNA expression of

cyclin A2, cyclin B1, cyclin D1, and cyclin D2 in PFI-3 treated samples from DM 48h as compared

to DMSO controls further correlates with continued cell cycle (Fig 4c). These results show that

bromodomain inhibition allows some of the myoblasts to overcome the signals to exit cycle that

are normally provided by the low mitogen media and by contact inhibition. Thus, PFI-3 induced

bromodomain inhibition may be affecting two aspects of myogenesis: timely exit from the cell

cycle and the expression of myogenic genes.

BAF180 is dispensable for C2C12 myoblast differentiation

The composition of mSWI/SNF complexes is variable depending on function, cell-type and

context. Every functional mSWI/SNF complex contains either the BRG1 or the BRM ATPase,

while one major sub-class of mSWI/SNF complexes also contains BAF180 (10, 13). Thus, PFI-3

treatment affects all mSWI/SNF complexes. BRG1 and BRM have been shown to be required for

skeletal muscle differentiation (25, 26, 28, 30, 56, 57), but the requirement for BAF180 in this

process has not been evaluated.

We knocked down BAF180 using siRNA. C2C12 cells depleted for BAF180 were induced for

differentiation alongside cells with scrambled siRNA treatment (Fig 5a, 5b). The cells showed no

phenotypic defect and differentiated normally. This result suggests that BAF180 may be

dispensable for myogenesis and implies that PFI-3 induced inhibition of myoblast differentiation

is mediated through inactivation of BRG1 and/or BRM bromodomain function.

Gene targets of PFI-3 inhibition of mSWI/SNF bromodomains predominantly overlap with

targets of BRG1 knockdown during myogenesis

The importance of BRG1 and BRM in skeletal muscle differentiation has been shown previously

by multiple groups. These studies have looked at muscle-specific gene expression profiles and

promoter binding of selected mSWI/SNF subunits on myogenic regulatory sequences. In a recently

published study, the authors performed siRNA-mediated knockdown of BRG1 in C2C12 cells

differentiated for 48h followed by RNA-sequencing analysis (58). We compared the siBRG1

dataset from this study with our RNA-sequencing dataset generated from PFI-3 treated C2C12

cells differentiated under similar conditions (Fig 6a). The rationale behind this comparison was to

get an understanding of the relative importance of the BRG1 bromodomain. The analysis showed

that 46% of the gene targets downregulated due to PFI-3 treatment overlapped with downregulated

genes in the siBRG1 dataset (Fig 6a). Similarly, 46% of the upregulated genes due to PFI-3

inhibition were common with upregulated genes from siBRG1 dataset. The results show that a

subset of BRG1-dependent gene expression in differentiating myoblasts requires bromodomain

function. GO analysis of the common overlapping genes was conducted. Common downregulated

genes belonged to muscle differentiation related processes while the common upregulated targets

fell into cell-cycle related categories. (Fig 6a-b; Supp. Table 5). Promoters of the common

upregulated and downregulated genes were also analyzed using HOMER to search for the presence

of known consensus motifs within 1kb upstream of their TSS (Fig 6a-b; Supp. Table 5). Promoters

of upregulated genes contained motifs known to be bound by E2F, KLF, NFY and Sp1 TF families

while those of downregulated genes were enriched for motifs corresponding to MEF and MRF

muscle-specific transcription factor families (Fig 6a-b; Supp. Table 5). Thus, the results from the

GO and HOMER motif enrichment analyses of overlapping genes and their promoters are similar

to those from PFI-3 treatment as shown in Fig 3c-d and therefore point towards a crucial role

played by BRG1 bromodomain in skeletal muscle differentiation and cell cycle regulation.

We examined the genes that were dependent on BRG1 for expression but independent of PFI-3-

mediated inhibiton of bromodomain function (Supp. Fig 3; Supp. Table 6). Genes that are up- and

down-regulated predominantly represented targets involved in metabolic processes and do not

include genes that control skeletal muscle differentiation or control of cell cycle. This suggests

that bromodomain-dependent regulation of gene expression is critical for myogenesis. This result

also is consistent with prior studies showing that ATPase domain function is required for BRG1-

mediated regulation of metabolism (59, 60) and that PFI-3 treatment did not affect cancer cell

proliferation dependent on BRG1 and/or BRM (40, 60).

Microarray analysis of gene expression in C2C12 cells upon siRNA-mediated knockdown of

BRG1 or BRM has also been done in a prior study by Albini et. al (30). Despite the difference in

methodologies, we overlapped the DEGs from that study with our RNA-seq data from PFI-3

treated C2C12 cells at comparable timepoints (48h post-differentiation) to evaluate if the outcome

from this comparison is in consonance with the previous results. The analysis showed that more

than one-third of the genes identified by Albini et al. as unique BRG1 targets overlapped with

differentially expressed genes from our PFI-3 RNA-seq dataset (Supp Fig 4a). Upon looking

individually at upregulated and downregulated targets uniquely regulated by BRG1, there was a

39% and 34% overlap, respectively (Supp Fig 4b). These common overlapping target genes were

then characterized using GO analysis (Supp Fig 4a-b). Upregulated genes fell into cell-cycle

related categories, and downregulated genes belonged to muscle differentiation related categories

(complete analysis in Supp. Table 7). This is in agreement with the overall results from PFI-3

treatment indicating the importance of active bromodomain for BRG1 function. Of the genes

identified by Albini et al. as unique BRM targets, only about 20% were common with PFI-3 DEGs

(Supp Fig 4b). These numbers show that more genes may be regulated by the BRG1 bromodomain

as compared to the BRM counterpart. Albini et al. also identified a set of genes that were

coregulated by both BRG1 and BRM. About 32% of these upregulated targets and 16% of the

downregulated targets were found in the corresponding list of differentially expressed genes due

to PFI-3 induced bromodomain inhibition (Supp Fig 4b; Supp. Table 7). This comparative analysis

further confirms that both BRG1 and BRM bromodomains play a role in skeletal myogenesis.

PFI-3 inhibition of mSWI/SNF bromodomains decreased binding of BRG1 and BRM to

target gene promoters.

BRG1-containing mSWI/SNF complexes are recruited to regulatory regions of myogenic genes

upon induction of muscle differentiation (25, 26, 29, 30, 61, 62). This step is required to induce

remodeling of chromatin at myogenic gene loci, thus allowing activation of muscle-specific gene

expression.

We performed ChIP experiments to determine whether the occupancy of BRG1 was affected at

the myogenic gene regulatory regions in response to bromodomain inhibition by PFI-3. As

expected, BRG1 occupancy at myogenic regulatory regions increased as a function of

differentiation in DMSO-treated samples (Fig 7a). However, BRG1 occupancy at the tested

regulatory sequences was partly inhibited in PFI-3 treated cells. This loss of binding correlates

with the decreased expression of these genes as seen in previous results (Fig. 2). We also looked

at binding of BRG1 and BRM on cyclin D1 and cyclin D2 promoters. It is known that in C2C12

myoblasts differentiated for 48h, BRM binding to the cyclin D1 gene promoter is crucial for its

repression and cell-cycle exit (30). In that study, the authors showed that the expression of cyclin

D1 was co-regulated by both BRG1 and BRM during the later stages of differentiation (30). In

MCF-7 cells, BRG1 has been shown to bind to cyclin D1 promoter and regulate its expression

(63). In PFI-3 treated C2C12 cells, chromatin IPs at the cyclin D1 promoter showed a decrease in

occupancy of BRM and BRG1 (Fig 7b). Similarly, we saw a significant decrease in binding of

BRG1 and BRM at the cyclin D2 promoter due to PFI-3 treatment. These results show that the

bromodomain function of BRG1 and BRM contributes to their binding at target gene promoters.

DISCUSSION

Bromodomains in mSWI/SNF proteins

Bromodomains are a conserved structural motif found in only 46 human proteins, and they are

classified into eight families (38). Bromodomains bind to acetylated lysines, which facilitates

protein-protein interactions (37). The ability of bromodomains to target proteins to acetylated

nucleosomes containing acetylated histones has been predicted to be a mechanism by which

chromatin epigenetic modifications are read, thereby enabling translation of the histone mark via

the bromodomain protein or via proteins associated with the bromodomain-containing protein (37,

64, 65). Family VIII bromodomains include the 6 bromodomains found in the N-terminal portion

of the BAF180 protein, the bromodomains present in the BRG1 and BRM ATPases, and the

bromodomain found in the histone lysine N-methyltransferase ASH1L (38).

Work to date on mSWI/SNF bromodomains has consisted of structural analyses (66–70), in vitro

studies of bromodomain binding to histones, DNA and nucleosomes (67, 70–74) and a limited

number of functional analyses in higher eukaryotes (75, 76). Of particular note, deletion of the

bromodomain in *Drosophila* BRM, which is the only SWI/SNF ATPase, had no effect on

developing or adult flies (77). In human cells, reconstitution of BRG1-deficient tumor cells with

wildtype or mutant versions of BRG1 determined that sequences C-terminal to the ATPase

domain, which includes the bromodomain, were not required for BRG1-mediated co-activation

of transcription by the glucocorticoid receptor (78). Similarly, BRG1 containing a bromodomain

mutation was capable of co-activating myocardin to promote smooth muscle-specific gene

expression (79). However, the BRG1 bromodomain directly contributes to the ability of

Repressor Element 1-silencing Transcription Factor (REST) to bind chromatin and repress target

genes (80). Mutation of each of the six BAF180 bromodomains revealed that four of the six

promoted tumor suppressor function, gene regulation, and chromatin affinity in clear cell renal

cell carcinoma cells (81). Other work showed that the 4[th] bromodomain of BAF180 mediated

interaction with acetylated p53, which promotes p53 binding to and transcriptional activation of

its target promoters (82). Thus, the requirement for functional mSWI/SNF protein

bromodomains is variable. It may be cell-type dependent as well. Here we demonstrate that the

BRG1 and/or BRM bromodomains, but not the BAF180 bromodomains, contribute to myogenic

differentiation. Both BRG1 and BRM are required for myogenic differentiation; this work is the

first to identify a contribution to myogenesis by any domain other than the ATPase domain. The

work further supports the idea of context-dependent requirements for mSWI/SNF bromodomain

functions.

Use of PFI-3 to probe biological function of mSWI/SNF subunits containing bromodomains

Efforts to identify pharmacological inhibitors of bromodomains identified salicylic acid as a specific interactor of BRG1, BRM, and BAF180 bromodomains (41, 83). This led to a series of structure-guided design steps that resulted in the PFI-3 inhibitor that is specific for the second and fifth bromodomain of BAF180 and the bromodomains of BRG1 and BRM (41, 42).

To date PFI-3 has been used to probe biological function in a number of contexts. Because BRG1 and other subunits of the mSWI/SNF enzyme have been shown to be required for proliferation of some cancer cells (84–86), PFI-3 was tested for inhibitory effects on cancer cell proliferation, with no effect observed on many different cancer cell types, including the NCI-60 tumor cell panel (40, 41, 87). These results demonstrate that mSWI/SNF bromodomains are not required for cancer cell proliferation. An inhibitory effect of PFI-3 was observed in PTEN-depleted prostate cancer cells in culture, in xenografts and in PTEN deficient mouse model susceptible to prostate cancer (88). PTEN is a tumor suppressor that normally regulates the AKT/PKB signaling pathway (89). Thus, the importance of mSWI/SNF bromodomains, and specifically, the BRG1 bromodomain, is enhanced in the absence of PTEN and inhibition of AKT/PKB signaling, suggesting a complex mode of regulation of mSWI/SNF protein bromodomain function. PFI-3 treatment increased the accessibility of an mSWI/SNF-repressed promoter and its gene expression (90). In these and other studies, PFI-3 altered both gene expression patterns dependent on mSWI/SNF bromodomain-containing proteins and the cellular and organismal phenotypes controlled by those genes (87, 88, 91–96). The data indicate that the pleiotropic effects of PFI-3 and inhibition of mSWI/SNF bromodomains links to the ability of the chromatin remodeling enzyme to modulate gene expression.

In the realm of tissue specification, PFI-3 treatment caused a loss of "stemness" and promoted

differentiation of ESCs and trophoblast and neural stem cells in the absence of differentiation

signaling (41, 97). In other contexts, PFI-3 inhibited differentiation, blocking the ability of

myoblasts and pre-adipocytes to form myotubes and adipocytes, respectively, in the presence of

differentiation signaling (42). In this report, we investigated the mechanisms responsible for the

inhibitory effects of PFI-3 on myogenesis. PFI-3 treatment affected bromodomain function of

BRG1 and BRM, but BAF180, as BAF180 was dispensable for differentiation. BRG1/BRM

bromodomain function was required for appropriate regulation of cell cycle withdrawal and

initiation of the myogenic gene expression program, with mis-regulation of a subset of the genes

regulated by BRG1 and BRM. Deficient gene regulation was linked to the partial inhibition of the

ability of BRG1 and BRM to bind to target gene regulatory sequences, reflecting a necessary

contribution of these bromodomains to promote interaction of the mSWI/SNF enzymes with

chromatin.

BAF180 is dispensable for myogenesis

mSWI/SNF complexes are a family of enzyme complexes marked by diversity of subunit

composition (6, 11, 98). Initial descriptions of mSWI/SNF complexes reported two separable

biochemical fractions that showed ATP-dependent chromatin remodeling activity (8–10). These

complexes have become known as BAF (BRG1/BRM-associated factors) and PBAF

(Polybromo-associated BAF), the latter taking its name from the presence of the BAF180 protein

that is specific to this complex. However, both BAF and PBAF complexes themselves are merely

separable groups of complexes that contain both shared and unique subunits (99, 100). A third

family of complexes, called ncBAF (noncanonical BAF) was identified more recently (13, 101).

Although there are many subtypes of functional mSWI/SNF complexes in the cell, all of them

necessarily have at least one of the proteins from the family VIII of bromodomains (102). BAF

and ncBAF complexes contain either BRG1 or BRM, while PBAF complexes contain BRG1 and

BAF180 (13). BAF180 has been implicated in DNA damage repair (103, 104) and is also

required for cardiac development (105, 106); knockout in mice caused severe hypoplastic

ventricle development and trophoblast placental defects (105). However, adult mice with

BAF180 depletion were phenotypically normal except for a hematopoietic stem cell defect

observed in aged mice (107). Similarly, BAF180 shows tumor-suppressive properties in some

but not all cancer cell lines (107–111). These findings are consistent with context-specific

requirements for the BAF180 protein.

In our work, we used siRNA-mediated knockdown to show that BAF180 is dispensable for

skeletal muscle differentiation. This suggests that the PBAF family of mSWI/SNF complexes are

also dispensable in this differentiation program. Although there are many reports characterizing

the requirement for mSWI/SNF complexes in myogenesis, focus has been limited to the two

ATPase subunits, to BAF47/INI1, which is shared by BAF and PBAF complexes, and to the

BAF60 subunit that is shared by all subfamilies of mSWI/SNF complexes (25–27, 29–31, 56, 57,

112–125). A prior report documented the binding of BAF250A to myogenic promoters (115),

perhaps implicating BAF complexes as the relevant mSWI/SNF enzyme subfamily for myogenic

differentiation, but the requirement for BAF250A was not evaluated. Nevertheless, a requirement

for specialized complexes for specific gene regulation events is one of the main hypotheses for

existence of diverse families of mSWI/SNF complexes.

BRG1 and BRM bromodomain function contribute to the regulation of myogenic

differentiation

BRG1 and BRM contribute to the activation of the myogenic gene expression program and BRM

contributes to the cell cycle arrest of myoblasts that precedes differentiation (25–30, 56). Inhibition

of the mSWI/SNF bromodomains by PFI-3 recapitulated these findings, indicating that the

bromodomains of BRG1 and BRM are needed to both regulate cell cycle exit and for the initiation

of tissue-specific gene expression. RNA-seq analysis of PFI-3 treated cells provided evidence of

global disruption of the regulation of cell cycle exit and the initiation of myogenic gene expression.

A recently published RNA-seq study investigating the role of chromatin remodeling in skeletal

myogenesis performed knockdown of BRG1 and evaluated gene expression at timepoints

comparable with our study (58). Comparison of this dataset with ours identified a large overlapping

subset of gene targets involved in cell cycle exit and myogenesis indicating that BRG1

bromodomain plays a crucial role in regulation of BRG1-dependent events in skeletal muscle

differentiation.

Additionally, prior microarray-based studies of myogenic gene expression upon BRG1 and BRM

knockdown also identified these processes as being BRG1- and BRM-dependent (30). Despite the

differences in platforms, we integrated the two datasets. There was a considerably greater overlap

between PFI-3 affected genes and genes mis-regulated by BRG1 knockdown than there was

between PFI-3 affected genes and genes mis-regulated by BRM knockdown. This may suggest

that a greater percentage of genes that require BRG1 are also dependent on the BRG1

bromodomain than is true for the set of genes that require BRM.

Regardless, our ChIP experiments demonstrated that bromodomain inhibition resulted in a

decreased ability of BRG1 to bind to genes activated during the myogenic differentiation protocol

and a decreased ability of BRG1 and BRM to bind to genes controlling cell cycle. The chromatin

interacting properties of the BRG1 and BRM bromodomains therefore likely contribute to gene

expression. The principles determining the variable requirements for BRG1, BRM and BAF180

bromodomains in different cellular contexts remain to be investigated. However, many of the

mSWI/SNF subunits have additional domains that could promote chromatin interaction. For

instance, BRG1 and BRM proteins have AT hooks, BAF180 and the BAF57 protein have an HMG

box-like domain, the ARID1A/1B and ARID2 proteins have ARID domains that may mediate

nucleic acid interactions, and BRD7 and BRD9 also have bromodomains. The requirement for any

of these domains may be dependent on the protein makeup of the particular mSWI/SNF complex

and the presence or absence of these other domains. Such a scenario suggests that these largely

non-sequence specific DNA-binding domains function in an additive or cooperative manner to

help facilitate chromatin interactions and remodeling events. Additional characterization of the

putative chromatin interactions domains in the context of differentiation will be necessary to

determine whether they are required and act in concert to promote mSWI/SNF interaction with

chromatin and function.

MATERIALS AND METHODS

Antibodies and Chemicals

Antibodies were purchased from Santa Cruz Biotech, USA (anti-BRG1, sc-17796; anti-BrdU, sc-

32323). Myosin Heavy chain antibody (#MF20) was purchased from the Developmental Studies

Hybridoma Bank, University of Iowa, USA. BRM antisera was described previously (126). Lysis

buffers for ChIP assays were purchased from Cell Signaling Technology, USA (SimpleChIP®

Enzymatic Cell Lysis Buffers A & B, 14282; SimpleChIP® Chromatin IP Buffers, 14231).

Dulbecco's modified Eagle's medium (DMEM) was purchased from ThermoFisher Scientific

(#11965118).

Cell culture

C2C12 cells were purchased from ATCC (Manassas, VA) and maintained at sub-confluent

densities in DMEM supplemented with 10% FBS and 1% penicillin/streptomycin in a humidified

incubator at 37°C in 5% CO_2.

Mouse satellite cells were isolated from leg muscles of 2 week old C57BL6/J mice using Percoll

sedimentation followed by differential plating as described previously (114). Mice were housed in

the animal care facility at the University of Massachusetts Medical School and used in accordance

with a protocol approved by the Institutional Animal Care and Use Committee (IACUC).

For differentiation, cells at > 70% confluency were switched to DMEM medium supplemented

with 2% horse serum and 2 μg/ml of bovine insulin (Sigma-Aldrich, St. Louis, MO). Where

indicated, cells were pre-treated with DMSO or PFI-3 (Cayman Chemical, Ann Arbor, MI) for

24h before inducing differentiation. PFI-3 was maintained during the course of the experiment at

50μM and the medium was replaced every 24 hours.

siRNA transfection

C2C12 cells were plated on 24-well plates in DMEM medium 24h before transfection. Cells were

transfected at 30-40% confluence using the Lipofectamine 2000 (Invitrogen) reagent with 50 nM

siRNA SMARTpool ON-TARGETplus (Dharmacon, Scrambled non-targeting Pool # D-001810-

10-20 and Pbrm1 #L-044781-00-0005). 48h post-transfection, the cells were induced for

differentiation and samples were harvested at indicated times for further analysis.

Immunocytochemistry

Cells were seeded on 22mm x 22mm size coverslips in 35mm dishes and were harvested after the

indicated treatments at the specified timepoints. The samples were washed with PBS twice and

then fixed with ice-cold fixative (2% formaldehyde and 0.2% glutaraldehyde in PBS) for 10

minutes on ice. The cells were washed with PBS twice and permeabilized with ice-cold

permeabilization buffer (0.2% Triton-X 100 in PBS) for 5 minutes on ice. Samples were then

washed once with PBS and blocked using freshly prepared blocking reagent (5% BSA in PBS) for

30 minutes at room temperature (RT). The cells were washed twice with PBS and incubated with

primary antibody cocktail diluted to the desired concentration in blocking reagent (2% BSA in

PBS) for 2 hours at room temperature. Post-incubation, samples were washed thrice with PBS 10

minutes each to remove non-specific binding. Cells were then incubated with fluorophore-

conjugated secondary antibody (1:100) and DAPI (2μg/ml), both diluted in blocking reagent (2%

BSA in PBS) for 45 minutes at RT followed by 3 washes with PBS to remove non-specific staining.

The stained cells on coverslips were then inverted-mounted on glass slides in 70% glycerol and

the sides were sealed with nail paint. Confocal imaging analysis was performed using Leica TCS

SP5 II laser scanning confocal microscope and analyzed with Leica Lite software.

Fusion Index

For calculation of fusion index, cells were harvested at the specified timepoints after the indicated

treatments and were immunostained with myosin heavy chain (MF20, DSHB) and DAPI (nuclear

staining) as described above. The images were captured at 40X magnification using a Leica TCS

SP5 II laser scanning confocal microscope. Analysis was performed by scoring cells for number

of nuclei and MHC staining. Fusion index was calculated as the percentage of nuclei/cells stained

with myosin heavy chain as compared to total number of nuclei/cells (43).

RIPA buffer Lysis

Cells were harvested after the indicated treatments at specific timepoints and were washed twice

with ice-cold PBS. After draining all residual PBS, cells were scraped into 1ml ice-cold PBS with

1X protease inhibitor cocktail (Sigma Aldrich, P8340) and pelleted at 1500 X g for 5 minutes at

4°C. The pellets were lysed in 500ul RIPA buffer (50 mM Tris-HCl, pH7.4, 150 mM NaCl, 1mM

EDTA, 1% NP-40 and 0.25% sodium deoxycholate) supplemented with 1X protease inhibitor

cocktail (Sigma Aldrich, P8340). Samples were incubated on ice for 30 minutes and whole cell

extracts were prepared by passing the lysed pellets through a 27-gauge needle at least 4-5 times.

Samples were centrifuged at 14000 X g for 10 minutes at 4°C and supernatants were collected.

Western Blot Analysis

Protein concentrations were determined using a Pierce[TM] BCA protein assay kit (ThermoFisher

Scientific, USA) according to the manufacturer's protocol. Equal amounts of protein from each

sample were aliquoted and mixed with 4X Laemmli Sample Buffer (BioRad) and boiled at 95°C

for 10 minutes. The samples were electrophoresed on denaturing SDS-polyacrylamide gels and

transferred onto Immobilon-P PVDF membranes (Merck Millipore, USA). The membranes were

then blocked using 5% non-fat milk in PBS for 30 minutes followed by overnight incubation at

4°C with primary antibody against protein of interest at the desired dilution in 2% non-fat milk

prepared in PBS. This was followed by 3 washes with TBS containing 0.1% Tween-20 for 5

minutes each at room temperature. The membranes were then incubated with HRP-conjugated

anti-mouse or anti-rabbit secondary antibodies (1:2500, GE Healthcare Life Sciences) diluted in

2% non-fat milk prepared in PBS for 1 hour at RT followed by 3 washes with TBS containing

0.1% Tween-20 for 5 minutes each at room temperature. Chemiluminescent detection was

performed with ECL Plus (GE Healthcare Life Sciences) using an Amersham Imager 600 (GE

Healthcare Life Sciences). Representative blots from 3 independent experiments are shown. Band

signal intensities were quantified using ImageJ software (NIH) (127).

RNA isolation and quantitative RT-PCR

For RNA isolation, cells were grown in 35mm dishes and harvested after the indicated treatments

at specified timepoints. The media were removed, and cells were washed twice with PBS before

adding 1ml of Trizol (ThermoFisher) to each sample. RNA extraction was performed as per the

manufacturer's protocol. The final RNA pellet was resuspended in 50μl nuclease-free water. RNA

concentrations were quantified using a Nanodrop1000 spectrophotometer (ThermoFisher

Scientific). cDNA was prepared using 2μg of total RNA using Superscript III First Strand

Synthesis Kit (Invitrogen) according to manufacturer's protocol.

For qRT-PCR, 15μl reactions were prepared in duplicate for all desired samples using 1μl each of

forward and reverse primers (10μM stocks) and 2μl cDNA template, and the volume was brought

to 7.5μl using UltraPure distilled water. 7.5μl of Fast SYBR Green 2X Master Mix (Applied

Biosystems) was added to the reaction. The samples were run using the default protocol in

QuantStudio 3 RT-PCR machine (Applied Biosystems). Fold-changes were calculated using the

$2^{-\Delta\Delta Ct}$ method (128). Primer sequences are listed in Supp. Table 8.

RNA-sequencing analysis

For RNA sequencing, RNA samples were prepared as described above. Duplicate samples for each

timepoint were evaluated for quality and concentration at the UMass Medical School MBCL

Fragment Analyzer services. QC-approved samples were sent to BGI Americas Corporation for

library preparation and RNA sequencing (129). Libraries were sequenced using the BGISEQ-500

platform and reads were filtered to remove adaptor-polluted, low quality and high content of

unknown base reads. About 99% of the raw reads were identified as clean reads (~65M). Of these,

about 70% of the reads were uniquely mapped to mouse reference genome mm10 using HISAT

(130). Transcripts were reconstructed using StringTie (131), and novel transcripts were identified

using Cufflinks (132). All transcripts were then combined and mapped to the mm10 reference

transcriptome using Bowtie2 (133). Gene expression levels were calculated using RSEM (134).

DEseq2 (135) and PoissonDis (136) algorithms were used to detect differentially expressed genes

(DEG). GO analysis was performed on DEGs to cluster genes into function-based and pathway-

based categories (48, 49). Motif analysis was performed using HOMER motif discovery software

as described previously (49). For each differentially expressed gene, motif enrichment analysis

was performed at promoters for locations up to 1 kb upstream of the TSS.

BrdU incorporation and immunostaining

Cell were grown on coverslips in 35mm dishes with the indicated treatments. The culture media

was replaced with fresh media containing 10mM BrdU 30 minutes prior to harvesting to allow the

cells in S-phase to incorporate the nucleoside analog. Harvested samples were fixed and

permeabilized as described above for immunocytochemistry. For DNA hydrolysis, samples were

then incubated with 1.2N HCl diluted in PBS for 1 hour at 37°C, followed by neutralization in

0.1M sodium borate for 5 minutes at RT. The samples were washed with PBS thrice and

immunostained as mentioned above.

Chromatin Immunoprecipitation assay

Chromatin immunoprecipitation assays were performed as described previously (115).

Quantification was performed using the fold enrichment method ($2^{-(Ct\ sample\ -\ Ct\ IgG)}$) and shown as

relative to a control region. Sequences of primers used for ChIP assays are listed in Supp. Table 8.

Statistical analysis

All quantitative data for gene expression and chromatin immunoprecipitation are shown as mean

+/- the standard deviation of at least three independent biological replicates. In the fusion index

and BrdU incorporation assays, two independent experiments were performed in duplicate. A

minimum of 200 nuclei per sample were counted and the results were expressed as the mean of

calculated fusion index +/- standard deviation. Statistical analyses were performed using Graphpad

Prism8 Student's t-test with two-tailed distribution and equal variance (Graphpad Prism Software

Inc.). Significance is displayed with $*p<0.05$, $**p<0.01$ and $***p<0.005$.

Data Availability

The data from this RNA-seq study has been deposited in NCBI's Gene Expression Omnibus and

are accessible through GEO accession number GSE151218.

ACKNOWLEDGEMENTS

We thank T Padilla-Benavides, S Syed, and J Dilworth for comments and suggestions. This work

was funded by NIH grants GM56244 and GM136393 to ANI.

REFERENCES

1. Allshire RC, Madhani HD. 2018. Ten principles of heterochromatin formation and

function. Nat Rev Mol Cell Biol 19:229–244.

2. Janssen A, Colmenares SU, Karpen GH. 2018. Heterochromatin: Guardian of the

Genome. Annu Rev Cell Dev Biol 34:265–288.

3. Hota SK, Bruneau BG. 2016. ATP-dependent chromatin remodeling during mammalian

development. Development 143:2882–2897.

4. De La Serna IL, Ohkawa Y, Imbalzano AN. 2006. Chromatin remodelling in mammalian

differentiation: Lessons from ATP-dependent remodellers. Nat Rev Genet 7:461–473.

5. Clapier CR, Cairns BR. 2009. The Biology of Chromatin Remodeling Complexes. Annu

Rev Biochem 78:273–304.

6. Hargreaves DC, Crabtree GR. 2011. ATP-dependent chromatin remodeling: genetics,

genomics and mechanisms. Cell Res 21:396–420.

7. Wu JI. 2012. Diverse functions of ATP-dependent chromatin remodeling complexes in

development and cancer. Acta Biochim Biophys Sin 44:54–69.

8. Kwon H, Imbalzano AN, Khavari PA, Kingston RE, Green MR. 1994. Nucleosome

disruption and enhancement of activator binding by a human SW1/SNF complex. Nature

370:477–481.

9. Imbalzano AN, Kwon H, Green MR, Kingston RE. 1994. Facilitated binding of TATA-

binding protein to nucleosomal DNA. Nature 370:481–485.

10. Wang W, Côté J, Xue Y, Zhou S, Khavari PA, Biggar SR, Muchardt C, Kalpana G V.,

Goff SP, Yaniv M, Workman JL, Crabtree GR. 1996. Purification and biochemical

heterogeneity of the mammalian SWI-SNF complex. EMBO J 15:5370–5382.

11.   Wang W, Xue Y, Zhou S, Kuo A, Cairns BR, Crabtree GR. 1996. Diversity and

586   specialization of mammalian SWI/SNF complexes. Genes Dev 10:2117–2130.

587   12.   Wu JI, Lessard J, Olave IA, Qiu Z, Ghosh A, Graef IA, Crabtree GR. 2007. Regulation of

588   Dendritic Development by Neuron-Specific Chromatin Remodeling Complexes. Neuron

589   56:94–108.

590   13.   Mashtalir N, D'avino AR, Michel BC, Cassel SH, Ranish JA, Kadoch Correspondence C.

591   2018. Modular Organization and Assembly of SWI/SNF Family Chromatin Remodeling

592   Complexes. Cell 175:1272-1288.e20.

593   14.   Chal J, Pourquié O. 2017. Making muscle: skeletal myogenesis in vivo and in vitro.

594   Development 144:2104–2122.

595   15.   Comai G, Tajbakhsh S. 2014. Molecular and cellular regulation of skeletal myogenesis.

596   Curr Top Dev Biol, 1st ed. 110:1–73.

597   16.   Motohashi N, Asakura A. 2014. Muscle satellite cell heterogeneity and self-renewal. Front

598   Cell Dev Biol 2:1–14.

599   17.   Chang NC, Rudnicki MA. 2014. Satellite Cells: The Architects of Skeletal Muscle. Curr

600   Top Dev Biol 107:161–181.

601   18.   Sambasivan R, Tajbakhsh S. 2015. Adult skeletal muscle stem cells. Results Probl Cell

602   Differ 56:191–213.

603   19.   Hernández-Hernández JM, García-González EG, Brun CE, Rudnicki MA. 2017. The

604   myogenic regulatory factors, determinants of muscle development, cell identity and

605   regeneration. Semin Cell Dev Biol 72:10–18.

606   20.   Moncaut N, Rigby PWJ, Carvajal JJ. 2013. Dial M(RF) for myogenesis. FEBS J

607   280:3980–3990.

21. Asfour HA, Allouh MZ, Said RS. 2018. Myogenic regulatory factors: The orchestrators of

myogenesis after 30 years of discovery. Exp Biol Med 243:118–128.

22. Blackwell T, Weintraub H. 1990. Differences and similarities in DNA-binding

preferences of MyoD and E2A protein complexes revealed by binding site selection.

Science 250:1104–1110.

23. Taylor M V., Hughes SM. 2017. Mef2 and the skeletal muscle differentiation program.

Semin Cell Dev Biol 72:33–44.

24. Pon JR, Marra MA. 2016. MEF2 transcription factors: Developmental regulators and

emerging cancer genes. Oncotarget 7:2297–2312.

25. de la Serna IL, Ohkawa Y, Berkes C a, Bergstrom D a, Dacwag CS, Tapscott SJ,

Imbalzano AN. 2005. MyoD targets chromatin remodeling complexes to the myogenin

locus prior to forming a stable DNA-bound complex. Mol Cell Biol 25:3997–4009.

26. Ohkawa Y, Tachibana T, Imbalzano AN, Yoshimura S, Higashi C, Marfella CGA,

Dacwag CS, Tachibana T, Imbalzano AN. 2007. Myogenin and the SWI / SNF ATPase

Brg1 Maintain Myogenic Gene Expression at Different Stages of Skeletal Myogenesis *. J

Biol Chem 282:6564–6570.

27. Simone C, Forcales SV, Hill DA, Imbalzano AN, Latella L, Puri PL. 2004. p38 pathway

targets SWI-SNF chromatin-remodeling complex to muscle-specific loci. Nat Genet

36:738–743.

28. Ohkawa Y, Marfella CGA, Imbalzano AN. 2006. Skeletal muscle specification by

myogenin and Mef2D via the SWI/SNF ATPase Brg1. EMBO J 25:490–501.

29. Forcales S V., Albini S, Giordani L, Malecova B, Cignolo L, Chernov A, Coutinho P,

Saccone V, Consalvi S, Williams R, Wang K, Wu Z, Baranovskaya S, Miller A, Dilworth

FJ, Puri PL. 2012. Signal-dependent incorporation of MyoD-BAF60c into Brg1-based

SWI/SNF chromatin-remodelling complex. EMBO J 31:301–316.

30. Albini S, Toto PC, Dall'agnese A, Malecova B, Cenciarelli C, Felsani A, Caruso M,

Bultman SJ, Puri PL. 2015. Brahma is required for cell cycle arrest and late muscle gene

expression during skeletal myogenesis. EMBO Rep 16:1037–1050.

31. de la Serna IL, Roy K, Carlson KA, Imbalzano AN. 2001. MyoD can induce cell cycle

arrest but not muscle differentiation in the presence of dominant negative SWI/SNF

chromatin remodeling enzymes. J Biol Chem 276:41486–91.

32. Khavari PA, Peterson CL, Tamkunt JW, Mendel DB, Crabtree GR. 1993. BRG1 contains

a conserved domain of the SWI2 / SNF2 family necessary for normal mitotic growth and

transcription. Lett to Nat 366:170–174.

33. Muchardt C, Yaniv M. 1993. A human homologue of Saccharomyces cerevisiae

SNF2/SWI2 and Drosophila brm genes potentiates transcriptional activation by the

glucocorticoid receptor. EMBO J 12:4279–4290.

34. Chiba H, Muramatsu1 M, Nomoto A, Kato H. 1994. Two human homologues of

Saccharomyces cerevisiae SWI2ISNF2 and Drosophila brahma are transcriptional

coactivators cooperating with the estrogen receptor and the retinoic acid receptor. Nucleic

Acids Res 22:1815–1820.

35. Meslamani J, Smith SG, Sanchez R, Zhou M-M. 2016. Structural features and inhibitors

of bromodomains. Drug Discov Today Technol 19:3–15.

36. Moustakim M, Clark PGK, Hay DA, Dixon DJ, Brennan PE. 2016. Chemical probes and

inhibitors of bromodomains outside the BET family. Medchemcomm 7:2246–2264.

37. Fujisawa T, Filippakopoulos P. 2017. Functions of bromodomain-containing proteins and

their roles in homeostasis and cancer. Nat Rev Mol Cell Biol 18:246–262.

38.   Filippakopoulos P, Picaud S, Mangos M, Keates T, Lambert JP, Barsyte-Lovejoy D,

Felletar I, Volkmer R, Müller S, Pawson T, Gingras AC, Arrowsmith CH, Knapp S. 2012.

Histone recognition and large-scale structural analysis of the human bromodomain family.

Cell 149:214–231.

39.   Xue Y, Canman JC, Lee CS, Nie Z, Yang D, Moreno GT, Young MK, Salmon ED, Wang

W. 2000. The human SWI/SNF-B chromatin-remodeling complex is related to yeast Rsc

and localizes at kinetochores of mitotic chromosomes. Proc Natl Acad Sci U S A

97:13015–13020.

40.   Vangamudi B, Paul TA, Shah PK, Kost-Alimova M, Nottebaum L, Shi X, Zhan Y, Leo E,

Mahadeshwar HS, Protopopov A, Futreal A, Tieu TN, Peoples M, Heffernan TP,

Marszalek JR, Toniatti C, Petrocchi A, Verhelle D, Owen DR, Draetta G, Jones P, Palmer

WS, Sharma S, Andersen JN. 2015. The SMARCA2/4 ATPase Domain Surpasses the

Bromodomain as a Drug Target in SWI/SNF-Mutant Cancers: Insights from cDNA

Rescue and PFI-3 Inhibitor Studies. Cancer Res 75:3865–3878.

41.   Fedorov O, Castex J, Tallant C, Owen DR, Martin S, Aldeghi M, Monteiro O,

Filippakopoulos P, Picaud S, Trzupek JD, Gerstenberger BS, Bountra C, Willmann D,

Wells C, Philpott M, Rogers C, Biggin PC, Brennan PE, Bunnage ME, Schüle R, Günther

T, Knapp S, Müller S. 2015. Selective targeting of the BRG/PB1 bromodomains impairs

embryonic and trophoblast stem cell maintenance. Sci Adv 1:1–11.

42.   Gerstenberger BS, Trzupek JD, Tallant C, Fedorov O, Filippakopoulos P, Brennan PE,

Fedele V, Martin S, Picaud S, Rogers C, Parikh M, Taylor A, Samas B, O'Mahony A,

Berg E, Pallares G, Torrey AD, Treiber DK, Samardjiev IJ, Nasipak BT, Padilla-

Benavides T, Wu Q, Imbalzano AN, Nickerson JA, Bunnage ME, Müller S, Knapp S,

Owen DR, Mahony AO, Berg E, Pallares G, Torrey AD, Treiber DK, Samardjiev IJ,

Nasipak BT, Padilla-Benavides T, Wu Q, Imbalzano AN, Nickerson A, Bunnage ME, Mu

S, Knapp S, Owen DR. 2016. Identification of a Chemical Probe for Family VIII

Bromodomains through Optimization of a Fragment Hit. J Med Chem 59:4800–4811.

43. Metzinger L, Passaquin AC, Warter JM, Poindron P. 1993. α-Methylprednisolone

promotes skeletal myogenesis in dystrophin-deficient and control mouse cultures.

Neurosci Lett 155:171–174.

44. Millay DP, O'Rourke JR, Sutherland LB, Bezprozvannaya S, Shelton JM, Bassel-Duby R,

Olson EN. 2013. Myomaker is a membrane activator of myoblast fusion and muscle

formation. Nature 499:301–5.

45. Leikina E, Gamage DG, Prasad V, Kozlov MM, Chernomordik L V, Millay

Correspondence DP, Goykhberg J, Crowe M, Diao J, Kozlov MM, Chernomordik L V,

Millay DP. 2018. Myomaker and Myomerger Work Independently to Control Distinct

Steps of Membrane Remodeling during Myoblast Fusion. Dev Cell 46:767–780.

46. Bi P, Ramirez-Martinez A, Li H, Cannavino J, McAnally JR, Shelton JM, Sánchez-Ortiz

E, Bassel-Duby R, Olson EN. 2017. Control of muscle formation by the fusogenic

micropeptide myomixer. Science 356:323–327.

47. Zhang Q, Vashisht AA, O'Rourke J, Corbel SY, Moran R, Romero A, Miraglia L, Zhang

J, Durrant E, Schmedt C, Sampath SC, Sampath SC. 2017. The microprotein Minion

controls cell fusion and muscle formation. Nat Commun 8:1–15.

48. Wei Huang D, Sherman BT, Lempicki RA. 2008. Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. Nucleic Acids Res 37:1–

13.

49.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H,

Glass CK. 2010. Simple Combinations of Lineage-Determining Transcription Factors

Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. Mol Cell

38:576–589.

50.   Müller H, Helin K. 2000. The E2F transcription factors: Key regulators of cell

proliferation. Biochim Biophys Acta - Rev Cancer 1470:1–12.

51.   Helin K. 1998. Regulation of cell proliferation by the E2F transcription factors. Curr Opin

Genet Dev 8:28–35.

52.   Dong JT, Chen C. 2009. Essential role of KLF5 transcription factor in cell proliferation

and differentiation and its implications for human diseases. Cell Mol Life Sci 66:2691–

2706.

53.   Soufi A, Dalton S. 2016. Cycling through developmental decisions: How cell cycle

dynamics control pluripotency, differentiation and reprogramming. Dev 143:4301–4311.

54.   Id YM, Mckay Id DJ, Buttitta Id L. 2019. Changes in chromatin accessibility ensure

robust cell cycle exit in terminally differentiated cells. Plos Biol 17:1–29.

55.   Politis PK, Thomaidou D, Matsas R. 2008. Coordination of cell cycle exit and

differentiation of neuronal progenitors. Cell Cycle 7:691–697.

56.   de la Serna IL, Carlson KA, Imbalzano AN. 2001. Mammalian SWI/SNF complexes

promote MyoD-mediated muscle differentiation. Nat Genet 27:187–190.

57.   Roy K, de la Serna IL, Imbalzano AN. 2002. The myogenic basic helix-loop-helix family

of transcription factors shows similar requirements for SWI/SNF chromatin remodeling

enzymes during muscle differentiation in culture. J Biol Chem 277:33818–24.

58.   Zhu X, Lan B, Yi X, He C, Dang L, Zhou X, Lu Y, Sun Y, Liu Z, Bai X, Zhang K, Li B,

Li MJ, Chen Y, Zhang L. 2020. HRP2-DPF3a-BAF complex coordinates histone

modification and chromatin remodeling to regulate myogenic gene transcription. Nucleic

Acids Res 48:6563–6582.

59.   Barutcu AR, Lajoie BR, Fritz AJ, McCord RP, Nickerson JA, Van Wijnen AJ, Lian JB,

Stein JL, Dekker J, Stein GS, Imbalzano AN. 2016. SMARCA4 regulates gene expression

and higherorder chromatin structure in proliferating mammary epithelial cells. Genome

Res 26:1188–1201.

60.   Wu Q, Madany P, Dobson JR, Schnabl JM, Sharma S, Smith TC, van Wijnen AJ, Stein

JL, Lian JB, Stein GS, Muthuswami R, Imbalzano AN, Nickerson JA. 2016. The BRG1

chromatin remodeling enzyme links cancer cell metabolism and proliferation. Oncotarget

7:38270–38281.

61.   Harada A, Okada S, Konno D, Odawara J, Yoshimi T, Yoshimura S, Kumamaru H,

Saiwai H, Tsubota T, Kurumizaka H, Akashi K, Tachibana T, Imbalzano AN, Ohkawa Y.

2012. Chd2 interacts with H3.3 to determine myogenic cell fate. EMBO J 31:2994–3007.

62.   Ohkawa Y, Mallappa C, Vallaster CSD, Imbalzano AN. 2012. Isolation of nuclei from

skeletal muscle satellite cells and myofibers for use in chromatin immunoprecipitation

assays. Methods Mol Biol 798:517–530.

63.   Rao M, Casimiro MC, Lisanti MP, Amico MD', Wang C, Shirley LA, Leader JE, Liu M,

Stallcup M, Engel DA, Murphy DJ, Pestell RG. 2008. Inhibition of cyclin D1 gene

transcription by Brg-1. Cell Cycle 7:647–655.

64.   Haynes SR, Dollard C, Winston F, Beck S, Trowsdale J, Dawid IB. 1992. The

bromodomain: a conserved sequence found in human, Drosophila and yeast proteins.

Nucleic Acids Res 20:246–257.

65.   Dhalluin C, Carlson JE, Zeng L, He C, Aggarwal AK, Zhou MM. 1999. Structure and

ligand of a histone acetyltransferase bromodomain. Nature 399:491–496.

66.   Singh M, Popowicz GM, Krajewski M, Holak TA. 2007. Structural ramification for

acetyl-lysine recognition by the bromodomain of human BRG1 protein, a central ATPase

of the SWI/SNF remodeling complex. ChemBioChem 8:1308–1316.

67.   Singh M, D'Silva L, Holak TA. 2006. DNA-binding properties of the recombinant high-

mobility-group-like AT-hook-containing region from human BRG1 protein. Biol Chem

387:1469–1478.

68.   Chandrasekaran R, Thompson M. 2007. Polybromo-1-bromodomains bind histone H3 at

specific acetyl-lysine positions. Biochem Biophys Res Commun 355:661–666.

69.   Shen W, Xu C, Huang W, Zhang J, Carlson JE, Tu X, Wu J, Shi Y. 2007. Solution

Structure of Human Brg1 Bromodomain and Its Specific Binding to Acetylated Histone

Tails. Biochemistry 46:2100–2110.

70.   Charlop-Powers Z, Zeng L, Zhang Q, Zhou MM. 2010. Structural insights into selective

histone H3 recognition by the human Polybromo bromodomain 2. Cell Res 20:529–538.

71.   Liao L, Alicea-Velázquez NL, Langbein L, Niu X, Cai W, Cho EA, Zhang M, Greer CB,

Yan Q, Cosgrove MS, Yang H. 2019. High affinity binding of H3K14ac through

collaboration of bromodomains 2, 4 and 5 is critical for the molecular and tumor

suppressor functions of PBRM1. Mol Oncol 13:811–828.

72.   Sanchez JC, Zhang L, Evoli S, Schnicker NJ, Nunez-Hernandez M, Yu L, Wereszczynski

J, Pufall MA, Musselman CA. 2020. The molecular basis of selective DNA binding by the

BRG1 AT-hook and bromodomain. Biochim Biophys Acta - Gene Regul Mech 1863:1–

14.

73.   Morrison EA, Sanchez JC, Ronan JL, Farrell DP, Varzavand K, Johnson JK, Gu BX,

771   Crabtree GR, Musselman CA. 2017. DNA binding drives the association of BRG1/hBRM

772   bromodomains with nucleosomes. Nat Commun 8:1–14.

773   74.   Slaughter MJ, Shanle EK, McFadden AW, Hollis ES, Suttle LE, Strahl BD, Davis IJ.

774   2018. PBRM1 bromodomains variably influence nucleosome interactions and cellular

775   function. J Biol Chem 293:13592–13603.

776   75.   Wang X, Wang S, Troisi EC, Howard TP, Haswell JR, Wolf BK, Hawk WH, Ramos P,

777   Oberlick EM, Tzvetkov EP, Vazquez F, Hahn WC, Park PJ, Roberts CWM. 2019. BRD9

778   defines a SWI/SNF sub-complex and constitutes a specific vulnerability in malignant

779   rhabdoid tumors. Nat Commun 10:1–11.

780   76.   Dutta A, Gogol M, Kim JH, Smolle M, Venkatesh S, Gilmore J, Florens L, Washburn

781   MP, Workman JL. 2014. Swi/Snf dynamics on stress-responsive genes is governed by

782   competitive bromodomain interactions. Genes Dev 28:2314–2330.

783   77.   Elfring LK, Daniel C, Papoulas O, Deuring R, Sarte M, Moseley S, Beek SJ, Ross

784   Waldrip W, Daubresse G, Depace A, Kennison JA, Tamkun JW. 1998. Genetic Analysis

785   of brahma : The Drosophila Homolog of the Yeast Chromatin Remodeling Factor

786   SWI2/SNF2. Genetics 148:251–265.

787   78.   Trotter KW, Fan H-Y, Ivey ML, Kingston RE, Archer TK. 2008. The HSA Domain of

788   BRG1 Mediates Critical Interactions Required for Glucocorticoid Receptor-Dependent

789   Transcriptional Activation In Vivo. Mol Cell Biol 28:1413–1426.

790   79.   Zhou J, Zhang M, Fang H, El-Mounayri O, Rodenberg JM, Imbalzano AN, Herring BP.

791   2009. The SWI/SNF chromatin remodeling complex regulates myocardin-induced smooth

muscle-specific gene expression. Arterioscler Thromb Vasc Biol 29:921–8.

80.   Ooi L, Belyaev ND, Miyake K, Wood IC, Buckley NJ. 2006. BRG1 chromatin

remodeling activity is required for efficient chromatin binding by repressor element 1-

silencing transcription factor (REST) and facilitates REST-mediated repression. J Biol

Chem 281:38974–38980.

81.   Porter EG, Dykhuizen EC. 2017. Individual bromodomains of Polybromo-1 contribute to

chromatin association and tumor suppression in clear cell renal carcinoma. J Biol Chem

292:2601–2610.

82.   Cai W, Su L, Liao L, Liu ZZ, Langbein L, Dulaimi E, Testa JR, Uzzo RG, Zhong Z, Jiang

W, Yan Q, Zhang Q, Yang H. 2019. PBRM1 acts as a p53 lysine-acetylation reader to

suppress renal tumor growth. Nat Commun 10:1–15.

83.   Fedorov O, Niesen FH, Knapp S. 2012. Kinase Inhibitor Selectivity Profiling Using

Differential Scanning Fluorimetry. Methods Mol Biol 795:109–118.

84.   Baylin SB, Jones PA. 2016. Epigenetic Determinants of Cancer. Cold Spring Harb

Perspect Biol 8:1–35.

85.   Feinberg AP, Tycko B. 2004. The history of cancer epigenetics. Nat Rev Cancer 4:143–

153.

86.   Zaidi SK, Van Wijnen AJ, Lian JB, Stein JL, Stein GS. 2013. Targeting deregulated

epigenetic control in cancer. J Cell Physiol 228:2103–2108.

87.   Wu Q, Sharma S, Cui H, LeBlanc SE, Zhang H, Muthuswami R, Nickerson JA,

Imbalzano AN. 2016. Targeting the chromatin remodeling enzyme BRG1 increases the

efficacy of chemotherapy drugs in breast cancer cells. Oncotarget 7:27158–75.

88.   Ding Y, Li N, Dong B, Guo W, Wei H, Chen Q, Yuan H, Han Y, Chang H, Kan S, Wang

815  X, Pan Q, Wu P, Peng C, Qiu T, Li Q, Gao D, Xue W, Qin J. 2019. Chromatin remodeling

816  ATPase BRG1 and PTEN are synthetic lethal in prostate cancer. J Clin Invest 129:759–

817  773.

818 89. Georgescu MM. 2010. Pten tumor suppressor network in PI3K-Akt pathway control.

819  Genes and Cancer 1:1170–1177.

820 90. Gillis NE, Taber TH, Bolf EL, Beaudet CM, Tomczak JA, White JH, Stein JL, Stein GS,

821  Lian JB, Frietze S, Carr FE. 2018. Thyroid Hormone Receptor β Suppression of RUNX2

822  Is Mediated by Brahma-Related Gene 1–Dependent Chromatin Remodeling.

823  Endocrinology 159:2484–2494.

824 91. Li Z, Xia J, Fang M, Xu Y. 2019. Epigenetic regulation of lung cancer cell proliferation

825  and migration by the chromatin remodeling protein BRG1. Oncogenesis 8:1–14.

826 92. Minderjahn J, Schmidt A, Fuchs A, Schill R, Raithel J, Babina M, Schmidl C, Gebhard C,

827  Schmidhofer S, Mendes K, Ratermann A, Glatz D, Nützel M, Edinger M, Hoffman P,

828  Spang R, Längst G, Imhof A, Rehli M. 2020. Mechanisms governing the pioneering and

829  redistribution capabilities of the non-classical pioneer PU.1. Nat Commun 11:1–16.

830 93. Li H, Lan J, Han C, Guo K, Wang G, Hu J, Gong J, Luo X, Cao Z. 2018. Brg1 promotes

831  liver fibrosis via activation of hepatic stellate cells. Exp Cell Res 364:191–197.

832 94. Li Z, Lv F, Dai C, Wang Q, Jiang C, Fang M, Xu Y. 2019. Activation of Galectin-3

833  (LGALS3) Transcription by Injurious Stimuli in the Liver Is Commonly Mediated by

834  BRG1. Front Cell Dev Biol 7:1–13.

835 95. Wang Z-J, Martin JA, Mueller LE, Caccamise A, Werner CT, Neve RL, Gancarz AM, Li

836  J-X, Dietz DM. 2016. BRG1 in the Nucleus Accumbens Regulates Cocaine-Seeking

837  Behavior-Supplementary info 80:652–660.

838   96.   Ganguly D, Sims M, Cai C, Fan M, Pfeffer LM. 2018. Chromatin Remodeling Factor

839   BRG1 Regulates Stemness and Chemosensitivity of Glioma Initiating Cells. Stem Cells

840   36:1804–1815.

841   97.   Güneş C, Paszkowski-Rogacz M, Rahmig S, Khattak S, Camgöz A, Wermke M, Dahl A,

842   Bornhäuser M, Waskow C, Buchholz F. 2019. Comparative RNAi Screens in Isogenic

843   Human Stem Cells Reveal SMARCA4 as a Differential Regulator. Stem cell reports

844   12:1084–1098.

845   98.   Kadoch C, Crabtree GR. 2015. Mammalian SWI/SNF chromatin remodeling complexes

846   and cancer: Mechanistic insights gained from human genomics. Sci Adv 1:1–17.

847   99.   Pulice JL, Kadoch C. 2016. Composition and function of mammalian SWI/SNF chromatin

848   remodeling complexes in human disease. Cold Spring Harb Symp Quant Biol 81:53–60.

849   100.   Hodges C, Kirkland JG, Crabtree GR. 2016. The many roles of BAF (mSWI/SNF) and

850   PBAF complexes in cancer. Cold Spring Harb Perspect Med 6:1–25.

851   101.   Michel BC, D'Avino AR, Cassel SH, Mashtalir N, McKenzie ZM, McBride MJ, Valencia

852   AM, Zhou Q, Bocker M, Soares LMM, Pan J, Remillard DI, Lareau CA, Zullow HJ,

853   Fortoul N, Gray NS, Bradner JE, Chan HM, Kadoch C. 2018. A non-canonical SWI/SNF

854   complex is a synthetic lethal target in cancers driven by BAF complex perturbation. Nat

855   Cell Biol 20:1410–1420.

856   102.   Tang L, Nogales E, Ciferri C. 2010. Structure and function of SWI/SNF chromatin

857   remodeling complexes and mechanistic implications for transcription. Prog Biophys Mol

858   Biol 102:122–128.

859   103.   Brownlee PM, Chambers AL, Cloney R, Bianchi A, Downs JA. 2014. BAF180 Promotes

860   Cohesion and Prevents Genome Instability and Aneuploidy. Cell Rep 6:973–981.

861   104.   Kakarougkas A, Ismail A, Chambers AL, Riballo E, Herbert AD, Künzel J, Löbrich M,

862   Jeggo PA, Downs JA. 2014. Requirement for PBAF in Transcriptional Repression and

863   Repair at DNA Breaks in Actively Transcribed Regions of Chromatin. Mol Cell 55:723–

864   732.

865   105.   Wang Z, Zhai W, Richardson JA, Olson EN, Meneses JJ, Firpo MT, Kang C, Skarnes

866   WC, Tjian R. 2004. Polybromo protein BAF180 functions in mammalian cardiac chamber

867   maturation. Genes Dev 18:3106–3116.

868   106.   Huang X, Gao X, Diaz-Trelles R, Ruiz-Lozano P, Wang Z. 2008. Coronary development

869   is regulated by ATP-dependent SWI/SNF chromatin remodeling component BAF180. Dev

870   Biol 319:258–266.

871   107.   Lee H, Dai F, Zhuang L, Xiao ZD, Kim J, Zhang Y, Ma L, You MJ, Wang Z, Gan B.

872   2016. BAF180 regulates cellular senescence and hematopoietic stem cell homeostasis

873   through p21. Oncotarget 7:19134–19146.

874   108.   Chowdhury B, Porter EG, Stewart JC, Ferreira CR, Schipma MJ, Dykhuizen EC. 2016.

875   PBRM1 regulates the expression of genes involved in metabolism and cell adhesion in

876   renal clear cell carcinoma. PLoS One 11:1–20.

877   109.   Huang L, Peng Y, Zhong G, Xie W, Dong W, Wang B, Chen X, Gu P, He W, Wu S, Lin

878   T, Huang J. 2015. PBRM1 suppresses bladder cancer by cyclin B1 induced cell cycle

879   arrest. Oncotarget 6:16366–16378.

880   110.   Xia W, Nagase S, Montia AG, Kalachikov SM, Keniry M, Su T, Memeo L, Hibshoosh H,

881   Parsons R. 2008. BAF180 is a critical regulator of p21 induction and a tumor suppressor

882   mutated in breast cancer. Cancer Res 68:1667–1674.

883   111.   Gao W, Li W, Xiao T, Liu XS, Kaelin WG. 2017. Inactivation of the PBRM1 tumor

884   suppressor gene amplifies the HIF-response in VHL-/- clear cell renal carcinoma. Proc

885   Natl Acad Sci U S A 114:1027–1032.

886 112. Albini S, Coutinho P, Malecova B, Giordani L, Savchenko A, Forcales SV, Puri PL. 2013.

887   Epigenetic Reprogramming of Human Embryonic Stem Cells into Skeletal Muscle Cells

888   and Generation of Contractile Myospheres. Cell Rep 3:661–670.

889 113. Padilla-Benavides T, Nasipak BT, Imbalzano AN. 2015. Brg1 Controls the Expression of

890   Pax7 to Promote Viability and Proliferation of Mouse Primary Myoblasts. J Cell Physiol

891   230:2990–2997.

892 114. Nasipak BT, Padilla-Benavides T, Green KM, Leszyk JD, Mao W, Konda S, Sif S,

893   Shaffer S a., Ohkawa Y, Imbalzano AN. 2015. Opposing calcium-dependent signalling

894   pathways control skeletal muscle differentiation by regulating a chromatin remodelling

895   enzyme. Nat Commun 6:1–12.

896 115. Witwicka H, Nogami J, Syed SA, Maehara K, Padilla-Benavides T, Ohkawa Y,

897   Imbalzano AN. 2019. Calcineurin Broadly Regulates the Initiation of Skeletal Muscle-

898   Specific Gene Expression by Binding Target Promoters and Facilitating the Interaction of

899   the SWI/SNF Chromatin Remodeling Enzyme. Mol Cell Biol 39:1–21.

900 116. Ochi H, Hans S, Westerfield M. 2008. Smarcd3 regulates the timing of zebrafish

901   myogenesis onset. J Biol Chem 283:3529–3536.

902 117. Lickert H, Takeuchi JK, Von Both I, Walls JR, McAuliffe F, Adamson SL, Henkelman

903   RM, Wrana JL, Rossant J, Bruneau BG. 2004. Baf60c is essential for function of BAF

904   chromatin remodelling complexes in heart development. Nature 432:107–112.

905 118. Harada A, Mallappa C, Okada S, Butler JT, Baker SP, Lawrence JB, Ohkawa Y,

906   Imbalzano AN. 2015. Spatial re-organization of myogenic regulatory sequences

temporally controls gene expression. Nucleic Acids Res 43:2008–2021.

119. Padilla-Benavides T, Nasipak BT, Paskavitz AL, Haokip DT, Schnabl JM, Nickerson JA,

Imbalzano AN. 2017. Casein kinase 2-mediated phosphorylation of Brahma-related gene

1 controls myoblast proliferation and contributes to SWI/SNF complex composition. J

Biol Chem 292:18592–18607.

120. Padilla-Benavides T, Haokip DT, Yoon Y, Reyes-Gutierrez P, Rivera-Pérez JA,

Imbalzano AN. 2020. CK2-Dependent Phosphorylation of the Brg1 Chromatin

Remodeling Enzyme Occurs during Mitosis. Int J Mol Sci 21:1–21.

121. Giacinti C, Bagella L, Puri PL, Giordano A, Simone C. 2006. MyoD recruits the

cdk9/cyclin T2 complex on myogenic-genes regulatory regions. J Cell Physiol 206:807–

813.

122. Joliot V, Ait-Mohamed O, Battisti V, Pontis J, Philipot O, Robin P, Ito H, Ait-Si-Ali S.

2014. The SWI/SNF Subunit/Tumor Suppressor BAF47/INI1 Is Essential in Cell Cycle

Arrest upon Skeletal Muscle Terminal Differentiation. PLoS One 9:1–11.

123. Serra C, Palacios D, Mozzetta C, Forcales S V., Morantte I, Ripani M, Jones DR, Du K,

Jhala US, Simone C, Puri PL. 2007. Functional Interdependence at the Chromatin Level

between the MKK6/p38 and IGF1/PI3K/AKT Pathways during Muscle Differentiation.

Mol Cell 28:200–213.

124. Ohkawa Y, Imbalzano AN. 2006. Skeletal muscle specification by myogenin. EMBO J

25:490–501.

125. Mallappa C, Nasipak BT, Etheridge L, Androphy EJ, Jones SN, Sagerstro CG, Ohkawa Y,

Imbalzano AN. 2010. Myogenic MicroRNA Expression Requires ATP-Dependent

Chromatin Remodeling Enzyme Function. Mol Cell Biol 30:3176–3186.

126.   de la Serna IL, Carlson KA, Hill DA, Guidi CJ, Stephenson RO, Sif S, Kingston RE,

Imbalzano AN. 2000. Mammalian SWI-SNF Complexes Contribute to Activation of the

hsp70 Gene. Mol Cell Biol 20:2839–2851.

127.   Abràmoff MD, Magalhaes PJ, Ram SJ. 2004. Image Processing with ImageJ.

Biophotonics Int 11:36–42.

128.   Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time

quantitative PCR and the 2-ΔΔCT method. Methods 25:402–408.

129.   Huang J, Liang X, Xuan Y, Geng C, Li Y, Lu H, Qu S, Mei X, Chen H, Yu T, Sun N, Rao

J, Wang J, Zhang W, Chen Y, Liao S, Jiang H, Liu X, Yang Z, Mu F, Gao S. 2017. A

reference human genome dataset of the BGISEQ-500 sequencer. Gigascience 6:1–9.

130.   Kim D, Langmead B, Salzberg SL. 2015. HISAT: A fast spliced aligner with low memory

requirements. Nat Methods 12:357–360.

131.   Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015.

StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat

Biotechnol 33:290–295.

132.   Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, Van Baren MJ, Salzberg SL,

Wold BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq reveals

unannotated transcripts and isoform switching during cell differentiation. Nat Biotechnol

28:511–515.

133.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods

9:357–359.

134.   Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome Bo. BMC Bioinformatics 12:1–16.

135.   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion

954   for RNA-seq data with DESeq2. Genome Biol 15:1–21.

955   136.   Audic S, Claverie JM. 1997. The significance of digital gene expression profiles. Genome

956   Res 7:986–995.

     Figure Legends:

     Fig. 1 (a) Confocal (top, scale 5μm) and bright field (bottom, scale 20μm) images for C2C12

     myoblasts treated with DMSO or PFI-3 and stained for myosin heavy chain (green) and with

     DAPI (blue) at the indicated timepoints. (b) Quantification of fusion index. (c) Differentiated

     myoblasts at respective timepoints were analyzed for number of nuclei per myotube. *p<0.05,

     **p<0.01 and ***p<0.005

     Fig. 2 mRNA expression levels at the indicated timepoints for (a) the fusion regulator genes

     myomaker and myomixer and (b) the myogenic genes myogenin, muscle creatine kinase, myosin

     light chain 1, Caveolin 3 and Integrin 7α in C2C12 myoblasts treated with DMSO or PFI-3.

     Expression was normalized to a control gene (EEF1A1). 100% expression is defined as the

     timepoint at which maximal expression was observed. ns, not significant, *p<0.05, **p<0.01 and

     ***p<0.005. (c) Representative western blot for MHC expression at the indicated times in

     C2C12 cells treated with DMSO or PFI-3. The indicated numbers are the pixel counts

     normalized to β-tubulin expression calculated using ImageJ.

     Fig. 3 (a) Heat maps showing results from RNA-seq analysis of PFI-3 treated C2C12 cells

     assayed while in the proliferative stage in growth media (GM) and while in differentiation media

     (DM) for 24h and 48h. (b) Venn diagram showing the number of genes affected at different

     timepoints. There were 572 differentially expressed genes (DEGs) in GM (blue), 1319 DEGs in

     DM 24h (peach) and 1681 DEGs in DM 48h (green). (c) GO analysis of downregulated genes at

     48h post induction of differentiation shows downregulation of muscle related genes. A HOMER

motif search shows enrichment of motifs corresponding to muscle-specific transcription factor

families. (d) GO analysis of upregulated genes at 48h post induction of differentiation shows

upregulation of cell cycle related processes. A HOMER motif search shows enrichment of

transcription factor motifs associated with cell cycle regulators.

Fig. 4 (a) C2C12 cells treated with PFI-3 show continued BrdU incorporation after 36h and 48 h

post-induction of differentiation as compared to control cells, scale 5 μm. (b) Quantification of

confocal images for BrdU incorporation assay in DMSO or PFI-3 treated C2C12 cells at the

indicated timepoints. (c) mRNA expression levels of cyclin A2, cyclin B1, cyclin D1, and cyclin

D2 in C2C12 myoblasts treated with DMSO or PFI-3 for indicated timepoints. Expression was

normalized to a control gene (EEF1A1). 100% expression is defined as the timepoint at which

maximal expression was observed. ns, not significant, $*p<0.05$, $**p<0.01$ and $***p<0.005$

Fig. 5 (a) Western blot analysis showed siRNA-mediated silencing of endogenous BAF180 in

proliferating C2C12 cells. A scramble siRNA (siScr) was used as a control. The indicated

numbers are the pixel counts normalized to Laminβ expression calculated using ImageJ. (b)

Representative images of myosin heavy chain staining in 48h and 96h differentiated cells

transfected with the control or BAF180-targeting siRNAs. The cells were fixed and analyzed by

immunofluorescence using an anti-myosin heavy chain mAb MF20 (*green*). The nuclei were

visualized by DAPI staining (*blue*). Scale bar, 20 μm.

1002

Fig. 6 Comparative analysis of RNA-seq datasets from BRG1 knockdown performed by Zhu et

al. (NAR, 2020) and from PFI-3 treatment. Venn diagrams represent DEGs in corresponding

datasets. The overlap represents genes common to both datasets. GO and HOMER motif

enrichment analyses show biological process categories and motifs identified within 1kb

upstream of the TSS in the promoters of (a) common downregulated genes and (b) common

upregulated genes.

1009

Fig. 7 ChIP assays show decreased occupancy of BRG1 and BRM on regulatory regions of

target genes upon PFI-3 inhibition. (a) Bar plots for BRG1 occupancy on the myogenin promoter

(Myog P), myosin heavy chain promoter (MHCIIb P), creatine kinase promoter (Ckm P) and

creatine kinase enhancer (Ckm E) are shown. (b) Bar plots for BRG1 and BRM occupancies on

the cyclin D1 and cyclin D2 promoters. The values have been normalized to an IgG experimental

control. These values are also normalized for binding at a non-specific region. ns, not significant,

*p<0.05, **p<0.01 and ***p<0.005

1017

Fig. 8 Graphical summary showing the effects of PFI-3 induced bromodomain inhibition on

skeletal myogenesis. In normal conditions, BRG1/BRM with active bromodomains can bind to

promoters of target genes when muscle differentiation is induced. This in turn affects two

important aspects of skeletal myogenesis: cell cycle exit and the formation of differentiated

multinucleated myotubes. In the presence of PFI-3, BRG1 and BRM show reduced binding to

target gene promoters leading to continued cell-cycle and incomplete differentiation resulting in

shorter myotubes.

Fig. 1 (a) Confocal (top, scale 5μm) and bright field (bottom, scale 20μm) images for C2C12 myoblasts treated with DMSO or PFI-3 and stained for myosin heavy chain (green) and with DAPI (blue) at the indicated timepoints. (b) Quantification of fusion index. (c) Differentiated myoblasts at respective timepoints were analyzed for number of nuclei per myotube. *p<0.05, **p<0.01 and ***p<0.005

Fig. 2 mRNA expression levels at the indicated timepoints for (a) the fusion regulator genes myomaker and myomixer and (b) the myogenic genes myogenin, muscle creatine kinase, myosin light chain 1, Caveolin 3 and Integrin 7α in C2C12 myoblasts treated with DMSO or PFI-3. Expression was normalized to a control gene (EEF1A1). 100% expression is defined as the timepoint at which maximal expression was observed. ns, not significant, *p<0.05, **p<0.01 and ***p<0.005. (c) Representative western blot for MHC expression at the indicated times in C2C12 cells treated with DMSO or PFI-3. The indicated numbers are the pixel counts normalized to β-tubulin expression calculated using ImageJ.

Fig. 3 (a) Heat maps showing results from RNA-seq analysis of PFI-3 treated C2C12 cells assayed while in the proliferative stage in growth media (GM) and while in differentiation media (DM) for 24h and 48h. (b) Venn diagram showing the number of genes affected at different timepoints. There were 3144 differentially expressed genes (DEGs) in GM (blue), 4674 DEGs in DM 24h (peach) and 5261 DEGs in DM 48h (green). (c) GO analysis of downregulated genes at 48h post induction of differentiation shows downregulation of muscle related genes. A HOMER motif search shows enrichment of motifs corresponding to muscle-specific transcription factor families. (d) GO analysis of upregulated genes at 48h post induction of differentiation shows upregulation of cell cycle related processes. A HOMER motif search shows enrichment of transcription factor motifs associated with cell cycle regulators.

Fig. 4 (a) C2C12 cells treated with PFI-3 show continued BrdU incorporation after 36h and 48h post-induction of differentiation as compared to control cells, scale 5 μm. (b) Quantification of confocal images for BrdU incorporation assay in DMSO or PFI-3 treated C2C12 cells at the indicated timepoints. (c) mRNA expression levels of cyclin A2, cyclin B1, cyclin D1, and cyclin D2 in C2C12 myoblasts treated with DMSO or PFI-3 for indicated timepoints. Expression was normalized to a control gene (EEF1A1). 100% expression is defined as the timepoint at which maximal expression was observed. ns, not significant, *p<0.05, **p<0.01 and ***p<0.005

a

siRNA

WT | scr | BAF180

200kDa -   BAF180

0.88   1.19   0.15

70kDa -   Laminβ

b

48h DM   96h DM

siScr

siBAF180

Fig. 5 (a) Western blot analysis showed siRNA-mediated silencing of endogenous BAF180 in proliferating C2C12 cells. A scramble siRNA (siScr) was used as a control. The indicated numbers are the pixel counts normalized to Laminβ expression calculated using ImageJ. (b) Representative images of myosin heavy chain staining in 48h and 96h differentiated cells transfected with the control or BAF180-targeting siRNAs. The cells were fixed and analyzed by immunofluorescence using an anti-myosin heavy chain mAb MF20 (green). The nuclei were visualized by DAPI staining (blue). Scale bar, 20 µm.

Fig. 6 Comparative analysis of RNA-seq datasets from BRG1 knockdown performed by Zhu et al. (NAR, 2020) and from PFI-3 treatment. Venn diagrams represent DEGs in corresponding datasets. The overlap represents genes common to both datasets. GO and HOMER motif enrichment analyses show biological process categories and motifs identified within 1kb upstream of the TSS in the promoters of (a) common downregulated genes and (b) common upregulated genes.

Fig. 7 ChIP assays show decreased occupancy of BRG1 and BRM on regulatory regions of target genes upon PFI-3 inhibition. (a) Bar plots for BRG1 occupancy on the myogenin promoter (Myog P), myosin heavy chain promoter (MHCIIb P), creatine kinase promoter (Ckm P) and creatine kinase enhancer (Ckm E) are shown. (b) Bar plots for BRG1 and BRM occupancies on the cyclin D1 and cyclin D2 promoters. The values have been normalized to an IgG experimental control. These values are also normalized for binding at a non-specific region. ns, not significant, *p<0.05, **p<0.01 and ***p<0.005

Graphical Summary:

Fig. 8 Graphical summary showing the effects of PFI-3 induced bromodomain inhibition on skeletal myogenesis. In normal conditions, BRG1/BRM with active bromodomains can bind to promoters of target genes when muscle differentiation is induced. This in turn affects two important aspects of skeletal myogenesis: cell cycle exit and the formation of differentiated multinucleated myotubes. In the presence of PFI-3, BRG1 and BRM show reduced binding to target gene promoters leading to continued cell-cycle and incomplete differentiation resulting in shorter myotubes.