Title: Induction of selective cell death in HIV-1-infected cells by DDX3

inhibitors leads to depletion of the inducible reservoir

Authors: Shringar Rao[1], Cynthia Lungu[2], Thijs H. Steijaert[1], Raquel Crespo[1], Robert-Jan

Palstra[1], Henrieke A. B. Prins[3], Wilfred van Ijcken[4], Casper Rokx[3], Yvonne M. Mueller[5], Jeroen

J. A. van Kampen[2], Annelies Verbon[3], Peter D. Katsikis[5], Charles A. B. Boucher[2], Rob A.

Gruters[2], Tokameh Mahmoudi[1]*

Affiliations:

[1]Department of Biochemistry, Erasmus University Medical Center, PO Box 2040, 3000 CA

Rotterdam, The Netherlands.

[2]Department of Viroscience, Erasmus Medical Centre, PO Box 2040, 3000 CA Rotterdam, The

Netherlands.

[3]Department of Internal Medicine, section of Infectious Diseases, Erasmus University Medical

Center, Rg-530, PO Box 2040 3000CA Rotterdam, The Netherlands.

[4]Erasmus Center for Biomics, Erasmus University Medical Center, Ee-671, PO Box 2040,

3000CA Rotterdam, Netherlands.

[5]Department of Immunology, Erasmus University Medical Center, PO Box 2040, 3000 CA

Rotterdam, The Netherlands.

*To whom correspondence should be addressed: t.mahmoudi@erasmusmc.nl.

ABSTRACT

An innovative approach to eliminate HIV-1-infected cells emerging out of latency, the major hurdle to HIV-1 cure, is to pharmacologically reactivate viral expression and concomitantly trigger intracellular pro-apoptotic pathways in order to selectively induce cell death (ICD) of infected cells, without reliance on the extracellular immune system. In this work we demonstrate the effect of DEAD-box polypeptide 3, X-Linked (DDX3) inhibitors on selectively inducing cell death in latent HIV-1-infected cell lines, primary CD4+ T cells and in CD4+ T cells from cART-suppressed people living with HIV-1 (PLWHIV). We used single-cell FISH-Flow technology to characterise the contribution of viral RNA to inducing cell death; pharmacological targeting of DDX3 reversed HIV-1 latency and selectively induced apoptosis in viral RNA-expressing CD4+ T cells from PLWHIV but not bystander cells. DDX3 inhibitor treatment of CD4+ T cells from PLWHIV in an *in vitro* culture model over five days resulted in an approximately 30% reduction of the inducible latent HIV-1 reservoir as determined by quantitation of CA HIV-1 RNA, by TILDA, as well as by FISH-Flow technology. RNA sequencing analysis revealed that while overall gene expression was minimally dysregulated, treatment of independent donor CD4+ T cells with DDX3 inhibitors led to significant downregulation of BIRC5 and HSPB1A, genes critical to cell survival during HIV-1 infection, providing mechanism for the observed selective cell death. Our data support the translation of DDX3 inhibitor class compounds into HIV-1 curative strategies and provide proof of concept for pharmacological reversal of latency coupled to induction of apoptosis towards elimination of the inducible reservoir.

INTRODUCTION

The main obstacle towards a Human Immunodeficiency Virus Type-1 (HIV-1) cure is the presence of a latent viral reservoir that does not actively produce viral particles, but retains the ability to do so and is not eliminated by combination antiretroviral therapy (cART) [1,2]. One strategy to eliminate the latent HIV-1 reservoir is the 'Shock and Kill' approach that involves the use of 'latency reversing agents' (LRAs) to reactivate viral gene expression (Shock), followed by elimination of the cells harbouring reactivated provirus (Kill) [3]. The 'Kill' aspect of this approach has largely depended on an effective host adaptive immune response. Although clinical studies with LRAs demonstrated induced reactivation *in vivo*, limited to no reduction in the size of the viral reservoir or time to viral rebound post cessation of cART was observed, indicating that elimination of reactivating cells warrants more attention [4-6]. So far, broadly neutralising antibodies, therapeutic vaccines, chimeric antigen receptors, checkpoint inhibitors and engineered bispecific antibodies have been investigated to eliminate reactivated HIV-1-infected cells (reviewed in [7]). However, dependency on cell-mediated and humoral immune responses that are already compromised in HIV-infected individuals because of impaired CD4+ T cell help and exhausted immune compartments poses a considerable obstacle to these strategies. Indeed, early results from combinatorial clinical trials which combine LRAs and immune-enhancing clearance strategies demonstrated that although HIV-1-specific immune responses were induced, no decrease in the size of viral reservoir or viral control in the absence of cART were observed ([8-10], reviewed in [11]).

A parallel approach to eliminate the viral reservoir independently of the extracellular immune system is to pharmacologically trigger pathways that intrinsically induce cell death (ICD) in reactivating HIV-1-infected reservoir cells. For example, using compounds to activate innate

intracellular signalling pathways could trigger apoptosis. Several HIV-1 proteins have been

demonstrated to have both anti-apoptotic activity in the early stages of the viral replication cycle

and pro-apoptotic activity in the later replication stages [12]. By inhibiting viral anti-apoptotic

activity or by promoting pro-apoptotic functions, HIV-1-infected cells can be selectively

eliminated, leading to a reduction in the size of the viral reservoir. Multiple pro-apoptotic

compounds that have been developed in context of cancer research could potentially be

repurposed for viral reservoir elimination and some have been investigated for their ability to

induce cell death of HIV-1-infected cells (reviewed in [13,14]). Interestingly, an important

characteristic of the latent viral reservoir that has recently come to light is that while it is

transcriptionally competent, it does not necessarily generate viral proteins due to post-

transcriptional blocks [15,16]. Induced viral RNA does however retain the ability to trigger innate

antiviral signalling pathways [17,18] and possibly induce cell death in reactivating cells. The

potential contribution of inducing viral transcription as a strategy to induce death of the infected

cells has been thus far under-investigated [19].

Advanced single-cell fluorescence in situ hybridisation Flow cytometry (FISH-Flow) [15,16,20,21]

provides a powerful tool to examine the specific contribution of HIV-1 genomic RNA (vRNA) in

inducing cell death. Unlike previous single cell HIV-1 flow cytometry-based studies that could

detect only viral protein production and the translationally competent reservoir, FISH-Flow

allows for the delineation of the contribution of the previously under characterised

transcriptionally competent cells that produce vRNA, can trigger innate immune responses and

may or may not express viral proteins [19,22]. The single cell analysis thus provides an additional

layer of insight when studying the contribution of vRNA to apoptosis.

DEAD-box polypeptide 3, X-Linked, herein referred to as DDX3, is a host protein belonging to

the DEAD-box (Asp-Glu-Ala-Asp) family of ATP-dependent RNA helicases. It is involved in all

aspects of RNA metabolism including processing, transport and translation, as well as in cell

cycle progression, stress response, innate immune sensing and apoptosis [23-25]. DDX3 has been

implicated in several cancers due to its oncogenic function, but also plays a dual role in cancer

progression because of its tumour suppressive activity [26-30]. Moreover, DDX3 plays distinct roles

in HIV-1 infection by enhancing viral RNA gene expression, but also in activating components

of the innate antiviral signalling pathway [31]. During HIV-1 replication, DDX3 has an integral

function in the nucleocytoplasmic export of the genomic unspliced viral RNA (vRNA) [32,33] as

well as in vRNA translation in the cytoplasm [34-36]. In dendritic cells, DDX3 senses abortive HIV-

1 RNA transcripts to induce type-I interferon immune responses via mitochondrial antiviral

signalling protein (MAVS) [37]. Because of its widely described roles in viral replication as well as

in tumorigenesis, DDX3 has thus emerged as an attractive target for both anti-cancer and

antiviral drugs [27,38].

Due to DDX3's roles in both HIV-1 RNA metabolism as well as in the regulation of apoptosis,

we hypothesized that treatment of HIV-1-infected cells with small molecule inhibitors of DDX3

would inhibit vRNA export and translation; while the residual vRNA would activate innate

antiviral immune signalling pathways and trigger apoptosis of vRNA-expressing cells, thereby

resulting in a depletion of the viral reservoir.

Here we evaluated the effects of DDX3-inhibition on the latent HIV-1 reservoir using two

clinically advanced inhibitors of DDX3, RK-33 and FH1321, which target the ATPase and

RNA-binding domains of DDX3 respectively. Pharmacological inhibition of DDX3 reversed

HIV latency in the latent HIV-1-infected J-Lat 11.1 cell line, in ex vivo latently infected CD4+ T

cells and in CD4+ T cells obtained from cART-suppressed HIV-1-infected donors via NF-kB

activation. DDX3 inhibitor-mediated HIV-1 latency reversal was observed predominantly at the

RNA level, which is consistent with the role of DDX3 as an inhibitor of vRNA export and

translation. RNA sequencing analysis of primary CD4+ T cells obtained from independent

donors revealed significant downregulation of the inhibitor of apoptosis protein BIRC5 and Heat

Shock protein 70 (HSP70) HSPA1B upon DDX3 inhibition. Importantly, treatment of latent

HIV-1-infected cells with DDX3 inhibitors resulted in selective induction of apoptosis in vRNA-

expressing cells but not in the uninfected/bystander cells. We quantitated the impact of treatment

with DDX3 inhibitors on the inducible latent HIV-1 reservoir in CD4+ T cells obtained from

cART-suppressed people living with HIV-1 (PLWHIV) in an *in vitro* culture model using three

different methods of reservoir quantitation. Strikingly, treatment with DDX3 inhibitors over five

days resulted in an approximately 30% reduction of the inducible reservoir as determined by

quantitation of cell associated (CA) HIV-1 RNA, by tat/rev induced limiting dilution assay, as

well as at the single cell level using FISH-Flow technology. These results demonstrate that

pharmacological inhibition of DDX3 induces latency reversal and selective apoptosis of latent

HIV-1 infected cells and serve as proof-of-concept that ICD inducers can decrease the inducible

HIV-1 reservoir in infected patient cells *ex vivo*.

RESULTS

DDX3 inhibition induces reactivation and cell death in J-Lat 11.1 cells

Our initial screening studies focussed on examining whether pharmacological inhibition of DDX3 can affect proviral reactivation in a cell line model of HIV-1 latency, J-Lat 11.1. This model is a Jurkat-derived cell line that contains a latent integrated copy of the HIV-1 provirus defective in *env* with a GFP reporter in place of the *nef* open reading frame [39]. J-Lat 11.1 cells harboured a basal frequency of GFP positive cells at <7.5 % that was reactivated by treatment with the PKC agonist phorbol myristate acetate (PMA) to >80% (Supplementary Fig. 1a). J-Lat 11.1 cells were treated with increasing concentrations (0.5 – 2 µM) of the DDX3 inhibitor RK-33 that has been developed for cancer treatment. RK-33 binds to the ATPase domain of DDX3 and is reported to improve radiation sensitisation of tumours in models of lung cancer, colorectal cancer, prostate cancer and Ewing's carcinoma [27,28,40-44]. Interestingly, treatment with RK-33 alone induced viral reactivation in a dose dependent manner, with approximately 18% of cells producing GFP when treated with 2µM RK-33 (Fig. 1a), thus unravelling a novel role for DDX3 inhibitors as LRAs. A larger latency reversal activity of RK-33 was observed at the RNA level with a 70 fold and 25 fold increase in the relative expression of the HIV-1-promoter-controlled GFP and HIV-1 genomic vRNA (*Pol*) transcripts respectively as quantified by RT-qPCR (Fig. 1b and c). Since DDX3 has been reported to play important roles in the nucleocytoplasmic export and translation of HIV-1 vRNA [35, 36, 38], we hypothesized that RK-33 could also influence HIV-1 gene expression at a post-transcriptional level by impairing either vRNA export or translation. To investigate this, we conducted FISH-Flow after treatment of J-Lat 11.1 cells with 2 µM RK-33 for 18 hours. Using probes that amplify and allow identification and quantification of viral RNA expression in cells, FISH-Flow allows for the simultaneous detection

of distinct populations: i) latent cells, which express neither viral RNA nor viral protein (vRNA-

/GFP-), ii) efficiently reactivated cells, which express both HIV-1 viral RNA and proteins

(vRNA+/GFP+), iii) transcriptionally competent viral RNA expressing cells with either

untranslated vRNA or a block in post-transcriptional vRNA processing and viral protein

expression (vRNA+/GFP-), and iv) cells only expressing GFP from multiply spliced viral RNA

transcripts (vRNA-/GFP+) (Fig. 1d). On average, under uninduced conditions, only 3% of cells

constitutively produced vRNA, whereas vRNA was produced in more than 80 % of cells upon

PMA stimulation (Fig. 1d and e). Treatment with 2 µM RK-33 resulted in expression of vRNA

in approximately 15% of J-Lat cells (Fig. 1d and e), corroborating its latency reversal activity.

Interestingly, 4% of cells were observed in the vRNA+/GFP- quadrant, corresponding to ~28%

of all vRNA+ cells (Fig. 1d and e). In contrast, only 0.02% of the vRNA-expressing PMA-

treated cells did not express GFP (Fig. 1d and e). This data indicates a post-transcriptional block

in vRNA metabolism upon DDX3 inhibition. In order to determine if this block was at the level

of nucleocytoplasmic export, we seeded the cells that underwent FISH-Flow analysis on a

coverslip and observed them by confocal microscopy. We observed that the vRNA+/GFP- cells

in RK-33-treated conditions contained vRNA sequestered in the nucleus, while in double

positive cells in both the PMA and RK-33-treated condition, vRNA was present in the cytoplasm

of cells (Supplementary Fig. 1b). Therefore, DDX3 inhibition by RK-33 resulted in impaired

nucleocytoplasmic export of the vRNA and reduced GFP expression, thereby highlighting the

importance of using vRNA expression and not viral protein production as a marker for HIV-1

infection when investigating the effects of DDX3 inhibitors.

DDX3 has previously been reported to have cell-type dependent differential effects on the

regulation of the NF-κB pathways, and has been implicated in binding to the p65 subunit thus

inhibiting NF-κB-mediated transcription [45,46]. In order to determine if RK-33 is capable of

activating NF-κB, Jurkat cells were nucleofected with firefly luciferase reporter plasmids either

containing or lacking NF-κB consensus sites upstream of a minimal promoter, along with an

internal Renilla control plasmid. Cells were then either mock-treated or treated with 2 μM RK-33

for 18 hours. Treatment with RK-33 resulted in a modest increase in relative luciferase

expression in the NF-κB reporter transfected cells as compared to the mock-treated cells

(Supplementary Fig. 1c). The NF-κB pathway was also activated in J-Lat 11.1 cells treated with

RK-33 as is evident by an upregulation of the NF-κB target genes IL-10 and MMP9 by 8 and 19-

fold respectively. (Fig. 1f and Supplementary Fig. 1d). Our data therefore suggests that the

DDX3 inhibitor RK-33 acts as a latency reversing agent that reactivates the HIV-1 provirus via

activation of NF-κB. Considering RK-33's previously described function in inducing apoptosis,

we also assessed whether the vRNA produced upon latency reversal in the RK-33-treated cells

were capable of activating innate antiviral signalling pathways and could further contribute to

inducing cell death [40]. J-Lat 11.1 cells and parental Jurkat cells that do not contain the integrated

HIV-1 provirus were treated with increasing concentrations of RK-33 (0.5 – 2 μM). Eighteen

hours post treatment, cells were stained with Hoescht 33342 and the percentage of viable cells

was quantified by flow cytometry. We observed a dose-dependent increase in cell death in RK-

33-treated J-Lat cells as compared to parental Jurkat cells, most strikingly at 2 μM RK-33 (Fig.

1g). In this condition, cell death was observed in 70 % of J-Lat 11.1 cells, but only in 30 % of

Jurkat cells, representing a significant increase in cell death in HIV-1-infected cells. Since they

are a cancer-derived cells line, the viability of Jurkat cells was also affected upon treatment with

the RK-33, albeit not as much as the J-Lat 11.1 cells. Using Flow cytometry activated cell

sorting we sorted and compared the reactivated GFP+ and GFP- J-Lat 11.1 cells that were treated

with 2 µM RK-33. As expected, the GFP+ J-Lat 11.1 sorted cells contain ~250 fold more vRNA

than the GFP- population since the latter contains the vRNA-/GFP- population (Fig. 1h).

Interestingly, concomitant with the observed increased viral RNA expression, we also observed

an approximately 35% decrease in the ratio of the anti-apoptotic BCl2 to the pro-apoptotic Bax

mRNA expression levels in the GFP+ population compared to the RK-33-treated GFP negative

population, indicating that the vRNA expressing cells have a higher susceptibility to undergo

apoptosis (Fig. 1i). We also observed an upward trend in the mRNA expression of the innate

antiviral signalling pathway components RIG-I and MAVS in the vRNA high fraction, consistent

with previous work demonstrating HIV-1 vRNA's ability to activate these pathways

(Supplementary Fig. 1e and f) [17,18]. Our data therefore indicates that in a preliminary screening

model of HIV-1 latency, DDX3 inhibition with RK-33 results in latency reversal and that the

RK-33-treated HIV-1 expressing cells harbour markers for apoptosis and are more susceptible to

cell death by RK-33 than the vRNA-/GFP- J-Lat 11.1 cells or uninfected parental Jurkat cells.

Fig. 1

Fig. 1. DDX3 inhibition reverses HIV-1 latency and induces cell death in J-Lat 11.1 cells

a J-Lat 11.1 cells were treated with increasing concentrations of the DDX3 inhibitor RK-33. %

of reactivation, monitored by GFP production, was quantified by Flow cytometry. Error bars

represent mean ± SD from six independent experiments with at least 10000 cells counted per

treatment (Paired, two-tailed *t*-test). The relative mRNA expression levels of **b** GFP and **c** Pol in

the RK-33-treated J-Lat cells were quantified by RT-qPCR. Error bars represent mean ± SD

from three independent experiments (Unpaired, two-tailed *t*-test). **d** Representative FISH-Flow

dot-plots from one independent experiment of J-Lat cells treated with DMSO (Mock), PMA or

2μM RK-33. **e** The % of vRNA expressing cells (GFP- or GFP+) as treated in (d) were

quantified. Error bars represent mean ± SD from three independent experiments with at least

10000 cells counted per treatment (Unpaired, two-tailed *t*-test). **f** Relative MMP9 mRNA

expression levels from cells treated as in (a) were quantified by RT-qPCR. Error bars represent

mean ± SD from independent experiments from 5 different donors (Unpaired, two-tailed *t*-test).

g % Cell death was measured in Jurkat vs. J-Lat cells treated with increasing concentrations of

RK-33 by flow cytometry. Error bars represent mean ± SD from five independent experiments

with at least 10000 cells counted per treatment (Paired, two-tailed *t*-test). J-Lat cells were sorted

into GFP- (vRNA low) and GFP+ (vRNA high) fractions and **h** the relative mRNA expression of

vRNA using primers against Pol and **i** the relative expression ratios of Bcl2 and Bax mRNA

expression were quantified by RT-qPCR. Error bars represent mean ± SD from three

independent experiments (Unpaired, two-tailed *t*-test).

DDX3 inhibition downregulates genes involved in cell survival

The inhibition of DDX3 helicase function, i.e., dsRNA unwinding, can be achieved by competitive inhibition of the ATP or RNA binding sites. RK-33 was demonstrated to inhibit RNA helicase activity in a concentration-dependent manner reducing ATP consumption in enzymatic assays [40]. A new class of inhibitors competitive for the DDX3 RNA binding site have been reported by Radi et. al. [47] and were further developed leading to identification of the potent and selective compound FH1321 (compound 16d in [38]), that specifically targets the DDX3 RNA binding site with no effect in its ATPase activity [38]. A hypothetical binding model obtained by molecular docking calculation for both RK-33 and FH1321 depicts the predicted binding of RK-33 to the ATP binding site of DDX3 while, FH1321 binds the RNA binding site (Fig. 2a, Supplementary Fig. 2a and b). For both molecules, the downstream effect is inhibition of the RNA helicase activity of DDX3 [40,47]. We therefore investigated the potential function and mechanism of action of RK-33 and FH1321 in HIV-1 reactivation and reservoir dynamics in the more relevant primary CD4+ T cells. First, we observed no significant decrease in viability when primary CD4+ T cells from three healthy donors were treated with up to 10 μM of either compound (Supplementary Fig. 2c). We also cultured CD4+ T cells in the presence or absence of 2 μM RK-33 or 1 μM FH-1321, for 5 days and observed no significant differences in cell viability between the DDX3 inhibitor-treated and mock-treated conditions (Supplementary Fig. 2d). To gain insight into the molecular mechanism by which DDX3 inhibition may induce cell death in HIV-1-infected cells, we treated primary CD4+ T cells obtained from two healthy donors with RK-33 and FH-1321 and performed RNA-sequencing. Minimal changes in gene expression were observed 4 hours post treatment with 2 μM RK-33 or 1 μM FH-1321 with 1 and 5 differentially expressed genes respectively with a 1.5-fold difference in expression and a false

discovery cut-off rate set to 0.05 (Fig. 2b and Supplementary Fig. 2e and f). At 18 hours post

treatment: primary CD4+ T cells treated with 1 µM FH-1321 displayed only 34 genes

differentially expressed while treatment with RK-33 (2 µM) resulted in 39 genes differentially

expressed in both donors (Fig. 2b-d and Tables 1 and 2). Interestingly, the majority of the

differentially expressed genes were common to both the RK-33 and FH-1321 treatments (n=23).

Since these effects are common between compounds belonging to two different classes of DDX3

inhibitors they are bound to be specific to inhibition of DDX3 helicase activity (Fig. 2e).

Notably, the overall change in gene expression was very limited, consistent with the minimal

potential for off target or cytotoxic effects associated with use of these compounds. Interestingly,

two of the differentially expressed genes significantly downregulated in CD4+ T cells in

response to RK-33, the inhibitor of apoptosis protein BIRC5 and HSPA1B (heat shock protein

70 (Hsp70)) have been previously implicated to play a role in HIV-1 induced cell death. BIRC5,

which is also significantly downregulated in response to FH-1321 treatment, has been previously

demonstrated to be upregulated in latent HIV-1-infected cells, and its inhibition resulted in

increased cell death of HIV-1-infected cells [48]. Also involved in regulating apoptosis in infected

cells, HSPA1B was shown to inhibit apoptosis induced by the HIV-1 protein Vpr [49]. Consistent

with the observed DDX3-mediated induction of NF-κB, HSPA1B was also shown to inhibit

HIV-1 reactivation via suppression of NF-κB signalling [50]. RK-33 and FH-1321-mediated

downregulation of BIRC5 and RK-33-mediated HSPA1B downregulation was confirmed by RT-

qPCR in six additional donor CD4+ T cells (Fig. 2f-g). To ensure that DDX3 inhibitors do not

induce T cell activation, we examined CD25 expression upon treatment with RK-33 and FH-

1321 (Supplementary Fig. 3a). Consistent with the minimal effects observed with the RNA

sequencing, treatment with DDX3 inhibitors did not result in induction of T cell activation, nor

did DDX3 inhibition compromise effector function in CD4+ T cells (Fig. 2h and i) or CD8+ T

cells (Supplementary fig. 3e and c) as determined by flow cytometry measuring IL-2 and IFNγ

upon PMA-induced activation. Moreover, treatment with DDX3 inhibitors did not affect

capacity for proliferation in either CD4+ T cells (Fig. 2j and Supplementary Fig. 3d) or CD8+ T

cells (Supplementary Fig. 3e and f) as determined after induction/stimulation with CD3/CD28.

We also confirmed that DDX3 inhibition at the concentrations used does not affect *de novo* host

cell translation in primary CD4+ T cells (Supplementary Fig. 3g and h). Finally, we determined

if treatment of CD4+ T cells with DDX3 inhibitors could induce NF-κB activation by

quantifying levels of target genes by RT-qPCR. Treatment with RK-33 and FH-1321 resulted in

induction of the targets IL-8 and MMP9 mRNA expression, and whereas FH-1321 treatment

induced expression of IL-8 and IL-10 mRNA expression (Supplementary Fig. 3i-k).

Fig. 2

b

Condition	Upregulated	Downregulated	No change
RK-33 - 4 hours post treatment	1	0	58301
FH1321 - 4 hours post treatment	5	0	58297
RK-33 - 18 hours post treatment	2	37	58263
FH1321 - 18 hours post treatment	2	32	58268

Fig. 2. Effects of DDX3 inhibitors RK-33 and FH-1321 on CD4+ T cells

a A. Molecular docking of RK-33 and FH-1321 to pre-unwound DDX3 confirmation as

predicted by AutoDock Vina. DDX3 domain 1 is shown as pale green cartoon, domain 2 is

shown as pale blue cartoon and surface color code is pink for ATP-binding residues and orange

for RNA-binding residues. Color code for RK-33 is yellow for carbon, red for oxygen, dark blue

for nitrogen. Color code for FH-1321 is green for carbon, red for oxygen, dark blue for nitrogen.

The predicted hydrogen bond between FH-1321 and Lysine 541 from DDX3 RNA-binding

pocket is indicated with a dashed line. **b** Table of differentially expressed genes upon treatment

of CD4+ T cells with DDX3 inhibitors. Volcano plot of differentially expressed genes 18 hours

post-treatment with **c** RK-33 and **d** FH-1321. **e** Venn diagram of overlapping differentially

expressed genes 18 hours post treatment with DDX3 inhibitors. CD4+ T cells from healthy

donors were treated with DMSO (Mock), RK-33, FH1321 or PMA-Ionomycin and the relative

expression of **f** BIRC5 and **g** HSPA1B mRNAs were quantified by RT-qPCR. Error bars

represent mean ± SD from independent experiments from 6 different donors (Unpaired, two-

tailed *t*-test). Percentage of IL-2 **h** and IFNγ **i** producing cells in unstimulated and stimulated

primary CD4+ lymphocytes after treatment with RK-33 and FH-1321 (n=3). **j** Percentage of

proliferated CD4+ T cells from 3 healthy donors upon treatment with DDX3 inhibitors.

DDX3 inhibitors reactivate and selectively kill HIV-1-infected cells in a primary model of

latency

Given our observations that DDX3 inhibition reverses latency, induces more cell death in the J-

lat model and doesn't affect primary CD4+ T cell function, we wondered if we could use the

DDX3 inhibitors to reactivate the provirus and induce cell death in a primary CD4+ T cell model

of HIV-1 latency, developed by Lassen and Greene [51-53] (Fig. 3a). In this model of HIV-1

latency, the virus used to infect the primary CD4+ T cells contained a luciferase reporter inserted

in the *nef* open reading frame, which is encoded by the multiply-spliced vRNA transcript (similar

to the GFP in the J-Lat cells). We treated latently-infected cells with 2 µM RK-33, 1 µM

FH1321 or PMA-ionomycin (as a positive control) and monitored latency reversal 18 hours post

treatment by FISH-Flow [22], RT-qPCR using primers against the Pol region of the vRNA and

relative luciferase expression (Fig. 3a). To quantify ICD, we also co-stained cells with a fixable

viability dye and either with a fluorophore-conjugated antibody that detects AnnexinV or cleaved

Caspase3/7, both markers for host cell apoptosis (Gating strategy in Supplementary Fig. 4a).

FISH-Flow has a high specificity for the target HIV-1 vRNA with negligible background in the

uninfected CD4+T cells and a sensitivity to detect up to 10 HIV-1 vRNA+ cells per million cells

[16,22] (Fig. 3b). In latent HIV-1 infected CD4+ T cells, basal levels of vRNA were produced in the

uninduced (Mock) condition, with a 4.8 (s.d. ±3.3) fold increase in the frequency of vRNA-

expressing cells upon PMA-ionomycin stimulation (Fig. 3b and c). Treatment with the DDX3

inhibitors RK-33 and FH1321 resulted in a 2.8 (s.d. ± 1.1) fold and a 2.5 (s.d. ± 1.4) fold

increase in the frequency of vRNA-expressing cells respectively as compared to the mock-

treated condition (Fig. 3b and c). We also measured intracellular vRNA expression from cell

lysates by RT-qPCR and observed a 12.4-fold increase with RK-33, a 15.7-fold increase with

FH1321 and a 66.8 fold increase with PMA-ionomycin as compared to the mock-treated cells

(Fig. 3d). When treated with DDX3-inhibitors, a 1.5-fold increase in relative luciferase units as

compared to the mock-treated condition was observed in both RK-33 and FH1321-treated

conditions (Fig. 3e). Together, these data demonstrate that the DDX3 inhibitors also induce

latency reversal in a primary CD4+T cell model of HIV-1 latency.

Based on our observations of increased cell death in J-Lat 11.1. cells in comparison to the Jurkat

cells and the lack of cell death in uninfected primary CD4 T-cells (Fig. 1g, Supplementary Fig.

2c and d), we sought to determine if increased cell apoptosis could be observed specifically in

the vRNA-expressing primary CD4+ T cells in the *in vitro* model of HIV-1 latency. For this we

quantified the percentage of AnnexinV positive cells in cultures treated with mock, RK-33,

FH1321 or PMA-Ionomycin. Because FISH-Flow is a single cell technique, we could distinctly

quantify the differential expression of the vRNA+ cells or the vRNA-/bystander cells from our

latency model. In order to validate the selectivity of cell death upon drug treatment, we also

quantified the percentage of AnnexinV expression in uninfected CD4+ T cells. Basal levels of

AnnexinV positivity (<1%) were observed in the uninfected cells or the vRNA- bystander cells

(Fig. 3f and g). Of the vRNA+ cells, 4.8 (s.d. ± 4.4) % were AnnexinV positive in the cultures

treated with PMA-ionomycin, indicating that viral reactivation might already make the cells

more susceptible to cell death (Fig. 3f). Interestingly, in the cultures treated with RK-33 and FH-

1321, 11.7 (s.d. ± 8.8) % and 10.3 (s.d. ± 4.6) % of vRNA-expressing cells respectively, were

AnnexinV positive (Fig. 3f). In order to further validate that the vRNA+ cells are indeed

undergoing apoptosis, we repeated these studies in three additional donors and measured the

expression of cleaved Caspase3/7 by FISH-Flow. We observed that 40.5 (s.d. ± 8.5) % and 38.9

(s.d. ± 15.5) % of RK-33 and FH-1321 treated cells were cleaved Caspase3/7 positive as

compared to the 13.3 (s.d. ± 5.9) % positive cells in the Mock-treated condition (Fig. 3g). We

were unable to observe vRNA+ mediated effects on the relative expression of the anti-apoptotic

protein Bcl2 and the pro-apoptotic protein Bax by RT-PCR (Supplementary Fig. 4b) since only a

small fraction of cells <1% is HIV-1 infected within the total population, further highlighting the

importance of FISH-Flow which enables investigation of HIV-1 induced selective cell death at

the single cell level. Together, this data shows that DDX3 inhibitors exacerbate apoptosis

triggered by the vRNA and selectively induce cell death in vRNA-expressing cells, while

uninfected cells are not affected.

Fig. 3. DDX3 inhibition has LRA and ICD activity in a primary model of HIV-1 latency

a Schematic for latency establishment in primary human CD4 + T cells using the Lassen and

Greene model. Latent HIV-1-infected cells were treated on Day 5 with DMSO (Unstimulated),

PMA/Ionomycin, and the DDX3 inhibitors RK-33 and FH-1321. Uninfected cells were used as a

negative control. **b** Representative FISH-Flow dot plots of cells as treated in (a) representing

50000 cells from each condition. **c** The % of vRNA expressing cells as depicted in (b) were

quantified and normalised to the unstimulated condition. Error bars represent mean ± SD from

independent experiments from 5 different donors, with a minimum of 100,000 events acquired

for each condition (Unpaired, two-tailed *t*-test). **d** The relative Pol mRNA expression levels from

cells as treated in (a) were quantified by RT-qPCR. Error bars represent mean ± SD from

independent experiments from 5 different donors in duplicate (Unpaired, two-tailed *t*-test). **e**

Relative luciferase activity of cells as treated in (a) was measured and normalised to the mock

treated condition. Error bars represent mean ± SD from independent experiments from 5

different donors in duplicate (Unpaired, two-tailed *t*-test). **f** The percentages of AnnexinV

positive cells from the uninfected, infected vRNA- fractions, or infected vRNA+ cells as

quantified by FISH-Flow. Error bars represent mean ± SD from independent experiments from 5

different donors, with a minimum of 100,000 events acquired for each condition (Paired, two-

tailed *t*-test). **g** The percentages of cleaved Caspase3 positive cells from the infected vRNA-

fractions or infected vRNA+ cells as quantified by FISH-Flow. Error bars represent mean ± SD

from independent experiments from at 3 different donors, with a minimum of 100,000 events

acquired for each condition (Paired, two-tailed *t*-test).

DDX3 inhibitors induce latency reversal in CD4+ T cells from HIV-1-infected donors

To further investigate the clinical potential of DDX3 inhibitors we examined their ability to

induce latency reversal and selective cell death in CD4+ T cells from chronic HIV-1-infected

donors on suppressive cART with undetectable viral loads for at least 12 months (n=3).

Following isolation from cryopreserved PBMCs obtained by leukapheresis of HIV-1-infected

donors, CD4+ T cells were treated according to the schematic in Fig. 4a. Briefly, cells were

cultured either with mock treatment, 2 µM RK-33 or 1 µM FH1321 overnight in the presence of

an HIV-1 integrase inhibitor. Cells were collected 18 hours post treatment and analysed by

FISH-Flow using probes against the vRNA, a viability stain, and antibodies directed against

AnnexinV to quantify viral reactivation and selective early apoptosis using a similar gating

strategy as described in Supplementary Fig. 4a. A nested RT-qPCR was also performed to

measure cell-associated vRNA with primers targeting the Gag region of the vRNA. A 4.7 (s.d. $\pm$

1.3) fold and a 3.1 (s.d. $\pm$ 0.5) fold increase in the percentage of vRNA-producing cells as

compared to the mock-treated cells was observed in the RK-33 and FH1321-treated conditions

respectively (Fig. 4b and c). Importantly, 36.4 (s.d. $\pm$ 14.3) % and 39.8 (s.d. $\pm$ 8.9) % of vRNA+

cells in the RK-33 and FH-1321 conditions were AnnexinV positive, while all the vRNA-

populations had <1% AnnexinV positivity (Fig. 4d). When LRA activity was measured by RT-

qPCR, a 6.7 (s.d. $\pm$ 5.2) fold and a 7.5 (s.d. $\pm$ 6.7) fold increase in the relative copies of Gag

vRNA in the RK-33 and FH1321 conditions (Fig. 4e, Supplementary Fig. 5a) was observed. We

also validated that both BIRC5 and HSPA1B mRNA expression was significantly downregulated

upon RK-33 treatment while FH-1321 treatment resulted in decreased expression of BIRC5

(Supplementary Fig. 5b and c). Therefore, treatment with DDX3 inhibitors also demonstrated

latency reversal activity and selective cell death in vRNA-containing CD4+ T cells from HIV-1-

infected donors.

Fig. 4

Fig. 4. DDX3 inhibition reverses latency in CD4+ T cells from HIV-1-infected donors and

selectively induces cell death

a Schematic of experiment to test LRA and ICD activity of DDX3 inhibitors in samples from

PLWHIV. CD4+ T cells isolated from HIV-1-infected donors were treated with DMSO

(Unstimulated), PMA/Ionomycin, and the DDX3 inhibitors RK-33 and FH-1321. For all graphs,

three different donors are represented by blue, red and green symbols consistently. **b**

Representative FISH-Flow dot plots from an HIV-1-infected donor as treated in (a) representing

100000 cells from each condition. **c** The % of vRNA expressing cells as depicted in (b) were

quantified and normalised to the unstimulated condition. Error bars represent mean ± SD from

independent experiments from 3 different donors, with a minimum of 100,000 events acquired

for each condition (Unpaired, two-tailed *t*-test). **d** The percentages of AnnexinV positive cells

from the vRNA- or the vRNA+ cells as quantified by FISH-Flow. Error bars represent mean ±

SD from independent experiments from 3 different donors, with a minimum of 100,000 events

acquired for each condition (Unpaired, two-tailed *t*-test). **e** The relative Gag mRNA expression

levels from cells as treated in (a) were quantified by nested RT-qPCR. Error bars represent mean

± SD from independent experiments from at 3 different donors in duplicate (Unpaired, two-tailed

t-test).

DDX3 inhibition results in a reduction in the size of the inducible viral reservoir

The induction we observed of the pro-apoptotic AnnexinV marker in HIV-1-expressing cells

upon DDX3 inhibitor treatment was consistent with the notion that DDX3 treatment could result

in depletion of the inducible HIV reservoir. We therefore determined whether the DDX3

inhibitors-treated reactivated and pro-apoptotic cells were actually eliminated over time,

resulting in a decrease in the size of the inducible viral reservoir *in vitro*. CD4+ T cells isolated

from PLWHIV (n=3) were cultured in the presence of the DDX3 inhibitors for five days and

then treated with PMA-Ionomycin to reactivate them to maximal capacity (Fig. 5A). The size of

the inducible viral reservoir upon PMA-stimulation after culturing with DDX3 inhibitors was

measured by three parallel methods of reservoir quantitation: cell-associated vRNA

quantification by nested Gag RT-qPCR, FISH-flow and TILDA (Fig. 5a). We first confirmed

that culturing patient CD4+ T cells with DDX3 inhibitors for five days did not significantly

impair viability (Supplementary Fig. 5d), consistent with the mild change in overall gene

expression observed by RNA sequencing. Strikingly, as determined by FISH-Flow, CD4+ T

cells cultured in presence of the DDX3 inhibitors RK-33 and FH-1321 showed a 31.1 (s.d. ±

28.2) % and 31.8 (s.d. ± 18.2) % decrease in the frequency of vRNA expressing cells

respectively as compared to the mock-treated cells (Fig. 5b and c). Similarly, upon treatment

with PMA-Ionomycin, a significant decrease in cell-associated Gag vRNA production was

observed by nested RT-PCR in the RK-33 and FH-1321-treated cells as compared to the Mock-

treated cells (Fig. 5d and e). For one representative donor, we also conducted TILDA to quantify

the inducible viral reservoir and observed a decrease in the size of the viral reservoir after five

days of treatment with RK-33 (Fig. 5f and g). To confirm that RK-33 and FH-1321 do not impair

PMA-induced HIV-1 reactivation, we examined the effect of co-treatment with PMA and DDX3

inhibitors in the Lassen and Greene *in vitro* CD4+ T cell model of HIV-1 latency. Co-treatment

of cells with PMA together with RK-33 or FH-1321 did not impair PMA-induced reactivation,

confirming that the decrease in reactivation observed is indeed due to a reduced number of

latently-infected cells (Supplementary Fig. 5e). Altogether, our data demonstrates that not only

do DDX3 inhibitors function as latency reversing agents in primary CD4+ T cells from HIV-1-

infected donors, they also selectively induce cell death in the vRNA-expressing cells, resulting in

their elimination and a decrease in the size of the inducible viral reservoir (Fig. 6).

Fig. 5. Prolonged DDX3 inhibition can reduce the size of the inducible viral reservoir

a Schematic of experiment to test the size of the inducible viral reservoir after culturing CD4+ T

cells from PLWHIV with DDX3 inhibitors. CD4+ T cells isolated from HIV-1-infected donors

were cultured in the presence of the DDX3 inhibitors RK-33 and FH-1321 or mock-treated for 5

days. On Day 5, basal CA-RNA levels were first measured and then cells were maximally

stimulated with PMA-Ionomycin for 12-18 hours. On Day 6, the inducible viral reservoir was

measured by FISH-Flow, CA-RNA and TILDA. For all graphs, three different donors are

represented by blue, red and green symbols consistently. **b** Representative FISH-Flow dot plots

from Day 6 of an HIV-1-infected donor as treated in (a) representing 50000 cells from each

condition. **c** The % of vRNA expressing cells on Day 6 as depicted in (b) were quantified and

normalised to the unstimulated condition. Error bars represent mean ± SD from independent

experiments from 3 different donors in duplicate, with a minimum of 100,000 events acquired

for each condition (Unpaired, two-tailed *t*-test). **d** Copies Gag/µg total RNA on Day 5 and Day 6

for each condition was quantified by RT-qPCR **e** The fold decrease of Gag mRNA expression

levels as compared to the mock-treated condition from cells on Day 6 as treated in (a) and

depicted in (d). Error bars represent mean ± SD from independent experiments from at 3

different donors in duplicate (Unpaired, two-tailed *t*-test). **f** Representative TILDA results from

cells treated with DDX3 inhibitors. **g** Fold change of the size of the viral reservoir upon

treatment with RK-33 and FH-1321 as measured by TILDA normalised to the Mock-treated

cells. Error bars represent mean ± SD from three independent experiments from one donor

(Unpaired, two-tailed *t*-test).

Fig. 6

SELECTIVE KILLING OF VIRAL RESERVOIR

Fig. 6. Model for latency reversal and cell death induced by DDX3 inhibition

DDX3 inhibition results in NF-κB activation and BIRC5 and HSPA1B downregulation in both

uninfected bystander and latent proviral-containing cells. The HIV-1 provirus is reactivated

generating vRNA that can activate innate antiviral signaling pathways which when combined

with BIRC5 or HSPA1B downregulation, results in upregulated Bax, downregulated BCl2,

increased cleaved caspase 3 and apoptosis of the vRNA-expressing cells, but leaving the

uninfected cells unharmed, thereby resulting in a reduction in the viral reservoir.

DISCUSSION

We demonstrate that the treatment of latent HIV-1-infected cells with pharmacological inhibitors of the host RNA binding protein DDX3, modestly reverses viral latency and impairs vRNA translation. This leads to accumulation of vRNA in infected cells which activates the innate antiviral signalling pathways [17,18]. In combination with the DDX3 inhibition-induced downregulation of the anti-apoptotic proteins BIRC5 and HSPA1B, this results in the selective induction of apoptosis in HIV-1 vRNA-expressing cells, with minimal toxicity to uninfected/bystander cells (Fig. 6). Importantly, we demonstrate a reduction in the size of the inducible viral reservoir when latently infected cells were cultured in presence of DDX3 inhibitors. Our data supports the proof of concept, that pharmacological induction of selective cell death in HIV-1 infected cells subsequent to latency reversal, may be a viable HIV-1 curative strategy.

The ability of pro-apoptotic molecules to induce NF-κB pathways and reactivate HIV-1 latency is not without precedence. Second mitochondria-derived activator of caspase (SMAC) mimetics such as LCL-161, SBI-0637142, birinipant, Debio1143 and AZD5582 that target the host anti-apoptotic factors XIAP and cIAP1/BIRC2 have been demonstrated to be potent LRAs via the activation of the non-canonical NF-κB pathways [54-56]. Some LRAs have also been demonstrated to promote cell death, as seen with benzolactam derivatives that are PKC agonists and induce apoptosis in latent HIV-1-infected cells [57,58]. A recent report also demonstrated that romidepsin treatment in a clinical trial resulted in an increase in the frequency of apoptotic T cells [59]. Therefore, the link between apoptosis and latency reversal is highly promising to exploit as a strategy to reactivate the viral reservoir and induce its selective elimination and is worthy of

further exploitation. The DDX3 inhibitors described in this report are such compounds that have

both LRA and ICD capability and were able to decrease the size of the inducible viral reservoir.

DDX3 inhibitors are especially interesting as a potential therapeutic class of compounds for use

in curative strategies against HIV-1 because they target multiple steps of the HIV-1 life cycle. In

this report, we describe the effects of DDX3 inhibition on viral reactivation via NF-κB

activation, as also previously characterised [46] and in agreement with a previous study, we found

DDX3 inhibitor–mediated impairment of vRNA export [32,35]. Additionally, DDX3 inhibitors may

also impair the translation of the vRNA [34,36]. Recent studies have demonstrated that the treatment

of PBMCs with DDX3 inhibitors targeting the RNA-binding site results in a decrease in released

infectious viral particles and that these compounds are also effective against HIV-1 strains

harbouring resistant mutations to currently approved antiretroviral drugs [38,60]. This characteristic

impairment of host cell translation is an advantage when applying DDX3 inhibitors towards an

HIV-1 cure strategy since the generation of infectious viral particles is impaired and therefore

treatment with DDX3 inhibitors would prevent both the unnecessary activation of the cell

mediated and humoral immune responses or inflammation induced by viral production.

Additionally, we demonstrate a newly characterised role for DDX3 inhibitors as ICD inducers to

eliminate the viral reservoir. DDX3 inhibitors demonstrated the ability to target the HIV-1-

infected cells selectively, an important characteristic when developing ICD inducers. In this

work, we demonstrate that the DDX3 inhibitors have negligible effects on the viability of

uninfected cells as well as on non HIV-1 vRNA containing cells from PLWHIV at the

concentrations effective in reversing latency and selectively inducing apoptosis in cells

harbouring HIV-1. We hypothesise that this selectivity may arise from the contribution of several

factors: firstly, DDX3 inhibition downregulates the expression of the anti-apoptotic protein

BIRC5 in both healthy and HIV-1-infected cells. Consistent with our observations, BIRC5 has

been shown to be involved in the survival of latent HIV-1-infected cells and treatment with

BIRC5 inhibitors resulted in cell death of HIV-1 protein expressing cells [48]. Our data is thus

consistent with a model whereby BIRC5 is essential to protect cells from the additional stress

induced by the presence of HIV-1. Secondly, HSPA1B that codes for Heat Shock protein 70

(Hsp70) is another gene that we found significantly downregulated in CD4+ T cells in response

to treatment with the DDX3 inhibitor RK-33. Hsp70 has been shown to protect cells from

apoptosis induced by the HIV-1 protein Vpr [49]. Our data thus suggest that one of the mechanisms

by which RK-33 promotes HIV-1 selective cell death is through downregulation of Hsp70,

whose absence leads to induction of cell death in the Vpr-expressing infected cells and not in the

uninfected bystander cells. Additionally, depletion of Hsp70 has been shown to reactivate HIV-1

via NF-κB activation [50], providing another possible mechanism for the observed latency reversal

property of RK-33. Moreover, DDX3 inhibition interferes with vRNA metabolism and the

residual vRNA that does not efficiently generate infectious viral particles could still induce

cellular stress [18], activate innate antiviral signalling pathways [17] and, in combination with the

downregulation of factors important for survival of latently infected cells, result in the selective

killing of HIV-1-infected cells. This selective toxicity is also important when promoting the

translation of these compounds into clinical settings. We validated that both RK-33 and FH-1321

have negligible effect on cell viability and gene expression in primary cells and that they do not

impair the effector function and proliferation of CD4+ T cells. Importantly, these compounds

also do not affect CD8+ T cell effector function and proliferation so that the essential cell-

mediated immune responses would not be affected during treatment. Extensive toxicology,

biodistribution and pharmacokinetics of RK-33 [40] and FH-1321 [38] have been conducted in mice

and rats respectively, and both demonstrate a good toxicology profile in these pre-clinical

studies, thereby advocating for their translation into clinical studies. Next-generation DDX3

inhibitors with significantly lower IC50s have also recently been developed with a very high

potential for clinical advancement [60].

We also present in this work a novel primary CD4+ T cell model that we have established as a

pipeline to further investigate compounds that induce cell death and to test their abilities to

reduce the size of the inducible viral reservoir using three parallel methods of reservoir

quantitation (Fig. 5a). Since the ICD strategy does not depend on other components of the host

immune system such as CD8+ T cells or NK cells, this system can be used to measure pro-

apoptotic compounds that have previously been studied to selectively eliminate HIV-1-

expressing cells. Some examples of compounds that can be tested include Bcl2 agonists such as

venetoclax [61,62] and SMAC mimetics like birinipant, GDC-0152, and embelin [63] that have been

demonstrated to induce cell apoptosis in HIV-1-infected CD4+ T cells, the latter in an

autophagy-dependent manner. Benzolactam derivatives such as BL-V8-310 are PKC agonists

and have been shown to function as LRAs, but also induce apoptosis of reactivating cells [57]. The

activation of p38/JNK pathways signalling by anisomycin resulted in increased cell death of

HIV-1-infected cells [18]. Specific autophagy-inducing peptides Tat-Beclin 1 and Tat-vFLIP-α2

induce a novel Na+/K+-ATPase dependent form of cell death (autosis), to kill latently HIV-

infected CD4+ T cells and prevent viral rebound [64]. In macrophages and microglial cells,

PI3K/Akt inhibitors restrict viral production and promote cell death of infected cells [65]. Another

strategy to induce apoptosis in infected cells is to inhibit viral trafficking and budding of Gag

with synthesized compounds such as L-HIPPO [66]. Pharmacological inhibition of BIRC5, also

one of the genes downregulated by DDX3 expression, has been demonstrated to target HIV-1-

infected cells and reduce the frequency of proviral containing cells *in vitro* [48]. However, some of

these treatments have been associated with global T cell activation and toxicity, selective

myeloid cell type dependent activity or non-reproducibility. Moreover, these studies have not

specifically investigated the contribution of the HIV-1 vRNA in inducing cell death, made

possible here using FISH-Flow technology. Further investigation of these compounds, possibly

in combination with DDX3 inhibitors, will be important to assess their ability to decrease the

size of the inducible viral reservoir in CD4+ T cells obtained from PLWHIV using an *in vitro*

model such as the one we describe in Fig. 5a and also to investigate the contribution of the

vRNA to cell death. These studies suggest that pharmacological induction of selective cell death

in HIV-1-infected cells, alone or in combination with other approaches, including those that

promote extracellular killing of HIV-1-infected cells may prove effective towards an HIV-1 cure.

One of the caveats of this study is that DDX3 inhibitors do not maximally reactivate the HIV-1

provirus as compared to the PMA-treated cells, thereby resulting in reactivation of only a

fraction of the latently infected cells and hence a partial decrease of the viral reservoir but not its

elimination. Therefore, a combinatorial strategy will be important to test in future work,

combining different classes of LRAs to maximally reactivate the latent HIV-1-infected reservoir

together with DDX3 inhibitors with the ultimate aim to induce cell death in all cells harbouring

inducible replication competent HIV-1. Attractive candidate LRAs for inclusion in this

combinatorial approach are de-repressors of HIV-1 LTR chromatin (BAF and HDAC inhibitors)

[53], activators of transcription (PKC agonists), as well as facilitators of HIV-1 transcription

elongation such as the newly characterised Gliotoxin [67]. Using a pipeline to measure vRNA

production and cell death at a single cell level, may enable identification of an optimal

combination of LRA-ICDs that lead to maximal reduction of the viral reservoir.

So far more than 160 compounds have been identified with latency reversing activity, and our

work has identified DDX3 inhibitors as another category of compounds to reverse viral latency

[68]. In addition to latency reversal, DDX3 inhibitors target multiple steps of the HIV-1 replication

cycle: transcription, nucleocytoplasmic export of the vRNA, translation and generation of

infectious viral particles. Importantly, they also promoted ICD and were selectively able to

induce apoptosis in vRNA expressing cells resulting in the decrease in size of the inducible viral

reservoir. Previous combinatorial studies and clinical trials with LRAs and immune-enhancing

clearance strategies demonstrated minimal effects on the clearance of the viral reservoir [4-6].

Although multiple compounds have been identified to induce apoptosis of HIV-1-infected cells

[13], no ICD inducer has been evaluated for capability to reduce the viral reservoir in PLWHIV yet.

Our data support DDX3 inhibitors as a promising new class of ICD inducers that can be utilized

in HIV-1 curative therapies.

MATERIALS AND METHODS

Cell culture and Reagents

Jurkat cells and J-Lat 11.1 cells (Jurkat cells containing integrated full-length HIV-1 genome

mutated in *env* gene and GFP replacing *nef*) [39] were maintained in RPMI-1640 medium (Sigma

Aldrich) supplemented with 7% FBS and 100 μg/ml penicillin-streptomycin at 37 °C in a

humidified, 5% CO_2 atmosphere. J-Lat latent proviruses were reactivated by adding 10 μM of

phorbol 12-myristate 13-acetate (PMA) (Sigma-Aldrich) to the culture media for 18 hours.

Primary CD4+ T cells were isolated from buffy coats from healthy donors by Ficoll gradient

followed by separation via negative selection with RosetteSep Human CD4+ T cell Enrichment

kit (StemCells Technologies) according to manufacturer's instructions. After isolation, CD4+ T

cells were maintained in RPMI-1640 medium (Sigma Aldrich) supplemented with 10% FBS and

100 μg/ml penicillin-streptomycin at 37 °C in a humidified, 5% CO_2 atmosphere. Cells were

activated with 100 ng/ml PMA and 1 μg/ml Ionomycin (Sigma Aldrich). HEK293T cells were

cultured in DMEM medium (Sigma Aldrich) supplemented with 10% FBS and 100 μg/ml

penicillin-streptomycin at 37°C in a humidified, 5% CO_2 atmosphere. RK-33 (SelleckChem)

was used at a concentration of 2 μM, unless indicated otherwise. FH1321 was provided by First

Health Pharmaceuticals and used at concentrations of 1 μM.

Total RNA Isolation and Quantitative RT-PCR (RT-qPCR)

Total RNA was isolated from the cells using TRIzol reagent (Thermo Fisher Scientific)

according to manufacturer's instructions. cDNA synthesis was performed using Superscript II

Reverse Transcriptase (Life Technologies) kit following the manufacturer's protocol. RT-qPCR

was performed using GoTaq qPCR Master Mix (Promega) following the manufacturer's

protocol. Amplification was performed on the CFX Connect Real-Time PCR Detection System

thermocycler (BioRad) using the following thermal cycling program starting with 3 min at 95

°C, followed by 40 cycles of 95 °C for 10 s and 60 °C for 30 s. Specificity of the RT-qPCR

products was assessed by melting curve analysis. Primers used for real-time PCR are listed in

Table 3. Expression data was calculated using 2-ΔΔCt method [69]. GAPDH housekeeping gene

expression was used for normalisation for J-Lat cell lines and Cyclophilin A was used for

primary cells.

NF-κB reporter assay

To generate the plasmids used in this assay, custom gBlock gene fragments (Integrated DNA

technologies) containing either the minimal mouse heat shock protein (Hspa1a) promoter, or four

NF-κB target sites followed by an Hspa1a promoter were designed with flanking NheI-NcoI

restriction sites and then cloned upstream of the luciferase gene in the pGL3 luciferase reporter

plasmid at the NheI-NcoI sites (Promega). Jurkat cells were transfected with pBluescript II

SK(+) (empty vector), pRL Renilla Luciferase Control Reporter Vector (Promega) to normalize

for transfection efficiency, and either the HSP-Luc or the NF-κB-HSP-Luc plasmids by

nucleofection using Amaxa Nucleofector (Amaxa AG-Lonza, Cologne, Germany) as previously

described [52]. Briefly, 10 million cells were centrifuged at 200 g for 10 min at room temperature,

re-suspended in 100 μl of solution R, and nucleofected with 2 μg plasmid DNA (1.5 μg

pBluescript II SK(+), 200 ng pRL and either Hspa1a-pGL3 or NF-κB-Hsp1a-pGL3) using

program O28. Nucleofected cells were re-suspended in 5 ml of pre-warmed, serum-free

antibiotic-free RPMI at 37 °C for 15 min and then plated in 6 ml of pre-warmed complete media.

Twenty-four hours post-nucleofection cells were either mock-treated or treated with 2 μM RK-

33. Luciferase expression was measured 24 hours post treatment in a GloMax 96 microplate

luminometer (Promega) using the Dual-Glo Luciferase assay system (Promega) and normalized

to *Renilla* luciferase expression. Data represent at least three independent experiments.

FISH-Flow

Cells were collected, fixed, permeabilised and subjected to the PrimeFlow RNA assay (Thermo

Fisher Scientific) following the manufacturer's instructions and as described in [20,21]. In primary

CD4+ T cells, cells were first stained in Fixable Viability dye 780 (BD Biosciences) for 20

minutes at room temperature (1:1000 in dPBS) followed by either AnnexinV-450 (BD

Biosciences) or with 2µM CellEvent™ Caspase-3/7 Green Detection Reagent (Thermo Fisher

Scientific) for 30 min at room temperature (1:150 in AnnexinV binding buffer-Biolegend). For

both CD4+ T cells and J-Lat 11.1 cells, mRNA was labelled with a set of 40 probe pairs against

the GagPol region of the vRNA (catalogue number GagPol HIV-1 VF10-10884, Thermo Fisher

Scientific) diluted 1:5 in diluent provided in the kit and hybridized to the target mRNA for 2

hours at 40°C. Positive control probes against the house-keeping gene RPL13A (VA1-13100,

Thermo Fisher Scientific) were used to determine assay efficiency. Samples were washed to

remove excess probes and stored overnight in the presence of RNAsin. Signal amplification was

then performed by sequential 1.5 hour, 40°C incubations with the pre-amplification and

amplification mix. Amplified mRNA was labelled with fluorescently-tagged probes for 1 hour at

40°C. Samples were acquired on a BD LSR Fortessa Analyzer and gates were set using the

unstimulated J-Lat 11.1 control sample or uninfected CD4+ T cells. Analysis was performed

using the FlowJo V10 software (Treestar).

Confocal Microscopy following FISH-Flow

J-Lat 11.1 cells that underwent the FISH-Flow assay were seeded on 18 mm diameter coverslips

and air dried. Coverslips were mounted in ProLong Gold Antifade Reagent (Life Technologies).

Immunofluorescence images were acquired using a confocal microscope (Leica, SP5). All phase

contrast pictures were acquired using a Leica DMIL microscope and a DFC420C camera. Images

were analysed and processed using Fiji / ImageJ (NIH).

Molecular docking

Molecular docking analysis was used to determine the most likely binding mode of RK-33 and

FH1321 to pre-unwound DDX3 conformation. Crystal structure of human DDX3 in complex

with dsRNA (PDB ID code 6O5F) in the pre-unwound confirmation was used as a template to

define the receptor for the docking simulation by elimination or protein chain B and RNA

domain C and D. RK-33 and FH1321 ligand structure were built and energy minimized using the

program Chimera [70]. Molecular docking of RK-33 and FH1321 to pre-unwound DDX3

confirmation was performed using Chimera's AutoDock Vina function [71]. The resulting solutions

were ranked based on highest binding affinity (or lowest binding energy). Figures were created

using PyMol Molecular Graphics System, Version 1.8. 2015 (Schrödinger L).

RNA-seq

RNA was isolated using TRI Reagent (Sigma-Aldrich). RNA-seq was performed according to

the manufacturer's instructions (Illumina) using the TruSeq Stranded mRNA Library Prep kit.

The resulting DNA libraries were sequenced according to the Illumina TruSeq v2 protocol on an

Illumina HiSeq 2500 sequencer. Reads of 50 bp in length were generated. Reads were mapped

against the UCSC genome browser hg19 reference genome with HiSat2 (version 2.1.0). Gene

expression was quantified using htseq-count (version 0.11.2). For all samples, at least 14.4

million reads were generated with counts on 22.6-24.2 thousand expressed genes. Differential

expression analysis of the RNA-seq data was performed using edgeR package run under Galaxy

(https://bioinf-galaxian.erasmusmc.nl/galaxy/). False discovery rate cut-off was set to 0.05 and a

fold change of ±1.5 was counted as differentially expressed. Heat maps were generated using

MORPHEUS (https://software.broadinstitute.org/morpheus/index.html).

Viability and Activation assays

In Jurkat and J-Lat cells, 10 ng/ml Hoescht 33342 was added to the cells and 30 min later, the

viability and reactivation were quantified by measuring 405nm+ and GFP+ expression

respectively by flow cytometry. In primary CD4+ T cells, to determine viability and cell

activation following DDX3 inhibition, cells were collected and stained with Fixable Viability

Dye eFluor® 780 (supplier) and anti-CD25-PE (Becton Dickinson). Mock-treated and cells

treated with PMA-Ionomycin were used as negative and positive controls, respectively. Cells

were stained for 30 min at 4 °C, washed with PBS, and resuspended for flow cytometric analysis.

T cell proliferation and functionality assay

To analyze the effect of the LRA on CD8+ and CD4+ T cells, proliferation and cytokine

expression were analyzed by flow cytometry. Primary CD8+ and CD4+ T cells were isolated

from buffy coats form healthy donors by Ficoll gradient (GE Healthcare) followed by negative

selection with RosetteSep Human CD8+ T Cell Enrichment Cocktail or RosetteSep Human

CD4+ T Cell Enrichment Cocktail (StemCell Technologies) respectively. Isolated cells were left

overnight for recovery. To analyze T cell proliferation capacity, 1 million CD8+ or CD4+ T cells

were stained with 0,07 uM CellTrace Far Red Cell Proliferation dye (Thermo Fisher Scientific)

following manufacturer's instructions. Cells were then cultivated for 72 hours with either

unstimulated or stimulated conditions in the presence of the LRA, and analyzed by flow

cytometry. Stimulation of T cells was performed using Anti-CD3/CD28 coated Dynabeads

(Thermo Fisher Scientific) following manufacturer's protocol. Proliferation capacity was

determined by a decrease in proliferation dye intensity in daughter cells upon cell division. To

analyze T cell functionality by means of cytokine expression 1 million CD8+ or CD4+ T cells

were left untreated or treated with the LRA for 18 hours. Cells were then left unstimulated or

stimulated with 50 ng/mL PMA and 1uM Ionomycin for 7 hours in the presence of a protein

transport inhibitor (BD GolgiPlug^{TM}, BD Biosciences). To stain for intracellular cytokines cells

were washed with PBS supplemented with 3% FBS followed by a fixation and permeabilisation

step with FIX & PERM Kit (Invitrogen) following manufacturer's protocol and incubated with

1:25 BV510 anti-IL2 (563265, BD Biosciences) and PE-Cy7 anti-IFNγ (27-7319-41,

eBioscience) in permeabilisation buffer for 45 minutes at 4C. Stained cells were washed with

PBS supplemented with 3% FBS and analyzed by flow cytometry.

Measurement of de novo protein synthesis

De novo protein synthesis upon DDX3 inhibition was measured by the incorporation of

puromycin into peptide chains [72]. Briefly, cells were incubated with 10 µg/mL puromycin

(Sigma Aldrich) for 10 min before cell lysis. Cells were washed with dPBS and lysed in ice-cold

lysis buffer (100 mM NaCl, 10 mM Tris, pH 7.5, 1 mM EDTA, 0.5% Nonidet P-40, protease and

phosphatase inhibitor cocktail [Roche]). Cell lysates were denatured in Laemmli sample buffer

and incubated for 5 min at 95°C. The proteins were separated by SDS-PAGE and transferred

onto nitrocellulose membranes (Bio-Rad). Membranes were blocked with 5% nonfat milk in

Tris-buffered saline pH 7.4 and 0.5% Tween 20 (TBST) and then incubated with primary

antibodies. After washes with TBST, the membranes were incubated with horseradish

peroxidase-conjugated secondary antibodies (Rockland Immunochemicals) and detected using

Western Lightning Plus-ECL reagent (Perkin-Elmer). Signal intensity and densitometry analyses

were conducted using ImageJ (NIH). Cell extracts were blotted with anti-Puromycin antibody

(12D10 Sigma Aldrich), anti-GAPDH (ab8245, Abcam) and anti-DDX3 (ab128206, Abcam) and

puromycin incorporation was assessed by summating the immunoblot intensity of all *de novo*

synthesized protein bands normalised to the signal intensity of the GAPDH band. CD4+ T cells

were treated with 10 µg/mL Cycloheximide (Sigma Aldrich) for 4 hours to inhibit translation as

a control for the technique.

Primary CD4 + T Cell Infection and *In vitro* HIV-1 latency model generation

Infections were performed using a pseudotyped virus expressing luciferase that were generated

by co-transfecting HXB2 Env together with the HIV-1 backbone plasmid with a luciferase

reporter (pNL4.3.Luc.R-E-) into HEK 293T cells using PEI (Polyethylenimine) transfection

reagent. 48 and 72 hours post-transfection, the pseudovirus-containing supernatant was collected,

filtered through a 0.45 µm filter, aliquoted and concentrated by ultracentrifugation (20 000 g for

1 hour at 4°C). The pellet was resuspended in RPMI-1640 and stored at -80 °C. The *in vitro*

model of HIV-1 latency was set up using the Lassen and Greene method [51] as follows: CD4+ T

cells were infected with the pNL4.3.Luc-R-E- virus by spinoculation (2 hours at 1200 g) 24

hours after isolation and cultured overnight in RPMI 10% FBS and 100 µg/ml penicillin-

streptomycin in presence of Saquinavir Mesylate (5 µM). Eighteen hours after spin-infection

cells were washed and cultured in growth media supplemented with 5 µM Saquinavir Mesylate.

Three days after infection cells were treated with DDX3 inhibitors in presence of Raltegravir (30

µM). HIV-1 molecular clone pNL4.3.Luc.R-E-, HIV-1 HXB2-Env expression vector, Saquinavir

Mesylate and Raltegravir were kindly provided by the NIH AIDS Reagents. HIV-1 molecular

clone pNL4.3.Luc.R-E- and HIV-1 HXB2-Env expression vector were donated by Dr. Nathaniel

Landau and Drs Kathleen Page and Dan Littman, respectively. Cells were harvested 24 hours

after stimulation with DDX3 inhibitors, washed once in PBS and either lysed for RNA

extraction, used for FISH-Flow analysis or lysed in passive lysis buffer to measure luciferase

activity using Luciferase Assay System (Promega). Relative light units were normalized to

protein content determined by Bradford assay (Bio Rad).

Assays with HIV-1-infected donor samples

PBMCs from HIV-1-infected donors were obtained by Ficoll density gradient isolation of

leukapheresed material from donors that were older than 18 years, cART-treated for at least 3

years, with viral loads below 50 copies/ml for more than 12 months and no blips in the past two

years. This study was conducted in accordance with the ethical principles of the Declaration of

Helsinki. HIV-1-infected patient volunteers were informed and provided signed consent to

participate in the study. The study protocol was approved by the Erasmus Medical Centre

Medical Ethics Committee (MEC-2012–583). CD4+ T cell isolation was performed using the

EasySep Human CD4+ T cell Enrichment kit (StemCells Technologies) according to

manufacturer's instructions. After isolation, CD4+ T cells were allowed to rest for at least 4

hours, maintained in RPMI-1640 medium (Sigma Aldrich) supplemented with 10% FBS, 100

μg/ml penicillin-streptomycin and Raltegravir (30 μM) at 37 °C, in a humidified, 5% CO_2

atmosphere. Cells were treated with DDX3 inhibitors or with PMA-Ionomycin as a positive

control for viral reactivation (100 ng/ml PMA and 1 μg/ml Ionomycin). Eighteen hours post

treatment, cells were either lysed in TRIzol for RNA extraction and RT-qPCR or analysed by

FISH-Flow. To measure the depletion in the size of the inducible viral reservoir, cells were

cultured for 5 days in 2 μM RK-33 and 1 μM FH1321 with a change in media and retreatment

after 48 hours. Five days post treatment, cells were treated with PMA-Ionomycin for 12 hours

and collected for TILDA, or after 18 hours for RT-qPCR and FISH-Flow.

812   Cell-associated HIV RNA measurement by nested RT-qPCR

Cell-associated HIV-1 RNA was measured as previously described with minor modifications [73].

Briefly, the first round of the PCR was performed using the primers Gag1 (5'

TCAGCCCAGAAGTAATACCCATGT 3') and SK431 (5'

TGCTATGTCAGTTCCCCTTGGTTCTCT 3') that amplifies a region within the HIV-1 gag

gene in 25 μl of PCR mix containing 4 μl of cDNA, 2.5 μl 10X Platinum Taq buffer, 2 mM

MgCl2, 0.4 mM concentrations of deoxynucleoside triphosphates, 0.2 μl Platinum Taq and 0.3

μM each of both primers. The PCR settings were as follows: 95°C for 5 min, followed by 15

cycles of 95°C for 30 s, 55°C for 30 s, and 72°C for 1 min. The product of the first PCR was

subsequently used as a template in the second, seminested, real-time, PCR amplification. A total

of 2 μl of the first PCR product was added to a PCR mix with final volume 25 μl containing 2.5

μl 10X Platinum Taq buffer, 2 mM MgCl2, 0.4 mM concentrations of deoxynucleoside

triphosphates, 0.2 μl Platinum Taq and 0.2 μM of the primers Gag1 (5'

TCAGCCCAGAAGTAATACCCATGT 3') and Gag2 (5' CACTGTGTTTAGCATGGTGTTT

3'), and 0.075 μM TaqMan dual-labeled fluorescent probe Gag3 (5'

[6FAM]ATTATCAGAAGGAGCCACCCCACAAGA[BHQ1] 3'). Real-time PCR settings were

as follows: 95°C for 5 min, followed by 45 cycles of 95°C for 30s and 60°C for 1 min. The

amplicon sizes were 221 bp for the first PCR and 83 bp for the second (real-time) PCR. Serial

dilutions of plasmid DNA standards were used to generate a standard curve to calculate the

copies of vRNA and then normalised to the amount of input RNA.

Tat/rev induced limiting dilution assay (TILDA)

Total CD4+ T cells were isolated from PBMCs by negative magnetic separation using EasySep

Human CD4+ T cell isolation kit (StemCell Technology). Following isolation, 2 x 10^6 cells/ml

were rested for 3 hours in complete RPMI-1640 (Lonza) at 37°C and 5% CO_2 and then

stimulated for 12 hours with 100 ng/ml phorbol 12-myristate 13-acetate (PMA) and 1 μg /ml

ionomycin (both from Sigma Aldrich). After stimulation, CD4+ T cells were washed and

resuspended in RPMI-1640 at serial dilutions; 1.8 x 10^6 cells/ml, 9 x 10^5 cells/ml, 3 x 10^5

cells/ml and 1 x 10^5 cells/ml. Using modified TILDA PCR conditions, 10 µl of the cell

suspension from each dilution was dispensed into 24 wells of a 96 well plate containing 2 µl

One-step RT-PCR enzyme (Qiagen), 10 µl 5X One-step RT-PCR buffer (Qiagen), 10 µl Triton-

x100 (0.3%), 0.25 µl RNAsin (40U/µl), 2 µl dNTPs (10 mM each), 1 µl of tat 1.4 forward

primer (20 µM) and rev reverse primer (20 µM) (as published in [74]), and nuclease free water to a

final reaction volume of 50 µl. The one-step RT-PCR was run using the following thermocycling

conditions: 50°C for 30 minutes, 95°C for 15 minutes, 25 cycles of 95°C for 1 minute, 55°C for

1 min and 72°C for 2 minutes, and a final extension at 72°C for 5 minutes. Afterwards 2 µl of the

1[st] PCR products were used as input for real time PCR to detect *tat/rev* msRNA. The 20 µl

reaction volume consisted of 5 µl Taqman Fast Advanced Mastermix 4x (Thermo Fisher

Scientific), 0.4 µl of tat 2.0 forward primer and rev reverse primer (each at 20 µM) and *tat/rev*

probe (5 µM) [74]. The real time PCR was performed in a LightCycler 480 Instrument II (Roche)

using the following programme: 95°C for 10 minutes, 45 cycles for 95°C for 10 seconds and

60°C for 30 seconds. Positive wells at each dilution were recorded and used to determine the

frequency of cells expressing *tat/rev* msRNA by maximum likelihood method using ELDA

software [75].

Statistical Analysis

All data are means ± SD of three or more independent biological replicates. Statistical

significance was calculated using two-tailed *t* test and indicated in Fig. legends. Analyses were

performed using Prism version 8.3.0 (GraphPad software).

REFERENCES

1 Finzi, D. *et al.* Identification of a reservoir for HIV-1 in patients on highly active antiretroviral
therapy. *Science* **278**, 1295-1300 (1997).
2 Chun, T. W., Moir, S. & Fauci, A. S. HIV reservoirs as obstacles and opportunities for an HIV cure.
Nat Immunol **16**, 584-589 (2015).
3 Deeks, S. G. HIV: Shock and kill. *Nature* **487**, 439-440 (2012).
4 Delagrèverie, H. M., Delaugerre, C., Lewin, S. R., Deeks, S. G. & Li, J. Z. Ongoing Clinical Trials of
Human Immunodeficiency Virus Latency-Reversing and Immunomodulatory Agents. *Open*
Forum Infect Dis **3**, ofw189-ofw189, doi:10.1093/ofid/ofw189 (2016).
5 Rasmussen, T. A. & Lewin, S. R. Shocking HIV out of hiding: where are we with clinical trials of
latency reversing agents? *Curr Opin HIV AIDS* **11**, 394-401 (2016).
6 Zerbato, J. M., Purves, H. V., Lewin, S. R. & Rasmussen, T. A. Between a shock and a hard place:
challenges and developments in HIV latency reversal. *Curr Opin Virol* **38**, 1-9 (2019).
7 Thorlund, K., Horwitz, M. S., Fife, B. T., Lester, R. & Cameron, D. W. Landscape review of current
HIV 'kick and kill' cure research - some kicking, not enough killing. *BMC Infect Dis* **17**, 595-595,
doi:10.1186/s12879-017-2683-3 (2017).
8 Leth, S. *et al.* Combined effect of Vacc-4x, recombinant human granulocyte macrophage colony-
stimulating factor vaccination, and romidepsin on the HIV-1 reservoir (REDUC): a single-arm,
phase 1B/2A trial. *Lancet HIV* **3**, e463-472 (2016).
9 Fidler, S. *et al.* Antiretroviral therapy alone versus antiretroviral therapy with a kick and kill
approach, on measures of the HIV reservoir in participants with recent HIV infection (the RIVER
trial): a phase 2, randomised trial. *Lancet* **395**, 888-898 (2020).
10 Gay, C. L. *et al.* Assessing the impact of AGS-004, a dendritic cell-based immunotherapy, and
vorinostat on persistent HIV-1 Infection. *Sci Rep* **10**, 5134 (2020).
11 Margolis, D. M. *et al.* Curing HIV: Seeking to Target and Clear Persistent Infection. *Cell* (2020).
12 Cummins, N. W. & Badley, A. D. Mechanisms of HIV-associated lymphocyte apoptosis: 2010. *Cell*
Death Dis **1**, e99-e99, doi:10.1038/cddis.2010.77 (2010).
13 Kim, Y., Anderson, J. L. & Lewin, S. R. Getting the "Kill" into "Shock and Kill": Strategies to
Eliminate Latent HIV. *Cell Host Microbe* **23**, 14-26 (2018).
14 Cooney, J., Allison, C., Preston, S. & Pellegrini, M. Therapeutic manipulation of host cell death
pathways to facilitate clearance of persistent viral infections. *J Leukoc Biol* **103**, 287-293 (2018).
15 Baxter, A. E. *et al.* Single-Cell Characterization of Viral Translation-Competent Reservoirs in HIV-
Infected Individuals. *Cell Host Microbe* **20**, 368-380 (2016).
16 Grau-Exposito, J. *et al.* A Novel Single-Cell FISH-Flow Assay Identifies Effector Memory CD4(+) T
cells as a Major Niche for HIV-1 Transcription in HIV-Infected Patients. *mBio* **8** (2017).
17 Berg, R. K. *et al.* Genomic HIV RNA induces innate immune responses through RIG-I-dependent
sensing of secondary-structured RNA. *PLoS One* **7**, e29291 (2012).
18 Fong, L. E., Sulistijo, E. S. & Miller-Jensen, K. Systems analysis of latent HIV reversal reveals
altered stress kinase signaling and increased cell death in infected T cells. *Sci Rep* **7**, 16179-
16179, doi:10.1038/s41598-017-15532-0 (2017).
19 Baxter, A. E., O'Doherty, U. & Kaufmann, D. E. Beyond the replication-competent HIV reservoir:
transcription and translation-competent reservoirs. *Retrovirology* **15**, 18 (2018).
20 Baxter, A. E. *et al.* Multiparametric characterization of rare HIV-infected cells using an RNA-flow
FISH technique. *Nat Protoc* **12**, 2029-2049 (2017).

21 Rao, S., Amorim, R., Niu, M., Temzi, A. & Mouland, A. J. The RNA surveillance proteins UPF1,
UPF2 and SMG6 affect HIV-1 reactivation at a post-transcriptional level. *Retrovirology* **15**, 42
(2018).
22 Grau-Expósito, J. *et al.* Latency reversal agents affect differently the latent reservoir present in
distinct CD4+ T subpopulations. *PLoS Pathog* **15**, e1007991 (2019).
23 Linder, P. & Jankowsky, E. From unwinding to clamping — the DEAD box RNA helicase family.
Nature Reviews Molecular Cell Biology **12**, 505-516, doi:10.1038/nrm3154 (2011).
24 Zhao, L. *et al.* Multifunctional DDX3: dual roles in various cancer development and its related
signaling pathways. *Am J Cancer Res* **6**, 387-402 (2016).
25 Soto-Rifo, R. & Ohlmann, T. The role of the DEAD-box RNA helicase DDX3 in mRNA metabolism.
Wiley Interdiscip Rev RNA **4**, 369-385 (2013).
26 Lin, T. C. DDX3X Multifunctionally Modulates Tumor Progression and Serves as a Prognostic
Indicator to Predict Cancer Outcomes. *Int J Mol Sci* **21** (2019).
27 Bol, G. M., Xie, M. & Raman, V. DDX3, a potential target for cancer treatment. *Mol Cancer* **14**,
188 (2015).
28 Heerma van Voss, M. R., van Diest, P. J. & Raman, V. Targeting RNA helicases in cancer: The
translation trap. *Biochim Biophys Acta Rev Cancer* **1868**, 510-520 (2017).
29 He, Y. *et al.* A double-edged function of DDX3, as an oncogene or tumor suppressor, in cancer
progression (Review). *Oncol Rep* **39**, 883-892 (2018).
30 Kukhanova, M. K., Karpenko, I. L. & Ivanov, A. V. DEAD-box RNA Helicase DDX3: Functional
Properties and Development of DDX3 Inhibitors as Antiviral and Anticancer Drugs. *Molecules* **25**
(2020).
31 Valiente-Echeverria, F., Hermoso, M. A. & Soto-Rifo, R. RNA helicase DDX3: at the crossroad of
viral replication and antiviral immunity. *Rev Med Virol* **25**, 286-299 (2015).
32 Yedavalli, V. S., Neuveut, C., Chi, Y. H., Kleiman, L. & Jeang, K. T. Requirement of DDX3 DEAD box
RNA helicase for HIV-1 Rev-RRE export function. *Cell* **119**, 381-392 (2004).
33 Mahboobi, S. H., Javanpour, A. A. & Mofrad, M. R. The interaction of RNA helicase DDX3 with
HIV-1 Rev-CRM1-RanGTP complex during the HIV replication cycle. *PLoS One* **10**, e0112969
(2015).
34 Soto-Rifo, R. *et al.* DEAD-box protein DDX3 associates with eIF4F to promote translation of
selected mRNAs. *Embo J* **31**, 3745-3756 (2012).
35 Frohlich, A. *et al.* DEAD-box RNA helicase DDX3 connects CRM1-dependent nuclear export and
translation of the HIV-1 unspliced mRNA through its N-terminal domain. *Biochim Biophys Acta*
1859, 719-730 (2016).
36 Soto-Rifo, R., Rubilar, P. S. & Ohlmann, T. The DEAD-box helicase DDX3 substitutes for the cap-
binding protein eIF4E to promote compartmentalized translation initiation of the HIV-1 genomic
RNA. *Nucleic Acids Res* **41**, 6286-6299 (2013).
37 Gringhuis, S. I. *et al.* Erratum: HIV-1 blocks the signaling adaptor MAVS to evade antiviral host
defense after sensing of abortive HIV-1 RNA by the host helicase DDX3. *Nat Immunol* **18**, 474
(2017).
38 Brai, A. *et al.* Human DDX3 protein is a valuable target to develop broad spectrum antiviral
agents. *Proc Natl Acad Sci U S A* **113**, 5388-5393 (2016).
39 Jordan, A., Bisgrove, D. & Verdin, E. HIV reproducibly establishes a latent infection after acute
infection of T cells in vitro. *Embo J* **22**, 1868-1877 (2003).
40 Bol, G. M. *et al.* Targeting DDX3 with a small molecule inhibitor for lung cancer therapy. *EMBO*
Mol Med **7**, 648-669 (2015).
41 Kondaskar, A. *et al.* Novel, Broad Spectrum Anti-Cancer Agents Containing the Tricyclic 5:7:5-
Fused Diimidazodiazepine Ring System. *ACS Med Chem Lett* **2**, 252-256 (2010).

42   Heerma van Voss, M. R. *et al.* Identification of the DEAD box RNA helicase DDX3 as a therapeutic
target in colorectal cancer. *Oncotarget* **6**, 28312-28326 (2015).
43   Xie, M. *et al.* RK-33 Radiosensitizes Prostate Cancer Cells by Blocking the RNA Helicase DDX3.
Cancer Res **76**, 6340-6350 (2016).
44   Heerma van Voss, M. R. *et al.* Targeting mitochondrial translation by inhibiting DDX3: a novel
radiosensitization strategy for cancer treatment. *Oncogene* **37**, 63-74 (2018).
45   Wang, X. *et al.* (DEAD)-box RNA helicase 3 modulates NF-kappaB signal pathway by controlling
the phosphorylation of PP2A-C subunit. *Oncotarget* **8**, 33197-33213 (2017).
46   Xiang, N. *et al.* The DEAD-Box RNA Helicase DDX3 Interacts with NF-κB Subunit p65 and
Suppresses p65-Mediated Transcription. *PloS one* **11**, e0164471-e0164471,
doi:10.1371/journal.pone.0164471 (2016).
47   Radi, M. *et al.* Discovery of the first small molecule inhibitor of human DDX3 specifically
designed to target the RNA binding site: towards the next generation HIV-1 inhibitors. *Bioorg*
Med Chem Lett **22**, 2094-2098 (2012).
48   Kuo, H. H. *et al.* Anti-apoptotic Protein BIRC5 Maintains Survival of HIV-1-Infected CD4(+) T Cells.
Immunity **48**, 1183-1194 e1185 (2018).
49   Iordanskiy, S. *et al.* Heat shock protein 70 protects cells from cell cycle arrest and apoptosis
induced by human immunodeficiency virus type 1 viral protein R. *J Virol* **78**, 9697-9704,
doi:10.1128/jvi.78.18.9697-9704.2004 (2004).
50   Chaudhary, P. *et al.* HSP70 binding protein 1 (HspBP1) suppresses HIV-1 replication by inhibiting
NF-κB mediated activation of viral gene expression. *Nucleic acids research* **44**, 1613-1629,
doi:10.1093/nar/gkv1151 (2016).
51   Lassen, K. G., Hebbeler, A. M., Bhattacharyya, D., Lobritz, M. A. & Greene, W. C. A flexible model
of HIV-1 latency permitting evaluation of many primary CD4 T-cell reservoirs. *PloS one* **7**,
e30176-e30176, doi:10.1371/journal.pone.0030176 (2012).
52   Stoszko, M. *et al.* Small Molecule Inhibitors of BAF; A Promising Family of Compounds in HIV-1
Latency Reversal. *EBioMedicine* **3**, 108-121, doi:10.1016/j.ebiom.2015.11.047 (2015).
53   Marian, C. A. *et al.* Small Molecule Targeting of Specific BAF (mSWI/SNF) Complexes for HIV
Latency Reversal. *Cell Chem Biol* **25**, 1443-1455 e1414 (2018).
54   Pache, L. *et al.* BIRC2/cIAP1 Is a Negative Regulator of HIV-1 Transcription and Can Be Targeted
by Smac Mimetics to Promote Reversal of Viral Latency. *Cell Host Microbe* **18**, 345-353 (2015).
55   Nixon, C. C. *et al.* Systemic HIV and SIV latency reversal via non-canonical NF-kappaB signalling in
vivo. *Nature* **578**, 160-165 (2020).
56   Bobardt, M. *et al.* The inhibitor apoptosis protein antagonist Debio 1143 Is an attractive HIV-1
latency reversal candidate. *PLoS One* **14**, e0211746 (2019).
57   Matsuda, K. *et al.* Benzolactam-related compounds promote apoptosis of HIV-infected human
cells via protein kinase C-induced HIV latency reversal. *J Biol Chem* **294**, 116-129,
doi:10.1074/jbc.RA118.005798 (2019).
58   Hattori, S. I. *et al.* Combination of a Latency-Reversing Agent With a Smac Mimetic Minimizes
Secondary HIV-1 Infection in vitro. *Front Microbiol* **9**, 2022 (2018).
59   Rosás-Umbert, M. *et al.* In vivo Effects of Romidepsin on T-Cell Activation, Apoptosis and
Function in the BCN02 HIV-1 Kick&Kill Clinical Trial. *Frontiers in Immunology* **11**,
doi:10.3389/fimmu.2020.00418 (2020).
60   Brai, A. *et al.* DDX3X inhibitors, an effective way to overcome HIV-1 resistance targeting host
proteins. *Eur J Med Chem* **200**, 112319 (2020).
61   Cummins, N. W. *et al.* Maintenance of the HIV Reservoir Is Antagonized by Selective BCL2
Inhibition. *J Virol* **91**, e00012-00017, doi:10.1128/jvi.00012-17 (2017).

62 Chandrasekar, A. P., Cummins, N. W. & Badley, A. D. The Role of the BCL-2 Family of Proteins in
HIV-1 Pathogenesis and Persistence. *Clin Microbiol Rev* **33** (2019).
63 Campbell, G. R., Bruckman, R. S., Chu, Y. L., Trout, R. N. & Spector, S. A. SMAC Mimetics Induce
Autophagy-Dependent Apoptosis of HIV-1-Infected Resting Memory CD4+ T Cells. *Cell Host*
Microbe **24**, 689-702 e687 (2018).
64 Zhang, G. *et al.* Selective cell death of latently HIV-infected CD4(+) T cells mediated by autosis
inducing nanopeptides. *Cell Death Dis* **10**, 419 (2019).
65 Lucas, A. *et al.* Targeting the PI3K/Akt Cell Survival Pathway to Induce Cell Death of HIV-1
Infected Macrophages with Alkylphospholipid Compounds. *PLOS ONE* **5**, e13121 (2010).
66 Tateishi, H. *et al.* A clue to unprecedented strategy to HIV eradication: "Lock-in and apoptosis".
Sci Rep **7**, 8957-8957, doi:10.1038/s41598-017-09129-w (2017).
67 Stoszko, M. *et al.* Gliotoxin, identified from a screen of fungal metabolites, disrupts 7SK snRNP,
releases P-TEFb and reverses HIV-1 latency. *bioRxiv*, 848929, doi:10.1101/848929 (2019).
68 Stoszko, M., Ne, E., Abner, E. & Mahmoudi, T. A broad drug arsenal to attack a strenuous latent
HIV reservoir. *Current Opinion in Virology* **38**, 37-53,
doi:https://doi.org/10.1016/j.coviro.2019.06.001 (2019).
69 Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C(T) method.
Nat Protoc **3**, 1101-1108, doi:https://doi.org/10.1038/nprot.2008.73 (2008).
70 Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and analysis.
J Comput Chem **25**, 1605-1612 (2004).
71 Trott, O. & Olson, A. J. AutoDock Vina: improving the speed and accuracy of docking with a new
scoring function, efficient optimization, and multithreading. *J Comput Chem* **31**, 455-461 (2010).
72 Schmidt, E. K., Clavarino, G., Ceppi, M. & Pierre, P. SUnSET, a nonradioactive method to monitor
protein synthesis. *Nature Methods* **6**, 275-277, doi:10.1038/nmeth.1314 (2009).
73 Pasternak, A. O. *et al.* Highly Sensitive Methods Based on Seminested Real-Time Reverse
Transcription-PCR for Quantitation of Human Immunodeficiency Virus Type 1 Unspliced and
Multiply Spliced RNA and Proviral DNA. *Journal of Clinical Microbiology* **46**, 2206,
doi:10.1128/jcm.00055-08 (2008).
74 Procopio, F. A. *et al.* A Novel Assay to Measure the Magnitude of the Inducible Viral Reservoir in
HIV-infected Individuals. *EBioMedicine* **2**, 874-883 (2015).
75 Hu, Y. & Smyth, G. K. ELDA: extreme limiting dilution analysis for comparing depleted and
enriched populations in stem cell and other assays. *J Immunol Methods* **347**, 70-78 (2009).

1033

1034

ACKNOWLEDGEMENTS

We would like to thank Jan-Willem Bakker, Alessia Tarditi and Matteo Andreini from First Health Pharmaceuticals Amsterdam for generous provision of the DDX3 inhibitor FH-1321; and Alicja Gorska and Tsung Wai Kan for technical help.

FUNDING

The research leading to these results has received funding from the Dutch AIDS Fonds (grants P-53302 and P-53601), Health Holland (grants LSHM19100-SGF and EMCLSH19023), ZonMW (grant 40-44600-98-333), the Federation of Medical Specialists (grant 59825822) and the EHVA T01 consortium, which is supported by the European Union's Horizon 2020 Research and Innovation Programme (grant 681032).

AUTHOR CONTRIBUTIONS

S.R. and T.M designed the studies; S.R., C.L., T.H.S., and R.C conducted experiments and performed data analysis; S.R., R.J.P. and W.V.I participated in gene expression analysis; H.P., C.R. and A.V. recruited patients for the study; Y.M.M., P.D.K., J.V.K., C.B., R.G. and T.M. provided expertise and supervision; S.R. and T.M. wrote the manuscript with input from all authors. All authors read and approved the final manuscript.

COMPETING INTERESTS

1054

1055   The authors declare that they have no competing interests.

1056

DATA AND MATERIALS AVAILABILITY

1057

1058   All data needed to evaluate the conclusions in the paper are present in the paper and/or the

1059   Supplementary Materials. Additional data related to this paper may be requested from the

1060   authors.

1061

1062

 Table 1: EdgeR report of differentially expressed genes 18h post treatment with 2µM RK-

 33

GeneID	Name	log2 Fold Change	log2 Counts Per Million	F Value	P Value	False Discovery Rate
ENSG00000133124	IRS4	-9.220240267	1.229125796	149.1749	2.89E-34	1.69E-29
ENSG00000096696	DSP	-8.484792295	0.549901848	93.17747	4.97E-22	9.65E-18
ENSG00000145147	SLIT2	-8.122797551	0.215628794	69.49186	7.84E-17	1.14E-12
ENSG00000261409	AL035425.3	-8.022265953	0.113605342	60.57704	7.19E-15	5.99E-11
ENSG00000138829	FBN2	-7.721874417	-0.141878891	49.16692	2.38E-12	1.39E-08
ENSG00000083857	FAT1	-7.711916601	-0.142206665	50.58668	1.15E-12	7.47E-09
ENSG00000154277	UCHL1	-7.629659662	-0.220071849	45.5358	1.51E-11	7.35E-08
ENSG00000139219	COL2A1	-7.594048405	-0.247211954	44.80238	2.20E-11	9.86E-08
ENSG00000229807	XIST	-7.425086724	4.779411144	281.6285	8.79E-06	0.013432
ENSG00000089685	BIRC5	-7.372112695	-0.421255852	38.53409	5.42E-10	2.26E-06
ENSG00000116962	NID1	-7.05272162	-0.69051715	28.24878	1.07E-07	0.000297
ENSG00000038427	VCAN	-7.017959819	-0.727725687	25.70546	3.99E-07	0.000861
ENSG00000176887	SOX11	-7.004754934	-0.728056321	27.13324	1.91E-07	0.000483
ENSG00000117114	ADGRL2	-6.95027203	-0.766705238	26.38183	2.81E-07	0.000655
ENSG00000054598	FOXC1	-6.756514098	-0.931471077	19.71939	8.99E-06	0.013432
ENSG00000188153	COL4A5	-6.756514098	-0.931471077	19.71939	8.99E-06	0.013432
ENSG00000197565	COL4A6	-6.610183248	-1.02235097	19.90055	8.17E-06	0.013236
ENSG00000169855	ROBO1	-6.551036428	-1.069611081	18.62718	1.59E-05	0.021095
ENSG00000170558	CDH2	-6.544969004	-1.069721671	18.90017	1.38E-05	0.019153
ENSG00000159217	IGF2BP1	-5.889308666	0.348421941	64.80318	8.72E-16	1.02E-11
ENSG00000135862	LAMC1	-5.486512767	0.002490741	48.29217	3.71E-12	1.97E-08
ENSG00000102024	PLS3	-4.323906265	1.49827749	117.2158	3.02E-27	8.81E-23
ENSG00000171246	NPTX1	-3.992292706	-0.550303193	23.25519	1.42E-06	0.002675
ENSG00000154734	ADAMTS1	-3.971030294	0.580596315	60.82655	6.34E-15	5.99E-11
ENSG00000145555	MYO10	-3.696600515	-0.303481708	26.65852	2.44E-07	0.000592
ENSG00000154380	ENAH	-3.64365296	-0.274861002	25.06365	5.59E-07	0.001124
ENSG00000142949	PTPRF	-3.553235332	-0.021528142	33.79895	6.35E-09	1.95E-05

ENSG00000163430	FSTL1	-3.20753747	-0.332407149	22.31002	2.33E-06	0.003987
ENSG00000117399	CDC20	-3.098292124	-0.420984699	17.47004	2.92E-05	0.034778
ENSG00000176788	BASP1	-3.03195077	-0.24810378	22.38461	2.24E-06	0.003952
ENSG00000111206	FOXM1	-3.015861999	0.001271235	28.7242	8.38E-08	0.000244
ENSG00000046604	DSG2	-2.684610226	0.484772447	34.90713	3.48E-09	1.19E-05
ENSG00000101057	MYBL2	-2.316037106	0.025267442	17.38763	3.06E-05	0.035688
ENSG00000148773	MKI67	-1.384356143	1.091562301	18.66531	1.56E-05	0.021095
ENSG00000083444	PLOD1	-1.18441556	1.399169378	17.50631	2.88E-05	0.034778
ENSG00000131747	TOP2A	-1.054906153	1.866051914	19.13111	1.25E-05	0.017772
ENSG00000204388	HSPA1B	-0.741213605	3.325135537	27.77549	1.39E-07	0.000367
ENSG00000178429	RPS3AP5	2.015230135	1.279297836	36.33718	3.41E-09	1.19E-05
ENSG00000234648	AL162151.2	4.827847593	0.344327701	50.73859	1.07E-12	7.47E-09

1065

1066

Table 2: EdgeR report of differentially expressed genes 18h post treatment with 1µM FH-

 1321

GeneID	Name	log2 Fold Change	log2 Counts Per Million	F value	PValue	False Discovery Rate
ENSG00000154277	UCHL1	-4.565471687	-0.206793672	38.67266739	5.05E-10	3.31E-06
ENSG00000261409	AL035425.3	-4.424650789	0.138068277	48.33716477	3.63E-12	3.52E-08
ENSG00000138829	FBN2	-4.131894042	-0.10684436	38.38083941	5.86E-10	3.42E-06
ENSG00000116962	NID1	-3.999181717	-0.658930046	22.77369298	1.83E-06	0.0053254
ENSG00000117114	ADGRL2	-3.832053102	-0.731611642	20.4765316	6.05E-06	0.0125945
ENSG00000159217	IGF2BP1	-3.68912559	0.398606752	48.96044087	2.64E-12	3.08E-08
ENSG00000229807	XIST	-3.488985213	4.901654584	651.1533715	7.66E-143	4.47E-138
ENSG00000038427	VCAN	-3.429296401	-0.658534387	18.19677341	2.00E-05	0.0323138
ENSG00000154734	ADAMTS1	-3.363272907	0.603352768	54.52344803	1.56E-13	2.27E-09
ENSG00000212402	SNORA74B	-3.359506375	0.031288549	24.79248338	5.34E-06	0.0115336
ENSG00000145147	SLIT2	-3.051459541	0.34517457	38.64546164	5.12E-10	3.31E-06
ENSG00000101057	MYBL2	-2.996889004	-0.083099003	26.07056348	3.34E-07	0.0010805
ENSG00000133124	IRS4	-2.98565677	1.371029055	71.98000489	2.22E-17	6.48E-13
ENSG00000083857	FAT1	-2.913815462	-0.014539277	28.05815995	1.18E-07	0.0004052
ENSG00000145555	MYO10	-2.805586002	-0.261404374	22.18212809	2.49E-06	0.0066275
ENSG00000139219	COL2A1	-2.783583292	-0.107869982	24.02115361	9.55E-07	0.0029316
ENSG00000201098	RNY1	-2.770572459	0.413772602	35.10682622	4.80E-09	2.33E-05
ENSG00000135862	LAMC1	-2.712320987	0.135877854	28.32916412	1.03E-07	0.0003743
ENSG00000102024	PLS3	-2.681318527	1.605768497	71.57130553	3.77E-17	7.32E-13
ENSG00000096696	DSP	-2.636756013	0.730407853	42.41260665	7.45E-11	6.21E-07
ENSG00000089685	BIRC5	-2.490957592	-0.234929297	18.96212263	1.34E-05	0.02512
ENSG00000238741	SCARNA7	-2.387719434	0.361869871	24.20859794	2.61E-06	0.0066275
ENSG00000163430	FSTL1	-2.380561336	-0.26218126	17.71426598	2.57E-05	0.0405086
ENSG00000200785	SNORD8	-2.299517907	-0.084333496	18.58796384	1.64E-05	0.0290541
ENSG00000142949	PTPRF	-2.256619192	0.113519171	22.13114508	2.55E-06	0.0066275
ENSG00000111206	FOXM1	-2.226389714	0.092218251	21.57249486	3.41E-06	0.0082946
ENSG00000212283	SNORD89	-1.860714666	1.311190667	36.42603494	1.90E-09	1.01E-05

ENSG00000208772	SNORD94	-1.822933237	1.022233292	29.05095049	7.20E-08	0.0002797
ENSG00000166165	CKB	-1.686278111	1.500619457	33.19463106	9.30E-09	4.17E-05
ENSG00000046604	DSG2	-1.609604699	0.644434181	18.28993873	1.90E-05	0.031651
ENSG00000202058	RN7SKP80	-1.234932643	1.641387087	21.21111392	4.96E-06	0.011119
ENSG00000126016	AMOT	-0.773124251	3.662624092	30.45093151	3.53E-08	0.0001469
ENSG00000130164	LDLR	-0.411325614	5.764550633	21.07423587	4.43E-06	0.0103253
ENSG00000101745	ANKRD12	0.339396244	7.502647544	18.65815618	1.57E-05	0.0285379
ENSG00000170881	RNF139	0.423776234	5.074578288	18.46989874	1.74E-05	0.0297963
ENSG00000213757	AC020898.1	0.876931574	2.474353677	19.06819338	1.27E-05	0.0246334
ENSG00000198064	NPIPB13	2.365643737	0.043629688	19.87731825	8.31E-06	0.0167005

1069

1070

1071 **Table 3: List of primers used in this study**

Target gene:	Forward:	Reverse:
GFP	5'-GAAGCAGCACGACTTCTTCAA-3'	5'-GCTTGTCGGCCATGATATAGA-3'
Pol	5'-GGTTTATTACAGGGACAGCAGAGA-3'	5'-ACCTGCCATCTGTTTTCCATA-3'
MMP9	5'-TGGTCCTGGTGCTCCTGGTG-3'	5'-GCTGCCTGTCGGTGAGATTGG-3'
BCL2	5'-GGTGGGGTCATGTGTGTGG -3'	5'-CGGTTCAGGTACTCAGTCATCC-3'
Bax	5'-CCCGAGAGGTCTTTTTCCGAG -3'	5'-CCAGCCCATGATGGTTCTGAT -3'
IL10	5'-CATCGATTTCTTCCCTGTGAA-3'	5'-TCTTGGAGCTTATTAAAGGCATTC-3'
GAPDH	5'-CAAGAAGGTGGTGAAGCAG-3'	5'-GCCAAATTCGTTGTCATACC-3'
RIG-I	5'-CCAAGCCAAAGCAGTTTTCAAG -3'	5'-CATGGATTCCCCAGTCATGG-3'
MAVS	5'-TGATTTCCTCGCAATCAGACG -3'	5'-GAAGCCGATTTCCAGCTGTATG-3'
BIRC5	5'-CAAGGACCACCGCATCTCTAC-3'	5'-AGTCTGGCTCGTTCTCAGTGG-3'
HSPA1B	5'-ACCTTCGACGTGTCCATCCTGA -3'	5'-TCCTCCACGAAGTGGTTCACCA -3'
IL8	5'-CTCTCTTGGCAGCCTTCC -3'	5'-TCCACTCTCAATCACTCTCAG-3'
Cyclophilin A	5'-TCATCTGCACTGCCAAGACTG-3'	5'-CATGCCTTCTTTCACTTTGCC-3'

1072

1073

1074

1075

1076

1077