Gene dosage screens in yeast reveal core signalling pathways controlling heat adaptation

Cosimo Jann[1,2,*], Andreas Johansson[1], Justin D. Smith[3,4], Leopold Parts[5,6,*], Lars M. Steinmetz[1,3,4,*]

[1]European Molecular Biology Laboratory (EMBL), Genome Biology Unit, Heidelberg, Germany.

[2]ETH Zurich, Department of Biology, Institute of Biochemistry, Zurich, Switzerland.

[3]Department of Genetics, Stanford University School of Medicine, Stanford, California, USA.

[4]Stanford Genome Technology Center, Stanford University, Palo Alto, California, USA.

[5]Wellcome Sanger Institute, Hinxton, UK.

[6]Department of Computer Science, University of Tartu, Tartu, Estonia.

*Correspondence to C.J. (cosimo.jann@embl.de) or L.P. (leopold.parts@sanger.ac.uk) or L.M.S.
(lars.steinmetz@embl.de).

Abstract

Heat stress causes proteins to unfold and lose their function, jeopardizing essential cellular processes.
To protect against heat and proteotoxic stress, cells mount a dedicated stress-protective programme,
the so-called heat shock response (HSR). Our understanding of the mechanisms that regulate the HSR
and their contributions to heat resistance and growth is incomplete. Here we employ CRISPRi/a to
down- or upregulate protein kinases and transcription factors in *S. cerevisiae*. We measure gene
functions by quantifying perturbation effects on HSR activity, thermotolerance, and cellular fitness at
23, 30 and 38°C. The integration of these phenotypes allowed us to identify core signalling pathways
of heat adaptation and reveal novel functions for the high osmolarity glycerol, unfolded protein
response and protein kinase A pathways in adjusting both thermotolerance and chaperone expression.
We further provide evidence for unknown cross-talk of the HSR with the cell cycle-dependent kinase
Cdc28, the primary regulator of cell cycle progression. Finally, we show that CRISPRi efficiency is
temperature-dependent and that different phenotypes vary in their sensitivity to knock-down. In
summary, our study quantifies regulatory gene functions in different aspects of heat adaptation and
advances our understanding of how eukaryotic cells counteract proteotoxic and other heat-caused
damage.

Keywords: CRISPRi/a, CRISPR/dCas9 screen, heat shock response, thermotolerance, Hsf1, HOG pathway,
PKA signalling.

Introduction

When exposed to high temperature, cells need to resist the proteotoxicity due to protein misfolding and aggregation[1,2] and other heat-caused damage[3]. They do so by eliciting a series of stress-protective events, referred to as the heat shock response (HSR)[4]. The HSR is highly conserved across eukaryotes and characterized by the fierce production of heat shock proteins (HSPs) which mostly function in maintaining protein homeostasis[5,6]. Dysregulation of the HSR results in altered chaperone capacity and is linked to neurodegenerative diseases[7,8] and aging[9] where decreased HSR activity aggravates proteotoxicity. Cancers also hi-jack and increase HSR activity to cope with their proteotoxic burden[10]. In the last two decades, heat-induced changes to the transcriptome and proteome have been well characterized[11–17]. In budding yeast, a shift from 30 to 37°C causes around a thousand genes to change transcription[11,17], while exposure to 42°C results in expression changes for more than 50% of the yeast genome (~3100 genes), with higher magnitude and longer upkeep compared to 37°C[17]. Much less is known about the mechanisms that enable regulation of this response which is essential to safeguard cellular survival.

Heat shock factor 1 (Hsf1) is considered the master HSR regulator, in yeast acting together with the general stress response factors Msn2 and Msn4[18,19]. Hsf1 is regulated through titration by chaperones[20–23] and hyperphosphorylation[24,25]. Dissection of the Hsf1-driven HSR in yeast[26] and human cells[27] revealed new mechanisms controlling Hsf1. However, recent studies demonstrate that the HSR remains largely unchanged when Hsf1 is absent in yeast and mammalian cells, and mainly driven by other transcriptional regulators[15,28]. In addition, even for Hsf1, Msn2 and Msn4, the most prominent transcription factors (TFs) of the yeast HSR, the protein kinases (PKs) mediating their heat-induced hyperphosphorylation and activation remain elusive[29]. Individual TFs are likely controlled by an interplay of signalling pathways that, apart from transcription[15], may also affect mRNA localization[30], stability[31] and translation[32].

HSR overlaps with oxidative and general stress responses[11,19] which trigger cell cycle arrest and the cell wall integrity (CWI) pathway[33,34]. The HSR also inhibits target of rapamycin (TOR) signalling[35] and is itself repressed by protein kinase A (PKA)[36,37]. High temperature further activates the high osmolarity glycerol (HOG) pathway, although its role is unknown[38–40]. A comprehensive understanding of how signalling programmes integrate to regulate the HSR is missing. In addition, it is unclear which molecular branches of the HSR contribute to cellular protection, given that the bulk of heat-induced genes[17,28] is dispensable for tolerance to both acute and anticipated stress[41–43].

Here we dissect HSR regulation by CRISPR interference and activation (CRISPRi/a) systems that employ a catalytically inactive Cas9 nuclease fused to transcriptional repression or activation domains[44,45]. Only a handful of studies reported the use of these technologies for functional genomic screens in *S. cerevisiae*, mostly assaying effects on growth[46–50]. We employ inducible CRISPRi/a[46,51] to modulate the abundance of protein kinases (PKs) and transcription factors (TFs), key regulators of almost every cellular pathway and trait, and screen for gene functions in cellular fitness at

temperatures 23°C, 30°C and 38°C, HSR activity and thermotolerance. We discover a handful of genes

capable of tuning thermotolerance by altering chaperone expression, including principal regulators of

the HSR, the unfolded protein response (UPR), as well as the HOG and PKA pathways. We further find

that CRISPRi effect size is temperature-dependent and that diverse traits are differentially sensitive to

knock-down. Altogether, our study reveals the HSR as a complex programme, regulated by multiple

molecular pathways and coupled with diverse cellular mechanisms to confer a rapid and precisely

tuned adaptation to heat.

Results

CRISPRi/a efficiently modulate gene expression

We first validated the performance of the employed perturbation and reporter systems. To confirm

CRISPRi effects on growth, we repressed the essential *HSF1* gene. As expected, this resulted in

decreased growth rate, with varied effect size for three gRNAs differing in target sequence and distance

to the transcription start site (TSS) (Fig. 1a and b). Effects were specifically observed in the presence

of the gRNA-inducing compound anhydrotetracycline (ATc) (Supplementary Fig. S1).

We quantify HSR activity with a heat-responsive reporter based on the truncated promoter of the *SSA1*

HSP70 gene[52,53] driving expression of an ultra-fast maturing GFP[54]. We validated our CRISPRi/a

systems by targeting this promoter, achieving efficient knock-down and overexpression of Hsp70

protein, respectively (Fig. 1c). The ATc-induced CRISPRa strain had fivefold increased reporter signal

already before heat shock (t=0), as expected for strong activation (dark green curve in Fig. 1c).

Interestingly, heat exposure (t>0) resulted in ~30 min delayed *SSA1* expression compared to the non-

induced CRISPRa strain. The *SSA1* promoter was thus not immediately induced if Hsp70 protein levels

were already elevated, in line with its ability to inhibit Hsf1[20,25].

As a proof of concept for using the HSR reporter for functional genomics, we tested its responsiveness

to Hsf1. Repression of Hsf1 decreased Ssa1 protein levels in heat and non-stress conditions (Fig. 1d),

as expected from *SSA1* mRNA changes after Hsf1 depletion[28] and chromatin-immunoprecipitation

(ChIP) of the *SSA1* promoter together with Hsf1 protein[55]. The GFP-based reporter is selectable by

Fluorescence-Automated Cell Sorting (FACS) and CRISPRi effects were inherited over at least 20

generations (Fig. 1e), indicating excellent suitability for genetic screens.

Figure 1. Evaluation of CRISPRi/a effects and the HSR reporter. a, Growth curves for repression of
HSF1 with three different gRNAs. The optical density at 600 nm (OD600) of strains grown with ATc to
induce CRISPRi was measured over time. Lines denote linear fits. **b,** Generation times of different *HSF1*
CRISPRi strains (x-axis), measured in n=3 replicate well cultures. **c,** Normalized GFP signal of *SSA1*
CRISPRa (act) and CRISPRi (rep) strains over time. Cultures were grown at 30°C, calibrated at

OD600=0.3 and exposed to 40°C throughout the experiment. The y-axis denotes GFP intensity

normalized by OD600. Lines denote means and ribbons denote standard deviations of n=6 replicate

well cultures, respectively. The chromosomally inserted HSR reporter is depicted as inlay. **d,** Cellular

reporter signal of CRISPRi strains at 30°C and exposed to 42°C for 5h. Dots denote single cells imaged

by fluorescence microscopy. Horizontal lines mark medians. **e,** Cellular GFP intensity of CRISPRi

strains (merged data for TF and PK libraries) sorted for high or low cellular GFP intensity after heat

shock, regrown for more than 20 generations, and imaged by microscopy. The cellular GFP intensity

(x-axis) is shown for bins of size 10. Dashed lines denote medians.

Gene dosage effects on cellular fitness

We first characterised CRISPRi effects on fitness (Fig. 2a). We repressed sets of either 129 protein

kinases[56] or 161 transcription factors[57] with up to six gRNAs per gene (Supplementary Fig. S2). Out of

1573 gRNAs in both libraries, 271 were significantly depleted (two-fold depletion, FDR<0.05) after

two days of competitive growth at 30°C (Supplementary Fig. S3). Approximately 40% of gRNAs were

effective (Fig. 2b), based on the dropout of essential genes defined as non-viable deletions according

to the *Saccharomyces* Genome Database (SGD)[58]. CRISPRi efficiency depends on the GC content and

secondary structure of gRNAs (Supplementary Fig. S4), and the chromatin accessibility at the targeted

genomic locus (Supplementary Fig. S5) which supports and complements previous findings[46,47]. Based

on the distance between TSS and gRNA target locus, the optimal range is between TSS-150 to TSS+25

nucleotides, with minor variation between target strands (Supplementary Fig. S4b).

Cellular fitness decreased upon repression of 68 genes (Fig. 2c, Supplementary Tab. 1). These were

enriched for essential functions (34% compared to 12% in the background) with roles in ribosome

biogenesis, cell cycle and chromosome segregation (Supplementary Fig. S6). Out of the 34 targeted

essential genes, 31 had at least one effective gRNA, and 23 were depleted with two or more supporting

gRNAs (Supplementary Fig. S7). In general, fitness effects correlated with knock-out (KO) screens,

despite differences in assay conditions and readouts (Supplementary Fig. S8), and outperformed

heterozygous deletions in detecting gene essentiality (Supplementary Fig. S9). We measured novel

fitness-modulatory roles for eight open reading frames (ORFs) (*CAD1, FPK1, IKS1, NHP6A, RSC30, SCH9,*

TEA1, TPK2) and two ambiguous loci where multiple TSS were potentially targeted (*FUS3|PEP1*,

MMO1|PHD1).

To validate screen performance, we selected ten ORFs for further characterization; six known to affect

growth and four measured with new functions. We observed high correlation between screen fold

changes and individually determined growth rates (Spearman *r*=0.77; Supplementary Fig. S10), and

half-maximal OD intervals ($t_{OD0.5}$) which additionally report on lag time (Spearman *r*=-0.87; Fig. 2d).

This follow-up confirmed novel roles in fitness regulation for all four genes included (*FPK1, RSC30,*

TEA1, TPK2).

Figure 2. CRISPRi effects are reproducible, capture positive controls and novel fitness modulators. a, Experimental set-up for competitive growth screens. Briefly, a computationally designed oligonucleotide library was cloned into plasmids and transformed into yeast. CRISPRi induced (+ATc) and reference yeast cultures (-ATc) were then profiled for fitness, followed by plasmid DNA extraction and gRNA barcodes sequencing. **b,** Frequency of CRISPRi strain dropout from populations before selection (after 10 h pre-growth, see Methods) and after two days selection at 30°C. Fractions of gRNAs with log2FC<=-1 and FDR<0.05 are shown for total and essential genes (merged for PK and TF library data). Numbers denote displayed ratios. **c,** Depletion of essential (green) and

non-essential genes (blue) for fitness at 30°C. Histogram bins have size 0.2 and dashed lines denote

medians. Gene log2FCs represent mean gRNA log2FCs per gene. **d,** Comparison of screen gRNA log2FC

with monoculture log2-scale effects on $t_{OD0.5}$ (time until cultures reach half-maximal OD) which reports

on growth rate and lag time. Error bars denote standard deviations of n=3 well cultures (x-axis) and

n=2 replicates for the fitness at 30°C screen (y-axis). Spearman correlation is shown. Repressed genes

are labelled. If multiple gRNAs were used, these are included in labels with g[1-9]. Dashed grey lines

are intercepts marking a fold change of 0. The linear model fit was generated with the R ggplot2

function geom_smooth, using default parameters and method="lm". Hsf1 was not included since

maximum OD values are not meaningful for severe growth defects (see gRNA1 in Fig. 1a). **e,** CRISPRi

effects on cellular fitness at 30 versus 38°C. Dots denote generation-normalized gene log2FCs coloured

by difference and Spearman correlation is shown. Genes with heat-sensitive phenotypes were labelled

in green (PKs) and orange boxes (TFs). Dashed purple lines mark diagonals indicating difference

thresholds at -/+ 0.04. Dashed grey lines are intercepts marking a fold change of 0 and the diagonal.

Genetic requirements for growth at high temperature

Having validated our assay at 30°C, we further screened for fitness effects at 23 and 38°C, detecting 18

and 30 depleted genes, respectively (Supplementary Tab. 1). Effect size and statistical power were

reduced compared to the 30°C screen due to less generations of selection, considering that doubling

time was lowered by ~60% at 23°C and by ~30% at 38°C (Supplementary Fig. S11-12). Generation-

normalized fold changes correlated well between temperatures, demonstrating high reproducibility

not only for read counts (Supplementary Fig. S13) but also for CRISPRi effects (Supplementary Fig.

S14). We detected 15 genes causing heat sensitivity upon repression, derived from reduced fitness at

38 compared to 30°C (Fig. 2e), nine of which were known, such as Hsf1 and Swi4 which control HSR

transcription[59,60], Reb1 which enhances Hsf1 transactivation[61], Ume6 which promotes Msn2/4-

dependent transcription as part of the Rpd3L histone deacetylation complex[62], and Cst6 with a yet

unknown but predicted role in the HSR[63]. Quantification of mRNA and gRNA levels in Hsf1 and Ume6

CRISPRi strains over time and temperatures confirmed efficient repression in all conditions, showing

that heat sensitivity is not simply due to stronger repression at higher temperature (Supplementary

Fig. S15). Novel roles in heat sensitivity were detected for Cep3, Gcr1, Tor2 and Rsc30. Supporting this,

Gcr1 controls the expression of glycolysis genes[64] and Tor2 regulates cytoskeleton organization[65], both

important for thermal adaptation[3]. Rsc30 is part of the RSC complex which translocates from ORFs to

promoters in heat to facilitate nucleosome dissociation[66] and Hsf1-mediated transcription[61], akin to

the known hit Rsc3 (Fig. 2e).

Figure 3. Modulators of HSR activity. a, Schematic of the reporter-based HSR screen. After heat exposure, cells with extremely high and low GFP intensity were collected by FACS and identified through barcode sequencing. **b,** Comparison of gene log2 fold changes for cellular fitness at 38°C versus HSR with Spearman correlation. Dots denote genes (n=290), coloured by screen effects as indicated in figure legend. Selected HSR modulators are labelled. **c,** Cellular *SSA1pr-GFP* reporter intensity (y-axis) of individual CRISPRa strains cultured with or without ATc (x-axis) after 5h exposure to 42°C, and imaged with fluorescence microscopy. Dots denote single cells. Used gRNAs are Hog1_g1, Iks1_g5, Hsf1_g1. Two-sided Wilcoxon adjusted p-values are depicted for tests between samples.

Modulators of the heat shock response

Most transcripts induced during heat shock appear to not serve protective functions and it is therefore debated if they compensate for loss-of-function effects due to protein instability[41]. Combining published RNAseq time course with thermal proteome profiling data, we found that heat-induced transcripts encode proteins with higher than average thermal stability, while proteins with low stability are down-regulated (Supplementary Fig. S16). This suggests that the HSR is a purposeful programme to enhance heat resistance rather than overproducing proteins that go astray.

To identify components controlling the HSR pathway, we established a flow cytometry assay (Fig. 3a) based on the Hsp70 reporter (introduced in Fig. 1c). This allowed us to measure impacts on chaperone expression independent of fitness effects (Fig. 3b & Supplementary Fig. S11). Out of the 290 TFs and

PKs, we found twenty to decrease and seven to increase HSR activity upon repression, implying

functions in promoting and inhibiting the HSR, respectively (Supplementary Tab. 1). CRISPRi effects

on the Hsp70 promoter cannot be fully explained by previous screens with artificial promoters based

on either heat shock elements (HSEs) recognized by Hsf1 or stress response elements bound by

Msn2/4, using deletion mutants and a one hour heat shock at 37°C (Supplementary Fig. S17)[26]. Most

of our hits were thus not known to tune chaperone expression during the HSR, although we found that

individual roles were supported by studies that probed 35 or 68 gene deletions with an *HSP12-GFP*

reporter gene[67,68].

Hsf1 was the most potent HSR activator (Fig. 3b), in agreement with single cell microscopy results (Fig.

1d). Additionally, Msn2 and Msn4 were measured as strong HSR stimulators, further validating the

experimental setup (Fig. 3b). The two regulators of the environmental stress response are redundant

for growth and thermotolerance, as discovered by Martínez-Pastor et al. (1996)[69] and confirmed by

our study. In contrast, repression of either factor decreased Hsp70 expression which demonstrates

their non-redundant roles in stimulating transcription as part of the HSR[11]. Notably, the second

strongest HSR stimulator in the panel was Iks1, a putative kinase of unknown function which is

transcriptionally induced at 37°C[70] and during proteotoxic stress[71] (Fig. 3b). We measured further

genes that promote HSR activity, derived from decreased reporter signal upon repression, that encode

chromatin remodellers (Rsc30, Nhp6A), activators of the plasma membrane ATPase (Ptk2, Hrk1)

stress-related kinases (Mck1, Rim15, Yak1), and surprisingly also central components of the UPR

(Ire1), the high osmolarity glycerol (HOG) pathways (Hog1, Pbs2, Rck2) and the protein kinase A (PKA)

subunit Tpk2. Additionally, we determined found four kinases (Cdc28, Hrr25, Mps1, Sch9) and three

TFs (Rim101, Sok2, Ume6) with roles in alleviating HSR activity. We confirmed that Cdc28 counteracts

the HSR by FACS (Supplementary Fig. S18a) which further agrees with screen measurements for its

regulators (Supplementary Fig. S18b). All three HSR-antagonizing TFs act as transcriptional

repressors[72–74], in line with their inhibiting roles.

Microscopy follow-ups confirmed screen results for the repression of Hog1, Iks1 and Ume6

(Supplementary Fig. 19). In addition, we show that CRISPRi screen phenotypes can be reversed using

CRISPRa strains for Hog1, Iks1 and Hsf1 by CRISPRa (Fig. 3e). HSR-stimulating kinases, such as Iks1,

Hog1, Rim15 and Yak1 potentially activate a potent TF. Rim15 and Yak1 phosphorylate both Hsf1 and

Msn2 upon glucose starvation[75–77] and our results suggest these roles also as part of the HSR. The

opposite phenotypes measured for Sch9 CRISPRi strains further agree with the role of Sch9 in

inhibiting Rim15[78], Yak1[79] and Hsf1 in starvation stress[80]. In line with our findings, Hog1 has recently

been reported to phosphorylate Hsf1 in osmostress[81]. Interestingly, the human Hog1 MAPK

orthologue p38 also phosphorylates and activates Hsf1 upon treatment with an Hsp90 inhibiting

compound[82].

To get insights into HSR-regulated processes, we determined PK interactors from phospho-

proteomics[83] and TF target genes from ChIP data[84] (Supplementary Fig. S20). Interactors of HSR-

modulating PKs were enriched for functions in mitogen-activated protein kinase (MAPK) signalling (p-

value=3.3e-05) and cell cycle regulation (p-value=1.8e-03) (Supplementary Fig. S20d). Target genes

of HSR-regulating TFs had roles in responses to heat and oxidative stress (p-value=7.2e-04), the fungal

cell wall (p-value=6.5e-03) and trehalose metabolism (p-value=4.3e-03) (Supplementary Fig. S20e).

This target-based analysis thus not only proved useful in recapitulating paramount mechanisms of

heat resistance that are remodelled as part of the HSR[18], but also implies that these processes are, at

least partially, controlled by the same TFs that regulate chaperone expression.

Modulators of thermotolerance

We were curious if genes adjusting the HSR pathway have stress-protective roles. We thus screened

for thermotolerance, the cells' ability to survive a sudden and lethal heat shock (Fig. 4a). Briefly, yeast

populations were selected at 50°C for 150 min (Supplementary Fig. S21) and recovered without

maintaining CRISPRi perturbations. Comparing sequencing barcodes before and after heat shock, we

identified twelve genes to promote and three to counteract thermotolerance, with minor or no impact

on fitness (Supplementary Fig. S22 & Supplementary Tab. 1). Most thermotolerance modulators also

physically interact with each other (14 of 15) holding the potential for cross-talk to fine-tune mutual

activities (Fig. 4b).

Decreased thermotolerance was observed for repression of three TFs that also had heat sensitive

fitness (Hsf1, Reb1, Cst6), and for PKs in stress signalling (Yak1, Rim15) and the HOG (Hog1, Pbs2),

UPR (Ire1) and PKA pathways (Tpk2) (Supplementary Fig. S22). Increased thermotolerance was

measured for CRISPRi strains of Sch9, a PK controlled by target of rapamycin (TOR) signalling[85], the

Sln1 sensor kinase of the HOG pathway[86], and the uncharacterized kinase Ypl150W[58]. Only three of the

measured thermotolerance effects were known according to SGD, including the enhanced heat

resistance of *sch9Δ* and the reduced tolerance of *pbs2Δ* and *rim15Δ* strains[43]. Strikingly, dilution spot

plating of individual CRISPRi strains confirmed thermotolerance effects of Hsf1 and Hog1 (Fig. 4e), as

well as for Tpk2, Pbs2, Cst6 and Rsc30 (Supplementary Fig. S23). Repression of the *TOR1/ARP3* locus

was used as negative control that strongly decreased growth, but not thermotolerance. Interestingly,

CRISPRa strains of Hsf1 had wildtype thermotolerance, suggesting that increased Hsf1 abundance may

not alter heat resistance, and thermotolerance was decreased for activation of Pbs2 (Supplementary

Fig. S23).

Genes found to modulate both thermotolerance and the HSR reporter modulators encode for Hsf1,

stress-related kinases (Sch9, Rim15) and components of the UPR (Ire1) and HOG pathways (Hog1,

Pbs2) (Fig. 4c & d). The Yak1 and Ypl150W PKs (Supplementary Fig. S24a & b), as well as the Tpk1/2/3

PKA subunits (Supplementary Fig. S25) were likely part of this overlap, although not fulfilling the strict

significance requirements.

Figure 4. Shared regulation of thermotolerance and HSR. a, Schematic of the thermotolerance screen. **b,** Protein-protein interaction network of thermotolerance modulators based on STRING[108]. Line thickness indicates interaction confidence, and node size the screen p-value. Node colour denotes increasing (blue) or decreasing (red) roles on thermotolerance based on screen results with a light to dark gradient indicating low to strong effect size. **c,** Gene fold change comparison between the thermotolerance and HSR screens. Significant modulators are labelled in orange (TFs) or green boxes (PKs). Genes in grey boxes encode PKs that missed the stringent significance thresholds. Dashed lines are intercepts marking a fold change of 0 and the diagonal. **d,** Venn diagram showing overlap of genes altering thermotolerance and Hsp70 expression. **e,** Dilution spot plating of individual CRISPRi strains grown with and without ATc at 30°C, or exposed to 50°C for 150 min.

CRISPRi effect magnitude depends on temperature and phenotype

We clustered effects across phenotypes to group genes by function (Fig. 5a), such as modulating fitness

with moderate (clusters II, V, VI) or severe impact (VII & VIII) or adjusting the HSR pathway (II, III, IV,

VII). Comparing CRISPRi magnitude across temperature, we found that effect size increased with

temperature (Fig. 5b; Supplementary Fig. S26). Notably, the gRNA sequence GC content and secondary

structure affects strain fitness with temperature-dependent contributions (Supplementary Fig. S27).

Finally, we observed that phenotypes vary in their sensitivity to knock-down, as shown for Hsf1 (Fig.

5c). All six *HSF1*-targeting gRNAs severely decreased chaperone expression, implying that Hsf1

abundance is diminished strong enough to indirectly affect the assayed reporter. However, only the

three most potent gRNAs affected fitness at 30°C, despite the essentiality of *HSF1* (Fig. 5c). Similar

effect size gradients were observed for *UME6* and *RSC30* with only a few strong gRNAs (derived from

HSR reporter impact) altering high temperature growth (Supplementary Fig. S28). Repression effects,

even if they translate to protein level changes, do therefore not necessarily impact a robust

downstream trait.

Figure 5. Knock-down sensitivity depends on growth and phenotype. **a,** Comparison of repression

effects between screens as Z-score gene log2FCs (n=290). The hierarchical tree of rows is arbitrarily

cut to form 8 clusters to group genes with shared effects, such as decreasing or increasing HSR activity
(II or III, respectively), decreasing thermotolerance and HSR activity (IV), or severe fitness defects
(VIII) upon repression. **b,** Temperature-dependent CRISPRi effects. Generation-normalized gene
log2FCs of fitness screens at different temperatures, grouped for essential (n=34) and non-essential
genes (n=255). c, Log2FCs (y-axis) of six HSF1-targeting gRNAs (x-axis) across screens. Dashed blue
lines denote log2FC cut-offs 1 and -1.

Discussion

We used CRISPRi screens to measure gene functions in temperature-associated growth, the heat shock

response (HSR) pathway and thermotolerance. The integration of these diverse phenotypic readouts

allowed us to reveal novel regulators of chaperone expression and link these to heat adaptation,

discovering unknown functions of cell cycle regulators and the HOG, UPR and PKA pathways.

Figure 6. The HSR is regulated by multiple molecular pathways that sense diverse
perturbations in different cellular compartments. Schematic overview of pathways controlling
HSR activity based on screen-derived roles illustrated as stimulating (green arrows) or inhibiting (red
stop indicators), and complemented by known processes from literature (in grey). Heat causes
proteotoxic stress which is counteracted by cellular mechanisms, such as chaperone-mediated re-
folding of misfolded proteins, and their degradation. Signalling pathway components of the HOG, UPR

and PKA pathways modulate chaperone expression and thermotolerance. In addition, the roles of

Cdc28 in promoting cell cycle progression and inhibiting HSR activity are shown. Further genetic roles

measured in screens are omitted for clarity.

Our screens allowed us to pinpoint core regulatory components of the HSR capable of tuning

thermotolerance by altering chaperone expression. We found central MAP kinases of the HOG pathway

to stimulate thermotolerance and chaperone expression, thus explaining Hog1 activation in heat[38–40]

with a role in protein homeostasis (Fig. 6). We showed that the UPR-triggering sensor kinase Ire1[87]

increases the expression of a cytosolic Hsp70 (Fig. 4c, Supplementary Fig. S24). This suggests a role

for the UPR in stimulating the HSR pathway, in line with the lowered activity of an artificial HSE

reporter in the heat-exposed *ire1Δ* strain[88]. However, cytosolic HSPs were not previously known as

UPR targets[89]. While PKA signalling is thought to counteract the HSR through inhibiting Msn2/4[90–92]

and by indirectly repressing Hsf1 presumably via Yak1 and Rim15[36,76,77], we show that only the Tpk1

and Tpk3 subunits inhibit, while Tpk2 enhances Hsp70 production and thermotolerance

(Supplementary Fig. S25a & b). This agrees with the increased Hsp12 expression and oxidative stress

survival of *tpk1Δ* and *tpk3Δ* mutants, while both are decreased in *tpk2Δ*[67]. Additionally, differential

Tpk1/2/3 roles are expected from their distinct physical interactors (Supplementary Fig. S25c) and

phosphorylation patterns[40].

Notably, the strongest HSR-antagonizing PK (Cdc28) is the only cyclin-dependent kinase (CDK) both

necessary and sufficient to drive the cell cycle in *S. cerevisiae*[93]. While cell cycle arrest is a known

consequence of the HSR[94], we show that key cell cycle regulators also signal back to the stress response

in a mutually inhibitory manner (Supplementary Fig. S18; Fig. 6). During stress recovery, CDK

promotes re-entry into mitosis and could simultaneously shut down the remains of a stress response

to quickly resume proliferation. Supporting this, the HSP abundance is frequently regulated depending

on cell cycle stage[95,96]. Additionally, given that Hog1 blocks mitosis by suppressing Cdc28-activators[97]

and its interacting cyclins in hyperosmotic conditions[98], it presumably contributes to growth arrest

also as part of the HSR.

We found repression effects to increase with temperature (Fig. 5b; Supplementary Fig. S26) which is

likely due to both technical (CRISPRi system performance) and biological factors (sensitivity to knock-

down caused by changes in transcription, translation and metabolism). We employed the

Streptococcus pyogenes Cas9 fused to the human Mxi1 repressor which both evolved to operate around

37°C. Accordingly, Cas9 has higher *in vitro* DNA cleavage efficiency at 37°C compared to 22°C[99].

Temperature may also affect gRNA hybridization, as proposed for microRNAs[100], in line with

temperature-dependent contributions of gRNA GC content and secondary structure (Supplementary

Fig. S27). This needs to be considered when using CRISPRi at extreme temperatures in yeast and other

microorganisms. In our study, we paid special attention to potential biases in heat sensitivity, defined

by contrasting fitness effects at 38 vs. 30°C. However, differences in strain dropout at these
temperatures are minor compared to 23°C. The comparable mRNA depletion across temperatures in
heat sensitive *HSF1* and *UME6* CRISPRi strains (Supplementary Fig. S15) and strong support by
deletion strains reassured of measuring biological functions.

While repression is complementary to knock-out, we also measured novel fitness-modulatory roles
(~11% of 30°C fitness screen hits) which are unlikely to result from off-targets due to their support
by two or more gRNAs. Additionally, effects were reproduced in screens at varied temperatures
(Supplementary Fig. S14). We hypothesize that the sudden mRNA depletion can yield a stressed cell
state while deletion mutants may trigger bypassing mechanisms when outgrown over several
generations, such as acquiring adaptive mutations[101]. In CRISPR-KO screens, selection is usually
performed immediately after introducing mutations and thus yield effects that are more severe[102] or
comparable to CRISPRi in human cells[103], depending on assay setup. In future studies, it will be
interesting to explore differential effects of knock-out and knock-down over time to explain the
strengths and drawbacks of both techniques. Key advantages of the presented CRISPRi/a platforms
are the inducible, reversible, and tunable modulation of transcription without altering genomic
sequence and the ability to probe essential ORFs as demonstrated for *HSF1* (Fig. 1d & Fig. 5c).

A limitation of our HSR screens is given by the specificity of our reporter which monitors chaperone
expression as a characteristic aspect of the HSR and may not report on other aspects. However,
functional insights in PK phosphorylation targets and TF target genes illustrated the relevance of our
screen hits for a variety of HSR-connected processes beyond HSP expression. Gene functions
discovered by our screens can be directly applied in yeast biotechnology to generate strains with
enhanced heat resistance or high temperature growth, i.e. to facilitate ethanol production[104,105]. If HSR-
modulatory roles are conserved in human cells, they can be evaluated as therapeutic targets for disease
treatment[7,10]. We anticipate that the established screen workflows and tools provide a basis to study
diverse other reporters and molecular pathways, potentially expanding them to genome-wide scale
and multiplexed application to infer genetic interactions and pathway connectivity.

Material and Methods

Chemicals, oligonucleotides, plasmids and strains
All chemical compounds, oligonucleotides, plasmids, and bacterial or yeast strains are listed in
Additional File A1.

Plasmid and strain construction
The Tet-inducible dCas9-MxiI and dCas9-nGal4-VP64 plasmids are available on AddGene (#73796 and
#71128). Chemically synthesized gRNA oligonucleotide libraries were purchased from CustomArray,

Inc. (GenScript) amplified by PCR and integrated into the NotI site of the pRS416 dCas9-Mxi1 plasmid
via Gibson Assembly[106] with 30bp homology regions, or ligation with T4 DNA Ligase. Single gRNA
oligos and primers were purchased from Sigma Aldrich, and cloned into the NotI site of plasmids.
Plasmids were transformed into *E. coli* NEB10beta chemo- or electrocompetent cells (New England
Biolabs). Plasmid extraction from *E. coli* cultures was performed with the QIAprep Spin Miniprep kit
(Qiagen) and was combined with FastPrep (MP Biomedicals) to break the cell wall of *S. cerevisiae* using
~50µl volume of autoclaved glass beads (Sigma Aldrich) per cell pellet. FastPrep was run three-times
at 5,5 m/s for 20s with 1min pausing. PCR was done using Phusion high fidelity polymerase (Thermo
Fisher Scientific). Quality control of isolated DNA was performed with NanoDrop1000 (Thermo Fisher
Scientific), Qubit spectro-fluorometer (Invitrogen) and High Sensitivity DNA Bioanalyzer chips
(Agilent). For confirmation of sequence identity, plasmids and PCR products were submitted for
Sanger sequencing to Eurofins Genomics. Template plasmids of PCR reactions were digested with DpnI
for removal of bacterial DNA, and products were then used for Gibson Assembly or ligation. For library
generation, multiple *E. coli* transformations were performed in parallel, pooled, and colonies of several
selection plates were scraped together in a dense culture with LB-Ampicillin to ensure >20x coverage
of libraries. Chemical transformation of *S. cerevisiae* was done as described previously[107].
Transformants were selected on synthetic complete uracil-dropout media (SC-Ura) agar plates. Single
colonies were picked, confirmed by colony PCR and cultured for individual strain experiments.
Transformations were sequence-verified by Sanger sequencing. Cell libraries were generated by
washing off transformant colonies when reaching small size ~36h after plating. For long-term storage
at -80°C, 25% glycerol stocks were prepared.
421
Plate reader growth and fluorescence assays
Sequence-verified strains were cultured in 96 well round bottom plates (Thermo Fisher Scientific)
filled with 100 µL in SC-Ura dropout media with or without 250 ng/ml ATc. Yeast cultures were
inoculated with OD600=0.005 for growth measurements or OD600=0.3 for fluorescence
measurements (GFP channel: 488 nm excitation and 512 nm emission wavelength) in 15 min intervals
with Genios (Tecan) or Synergy HTX (Biotek) plate readers according to the manufacturer's
instructions. The maximum growth rate and the time until half-maximum OD600 were determined by
fitting a linear model and calculating its slope using the cellGrowth R package (V. 1.30.0)[108].
430
RNA extraction from yeast cells and qRT-PCR
Exponential phase yeast cultures were diluted to OD600=0.3 and cultured with 250 ng/ml ATc or
without ATc for either 4 or 6 hours at temperatures as indicated. Cells were collected using a vacuum
filter device and instantly frozen in liquid nitrogen. Cells were then resuspended in Trizol (Invitrogen)
and RNA was extracted using the Quick RNA Kit (Zymo Research) according to manufacturer's
instructions. RNA was reverse-transcribed to cDNA using SuperScript III (Invitrogen) with RNasin
(Promega, Life Technologies), Oligo(dT)18 (Thermo Fisher Scientific) and a reverse primer specific for

the 3' region of guide RNAs (see Additional File A1). This cDNA was diluted 1:10 or 1:20 and then used

for SYBR Green quantitative reverse transcription PCR (qRT-PCR) using PowerUp SYBR Green PCR

Master Mix (Thermo Fisher Scientific) and the Applied Bio-Systems QuantStudio 6 Flex Real-Time PCR

System (Thermo Fisher Scientific). Analysis was performed with R code as follows: The log2 fold

change between +ATc samples and -ATc reference samples was computed as the negative delta delta

Ct (-ddCt) with ddCt = ((average transcript Ct) - (average *ACT1* house-keeping control Ct) of +ATc

condition) – ((average transcript Ct) - (average *ACT1* house-keeping control Ct) of -ATc reference

condition). Average Ct values were calculated from triplicates. Primers were designed to yield

products of 75-130 nucleotides and are listed in Additional File A1.

Microscopy and image analysis

To quantify cellular reporter gene expression, cells were imaged with the Zeiss CellObserver

microscope (Carl Zeiss AG) during cultivation at 30°C and after 5h exposure to 42°C. Images were

acquired with the ZEN Black software, and analysis was performed with KNIME to quantify cellular

GFP signals[109] and R[110] for data visualization.

Choice of HSR reporter

In search of a suitable reporter for HSR activity, we prioritized genes by their stress-related

expression[11, 111]. We evaluated 44 Green Fluorescent Protein (GFP) tag strains[112] on fluorescence

induction after heat shock and found the *SSA1-GFP* fusion strain to yield the highest and most stable

reporter signal in this panel (Supplementary Fig. S29). To quantify *in vivo* HSR activity, and for all

screens presented, we employed a diploid BY4743 strain[113] harbouring a chromosomally integrated

reporter, consisting of the highly heat-responsive *SSA1* promoter with a Δ-280 bp truncation[52] which

controls expression of a fast-maturing GFP[54]. This reporter was chosen over artificial TF-specific

promoters[26,114] to measure effects of various transcription factors that affect Hsp70 expression.

Design of gRNA libraries

Guide RNA oligonucleotides were designed to target 161 TFs[57] (retrieved from

yetfasco.ccbr.utoronto.ca on 16/04/2014 using DNA-binding=1 and dubious=false parameters) and

129 PKs[56] (retrieved from yeastkinome.org on 16/04/2014). Libraries consist of 885 and 668 gRNAs,

respectively. Each gene was covered by up to six different gRNAs to minimize off-target effect calling

(Supplementary Fig. S2a). Guide RNAs were designed considering the distance of their midpoint (of

the 20nt target sequence) to the respective TSS[115] and nucleosome occupancy[116]. Blast

(blast.ncbi.nlm.nih.gov/Blast.cgi) and ECRISP (version 5.4)[117] were used to check potential off-target

binding sites in the yeast genome, allowing for two mismatches at most. The gRNA design pipeline was

published as part of Smith et al. (2016) and is available at lp2.github.io/yeast-crispri/. Potentially

regulated TSS in close proximity to the intentional target TSS (Supplementary Fig. S2b) are included

in the gene name. Specifically, if two or more gRNAs designed for "gene1" potentially targeted the TSS

of another "gene2" within 150 nt distance[47], the target locus is annotated as "gene1|gene2".

Screens

Plasmid libraries targeting sets of either TFs or PKs were transformed and profiled individually.

Screens were performed with 30 ml bulk yeast populations in 150 ml flasks. Populations were pre-

grown at 30°C with or without 250 ng/ml ATc for 10 h (~3 generations) before selection to enable

acquisition of CRISPRi-mediated changes on protein and phenotype level. This pre-growth had minor

effects on strain composition of populations (Supplementary Fig. S3) so that almost every gRNA

barcode in the design (>98%) was probed.

Competitive growth screens

For competitive growth selection, pre-grown cultures were diluted to OD600=0.005 and grown over

1.5–2 days at temperatures 23, 30 or 38°C. Fitness screens were performed with two replicates. Due

to anhydrotetracycline instability at high temperature, we confirmed that the compound maintains its

biological activity in the used concentration over at least 3 days at 38°C (Supplementary Fig. S30).

HSR screens and FACS

For HSR screens, pre-grown cultures were diluted to OD600=0.3, exposed to 42°C for 5 h to induce the

SSA1pr-GFP reporter and sorted with flow cytometry. Sorting was performed immediately after heat

shock to measure effects during the stress as opposed to recovery. 250.000 – 500.000 cells within the

top and bottom 5% of cellular GFP reporter intensity were collected by FACS. Gating was used to select

cells representing the bulk population in forward and sideward scattering, and to exclude dividing

cells (Supplementary Fig. S31). Sorted cells were recovered for ~5 generations without CRISPRi

induction and accounting for growth arrest. HSR screens were performed in three replicates. Flow

cytometry was performed using a MoFlo cell sorter (Beckman Coulter Inc.), equipped with a 70 μm

nozzle. A Sabre argon ion laser (Coherent Inc.), tuned to 488nm (200mW) was used as primary laser.

Laser illumination, optical configuration and sorting parameters were optimized with Flow-Check

fluorospheres (Beckman Coulter Inc.). Single cells were measured and sorted in purify one-drop mode.

Data was acquired with Moflo Summit and analysed with R code[110].

Thermotolerance screens

For thermotolerance selection, cultures were diluted to OD600=0.3, exposed to 50°C for 150 min and

recovered for ~7 generations considering growth arrest and rate during recovery (Supplementary Fig.

S21). Two samples did not pass quality control and were excluded from analysis (PK after heat shock

+ATc Rep2 & TF after heat shock -ATc Rep2 in Supplementary Fig. S13). Thermotolerance screens

were performed in two replicates for the TF and four replicates for the PK library.

 Next generation sequencing

QuBit (Thermo Fisher Scientific) was used to quantify extracted plasmid DNA. To amplify gRNA
barcodes, PCR was performed with ~5 ng plasmid DNA as template and primers that add inline
barcodes and Illumina P5 and P7 adapters (listed in Additional file A1). PCR products of all samples
were run on 1% Agarose gel with SYBR Safe (Invitrogen) to control DNA amount, size and purity. Gel
bands of PCR products were excised and DNA purified with the MinElute kit (Qiagen). After DNA
quantification by QuBit, equal amounts of PCR products were pooled. This sequencing library was size-
selected with an eGel (Thermo Fisher Scientific) and controlled for purity on a DNA-Bioanalyzer high
sensitivity chip (Agilent). Illumina sequencing was performed in paired-end and 75-100 base pairs
read length on NextSeq500 machines with 15% PhiX spike-in.

Sequencing data analysis
Sequencing data was demultiplexed with Jemultiplexer. Base calling quality was controlled with
FastQC. Reads were trimmed and aligned to a reference FASTA file with DNA barcodes using the
Burrows-Wheeler algorithm to compute read counts. Computational analysis was performed with the
edgeR R package[118] although other count-based packages can alternatively be used. Fold changes of
fitness screens were calculated as contrasts of read counts between +ATc and -ATc populations that
both underwent competitive growth selection (Fitness effect). Fold changes of the thermotolerance
screen are ratios between +ATc samples after versus before heat shock (thermotolerance effect), and
for the HSR screen denote ratios between +ATc samples sorted for high versus low cellular reporter
intensity (HSR effect). For all screens, we provide sample correlations of read counts (Supplementary
Fig. S13), as well as for the computed gRNA log2FCs (Supplementary Fig. S32) and gene log2FCs
(Supplementary Fig. S33). Gene log2FCs were computed as the mean log2FC of gRNAs per gene and
also be calculated without prior calculation of gRNA log2FCs directly from the geometric mean of gRNA
reads per gene. A single analysis workflow thus enables computation of log2FCs and adjusted p-
values/FDRs for genes and individual gRNAs (Supplementary Fig. S34). We benchmarked approaches
to calculate gene scores. We report gene scores as mean log2FC of all gRNAs per gene since it
performed better or as well as other measures, including median and rank-based scores (see
Supplementary Fig. S35 for correlation plots and Supplementary Fig. S36 for receiver operating
characteristics and precision-recall curves) and has higher robustness against noise and off-targets.
Significant gene functions are supported by a gene log2FC with FDR<0.05 and at least two gRNAs with
an absolute log2FC>=1 and FDR<0.05. Fold changes between non-induced samples (-ATc) are helpful
to determine the background variation without CRISPR-based perturbation for each screen
(Supplementary Fig. S37).

Gene ontology (GO) enrichments and target gene analysis
GO enrichment analyses were performed using the gProfiler[119] and gProfiler2 R packages[120].
Significant genes are queried with all target genes of the library as a statistical background. Major

cellular processes and functions are reported as enriched with adjusted Benjamini Hochberg FDRs
specified. TF target genes were determined based on Chromatin Immuno-Precipitation on chip (ChIP-
chip) data[84]. Target genes bound by at least two TFs identified as significant modulators were used for
GO enrichment, using the *S. cerevisiae* genome as background. Phosphorylation targets of protein
kinases were determined using the phosphogrid 2.0 database[83]. Physical interactors of significant PKs
were used for enrichment analysis with a background consisting of all identified *S. cerevisiae* protein
kinase targets.
558
Dilution spot plating thermotolerance assays
Individual CRISPRi/a strains were cultured in SC-Ura with or without 250 ng/ml ATc for 1 day, diluted
to OD600=0.3 and exposed to a 50°C in a table incubator (Eppendorf AG) for either 150 min or 90 min
as indicated. A dilution series was prepared in SC-Ura and 10 µl of each dilution was plated on SC-Ura
agar plates for the recovery of cells that survived the treatment without CRISPRi induction. After 2-3
days incubation at 30°C, photographs of plates were taken and colonies counted.
565
Computational preparation and visualization
Figures were prepared using Adobe Illustrator 2019. Data was processed in R (V. 3.4.1)[110] with the
tidyverse[121] and dplyr[122] packages, and plots were generated with the LSD (V. 3.0) [123], ggplot2 (V.
3.1.0)[124] and ggally (V. 1.3)[125] packages. Minimum free energies of RNA secondary structure were
computed using the ViennaRNA package (V. 2.0)[126]. Networks were generated with Gephi[127], using
protein-protein interaction data from the STRING database[128].
572
Statistics
For screens, multiple testing adjusted p-values (Benjamini Hochberg FDRs) were calculated with
standard edgeR functions[118] as described. For microscopy data comparisons between sample
populations, a two-sided Wilcoxon adjusted p-value was computed using the ggpubr (V. 0.3.0) R
package[129]. Boxplots are shown with a middle line corresponding to the median, and the lower and
upper hinges denoting the first and third quartiles, respectively. For all experiments with multiple data
points, these represent distinct samples and not repeated measurements.
580
581 *Code availability*
582 The KNIME image analysis workflow is available on request. The R code for screen analysis can be
583 downloaded from https://github.com/IAmTheMatrix/CRISPRi_Screen_Analysis/.
584
585 *Data availability*
586 Demultiplexed Illumina sequencing data has been uploaded to Gene Expression Omnibus, available
587 through GSE155455. The raw read counts and computed fold changes of gRNA barcodes and genes are

588 provided as Additional Files A2, A3 and A4, respectively. Source data underlying figures is provided in
589 Additional File A5.
590

591 **Supplementary Information**

592 Supplementary Figures S1-S37 and Supplementary Table 1 are provided as two pdf files. Additional
593 Files A1-A5 are xlsx files listing reagents, strains, DNA sequences, providing raw and processed
594 sequencing data and source data to figures.
595

Acknowledgements

The authors are grateful to Tanja Specht and Marko Kaksonen (University of Geneva) for sharing yeast
GFP collection strains, and to Marc Sherman and Barak Cohen (Washington University, St. Louis) for
kindly providing the SSA1pr-GFP reporter strain. The authors thank Mart Loog (University of Tartu),
Timo Mühlhaus and Michael Schroda (University of Kaiserslautern) for productive discussions, Lin Gen
and Aaron Brooks for computational support, and the EMBL Advanced Light Microscopy, Genomics
and Flow Cytometry facilities for their excellent support, especially Diana Ordonez, Malte Paulsen and
Vladimir Benes. This work was supported by an Advanced Investigator grant from the European
Research Council (ERC) under the European Union's Horizon 2020 research and innovation
programme (AdG-742804 to L.M.S.) by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation; project STE 1422/4-1 to L.M.S.), by Estonian Research Council (IUT 34-4 to L.P.), a Marie
Curie International Outgoing Fellowship to L.P., and Wellcome (grant number 206194 supporting L.P.).
C.J. was supported by an Interdisciplinary Sciences Fellowship of the Joachim Herz Foundation.

Competing Interests

The authors declare no competing interests.
612

613 **Contributions**

614 C.J. conceived the study, designed and performed experiments, analysed data and wrote the
615 manuscript. A.J. assisted with thermotolerance screens. L.P. designed and J.D.S generated gRNA
616 libraries. A.J., J.D.S., L.P. and L.M.S. provided valuable advice and revised the manuscript. L.P. and L.M.S.
617 supervised the study.
618

References

1.   Ananthan, J., Goldberg, A. L. & Voellmy, R. Abnormal proteins serve as eukaryotic stress signals and
trigger the activation of heat shock genes. *Science (80-.).* **232**, 522–524 (1986).
2.   Hartl, F. U. Molecular chaperones in cellular protein folding. *Nature* vol. 381 571–580 (1996).
3.   Piper, P. W. Molecular events associated with acquisition of heat tolerance by the yeast
Saccharomyces cerevisiae. *FEMS Microbiol. Rev.* **11**, 339–355 (1993).

4.   Ritossa, F. A new puffing pattern induced by temperature shock and DNP in drosophila. *Experientia*
18, 571–573 (1962).

5.   Richter, K., Haslbeck, M. & Buchner, J. The Heat Shock Response: Life on the Verge of Death. *Mol. Cell*
40, 253–266 (2010).

6.   Hartl, F. U., Bracher, A. & Hayer-Hartl, M. Molecular chaperones in protein folding and proteostasis.
Nature **475**, 324–332 (2011).

7.   Kampinga, H. H. & Bergink, S. Heat shock proteins as potential targets for protective strategies in
neurodegeneration. *The Lancet Neurology* vol. 15 748–759 (2016).

8.   Maiti, P. & Manna, J. Dysregulation of Heat Shock Proteins in Neurodegenerative Diseases:
Restorative Roles of Small Molecules and Natural Compounds. in *Heat Shock Proteins in Neuroscience*
85–114 (Springer, 2019). doi:10.1007/978-3-030-24285-5_7.

9.   Ben-Zvi, A., Miller, E. A. & Morimoto, R. I. Collapse of proteostasis represents an early molecular
event in Caenorhabditis elegans aging. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 14914–14919 (2009).

10.   Dong, B., Jaeger, A. M. & Thiele, D. J. Inhibiting Heat Shock Factor 1 in Cancer: A Unique Therapeutic
Opportunity. *Trends in Pharmacological Sciences* vol. 40 986–1005 (2019).

11.   Gasch, A. P. *et al.* Genomic expression programs in the response of yeast cells to environmental
changes. *Mol. Biol. Cell* **11**, 4241–4257 (2000).

12.   Causton, H. C. *et al.* Remodeling of yeast genome expression in response to environmental changes.
Mol. Biol. Cell **12**, 323–337 (2001).

13.   Hahn, J.-S., Hu, Z., Thiele, D. J. & Iyer, V. R. Genome-Wide Analysis of the Biology of Stress Responses
through Heat Shock Transcription Factor. *Mol. Cell. Biol.* **24**, 5249–5256 (2004).

14.   Nagaraj, N. *et al.* System-wide perturbation analysis with nearly complete coverage of the yeast
proteome by single-shot ultra HPLC runs on a bench top orbitrap. *Mol. Cell. Proteomics* **11**, 1–11
(2012).

15.   Mahat, D. B., Salamanca, H. H., Duarte, F. M., Danko, C. G. & Lis, J. T. Mammalian Heat Shock Response
and Mechanisms Underlying Its Genome-wide Transcriptional Regulation. *Mol. Cell* **62**, 63–78
(2016).

16.   Määttä, T., Rettel, M., Helm, D., Stein, F. & Savitski, M. Aggregation and Disaggregation Features of
the Human Proteome. *bioRxiv* (2020) doi:10.1101/2020.02.05.931675.

17.   Mühlhofer, M. *et al.* The Heat Shock Response in Yeast Maintains Protein Homeostasis by
Chaperoning and Replenishing Proteins. *Cell Rep.* **29**, 4593-4607.e8 (2019).

18.   Verghese, J., Abrams, J., Wang, Y. & Morano, K. A. Biology of the Heat Shock Response and Protein
Chaperones: Budding Yeast (Saccharomyces cerevisiae) as a Model System. *Microbiol. Mol. Biol. Rev.*
76, 115–158 (2012).

19.   Morano, K. A., Grant, C. M. & Moye-Rowley, W. S. The response to heat shock and oxidative stress in
saccharomyces cerevisiae. *Genetics* **190**, 1157–1195 (2012).

20.   Abravaya, K., Myers, M. P., Murphy, S. P. & Morimoto, R. I. The human heat shock protein hsp70
interacts with HSF, the transcription factor that regulates heat shock gene expression. *Genes Dev.* **6**,
1153–1164 (1992).

21.   Baler, R., Welch, W. J. & Voellmy, R. Heat shock gene regulation by nascent polypeptides and

665   denatured proteins: hsp70 as a potential autoregulatory factor. *J. Cell Biol.* **117**, 1151–1159 (1992).

666 22. Wang, Y., Gibney, P. A., West, J. D. & Morano, K. A. The yeast Hsp70 Ssa1 is a sensor for activation of
667   the heat shock response by thiol-reactive compounds. *Mol. Biol. Cell* **23**, 3290–3298 (2012).

668 23. Krakowiak, J. *et al.* Hsf1 and Hsp70 constitute a two-component feedback loop that regulates the
669   yeast heat shock response. *Elife* **7**, 1–17 (2018).

670 24. Sorger, P. K. & Pelham, H. R. B. Yeast heat shock factor is an essential DNA-binding protein that
671   exhibits temperature-dependent phosphorylation. *Cell* **54**, 855–864 (1988).

672 25. Zheng, X. *et al.* Dynamic control of Hsf1 during heat shock by a chaperone switch and
673   phosphorylation. *Elife* **5**, (2016).

674 26. Brandman, O. *et al.* A ribosome-bound quality control complex triggers degradation of nascent
675   peptides and signals translation stress. *Cell* **151**, 1042–1054 (2012).

676 27. Raychaudhuri, S. *et al.* Interplay of acetyltransferase EP300 and the proteasome system in regulating
677   heat shock transcription factor 1. *Cell* **156**, 975–985 (2014).

678 28. Solís, E. J. *et al.* Defining the Essential Function of Yeast Hsf1 Reveals a Compact Transcriptional
679   Program for Maintaining Eukaryotic Proteostasis. *Mol. Cell* **63**, 60–71 (2016).

680 29. Veri, A. O., Robbins, N. & Cowen, L. E. Regulation of the heat shock transcription factor Hsf1 in fungi:
681   Implications for temperature-dependent virulence traits. *FEMS Yeast Research* vol. 18 (2018).

682 30. Saavedra, C., Tuug, K. S., Amberg, D. C., Hopper, A. K. & Cole, C. N. Regulation of mRNA export in
683   response to stress in Saccharomyces cerevisiae. *Genes Dev.* **10**, 1608–1620 (1996).

684 31. Castells-Roca, L. *et al.* Heat shock response in yeast involves changes in both transcription rates and
685   mRNA stabilities. *PLoS One* **6**, (2011).

686 32. Zid, B. M. & O'Shea, E. K. Promoter sequences direct cytoplasmic localization and translation of
687   mRNAs during starvation in yeast. *Nature* **514**, 117–121 (2014).

688 33. Ketela, T., Green, R. & Bussey, H. Saccharomyces cerevisiae Mid2p is a potential cell wall stress
689   sensor and upstream activator of the PKC1-MPK1 cell integrity pathway. *J. Bacteriol.* **181**, 3330–
690   3340 (1999).

691 34. Bermejo, C. *et al.* The sequential activation of the yeast HOG and SLT2 pathways is required for cell
692   survival to cell wall stress. *Mol. Biol. Cell* **19**, 1113–1124 (2008).

693 35. Bandhakavi, S. *et al.* Hsf1 activation inhibits rapamycin resistance and TOR signaling in yeast
694   revealed by combined proteomic and genetic analysis. *PLoS One* **3**, (2008).

695 36. Ferguson, S. B. *et al.* Protein kinase A regulates constitutive expression of small heat-shock genes in
696   an Msn2/4p-independent and Hsf1p-dependent manner in Saccharomyces cerevisiae. *Genetics* **169**,
697   1203–1214 (2005).

698 37. Zou, J., Guo, Y., Guettouche, T., Smith, D. F. & Voellmy, R. Repression of heat shock transcription factor
699   HSF1 activation by HSP90 (HSP90 complex) that forms a stress-sensitive complex with HSF1. *Cell*
700   **94**, 471–480 (1998).

701 38. Winkler, A. *et al.* Heat Stress Activates the Yeast High-Osmolarity Glycerol Mitogen-Activated
702   Protein Kinase Pathway , and Protein Tyrosine Phosphatases. *Society* **1**, 163–173 (2002).

703 39. Dunayevich, P. *et al.* Heat-stress triggers MAPK crosstalk to turn on the hyperosmotic response
704   pathway. *Sci. Rep.* **8**, 1–15 (2018).

40.   Kanshin, E., Kubiniok, P., Thattikota, Y., D'Amours, D. & Thibault, P. Phosphoproteome dynamics of
Saccharomyces cerevisiae under heat shock and cold stress. *Mol. Syst. Biol.* **11**, 813 (2015).

41.   Berry, D. B. & Gasch, A. P. Stress-activated genomic expression changes serve a preparative role for
impending stress in yeast. *Mol. Biol. Cell* **19**, 4580–4587 (2008).

42.   Gibney, P. A., Lu, C., Caudy, A. A., Hess, D. C. & Botstein, D. Yeast metabolic and signaling genes are
required for heat-shock survival and have little overlap with the heat-induced genes. *Proc. Natl.*
Acad. Sci. U. S. A. **110**, (2013).

43.   Jarolim, S. *et al.* Saccharomyces cerevisiae genes involved in survival of heat shock. *G3 Genes,*
Genomes, Genet. **3**, 2321–2333 (2013).

44.   Gilbert, L. A. *et al.* Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation. *Cell*
159, 647–661 (2014).

45.   Gilbert, L. A. *et al.* CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes.
Cell **154**, 442 (2013).

46.   Smith, J. D. *et al.* Quantitative CRISPR interference screens in yeast identify chemical-genetic
interactions and new rules for guide RNA design. *Genome Biol.* **17**, 1–16 (2016).

47.   Smith, J. D. *et al.* A method for high-throughput production of sequence-verified DNA libraries and
strain collections. *Mol. Syst. Biol.* **13**, 913 (2017).

48.   Jaffe, M. *et al.* Improved discovery of genetic interactions using CRISPRiSeq across multiple
environments. *Genome Res.* **29**, 668–681 (2019).

49.   McGlincy, N. J. *et al.* A genome-scale CRISPR interference guide library enables comprehensive
phenotypic profiling in yeast. *bioRxiv* 2020.03.11.988105 (2020) doi:10.1101/2020.03.11.988105.

50.   Momen-Roknabadi, A., Oikonomou, P. & Tavazoie, S. An inducible CRISPR-interference library for
genetic interrogation of Saccharomyces cerevisiae biology. *bioRxiv* 2020.03.05.978619 (2020)
doi:10.1101/2020.03.05.978619.

51.   Farzadfard, F., Perli, S. D. & Lu, T. K. Tunable and multifunctional eukaryotic transcription factors
based on CRISPR/Cas. *ACS Synth. Biol.* **2**, 604–613 (2013).

52.   Slater, M. R. & Craig, E. A. Transcriptional regulation of an hsp70 heat shock gene in the yeast
Saccharomyces cerevisiae. *Mol. Cell. Biol.* **7**, 1906–1916 (1987).

53.   Bush, G. L. & Meyer, D. I. The refolding activity of the yeast heat shock proteins Ssa1 and Ssa2 defines
their role in protein translocation. *J. Cell Biol.* **135**, 1229–1237 (1996).

54.   Sherman, M. S., Lorenz, K., Lanier, M. H. & Cohen, B. A. Cell-to-Cell Variability in the Propensity to
Transcribe Explains Correlated Fluctuations in Gene Expression. *Cell Syst.* **1**, 315–325 (2015).

55.   Jonge, W. J. *et al.* Molecular mechanisms that distinguish TFIID housekeeping from regulatable SAGA
promoters. *EMBO J.* **36**, 274–290 (2017).

56.   Breitkreutz, A. *et al.* A global protein kinase and phosphatase interaction network in yeast. *Science*
(80-.). **328**, 1043–1046 (2010).

57.   De Boer, C. G. & Hughes, T. R. YeTFaSCo: A database of evaluated yeast transcription factor sequence
specificities. *Nucleic Acids Res.* **40**, (2012).

58.   Cherry, J. M. *et al.* SGD: Saccharomyces genome database. *Nucleic Acids Res.* (1998)
doi:10.1093/nar/26.1.73.

59.   Hahn, J. S. & Thiele, D. J. Activation of the Saccharomyces cerevisiae Heat Shock Transcription Factor
under Glucose Starvation Conditions by Snf1 Protein Kinase. *J. Biol. Chem.* **279**, 5169–5176 (2004).

60.   Kim, K. Y. & Levin, D. E. Transcriptional reporters for genes activated by cell wall stress through a
non-catalytic mechanism involving Mpk1 and SBF. *Yeast* **27**, 541–548 (2010).

61.   Pincus, D. *et al.* Genetic and epigenetic determinants establish a continuum of Hsf1 occupancy and
activity across the yeast genome. *Mol. Biol. Cell* **29**, 3168–3182 (2018).

62.   Ruiz-Roig, C., Viéitez, C., Posas, F. & De Nadal, E. The Rpd3L HDAC complex is essential for the heat
stress response in yeast. *Mol. Microbiol.* **76**, 1049–1062 (2010).

63.   Wu, W. S. & Li, W. H. Identifying gene regulatory modules of heat shock response in yeast. *BMC*
Genomics **9**, 1–15 (2008).

64.   Clifton, D., Weinstock, S. B. & Fraenkel, D. G. Glycolysis mutants in Saccharomyces cerevisiae.
Genetics **88**, 1–11 (1978).

65.   Schmidt, A., Kunz, J. & Hall, M. N. TOR2 is required for organization of the actin cytoskeleton in yeast.
Proc. Natl. Acad. Sci. U. S. A. **93**, 13780–13785 (1996).

66.   Vinayachandran, V. *et al.* Widespread and precise reprogramming of yeast protein-genome
interactions in response to heat shock. *Genome Res.* **28**, 357–366 (2018).

67.   Sadeh, A., Movshovich, N., Volokh, M., Gheber, L. & Aharoni, A. Fine-tuning of the Msn2/4-mediated
yeast stress responses as revealed by systematic deletion of Msn2/4 partners. *Mol. Biol. Cell* **22**,
3127–3138 (2011).

68.   Gutin, J., Sadeh, A., Rahat, A., Aharoni, A. & Friedman, N. Condition-specific genetic interaction maps
reveal crosstalk between the cAMP / PKA and the HOG MAPK pathways in the activation of the
general stress response . *Mol. Syst. Biol.* **11**, 829 (2015).

69.   Martínez-Pastor, M. T. *et al.* The Saccharomyces cerevisiae zinc finger proteins Msn2p and Msn4p
are required for transcriptional induction through the stress response element (STRE). *EMBO J.* **15**,
2227–2235 (1996).

70.   Sakaki, K., Tashiro, K., Kuhara, S. & Mihara, K. Response of Genes Associated with Mitochondrial
Function to Mild Heat Stress in Yeast Saccharomyces cerevisiae. *J. Biochem.* **134**, 373–384 (2003).

71.   Trotter, E. W. *et al.* Misfolded proteins are competent to mediate a subset of the responses to heat
shock in Saccharomyces cerevisiae. *J. Biol. Chem.* **277**, 44817–44825 (2002).

72.   Lamb, T. M. & Mitchell, A. P. The Transcription Factor Rim101p Governs Ion Tolerance and Cell
Differentiation by Direct Repression of the Regulatory Genes NRG1 and SMP1 in Saccharomyces
cerevisiae. *Mol. Cell. Biol.* **23**, 677–686 (2003).

73.   Ward, M. P., Gimeno, C. J., Fink, G. R. & Garrett, S. SOK2 may regulate cyclic AMP-dependent protein
kinase-stimulated growth and pseudohyphal development by repressing transcription. *Mol. Cell.*
Biol. **15**, 6854–6863 (1995).

74.   Carrozza, M. J. *et al.* Stable incorporation of sequence specific repressors Ash1 and Ume6 into the
Rpd3L complex. *Biochim. Biophys. Acta - Gene Struct. Expr.* **1731**, 77–87 (2005).

75.   Yamamoto, N., Maeda, Y., Ikeda, A. & Sakurai, H. Regulation of thermotolerance by stress-induced
transcription factors in Saccharomyces cerevisiae. *Eukaryot. Cell* **7**, 783–790 (2008).

76.   Lee, P., Cho, B. R., Joo, H. S. & Hahn, J. S. Yeast Yak1 kinase, a bridge between PKA and stress-

responsive transcription factors, Hsf1 and Msn2/Msn4. *Mol. Microbiol.* **70**, 882–895 (2008).

77.   Lee, P. *et al.* Rim15-dependent activation of Hsf1 and Msn2/4 transcription factors by direct
phosphorylation in Saccharomyces cerevisiae. *FEBS Lett.* **587**, 3648–3655 (2013).

78.   Pedruzzi, I. *et al.* TOR and PKA Signaling Pathways Converge on the Protein Kinase Rim15 to Control
Entry into G0. *Mol. Cell* **12**, 1607–1613 (2003).

79.   Zhang, A. & Gao, W. Mechanisms of protein kinase Sch9 regulating Bcy1 in Saccharomyces cerevisiae.
FEMS Microbiol. Lett. **331**, 10–16 (2012).

80.   Mudholkar, K., Fitzke, E., Prinz, C., Mayer, M. P. & Rospert, S. The Hsp70 homolog Ssb affects
ribosome biogenesis via the TORC1-Sch9 signaling pathway. *Nat. Commun.* **8**, (2017).

81.   Janschitz, M. *et al.* Novel interconnections of HOG signaling revealed by combined use of two
proteomic software packages. *Cell Commun. Signal.* **17**, (2019).

82.   Dayalan Naidu, S. *et al.* Heat Shock Factor 1 Is a Substrate for p38 Mitogen-Activated Protein Kinases.
Mol. Cell. Biol. (2016) doi:10.1128/mcb.00292-16.

83.   Sadowski, I. *et al.* The PhosphoGRID Saccharomyces cerevisiae protein phosphorylation site
database: Version 2.0 update. *Database* **2013**, (2013).

84.   Gonçalves, E. *et al.* Systematic Analysis of Transcriptional and Post-transcriptional Regulation of
Metabolism in Yeast. *PLoS Comput. Biol.* **13**, 1–20 (2017).

85.   Birney, E. *et al.* Identification and analysis of functional elements in 1% of the human genome by the
ENCODE pilot project. *Nature* **447**, 799–816 (2007).

86.   Saito, H. & Posas, F. Response to hyperosmotic stress. *Genetics* **192**, 289–318 (2012).

87.   Cox, J. S., Shamu, C. E. & Walter, P. Transcriptional induction of genes encoding endoplasmic
reticulum resident proteins requires a transmembrane protein kinase. *Cell* **73**, 1197–1206 (1993).

88.   Weindling, E. & Bar-Nun, S. Sir2 links the unfolded protein response and the heat shock response in
a stress response network. *Biochem. Biophys. Res. Commun.* **457**, 473–478 (2015).

89.   Travers, K. J. *et al.* Functional and genomic analyses reveal an essential coordination between the
unfolded protein response and ER-associated degradation. *Cell* **101**, 249–258 (2000).

90.   Garreau, H. *et al.* Hyperphosphorylation of Msn2p and Msn4p in response to heat shock and the
diauxic shift is inhibited by cAMP in Saccharomyces cerevisiae. *Microbiology* **146**, 2113–2120
(2000).

91.   Görner, W. *et al.* Nuclear localization of the C2H2 zinc finger protein Msn2p is regulated by stress
and protein kinase A activity. *Genes Dev.* **12**, 586–597 (1998).

92.   Smith, A., Ward, M. P. & Garrett, S. Yeast PKA represses Msn2p/Msn4p-dependent gene expression
to regulate growth, stress response and glycogen accumulation. *EMBO J.* **17**, 3556–3564 (1998).

93.   Beach, D., Durkacz, B. & Nurse, P. Functionally homologous cell cycle control genes in budding and
fission yeast. *Nature* **300**, 706–709 (1982).

94.   Rowley, A., Johnston, G. C., Butler, B., Werner-Washburne, M. & Singer, R. A. Heat shock-mediated
cell cycle blockage and G1 cyclin expression in the yeast Saccharomyces cerevisiae. *Mol. Cell. Biol.*
13, 1034–1041 (1993).

95.   Kao, H. T., Capasso, O., Heintz, N. & Nevins, J. R. Cell cycle control of the human HSP70 gene:
implications for the role of a cellular E1A-like function. *Mol. Cell. Biol.* **5**, 628–633 (1985).

96.   Santos, A., Wernersson, R. & Jensen, L. J. Cyclebase 3.0: A multi-organism database on cell-cycle
826   regulation and phenotypes. *Nucleic Acids Res.* **43**, D1140–D1144 (2015).

827   97.   Escoté, X., Zapater, M., Clotet, J. & Posas, F. Hog1 mediates cell-cycle arrest in G1 phase by the dual
828   targeting of Sic1. *Nat. Cell Biol.* **6**, 997–1002 (2004).

829   98.   González-Novo, A. *et al.* Hog1 Targets Whi5 and Msa1 Transcription Factors To Downregulate Cyclin
830   Expression upon Stress. *Mol. Cell. Biol.* **35**, 1606–1618 (2015).

831   99.   LeBlanc, C. *et al.* Increased efficiency of targeted mutagenesis by CRISPR/Cas9 in plants using heat
832   stress. *Plant J.* **93**, 377–386 (2018).

833   100.   Carmel, I., Shomron, N. & Heifetz, Y. Does base-pairing strength play a role in microRNA repression?
834   *Rna* **18**, 1947–1956 (2012).

835   101.   Puddu, F. *et al.* Genome architecture and stability in the Saccharomyces cerevisiae knockout
836   collection. *Nature* **573**, 416–420 (2019).

837   102.   Evers, B. *et al.* CRISPR knockout screening outperforms shRNA and CRISPRi in identifying essential
838   genes. *Nat. Biotechnol.* **34**, 631–633 (2016).

839   103.   Sanson, K. R. *et al.* Optimized libraries for CRISPR-Cas9 genetic screens with multiple modalities.
840   *Nat. Commun.* **9**, (2018).

841   104.   Banat, I. M., Nigam, P., Singh, D., Marchant, R. & McHale, A. P. Review: Ethanol production at elevated
842   temperatures and alcohol concentrations: Part I - Yeasts in general. *World Journal of Microbiology*
843   *and Biotechnology* vol. 14 809–821 (1998).

844   105.   Caspeta, L. *et al.* Engineering high-gravity fermentations for ethanol production at elevated
845   temperature with Saccharomyces cerevisiae. *Biotechnol. Bioeng.* **116**, 2587–2597 (2019).

847   **Methods References**

848   106.   Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat.*
849   *Methods* **6**, 343–345 (2009).

850   107.   Gietz, R. D. & Schiestl, R. H. Quick and easy yeast transformation using the LiAc/SS carrier DNA/PEG
851   method. *Nat. Protoc.* **2**, 31–34 (2007).

852   108.   Gagneur J and Neudecker A, Gagneur, J. & Neudecker, A. cellGrowth: Fitting cell population growth
853   models. R package version 1.16.0. *R Packag. version* **1**, (2012).

854   109.   Berthold, M. R. *et al.* KNIME-the Konstanz information miner: version 2.0 and beyond. *AcM SIGKDD*
855   *Explor. Newsl.* **11**, 26–31 (2009).

856   110.   R Core Team. R 3.5.1. A Language and Environment for Statistical Computing. *R Foundation for*
857   *Statistical Computing* vol. 2 https://www.R-project.org (2018).

858   111.   Soste, M. *et al.* A sentinel protein assay for simultaneously quantifying cellular processes. *Nat.*
859   *Methods* **11**, 1045–1048 (2014).

860   112.   Huh, W. K. *et al.* Global analysis of protein localization in budding yeast. *Nature* **425**, 686–691
861   (2003).

862   113.   Brachmann, C. B. *et al.* Designer deletion strains derived from Saccharomyces cerevisiae S288C: A
863   useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast*

(1998) doi:10.1002/(SICI)1097-0061(19980130)14:2<115::AID-YEA204>3.0.CO;2-2.

114.   Sorger, P. K. Yeast heat shock factor contains separable transient and sustained response
transcriptional activators. *Cell* **62**, 793–805 (1990).

115.   Pelechano, V., Wei, W. & Steinmetz, L. M. Extensive transcriptional heterogeneity revealed by
isoform profiling. *Nature* **497**, 127–131 (2013).

116.   Schep, A. N. *et al.* Structured nucleosome fingerprints enable high-resolution mapping of chromatin
architecture within regulatory regions. *Genome Res.* **25**, 1757–1770 (2015).

117.   Heigwer, F., Kerr, G. & Boutros, M. E-CRISP: Fast CRISPR target site identification. *Nature Methods*
vol. 11 122–123 (2014).

118.   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: A Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2009).

119.   Reimand, J. *et al.* g:Profiler-a web server for functional interpretation of gene lists (2016 update).
Nucleic Acids Res. 44, W83–W89 (2016).

120.   Kolberg, L., Raudvere, U., Kuzmin, I., Vilo, J. & Peterson, H. gprofiler2 -- an R package for gene list
functional enrichment analysis and namespace conversion toolset g:Profiler. *F1000Research* **9**, 709
(2020).

121.   Wickham, H. tidyverse: Easily Install and Load the 'Tidyverse'. *Bioinformatics* (2011).

122.   R Core Team. Package 'dplyr' - A Grammar of Data Manipulation. *CRAN Repository* (2019).

123.   B. Schwalb, A. Tresch, and R. F. LSD: Lots of Superior Depictions. *Compr. R Arch. Netw.* **3**, (2011).

124.   Ginestet, C. ggplot2: Elegant Graphics for Data Analysis. *J. R. Stat. Soc. Ser. A (Statistics Soc.* **174**, 245–
246 (2011).

125.   Schloerke, B. *et al.* GGally: Extension to 'ggplot2'. R package version 2.0.0. https://CRAN.R-
project.org/package=GGally (2020).

126.   Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, (2011).

127.   Bastian, M., Heymann, S. & Jacomy, M. Gephi: An Open Source Software for Exploring and
Manipulating Networks. *Third Int. AAAI Conf. Weblogs Soc. Media* (2009).

128.   Snel, B., Lehmann, G., Bork, P. & Huynen, M. A. String: A web-server to retrieve and display the
repeatedly occurring neighbourhood of a gene. *Nucleic Acids Res.* **28**, 3442–3444 (2000).

129.   Kassambara, A. Package ' ggpubr '. *https://rpkgs.datanovia.com/ggpubr/ BugReports* (2020).

130.   Baryshnikova, A. *et al.* Quantitative analysis of fitness and genetic interactions in yeast on a genome
scale. *Nat. Methods* (2010).

131.   Breslow, D. K. *et al.* A comprehensive strategy enabling high-resolution functional analysis of the
yeast genome. *Nat. Methods* (2008).

132.   Deutschbauer, A. M. *et al.* Mechanisms of haploinsufficiency revealed by genome-wide profiling in
yeast. *Genetics* (2005).

133.   Gu, Z. *et al.* Role of duplicate genes in genetic robustness against null mutations. *Nature* (2003).

134.   Qian, W., Ma, D., Xiao, C., Wang, Z. & Zhang, J. The Genomic Landscape and Evolutionary Resolution
of Antagonistic Pleiotropy in Yeast. *Cell Rep.* (2012).

135.   Jarzab, A. *et al.* Meltome atlas—thermal proteome stability across the tree of life. *Nat. Methods* **17**,
495–503 (2020).

136.   Guarente, L. & Mason, T. Heme regulates transcription of the CYC1 gene of S. cerevisiae via an
905   upstream activation site. *Cell* 32, 1279–1286 (1983).
906   137.   Sorger, P. K. & Pelham, H. R. Purification and characterization of a heat-shock element binding
907   protein from yeast. *EMBO J.* **6**, 3035–3041 (1987).
908   138   Brachmann, C. B. *et al.* Designer deletion strains derived from Saccharomyces cerevisiae S288C: A
909   useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast*
910   (1998).
911   139.   Jimeno-González, S., Gómez-Herreros, F., Alepuz, P. M. & Chávez, S. A Gene-Specific Requirement for
912   FACT during Transcription Is Related to the Chromatin Organization of the Transcribed Region. *Mol.*
913   *Cell. Biol.* **26**, 8710–8721 (2006).