Compositional Variability and Mutation Spectra of Monophyletic SARS-CoV-2 Clades

Xufei Teng[1,2,3,#,a], Qianpeng Li[1,2,3,#,b], Zhao Li[1,2,3,#,c], Yuansheng Zhang[1,2,3,#,d], Guangyi Niu[1,2,3,e], Meng Zhang[4,f], Jingfa Xiao[1,2,3,g], Jun Yu[1,2,3,h,*], Zhang Zhang[1,2,3,I,*], Shuhui Song[1,2,3,j,*]

[1] *China National Center for Bioinformation, Beijing 100101, China*

[2] *National Genomics Data Center & CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China*

[3] *University of Chinese Academy of Sciences, Beijing 100049, China*

[4] *Cixi Institutes for Life Science Innovation, Cixi, Zhejiang 315999, China*

[#] These authors contributed equally

*Correspondence:
  songshh@big.ac.cn (Song S), zhangzhang@big.ac.cn (Zhang Z), junyu@big.ac.cn (Yu J).

Abstract

COVID-19 and its causative pathogen SARS-CoV-2 have rushed the world into a staggering pandemic in a few months and a global fight against both is still going on. Here, we describe an analysis procedure where genome composition and its variables are related, through the genetic code, to molecular mechanisms based on understanding of RNA replication and its feedback loop from mutation to viral proteome sequence fraternity including effective sites on replicase-transcriptase complex. Our analysis starts with primary sequence information and identity-based phylogeny based on 22,051 SARS-CoV-2 genome sequences and evaluation of sequence variation patterns as mutation spectrum and its 12 permutations among organized clades tailored to two key mechanisms: strand-biased and function-associated mutations. Our findings include: (1) The most dominant mutation is C-to-U permutation whose abundant second-codon-position counts alter amino acid composition toward higher molecular weight and lower hydrophobicity albeit assumed most slightly deleterious. (2) The second abundance group includes: three negative-strand mutations U-to-

C, A-to-G, G-to-A and a positive-strand mutation G-to-U generated through an identical

mechanism as C-to-U. (3) A clade-associated and biased mutation trend is found attributable

to elevated level of the negative-sense strand synthesis. (4) Within-clade permutation

variation is very informative for associating non-synonymous mutations and viral proteome

changes. These findings demand a bioinformatics platform where emerging mutations are

mapped on to mostly subtle but fast-adjusting viral proteomes and transcriptomes to provide

biological and clinical information after logical convergence for effective pharmaceutical and

diagnostic applications. Such thoughts and actions are in desperate need, especially in the

middle of the *War against COVID-19*.

KEYWORDS: SARS-CoV-2; Nucleotide composition; Mutation spectrum; Viral replication

44   Introduction

COVID-19, a novel pneumonia epidemic causing an outbreak first identified and reported in

Dec 2019 from China [1] and subsequently spread to other countries swiftly, has been posing

enormous professional, economic, and political challenges to global health services and

hazardous control systems. As of 12 June 2020, there have been 7,410,510 confirmed cases

and 418,294 deaths reported [2]. COVID-19 is of great contagious (even at incubation period)

and has lower mortality to our current understanding [3-5]. The novel betacoronavirus

identified through *de novo* sequencing from patients with COVID-19 is designated as

"Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2)" by International

Committee on Taxonomy of Viruses (ICTV) [1, 6, 7].

The recent threats from SARS-CoV-2, SARS-CoV, and MERS-CoV are different from

those of earlier human coronaviruses (CoVs), including alphacoronaviruses, such as hsa-

CoV-229E, hsa-CoV-NL63, hsa-CoV-OC43 and hsa-CoV-HKU1 [8-10], in at least two

aspects. First, the recent groups of betacoronaviruses appears to come more frequently in the

past two decades as compared to the early comers where new members may be discovered as

technology become more efficient and accurate [11]. The current SARS-CoV-2 is also

different from both SARS-CoV and MERS-CoV as its genome composition is most closely

for "living with mammals and humans", where a much lower G+C content has been evolved

and is closer to two other human-adapted CoVs, hsa-CoV-229E and hsa-CoV-OC43, than its

members of the recent group, although it shares higher sequence identities with the two new

CoVs, 80.12% and 60.06%, respectively [11]. Second, it has been infecting far larger

populations, as compared to the two recent outbreaks, with variable yet more complex

symptoms [12]. The causative factors of such an unprecedented disease potency remain to be

elucidated for the days and months to come [1, 3–7].

Genomes of coronaviruses mutate in a unique way where signatures of DNA pairing and

repairing mechanisms are absent completely, and instead, they possess an error-prone

synthesis of single-stranded full or partial genomic sequences by multi-component

membrane-associated enzymatic structure known as the replicase-transcriptase complexes

(RTCs) and double-membrane vesicles (DMVs) although they do have certain enzymatic

activity resembling repair mechanisms of cellular organisms, such as proofreading [13], and

other possible cellular mechanisms may also be involved, such as RNA editing as recently

proposed [14,15]. Here we define a series of displays to understand compositional dynamics

or variability that ultimately interconnects to proteomic variability including RTCs and

DMVs (of course also other omics) through the organization of the genetic code [16–19]. We

subsequently compare SARS-CoV-2 with other human CoVs for between-population

variation analysis to point out that it is not a direct descendant of the previous human-

infecting CoVs. We finally make efforts to decipher the SARS-CoV-2 clades for its

variations and suggest that what we have seen now are not the natural picture of the

pandemics and the missing-links are not among human populations but the wildlife close to

human habitats in Southeast Asian territories, including islands and shorelines, not just

limited to bats and pangolins. We also show how to examine clade-associated permutation

variations and relate genetic variations to protein structures and phenotypic data. Nailing

down a single animal of human origin of the virus will not be the goals of this genomics-

based study but to provide information for smarter drug design, effective vaccine

development, accurate diagnostics.

Results and Discussion

Compositional dynamics and its parameters are essential and useful features for

evaluating the evolutionary status and molecular mechanisms of SARS-CoV-2 towards

pandemics

RNA genomics is very different from DNA genomics in several ways [11]. First, in the RNA

genome, the A:U basepair is actually 2 Daltons heavier than that of the G:C basepair due to a

larger molecular weight of uridine, whereas in the DNA genome the GC pair is a single

99 Dalton heavier than that of the A:T basepair. In CoVs, the G+C content is actually in a trend

100 of reducing so that the virus is in turn becoming heavier due to the increased U content [11].

101 Second, single-stranded RNA genomes are synthesized without stable double-stranded

102 intermediates that allow mismatch repair albeit existence of short and extremely rare double-

103 stranded RNA fragments involved in interference-based immunity [20, 21]. Third, the

104 existence of Wobble basepairing for secondary structures is of essence for operational

105 functions of all RNA molecules in addition to genetic information inheritance [22]. That said,

106 we can now look at how the RNA genome of SARS-CoV-2 and related CoVs take advantage

107 of these RNA-centric features.

108  As a positive-sense single-stranded RNA virus, SARS-CoV-2 has a genome length of ~

109 29903 nucleotides (nt) (GenBank: NC_045512.2). It encodes two large polypeptides, ORF1a

110 and ORF1b, along with other 15 non-structure proteins (nsps; **Figure 1**A). In order to

111 propagate and complete the life cycle, its positive-sense genome is first replicated to

112 synthesize full-length negative-sense antigenomes and 10 shorter subgenomes (sgRNAs),

113 executed by RTCs and DMVs, and the sgRNAs encode four structural proteins (S, spike; E,

114 envelope; M, membrane; and N, nucleocapsid) and six accessory proteins (ORF3a, ORF6,

115 ORF7a, ORF7b, ORF8, and ORF10) arranged among structural proteins depending on the

116 current annotation (GenBank: NC_045512.2).

117  Traditionally, we use three basic plots to display composition dynamics based on primary

118 genomic parameters over genome length: G+C contents at three codon positions (GC1, GC2

119 and GC3); purine content (A+G content); and GC skew (the content of (G-C)/(G+C)). Here,

120 we use a 300-nt sliding window with a step size of 21 nt, as the majority of viral sequences

121 are protein-coding, to illustrate the dynamics of the composition parameters, G+C and purine

122 contents (Figure 1B). The G+C content of SARS-CoV-2 varies in a narrow but significant

123 window of 18.00% (31.67%–49.67%) and the purine content in a slightly narrower window

124 of 15.50% (41.67%–57.17%) in average over the entire genome length. The GC skew of the

125 SARS-CoV-2 genome indicates the G+C ratio is relatively higher in structural proteins than

126 ORF1ab and this imbalance is a signature of distinct mutational biases caused by viral

127 replication machinery, known as RTCs. It is also variable as a frequent shift toward negative

128 values are often seen in individual ORFs and defined proteins. Such minor anomalies suggest

129 either recombination or selection events, which are species- or isolate-specific. The

130 differences become obvious when SARS-CoV-2 is compared to the closely-related bat and

131 pangolin CoVs (raf-betaCoV-RaTG13 and mja-betaCoV-P4L; Figure S1A and S1B), and the

132 authentic within-species variation is exemplified when SARS-CoV and MERS-CoV are

matched up with civets and camels in similar parameters, respectively (Figure S1C, S1D,

S1E and S1F). The G+C content of different codon positions is also very informative, where

GC3 is very characteristic of mutation pressure as it is obvious that all GC3 values of the

viral proteins are biased toward lower G+C contents. GC3-associated mutations often reflect

directional mutation patterns as observed strongly in certain lineages of plants and warm-

blooded vertebrates as negative gradients from the transcription starts, and such trends are

attributable to a special DNA repair mechanism, transcription-coupled DNA repair [23–25].

The notion here is to remind ourselves that transcription-centric mutations may be accounted

for some of the mutation events in RNA viruses in their replication-transcription processes.

Occasional twists from the trend often indicate selective pressures, such as in the case of S,

M, and N proteins, and weaker GC3 or stronger GC1 or GC2 selections. Codon-associated

G+C content trends are less informative for small ORFs, such as the case of ORF10. Most of

the sequence signatures are indicative rather than proven functional relevance of proteins but

very useful for providing clues of sequence anomaly.

For studying RNA viral genomes, in addition to previously-defined parameters, we need

to introduce the concept of single nucleotide (A, U, G and C) contents at three codon

positions (such as U1, U2 and U3 for uridines) (**Figure 2**) and to plot out compositional

dynamics for positive-sense or the genome and negative-sense strands, which include both

templates for synthesizing new positive-sense genomes or antigenome and subgenomes. All

viral mRNAs are transcribed from an antigenome and other 9 subgenomes (Figure 2B). For

compositional dynamics of RNA genomes, uridine is the star nucleotide, and A+U content

becomes the most important. Just for the sake of convenience, we would like to keep the

concept of G+C content since it has been known to be a useful variable for DNA

compositional dynamics [26] and provide an approximation for less selected nucleotide

position. For both DNA and RNA viruses, this variable G+C content remains similar to their

hosts mostly, except those that are not well adapted to their hosts, such as SARS-CoV or

MERS-CoV (Figure S2A; [11]).

As shown in a phylogenetic tree constructed based on 15 representative coronaviruses

(Figure 2A), the nucleotide content of SARS-CoV-2 is most similar to those of raf-betaCoV-

RaTG13 and mja-betaCoV-P4L, which are considered to be distantly related but most closely

related so-far-found host of SARS-CoV-2. Other known zoonotic and corresponding human

counterpart CoVs are rather close to each other in their compositions. We have made a few

interesting observations here. First, the single nucleotide content is more informative than

G+C content, especially for genome analysis on RNA viruses. The former points out only

how G+C content drifts toward richness or poorness but the latter narrows it down to single

nucleotide effect. In our case, U stands out at cp3, which alters the overall nucleotide

contents, and it drives the G+C content so low that even its partner A content has gone to the

same extremity, so that the low G+C content is a result of both lowering U and A. If the

organization principles are considered here, half of the codons are not sensitive to cp3

changes, and most of them are smaller amino acids (Figure S2B; [16–19]). Second, at the cp1,

G and C contents are both pulled apart toward extremity but not A or U, while the two

pyrimidines and two purines appear stretched to separate directions; these trends suggest

strong selective pressure at the first codon position over the entire genome. It is indeed that

cp1 codons shoulder the most mutation pressures since they fall into all 4 negative-sense

strand permutations (known as R1-derived permutations, C-to-U, G-to-A, U-to-C and A-to-

G). Third, the cp2 contents are most row-flipping changes referenced to the genetic code

organization [18]. These alterations are very useful for alternating chemical characteristics

between related amino acids, and in terms of flexibility, cp2 codons are less stringent than

cp3 but more flexible than cp1. The balancing power becomes more obvious when ORFs or

proteins are examined individually for their composition dynamics (Figure 1C). Finally, it is

conclusive that the more similar the CoVs in composition dynamic parameters, the closer

they are genetically and phylogenetically in principle. However, primary parameters, such as

G+C and purine contents are necessary but may not be sufficient. For instance, there has been

a CoV genome isolate from a wild vole captured in northeastern China, whose G+C and

purine contents overlap with SARS-CoV-2 completely (Rodent coronavirus isolate RtMruf-

CoV-2/JL2014; 0.38, 0.496; [27]) but its genome sequence is different (sharing 61.87%

identity with SARS-CoV-2). Therefore, we have yet to find a within-population immediate

animal host of SARS-CoV-2 albeit best similarity of composition dynamics seen among them.

Our subsequent study is focused on composition dynamics within CoV genomes. It is

interesting to see uniformity among all codon position contents of all CoV genomes,

increased G+C content from antigenomes to subgenomes. However, this trend is an illusion

where the real trend is the lower G+C content of antigenomes but higher G+C contents of

subgenomes due to stronger selection over structural proteins. This observation becomes

clearer when all ORFs and proteins are scrutinized one by one (Figure 1C and Figures S1).

SARS-CoV-2 has an exceptionally short subgenome 9 (sg9) which only contains ORF10, but

we have no evidence that it is either functional or non-functional. These results collectively

remind us that SARS-CoV-2 and its two most-closely-related CoVs, unlike in the case of

many other known CoVs, have a unique genome composition and similar dynamics to the

early-adapted human CoVs [11], and CoV-borne bats and other mammals may already coexist with ability to jump on to humans and domestic animals but only limited by environmental and geographic constraints.

Mutation spectrum is composed of permutations that are distinct according to their strand specificity, order of synthesis, and ratio of positive-sense vs. negative-sense strands during propagation

We use 12 permutations to represent directional mutations and classify them according to strand-specific replication mechanisms (**Figure 3**) since they are readily related to codons [11] (Figure S2B). From a total of 5,054 point mutations, 1,416, 1,497, and 2,141 mutations fall on codon position 1, 2, and 3, respectively. The permutations are categorized into R1 (C-to-U, G-to-A, U-to-C and A-to-G), R2 (C-to-A, U-to-G, A-to-C and G-to-U), and R12 (C-to-G, U-to-A, A-to-U and G-to-C) derived according to their occurrence tailored to RTC-directed strand synthesis: R1 from the first negative-sense strand, R2 from the subsequent positive-sense strand, and R12 from R1 plus R2. The most abundant permutations are four R1 permutations and one R2 permutation, G-to-U (Figure 3A and 3B). What have we shown here is how sensitive are nucleotide content of cp3 to selective pressure, and most cp3 permutations disappear except the R2 G-to-U permutations at cp3, where all changes are transversions and more than half of all codons (all pro-diversity changes) are sensitive to them. Similar results are observed in our analysis on SARS-CoV and MERS-CoV (Figure S3A and S3B). There are slightly different patterns among SARS-CoV and MERS-CoV and their within-population mammals from the SARS-CoVs and close relatives, the higher U-to-C permutations. The predominate C-to-U represents a driving force of variation, and it manifests why both G+C and A+G contents of SARS-CoV-2 appear relatively lower against MERS-CoV and SARS-CoV and even more when compared to human CoVs, such as 229E and OC43 (Figure S2A).

Since most cp1 and cp2 related permutations are sensitive to selection, we have examined how individual permutations correlated to codon rearrangements in the two halves tables: pro-diversity and pro-robustness (Figure 3C and 3D) [16, 17]. Only two examples, C-to-U and A-to-G, are shown here and the rest are summarized in Figure S3C. Several observations are worthy of in-depth discussion. First, it is known that three amino acids and their codons are unique in balancing one of two purine-sensitive halves; they are Leu (leucine), Arg (arginine), and Ser (serine) [16–19]. The most abundant amino acid in protein coding sequences (known as codon usage) is Leu and it buffers C-to-U|U-to-C mutations at cp1. Arg

and Ser are also abundant as they both are 6-fold degenerate codons; Arg appears buffering

A-to-G|G-to-A at cp1 and Ser carries two: U-to-A|A-to-U at cp1 and G-to-C|C-to-G at cp2.

Second, amino acid exchanges are permissive in physiochemical properties [23–25]. For

instance, Ser has a very similar size to Ala (alanine) so that G+C content increase is buffered

by the two amino acids as G-to-U|U-to-G permutations. Third, other examples are codon

alterations among hydrophobic amino acids as they are mostly C-to-U changes at cp2 among

those in the pro-robustness half. The overall effects are displayed together in Figure 3D. It is

rather clear that changes toward lower G+C content and near the balanced purine content are

both beneficial for CoVs, especially SARS-CoV-2, as these changes are pro-diversity, in

favor of larger and more hydrophilic amino acids.

Clade-associated biased mutation trend in SARS-CoV-2 revealed physiochemical

features of replication machinery

Difficulties for analyzing CoV genomes are multifold. Since we have yet to identify the

natural hosts and mammalian intermediate hosts, if there is any, this massive dataset has to be

analyzed by stratifying the data into structured and non-structured clades; the former can be

analyzed first and the rest await further ideas. The next is even more troublesome. Assuming

that we have 5 or more genome sequences per CoV isolate and variations identified among

them are still a miniscule fraction of the total virions produced in a patient body (medians and

means of variations per CoV isolate among C01 to C09, see Table S3), since the viral load

per patient sample, such as sputum [28, 29], is equivalent to a 5-person or more sampling of

the entire human population on earth, 1 out of 10^9. Even so, we have still been able to find

shared variations among patient samples occasionally and even more lucky to have some

clade structures, by and large due to the relatedness of the patients in the transmission

network. Finally, we have to admit that many assumptions have to be made about these

samples and their genome sequences above sequence and assembly errors for phylogeny and

genetic studies.

Nevertheless, we have constructed a somewhat stable phylogenetic tree-and-branch

structure for further analysis (**Figure 4**A). It is composed of 8 monophyletic clades and 1

non-monophyletic clade based on both orders of sample collection date and highly-shared

mutations. Among the clades, C02 shares two landmark mutations, C8782U in ORF1ab and

U28144C in ORF8, and earlier date (2019/12/30). C04 shares three more mutations

(C17747U, A17858G, and C18060U in ORF1ab) than what C02 have, and a late collection

date (2020/02/20). Clades C03, C05, and C07 are also distinguishable by some major

mutations, so are C06, C08, and C09; the latter clades are clustered together based on four
shared and other clade-associated mutations. The leftover large number of isolates that lack
all landmark mutations are grouped into C01, which have the earliest collection date on
2019/12/24. According to the literature and our discussion, we have further grouped the
clades into three clusters, S (C02 and C04), G (C06, C08 and C09), and L (all the rest) since
phylogeny shows clear divergence among them. We have several notions about this imperfect
hierarchical structure. First, our within- and between-clade analysis of high major allele
frequency (MAF) variations reveals that some clade-associated signature mutations are also
shared among clades. For instance, C14805U in ORF1ab and A24034G in Cluster S have
recurred in other clades of different clusters, which are excellent landmarks for subclade
definition. Another notion is that higher MAF within-clade mutations (such as MAF>0.2) are
mostly non-synonymous mutations, indicating selection at work (Figure S4). Our neighbor-
joining tree based on distances from 9 clades suggests that SARS-CoV-2 appears originated
from multiple zoonotic reservoirs instead of a single direct ancestor (Figure S4). In addition,
our classification rationales are largely in agreement with published reports [30]; for example,
Cluster S is in accordance with previously defined S type [31] and Cluster G is in line with
GISAID [32] defined the G clade. Cluster L is similar to the V and L clades combined, of
GISAID. A maximum likelihood (ML) based unrooted phylogenetic tree is shown in Figure
4B.

To look for clade-associated compositional and functional features, we have first built a
consensus sequence for each clade and subsequently calculated frequencies for each within-
clade permutation (Table S2; **Figure 5**A and 5B). A key assumption behind this is that
certain functional mutations may have clade-specific effects on mutation spectrum, to close a
loop where sequence mutations through genetic coding principles alter the viral proteome
function. Our observations are of importance in establishing logics about compositional
dynamics between nucleic acids and proteins. First, permutations among clades are indeed
variable according to their proportions calculated from genome variants, and aside from 5
high-proportion permutations, 4 R1 and 1 R2 permutations, two other R2 and one R12
permutations appear also joining in, which are U-to-G and A-to-C, as well as A-to-U,
respectively. Second, the variable permutations, where some may represent effect of mutation
pressure and others may exaggerate selection pressure, are unique to clades and clade clusters.
For instance, clade cluster S has the lowest G-to-U fraction as compared to those of L and G;
in addition, among the S clades, C04 has the lowest value of G-to-U. Similarly, C03, C05,
C06, C08, and C09 have relatively higher G-to-U permutations. Third, based on the disparity

of permutations or simply mutation spectra, we have taken a rather radical step to assume

RTC statuses in favor of either *tight* or *loose* statuses for binding to purines and pyrimidines

(see Figure S5). Since purines are larger than pyrimidines in size, the purine- or R-tight must

be different from pyrimidine- or Y-tight. The results are strikingly predictable in that the R-

tight status suggests a tighter binding pocket where a descending trend for tight permutations

(C-to-U, G-to-U, and U-to-A) reverses into the opposite trend for Y-tight permutations. It

indicates that the RTC structure and conformation variables may be definable in principle. At

this point, we do not have discrete definitions for these so-called tight statuses but the less

trendy R-loose and Y-loose statuses also support a similar idea.

We have further examined the compositional subtleties among the clades and clusters

with a focus on G+C and purine content variability as both contents appear drifting toward

optima in SARS-CoV-2 and its relatives (Figure 5C and 5D). Different clades exhibit distinct

compositional features and such dynamics are very indicative for the existence of feedback

loops connecting RNA variables to protein variables. Two directions have to be advised for

understanding these features albeit in absence of between-clade statistics. The first direction

is driven by strong mutations, perhaps coupled to tight-loose switches in the catalytic pocket

of RdRPs in RTCs. It is clear that except C01, the G or C06-C08-C09 cluster has the lowest

G+C (0.37929, based on a C08 CoV sampled in Australia) and the lowest purine contents

(0.49527, based on a C08 CoV collected in Bangladesh and a C09 CoV collected in England).

Both lower G+C and purine contents are indicative of mutation pressure and signal this fast-

evolving cluster of CoVs. Since this cluster has the largest collection of CoVs, it is also not

surprising to see a more complex median diversification within clades (Figure 5C and 5D).

The second direction is the drive from selection or both selection and mutation in balance or

imbalance, as well as in modes of fine-tuning or quick-escaping. Some results from our

analyses are shown here for briefing purposes (Figure 5E to 5G). For instance, G+C and

purine contents at cp3 are informative for mutation drives and other measures are less clear

cut (Figure 5F), given the evidence that even MAFs among clades are not stably distributed

among clades as lower MAF variations are rather sporadic and hard to analyze even binned

into groups (data not shown).

Based on our clade and clade cluster analysis, it is tempting for us to speculate that there

are plenty of rooms for further investigations into mutation spectra among large clades and

even smaller clades or closely related individual CoV genomes for several reasons. First, all

high-frequency MAFs should be identified and classified and these variations are candidates

for highly selected mutations. Second, all within-clade minor but not rare alleles (less than

1/10,000), such as those of MAFs in a range of 0.01% to 10% should also be identified; they

provide basis of within-clade sequence analysis. Third, all non-structured CoV genomes must

be also classified based on shared variations, as they are not only valuable for within-clade

but also for clade-cluster analyses since there is a large background of genome variations not

yet brought into the databases.

Within-clade variations and their implications for future SARS-CoV surveillance

Within-clade compositional dynamics can also be very informative, especially for covering

and predicting future functional changes, such as identifying mutated and diversified forms of

CoVs for drug and vaccine designs. It is also of essence for nucleic acid-based diagnostics,

such as clade-specific identifications. We are in a process of developing an interactive

database and mutation-function predicting algorithms based on our results to interpret novel

sequence variations in real time. Within-population variations are identified based on clade

consensus sequence after alignment and extracted from datasets that have hundreds and

thousands of genome sequences. The analysis of within-population variations relies on

structured phylogeny and proportion change of permutations. The changes, based on

functional relevance, can be classified into either copy number-related or RTC-specificity

related, or sometimes both.

We have taken two steps to extract information in order to distinguish the underlining

mechanisms (**Figure 6**). In the first approach, we identify key mutations based on MAF of

mutations with a consideration of relatively even distribution among subclades and name the

subclades in a sequential order based on the absence of a subset (Figure 6A). In the second

step, we plot out permutations to track changes among subclades (Figure 6B). For instance,

clade C02 can be divided into 8 subclades and its variable permutation fractions are clearly

recognizable. An immediate discovery is the trends of descending C-to-U, ascending A-to-G,

and wavy G-to-U that initially goes up with A-to-G but rides down with C-to-U afterward.

Taking the two smaller clades, such as C03 and C05, as examples (Figure 6D and 6E), we

first find that their trends of permutation variables show opposite directions, where the

increasing C-to-U accompanies with the decreasing G-to-U. A closer examination reveals

that the increasing C-to-U in C03 is also accompanied by descending U-to-C. The only

permutation showing an increasing trend in C03 is G-to-A. The take-home message from

these trends is that RNA synthesis of this subclade is biased toward producing more negative-

sense strands or its mutation spectrum exhibits increasing mutations generated during the

negative-sense strand synthesis. Such analysis can be carried out continuously when more

CoV genome data become available as other within-clade variations are not as informative as
C03 vs. C05 (Figure S6).

Several precautions are worth noting in such analysis. The most noticeable weakness is
the fact that we assume function-related mutations are discovered in our dataset. As we have
proposed an analogy before, chances are slim, dozens out of millions or even billions.
Furthermore, even if we see drastic changes in permutations and mutation spectra, the
mutations we identified still need validation empirically and based on different data types or
sources albeit rare and precious. Finally, most frequently encountered situations are those that
multiple mutations exhibit cofounding effects for a phenotypically identified functional or
structural feature, and undoubtedly, more and deep-sequencing data are still invaluable and
irreplaceable.

Conclusion

This COVID-19 pandemic provides once-in-a-lifetime opportunity for the fields of
biomedicine and other life sciences to work together on it as many facets as possible albeit
exchanging with lives and other massive losses. If lessons told, we had learned things in
serious ways in the last two CoV epidemics and we did prepare ourselves with vaccines and
medication since, we would not have suffered this much this time. If one assumes that the last
two outbreaks of SARS-CoV and MERS-CoV came surely by chances, this time SARS-CoV-
2 is here for real, and a worst-case scenario is that it may stay with us forever or until
effective vaccination is developed. Nevertheless, it certainly will stay with us for quite a
while for many reasons [11]. First, at least it and other within-population versions of
coronaviruses will definitely come again because we have not been able to trace its origin and
ways its transmission from the very beginning, neither the Wuhan outbreak nor the recent
Beijing outbreak in China even guided with very strict quarantine roles and prompt action
plans. Next, this particular virus, SARS-CoV-2, has evolved to a composition status where
some of its natural yet genetically distant hosts or possible intermediate mammalian hosts
have acquired similar status [33, 34]. Furthermore, we do not yet have enough data to really
map out the phylogenetic position that allows us to pinpoint its natural origin and human
transmission routes.

The number one needs for us is data, genomic and clinical data, which should be as
complete as possible and with characteristics including high-quality and high-coverage at
single-molecule resolution. We currently have been acquiring genomic data and the

specialized databases have collections over ten thousand non-redundant sequence variations,

but still not enough to address more than a few possible functional changes of some key

protein components [35–38], let alone understanding mutation-centric cellular mechanisms.

Based on median and mean estimates, we have on average a mutation accumulation rate of

half a dozen per patient. Although there have been data reported from single-molecule

sequencing platform but they are low in coverage [39].

Our final notion is to emphasize the importance of analysis strategies and supporting

platforms. Since questions always overwhelm what we can possibly address [40], prioritizing

tasks are of essence together with choices of strategies. The first platform to be established

concerns mutation-to-function interpretation, where we have present one in this report.

Another to be considered is mathematic modeling, such as cellular and disease transmissions

[41–46] and viral mutation-selection paradigm, for testing and evaluating different

parameters and prioritizing what kind of data to be acquired with high priorities. In addition,

cellular and molecular data, including different omics studies [47], all need to be incorporated

into a COVID-19 knowledgebase, where information from multi-disciplinary studies are

managed, organized, and mined.

421   Materials and Methods

422   SARS-CoV-2 and other related coronaviruses sequences

We used the public-available SARS-CoV-2 data collected worldwide among the major

databases, including CNCB/NGDC [48], CNGBdb [49], GISAID [32], GenBank [50] and

NMDC [51] on June 12[th], 2020. To ensure authenticity and reliability, our datasets must meet

the following criteria: (1) The genome sequence is labeled as complete that covers all coding

regions of the reference genome (GenBank accession NC_045512.2). (2) It has no more than

15 uncertain bases that often substituted as "N"s. (3) It has no more than 50 degenerate bases

that often labeled as discrete nucleotides. These high-quality genomes were aligned to the

reference using MUSCLE (version 3.8.31) with default parameter settings [52]. Further

analyses of SARS-CoV-2 and related CoV genomes are referenced to genome annotation of

the same reference genome (NC_045512.2) and other information provided by the RefSeq

database at NCBI.

Other closely related CoV genome sequences used include hsa-betaCoV-HKU1, hsa-

betaCoV-OC43, ave-gamaCoV, mga-gamaCoV, smu-alphaCoV-WS, hsa-alphaCoV-229E,

hsa-alphaCoV-NL63, taf-alphaCoV-NL63, MERS-CoV (from human and camel hosts), cdr-

betaCoV-B73, SARS-CoV (from human and civet hosts), pla-betaCoV-SZ3 and raf-

betaCoV-RaTG13 are retrieved from NCBI and mja-betaCoV-P4L are retrieved from NGDC.

A full listing of our sequences dataset including virus genre, strain name, accession number

and sources is provided in Table S3.

Calculation of genomic composition parameters

We display several genomic composition dynamics and its parameters (G+C content, A+G

content and GC skew) using different sliding windows. The first 300 nt are grouped as an

initial window, and subsequent windows are uniformly shifted in a 21-nt step. Within these

displays, the G+C contents referenced to the three codon positions of each open reading

frame or ORF are measured by adjusting the sliding window according to the ORF lengths

within viral genomes. As for ORFs longer than 2000 nt, a relatively large window size (300

nt) is adopted, and the step size is calculated via a custom formula $round\left(\frac{length_{ORF}-300}{600}\right) -$

$vb,\ 0 \leq vb \leq 2$ where $length_{ORF}$ denotes the length of ORF and vb varied from zero to two

bases to make sure the window size is divisible by 3; for ORFs with a medium size (longer

than 500 bases and shorter than 2000 nt), the window size is defined as $round\left(\frac{1}{4}*\right.$

$\left. length_{ORF}\right) - vb,\ 0 \leq vb \leq 2$, while the step size is simply defined as 3 nt; as for those

small ORFs (shorter than 500 nt) such as structural proteins, a constant 21-nt window size

and 3-nt step size is used for calculating genomic composition frequency.

The criteria for choosing the representative CoV genome sequences for constructing a

representative phylogenetic tree (Figure 2) are multi-fold. First, we include all 7 human-

infecting coronaviruses for the analysis, which are SARS-CoV, SARS-CoV-2, MERS-CoV,

hsa-alphaCoV-229E, hsa-betaCoV-OC43, hsa-betaCoV-HKU1, and hsa-alphaCoV-NL63 (a

prefix hsa- standing for *Homo sapiens* was used to label the unfamiliar human-infecting

CoVs). Second, we categorized all human-infecting, for simplicity, into 4 lineages: SARS-

CoV-2, SARS-CoV, MERS-CoV, and the older human CoV lineages. Therefore, their related

CoVs in the literature were also selected for the analysis, including a single closely-related

CoVs for each lineage (based on sequence identity): SARS-CoV-related (pla-betaCoV-SZ3),

MERS-CoV-related (cdr-betaCoV-B73), SARS-CoV-2 related (raf-betaCoV-RaTG13), and

NL63-related (taf-alphaCoV-NL63; both species and CoV genera were labelled for clarity).

Third, we also added more informative CoV genome sequences to enrich lineage-associated

information, which are a pangolin coronavirus genome (mja-betaCoV-P4L) reported to be

closed to SARS-CoV-2 and 3 non-beta-coronaviruses that infect animals (e.g., ave-gamaCoV

from gamma-coronavirus genus and smu-alphaCoV-WS from alpha-coronavirus genus).

Fourth, we only used complete protein-coding sequences from the CoVs to construct the

phylogenetic tree and to calculate genome parameter contents. The sequences were aligned

by using MUSCLE and the UPGMA tree was constructed by using MEGA-X [53]. The G+C

content and single nucleotide content of each virus genome at three codon positions was also

calculated. Subgenomes of SARS-CoV was obtained from Marra et al [54], and we annotated

the subgenomes of SARS-CoV-2, mja-betaCoV-P4L, and raf-betaCoV-RaTG13 based on the

annotation of NCBI (GenBank accession NC_045512.2). In addition, G+C and single

nucleotide contents of the complete genome and its subgenomes of these four viruses at three

codon positions were displayed to serve as sequence composition references.

Variation calling and categorization

All sequence variations are identified and categorized based on comparisons between the

query and the reference genomes, and files were generated by using an in-hoc Perl scripts

based on alignment results. The tailored annotation (gene, location and consequence on the

protein sequence) of each variant is determined with VEP (version 99.0) [55]. Since a large

number of gaps and low-quality sequences at the 3' and 5' ends, variations (substitutions,

insertions and deletions or indels) occurring 50 nt each at 5'- and 3'-ends of the genome are

not considered. Since the higher quartile of variations per genome among SARS-CoV-2

populations is 9 (based on the 22,051 sequences we analyzed in this study), we filtered out

the problematic sites that exceed 50 variations as compared to the reference genome. CoV

genome sequences have at least one mutation are used in this study. A full listing of

variations among coding regions identified in this study is provided in Table S4.

All continuously updated mutation files of the SARS-CoV-2 populations in variant call

format (version 4.2) are deposited at the variation page of the 2019nCoVR database

contributed by CNCB/NGDC (https://bigd.big.ac.cn/ncov/variation/).

Mutation spectrum analysis

A mutation spectrum for within-population variations is composed of two lines of

information; one concerns mutations that are referenced to a population consensus built based

on the entire collection, and the other contains frequencies of all mutations and their

directional changes, i.e., permutations. To reduce pitfalls of sequencing errors, we only

selected mutations that occur more than twice in the whole collection of SARS-CoV-2

populations (clades or clade clusters that are often defined based on phylogenetic analysis).

In theory, there are 16 possible permutations but 4 of them are unrecognizable so that 12

permutations (C-to-U, A-to-G, U-to-C, G-to-A, G-to-U, U-to-G, A-to-C, C-to-A, U-to-A, G-

to-C, C-to-G and A-to-U) are there as an informative set. When the number of CoV genomes

collected are limited, such as SARS-CoVs and MERS-CoVs, entire data sets are pooled

together without clades. In our analyses on SARS-CoVs and MERS-CoVs (Figure S3A), we

aligned sequences from these two lineages to their reference genomes (SARS-CoV:

NC_004718.3; MERS-CoV: NC_019843.3) to call variations. When aligned on overlapping

sequences, due to large deletions and additional ORFs, are encountered, we always choose

the largest or only one of the ORFs to represent the segment, respectively. For example, in

SARS-CoV lineage, if a mutation falls into the overlapping region of ORF9a (encoding the N

protein) and ORF9b, we have only used the ORF9a annotations to avoid redundancy.

Phylogeny constructing

Given the scale of SARS-CoV-2 sequence collections, we focused on genomes with unique

information contributing to phylogenetic analysis. First, mutations (including single-

nucleotide substitution and indels) at frequencies equal or greater than 10 in between-clade or

–population calls were selected. FastTree (version 2.1.11) [56] is used to construct maximum

likelihood phylogeny based on 5,121 genomes that have met our criteria, and iTol [57], an

interactive web server was employed for setting an unrooted format and annotating samples.

For Figure S4, the neighbor-joining method is used for constructing phylogeny from the

Euclidean distance of the mutation frequency matrix of clades, and the tree was generated

and visualized by R package phangorn [58] and ggtree [59].

Estimation of G+C and purine contents of genome sequences

528   G+C and purine (or A+G) contents of CoVs in general vary in a narrow range, and therefore,

529   subtleties among the content changes have to be scrutinized with low-quality sequences

530   excluded. A more sensitive approach is used in this study where two points are assumed; all

531   genomes are full-length and variant alleles in coding sequences are the varied composition.

532   The absolute frequencies of A+G and G+C content are defined as:

$$Genomic\ AG\ content = \frac{8954 + 5492 + \left(A_{alt} - A_{ref}\right) + \left(G_{alt} - G_{ref}\right)}{29903 - (Del_{alt} - Ins_{alt})} \#(1)$$

534   and

$$Genomic\ GC\ content = \frac{5492 + 5863 + \left(G_{alt} - G_{ref}\right) + \left(C_{alt} - C_{ref}\right)}{29903 - \left(Del_{alt} - Ins_{alt}\right)} \#(2)$$

where 8,954, 5,492, 5,863, and 29,903 are the frequencies of A, G, C and total length of the SARS-CoV-2 reference, respectively. For any sequence compared with the reference, the Del_{alt} and Ins_{alt} measures the deleted and inserted nucleotides of this sequence, respectively, and that is why $(Del_{alt} - Ins_{alt})$ means the variation of sequence length. For all the variant sites in this sequence, $\left(A_{alt} - A_{ref}\right) + \left(G_{alt} - G_{ref}\right)$ in Equation (1) measures the number of A and G variations in compared sequence, A_{alt} and G_{alt} denote the number of nucleotides mutated to A or G while A_{ref} and G_{ref} represent the number of nucleotides mutated from A or G. Similarly, $\left(G_{alt} - G_{ref}\right) + \left(C_{alt} - C_{ref}\right)$ in Equation (2) represents the varied number of G and C among compared sequences.

Clade subgrouping

To detect trend followers and disrupters in mutation spectra, a pipeline was developed to select such genomes and mutations within clades iteratively. The first step includes locating high-frequency mutations (major alleles, MA) in a clade and extracting all genomes without this MA mutation to form a subset of the clade. The second step is, within the new subclade, to iterate the process until such mutations are thoroughly identified and no more mutations exceed a manually set threshold of MAF. Since the number of unique variations among clades have been varying significantly over time, the thresholds are 0.05 in C01, C04, C06, C08 and C09; 0.1 in C02, C03, C05 and C07. The proportion of permutations in each subclade and the located gene and mutation type (synonymous or non-synonymous) of mutations are provided in Table S5.

Authors' contributions

JY designed, supervised, and coordinated the study. SS, ZZ and JX participated in the design of the study. XT, QL, ZL, YZ performed the data analysis. JY, SS, XT, QL, ZL, YZ designed and drew the figures. JY and XT drafted the manuscript. JY, ZZ, SS, QL and XT revised the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors have declared no competing interests.

Acknowledgements

This work was supported by grants from The Strategic Priority Research Program of the Chinese Academy of Sciences [XDA19090116 to S.S., XDA19050302 to Z.Z.], National Key R&D Program of China [2020YFC0848900, 2017YFC0907502], 13th Five-year Informatization Plan of Chinese Academy of Sciences [XXH13505-05], K. C. Wong Education Foundation to Z.Z., and International Partnership Program of the Chinese Academy of Sciences [153F11KYSB20160008]. The Youth Innovation Promotion Association of Chinese Academy of Science [2017141 to S.S.]; Funding for open access charge: The Youth Innovation Promotion Association of Chinese Academy of Science. We thank our colleagues and students for their hard working on the 2019nCoVR (https://bigd.big.ac.cn/ncov). We thank a number of users and CNCB/NGDC members for reporting bugs and sending comments.

References

[1] Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, et al. A new coronavirus associated with human respiratory disease in China. Nature 2020;579:265–9.

[2] World Health Organization. Coronavirus disease (COVID-2019) situation report - 144. https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports (Jun 12 2020, date last accessed).

[3] He X, Lau EHY, Wu P, Deng X, Wang J, Hao X, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. Nat Med 2020;26:672–5.

[4] Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. N Engl J Med 2020;382:1177-9.

[5] Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. JAMA 2020.

[6] Coronaviridae Study Group of the International Committee on Taxonomy of Viruses. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. Nat Microbiol 2020;5:536–44.

[7] Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. N Engl J Med 2020;382:727–33.

[8] Cui J, Li F, Shi Z-L. Origin and evolution of pathogenic coronaviruses. Nat Rev
Microbiol 2019;17:181–92.

[9] de Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights
into emerging coronaviruses. Nat Rev Microbiol 2016;14:523–34.

[10] Fung TS, Liu DX. Human Coronavirus: Host-Pathogen Interaction. Annu Rev Microbiol
2019;73:529–57.

[11] Yu J. From Mutation Signature to Molecular Mechanism in the RNA World: A Case of
SARS-CoV-2. Genomics Proteomics Bioinformatics 2020;in press.

[12] Guo Y-R, Cao Q-D, Hong Z-S, Tan Y-Y, Chen S-D, Jin H-J, et al. The origin,
transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak - an
update on the status. Mil Med Res 2020;7:11.

[13] Smith EC, Blanc H, Surdel MC, Vignuzzi M, Denison MR. Coronaviruses lacking
exoribonuclease activity are susceptible to lethal mutagenesis: evidence for proofreading and
potential therapeutics. PLoS Pathog 2013;9:e1003565.

[14] Simmonds P. Rampant C-->U Hypermutation in the Genomes of SARS-CoV-2 and
Other Coronaviruses: Causes and Consequences for Their Short- and Long-Term
Evolutionary Trajectories. mSphere 2020;5.

[15] Di Giorgio S, Martignano F, Torcia MG, Mattiuz G, Conticello SG. Evidence for host-
dependent RNA editing in the transcriptome of SARS-CoV-2. Sci Adv 2020;6:eabb5813.

[16] Xiao J, Yu J. A scenario on the stepwise evolution of the genetic code. Genomics
Proteomics Bioinformatics 2007;5:143–51.

[17] Yu J. A content-centric organization of the genetic code. Genomics Proteomics
Bioinformatics 2007;5:1–6.

[18] Zhang Z, Yu J. On the organizational dynamics of the genetic code. Genomics
Proteomics Bioinformatics 2011;9:21–9.

[19] Zhang Z, Yu J. The pendulum model for genome compositional dynamics: from the four
nucleotides to the twenty amino acids. Genomics Proteomics Bioinformatics 2012;10:175–80.

[20] DeWitte-Orr SJ, Collins SE, Bauer CMT, Bowdish DM, Mossman KL. An accessory to
the 'Trinity': SR-As are essential pathogen sensors of extracellular dsRNA, mediating entry
and leading to subsequent type I IFN responses. PLoS Pathog 2010;6:e1000829.

[21] Totura AL, Baric RS. SARS coronavirus pathogenesis: host innate immune responses
and viral antagonism of interferon. Curr Opin Virol 2012;2:264–75.

[22] Crick FH. Codon--anticodon pairing: the wobble hypothesis. J Mol Biol 1966;19:548–55.

[23] Cui P, Lin Q, Ding F, Hu S, Yu J. The transcript-centric mutations in human genomes.

Genomics Proteomics Bioinformatics 2012;10:11–22.

[24] Cui P, Ding F, Lin Q, Zhang L, Li A, Zhang Z, et al. Distinct contributions of replication

and transcription to mutation rate variation of human genomes. Genomics Proteomics

Bioinformatics 2012;10:4–10.

[25] Wong GK-S, Wang J, Tao L, Tan J, Zhang J, Passey DA, et al. Compositional gradients

in Gramineae genes. Genome Res 2002;12:851–6.

[26] Lobry JR. Asymmetric substitution patterns in the two DNA strands of bacteria. Mol

Biol Evol 1996;13:660–5.

[27] Wu Z, Lu L, Du J, Yang L, Ren X, Liu B, et al. Comparative analysis of rodent and

small mammal viromes to better understand the wildlife origin of emerging infectious

diseases. Microbiome 2018;6:178.

[28] Pan Y, Zhang D, Yang P, Poon LLM, Wang Q. Viral load of SARS-CoV-2 in clinical

samples. Lancet Infect Dis 2020;20:411–2.

[29] Wang W, Xu Y, Gao R, Lu R, Han K, Wu G, et al. Detection of SARS-CoV-2 in

Different Types of Clinical Specimens. JAMA 2020.

[30] Rambaut A, Holmes EC, O'Toole A, Hill V, McCrone JT, Ruis C, et al. A dynamic

nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. Nat

Microbiol 2020.

[31] Tang X, Wu C, Li X, Song Y, Yao X, Wu X, et al. On the origin and continuing

evolution of SARS-CoV-2. National Science Review 2020.

[32] Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative

contribution to global health. Glob Chall 2017;1:33–46.

[33] Zhou H, Chen X, Hu T, Li J, Song H, Liu Y, et al. A Novel Bat Coronavirus Closely

Related to SARS-CoV-2 Contains Natural Insertions at the S1/S2 Cleavage Site of the Spike

Protein. Curr Biol 2020;30:2196-203 e3.

[34] Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak

associated with a new coronavirus of probable bat origin. Nature 2020;579:270–3.

[35] Becerra-Flores M, Cardozo T. SARS-CoV-2 viral spike G614 mutation exhibits higher

case fatality rate. Int J Clin Pract 2020:e13525.

[36] Daniloski Z, Guo X, Sanjana NE. The D614G mutation in SARS-CoV-2 Spike increases

transduction of multiple human cell types. bioRxiv 2020:2020.06.14.151357.

[37] Korber B, Fischer W, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Spike
mutation pipeline reveals the emergence of a more transmissible form of SARS-CoV-2.
bioRxiv 2020:2020.04.29.069054.

[38] Zhang L, Jackson CB, Mou H, Ojha A, Rangarajan ES, Izard T, et al. The D614G
mutation in the SARS-CoV-2 spike protein reduces S1 shedding and increases infectivity.
bioRxiv 2020:2020.06.12.148726.

[39] Wang M, Fu A, Hu B, Tong Y, Liu R, Gu J, et al. Nanopore target sequencing for
accurate and comprehensive detection of SARS-CoV-2 and other respiratory viruses.
medRxiv 2020:2020.03.04.20029538.

[40] Teymoori-Rad M, Samadizadeh S, Tabarraei A, Moradi A, Shahbaz MB, Tahamtan A.
Ten challenging questions about SARS-CoV-2 and COVID-19. Expert Rev Respir Med 2020.

[41] Liu Q, Zhao S, Shi C-M, Song S-H, Zhu S, Su Y, et al. Population genetics of SARS-
CoV-2: disentangling sampling bias and clustering infections. Genomics Proteomics
Bioinformatics 2020;in press.

[42] Cotten M, Watson SJ, Kellam P, Al-Rabeeah AA, Makhdoom HQ, Assiri A, et al.
Transmission and evolution of the Middle East respiratory syndrome coronavirus in Saudi
Arabia: a descriptive genomic study. Lancet 2013;382:1993–2002.

[43] Gire SK, Goba A, Andersen KG, Sealfon RSG, Park DJ, Kanneh L, et al. Genomic
surveillance elucidates Ebola virus origin and transmission during the 2014 outbreak. Science
2014;345:1369–72.

[44] Lemey P, Suchard M, Rambaut A. Reconstructing the initial global spread of a human
influenza pandemic: A Bayesian spatial-temporal model for the global spread of H1N1pdm.
PLoS Curr 2009;1:RRN1031.

[45] Smith GJD, Vijaykrishna D, Bahl J, Lycett SJ, Worobey M, Pybus OG, et al. Origins
and evolutionary genomics of the 2009 swine-origin H1N1 influenza A epidemic. Nature
2009;459:1122–5.

[46] Yu W-B, Tang G-D, Zhang L, Corlett RT. Decoding the evolution and transmissions of
the novel pneumonia coronavirus (SARS-CoV-2 / HCoV-19) using whole genomic data.
Zool Res 2020;41:247–57.

[47] Sanders W, Fritch EJ, Madden EA, Graham RL, Vincent HA, Heise MT, et al.
Comparative analysis of coronavirus genomic RNA structure reveals conservation in SARS-
like coronaviruses. bioRxiv 2020:2020.06.15.153197.

[48] National Genomics Data Center M, Partners. Database Resources of the National
Genomics Data Center in 2020. Nucleic Acids Res 2020;48:D24–D33.

[49] Wang B, Liu F, Zhang EC, Wo CL, Chen J, Qian PY, et al. The China National

GeneBank horizontal line owned by all, completed by all and shared by all. Hereditas(Beijing)

2019;41:761-72.

[50] Sayers EW, Cavanaugh M, Clark K, Ostell J, Pruitt KD, Karsch-Mizrachi I. GenBank.

Nucleic Acids Res 2020;48:D84–D6.

[51] Wu L, Sun Q, Desmeth P, Sugawara H, Xu Z, McCluskey K, et al. World data centre for

microorganisms: an information infrastructure to explore and utilize preserved microbial

strains worldwide. Nucleic Acids Res 2017;45:D611–D8.

[52] Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 2004;32:1792–7.

[53] Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. Mol Biol Evol 2018;35:1547-9.

[54] Marra MA, Jones SJ, Astell CR, Holt RA, Brooks-Wilson A, Butterfield YS, et al. The

Genome sequence of the SARS-associated coronavirus. Science 2003;300:1399-404.

[55] McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GRS, Thormann A, et al. The Ensembl

Variant Effect Predictor. Genome Biol 2016;17:122.

[56] Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees

for large alignments. PLoS One 2010;5:e9490.

[57] Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new

developments. Nucleic Acids Res 2019;47:W256–W9.

[58] Schliep KP. phangorn: phylogenetic analysis in R. Bioinformatics 2011;27:592–3.

[59] Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Curr Protoc

Bioinformatics 2020;69:e96.

Figures

Figure 1 A display of genome compositional dynamics of SARS-CoV-2 and related

CoVs

A. The complete genome sequence of SARS-CoV-2 (NC_045512.2), including both

structural and non-structural components. **B.** We use a 300-nt sliding window with a 21-nt

step to show dynamic changes of genome G+C, purine, GC1 to GC3, and GC skew (G-

C/G+C) contents over the entire genomes. **C.** A similar procedure as described above is

applied to individual ORFs and proteins. Note that the GC skews are not uniform over the

genome length and the observation suggests possible recent recombination among closely
related CoVs.

Figure 2 Nucleobase contents of genomes and subgenomes of SARS-CoV-2 and related
CoVs
A. A schematic phylogenetic tree is used to cluster genome sequences and compositional
variables (15 CoVs genome sequences, from top to bottom, are: hsa-betaCoV-HKU1, hsa-
betaCoV-OC43, ave-gamaCoV, mga-gamaCoV, smu-alphaCoV-WS, hsa-alphaCoV-229E,
hsa-alphaCoV-NL63, taf-alphaCoV-NL63, MERS-CoV, cdr-betaCoV-B73, SARS-CoV, pla-
betaCoV-SZ3, mja-betaCoV-P4L, SARS-CoV-2, and raf-betaCoV-RaTG13). These
compositional variables include GC contents at three codon positions (codon positions 1, 2,
and 3 are denoted as GC1, GC2, and GC3) and single nucleotide contents at three codon
positions (A1, A2, A3; U1, U2, U3; G1, G2, G3; and C1, C2, C3). Nucleotides are labeled in
different shapes: purines, triangles; pyrimidines, open circles. A and U or G and C are
colored blue or red, respectively. It becomes obvious that the two closely-related CoV
genomes to SARS-CoV-2, the reported bat (raf-betaCoV-RaTG13) and the pangolin (mja-
betaCoV-P4L) have very similar codon G+C content as well as base contents. The cp1
(codon position 1) base content appears most characteristic of balanced purine content of
SARS-CoV-2 and its close relatives. The cp2 (codon position 2) base content of SARS-CoV-
2 and all other CoVs has higher and relatively balanced A+U content. The older human CoVs
have either lowest or higher G+C content and unbalanced purine contents. Note that G+C
contents in three codon positions are labeled differently from those of single nucleotide
contents in color codes. G+C content represents a single measure but single nucleotide
contents demonstrate trends of all four nucleotides. **B.** The G+C and single nucleotide
contents at different codon positions of complete genomes and subgenomes of SARS-CoV-2,
SARS-CoV, mja-betaCoV-P4L, and raf-betaCoV-RaTG13 are displayed to illustrate the
driving force for G+C content decrease toward 3' end of the genome, which is rather a result
of, in terms of mechanism, the increased U content and C-to-U permutation. The negative
gradient of U is also obvious from the 5' end to the 3' end.
Figure 3 Mutation spectra of SARS-CoV-2 in a context of G+C contents and codon
positions

A. The SARS-CoV-2 mutation spectrum is composed of 12 permutations and they are
divided by codon positions among all mutations. C-to-U (CU), U-to-C (UC), A-to-G (AG),
G-to-A (GA), and G-to-U (GU), are always dominant due to two principles; one is that the
first four permutations occur when positive-sense genome is synthesized, and the other is that
a G-by-A replacement is always preferred by RTCs so that G-to-U permutation as the most
dominant occurring when the antigenome serves as a template. **B.** C-to-U permutations at cp3
diminish among non-synonymous mutations and this phenomenon indicates that most protein
composition relevant variations are cp1 and cp2 variations. The remaining non-synonymous
mutations in G-to-U (GU) permutation may be a result of biased strand synthesis. **C.**
Displays of permutation-to-codon changes among non-synonymous mutations. The codon
table is divided into two halves: the pro-diversity half (blue) whose cp3 is sensitive to
transitional change and the pro-robust half (purple) whose cp3 position is insensitive to any
change. Two examples, C-to-U (1051 in counts) and A-to-G (314 in counts) permutations are
shown here. When a codon has a C-to-U change, the codon position varies, results of such
changes relative to codon positions are summarized on both sides of the codon flow chart.
Note that cp1 and cp2 changes appear more than those of cp3. The ratio between codons of
pro-robust half and pro-diversity half is displayed on each bar. **D.** All permutations are
plotted against the reference genome sequence to show how changes are related to amino
acids. In the molecular weight index, most cp1 and cp2 changes are most obvious, showing
an increasing trend. In the hydrophobicity index, most cp1 and cp2 changes increase toward
less hydrophobicity.

Figure 4 Sequence-variation-based phylogenies of SARS-CoV-2
A. CoV genomes are divided into clades and clade clusters based on high-frequency
mutations among the genome sequences. The shared variations are excellent indicators for
shared ancestors and those between clusters (blue half parentheses) and within clusters (red
half parentheses) are labeled with positions and nucleotide variations that are all referenced to
the SARS-CoV-2 genome (NC_045512.2), its positions, and relative frequencies (thin
vertical bars). The dates when each clade started are also indicated. **B.** The current collection
shows 9 clades (C01 to C09) in three clusters (S, L and G). An unrooted phylogenic tree of
the clades and clusters (color-coded), the tree scale is 0.01.

Figure 5 Mutation spectrum and composition dynamics among 9 SARS-CoV-2 clades
A. Plots showing permutation variation of each clade. Aside from the 5 dominant
permutations, A-to-C (R2 permutation), U-to-G (R2 permutation) and A-to-U (R12
permutation) changes appear also significant; such an increase in proportion of R2 and R12

permutations often indicates copy number (synthesis) bias between the two strands. **B.** When
permutations are grouped based on structure-conformation model (Figure S5) into tight and
loose groups (a four-parameter model), their trends of changes become obvious. The R-tight
discourages A-by-G replacement but encourages C-by-U replacement when the genome is
replicated. The loose statuses, regardless R-loose or Y-loose, place no pressure on
permutation variability. **C**. Violin plots showing the G+C content among clades. **D**. Violin
plots showing the purine content among the clades. C08 and C09 have been drifting both
contents toward lower ends. C03 has also been drifting in a greater extent of its purine
content and a lesser extent of its G+C content, comparatively. **E.** The mean (solid circles) and
median (solid triangles) of G+C and purine contents among clades. The same three more
expressive clades, as seen in (**C**) and (**D**), are indeed obvious (inset). **F.** The compositional
dynamics of cp3 nucleotides that are less selected and with a stringent cutoff value (> or =5).
G. Composition distributions based on major alleles, at frequencies equal or greater than 0.01
to emphasize the effect from selection. **H**. Composition distributions based on major alleles at
frequencies equal or greater than 0.05. **I**. Composition distributions based on major alleles at
frequencies equal or greater than 0.1.

Figure 6 Within-clade permutation variations are excellent indicators of functional
mutations

A. An example of permutation shifting of clade C02 and among its subclades. The number of
SARS-CoV-2 genomes is indicated in the parentheses. The clear trends are two-fold. First,
decreased C-to-U permutation is coupled with increased A-to-G and decreased G-to-U
permutations. Second, A-to-U permutation is also increased as expected based on the model
shown in Figure S5. These trends pf permutation changes suggest irrelevant to the ratio of
strand-biased synthesis (positive sense vs negative sense) but possible structural and/or
conformational variation in the RTCs. (**B**) – (**F**) show within-clade permutation changes of
C02, C04, C03, C05 and C07. In each display, the first column of the x-axis shows the
proportion of permutations calculated for each clade. Two opposite trends of permutation
variations are seen between C03 and C04, and C07 has a rather wavy pattern.

Supplementary material

Supplementary Figure S1 A display of genome compositional dynamics of SARS-CoV,
MERS-CoV and their within-population CoVs

We use a 300-bp sliding window with a 21-bp step to show dynamic changes of genome

G+C, purine, GC1 to GC3, and GC skew (G-C/G+C) contents. The complete genome

sequences and data sources are listed in Table S1.

Supplementary Figure S2

A. G+C and purine content plot to show how these contents distribute among human CoVs.

Note that all older human CoVs are drifted toward lower G+C and purine contents, and this

phenomenon indicates lower selection pressure or insensitivity on composition changes. Full

names of the human CoVs are listed in the legend of Figure 2. **B.** A genetic code table to

show how nucleotide permutations are related to codons. The table is divided into two halves

(colored and uncolored backgrounds) and cp1 and cp2 relative to their permutations

sensitivity and changes are indicated with half parentheses with color-coding: C-to-U|U-to-C

and A-to-G|G-to-A, red; G-to-U|U-to-G and A-to-C and C-to-U, blue; and A-to-U|U-to-A

and G-to-C|C-to-G, green. Note that cp1 and cp2 are sensitive to column and row codon

swaps, respectively. Cp3 is in a unique position where only half of the codons are sensitive to

its changes, and the other half is so organized that some codons are more permissive than

others.

Supplementary Figure S3   Mutation spectra of SARS-CoV-2 in a context of codon

positions

A. MERS-CoV and SARS-CoV mutation spectra are composed of 12 permutations and they

are divided by codon positions. C-to-U (CU) permutations are always as dominant as what

SARS-CoV-2 shows. The mutation counts are partitioned into synonymous and non-

synonymous mutations. **B.** C-to-U permutations at cp3 diminish among non-synonymous

mutations and this phenomenon indicates that most protein composition relevant variations

are cp1 and cp2 variations. Note that all older human CoVs are drifted toward lower G+C and

purine contents, and this phenomenon indicates lower selection or insensitivity on

composition changes. Full names of the human CoVs are listed in the legend of Figure 2.

Mutation counts are calculated from non-synonymous mutations. **C.** Displays of permutation-

to-codon changes among non-synonymous mutations. The permutations showed here contain

U-to-C, G-to-A, G-to-U, C-to-A, U-to-G, A-to-C, A-to-U, G-to-C, U-to-A and C-to-G.

Supplementary Figure S4   High-frequency within-clade mutations

Signature site information and frequency table of star mutations in each clade, with a

neighbor-joining tree based on the frequency data in the table.

Supplementary Figure S5

This table illustrates how 12 permutations are related to G+C and purine contents and the
subtle RTC specificity of CoVs. Mutations occur when the positive-sense RNA genome (R1,
mutation happens when the negative-sense genome is synthesized) or its negative-sense
subgenomes are synthesized (R2). The G+C content insensitive permutations occur after two
syntheses (R12). CU and A-to-G (AG) are preferred when RTC is in a status that encourages
a large-to-small substrate exchange in a higher ratio than the opposite, small-to-large
substrate exchange. We term this preferred exchange as "tight" status. This status is also
divided into R-tight (R, purine; to indicate that the mechanism is an A-by-G replacement) and
Y-tight (Y, pyrimidine; to indicate that the mechanism is a C-by-U replacement). When an
exchange of substrate happens from small-to-large, it is referred as a "loose" status that is
also divided into two, R-loose and Y-loose. Note that some permutations are not sensitive to
G+C and purine contents but others are sensitive. Arrow-headed dashed lines connect R1
permutation to R12 permutations, and note that cross-column relationship is rather striking,
which re-routes some structural principles, which navigates mutation forces on one hand and
leaves room for selection to work on, on the other hand.

Supplementary Figure S6 Within-clade permutation variations are excellent indicators
of functional mutations

(**A**) – (**D**) show within-clade permutation changes of C06, C08, C09 and C01.

Supplementary Table S1 The mutation counts in each clade and cluster

Supplementary Table S2 The proportion of permutations in each clade and clusters
based on different genomic regions

Supplementary Table S3 Selected CoV genome sequences used for this study

Supplementary Table S4 The SARS-Cov-2 mutation table on coding region (based on
data on June 12th 2020)

Supplementary Table S5 The proportion of permutations in each clade and their
subclades