Population Genomics of *Plasmodium vivax* in Panama to Assess the Risk of Case Importation on Malaria Elimination

Lucas E. Buyon[1,3], Ana Maria Santamaria [2], Angela M. Early[1,3], Mario Quijada [2], Itza

Barahona[4], Jose Lasso[4], Mario Avila[4], Sarah K. Volkman [1,3,5] Matthias Marti[6], Daniel E.

Neafsey[1,3], Nicanor Obaldia III[1,2,6]

[1] Department of Immunology and Infectious Diseases, Harvard TH Chan School of Public

Health, Boston, MA. USA, [2]Instituto Conmemorativo Gorgas de Estudios de la Salud, Panama

City, Panama, [3]The Broad Institute, Cambridge, MA., USA [4]Ministerio de Salud, Panama,

Panama, [5]Simmons University, College of Natural, Behavioral and Health Sciences, Boston,

MA, USA [6]University of Glasgow, Glasgow, Scotland, UK.

To whom correspondence should be addressed: nobaldia@gorgas.gob.pa and

neafsey@hsph.harvard.edu

Abstract

Malaria incidence in Panama has plateaued in recent years in spite of elimination efforts, with almost all cases caused by *Plasmodium vivax*. Notwithstanding, overall malaria prevalence remains low (fewer than 1 case per 1000 persons). We used selective whole genome amplification to sequence 96 *P. vivax* samples from Panama collected between 2007 and 2019 to study the population structure and transmission dynamics of the parasite. Imported cases resulting from increased levels of human migration could threaten malaria elimination prospects, and four of the samples evaluated came from individuals with travel history. We explored patterns of recent common ancestry among the samples and observed that a single highly genetically related lineage was dominant among the samples (47 out of 59 samples with good sequencing coverage), spanning the entire period of the collection (2007-2019) and all regions of the country. We also found a second, smaller clonal lineage of four parasites collected between 2017 and 2019. To explore the regional context of Panamanian *P. vivax* we conducted principal components analysis and constructed a neighbor-joining tree using these samples and samples collected worldwide from a previous study. Three of the four samples with travel history clustered with samples collected from their suspected country of origin (consistent with importation), while one appears to have been a result of local transmission. The small number of genetically unique Panamanian *P. viva*x samples clustered with samples collected from Colombia, suggesting they represent the genetically similar ancestral *P. vivax* population in Panama or were recently imported from Colombia. The low diversity we observe in Panama indicates that this parasite population has been previously subject to a severe bottleneck and may be eligible for elimination. Additionally, while we confirmed that *P. vivax* is imported to Panama from diverse geographic locations, the lack of impact from imported cases on the overall parasite

population genomic profile suggests that onward transmission from such cases is limited and that

imported cases may not presently pose a major barrier to elimination.

Author Summary

Panama has greatly reduced *P. vivax* incidence, however, this progress has plateaued.

Understanding parasite transmission patterns and identifying imported cases is critical to help

Panama and other countries succeed in their elimination efforts. Genomic epidemiology and

population genomics can help provide information needed to inform malaria control policy. In

this study, we collected 100 Panamanian *P. viva*x samples from two collection periods (2007-

2009 and 2017-2019), of which 59 yielded usable sequencing data. 4 samples had patient travel

history data associated with them. We found that the majority of samples belong to a single

highly related lineage, termed CL1. This lineage has persisted since at least 2007. We also

highlight how genomic epidemiology can be used to spotlight parasites that may be imported as

a result of human migration, as well as corroborate or refute the country of origin as suggested

by patient travel history. We observe no evidence of outcrossing between these potentially

imported parasites and the local Panamanian parasite population, suggesting that imported

parasites are not driving ongoing malaria transmission in Panama. The low diversity we observe

in Panama indicates that this parasite population has been previously subject to a severe

bottleneck and may be eligible for elimination.

Introduction

Malaria is a parasitic disease transmitted by the bite of female *Anopheles* mosquitoes. Malaria

parasites cause approximately 219 million cases and 435,000 deaths each year, the vast majority

in sub-Saharan Africa (1). *Plasmodium falciparum,* the most virulent of the five *Plasmodium*

species that infect humans (*P. falciparum*, *P. vivax, P. malariae, P. ovale, and P. knowlesi*)

causes the majority of these cases (1). Though billions of dollars have been devoted to the

control and eradication of malaria caused by *P. falciparum,* comparatively little attention is given

to *P. vivax,* the most prevalent malaria parasite outside Africa (2). The impact of *P. vivax* on

human health was once considered minimal relative to the more virulent *P. falciparum* (1,2).

However, recent studies suggest this parasite species causes a significant global health burden

(1,2). The *P .vivax* cycle includes a dormant liver "hypnozoite" stage, unique among human

malaria parasites. (2) The hypnozoite stage can cause a relapse of malaria years after initial

infection, thus beginning the cycle of infection and complicating control efforts. (2)

Sixty percent of the population of Central and South America live in areas with ongoing

malaria transmission, predominantly caused by *P. vivax* (3). The region experiences about

700,000 *P. vivax* cases each year (1). Between 2000 and 2015 the incidence rate of malaria fell

37% globally and 42% in Africa (1). In the Americas during the same period, malaria mortality

decreased by 72% (1). Unfortunately, recent evidence suggests that this trend has stalled, and in

some countries, malaria incidence has even increased (1). Panama eliminated the autochthonous

transmission of *P. falciparum* in 2010*,* outside of a small outbreak on the Colombian border in

2015 (4). Since 2010, *P. vivax* has caused almost all the malaria cases in Panama (1,5,6). *P.*

vivax cases in Panama have declined precipitously since 2005, from close to 1 case per 1000

persons, to under 0.25 cases per 1000 persons in 2017 (7). However, malaria incidence in

Panama has plateaued since 2008. This plateau in incidence could be due to low levels of

transmission and/or imported cases that are re-seeding infections.

Human movement leading to parasite migration is a potentially significant

epidemiological threat to malaria control in Panama. Parasite importation stemming from human

migration is a challenge to elimination programs in other countries around the world (8–10).

The unique geographic position of Panama makes it a crossroads for human migration to the

United States from South America (5,11). Migrants enter Panama through two paths: through

the Darien jungle region on the border with Colombia, and through the Kuna Yala Amerindian

reserve ('Comarca') on the Caribbean coast (5,6,11). Previous studies implicate these regions as

focal points of ongoing malaria transmission in Panama and suggest this is partly due to

imported parasites (5,11). It is estimated that approximately 60,000 continental and extra-

continental migrants crossed the southern border of Panama through the Darien jungle region in

2015 and 2016 (12).

To inform effective malaria elimination strategies in Panama, it is critical to

differentiate between imported infections or sustained local infection as the underlying cause of

persistent *P. vivax* incidence in Panama (4,5). Whole-genome sequencing can help paint a

detailed picture of parasite movement and transmission within and between countries (10,12,13).

However, *P. vivax* cannot be grown *in vitro*, and the difficulty of sequencing *P. vivax* from

clinical samples dominated by host DNA has hindered parasite population studies(14). Recent

advances such as hybrid capture (15) and selective whole genome amplification (SWGA)

mitigate this problem by allowing for parasite DNA to be selectively enriched before

sequencing(14). Both methods have allowed for population genomic studies of *P. vivax* using

samples directly from patients.

In this study, we describe the population genomics of *P. vivax in* Panama over a 12-year

time span, with the aim of understanding patterns of genetic variation and recent shared ancestry

(relatedness) at different geographical and temporal scales. We found the vast majority of *P.*

vivax cases in Panama belong to a single highly related lineage that has persisted for at least a

decade. Furthermore, we observed a second smaller clonal lineage concentrated near the

Panamanian-Colombian border. We also found several samples that shared no relatedness with

any other sample, which may represent either localized pockets of outbred *P. vivax* transmission

or imported cases. These findings suggest the Panamanian *P. vivax* population has extremely low

genetic diversity and is on the cusp of elimination. Revealing these patterns of relatedness among

parasite infections can help inform best strategies for targeting interventions or case investigation

methods to increase the likelihood of successful elimination. The results obtained from this study

will help inform future elimination efforts in Panama and the rest of Meso-America.

Results

Sample Extraction, Sequencing, and Data Filtering

We successfully generated usable sequencing data from 35/60 (58%) Panamanian *P. vivax*

samples collected between 2007-2009 and 24/40 (60%) collected between 2017-2019, for a total

of 59 samples (Fig 1A). All Panamanian samples had an F_{ws} statistic greater than 0.95, indicating

that they were all monoclonal (Fig 1B).

We next analyzed the 59 Panamanian genomes in the context of 109 previously

published *P. vivax* genomes, generating a filtered VCF consisting of 168 samples and 62,211

high-quality biallelic SNPs.

Recent Common Ancestry Analysis Reveals Single Highly Related Lineage of Parasites

We used hmmIBD (23) to estimate the proportion of the genome that is IBD among

Panamanian sample pairs to understand patterns of recent common ancestry. We subsetted the

VCF to contain only samples collected in Panama and Colombia to estimate pairwise IBD. We

strictly filtered sites on the basis of minimum and maximum read depth (five and thirty

respectively), resulting in a VCF with 89 samples and 15,788 sites for input into hmmIBD.

We observed a bimodal distribution of pairwise IBD in Panamanian samples, with peaks

near zero and 0.95 (Fig 2A). Forty-seven of the 59 Panamanian samples shared high IBD

(>0.875) with each other, indicating very recent common ancestry. Four other Panamanian

samples, all collected in the Kuna Yala Province, shared 100% IBD with each other, and 0-10%

IBD with any other sample, Panamanian or Colombian (Fig 2A and S1 Fig). Another four

Panamanian samples exhibited no IBD with each other nor any of the other Panamanian samples.

All four of these samples were collected in the Darien jungle region or Kuna Yala, which are the

two main points of entry for migrants traveling through Panama. These four samples drive the

modal peak of pairwise IBD at zero.

The variable degree of relatedness among the 47 samples sharing > 0.85 IBD suggested

that data quality potentially impacted the estimation of IBD. We plotted the relationship between

IBD and sample quality, measured by the average proportion of high coverage sites in each

sample pair, to determine if pairwise sample data quality affected the estimation of IBD (Fig

2B). We defined high coverage sites as sites with greater than 5x coverage (the cutoff for site

filtering). We observed that as the average proportion of high coverage sites for sample pairs

increased, pairwise IBD estimates correspondingly increased as well (Fig 2B). This relationship

suggests that poor data quality can lead to underestimation of IBD. It is possible that the majority

of the pairwise IBD estimations would be closer to one had the overall sample sequence quality

been higher. The prevalent highly genetically related lineage is referred to henceforth as cluster

one (CL1). CL1 samples share an IBD fraction of at least 0.875 with other samples in this

cluster. We also concluded that the four samples that shared 100% IBD with each other

constituted a second completely clonal lineage, henceforth referred to as cluster two (CL2).

Next, we examined how these two clusters and the other Panamanian samples not

belonging to either lineage were geographically distributed in Panama (Fig 3). Samples from

CL1 were found across Panama. Notably, samples collected from both 2007-2009 and 2017-

2019 were found in this lineage. The inclusion of samples from both collection periods

demonstrates that this lineage has persisted throughout Panama for at least a decade. We did not

find any evidence of structure in the *P. vivax* population by region or relative to the Panama

Canal, as was previously observed for *P. falciparum* (4).

We only observed samples belonging to CL2 in a specific locality, Puerto Obaldia, in the

Kuna Yala Amerindian territory (Comarca) along the Atlantic Coast. We lacked geographic

information for one of the four samples in CL2. Collection location data for the other three

samples show they were collected in roughly the same location, Puerto Obaldía in Kuna Yala.

Additionally, of four samples that shared no recent common ancestry with any sample in the

dataset, three were collected in Darien province, along the Colombian border and one was

collected in Kuna Yala.

After identifying two highly related lineages in Panama, we explored an approach for

determining whether the samples excluded from analysis due to low coverage could belong to

one of these lineages. We identified a set of 264 genotyped SNPs that were called in at least 80%

of samples across both Panama sample collection periods. We then calculated Nei's standard

genetic distance on all pairwise sample comparisons. The majority of excluded samples in both

collection periods (17/21 and 4/16 for the 2007-2009 and 2017-2019 collection periods

respectively) exhibited very low levels (0-1%) of genetic distance with CL1 samples (S2 Fig).

Seven of the 2017-2019 samples exhibited 0-1% genetic distance with CL1 also showed 0-1%

genetic distance with CL2, making the identity of these samples uncertain. Three excluded

samples from 2007-2009 collected in the Darien Jungle Region had relatively high genetic

distance from all other samples in the dataset.

Exploring the regional context of Panamanian *P. vivax*

We built a neighbor-joining tree using the Panamanian samples plus previously

sequenced samples (20) (Fig 4A) to understand the Panamanian *P. vivax* population in a global

context. As noted in previous studies (20,30), we observed clusters of samples corresponding to

different geographic regions, with a large cluster of Central and South American samples. CL1

and CL2 formed distinct clusters within the Central and South American cluster with 100%

bootstrap support. CL2 is situated in a cluster containing samples from Colombia, with 100%

bootstrap support at deep nodes. While these four samples are clustered together with 100%

support and short branch length, a long branch connects them to the rest of the Colombian

cluster. The Panamanian samples that shared zero IBD with either cluster also grouped with the

Colombian samples. These samples seem to share distant ancestry with each other and the rest of

the Colombian samples. The samples also formed their own sub-cluster within the Colombian

cluster.

PCA conducted with all samples worldwide showed tight clustering of all Central and

South American samples, with only one Panamanian sample falling outside this Central and

South American cluster (S3 Fig). PCA restricted to the samples collected from Central and South

America is heavily influenced by covariation among samples within the two clusters (S4 Fig).

PCA performed with a single consensus sequence representing each cluster revealed CL1

clusters with samples from Peru and Brazil and CL2 clusters with the Colombian samples (Fig

4B). All four outbred Panamanian samples that shared no recent ancestry with the other samples

also clustered with the Colombian samples. Principal component one differentiated CL1 and

samples from Brazil and Peru from CL2 and the rest of the Central and South American samples.

211
Genomic data both support and refute travel history data

Four of the 59 samples had travel history data associated with them. All of these samples

with travel history data were collected during the 2007-2009 period. Travel history information

suggested that two samples were originally from Brazil, one sample from India, and one sample

from China. The two samples with Brazilian travel history fell within the Brazilian cluster on the

NJ tree and clustered with Brazilian samples on PCA (Fig 2A and 2B). The one sample with

Indian travel history grouped with the other Indian samples on the NJ tree, and clustered with the

other Indian samples via PCA as well (Fig 2A and S4 Fig). This sample was the only one

collected in Panama to fall outside of the Central and South American cluster in the PCA with

the worldwide sample set. For the two samples with Brazilian travel history and one sample with

Indian travel history, genomic data supported the same country of origin as the travel history

information.

The sample with Chinese travel history had a discrepancy between the region of origin

suggested by its travel history information and its genomic data. This sample clusters with the

Central and South American samples on the worldwide PCA instead of with the samples from

China. This sample clustered with the Colombian samples in the PCA conducted with only the

Central and South American samples (Fig 4B). Similarly, on the NJ tree, this sample fell within

the Colombian cluster with 100% bootstrap support along with the four Panamanian samples that

shared zero IBD with other Panamanian samples in the dataset.

Discussion

Panama is on the cusp of eliminating malaria after several decades of intervention (5).

We found extremely low clonal diversity in the Panamanian *P. vivax* population, observing that

the majority of the successfully sequenced samples (47/59) belonged to a single highly related

lineage, CL1. CL1 has persisted throughout Panama for at least a decade, in spite of ongoing

elimination efforts. Sample contamination could not explain this pattern as samples were

collected in two collection periods 10 years apart and extracted, amplified, and sequenced

separately. Our study suggests that the Panamanian *P. vivax* population has been through a

strong bottleneck due to reduced transmission, resulting in the majority of the population

belonging to a single highly related lineage. Several studies of the relationship between *P.*

falciparum genetic diversity and transmission intensity from Senegal (30), Thailand (31, 32),

Colombia (32), have documented that when there is a decline in parasite transmission, there is

an increase in the clonal composition of the population. Furthermore, there is evidence of

persistence and transmission of *P. falciparum* clonal lineages in Colombia (33), Ecuador (34),

and Haiti (35). This study demonstrates a similar relationship in Panama between low

transmission and extremely low genetic diversity of the *P. vivax* population. Almost all CL1

samples (46/47) share IBD > 0.95 with at least one other CL1 sample, suggesting a substantial

fraction of this population is clonal. Interestingly, the CL1 consensus sequence clusters with

samples from Brazil and Peru on PCA, suggesting a greater degree of *P. vivax* population

panmixia across Central and South America than previously understood, and that the population

has become more fractured over time (20). Alternatively, CL1 could have been imported into

Panama at an earlier point in time and has supplanted the ancestral *P. vivax* population in the

country, becoming the predominant lineage. Both scenarios point to the need for further

longitudinal genomic studies of *Plasmodium* parasites to better understand population dynamics

over space and time.

Previous studies have indicated that Panama has focal transmission in indigenous regions

(Comarcas) (5,6,11) Malaria transmission in Panama is increasingly concentrated in the

Comarcas, with the proportion of total malaria cases in Panama reported from Comarcas rising

from 41.8% in 2005 to 90% in 2019 (36). Prior work shows that low transmission can lead to

population structure in part due to variable malaria transmission (31). The finding that CL1 is

distributed ubiquitously throughout Panama is unexpected given the concentration of the malaria

epidemic within spatially separated regions of the country. The geographic distribution of CL1

suggests that parasites have historically moved throughout the country, founding new

populations or supplanting small existing ones. Case investigations and understanding human

movement patterns throughout Panama will be critical to achieving elimination.

This study had some limitations. We were unable to generate high-quality sequencing

data from ~40% of the samples. Fortunately, we were able to extract a small subset of 264

genotypable SNPs from the excluded samples. We used this SNP set to calculate Nei's standard

genetic distance to determine if the excluded samples were genetically distant from CL1 or CL2.

We found that ~57% of the excluded samples were genetically similar to CL1 (S2 Fig). This

result indicates that our assessment of relatedness within the Panamanian *P. vivax* population is

not biased by parasitemia or other factors that could have affected sequencing success. This

finding also suggests that we did not miss additional genetically distinct circulating Panamanian

P. vivax strains by excluding these samples. Seven samples from 2017-2019 exhibited low

genetic distance from both CL1 and CL2. However, these seven samples were missing ~60-80%

of calls even after stringent filtration. The high missing rate in these seven samples potentially

explains the low genetic distance to most other samples in the dataset. Additionally, all samples

that did yield usable sequence data represented almost all localities across Panama. The

exception to this was a group of samples collected near the Panamanian-Costa-Rican border that

we were unable to generate usable sequence data from. However, most ongoing malaria

transmission in Panama occurs East of the Panama Canal, where most of the samples that

generated usable sequencing data were collected (6,11). Due to the geographic sampling

coverage of regions with ongoing malaria transmission, we believe these data are reflective of

the current state of the Panamanian *P. vivax* population. We also lacked geographic collection

data for two of the successfully sequenced samples, and they were excluded from the geographic

analysis. The lack of geographical data is unlikely to bias our conclusions since these samples

came from both different collection periods and regions. The two samples also constitute a small

proportion of samples in the final dataset.

Genomic epidemiology can help to support malaria elimination efforts in Panama in

multiple ways. First, genomic data can help identify genetically distinct cases that may be

imported. Panama sits at the crux of migration paths to the United States, and it is possible that

genetically distinct samples collected in Panama represent imported cases. Integrating travel

history information with genomic data can help solidify the identification of imported cases.

Four samples had patient travel history information, and genomic data supported the presumed

country of origin for three of them. The fourth sample was collected from a subject with travel

history from China. However, it clustered with Colombian samples on the NJ tree and the PCA,

suggesting the infection was likely acquired somewhere in Central or South America, rather than

China.

Second, genomic data will be critical to determine if imported parasites are contributing

to local transmission and/or admixing with the local parasite population. For example, we did not

observe evidence of admixture between the imported samples from India and Brazil and the

samples that comprise CL1, or evidence of onward clonal transmission of the imported samples.

Our data cannot distinguish whether CL2 is a native Panamanian parasite lineage or if it has been

imported from Colombia. However, the four CL2 samples displayed a genomic and

epidemiological pattern consistent with recent local transmission, as all CL2 samples are

virtually identical and were collected from the same municipality in 2019.

Overall, the existence of one main parasite genetic lineage exhibiting no recent evidence

of outcrossing with imported infections suggests that Panama is ripe for the elimination of *P.*

vivax. While case importation remains a threat, the lack of evidence of outcrossing suggests it

may not be sufficient to prevent elimination under present circumstances. Ongoing genomic

surveillance and case containment efforts will be needed, however, to mitigate the risk of

outbreaks resulting from imported cases and reversal of the impressive progress that has been

recently made towards malaria elimination in Panama.

Materials and Methods

Sample collection

We collected 96 *P. vivax* samples from infected consenting volunteers identified through passive

or active surveillance by technicians from the National Department of Vector Control from the

Ministry of Health (MINSA) of Panama. The Gorgas Memorial Institute of Health Studies

Bioethics Committee gave the study ethical approval (Permits: DIDETEC/06/ET-PA/02 and

154/CBI/ICGES/17). Two groups of DNA samples from infected patients were used in this

study: 1) 56 DNA samples collected during 2007-2009 and 2) 40 DNA samples collected during

2017-2019. The Gorgas Memorial Institutional Animal Use and Care Committee (CIUCAL)

(Permit: 002/CIUCAL-ICCES-2012) approved the use of Aotus *P. vivax* AMRU-1 and SAL-1

infected blood samples as a source of control DNA. Patient blood samples were collected by

finger-prick with a lancet and spotted into EBF 903 Five Spot Blood Cards (Eastern Business

Forms, INC, SC, USA). The samples were then transported at ambient temperature to the

laboratory and stored at –20℃ until processing. We also created thin and thick blood smears

from patient samples. The blood smears were stained with Giemsa for percent parasite density

determination, species identification, and stage differential counts. Each volunteer donated ~150

µL of blood.

Information survey

We collected demographic, geographic, socioeconomic, and epidemiological info from each

study subject using an epidemiological form developed for the Survey123 for ArcGIS online

survey program (Esri, Redlands, CA).

Malaria microscopy

Giemsa stained thick and thin blood smears were examined by light microscopy for parasite

density determinations, P*lasmodium* species confirmation, and parasite lifecycle stage count.

Parasite densities were calculated by quantifying the number of malaria-infected red blood cells

(iRBCs) among 500 – 2000 RBCs on a thin blood smear and expressing the result as %

parasitemia (% parasitemia = parasitized RBCs /total RBCs) x 100).

DNA extraction:

We extracted DNA from the filter paper blood spots using the Chelex method as described (16)

for samples obtained during 2007-2009 and with the Qiagen® DNA mini kit for samples

obtained during 2017-2019.

Molecular confirmation of *P. vivax* infection

We confirmed *P. vivax* infection for all samples collected during 2017-2019 by amplification of

the *P. vivax* PVX_18SrRNA gene using a qRT-PCR assay as described (17).

Selective whole genome amplification and sequencing

We carried out DNA pre-amplification as described (14). Briefly, the thermocycler was

preheated to 35℃. We dispensed aliquots of 37μl of Power SYBR Green Master Mix, plus 3μl

phi39 into each PCR tube, next adding DNA and water to achieve a final volume of 47μl.

Thermocycler settings were as follows: 35℃ x 10 min; 34℃ x 10 min; 33℃ x 10 min; 32℃ x 10

min; 31℃ x 10 min 30℃ x 16 hours, 65℃ x 10 min, 4 ℃ for infinity. SWGA reaction products

were diluted with 50 μl of water. We purified 50 μl of the diluted product using 50 μl AmPURE

beads according to the instructions of the manufacturer. We then eluted beads in 30 μl of water.

Approximately 60-120 ng/μl of DNA was obtained after bead purification of the SWGA

reaction. We measured DNA concentration using Nanodrop quantitation.

Whole-genome sequencing

We performed whole-genome sequencing (WGS) on all 96 *P. vivax* samples using Nextera

libraries and an Illumina HiSeq X platform. Sample reads were aligned to the P01 reference

genome assembly using BWA-MEM, version 0.7 (18). Illumina sequencing reads are available

through the NCBI Sequence Read Archive with BioProject accession numbers

SAMN15722613–SAMN15722671.

SNP Discovery and Quality Filtering

We marked duplicate reads using the MarkDuplicates tool from Picard tools. We next

performed local realignment around indels using the Genome Analysis Toolkit (GATK)

RealignerTargetCreator and GATK IndelRealigner (GATK Version 3.5.0). We called variants

using GATK HaplotypeCaller using best practices to call and filter SNPs and generate gVCFs

for each sample. We called samples in two batches, one containing samples collected in 2007-

2009 and one containing samples collected in 2017-2019. We performed joint variant calling on

the sets separately using GATK GenotypeGVCFs tool with GATK hard filters, including calls in

subtelomeric regions. The resulting VCFs consisted of 56 samples and 407,554 sites for the

2007-2009 samples, and 40 samples and 171,433 variants for the 2017-2019 samples. We

retained samples for analysis if they exhibited a minimum mean read depth of five and had calls

at more than 80% of variant sites in the VCF corresponding to their collection period, including

those in subtelomeric regions. We calculated and evaluated data quality measures using the

VCFtools package and custom R scripts (19). Thirty-five samples from 2007-2009 and 24

samples from 2017-2019 passed these filters and were kept for further analysis.

We next used GenotypeGVCFs tool to construct a joint VCF with the 59 Panamanian

samples plus a collection of previously collected global samples (Bioproject numbers

PRJNA240356-PRJNA240533 (20)). The joint VCF contained 168 samples and 2,4250,245

variants. We filtered sites on the basis of quality (GQ > 40), passing VQSR truth sensitivity level

of 0.99 or greater, missing rate (having a call at that site in > 85% of samples). We also

excluded any sites that were not bi-allelic and indels. The joint VCF generated after filtering

contained 168 samples and 62,211 sites.

Lastly, we generated a VCF containing SNPs found jointly in 80% of both the 2007-2009

and 2017-2019 samples. We also filtered sites in this VCF by excluding non-biallelic SNPs and

on the basis of quality (GQ > 30), and passing GATK filters. The resultant VCFs contained 56

samples and 2335 SNPs for the 2007-2009 samples, and 40 samples and 1301 SNPs for the

2017-2019 samples. For these samples, we generated a highly filtered variant set containing

biallelic SNPs that passed the GATK filters (GQ > 30, truth sensitivity level > 0.99, Mean DP >

five) and were called in at least 80% of the samples from both time periods (2007-2009; 2017-

2019). Calls from the two sample sets were merged to create a unified VCF of 96 samples and

264 genotyped SNPs.

Determination of Sample Clonality

We estimated sample clonality using the F_{ws} statistic. F_{ws} measures the within-sample

genetic diversity (measured by heterozygosity H_w) relative to the overall population genetic

diversity (H_s) (21). The underlying theory assumes that a monoclonal (single strain) infection has

extremely low genetic diversity relative to overall population genetic diversity. By contrast, a

polyclonal (multiple strain) infection has high diversity relative to overall population diversity

(compared to a monoclonal infection). By estimating the ratio between within-host diversity and

population diversity, we can distinguish between monoclonal and polyclonal infections (21). A

sample with an F_{ws} statistic of 0.95 or greater (>=0.95) is considered monoclonal. We calculated

F_{ws} using the R package moimix (22).

Analysis of Recent Common Ancestry

We used hmmIBD (23) to estimate the proportion of sites identical by descent (IBD) between

sample pairs to ascertain recent common ancestry among Panamanian and Colombian samples

collected previously from the global *P. vivax* population study (20). We estimated minor allele

frequency (MAF) for IBD inference using the genetically distinct Panamanian samples, a

representative sample from each of the two highly related Panamanian clusters, and the

Colombian samples. We included Colombian samples to improve MAF estimation given the

greater clonal diversity of the Colombian parasite population and presumed historical gene flow

with Panama. We subsetted the master VCF file to keep only samples collected in Panama and

Colombia. Sites were excluded sites on the basis of minimum and maximum read depth (five and

thirty respectively) to ensure that we were using only high-quality SNPs. The input VCF for

hmmIBD contained 89 samples (59 Panamanian samples and 30 Colombian samples) and 15,788

variant sites. We then re-formatted the data using a custom perl script for input into hmmIBD

along with the MAF estimates. We conducted analysis and visualization of the hmmIBD output

using custom R scripts.

Analysis of Population Structure

We employed principal components analysis (PCA) and a neighbor-joining tree to study

the population structure of Panamanian samples in the context of the worldwide *P. vivax*

population (20). We used a strictly filtered SNP set for PCA, keeping only variants with calls in

at least 95% of samples. This input VCF consisted of 168 samples and 2,428 variants. We used

the R package SNPRelate to conduct PCA (24). Covariation within the two clusters heavily

influenced the PCA of all samples, so we also performed PCA using a single consensus sequence

for each cluster.

We used the R packages ape, StAMPP, pegas, and adegenet, (25–28) to generate the

neighbor-joining tree and genetic distance statistics. First, we calculated Nei's distance for all

pairwise sample combinations using the master VCF consisting of 168 samples and 62,211 sites

to generate a distance matrix. The distance matrix was used to generate a tree. We used the

bootphylo function in the ape package to bootstrap the dataset 100 times to estimate nodal

support. We then visualized the final tree with support values using the FigTree program

(29). We used R software (R version 3.6.1) to carry out statistical analysis and data

visualization.

Acknowledgments

The authors thank Selina Bopp and Becky Kuzma at HSPH for conducting SWGA of the *P.*
vivax samples. We also thank at ICGES, Néstor Sosa, Director, José Calzada, Azael Saldaña and
Gladys Tuñón, for their administrative support; also at ICGES Marlon Núñez, Ariel Magallón,
and José C. Marín for their technical and logistic support; at the University of California-San
Diego, Elizabeth Winzeler and Annie Cowell for their technical advice on establishing SWGA
for *P. vivax*. Department of Research and Development (I+D), Secretary of Science,
Technology, and Innovation (SENACYT) of the Government of the Republic of Panamá, Iriela
Aguilar, and Milagros Mainieri for their administrative and technical support with project
management.

References

1. World Health Organization. World Malaria Report 2019. Vol. 1. WHO; 2019.

2. Price RN, Tjitra E, Guerra CA, Yeung S, White NJ, Anstey NM. Vivax malaria: neglected and not benign. Am J Trop Med Hyg. 2007 Dec;77(6 Suppl):79–87.

3. Arevalo-Herrera M, Quiñones ML, Guerra C, Céspedes N, Giron S, Ahumada M, et al. Malaria in selected non-Amazonian countries of Latin America. Acta Trop. 2012 Mar;121(3):303–14.

4. Obaldia N, Baro NK, Calzada JE, Santamaria AM, Daniels R, Wong W, et al. Clonal outbreak of Plasmodium falciparum infection in eastern Panama. J Infect Dis. 2015 Apr 1;211(7):1087–96.

5. Carrera LC, Victoria C, Ramirez JL, Jackman C, Calzada JE, Torres R. Study of the epidemiological behavior of malaria in the Darien Region, Panama. 2015–2017. PLOS ONE. 2019 Nov 15;14(11):e0224508.

6. Calzada JE, Marquez R, Rigg C, Victoria C, De La Cruz M, Chaves LF, et al. Characterization of a recent malaria outbreak in the autonomous indigenous region of Guna Yala, Panama. Malar J. 2015 Nov 17;14:459.

7. World Health Organization. Panama Malaria Country Profile 2017. WHO Malaria Country Profiles [Internet]. 2017;1(1). Available from: https://www.who.int/malaria/publications/country-profiles/profile_pan_en.pdf?ua=1

8. Ruktanonchai NW, DeLeenheer P, Tatem AJ, Alegana VA, Caughlin TT, Zu Erbach-Schoenberg E, et al. Identifying Malaria Transmission Foci for Elimination Using Human Mobility Data. PLoS Comput Biol. 2016 Apr;12(4):e1004846.

9. Marshall JM, Touré M, Ouédraogo AL, Ndhlovu M, Kiware SS, Rezai A, et al. Key traveller groups of relevance to spatial malaria transmission: a survey of movement patterns in four sub-Saharan African countries. Malar J. 2016 Apr 12;15:200.

10. Wesolowski A, Taylor AR, Chang H-H, Verity R, Tessema S, Bailey JA, et al. Mapping malaria by combining parasite genomic and epidemiologic data. BMC Medicine. 2018 Oct 18;16(1):190.

11. Lainhart W, Dutari LC, Rovira JR, Sucupira IMC, Póvoa MM, Conn JE, et al. Epidemic and Non-Epidemic Hot Spots of Malaria Transmission Occur in Indigenous Comarcas of Panama. PLoS Negl Trop Dis . 2016 May 16;10(5). doi: 10.1371/journal.pntd.0004718

12. José Arcia. Darién, convertido en un río imparable de migrantes irregulares [Internet]. La Estrella de Panamá. 2019 [cited 2020 Apr 29]. Available from: https://www.laestrella.com.pa/nacional/190709/rio-darien-imparable-migrantes-convertido

13. Neafsey DE, Volkman SK. Malaria Genomics in the Era of Eradication. Cold Spring Harb Perspect Med. 2017 Aug;7(8):a025544.

14. Cowell AN, Loy DE, Sundararaman SA, Valdivia H, Fisch K, Lescano AG, et al. Selective Whole-Genome Amplification Is a Robust Method That Enables Scalable Whole-Genome Sequencing of Plasmodium vivax from Unprocessed Clinical Samples. mBio. 2017 07;8(1).

15. Bright AT, Tewhey R, Abeles S, Chuquiyauri R, Llanos-Cuentas A, Ferreira MU, et al. Whole genome (1)sequencing analysis of Plasmodium vivax using whole genome capture. BMC Genomics. 2012 Jun 21;13(1):262.

16. Wooden J, Kyes S, Sibley CH. PCR and strain identification in Plasmodium falciparum. Parasitol Today (Regul Ed). 1993 Aug;9(8):303–5.

17. Obaldia N, Meibalan E, Sa JM, Ma S, Clark MA, Mejia P, et al. Bone Marrow Is a Major Parasite Reservoir in Plasmodium vivax Infection. mBio. 2018 08;9(3).

18. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009 Jul 15;25(14):1754–60.

19. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call format and VCFtools. Bioinformatics. 2011 Aug 1;27(15):2156–8.

20. Hupalo DN, Luo Z, Melnikov A, Sutton PL, Rogov P, Escalante A, et al. Population genomics studies identify signatures of global dispersal and drug resistance in Plasmodium vivax. Nat Genet. 2016;48(8):953–8.

21. Auburn S, Campino S, Miotto O, Djimde AA, Zongo I, Manske M, et al. Characterization of Within-Host Plasmodium falciparum Diversity Using Next-Generation Sequence Data. PLOS ONE. 2012 Feb 29;7(2):e32891.

22. Lee S, Harrison A, Tessier N, Tavul L, Miotto O, Siba P, Kwiatkowski D, Müller I, Barry AE and Bahlo M. Assessing clonality in malaria parasites using massively parallel sequencing data. in preparation. 2016;

23. Schaffner SF, Taylor AR, Wong W, Wirth DF, Neafsey DE. hmmIBD: software to infer pairwise identity by descent between haploid genotypes. Malar J. 2018 May 15;17(1):196.

24. Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. A high-performance computing toolset for relatedness and principal component analysis of SNP data. Bioinformatics. 2012 Dec 1;28(24):3326–8.

25. Pembleton LW, Cogan NOI, Forster JW. StAMPP: an R package for calculation of genetic differentiation and structure of mixed-ploidy level populations. Molecular Ecology Resources. 2013;13(5):946–52.

26. Paradis E. pegas: an R package for population genetics with an integrated-modular approach. Bioinformatics. 2010 Feb 1;26(3):419–20.

27. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics. 2019 Feb 1;35(3):526–8.

28. Jombart T, Ahmed I. adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. Bioinformatics. 2011 Nov 1;27(21):3070–1.

29. Rambaut A, Drummond AJ. FigTree version 1.4. 0. 2012.

30. Rougeron V, Elguero E, Arnathau C, Acuña Hidalgo B, Durand P, Houze S, et al. Human Plasmodium vivax diversity, population structure and evolutionary origin. PLoS Negl Trop Dis . 2020 Mar 9 ;14(3). doi: 10.1371/journal.pntd.0008072

31. Nkhoma SC, Nair S, Al-Saai S, Ashley E, McGready R, Phyo AP, et al. Population genetic correlates of declining transmission in a human pathogen. Mol Ecol. 2013 Jan;22(2):273–85.

32. Manske M, Miotto O, Campino S, Auburn S, Almagro-Garcia J, Maslen G, et al. Analysis of Plasmodium falciparum diversity in natural infections by deep sequencing. Nature. 2012 Jul 19;487(7407):375–9.

33. Echeverry DF, Nair S, Osorio L, Menon S, Murillo C, Anderson TJ. Long term persistence of clonal malaria parasite Plasmodium falciparum lineages in the Colombian Pacific region. BMC Genetics. 2013 Jan 7;14(1):2.

34. Sáenz FE, Morton LC, Okoth SA, Valenzuela G, Vera-Arias CA, Vélez-Álvarez E, et al. Clonal population expansion in an outbreak of Plasmodium falciparum on the northwest coast of Ecuador. Malar J. 2015 Dec 10;14. doi: 10.1186/s12936-015-1019-2

35. Charles M, Das S, Daniels R, Kirkman L, Delva GG, Destine R, et al. Plasmodium falciparum K76T pfcrt Gene Mutations and Parasite Population Structure, Haiti, 2006–2009. Emerg Infect Dis. 2016 May;22(5):786–93.

542   36.   Hurtado L, Cumbrera A, Rigg C, Perea M, Santamaría AM, Chaves LF, et al. Long-term
543   transmission patterns and public health policies leading to malaria elimination in Panamá.
544   Malar J. 2020 Jul 23 ;19. doi: 10.1186/s12936-020-03329-y
545

Fig 1. Sequencing and Sample Assessment at Variant Sites. A) Distribution of variant site read coverage for each sample stratified by the collection period. Coverage values > 100 were censored for visualization purposes. Samples within the red boxes were kept for analysis. B) Distribution of F_{ws} values for all samples, stratified by the collection period. We interpreted F_{ws} values > 0.95 as evidence of sample monoclonality.

Fig 2. IBD analysis of the Panamanian samples. A) The distribution of pairwise IBD estimates among the Panamanian samples. IBD values near zero indicate no recent common ancestry. Values closer to one indicate that the sample pair are clonal or essentially clonal. B) Depicts pairwise IBD estimates for all Panamanian sample pairs with IBD > 0.875 plotted against the mean proportion of high coverage sites (sites with > 5x coverage) in each sample pair. The line indicates a linear regression, the box displays the Pearson correlation coefficient between the two axes variables.

Fig 3. Map of Panamanian sample collection sites. Sample colors show which cluster (or neither) each sample belongs too. Shape indicates the sample collection period. The dotted line shows the location of the Panama Canal. The Blue Line shows the border of the Comarca Kuna Yala. The Darien Jungle Region is indicated by the green shaded area.

Fig 4. Population Structure. A) Neighbor-joining tree for all samples worldwide. Node symbols denote support values: circles indicate 100% support, triangles indicate > 50% support. Branch color indicates the country of collection for each sample. Panamanian samples with travel history are noted with the colored stars. B) PCA of Central and South American samples. Circle color indicates country of collection. Consensus sequences for cluster one and cluster two are noted as a square and triangle respectively. Panamanian samples with travel history are annotated.

Costa Rica

Puerto Obaldia

Panama City

Colombia

- - - - - - - Panama Canal   - - - - - - - Comarca Kuna Yala Boundary

 Darien Jungle Region

Clonal Lineage Grouping

 Cluster 1

 Cluster 2

 Not Part of Either Cluster   **Collection Period** ◯ 2007-2009 ☐ 2017-2019