Chronic hyperglycaemia drives functional impairment of lymphocytes in diabetic *INS*[C94Y] transgenic pigs

Isabella-Maria Giese [1], Simone Renner [2,3,4], Eckhard Wolf [2,3,4,5], Stefanie M. Hauck [4,6], Cornelia A. Deeg [1*]

[1] Chair of Physiology, Department of Veterinary Sciences, LMU Munich, Germany

[2] Chair for Molecular Animal Breeding and Biotechnology, Gene Center and Department of Veterinary Sciences, LMU Munich, Germany

[3] Center for Innovative Medical Models (CiMM), Department of Veterinary Sciences, LMU Munich, Germany

[4] German Center for Diabetes Research (DZD), Neuherberg, Germany

[5] Laboratory for Functional Genome Analysis (LAFUGA), Gene Center, LMU Munich, Germany

[6] Research Unit Protein Science, Helmholtz Center Munich, German Research Center for Environmental Health GmbH, Germany

*Correspondence:
Cornelia A. Deeg:
Cornelia.Deeg@lmu.de

Keywords: Animal model(s) / Annexin A1 / Diabetes mellitus / Cell metabolism / PHA / Proliferation / T cells / Impaired immune cell function

4024 words, 4 figures, 2 tables

Abstract

People with diabetes mellitus have an increased risk for infections, however, there is still a critical gap in precise knowledge about altered immune mechanisms in this disease. Since diabetic *INS*[C94Y] transgenic pigs exhibit elevated blood glucose and a stable diabetic phenotype soon after birth, they provide a favourable model to explore functional alterations of immune cells in an early stage of diabetes mellitus *in vivo*. Hence, we investigated peripheral blood mononuclear cells (PBMC) of these diabetic pigs compared to non-transgenic wild-type littermates. We found a 5-fold decreased proliferative response of T cells in *INS*[C94Y] tg pigs to polyclonal T cell mitogen phytohaemagglutinin (PHA). Using label-free LC-MS/MS, a total of 2,704 proteins were quantified, and distinct changes in protein abundances in CD4[+] T cells of early-stage diabetic pigs were detectable. Additionally, we found significant increases in mitochondrial oxygen consumption rate (OCR) and higher basal glycolytic activity in PBMC of diabetic *INS*[C94Y] tg pigs, indicating an altered metabolic immune cell phenotype in diabetics. Thus, our study provides new insights into molecular mechanisms of dysregulated immune cells triggered by permanent hyperglycaemia.

Introduction

Diabetes mellitus is a major risk factor regarding the outcome of bacterial and viral infections (Harding,
Pavkov, Magliano, Shaw, & Gregg, 2019). Among diabetic patients, increased numbers of highly
susceptible individuals suffering from tuberculosis (Kumar et al., 2019) or influenza (Marshall et al.,
2020) are common. Recently, meta-analyses revealed an enhanced severity and mortality of COVID-
19 infections in people with diabetes mellitus (Y. Hu et al., 2020; Zhou et al., 2020). Undoubtedly,
there is a link between the disease and an impaired immune system, but the molecular mechanisms
involved in altered immune cell function in diabetic patients are still unknown to date. Dysfunctional
T cells may play a pivotal role in immunological impairment in diabetes mellitus (Nicholas et al.,
2019). However, the precise effects of chronic hyperglycaemia on immune cell function are still
uncertain since contradictory findings of both hyperresponsive (Nyambuya, Dludla, Mxinwa, &
Nkambule, 2020) and attenuated (Lau et al., 2019; Lee et al., 2019) T cells were reported in type 2
diabetic patients. While numerous studies linked hyperglycaemia to enhanced T cell activation and
proliferation (Nyambuya et al., 2020), others suggest that an abnormal glucose homeostasis promotes
an insufficient T-cell response via an increased frequency of senescent T cells with proliferative
impairment (Lau et al., 2019; Lee et al., 2019). Thus, it has yet to be determined whether dysfunctional
T cells are impaired or highly activated in diabetes mellitus and whether hyperglycaemia facilitates
immune cell dysregulation. Alterations of cellular mechanisms and molecular networks in disease
states can be analysed by changes in proteome composition (Soongsathitanon, Umsa-Ard, &
Thongboonkerd, 2019). Still little is known about accurate protein profiles of T cells affected by
hyperglycaemia *in vivo*. Therefore, mass spectrometry-based studies are a powerful tool for obtaining
comprehensive information regarding cellular pathology (Lepper et al., 2018) and enable hypothesis-
generating approaches on the protein level to understand the molecular basis of altered proteome
composition and regulation caused by disease. Moreover, recent advances in the field of immuno-
metabolism recognized cellular metabolism as a primary driver and regulator of immune cell function,
leading to a wide variety of functionally different immune cells (Olenchock, Rathmell, & Vander
Heiden, 2017). Since there is still a lack of knowledge about hyperglycaemia-associated alterations of
immune cell metabolic phenotypes, there is an urgent need to understand whether and how high glucose
levels affect the metabolic phenotype of the cells and thereby change their immune cell function
(Olenchock et al., 2017). Diabetic *INS*^C94Y transgenic (tg) pigs were generated as a large animal model
of permanent neonatal diabetes mellitus (Renner et al., 2013). These pigs are characterized by impaired
insulin secretion with consecutive hypoinsulinaemia and increased fasting blood glucose levels
(Renner et al., 2013). Early on, *INS*^C94Y tg pigs show a stable diabetic phenotype and mirror several
disease-associated alterations of diabetes mellitus as seen in humans such as cataract, retinopathy,
impaired myocardial function and regeneration (Hinkel et al., 2017; Kleinwort et al., 2017). Since early
stages of disease in diabetic patients are often unperceived for a long time, initial immunological
alterations are difficult to observe due to the latent disease process. Thus, this diabetic pig model
enables the exploration of deviant immune cell function and underlying mechanisms in an early stage
of diabetes mellitus with hyperglycaemia and hypoinsulinaemia, offering valuable insights into disease
pathogenesis. Hence, we used PBMC of young *INS*^C94Y tg pigs and non-transgenic wild-type
littermates to investigate the impact of a permanent early-life hyperglycaemic condition on immune
cell function.

Materials and methods

Animal model and sample preparation

In this study, PBMC of 23 sex-matched diabetic *INS*[C94Y] tg pigs and 31 non-transgenic wild-type littermates at the age of 12 weeks were used. *INS*[C94Y] tg pigs were generated previously as described and exhibited significantly elevated random blood glucose levels within 24 h after birth compared with their non-transgenic wild-type littermates (Renner et al., 2013). All animals were housed under controlled conditions, had free access to water and were fed a commercial diet. For sampling, pigs were fasted overnight and heparinized venous whole blood was collected. PBMC were isolated by density gradient centrifugation (RT, 500 x g, 25 min, brake off) with Pancoll separating solution (PAN-Biotech, Aidenbach, Germany), and restored in PBS (pH 7.4) or RPMI medium (PAN-Biotech, Aidenbach, Germany), supplemented with 10% heat-inactivated foetal calf serum (FCS) and 1% penicillin/streptomycin (both Biochrom, Berlin, Germany). Blood withdrawal was performed according to the German Animal Welfare Act with permission from the responsible authority (Government of Upper Bavaria), following the ARRIVE guidelines and Directive 2010/63/EU.

Lymphocyte stimulation by mitogens

In addition to unstimulated controls, PBMC were either stimulated by pokeweed mitogen (PWM; 1µg/ml), concanavalin A (ConA; 1µg/ml) or phytohaemagglutinin (PHA; 1µg/ml). After an incubation period of 32 hours at 37°C, cells were incorporated with ^{3}H-thymidine (Perkin Elmer, Hamburg, Germany) and incubated for 16 more hours. After harvesting, ^{3}H-thymidine incorporation was quantified by detecting counts per minute (cpm), using a Microbeta (Perkin Elmer, Hamburg, Germany). Proliferation rate was expressed as factor of ^{3}H-thymidine incorporation with respect to unstimulated cells.

Flow cytometry

Staining of 2×10^5 cells per well was performed with mouse anti-human CD79α (clone HM57, Bio-Rad AbD Serotec, Puchheim, Germany, 1:100) for identification of B cells and Alexa Fluor 647-conjugated rat anti-human CD3ε (clone CD3-12, 1:200; cross-reactive to pig (Czajka et al., 2015)) for identification of T cells. We used FITC-conjugated mouse anti-pig CD4 alpha (clone MIL17, Bio-Rad AbD Serotec, Puchheim, Germany, 1:20) and Alexa Fluor 647-conjugated mouse anti-pig CD8α (clone 76-2-11, Becton Dickinson, Heidelberg, Germany, 1:400) to identify αβT cells and mouse anti-pig SWC5 (clone b37c10, Bio-Rad AbD Serotec, Puchheim, Germany, 1:100) for identification of γδT cell subpopulation. If necessary, a secondary antibody was used (Alexa Fluor 647-conjugated goat F(ab')2 anti-mouse IgG (Fc), Dianova, Hamburg, Germany, 1:1000). Analyses were performed with MACSQuant Analyzer 10 and Flowlogic Software (both Miltenyi Biotec, Bergisch Gladbach, Germany).

Magnetic activated cell sorting (MACS) for CD4[+] T cells

Briefly, a total of 6×10^7 cells was incubated in staining buffer with mouse anti-pig CD4 alpha (clone MIL17, Bio-Rad AbD Serotec, Puchheim, Germany, 1:50) at 4°C for 20 min. Staining buffer contained phosphate-buffered saline (pH 7.2) and was supplemented with 2 mM EDTA and 0.5% bovine serum albumin (BSA). In the next step, cells were resuspended in 480 µL staining buffer before adding 120 µL anti-mouse IgG$_{2a/b}$ MicroBeads (Miltenyi Biotec, Bergisch Gladbach, Germany) for an incubation period of 15 min. In further steps, BSA in staining buffer was omitted to prevent interference with mass

spectrometry. Magnetic separation was performed using LS columns (Miltenyi Biotec, Bergisch
Gladbach, Germany). Magnetically-labelled $CD4^+$ T cells were retained in the magnetic field, while
unwanted cells were eliminated by three washing steps. Positive $CD4^+$ T cell fraction was eluted by
removing the column from magnetic field and flushing with staining buffer. 6×10^5 positive selected
cells were pelleted and stored at -20 °C until filter-aided sample preparation (FASP). The isolation of
porcine $CD4^+$ T cell routinely achieved >90% purity, confirmed by flow cytometry.

Mass spectrometry and data analysis

Purified $CD4^+$ T cells of four diabetic *INS*^{C94Y} tg pigs and five littermate wildtypes were analyzed.
$CD4^+$ T cell pellets were lysed in urea buffer (8M in 0.1M Tris/HCl pH 8.5) and 10 µg total protein of
each sample was proteolysed with LysC and trypsin by a modified filter-aided sample preparation
(FASP) as described (Grosche et al., 2016). Acidified eluted peptides were analyzed in the data-
independent acquisition mode on a Q Exactive HF-X mass spectrometer (Thermo Fisher Scientific,
Waltham, MA, USA) online coupled to an ultra-high-performance liquid chromatography (UHPLC)
system (Ultimate 3000, Thermo Fisher Scientific). The spectral library was generated directly in
Spectronaut Pulsar X (Biognosys, Schlieren, Switzerland; version 12.0.20491.17.25792) as described
(Singh et al., 2020). Spectronaut was equipped with the Ensembl Pig Database (Release 75
(Sscrofa10.2), 25,859 sequences, https://www.ensembl.org). Peptide identification was filtered to
satisfy an FDR of 1% by the mProphet approach with q-value cut-off at 0.05 (Reiter et al., 2011).
Statistical analysis was performed on log2 transformed normalized abundance values using Student's
t-test. All proteins with $p < 0.05$ were considered significantly different and proteins with *INS*^{C94Y}/wt
ratio ≥ 2.0 were determined as differentially abundant. The heatmap of hierarchical cluster analysis
was created with open source software Cluster 3.0 and was illustrated via Java TreeView (version
1.1.6r4, http://jtreeview.sourceforge.net). GraphPad Prism Software (version 5.04) was used to design
Volcano plot, and pathway enrichment analysis was done with open source software Reactome
(Pathway Browser version 3.6, https://reactome.org).

Quantification of ANXA1 by immunofluorescence staining

To quantify ANXA1 in porcine PBMC, flow cytometric analysis was performed using rabbit anti-
human ANXA1 (Thermo Fisher Scienific, Ulm, Germany, 1:100) and Alexa Fluor 647-conjugated
goat anti-rabbit IgG H+L (Invitrogen, Karlsruhe, Germany; 1:500). Cross reactivity of anti-human
ANXA1 was proved and high sequence homology of porcine ANXA1 was confirmed via BLASTP
(https://blast.ncbi.nlm.nih.gov/Blast.cgi). For intracellular staining, PBMC were permeabilized (BD
Cytofix/Cytoperm, Becton Dickinson, Heidelberg, Germany) for 20 min at 4°C and washed with
washing buffer (BD Perm/Wash, Becton Dickinson, Heidelberg, Germany), diluted in PBS 1:10.
For immunofluorescence staining, purified $CD4^+$ T cells of wild-type and *INS*^{C94Y} tg pigs were
incubated with FITC-conjugated anti-pig CD4 alpha before staining of ANXA1. After fixation with 1
% PFA, cell nuclei were counterstained with 4′,6-diamidino-2-phenylindole (DAPI; Invitrogen,
Karlsruhe, Germany, 1:100) for 30 min at RT. 5×10^4 cells were transferred to microscope slides and
centrifugated (300 x g, 10 min) before coverslip using mounting medium (Serva, Rosenheim,
Germany). Visualization of stained targets was performed using a Leica Dmi8 microscope with
associated LAS-X-software (Leica, Wetzlar, Germany).

Measurement of OCR and ECAR by Seahorse XFe Analyzer

Metabolic phenotypes of porcine PBMC were determined using a Seahorse XFe Analyzer (Agilent
Technologies, Waldbronn, Germany) measuring oxygen consumption rate (OCR), which is attributed

to mitochondrial respiration and extracellular acidification rate (ECAR), which can be related to
glycolysis (van der Windt, Chang, & Pearce, 2016). In accordance with the manufacturer's instructions,
sterile XF assay buffer (Seahorse XF RPMI Medium supplemented with 10mM glucose, 2mM L-
glutamine, and 1mM pyruvate, pH 7.4; Agilent Technologies, Waldbronn, Germany) was used for
experiments. Prior to the start of the assay, sensor cartridges (Agilent Technologies, Waldbronn,
Germany) were prepared adding oligomycin, FCCP and rotenone & antimycin A. A total of 1×10^6
PBMC was seeded in 24-well XF24 cell culture microplates (Agilent Technologies, Waldbronn,
Germany), while four wells were kept free from cells as background correction. Baseline OCR and
ECAR were measured before adding oligomycin, FCCP and rotenone & antimycin A. OCR was
reported in units of pmol/minute and ECAR in mpH/minute.

Statistical Analysis

Kolmogorow-Smirnov (KS) test was performed for determination of Gaussian distribution. If KS test
indicated $p < 0.05$ (no normal distribution), Mann-Whitney U test was used for statistical analysis,
while student's *t*-test was used if KS test was $p > 0.05$ (normal distribution). In both tests, statistical
probabilities were considered significant at $p < 0.05$. Significances are indicated by asterisks with *p
< 0.05, **$p < 0.01$ and ***$p < 0.001$.

Results

Reduced proliferative capacity of lymphocytes in diabetic *INS*^{C94Y} transgenic pigs after
polyclonal stimulation with PHA

For assessment of the proliferation response *in vitro*, we used B- and T cell mitogen PWM (Nam et al.,
2019) and the specific T cell mitogens ConA (Rodríguez-Gómez et al., 2019) and PHA (Lin, Chen,
Liao, Lee, & Chien, 2016) to stimulate lymphocytes of diabetic *INS*^{C94Y} tg pigs and wild-type
littermates. Interestingly, while no differences could be observed between the two groups after
stimulation with PWM (Fig 1a) and ConA (Fig 1b), lymphocytes of diabetic pigs (n=11) revealed a
significantly 5-fold decreased proliferation rate in response to PHA compared to wildtypes (n=13) (Fig
1c; ***$p \leq 0.001$).

Lymphocyte subpopulation ratio did not differ between wildtypes and diabetic pigs

Next, we examined whether the decreased ability to proliferate correlated with altered lymphocyte
subpopulations in diabetic *INS*^{C94Y} tg pigs. There were no significant differences in the percentage of
B or T cell population between wildtypes and diabetic pigs (Supplementary Fig 1a, b). Moreover,
measurements of T cell subsets CD4⁺ (Supplementary Fig 1c), CD8α⁺ (Supplementary Fig 1d),
CD4⁺CD8α⁺ αβT cells (Supplementary Fig 1e) and SWC5⁺ γδT cells (Supplementary Fig 1f) revealed
no differences between both groups.

Proteomic profiling identified disease-associated protein ANXA1 in CD4⁺ T cells of diabetic
INS^{C94Y} transgenic pigs

We characterized the proteomes of porcine CD4⁺ T cells, which we hypothesized to be the key drivers
of impaired proliferation response. Purified CD4⁺ T cells were examined with differential proteome
analyses, using data-independent acquisition LC−MS/MS. A high-resolution proteome with 2,704
identified proteins was obtained (quantified with at least two unique peptides; Supplementary Table

1), with 80 proteins significantly different in abundance (*p < 0.05; Fig 2a). Two of these proteins had
a ≥ 2-fold difference in abundance, named MYL4 and ANXA1 (Fig 2b). In the whole sample set,
ANXA1 was highlighted with highest statistical significance in protein abundance differences between
wildtypes and diabetic pigs (***p < 0.001). Interestingly, higher levels of ANXA1 were also already
described in serum and plasma of type 1 (Purvis et al., 2018) and II diabetes mellitus patients (Gareth
S. D. Purvis et al., 2019). Therefore, we subsequently focused on ANXA1 to get further information.

ANXA1 was higher abundant at CD4[+]T cell outer cell membranes of diabetic *INS*[C94Y]
transgenic pigs

Flow cytometric analyses confirmed that ANXA1 expression was significantly increased in CD4[+] T
cells of diabetic *INS*[C94Y] tg pigs (n=9) compared to wildtypes (n=11) (*p ≤ 0.05; Fig 3 a-c).
Furthermore, we analysed ANXA1 subcellular localization in purified CD4[+] T cells with
immunocytology and flow cytometry. While intracellular ANXA1 levels were equivalent between both
groups (Fig 3 d-f; n= 4 per group), ANXA1 was significantly enhanced at the outer cell membrane of
CD4[+] T cells from diabetic pigs (*p<0.05; Fig 3 g-i; n= 6 per group).

Pathway enrichment analyses of CD4[+] T cells highlight lipophagy pathway in the diabetic
condition

To detect altered biological pathways in diabetic immune cells, we performed pathway enrichment
analyses via the Reactome database (Jassal et al., 2020). The most significant pathways enriched in
CD4[+] T cells of wildtypes were "Lagging strand synthesis", "Gap-filling DNA repair synthesis and
ligation in GG-NER" and "Polymerase switching" (Table 1). In contrast, proteins with significantly
higher abundances in CD4[+] T cells of *INS*[C94Y] tg pigs were associated with the pathways "Downstream
signal transduction", "RUNX1 regulates transcription of genes involved in differentiation of myeloid
cells" and "Lipophagy" (Table 2). Since lipophagy is linked to metabolic dysfunction of immune cells
(Kounakis, Chaniotakis, Markaki, & Tavernarakis, 2019; Tian et al., 2019), and it was recently shown
that functional different cell subtypes can be identified solely by their metabolic phenotypes
(Olenchock et al., 2017), we next analysed metabolic profiles of the T cells.

Increased basal glycolytic activity and mitochondrial respiration in PBMC of diabetic pigs

Interestingly, the basal glycolytic rate of PBMC from diabetic pigs was significantly higher compared
to wild-type PBMC (Fig 4a; *p<0.05), indicating increased lactate production and glycolytic activity
of diabetic PBMC. Furthermore, analyses of OCR revealed a significant increase in oxygen
consumption at all measured time points in diabetic pigs. Compared to wild-type PBMC, a rise of
basal, ATP-linked and maximal mitochondrial respiration (**p<0.01) as well as spare respiratory
capacity (*p<0.05) was present in PBMC from diabetic pigs (Fig 4b). These differences indicated a
fundamentally different metabolic phenotype of PBMC from wildtypes (n=15) and diabetic *INS*[C94Y] tg
pigs (n=10).

Discussion

To assess the impact of permanent hyperglycaemia on immune cell function *in vivo*, we characterised
lymphocytes of a transgenic pig model for permanent neonatal diabetes mellitus. Significant
differences in proliferative response, CD4$^+$ T cell proteomes and the metabolic phenotypes were
already detectable in these early diabetic pigs. Our findings point to main distinct changes in the T cell
compartment triggered by early-life high blood glucose and indicate that several immune alterations
already occur in the early stage of diabetes mellitus.
We observed no differences of lymphocyte proliferation between controls and diabetic pigs after
polyclonal stimulation with B and T cell mitogen PWM and T cell mitogen ConA. However, after
polyclonal stimulation with T cell mitogen PHA, we detected a distinct decreased proliferation of
lymphocytes from diabetic pigs, which points to an impaired immune response selectively induced by
this mitogen. Interestingly, a decreased response of T cells to PHA-stimulation was already
demonstrated in diabetic patients (Chang & Shaio, 1995; Richard et al., 2017). The significantly
reduced proliferation to PHA was correlated with a decreased interleukin (IL)-2 production (Richard
et al., 2017). The cytokine IL-2, as well as its IL-2 receptor, is a cytokine crucial for proliferation
processes in human T cells (Abbas, Trotta, D, Marson, & Bluestone, 2018) and, thus, is enhanced in
these cells after PHA stimulation leading to cell proliferation (S. Hu et al., 2019). These findings were
supported and extended by another transcriptomic study of PHA-activated human CD4$^+$ and CD8$^+$ T
cells with decreased gene expression of IL-2 and IL-2 receptor in diabetic type 2 patients (Stentz &
Kitabchi, 2007). Furthermore, T cells from these patients exhibited differences in expression of
additional genes and gene products like lower expression of genes of insulin signalling pathway and
enzymes of the glycolytic pathway (Stentz & Kitabchi, 2007), indicating that an impaired immune
response in the diabetic context induced by PHA is affected by more than IL-2 signalling. Little is
known about the precise signalling pathways and intracellular processes stimulated by PHA and ConA
in humans and pigs. However, the impaired immune response selectively induced by PHA in
hyperglycaemic conditions, which is not addressed by polyclonal T cell mitogen ConA (Rodríguez-
Gómez et al., 2019), points to PHA-associated cell signalling pathways specifically altered in the
diabetic condition. In murine T cells, ConA was shown to bind to surface glycoproteins and with high
affinity to the co-stimulatory molecule CD28, leading to enhanced T cell proliferation (Ando et al.,
2014). If mechanisms hold true for porcine T cells, hyperglycaemia does not affect CD28 signalling in
diabetic pigs. In contrast, PHA is able to mimic the stimulation of T cells via crosslinking of T cell
receptor and CD3 complex, which is followed by a sustained calcium influx as an important signal for
T cell proliferation (Lin et al., 2016). In the Jurkat T cell line, high glucose levels were shown to inhibit
T cell activation via an increase in non-enzymatic glycation of concentration-regulating calcium
channels and delayed specific activation of these cells with anti-CD3 monoclonal antibody (Boldizsár,
Berki, Miseta, & Németh, 2002). Since alterations of calcium homeostasis were shown in various cell
types of diabetic patients (Klec, Ziomek, Pichler, Malli, & Graier, 2019), insufficient calcium-mediated
signalling in T cells of diabetic pigs might point to impaired PHA-induced proliferation. Thus, the
immune cell calcium homeostasis of diabetic *INS*[C94Y] tg pigs merits further investigations in future
studies to examine possible correlations to the impaired T cell response in the diabetic cases observed
in our study.
To gain deeper insights into hyperglycaemia-associated molecular changes of CD4$^+$ T cells, we
analysed the proteome of these cells from diabetic pigs and wildtypes as a hypothesis-generating
approach for the dysregulated immune responses. Notably, we obtained a comprehensive proteomic
data set for porcine CD4$^+$ T cells with a high resolution of 2,704 identified proteins. Among these, we
found a total of 3% (80 proteins) with significantly different protein abundances between both groups,
indicating distinct proteome differences in CD4$^+$ T cells from early stage diabetic pigs, functionally
associated to DNA damage responses and cell protection, actin cytoskeleton organization and

regulation of cellular energy metabolism. While many studies of immune cells from diabetic type 2
patients were performed by a 2D-gelelectrophoresis approach, only significantly altered spots were
analysed subsequently with mass spectrometry (Giorgi et al., 2017; Soongsathitanon et al., 2019). To
our knowledge, there is no in-depth proteomic fingerprint of CD4$^+$ T cells in hyperglycaemic
conditions at initial stage of non-immune-mediated cases of diabetes mellitus. Interestingly, the protein
with the highest statistical significance and a 2-fold higher abundance in diabetic pigs was ANXA1.
This protein identification corresponds to findings in type 1 (Purvis et al., 2018) and type 2 (G. S. D.
Purvis, M. Collino, et al., 2019) diabetic patients, where ANXA1 was previously found with increased
levels in serum and plasma, pointing to a pivotal, disease-associated alteration of this protein. However,
the biological consequence of elevated ANXA1 in diabetes mellitus remains unexplained (Purvis,
Solito, & Thiemermann, 2019). Early studies in mice have shown that ANXA1 acts as a molecular T
cell tuner by increasing the strength of T cell receptor signaling and T cell activation (D'Acquisto et
al., 2007). Overexpressed ANXA1 induced the differentiation of naïve T cells to pro-inflammatory
Th1 cells with a high expression of INFγ (Huang, Zhou, Liu, & Zhang, 2016), promoting an
inflammatory milieu. Thus, the higher abundance of ANXA1 in CD4$^+$ T cells of diabetic pigs might
indicate an enhanced T cell activation with functional importance of a pro-inflammatory immune
response. However, also the opposite effect was shown in ANXA1-deficient mouse CD4$^+$ T cells
(Yang et al., 2013). These cells increased their activation, proliferation and inflammation in the absence
of ANXA1, suggesting that this protein rather has an anti-inflammatory role and attenuates an
exacerbated inflammatory response (Yang et al., 2013). One possible reason is that externalized
ANXA1 regulates the extracellular regulated protein kinases (ERK)/mitogen-activated protein kinases
(MAPK) signaling pathway via binding to its formyl peptide receptor (FPR) (Purvis et al., 2018). This
pathway is important for T cell activity and strongly contributes to T cell activation and proliferation
(Liu et al., 2019) and was shown to be activated in hyperglycaemia (Purvis et al., 2018). Treatment of
Anxa1$^{-/-}$ mice with human recombinant ANXA1 attenuated inflammatory processes in tissues by
reducing MAPK signaling (Purvis et al., 2018). Thus, a higher abundance of ANXA1 in diabetic CD4$^+$
T cells may temper signaling to protect the cell from hyperresponsiveness initiated by chronic
hyperglycaemia and, thus, exert anti-inflammatory effects. Although we cannot yet define the precise
role of ANXA1 in CD4$^+$ T cells in the porcine diabetic model, the identification of ANXA1 in humans
and pigs, independent of aetiology, is an important finding which underscores a crucial role of this
protein in T cell impairment accompanying diabetes mellitus. It furthermore highlights the potential
translational quality of this animal model for diabetes research, which is, in our opinion, recommended
for future studies to examine the role of this molecule for the observed dysregulated immune responses.
Interestingly, one of the most over-represented pathways of altered proteins in CD4$^+$ T cells of diabetic
pigs was lipophagy. Lipophagy is linked to metabolic dysfunctions of cells (Kounakis et al., 2019; Tian
et al., 2019), and thus, pointed to an altered metabolic phenotype in diabetic pigs. We therefore
characterised the metabolic immune cell phenotype of PBMC, since cellular metabolism can be a
primary driver and regulator of immune cell function (Olenchock et al., 2017). Metabolic properties of
diabetic PBMC in this study revealed an increase in mitochondrial respiration and basal glycolytic
activity. We found significant increases in mitochondrial oxygen consumption rate (OCR) in PBMC
of diabetic pigs. Interestingly, in line with our findings, increased oxygen consumption was also
demonstrated in PBMC of type 2 diabetes patients (Hartman et al., 2014). Previously, this group also
measured a higher production of reactive oxygen species (ROS) in these cells and hypothesised
mitochondrial damage in diabetic PBMC caused by enhanced oxidative stress (Widlansky et al., 2010).
Since ROS synthesis is essential for physiological immune activation of lymphocytes but lead to
cellular damage if concentration surpasses (Alfatni et al., 2020), this parameter should be determined
in PBMC of diabetic pigs, in order to evaluate whether immune cells are impaired by oxidative cell
stress. On the contrary, mitochondrial respiration with increased oxygen consumption is used by

immune cells to supplement increased glycolysis, which is the main metabolic pathway fueling effector
function upon T cell activation (Gaber, Chen, Krauss, & Buttgereit, 2019). Besides increased oxygen
consumption rates, we also found a higher basal glycolytic activity in early diabetic pigs, suggesting
that hyperglycaemia induced metabolic adaption to local conditions of immune cells and promoted a
more activated status in T cells (Bonacina, Baragetti, Catapano, & Norata, 2019). A rapid increase in
glycolysis is an important determinant in cellular energy production and required for effective cell
proliferation, size expansion, and differentiation (Gaber et al., 2019). Hence, it is also possible that
PBMC in young diabetic pigs are highly metabolically active to challenge hyperglycaemic conditions.
Our findings indicate that metabolic disturbances occurring at early stage diabetes mellitus, as observed
in *INS*^{C94Y} tg pigs, likely cause metabolic reprogramming in PBMC and may also influence their
activation status. Since further studies are absolutely required for a better understanding of bioenergetic
profiles of PBMC in diabetes mellitus, diabetic pigs enable to define mitochondrial properties and
cellular energy production in these cells at initial stage of disease.
Taken together, our findings point to an early, fundamental dysregulation of immune cells when
exposed to prolonged high blood glucose. Chronic hyperglycaemia leads to profound molecular and
functional changes of T cells reasoned by proteomic and metabolic alterations, and, moreover, affects
T cell immune responses *in vitro*. Diabetic pigs qualify for additional translational experiments to
explore the crucial role of altered ANXA1 levels in adaptive immunity in diabetes mellitus. The altered
metabolic properties of immune cells in young diabetic pigs provided novel information on distinct
immune cell metabolism at initial stage of disease. The exact meaning of this significantly altered
metabolic phenotype of the immune cells is not known so far, thus, *INS*^{C94Y} tg pigs are a valuable
source to further analyse respective changes for the diagnosis, or treatment or even prevention of early
stage diabetes mellitus and its immune system-related complications.

Acknowledgements

354 The authors thank F. Stetter, M. Weigand, M. Schilloks and B. Amann from Chair of Physiology, LMU
355 Munich for critical discussions and S. Zettler and A. Hinrichs from Chair for Molecular Animal
356 Breeding and Biotechnology, LMU Munich for assistance in blood withdrawal.

357 This manuscript previously appeared as a preprint on bioRxiv.org (see References).

Funding

359 This work was supported by grants from the Deutsche Forschungsgemeinschaft SPP project 2127
360 [DFG DE 719/7-1 to C.D., HA 6014/5-1 to S.M.H.] and by the German Center for Diabetes Research
361 [82DZD00802 to E.W and S.R.].

Duality of interest

363 The authors declare that there is no duality of interest associated with this manuscript.

Contribution statement

CD conceived and designed the experiments; IMG, SH and CD performed the experiments; IMG and
SH analysed the data; SR and EW developed and characterized the *INS*^{C94Y} tg pig model and
contributed reagents and materials; IMG and CD wrote the manuscript. All coauthors critically read
the manuscript and approved the final version to be published. CD is the guarantor of this work and,

as such, had full access to all the data in the study and takes responsibility for the integrity of the data
and the accuracy of the data analysis.

Data availability

The data are available on request from the authors.

References

Abbas, A. K., Trotta, E., D, R. S., Marson, A., & Bluestone, J. A. (2018). Revisiting IL-2: Biology
and therapeutic prospects. *Sci Immunol, 3*(25). doi:10.1126/sciimmunol.aat1482

Alfatni, A., Riou, M., Charles, A. L., Meyer, A., Barnig, C., Andres, E., et al. (2020). Peripheral
Blood Mononuclear Cells and Platelets Mitochondrial Dysfunction, Oxidative Stress, and
Circulating mtDNA in Cardiovascular Diseases. *J Clin Med, 9*(2). doi:10.3390/jcm9020311

Ando, Y., Yasuoka, C., Mishima, T., Ikematsu, T., Uede, T., Matsunaga, T., et al. (2014).
Concanavalin A-mediated T cell proliferation is regulated by herpes virus entry mediator
costimulatory molecule. *In Vitro Cell Dev Biol Anim, 50*(4), 313-320. doi:10.1007/s11626-
013-9705-2

Boldizsár, F., Berki, T., Miseta, A., & Németh, P. (2002). Effect of hyperglycemia on the basal
cytosolic free calcium level, calcium signal and tyrosine-phosphorylation in human T-cells.
Immunol Lett, 82(1-2), 159-164. doi:10.1016/s0165-2478(02)00032-9

Bonacina, F., Baragetti, A., Catapano, A. L., & Norata, G. D. (2019). The Interconnection Between
Immuno-Metabolism, Diabetes, and CKD. *Curr Diab Rep, 19*(5), 21. doi:10.1007/s11892-
019-1143-4

Chang, F.-Y., & Shaio, M.-F. (1995). Decreased cell-mediated immunity in patients with non-
insulin-dependent diabetes mellitus. *Diabetes Res Clin Pract, 28*(2), 137-146.
doi:https://doi.org/10.1016/0168-8227(95)00168-8

Czajka, A., Ajaz, S., Gnudi, L., Parsade, C. K., Jones, P., Reid, F., et al. (2015). Altered
Mitochondrial Function, Mitochondrial DNA and Reduced Metabolic Flexibility in Patients
With Diabetic Nephropathy. *EBioMedicine, 2*(6), 499-512. doi:10.1016/j.ebiom.2015.04.002

D'Acquisto, F., Merghani, A., Lecona, E., Rosignoli, G., Raza, K., Buckley, C. D., et al. (2007).
Annexin-1 modulates T-cell activation and differentiation. *Blood, 109*(3), 1095-1102.
doi:10.1182/blood-2006-05-022798

Gaber, T., Chen, Y., Krauss, P. L., & Buttgereit, F. (2019). Metabolism of T Lymphocytes in Health
and Disease. *Int Rev Cell Mol Biol, 342*, 95-148. doi:10.1016/bs.ircmb.2018.06.002

Giese, I.-M., Renner, S., Wolf, E., Hauck, S. M., & Deeg, C. A. (2020). Chronic hyperglycaemia
drives functional impairment of lymphocytes in diabetic INSC94Y transgenic pigs [Preprint].
bioRxiv. doi:10.1101/2020.08.26.267914

Giorgi, A., Tempera, I., Napoletani, G., Drovandi, D., Potestà, C., Martire, S., et al. (2017).
Poly(ADP-ribosylated) proteins in mononuclear cells from patients with type 2 diabetes
identified by proteomic studies. *Acta Diabetol, 54*(9), 833-842. doi:10.1007/s00592-017-
1013-y

Grosche, A., Hauser, A., Lepper, M. F., Mayo, R., von Toerne, C., Merl-Pham, J., et al. (2016). The
Proteome of Native Adult Muller Glial Cells From Murine Retina. *Mol Cell Proteomics,*
15(2), 462-480. doi:10.1074/mcp.M115.052183

Harding, J. L., Pavkov, M. E., Magliano, D. J., Shaw, J. E., & Gregg, E. W. (2019). Global trends in
diabetes complications: a review of current evidence. *Diabetologia, 62*(1), 3-16.
doi:10.1007/s00125-018-4711-2

Hartman, M. L., Shirihai, O. S., Holbrook, M., Xu, G., Kocherla, M., Shah, A., et al. (2014). Relation
of mitochondrial oxygen consumption in peripheral blood mononuclear cells to vascular
function in type 2 diabetes mellitus. *Vasc Med, 19*(1), 67-74. doi:10.1177/1358863x14521315

Hinkel, R., Howe, A., Renner, S., Ng, J., Lee, S., Klett, K., et al. (2017). Diabetes Mellitus-Induced
Microvascular Destabilization in the Myocardium. *J Am Coll Cardiol, 69*(2), 131-143.
doi:10.1016/j.jacc.2016.10.058

Hu, S., Chen, C. W., Chen, S. T., Tsui, K. H., Tang, T. K., Cheng, H. T., et al. (2019). Inhibitory
effect of berberine on interleukin-2 secretion from PHA-treated lymphocytic Jurkat cells. *Int*
Immunopharmacol, 66, 267-273. doi:10.1016/j.intimp.2018.11.020

Hu, Y., Sun, J., Dai, Z., Deng, H., Li, X., Huang, Q., et al. (2020). Prevalence and severity of corona
virus disease 2019 (COVID-19): A systematic review and meta-analysis. *J Clin Virol, 127*,
104371. doi:10.1016/j.jcv.2020.104371

Huang, P., Zhou, Y., Liu, Z., & Zhang, P. (2016). Interaction between ANXA1 and GATA-3 in
Immunosuppression of CD4(+) T Cells. *Mediators Inflamm, 2016*, 1701059.
doi:10.1155/2016/1701059

Jassal, B., Matthews, L., Viteri, G., Gong, C., Lorente, P., Fabregat, A., et al. (2020). The reactome
pathway knowledgebase. *Nucleic acids research, 48*(D1), D498-D503.
doi:10.1093/nar/gkz1031

Klec, C., Ziomek, G., Pichler, M., Malli, R., & Graier, W. F. (2019). Calcium Signaling in ß-cell
Physiology and Pathology: A Revisit. *Int J Mol Sci, 20*(24), 6110. doi:10.3390/ijms20246110

Kleinwort, K. J. H., Amann, B., Hauck, S. M., Hirmer, S., Blutke, A., Renner, S., et al. (2017).
Retinopathy with central oedema in an INS (C94Y) transgenic pig model of long-term
diabetes. *Diabetologia, 60*(8), 1541-1549. doi:10.1007/s00125-017-4290-7

Kounakis, K., Chaniotakis, M., Markaki, M., & Tavernarakis, N. (2019). Emerging Roles of
Lipophagy in Health and Disease. *Frontiers in Cell and Developmental Biology, 7*(185).
doi:10.3389/fcell.2019.00185

Kumar, N. P., Fukutani, K. F., Shruthi, B. S., Alves, T., Silveira-Mattos, P. S., Rocha, M. S., et al.
(2019). Persistent inflammation during anti-tuberculosis treatment with diabetes comorbidity.
Elife, 8. doi:10.7554/eLife.46477

Lau, E. Y. M., Carroll, E. C., Callender, L. A., Hood, G. A., Berryman, V., Pattrick, M., et al. (2019).
Type 2 diabetes is associated with the accumulation of senescent T cells. *Clin Exp Immunol,*
197(2), 205-213. doi:10.1111/cei.13344

Lee, Y.-h., Kim, S. R., Han, D. H., Yu, H. T., Han, Y. D., Kim, J. H., et al. (2019). Senescent T Cells
Predict the Development of Hyperglycemia in Humans. *Diabetes, 68*(1), 156.
doi:10.2337/db17-1218

Lepper, M. F., Ohmayer, U., von Toerne, C., Maison, N., Ziegler, A. G., & Hauck, S. M. (2018).
Proteomic Landscape of Patient-Derived CD4+ T Cells in Recent-Onset Type 1 Diabetes. *J*
Proteome Res, 17(1), 618-634. doi:10.1021/acs.jproteome.7b00712

Lin, V. H.-C., Chen, J.-J., Liao, C.-C., Lee, S.-S., & Chien, E. J. (2016). The rapid
immunosuppression in phytohemagglutinin-activated human T cells is inhibited by the
proliferative Ca2+ influx induced by progesterone and analogs. *Steroids, 111*, 71-78.
doi:https://doi.org/10.1016/j.steroids.2016.01.010

Liu, J., Guo, K., Hu, L., Luo, T., Ma, Y., Zhang, Y., et al. (2019). ZAP70 deficiency promotes
reverse cholesterol transport through MAPK/ERK pathway in Jurkat cell. *Mol Immunol, 107*,
21-28. doi:10.1016/j.molimm.2019.01.001

Marshall, R. J., Armart, P., Hulme, K. D., Chew, K. Y., Brown, A. C., Hansbro, P. M., et al. (2020).
Glycemic Variability in Diabetes Increases the Severity of Influenza. *mBio, 11*(2).
doi:10.1128/mBio.02841-19

Nam, J. H., Cha, B., Park, J. Y., Abekura, F., Kim, C. H., & Kim, J. R. (2019). Mitogen-Induced
Interferon Gamma Production in Human Whole Blood: The Effect of Heat and Cations. *Curr*
Pharm Biotechnol, 20(7), 562-572. doi:10.2174/1389201020666190528093432

Nicholas, D. A., Proctor, E. A., Agrawal, M., Belkina, A. C., Van Nostrand, S. C., Panneerseelan-
Bharath, L., et al. (2019). Fatty Acid Metabolites Combine with Reduced β Oxidation to
Activate Th17 Inflammation in Human Type 2 Diabetes. *Cell Metab, 30*(3), 447-461.e445.
doi:https://doi.org/10.1016/j.cmet.2019.07.004

Nyambuya, T. M., Dludla, P. V., Mxinwa, V., & Nkambule, B. B. (2020). T-cell activation and
cardiovascular risk in adults with type 2 diabetes mellitus: A systematic review and meta-
analysis. *Clinical Immunology, 210*, 108313. doi:https://doi.org/10.1016/j.clim.2019.108313

Olenchock, B. A., Rathmell, J. C., & Vander Heiden, M. G. (2017). Biochemical Underpinnings of
Immune Cell Metabolic Phenotypes. *Immunity, 46*(5), 703-713.
doi:10.1016/j.immuni.2017.04.013

Purvis, G. S. D., Chiazza, F., Chen, J., Azevedo-Loiola, R., Martin, L., Kusters, D. H. M., et al.
(2018). Annexin A1 attenuates microvascular complications through restoration of Akt
signalling in a murine model of type 1 diabetes. *Diabetologia, 61*(2), 482-495.
doi:10.1007/s00125-017-4469-y

Purvis, G. S. D., Collino, M., Loiola, R. A., Baragetti, A., Chiazza, F., Brovelli, M., et al. (2019).
Identification of AnnexinA1 as an Endogenous Regulator of RhoA, and Its Role in the
Pathophysiology and Experimental Therapy of Type-2 Diabetes. *Front Immunol, 10*, 571.
doi:10.3389/fimmu.2019.00571

Purvis, G. S. D., Collino, M., Loiola, R. A., Baragetti, A., Chiazza, F., Brovelli, M., et al. (2019).
Identification of AnnexinA1 as an Endogenous Regulator of RhoA, and Its Role in the
Pathophysiology and Experimental Therapy of Type-2 Diabetes. *Front Immunol, 10*(571).
doi:10.3389/fimmu.2019.00571

Purvis, G. S. D., Solito, E., & Thiemermann, C. (2019). Annexin-A1: Therapeutic Potential in
Microvascular Disease. *Front Immunol, 10*, 938. doi:10.3389/fimmu.2019.00938

Reiter, L., Rinner, O., Picotti, P., Huttenhain, R., Beck, M., Brusniak, M. Y., et al. (2011). mProphet:
automated data processing and statistical validation for large-scale SRM experiments. *Nat*
Methods, 8(5), 430-435. doi:10.1038/nmeth.1584

Renner, S., Braun-Reichhart, C., Blutke, A., Herbach, N., Emrich, D., Streckel, E., et al. (2013).
Permanent neonatal diabetes in INS(C94Y) transgenic pigs. *Diabetes, 62*(5), 1505-1511.
doi:10.2337/db12-1065

Richard, C., Wadowski, M., Goruk, S., Cameron, L., Sharma, A. M., & Field, C. J. (2017).
Individuals with obesity and type 2 diabetes have additional immune dysfunction compared
with obese individuals who are metabolically healthy. *BMJ Open Diabetes Res Care, 5*(1),
e000379. doi:10.1136/bmjdrc-2016-000379

Rodríguez-Gómez, I. M., Talker, S. C., Käser, T., Stadler, M., Reiter, L., Ladinig, A., et al. (2019).
Expression of T-Bet, Eomesodermin, and GATA-3 Correlates With Distinct Phenotypes and
Functional Properties in Porcine γδ T Cells. *10*(396). doi:10.3389/fimmu.2019.00396

Singh, J., Kaade, E., Muntel, J., Bruderer, R., Reiter, L., Thelen, M., et al. (2020). Systematic
Comparison of Strategies for the Enrichment of Lysosomes by Data Independent Acquisition.
J Proteome Res, 19(1), 371-381. doi:10.1021/acs.jproteome.9b00580

Soongsathitanon, J., Umsa-Ard, W., & Thongboonkerd, V. (2019). Proteomic analysis of peripheral
blood polymorphonuclear cells (PBMCs) reveals alteration of neutrophil extracellular trap
(NET) components in uncontrolled diabetes. *Mol Cell Biochem, 461*(1-2), 1-14.
doi:10.1007/s11010-019-03583-y

Stentz, F. B., & Kitabchi, A. E. (2007). Transcriptome and Proteome Expressions Involved in Insulin
Resistance in Muscle and Activated T-Lymphocytes of Patients with Type 2 Diabetes.
Genomics, Proteomics & Bioinformatics, 5(3), 216-235. doi:https://doi.org/10.1016/S1672-
0229(08)60009-1

Tian, Y., Yang, B., Qiu, W., Hao, Y., Zhang, Z., Yang, B., et al. (2019). ER-residential Nogo-B
accelerates NAFLD-associated HCC mediated by metabolic reprogramming of oxLDL
lipophagy. *Nat Commun, 10*(1), 3391. doi:10.1038/s41467-019-11274-x

van der Windt, G. J. W., Chang, C. H., & Pearce, E. L. (2016). Measuring Bioenergetics in T Cells
Using a Seahorse Extracellular Flux Analyzer. *Curr Protoc Immunol, 113*, 3.16b.11-
13.16b.14. doi:10.1002/0471142735.im0316bs113

Widlansky, M. E., Wang, J., Shenouda, S. M., Hagen, T. M., Smith, A. R., Kizhakekuttu, T. J., et al.
(2010). Altered mitochondrial membrane potential, mass, and morphology in the
mononuclear cells of humans with type 2 diabetes. *Transl Res, 156*(1), 15-25.
doi:10.1016/j.trsl.2010.04.001

Yang, Y. H., Song, W., Deane, J. A., Kao, W., Ooi, J. D., Ngo, D., et al. (2013). Deficiency of
annexin A1 in CD4+ T cells exacerbates T cell-dependent inflammation. *J Immunol, 190*(3),
997-1007. doi:10.4049/jimmunol.1202236

Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., et al. (2020). Clinical course and risk factors for
mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study.
The Lancet, 395(10229), 1054-1062. doi:https://doi.org/10.1016/S0140-6736(20)30566-3

Figure legends

Fig. 1 Proliferative capacity of porcine lymphocytes. Mitogens were used for stimulation of lymphocytes analyzed after 48 h of culture: **(A)** pokeweed mitogen (PWM) **(B)** concanavalin A (ConA) or **(C)** phytohaemagglutinin (PHA). Proliferation rate of stimulated lymphocytes did not differ between wild-type (wt, n=12) and INS^{C94Y} tg pigs (n=10) stimulated by ConA and PWM. After PHA-stimulation, lymphocytes of INS^{C94Y} tg pigs (n=11) revealed a significantly 5-fold decreased proliferation rate (***p≤0.001) compared to non-transgenic wild-type littermates (n=13). Data are represented as means ± SEM; horizontal bars indicate group means.

Fig. 2 Quantitative proteome analyses of porcine CD4$^+$ T cells. **(A)** Hierarchical cluster analysis of the porcine CD4$^+$ T cell proteome revealed significant differences in protein abundance (p < 0.05) in INS^{C94Y} tg pigs (n=4) and non-transgenic littermates (wt, n=5) indicated by their human orthologue gene names. Red color indicates increased and blue color decreased abundances of the identified proteins. **(B)** Volcano plot illustrates quantified proteins (≥2 unique peptides) in CD4$^+$ T cells of INS^{C94Y} tg pigs and wild-types (wt) plotted against the respective statistical significance. The horizontal line indicates the p-value cut-off at 0.05. Proteins with a mean fold change ≥ 2 and p <0.05 are shown in red (MYL4 and ANXA1).

Fig. 3 Differential ANXA1 expression in porcine CD4$^+$ T cells. **(A-C)** Representative flow cytometric images of ANXA1$^+$CD4$^+$ T cells of a wild-type **(A)** and an INS^{C94Y} tg pig **(B)** are shown with reduced ANXA1 expression in wild-type CD4$^+$ T cells. **(C)** Scatter plots summarising flow cytometry analyses of PBMC of eleven wild-type and nine INS^{C94Y} tg pigs. ANXA1 was significantly elevated in CD4$^+$ T cells of INS^{C94Y} tg pigs (*p < 0.05). **(D-F)** Similar intracellular ANXA1 levels in purified CD4$^+$ T cells of a wild-type **(D)** and an INS^{C94Y} tg pig **(E)** shown by immunofluorescence staining for ANXA1 (red), CD4 (green) and nucleus (blue, counterstained with DAPI) and confirmed by flow cytometry using n=4 per group **(F)**. **(G-I)** Representative images of immunofluorescence staining for ANXA1 on the outer cell membrane of porcine CD4$^+$ T cells demonstrate higher ANXA1 expression in CD4$^+$ T cells of the INS^{C94Y} tg pig **(H)** compared to the wild-type **(G)** (arrows). **(I)** A higher ANXA1 expression on the cell surface of diabetic CD4$^+$ T cells compared to CD4$^+$ T cells of wild-types was confirmed by flow cytometry (n=6 per group). Horizontal bars indicate group means.

Fig. 4 Basal glycolytic and mitochondrial respiratory profiles of porcine PBMC. **(A)** Basal glycolytic rate (Extracellular acidification rate, ECAR) was significantly elevated in PBMC of INS^{C94Y} tg pigs (n=10) compared to wild-type littermates (wt, n=15) (*p<0.05, Student's *t* test) **(B)** Oxygen consumption rate (OCR) was measured under basal conditions and after injection of oligomycin, FCCP, rotenone and antimycin A. Compared to PBMC from wild-types (wt, n=15), mitochondrial respiration of PBMC from diabetic pigs (n=10) yielded a significantly increased basal, ATP-linked and maximal respiration (**p<0.01) as well as a significantly increased spare respiratory capacity (*p<0.05). Data are represented as means ± SEM.

565   **Table 1** Pathway Enrichment Analysis of CD4$^+$ T cells from wild-type littermates.

Most significant pathways enriched in CD4$^+$ T cells of wild-types		
Pathway name	**p-value**	**Identified proteins**
Lagging Strand Synthesis	2.00e-12	POLD1, POLD2, POLD3, RFC2, RFC3
Gap -filling DNA repair synthesis and ligation in GG-NER	2.69e-12	XRCC1, POLD1,POLD2, POLD3, RFC2, RFC3
Polymerase switching	4.45e-12	POLD1,POLD2, POLD3, RFC2, RFC3

566   **Table 2** Pathway Enrichment Analysis of CD4$^+$ T cells from *INS*^{C94Y} tg pigs.

Most significant pathways enriched in CD4$^+$ T cells of *INS*^{C94Y} tg pigs		
Pathway name	**p-value**	**Identified proteins**
Downstream signal transduction	4.37e-04	NCK1, PTPN11
RUNX1 regulates transcription of genes involved in differentiation of myeloid cells	8.92e-04	PRKCB
Lipophagy	0.001	PLIN3, PRKAB1