The importance of animal weapons and fighting style in animal contests

Alexandre V. Palaoro[1,2,*] & Paulo Enrique Cardoso Peixoto[3]

[1]LUTA do Departamento de Ecologia e Biologia Evolutiva, Universidade Federal de São Paulo,

Brazil

[2]LAGE do Departamento de Ecologia, Universidade de São Paulo, Brazil

[3]LASEXIA, Departamento de Genética, Ecologia e Evolução, Universidade Federal de Minas

Gerais, Brazil

* Corresponding author: LUTA do Departamento de Ecologia e Biologia Evolutiva,

Universidade Federal de São Paulo, São Paulo, Brazil. Rua Artur Riedel, 275, Eldorado,

Diadema, CEP 09972-270. E-mail: alexandre.palaoro@gmail.com

Word count. 11,584

ABSTRACT

In many species that fight over resources, individuals use specialized structures to gain

mechanical advantage over their rivals during contests (i.e., weapons). Although weapons are

widespread across animals, how they affect the probability of winning contests is still debated.

According to theory, understanding the weapons' function in contests depends on identifying

differences in how weapons are measured (e.g., weapon length *versus* shape), and in how

weapons are used during fights. Here, we developed a meta-analysis spanning 1,138 studies,

from which were drawn 52 species and 107 effect sizes to identify: (1) what aspects of animal

weapons are measured in the literature, and how these measures bias our knowledge; (2) how

animals use their weapons during fights - i.e., weapon function; and (3) if weapon function

correlates to the magnitude of how weapons influence contest resolution. First, we found that

most of the literature focuses on linear measures of weapons, such as length. The few reports on

weapon performance (e.g., biting force) were found only for Crustacea and Squamata. This bias

highlights that measuring performance of weapons such as horns and spines might increase the

breadth of our knowledge on weapons. Furthermore, we also found that linear measures

showed stronger effects on contest success than performance measures. Second, we divided

weapon function into displays and fighting style (i.e., how the weapon is used during fights).

Regarding displays, most species displayed their weapons before contests (59.61%), rather than

the body (34.61%). A minority (three species, 5.76%) engaged in fights without any type of

display. Thus, species that bear weapons almost always perform displays before engaging in

physical contact, a common hypothesis in contest theory that was never tested across taxa until

now. Regarding fighting style, we found that most weapons were used for more than one

behaviour during fights (e.g., squeezing and pushing). Further, pushing seems to be the most

common behaviour among species, but it is usually accompanied by another behaviour, such as

lifting or squeezing. Thus, oversimplifying fighting style can bias results because some styles

might impose contrasting biomechanical pressures (e.g., pushing vs squeezing). Third, we

found that display type did not influence the importance of weapon size on contests. Fighting

style, on the other hand, influenced the effect of weapon size on contest outcome significantly.

Species that used their weapons to impact, pierce or squeeze showed smaller differences

between winners and losers when compared to pushing or lifting (and multifunctional

weapons). Thus, pushing and lifting seem important for selecting larger weapons – even though

some of them might also be used for squeezing, piercing or impacting. Overall, our results show

that we have a biased understanding of animal weapons, built mostly on weapon size alone.

Further, our analyses show that the importance of weapon size differs depending on the

fighting style. If we lessen those biases, we will have a better and broader understanding of how

weapons evolve and diversify.

Keywords. Animal contests, morphological evolution, animal fighting, sexual selection, weapon

biomechanics

CONTENTS

80
81

I. INTRODUCTION

Agonistic interactions have drawn attention from scientists since Darwin's seminal publication

of *The Descent of Man, and Selection in Relation to Sex* (Darwin, 1871). The field started mostly as

descriptions of how animals engaged in agonism (e.g., Archer, 1988), but then gained a strong

theoretical background by adding evolutionary game theory to its core (Maynard Smith & Price,

1973). Building on that theory, the field currently focus on how individuals make decisions

during agonistic interactions (Hardy & Briffa, 2013; Chapin, Peixoto, & Briffa, 2019), the

evolutionary consequences of winning or losing (Filice & Dukas, 2019), and what traits affect

winning chances (Vieira & Peixoto, 2013). Among these diverse topics, however, one subject

frequently raised is the existence of specialized morphological structures used during contests

(termed animal weapons, McCullough, Miller, & Emlen, 2016). A weapon can be seen as

specialized morphology used to gain mechanical advantage on the rivals (*sensu* Eberhard *et al.*,

2018; see Table 1 for definitions), and because of this effect on the rival, weapons can affect all

aspects of agonistic interactions. For instance, rivals may assess aspects of weapon size or

performance when deciding whether to stay in or give up on the contest (Palaoro & Briffa, 2017;

Pinto, Palaoro, & Peixoto, 2019). Weapons can also influence whether a contest escalate to

physical fighting or not (Számadó, 2008). A myriad of studies have already shown that weapons

are important for contest resolution, but most of them focus on single species (Vieira & Peixoto,

2013; Pinto *et al.*, 2019). Reviews on this topic provide diverse insights on weapon evolution

because they contain extensive knowledge on the shapes and sizes of animal weapons (Emlen,

2008; Rico-Guevara & Hurme, 2019). However, they lack quantitative information on selection

pressures that may act on weapons. Due to this, it is necessary to integrate data of a wider

diversity of species to estimate how weapons affect contests; hence providing a broader picture

of the relative importance of weapons for contest resolution.

One major hurdle to assess the relative importance of weapons is the large diversity of

measures used as proxies of weapon traits (Vieira & Peixoto, 2013). For instance, weapon traits

may be divided into three major categories: performance (such as a weapon's capacity to exert

force), size (such as weapon mass or length), and shape (such as the ratio between weapon

length and width). Although all of them may be important for determining the winner, there is

a debate on which trait(s) should influence contests the most (Eberhard *et al.*, 2018; Palaoro *et al.*,

2020). If one trait is more important than the others, we can expect stronger selective pressure

on that trait. However, a systematic review on how different weapon traits affect winning

chances is still lacking. A first step would be to identify the existing patterns on how studies are

measuring the traits of different weapons to assess if there are any gaps or potential biases in

our knowledge. The second step would be to quantify the relative importance of each weapon

trait for contest resolution. For example, if performance is more important than size for victory,

we expect larger differences between winners and losers on proxies of weapon performance

(such as weapon muscle mass) when compared to proxies of weapon size (such as weapon

length).

Another point frequently raised in studies of weapon evolution is related to how the

function of the weapon should influence the importance of each weapon trait on winning

contests (McCullough, Tobalske, & Emlen, 2014; McCullough *et al.*, 2016, see Table 1 for

definitions). Before physical contact, for instance, individuals may vary on whether they use

their weapons as displays or not. Rivals can either use their weapons before the fight as threat

displays (Lappin *et al.*, 2006), or use their weapons only during physical contact (Katsuki *et al.*,

2014). Weapons also vary in how they are used during fights, i.e., fighting style (*sensu*

McCullough *et al.*, 2016). For instance, individuals may use weapons to push (Graham, Padilla-

Perez, & Angilletta, 2020), squeeze (Dennenmoser & Christy, 2013), pierce (Candaten *et al.*,

2020), or even lift and throw their rivals away from the contested resource (Goyens, Dirckx, &

Aerts, 2015). Due to this variation, fighting styles have been notoriously difficult to assess

precisely. However, without a standardized framework to assess weapon function, we cannot

identify potential differences in the selective pressures on weapon traits that may be related to

the way rivals use their weapons during contests. For this, it is also necessary to perform a

systematic review of how individuals use their weapons during contests; then test whether

different fighting styles are able to predict how important each weapon trait is for contest

resolution.

A crucial step to perform a systematic review to estimate the importance of weapons in

contest resolution is to quantify how weapon traits affect winning chances. One way to obtain

such quantification is to estimate how much winners and losers differ in a given weapon trait

(e.g., weapon size or performance). However, since different species may differ in how much a

given weapon trait varies or in the type of measure that can be done, the estimation of the

relative difference in weapon traits among winners and losers must be standardized to be

comparable. In particular, the standardized mean difference in weapon traits between winners

and losers is a powerful metric frequently used in ecology and evolution (i.e., Hedges' g effect

size, Nakagawa & Cuthill, 2007): larger standardized mean differences between the groups

indicate a stronger effect of the measured trait or variable. When applied to animal contests, it

would be expected that traits that are more important in resolving contests should show

marked differences between winners and losers, while traits that are less important should be

more similar between rivals (e.g., Vieira & Peixoto, 2013). Therefore, as specific weapon traits

become more important to determine contest resolution, the greater should be the difference in

such traits between winners and losers.

Here, our goal is to quantify the importance of weapon traits and fighting style on

contest resolution. For this, we performed a meta-analytical review to answer the following

questions: 1) Are there biases in how weapons are measured and do distinct traits differentially

influence contest resolution? 2) How much variation exists in weapon function and are there

similarities among species? 3) Do similarities in displays and fighting style influence the relative

importance of weapon traits in contests? Below we provide a general description of how we

gathered information to answer these questions and then present our rationale and findings for

each question separately.

II. GENERAL PROCEDURES

Below we describe how we searched for the articles used in this review, how we extracted and transformed the information obtained from the articles and how we controlled for phylogenetic effects in each analysis. Such procedures were the same to obtain the data used to answer our three questions. Specific procedures adopted for each question are separately described in the corresponding section.

(1) Study selection and data gathering

We searched for articles using the Web of Science (https://www.webofknowledge.com) and Scopus (https://scopus.com) using their core collection databases from 1945 to 2019. For both searches we used the following keywords: "contest*", "fight*", "assessment*", "resource holding p*", "resource-holding-p*", "agonis*", "territory defen*e", "weapon*", "armament*", "sexual* trait*", "sexual*-selected trait*" "body size*", "antler*", "horn*", "jaw*", "claw*". All keywords were used with the "OR" Boolean operator. During our search, we excluded all studies in which the species did not bear a weapon (Table 1) such as butterflies and dragonflies. Since we had to classify species according to how individuals use them during contests, we excluded all species in which the behaviours adopted during the contests were undescribed.

For the selected studies, we collected information about mean values (and their corresponding variation) of weapon measures for winners and losers of contests. Within each study, we also recorded the pairing method used by researchers. We distinguished between studies in which fighting individuals were experimentally paired to have similar body sizes but differing weapon sizes and studies in which individuals were randomly chosen to contest (we also included in this second group studies in which one individual was free to choose their

rivals). Further, we found no studies that paired individuals by weapon size and let body sizes

differ; thus, we use 'paired contests' to refer to contests where individuals have similar body

sizes, but differing weapon sizes. Another confounding effect might come from resource value

(*sensu* Arnott & Elwood, 2008): individuals that value more a resource often are more motivated

to fight regardless of their fighting ability and have a higher chance of winning (e.g., Palaoro *et*

al., 2017). Therefore, to avoid any bias related to resource value, we only included the

treatments in which there was no evidence that individuals had different motivations to fight.

We also recorded whether the study was performed in a laboratory environment or in the wild.

We obtained a total of 1108 papers through those searches. We also added 30 relevant

papers cited in reviews that we did not find in our primary searches (Emlen, 2008; Vieira &

Peixoto, 2013; Pinto *et al.*, 2019; Rico-Guevara & Hurme, 2019). After excluding papers that did

not provide all necessary information (Fig. S1), our final data set comprise 48 papers that

contained 52 species, comprising both vertebrate and invertebrate species (Fig. S2). Within

these, we had information for 33 species involving randomly paired rivals and 27 species

involving rivals paired by size (Fig. 1). We performed all steps of the literature review following

the PRISMA protocol (Liberati *et al.*, 2009), and the flow diagram containing all steps can be

found in Fig. S1.

(2) Effect size calculations and phylogenetic tree

To compare the magnitude of trait differences between winners and losers across different

species, we used Hedges' *g*, which is a standardized dimensionless measure that allows

comparisons among different types of measurements and species (Koricheva, Gurevitch, &

Mengersen, 2013). To calculate Hedges' *g*, we used the mean trait values, their standard

deviations, and respective sample sizes for each trait measure (such as weapon mass or weapon

length) of winners and losers of each selected study. Since we always calculated the Hedges' *g*

as the difference between winner and loser traits, values greater than zero indicate that winners,

on average, have a greater trait value than losers; while values smaller than zero indicate that

losers have, on average, a lower trait value than winners. Unfortunately, mean values and

standard deviations were not always available on papers. In those cases, we gathered results

from statistical tests comparing winners and losers (e.g., *t*-values, F-values, degrees of freedom,

and sample sizes) to transform the statistical results into Hedges' *g* values using the package

"*compute.es*" in R (Del-Re, 2013). Whenever the mean and standard deviations were displayed

on graphs, we used the *webplotdigitizer* software to extract the values directly from the figures

(Rohatgi, 2019). If none of those options were available, we contacted the corresponding author

to request the data.

To control for the phylogenetic relatedness between the species in our sample, we built a

phylogeny comprising all 52 species (Fig. S2) using the Interactive Tree of Life online tree

generator (iTOL; https://itol.embl.de/). After generating the tree, we estimated branch length

using a Brownian motion model of evolution to simulate an ultrametric phylogenetic tree

(Paradis, Claude, & Strimmer, 2004). We then transformed the ultrametric tree into a variance-

covariance matrix that reflects the phylogenetic relatedness among the species. The variance-

covariance matrix was then imputed in our meta-analytical models (see below for a detailed

description of each model) as a random variable. We made these procedures using the packages

"*rtol*" (Michonneau, Brown, & Winter, 2016) and "*ape*" in R (Paradis *et al.*, 2004).

III. WHAT IS THE EFFECT OF DIFFERENT WEAPON TRAITS ON CONTEST RESOLUTION?

Weapons can be measured regarding their performance, size, and shape; each of these traits can be used as proxies for fighting ability (see Vieira & Peixoto, 2013 and Palaoro *et al.*, 2020b). This diversity of measures sparked interest on whether (and how) distinct weapon traits may affect contest resolution (Lappin & Husak, 2005; McCullough *et al.*, 2016; Eberhard *et al.*, 2018; Palaoro *et al.*, 2020). Given the important role weapons play in deciding a contest winner, understanding if each trait influences contest resolution is thus an important step to reveal how selective pressures may have acted on weapon evolution. In particular, if specific weapon traits are more important in determining victory, they should show greater differences between winners and losers when compared to weapon traits that are less important. For this reason, we used a meta-analytic review to assess how weapon traits are measured in studies on contest resolution; and tested if the difference between winners and losers changed according to the type of measurement made.

We divided the traits measured in six categories: weapon asymmetry, index of weapon size, weapon area, weapon performance, weapon linear measures (length or width), and weapon mass. Weapon asymmetry is often measured as the difference in size between the two sides of a bilateral weapon, e.g., the difference in the maxillae of a cricket (Briffa, 2008). The index of weapon size is used to calculate the size of a complex shape, i.e., to incorporate the complexity of branching of the antlers of a deer species into a metric of overall size (Hoem *et al.*, 2007). Within performance, we considered only measures related to force output, such as muscle size and bite force measurements. Regarding area, linear measures, and mass, although they might represent the same component of a weapon (i.e., size), they have different scaling properties that can add non-random biases to the estimates (see Houle *et al.*, 2011; Pélabon *et al.*,

2014). In addition, in Arthropods, while mass may vary with individual condition, size is a fixed

attribute in adults (e.g., Peixoto & Benson, 2008). Therefore, we separated linear from area and

mass measurements.

(1) Methods

After determining the type of trait measured for each species, we built a multilevel meta-

analytical model using the type of trait as our moderator, the Hedges' g effect size as our

response variable, and the inverse of the variance of Hedges' g as a weight. We also included

five variables as random effects. First, we used 'study ID' because sometimes we extracted more

than one effect size per study. In this random effect, we frequently had several traits of the same

weapon being measured (e.g., linear and mass measurements within the same study). Thus, we

build a correlation matrix for the 'study ID' random effect because the correlation between

effect sizes can bias the outcome (e.g., Weaver *et al.*, 2018; Mathot, Dingemanse, & Nakagawa,

2019). The correlation matrix had the Pearson's correlation coefficient between different

weapons traits to control for any allometric effect on our estimates. We found most of the

correlations in the same papers from which we found the effect sizes. But, for those that we did

not find, we searched the literature and used Pearson's coefficients in the papers cited in Table

S1. In the few cases in which no information was available, we used 0.5 as a coefficient value

(following Weaver *et al.*, 2018). The matrix we used can be assessed together with our codes and

dataset (check the Data Availability section). Second, we used 'species ID' to account for effect

sizes that came from the same species, but different studies. Third, we used the environment in

which the original study was performed (i.e., laboratory or wild). Fourth, we used a matrix

containing the phylogenetic relatedness among species. Finally, we used the pairing system

used in the original study (i.e., whether individuals were paired according to their body size or

not).

To estimate heterogeneity and biases in the model, we used two approaches. First, we

calculated the ratio of heterogeneity to the total variation observed across effect estimates in

multivariate studies (I^2, Borenstein, 2009). We also partitioned the I^2 into the contribution of

each random variable (Nakagawa & Santos, 2012). Thus, we had estimations of the within-

studies variance (I^2_{study}, similar to most mixed models), the species ID variance ($I^2_{species}$), the

phylogenetic variance ($I^2_{phylogeny}$), and the pairing method variance ($I^2_{pairing}$). The sum of these

different I^2 are equal to the total variance observed (I^2_{total}). To estimate the phylogenetic signal in

the effect size, we also calculated the phylogenetic heritability index, H^2, which is similar to

Pagel's λ (Nakagawa & Santos, 2012). Finally, to test for publication bias, we conducted a

modified version of the Egger's test, in which we used the residuals of our meta-analytic model

as the response variable and the standard deviation of the effect sizes as our predictor variable

(Egger *et al.*, 1997). If the intercept of this regression is not different from zero ($\alpha > 0.10$), then

there is little evidence for publication bias (Nakagawa & Santos, 2012).

(2) Results

Linear measures were the most common trait found in the literature (74 out of 107 effect sizes,

69.15%). Performance measures were the second most common with 23 effect sizes (21.49%),

followed by area (n = 4, 3.73%), asymmetry (n = 3, 2.88%), mass (n = 2, 1.87%), and index

measures (n = 1, 0.9%). Linear measures were found for most species in the sample, but

performance measures were concentrated on crustaceans and lizards. Of the 23 effect sizes on

performance, 11 came from crustaceans (47.8%), 11 from lizards (47.8%) and 1 from a cricket

(*Teleogryllus commodus*, Fig. 1). Therefore, there is a clear bias on the type of inference drawn for most groups: inferences still rely mostly on size measures, rather than performance, or other measures.

The low sample sizes for measures of weapon area, asymmetry, mass, and index would render any statistical assessment of their influence on contest success weak. Thus, we removed these measures from our sample and only tested the linear and performance measures. Overall, winners had greater weapon traits than losers regardless of whether linear or performance was measured (QM_2 = 117.05, p <0.0001). However, linear measures had a greater effect on contest success than performance measures (QM_1 = 23.29, p <0.0001; Fig. 2). The model showed low heterogeneity. Study ID and pairing method were responsible for most of that heterogeneity, while phylogeny and genus had negligible effects on variance (Table 2). We found evidence for publication bias on the effect sizes (Egger's test; intercept: -0.315, 95% CI: -0.569, -0.06, t = -2.461, p = 0.015).

(3) Discussion

Our results suggest that linear measures have a higher relative importance to determine contest outcome than performance measures. Therefore, different traits of the weapon may provide distinct information about the relative importance of weapons in contests. Given recent evidence (e.g., Palaoro *et al.*, 2020b; Emberts *et al.*, 2021), it is somewhat surprising to find that weapon size is more important than weapon performance on contest resolution. However, interpreting these results requires caution because of the biases and limitations of the current literature. For instance, performance measures are concentrated on crustaceans and lizards (only one performance measure was found outside those two groups). These measures provide

information on the performance of weapons that have very similar morphologies: claws and

jaws. Meanwhile, linear measures have a wider diversity of weapons, such as antlers,

mandibles, and horns, which are also distributed among more taxa. Therefore, to ensure that

the effect we found is not an artifact associated with differences in the diversity of weapons that

have linear and performance measures, we reran the analysis using only data gathered from

crustaceans and lizards. The result was the same to that observed in the analysis considering all

species: the effect size of linear components was higher than the effect size of performance

components (Fig. S4). Thus, despite the bias in the diversity of weapons, it seems that linear

measures are more important for contest resolution than performance measures.

One explanation for why linear measures seem more important than performance

measures may reside on engaging the rival from a safe distance (Eberhard *et al.*, 2018). Being

able to handle the rival without being exposed can also give the individual more time align the

weapon relative to the rival's weapon or body without incurring any extra costs (fighting skill,

Briffa & Lane, 2017). If that is true, then any weapon can benefit from being larger. However,

that may be particularly true for weapons that can push the rival (even if pushing co-occurs

with other behaviours, see the next session). That way, the individual that has more time to find

a better spot to lock on the rival and enjoy and increased chance to win the contest by pushing

the opponent better. Thus, individuals might benefit from having a larger weapon, regardless of

their weapon morphology. Alternatively, weapons might be used as visual displays and during

fights to gain mechanical advantage on the rival. These selective pressures combined could

favour bigger weapons if size influences contest success by getting rivals to retreat before

physical contact (McCullough *et al.*, 2016). We tackle this hypothesis in the next sessions.

Along with the lack of measurements of performance during fights, there is a clear

information gap on the performance of weapons that are used for other types of behaviours,

such as striking or ramming. The only example outside of biting performance we have on our

dataset are the bullet-fast strikes of mantis shrimps (Green & Patek, 2015). Most of the types of

weapons where we miss information on other measures are, in fact, weapons that do not have

specific muscles attached to them, such as antlers, horns, and spines. These weapons are used

for a multitude of behaviours during fights (see next session) and rather than having their own

movement (i.e., biting, striking), they are used simultaneously with body movement. A few

studies show how to measure the performance of horns (McCullough, 2014) and spines (Crofts

et al., 2019) during relevant tasks. Thus, expanding our knowledge to how much these

performances influence contests can broaden our understanding of weapon evolution.

IV. ARE THERE SIMILARITIES AMONG SPECIES IN WEAPON FUNCTION?

Weapon function may be composed of two moments. In the first moment (not present in some

species, see below), individuals may use their weapons as displays before physical contact

ensue. In the second moment, individuals use their weapons to manipulate the rival, which is

referred here as fighting style (*sensu* McCullough *et al.*, 2016). Both moments can vary broadly

within and between species. For instance, in the fiddler crab, *Austruca mjoebergi*, males use their

claws as threat displays before physical contact. If the contest escalates, males use their claws to

pinch rivals; while pinching, males are also trying to lift rivals off the ground and shove them

away (Dennenmoser & Christy, 2013). In other species of crustaceans, such as crayfish, claws

are used not only to pinch but also to push rivals (Graham *et al.*, 2020). Perhaps because of this

variation, weapon function has seldom been comparatively investigated across different taxa. In

the few groups this has been done (e.g., bovids, Caro *et al.*, 2003) such variation has been

eliminated by linking fighting style of a species to a single behaviour (and discarding displays).

However, since the behaviour adopted may change, ascribing fighting style to a single

behaviour may underestimate the importance of the weapon for contests success and restrict

our comprehension about the selective forces that act on weapon evolution. For this reason,

here we reviewed the behaviours adopted during the fight for the species selected in our meta-

analysis, which include both vertebrates and invertebrates. We also identified if there are

groups of species that show similar weapon functions based on more complete descriptions of

fighting behaviour.

(1) Contest descriptions

For each species selected in our meta-analysis, we searched for information on weapon

function. When the selected article did not provide descriptions of fighting behaviour, we

searched for additional articles that contained detailed descriptions of the fighting behaviour

(Supplementary File S2). Based on the exact descriptions provided in the literature, we were

able to identify general categories of how contests begin and how the weapon is used during

physical contact (i.e., fighting style).

(a) How do contests begin?

We classified species in three groups according to how contests begin. First, we found species in

which individuals perform behaviours that allow them to gather information about the rival's

body size before engaging in contests. For instance, in the cricket *Melanotes ornata,* males lash

their antennae toward the rival's body before deciding whether to use their legs to kick the rival

(Lobregat *et al.*, 2019). In the lizard *Carinascincus microlepidotus*, on the other hand, males turned

sideways towards the rival and performed dorso-lateral displays of the body before escalating

to physical contact (Olsson & Shine, 2000). We named this category 'body size estimation'

because individuals had a chance to assess their rivals body given their displays and antennae

touch, but they did not necessarily have threat displays exclusively involving the weapon

(following Számadó, 2008). Second, we found species in which individuals perform behaviours

that allow them to gather more precise information about their rivals' weapons before engaging

in physical contact. For example, in the crayfish *Cherax dispar*, individuals display their enlarged

front claws (which is used as a weapon) to one another before making physical contact, which

allow a visual estimation of weapon size (Wilson *et al.*, 2007). Grasshoppers also used similar

behaviours in which they flare their mandibles to the rivals before engaging in physical contact

(Umbers *et al.*, 2012). We named this category 'weapon display' because rivals had the

opportunity to assess the size of the rival's weapon (or be threatened by the rival's weapon, see

Számadó, 2008). Third, there were species that did not perform explicit behaviours that would

allow rivals to gather information about body or weapon size before engaging in physical

contact. An example is the *Sagra femorata* beetle, in which rivals do not use any type of display

before beginning a physical struggle (Katsuki *et al.*, 2014; O'Brien & Boisseau, 2018). We named

this category 'no display' because there was no evidence that rivals gather information or

threaten each other before contests ensue. Although we cannot exclude the possibility that

rivals gather information during the physical contact phase, we assure that the decision to begin

a contest in the group of 'no display' is little or not affected by mutual evaluations performed

before rivals engage in a physical struggle. The full descriptions of contest behaviours for all

species can be checked in Supplementary File S2.

(b) How are weapons used during physical contact?

We identified six categories which relate to fighting style: (i) to lift; (ii) to push; (iii) to pull; (iv)

to squeeze; (v) to deliver forceful impact blows; (vi) to pierce. Lifting weapons were mainly

used to lift the rival from the substrate to either disbalance or topple the rival. For instance, the

stag beetle *Cyclommatus metallifer* uses enlarged mandibles to hold and lift the rival off the tree

trunk in which they frequently fight on (Goyens *et al.*, 2015). Pushing weapons were used to

push the rival away from the bearer. For instance, dung beetles that fight inside tunnels use

their horns to push rivals off the entrance of the tunnels (McCullough & Simmons, 2016).

Pulling weapons were used to pull the rival near the bearer, frequently dislodging it. For

instance, *Aegla longirostri* freshwater anomurans use their claws to pinch and pull the rivals,

dislodging them from the substrate (Ayres-Peres, Araujo, & Santos, 2011). This behaviour is

rarely performed without squeezing. Squeezing weapons were mainly used to provide forceful

grasp on a rival. Crustaceans and lizards are the frequent examples of this category, using their

claws and jaws to squeeze rival's body parts (Husak, Lappin, & Van Den Bussche, 2009;

Dennenmoser & Christy, 2013). Impact weapons were used to deliver rapid or explosive strikes

to the rival. A noteworthy example is the raptorial appendages of mantis shrimps, which are

used to strike the rival's telson with a movement so fast that it can crack the abdomen's cuticle

(Green & Patek, 2015 but see Taylor & Patek, 2010). Lastly, piercing weapons were used mainly

to pierce the rival's skin or cuticle, typically with sharp, pointy structures. One example is the

bird *Ficedula hypoleuca* which uses its beak to pierce the rival's skin during physical contact

(Dale & Slagsvold, 1995).

It is important to note that, despite these six different functions, most species used their

weapons for two or, more rarely, three functions during the fight. For example, in the cricket

Loxoblemmus doenitzi, males have a flat head with horns on the edges that are used during fights

to push one another. However, males may also use the horns to rapidly beat the rival's horns or

body (Kim, Jang, & Choe, 2011). Therefore, for species in which more than one weapon usage

was described, we created categories with all the functions associated with that weapon. In our

example, we consider that males of *L. doenitzi* use their weapons for both pushing and

impacting their rivals.

(2) Results

Based on the combination between how contests begin and how weapons are used during

contests we identified 16 categories of weapon function distributed among the 52 species (Table

3). According to the descriptions we gathered, contests frequently began by individuals

displaying their weapons to rivals (n = 32 species out of 52, 61.53%); less frequently by

displaying their body size (n = 17 of 52 species, 32.69%); and rarely by not making any display

(n = 3 of 52 species, 5.76%). Regarding function during fights, most weapons are used for more

than one function (n = 36 of 52, 69.23%), while few are used for a single function (n = 16 of 52,

30.76%). Regardless of whether we count multifunctional weapons, or weapons with a single

function, squeezing is the most common function (35), followed by pushing (23), lifting (15),

impacting (15), pulling (3), and piercing (2). To see how each species was categorized, see Fig. 1;

for complete descriptions, see Table 3.

(3) Discussion

According to contest theory, displays should be favoured in animals that frequently engage in

contests as a mechanism to decrease the costs of aggression (Emlen, 2008). By displaying (the

weapon or the body), an individual might induce the rival to give up on the contest even before

they started - thus saving energy. That is why displays should be favoured across all animals:

they decrease the likelihood of injuries and increase the amount of energy saved (Hardy &

Briffa, 2013). Our result is the first to corroborate the theoretical prediction for a directional

selection for displays using a diverse group of animals: we showed that displays are

commonplace among fighting animals. That seems especially true for weapon displays, which

is well distributed among all functions (Table 3) and hence does not seem subject to

oversampling of any given taxa. Further, displaying the weapon is more common than body

displays. Therefore, we have shown that selection seems to be acting to decrease the costs of

aggression across animals by favouring contests that begin with displays, rather than

instantaneous aggression (Emlen, 2008).

In the three cases that we found no displays, three opposite explanations arise. First,

animals might escalate all contests if resource value is extremely high. In the case of the male

hermit crabs *Diogenes nitidimanus*, for instance, males guard the females by grasping the outer

rim of her shell and carrying her around (Yoshino, Koga, & Oki, 2011). If that male loses the

possession of the female in a contest, it is likely it will not mate until the next reproductive

cycle. Access to mature females is thought to be difficult in *D. nitidimanus* because females are

only receptive during very short time windows (only after molting, Asakura, 1987). Thus, males

holding females might opt to go all-in in a fight to keep the female, similar to predictions from

the 'Desperado' effect (Grafen, 1987). Second, if the costs of fighting are very low, there might

be no selection to avoid fighting and decrease the costs. However, the three species in our

sample (*Trypoxylus dichotomus*, *Sagra femorata*, *D. nitidimanus*) have enlarged weapons (i.e.,

disproportional size in relation to body size, Yoshino *et al.*, 2011; Johns *et al.*, 2014; O'Brien,

Katsuki, & Emlen, 2017). Thus, it is unlikely that these species are evolving weapons because

contests have very low costs. The third explanation resides on a possible lack of details in

behavioural descriptions of fighting behaviour. Those might be unconscious biases, such as

focusing on what animals are doing during the fight and not when the fight is starting, or subtle

signalling behaviours that are outside the observer's sensory cognition. For instance, rhinoceros

beetles can use acoustic signals during courtship that were unknown until recently (Hunt *et al.*,

2020). Thus, it is possible that these species indeed use displays, but we were not able to assess

them yet.

Squeezing was the most common behaviour, followed by pushing. While squeezing can

be considered a bias because of the oversampling of crustaceans and lizards (both with

squeezing weapons), pushing is also common behaviour during fights, although in many

instances it is associated with another behaviour. As shown in Table 3, pushing was found

associated with another behaviour in 31 (of 34 records), being used as the only fighting

behaviour only in three species. Thus, pushing is one of the main reasons most weapons are

multifunctional. It is interesting to note that, in most species, pushing is associated with

squeezing – which is a similar pattern to the vectorially opposite behaviour, pulling.

Individuals are unable to pull a rival without holding the rival. Pushing, however, can be done

simply by contacting the individual (e.g., interlocking antlers used to push). Thus, despite these

differences, it seems that squeezing might increase the chance that an individual also tries to

manipulate the rival by pulling or pushing.

Piercing, on the other hand, is the rarest behaviour in our sample. The only species that

displayed that behaviour in our sample was *Ficedula hypoleuca*, a bird that pecks its rivals during

fights (Dale & Slagsvold, 1995), and *Tetranychus urticae*, a mite species in which individuals use

their stylets to pierce rivals (Potter, Wrensch, & Johnston, 1976). Other species, such as

hummingbirds (Rico-Guevara & Araya-Salas, 2015) and coreid bugs (Emberts *et al.*, 2021), can

use piercing as their fighting style, but even among them few groups use piercing. Piercing is

indeed expected to be rare because it might be injurious. Over evolutionary time, theory

predicts that species would tend to evolve displays to decrease the likelihood of engaging in

injurious contests, unless resource value is extremely high (Hardy & Briffa, 2013). By evolving

displays, weapons would then tend to increase in size and complexity, which could decrease

piercing performance and change the function of the weapon altogether (Emlen, 2008;

Anderson, 2018). A similar route is believed to have occurred in cervids, where short, pointy

antlers started to increase in size and complexity as species evolved (Barrette, 1977; Caro *et al.*,

2003; Davis, Brakora, & Lee, 2011). Our results show that using displays before fights is indeed

a common strategy among animals (Table 3), but we still need to test whether these displays

decrease the injury capacity of weapons over evolutionary time.

Another important pattern is that the fighting style of most species (65%) are comprised

of multiple behaviours. Since fights tend to involve ritualized behaviours (Hardy & Briffa,

2013), it seems improbable that the description of a weapon being used in more than one

behaviour occurred by chance. Therefore, it seems that there is a higher tendency for weapons

to be used for more behaviours. Perhaps more complex fighting styles increase the winning

chances because it gives more options for individuals to inflict costs on their rivals. But at the

same time, depending on the combinations of behaviour, this may also generate opposing

selective forces on weapon morphology. For example, weapons used for lifting and pushing

will probably favour a single morphological type that provides an efficient way to fit the

weapon on the rival and a strong body to work as a lever for the weapon to work properly.

Weapons used for impacting and pushing, on the other hand, should favour a strong structure

that is capable of delivering high forces, but a different morphology for pushing the rival.

Perhaps, the occurrence of opposite selective forces acting on weapons explains why some

functional combinations are not described (Corn *et al.*, 2021). Further, if some behaviours tend to

involve multiple non-weapon parts (e.g., lifting uses the legs and body), while others are

essentially a weapon movement (e.g., squeezing), perhaps these behaviours should not be

considered equally when weapon evolution is concerned. On the one hand, the distinct selective

forces on some functions might promote weapon diversification (Wainwright *et al.*, 2005; Polly,

2020). On the other hand, the weapon is primarily used solely in some behaviours. The full

breadth of these possibilities remains to be investigated.

V. DOES DISPLAY AND FIGHTING STYLE INFLUENCE WHICH WEAPON

COMPONENTS ARE IMPORTANT FOR CONTEST RESOLUTION?

As shown in the previous section, species differ in how they begin contests and in how they use

their weapons during fights. Given this variation, it is possible that displays and fighting style

affect which traits are more important to increase contest success. In particular, it may be that,

in species that display their weapons before engaging in physical contact, greater weapon size

increases the chance of contest success because weapon size would deter most rivals from

fighting. For species that instead assess body size, not weapon size, differences in weapon size

might have a more important role during fight than before physical contact ensues. Therefore,

in species that use body size displays, the difference between winners and losers in weapon size

should be lower than in species that use weapon displays. Finally, in species that do not use any

type of display before fighting, weapons may still be important in determining victory. The

absence of a display prior to physical contact may relax the selective pressure on weapon size,

but may increase the selection acting on performance depending on fighting style.

Regarding fighting style, it is possible that differences in how a weapon is used affects

the relative importance of weapon size on contest resolution. For species that use weapons to

lift or push rivals, reach should be important to decide who wins the contest. Basically, larger

weapons allow the individual to touch its rival before being touched. That allows the individual

to attack without being exposed to a riposte. Thus, we can expect a large selective pressure on

the size of the weapon for these two types of fighting (up to a certain mechanical limit, see

McCullough, 2014). Squeezing, impact, or piercing, on the other hand, do not necessarily rely

on size. Although weapon size may still be important (to attack first and due to allometric

effects, Pélabon *et al.*, 2014), a larger weapon may not equate to a weapon that performs better.

Crayfish, for instance, bear large claws that can be relatively weak for their overall size

(Robinson & Gifford, 2019). Because weapon size may not determine its performance, it is

possible that an individual with a smaller but stronger weapon can cause more injuries than an

individual with a larger but weaker weapon. Therefore, we expect the mean difference in

weapon size between winners and losers for weapons used to squeeze, impact, or pierce to be

low.

(1) Methods

To analyse the differences in the type of display and fighting style, we performed two

multilevel meta-analytical models using only the effect sizes for linear measurements. As can be

seen in section III, the data on performance measures contain only Crustacea and Squamata

(and one cricket), which biases the types of display and fighting style we observe. Thus, we

used the type of display evaluated in the previous session as our moderator, the Hedges' *g*

effect size as our response variable, and the inverse of the variance of Hedges' *g* as a weight.

The rest of the model, such as its random effects, and how we assessed heterogeneity, are equal

to the procedures described in session 3.1.

For the fighting style analysis, we used the descriptions on the previous section to

categorize fighting style in three groups: (i) Size-emphasis; (ii) Performance-emphasis; (iii)

Intermediate. The size-emphasis group consisted of weapons used to pull, push and lift rivals,

including weapons with these two functions. The performance-emphasis group consisted of

weapons used to squeeze, impact, pierce, or pull rivals. Again, if a weapon had two of these

three behaviours simultaneously, we categorized it as 'performance-emphasis'. Any weapon

used for two or more behaviours that belonged to two different groups (i.e., 'size' and

'performance-emphasis'), was categorized as 'intermediate' (all species that used the weapons

for three behaviours were included in this last category). For this multilevel meta-analytical

model, we used the function group as our moderator. We used the same random effects and

heterogeneity assessments described in session 3.1.

(2) Results

(a) How do fights begin?

Winners had larger weapons than losers regardless of the type of display on average (QM_3 =

28.48, p < 0.0001, Fig. 3), but the confidence interval overlapped zero when males did not use

displays (Fig. 3). Further, the types of displays did not differ among themselves (QM_2 = 3.18, p

= 0.2). The model showed low heterogeneity. Study ID and pairing method were responsible for

most of that heterogeneity, while phylogeny and genus had negligible effects on variance (Table

4). We found evidence for publication bias on the effect sizes (Egger's test; intercept: -0.368, 95%

CI: -0.656, -0.081, t = -2.553, p = 0.012).

(b) How do weapons differ among fighting style?

Winners were larger than losers in all categories, even though some of them had confidence

intervals that slightly overlapped with zero (QM_3 = 107.72, p < 0.0001, Fig. 4). We also found

differences in the asymmetry between winners and losers among the categories. Weapons were

more important for contest success in size-emphasis and intermediate fighting style when

compared to performance-emphasis (QM_1 = 18.84, p < 0.001, QM_1 = 8.42, p = 0.003,

respectively). The model showed low heterogeneity. Study ID and pairing method were

responsible for most of that heterogeneity, while phylogeny and genus had negligible effects on

variance (Table 5). We found evidence for publication bias on the effect sizes (Egger's test;

intercept: -0.368, 95% CI: -0.656, -0.081, t = -2.553, p = 0.012).

(3) Discussion

Contrary to our expectations, there was no difference in weapon size between winners and

losers among different types of display. Therefore, the existence of a weapon or body display

before the physical contact phase does not seem to impose additional selective pressures on

weapon size. This is interesting because visual displays are thought to be one important driver

for increments in weapon size (O'Brien *et al.*, 2017; Eberhard *et al.*, 2018), and most of the species

in our sample perform visual displays (Fig. 3).

On the other hand, the fighting style seems to impose different pressures on weapon

evolution: the mean difference between winners and losers was higher for the "size-emphasis"

and "intermediate" groups when compared with the "performance-emphasis" group.

Therefore, it seems that, even if a weapon is also used for squeezing, piercing or impacting, the

presence of an additional behaviour related to lifting or pushing will favour increases in

weapon size. Perhaps lifting and pushing are important in multifunctional weapons because

they allow an individual to reach the rival before being reached or to physically expel a rival

from the defended resource. For instance, *Melanotes ornata* crickets, and fiddler crabs can use

their weapons to throw their rivals away from the contested resource (Supplementary File S2).

It is important to highlight, however, that weapons have many different aspects that

may affect how they impose costs on rivals during the contest (Palaoro & Briffa, 2017).

Therefore, to reach a more comprehensive picture of how different weapon traits change

according to fighting style it is necessary to gather information on many different weapon

aspects - and many different weapon types. What is clear from our results is that species show

marked differences in fighting style and that such differences have the potential to affect (at

least for size) the way weapons evolve.

VI. CONCLUDING REMARKS

(1) Our data on 52 species of both vertebrates and invertebrates shows biases on how

animal weapons are measured. Linear components, such as lengths and widths, are

responsible for 69.15% of the information on the literature; performance components,

such as bite force, for 21.49%; and any other measures such as weapon asymmetry,

mass, or volume are rare.

(2) We also found significant bias on the taxa in which performance components are measured. Lizards and crustaceans (and one cricket species) are the only taxa in which we found measures of weapon performance (e.g., bite force) associated with winning or losing fights. Thus, there is a significant gap on how we perceive the importance of weapon performance to solve contests. To have a better idea of how performance is important, we need more information on the performance of weapons such as horns, antlers, and spines.

(3) Linear components of weapons were more associated with victory than performance components. This can be a corollary of the bias mentioned above, but our sensitivity analysis suggests that the effect is robust. One possible explanation is that size can allow individuals to handle rivals from a safe distance, regardless of the weapon type. Handling rivals from a safe distance can give the individual more time to get a tighter grip and increase their chance of winning the contest.

(4) The majority of species tend to begin contests by displaying their weapons to the rival (59.61%); a smaller portion either displays their body through movements or touch the rival with antennae or any other mechanosensory morphology (34.61%). A minority begins contests with no display at all (5.76%). Hence, as theory suggests, most species do begin contests by using displays.

(5) Few weapons were used for a single behaviour during contests (30.8%), most of them were used for two or more behaviours (69.2%). From those multiple behaviours, pushing rivals with the weapon is the most frequent co-occurring behaviour, suggesting that selection may not be working solely on a weapon's sole behaviour, but rather on the fighting style. Further, piercing is the rarest behaviour in our sample with only two species described with those weapons.

(6) We found no strong effect of type of display on the importance of weapon size on

contest resolution. The only subtle difference we found was that, when displays were

absent, weapon size did not differ between winners and losers. Given that most species

in our sample perform some type of display, it suggests that displays may signal

weapon size. However, we need more data from species that do not perform any type of

display to understand the effect properly.

(7) Fighting style influences the importance of weapon size on contests. Weapons used to

squeeze, impact or pierce had a lower difference between winners and losers when

compared to weapon used for push, pull, or lift or weapons with multiple behaviours.

Once again, it seems that being able to reach the rival first is important for contest

success.

Acknowledgments. We thank Dr. Patrick Green for the suggestions on an earlier version of the

manuscript. We thank all the authors who shared the raw data of their paper with us:

Chatchawan Chaisuekul, Christine W. Miller, Devin O'Brien, Kate Umbers, Leilani Walker,

Marcelo M. Dalosto, Murray Fea, Sarah E. Reece, Zach Stahlschmidt, Zachary Emberts, and

Zackary Graham. We also thank Eduardo S.A. Santos for his help with the heterogeneity and

bias analyses. A.V.P. thanks FAPESP for the post-doctoral grant (process no: 2016/22679-3).

P.E.C.P. thanks Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq

produtividade em pesquisa 311212/2018-2) and Pesquisa e Desenvolvimento of Agência

Nacional de Energia Elétrica and Companhia Energética de Minas Gerais (P&D

ANEEL/CEMIG, PROECOS project GT- 599).

Authors' contribution. A.V.P and P.E.C.P. designed the study, A.V.P. collected the data and ran

the analysis, A.V.P. and P.E.C.P. wrote the manuscript.

Data availability. All data and codes will be shared on github

(https://github.com/alexandrepalaoro) upon acceptance.

VII. REFERENCES

ANDERSON, P.S.L. (2018) Making a point: shared mechanics underlying the diversity of
biological puncture. *Journal of Experimental Biology* **221**.
ARCHER, J. (1988) *The Behavioural Biology of Aggression.* CUP Archive.
ARNOTT, G. & ELWOOD, R.W. (2008) Information gathering and decision making about resource
value in animal contests. *Animal Behaviour* **76**, 529–542.
ASAKURA, A. (1987) Population Ecology of the Sand-Dwelling Hermit Crab *Diogenes nitidimanus*
Terao: 3. Mating System. *Bulletin of Marine Science* **41**, 282–288.
AYRES-PERES, L., ARAUJO, P.B. & SANTOS, S. (2011) Description of the Agonistic Behavior of
Aegla longirostri (Decapoda: Aeglidae). *Journal of Crustacean Biology* **31**, 379–388.
BARRETTE, C. (1977) Fighting Behavior of Muntjac and the Evolution of Antlers. *Evolution* **31**,
169–176.
BORENSTEIN, M. (2009) Effect sizes for continuous data. In *The Handbook of Research
Synthesis and Meta-Analysis* (eds H. COOPER, L.V. HEDGES & J.C. VALENTINE), pp. 221–236.
Russell Sage Foundation.
BRIFFA, M. (2008) Decisions during fights in the house cricket, *Acheta domesticus*: mutual or
self assessment of energy, weapons and size? *Animal Behaviour* **75**, 1053–1062.
BRIFFA, M. & LANE, S.M. (2017) The role of skill in animal contests: a neglected component of
fighting ability. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20171596.
CANDATEN, A., POSSENTI, A.G., MAINARDI, Á.A., DA ROCHA, M.C. & PALAORO, A.V. (2020)
Fighting scars: heavier gladiator frogs bear more injuries than lighter frogs. *Acta Ethologica*.
CARO, T.M., GRAHAM, C.M., STONER, C.J. & FLORES, M.M. (2003) Correlates of horn and antler
shape in bovids and cervids. *Behavioral Ecology and Sociobiology* **55**, 32–41.
CHAPIN, K.J., PEIXOTO, P.E.C. & BRIFFA, M. (2019) Further mismeasures of animal contests: a
new framework for assessment strategies. *Behavioral Ecology* **30**, 1177–1185.
CORN, K.A., MARTINEZ, C.M., BURRESS, E.D. & WAINWRIGHT, P.C. (2021) A Multifunction Trade-
Off has Contrasting Effects on the Evolution of Form and Function. *Systematic Biology*.
CROFTS, S.B., LAI, Y., HU, Y. & ANDERSON, P.S.L. (2019) How do morphological sharpness
measures relate to puncture performance in viperid snake fangs? *Biology Letters* **15**, 20180905.
DALE, S. & SLAGSVOLD, T. (1995) Female Contests for Nest Sites and Mates in the Pied
Flycatcher *Ficedula hypoleuca*. *Ethology* **99**, 209–222.
DARWIN, C. (1871) *The descent of man, and selection in relation to sex.* John Murray, London.

DAVIS, E.B., BRAKORA, K.A. & LEE, A.H. (2011) Evolution of ruminant headgear: a review.
Proceedings of the Royal Society B: Biological Sciences **278**, 2857–2865.
DEL-RE, A.C. (2013) *compute.es*: Compute Effect Size. R Package version 0.2-2. *R Package*.
DENNENMOSER, S. & CHRISTY, J.H. (2013) The Design of a Beautiful Weapon: Compensation for
Opposing Sexual Selection on a Trait with Two Functions. *Evolution* **67**, 1181–1188.
EBERHARD, W.G., RODRÍGUEZ, R.L., HUBER, B.A., SPECK, B., MILLER, H., BUZATTO, B.A. &
MACHADO, G. (2018) Sexual Selection and Static Allometry: The Importance of Function. *The*
Quarterly Review of Biology **93**, 207–250.
EGGER, M., SMITH, G.D., SCHNEIDER, M. & MINDER, C. (1997) Bias in meta-analysis detected by
a simple, graphical test. *BMJ* **315**, 629–634.
EMBERTS, Z., HWANG, W.S. & WIENS, J.J. (2021) Weapon performance drives weapon evolution.
Proceedings of the Royal Society B: Biological Sciences **288**, 20202898.
EMLEN, D.J. (2008) The Evolution of Animal Weapons. *Annual Review of Ecology, Evolution,*
and Systematics **39**, 387–413.
FILICE, D.C.S. & DUKAS, R. (2019) Winners have higher pre-copulatory mating success but
losers have better post-copulatory outcomes. *Proceedings of the Royal Society B: Biological*
Sciences **286**, 20182838.
GOYENS, J., DIRCKX, J. & AERTS, P. (2015) Stag Beetle Battle Behavior and its Associated
Anatomical Adaptations. *Journal of Insect Behavior* **28**, 227–244.
GRAFEN, A. (1987) The logic of divisively asymmetric contests: respect for ownership and the
desperado effect. *Animal Behaviour* **35**, 462–467.
GRAHAM, Z.A., PADILLA-PEREZ, D.J. & ANGILLETTA, M.J. (2020) Virile crayfish escalate
aggression according to body size instead of weapon size. *Animal Behaviour* **163**, 9–15.
GREEN, P.A. & PATEK, S.N. (2015) Contests with deadly weapons: telson sparring in mantis
shrimp (Stomatopoda). *Biology Letters* **11**, 20150558.
HARDY, I.C.W. & BRIFFA, M. (2013) *Animal Contests*. Cambridge University Press.
HOEM, S.A., MELIS, C., LINNELL, J.D.C. & ANDERSEN, R. (2007) Fighting behaviour in territorial
male roe deer *Capreolus capreolus*: the effects of antler size and residence. *European Journal*
of Wildlife Research **53**, 1–8.
HOULE, D., PÉLABON, C., WAGNER, G.P. & HANSEN, T.F. (2011) Measurement and Meaning in
Biology. *The Quarterly Review of Biology* **86**, 3–34.
HUNT, D., BARTON, N., CALDWELL, C., SANDER, C., BOISSEAU, R., ALLEN, C. & EMLEN, D. (2020)
Singing Beetles? Figuring out how male rhinoceros beetles produce their courtship songs.
[Undergraduate Thesis]. *University of Montana Conference on Undergraduate Research*
(UMCUR).
HUSAK, J.F., LAPPIN, A.K. & VAN DEN BUSSCHE, R.A. (2009) The fitness advantage of a high-
performance weapon. *Biological Journal of the Linnean Society* **96**, 840–845.
JOHNS, A., GOTOH, H., MCCULLOUGH, E.L., EMLEN, D.J. & LAVINE, L.C. (2014) Heightened
Condition-Dependent Growth of Sexually Selected Weapons in the Rhinoceros Beetle,
Trypoxylus dichotomus (Coleoptera: Scarabaeidae). *Integrative and Comparative Biology* **54**,
614–621.
KATSUKI, M., YOKOI, T., FUNAKOSHI, K. & OOTA, N. (2014) Enlarged Hind Legs and Sexual
Behavior with Male-Male Interaction in *Sagra femorata* (Coleoptera: Chrysomelidae).
Entomological News **124**, 211–220.

KIM, H., JANG, Y. & CHOE, J.C. (2011) Sexually dimorphic male horns and their use in agonistic
behaviors in the horn-headed cricket *Loxoblemmus doenitzi* (Orthoptera: Gryllidae). *Journal of*
Ethology **29**, 435–441.
KORICHEVA, J., GUREVITCH, J. & MENGERSEN, K. (2013) *Handbook of Meta-analysis in Ecology*
and Evolution. Princeton University Press.
LAPPIN, A.K., BRANDT, Y., HUSAK, J.F., MACEDONIA, J.M. & KEMP, D.J. (2006) Gaping Displays
Reveal and Amplify a Mechanically Based Index of Weapon Performance. *The American*
Naturalist **168**, 100–113.
LAPPIN, A.K. & HUSAK, J.F. (2005) Weapon Performance, Not Size, Determines Mating Success
and Potential Reproductive Output in the Collared Lizard (*Crotaphytus collaris*). *The American*
Naturalist **166**, 426–436.
LIBERATI, A., ALTMAN, D.G., TETZLAFF, J., MULROW, C., GØTZSCHE, P.C., IOANNIDIS, J.P.A.,
CLARKE, M., DEVEREAUX, P.J., KLEIJNEN, J. & MOHER, D. (2009) The PRISMA statement for
reporting systematic reviews and meta-analyses of studies that evaluate health care
interventions: explanation and elaboration. *Journal of Clinical Epidemiology* **62**, e1–e34.
LOBREGAT, G., GECHEL KLOSS, T., PEIXOTO, P.E.C. & SPERBER, C.F. (2019) Fighting in rounds:
males of a neotropical cricket switch assessment strategies during contests. *Behavioral Ecology*
30, 688–696.
MATHOT, K.J., DINGEMANSE, N.J. & NAKAGAWA, S. (2019) The covariance between metabolic
rate and behaviour varies across behaviours and thermal types: meta-analytic insights.
Biological Reviews **94**, 1056–1074.
MAYNARD SMITH, J. & PRICE, G.R. (1973) The Logic of Animal Conflict. *Nature* **246**, 15–18.
MCCULLOUGH, E.L. (2014) Mechanical limits to maximum weapon size in a giant rhinoceros
beetle. *Proceedings of the Royal Society B: Biological Sciences* **281**, 20140696.
MCCULLOUGH, E.L., MILLER, C.W. & EMLEN, D.J. (2016) Why Sexually Selected Weapons Are
Not Ornaments. *Trends in Ecology & Evolution* **31**, 742–751.
MCCULLOUGH, E.L. & SIMMONS, L.W. (2016) Selection on male physical performance during
male–male competition and female choice. *Behavioral Ecology* **27**, 1288–1295.
MCCULLOUGH, E.L., TOBALSKE, B.W. & EMLEN, D.J. (2014) Structural adaptations to diverse
fighting styles in sexually selected weapons. *Proceedings of the National Academy of Sciences*
111, 14484–14488.
MICHONNEAU, F., BROWN, J.W. & WINTER, D.J. (2016) *rotl*: an R package to interact with the
Open Tree of Life data. *Methods in Ecology and Evolution* **7**, 1476–1481.
NAKAGAWA, S. & CUTHILL, I.C. (2007) Effect size, confidence interval and statistical significance:
a practical guide for biologists. *Biological Reviews* **82**, 591–605.
NAKAGAWA, S. & SANTOS, E.S.A. (2012) Methodological issues and advances in biological meta-
analysis. *Evolutionary Ecology* **26**, 1253–1274.
O'BRIEN, D.M. & BOISSEAU, R.P. (2018) Overcoming mechanical adversity in extreme hindleg
weapons. *PLoS ONE* **13**.
O'BRIEN, D.M., KATSUKI, M. & EMLEN, D.J. (2017) Selection on an extreme weapon in the frog-
legged leaf beetle (*Sagra femorata*). *Evolution* **71**, 2584–2598.
OLSSON, M. & SHINE, R. (2000) Ownership influences the outcome of male-male contests in the
scincid lizard, *Niveoscincus microlepidotus*. *Behavioral Ecology* **11**, 587–590.

PALAORO, A.V. & BRIFFA, M. (2017) Weaponry and defenses in fighting animals: how allometry
can alter predictions from contest theory. *Behavioral Ecology* **28**, 328–336.
PALAORO, A.V., PEIXOTO, P.E.C., BENSO-LOPES, F., BOLIGON, D.S. & SANTOS, S. (2020) Fight
intensity correlates with stronger and more mechanically efficient weapons in three species of
Aegla crabs. *Behavioral Ecology and Sociobiology* **74**.
PALAORO, A.V., VELASQUE, M., SANTOS, S. & BRIFFA, M. (2017) How does environment influence
fighting? The effects of tidal flow on resource value and fighting costs in sea anemones. *Biology*
Letters **13**, 20170011.
PARADIS, E., CLAUDE, J. & STRIMMER, K. (2004) *APE*: Analyses of Phylogenetics and Evolution
in R language. *Bioinformatics* **20**, 289–290.
PEIXOTO, P.E.C. & BENSON, W.W. (2008) Body Mass and not Wing Length Predicts Territorial
Success in a Tropical Satyrine Butterfly. *Ethology* **114**, 1069–1077.
PÉLABON, C., FIRMAT, C., BOLSTAD, G.H., VOJE, K.L., HOULE, D., CASSARA, J., ROUZIC, A.L. &
HANSEN, T.F. (2014) Evolution of morphological allometry. *Annals of the New York Academy of*
Sciences **1320**, 58–75.
PINTO, N.S., PALAORO, A.V. & PEIXOTO, P.E.C. (2019) All by myself? Meta-analysis of animal
contests shows stronger support for self than for mutual assessment models. *Biological*
Reviews **94**, 1430–1442.
POLLY, P.D. (2020) Functional Tradeoffs Carry Phenotypes Across the Valley of the Shadow of
Death. *Integrative and Comparative Biology* **60**, 1268–1282.
POTTER, D.A., WRENSCH, D.L. & JOHNSTON, D.E. (1976) Aggression and Mating Success in
Male Spider Mites. *Science* **193**, 160–161.
RICO-GUEVARA, A. & ARAYA-SALAS, M. (2015) Bills as daggers? A test for sexually dimorphic
weapons in a lekking hummingbird. *Behavioral Ecology* **26**, 21–29.
RICO-GUEVARA, A. & HURME, K.J. (2019) Intrasexually selected weapons. *Biological Reviews*
94, 60–101.
ROBINSON, C.D. & GIFFORD, M.E. (2019) Sexual dimorphism in performance and muscle
allocation in the western painted crayfish *Faxonius palmeri longimanus* (Faxon, 1898)
(Decapoda: Astacidea: Cambaridae). *Journal of Crustacean Biology* **39**, 267–273.
ROHATGI, A. (2019) *WebPlotDigitizer* 4.2: HTML5 based online tool to extract numerical data
from plot images. *Https://Automeris.Io/WebPlotDigitizer.*
SZÁMADÓ, S. (2008) How threat displays work: species-specific fighting techniques, weaponry
and proximity risk. *Animal Behaviour* **76**, 1455–1463.
TAYLOR, J.R.A. & PATEK, S.N. (2010) Ritualized fighting and biological armor: the impact
mechanics of the mantis shrimp's telson. *Journal of Experimental Biology* **213**, 3496–3504.
UMBERS, K.D., TATARNIC, N.J., HOLWELL, G.I. & HERBERSTEIN, M.E. (2012) Ferocious fighting
between male grasshoppers. *PLoS One.*
VIEIRA, M.C. & PEIXOTO, P.E.C. (2013) Winners and losers: a meta-analysis of functional
determinants of fighting ability in arthropod contests. *Functional Ecology* **27**, 305–313.
WAINWRIGHT, P.C., ALFARO, M.E., BOLNICK, D.I. & HULSEY, C.D. (2005) Many-to-One Mapping
of Form to Function: A General Principle in Organismal Design? *Integrative and Comparative*
Biology **45**, 256–262.

WEAVER, R.J., SANTOS, E.S.A., TUCKER, A.M., WILSON, A.E. & HILL, G.E. (2018) Carotenoid
metabolism strengthens the link between feather coloration and individual quality. *Nature*
Communications **9**, 1–9.
WILSON, R.S., ANGILLETTA JR., M.J., JAMES, R.S., NAVAS, C. & SEEBACHER, F. (2007) Dishonest
Signals of Strength in Male Slender Crayfish (*Cherax dispar*) during Agonistic Encounters. *The*
American Naturalist **170**, 284–291.
YOSHINO, K., KOGA, T. & OKI, S. (2011) Chelipeds are the real weapon: cheliped size is a more
effective determinant than body size in male–male competition for mates in a hermit crab.
Behavioral Ecology and Sociobiology **65**, 1825.

Table 1. Definitions used throughout this paper.

Term	Definition
Weapon	Specialized morphology used to gain mechanical advantage on rivals during physical contact.
Display	Any behaviour or movement that can be used as a cue of fighting ability, or as a threat behaviour. In most species, it is the part of the contest that does not involve physical contact. In the species where rivals are in physical contact, it is associated with physical contact of low intensity, such as antennae touching.
Fight	Part of the contest where the individuals are within reach of one another and are in physical contact. Can be seen as a physical struggle between two rivals.
Contest	Aggressive competition between two rivals of the same species for an indivisible resource. Can be composed of two phases: a display phase and a physical contact phase.
Fighting style	Behaviours performed during physical contact that aim to gain mechanical advantage over the rival. Can be composed of either a single or multiple behaviour used simultaneously. Frequently associated with movements of the weapon itself (e.g., squeezing).
Weapon function	The combination of display and fighting style an individual uses during a contest. See Table 3 and Table S1 for a detailed list of weapon function.

868   **Table 2**. Heterogeneity of the multilevel meta-analytical model of differences in how weapons

869   are measured.

Random effect	I² (95% CI)
Study ID	10.93 (9.009, 12.85)
Pairing method	17.37 (15.44, 19.29)
Phylogeny	0.012 (0.00, 1.92)
Genus	0.0001 (0.00, 1.92)
Total	23.71 (21.79, 25.63)
H^2	<0.001 (0.00, 1.92)

871 **Table 3.** Functions (lines) and displays (columns) of fighting styles found across the 52 species

872 of animals used in our study. All possible combinations of the five functions were considered,

873 even though some of them were absent for our sample. To see how each species was

874 categorized, please see Fig. 1 and Supplementary File S2 for the descriptions.

FUNCTION	WEAPON DISPLAY	BODY SIZE ESTIMATION	NO DISPLAY	TOTAL
Squeezing	12	1	-	13
Impact	1	1	-	2
Pushing	1	-	-	1
Lift + Squeezing	1	1	1	3
Lift + Impact	3	-	1	4
Lift + Pushing	2	1	1	4
Squeezing + Impact	3	-	-	3
Squeezing + Piercing	-	1	-	1
Squeezing + Pushing	1	5	1	7
Squeeze + Pull	1	2	-	3
Impact + Pushing	-	3	-	3
Piercing + Pushing	1	-	-	1
Push + Impact + Lift	2	-	-	2
Push + Impact + Squeeze	1	1	-	2
Push + Lift + Squeeze	1	1	-	2
Push + Squeeze + Pull + Lift	1	-	-	1
TOTAL	31	18	3	52

877 **Table 4.** Heterogeneity of the meta-analytical model of differences in how contests begin (i.e., if

878 displays are used, and which type of displays).

Random effect	I² (95% CI)
Study ID	9.75 (7.83, 11.67)
Pairing method	17.93 (16.01, 19.85)
Phylogeny	5.84 (3.92, 7.76)
Genus	< 0.0001 (0.00, 1.92)
Total	33.54 (31.62, 35.46)
H²	0.17 (0.00, 2.09)

880 **Table 5.** Heterogeneity and variance of the meta-analytical model of differences in how

881 weapons are used during contests.

Random effect	I² (95% CI)
Study ID	6.91 (4.99, 8.83)
Pairing method	18.46 (16.54, 20.38)
Phylogeny	< 0.0001 (0.00, 1.92)
Genus	< 0.0001 (0.00, 1.92)
Total	25.38 (23.46, 27.30)
H²	< 0.0001 (0.00, 1.92)

FIGURE LEGENDS

Fig. 1. Hedges' g value (circle) and corresponding confidence intervals (lines) for species,

pairing method (whether contests were paired for body size or not), component measured (e.g.,

length, width, mass), type of display (i.e., displaying weapon, body size, or not displaying) and

function of the weapon during the contests (see topic 3). Study demonstrates which samples

were taken from the same study (references in Supplementary File S1). Larger grey circles

denote studies with larger sample sizes. Silhouettes were taken from phylopic. Grey rows were

selected randomly to facilitate visual separation of groups. For more information regarding the

studies and pairing methods, see Fig. S3.

Fig 2. Weapons had an overall positive effect on contest success with winners having greater

trait values than losers. Linear measures had a higher difference between winners and losers

than performance measures demonstrating that winners tend to have much larger, rather than

stronger weapons, than losers (QM_1 = 23.29, p < 0.001). The effect size, Hedges' g, represents the

mean standardized difference between winners and losers. Positive values denote that winners

were larger than losers, while negative values represent the opposite. Dots represent the

estimated values from a multilevel meta-analytic model considering Hedges' g as the response

variable, weapon component as a moderator variable, and study ID, species ID, phylogeny,

environment and pairing method as random variables. The error bars represent the 95%

confidence interval of the estimate. The numbers above the error bars represent the number of

effect sizes in each component.

Fig. 3. Winners had larger weapons than losers regardless of the type of display, but these

differences were similar among the types of display. The effect size, Hedges' g, represents the

mean standardized difference in linear measures between winners and losers. Positive values

denote that winners had larger weapons than losers, while negative values represent the

opposite. Dots represent the estimated values from a multilevel meta-analytic model

considering Hedges' g as the response variable and the type of display as a moderator variable.

The error bars represent the 95% confidence interval of the estimate. The numbers above the

error bars represent the number of effect size in each component.

Fig. 4. Winners had weapons that were, on average, larger than losers; and asymmetries

between winners and losers in performance-emphasis were lower than the other two fighting

styles. The effect size, Hedges' g, represents the mean standardized difference in the linear

measures of weapons between winners and losers. Positive values denote that winners had

larger weapons than losers, while negative values represent the opposite. Dots represent the

estimated values from a multilevel meta-analytic model considering Hedges' g as the response

variable and the fighting style as a moderator variable. The error bars represent the 95%

confidence interval of the estimate. The numbers above the error bars represent the number of

effect size in each component.

Species	Measure	Display	Fighting style	Hedges g [95% CI]
Teleogryllus commodus	Performance	Weapon	Impact & Lift	0.43 [−0.02, 0.89]
Pachyrhamma waitomoensis	Linear	Body	Push & Impact	1.00 [0.57, 1.44]
Melanotes ornata	Linear	Body	Push & Impact	0.64 [0.35, 1.09]
Loxoblemmus doenitzi	Linear	Body	Push & Impact	0.36 [−0.32, 1.04]
Kosciuscola tristis	Linear	Weapon	Squeeze	0.27 [−0.22, 0.75]
Hemideina crassidens	Linear	Weapon	Pull & Squeeze	1.03 [0.44, 1.62]
Gryllus pennsylvanicus	Linear	Weapon	Push & Squeeze	0.81 [0.13, 1.49]
Gryllus pennsylvanicus	Linear	Weapon	Push & Squeeze	0.67 [0.15, 1.19]
Gryllus pennsylvanicus	Linear	Weapon	Push & Squeeze	0.24 [−0.21, 0.68]
Gryllus pennsylvanicus	Linear	Weapon	Push & Squeeze	0.22 [−0.22, 0.66]
Gryllus pennsylvanicus	Linear	Weapon	Push & Squeeze	0.25 [−0.20, 0.70]
Gryllus pennsylvanicus	Asymmetry	Weapon	Push & Squeeze	0.00 [−0.44, 0.44]
Gryllus firmus		Weapon	Lift	0.53 [0.29, 0.77]
Acheta domesticus	Linear	Weapon	Impact & Lift	−0.05 [−0.53, 0.43]
Acheta domesticus	Linear	Weapon	Impact & Lift	0.29 [−0.19, 0.78]
Rangifer tarandus	Linear	Weapon	Push, Impact & Lift	3.29 [1.25, 5.33]
Rangifer tarandus	Linear	Weapon	Push, Impact & Lift	1.38 [0.48, 2.28]
Hipposideros armiger	Linear	Body	Impact	0.09 [−0.26, 0.44]
Capreolus capreolus	Index	Weapon	Push, Impact & Lift	2.34 [1.38, 3.30]
Gonatodes albogularis	Linear	Weapon	Squeeze	0.36 [−0.24, 0.96]
Gonatodes albogularis	Linear	Weapon	Squeeze	0.08 [−0.51, 0.68]
Gallotia galloti	Linear	Weapon	Squeeze	0.28 [−0.35, 0.90]
Gallotia galloti	Performance	Weapon	Squeeze	0.36 [0.09, 0.63]
Gallotia galloti	Linear	Weapon	Squeeze	0.69 [0.11, 1.28]
Gallotia galloti	Linear	Weapon	Squeeze	0.02 [−0.49, 0.54]
Ctenophorus maculosus	Performance	Weapon	Squeeze	0.41 [0.07, 0.75]
Ctenophorus maculosus	Linear	Weapon	Squeeze	0.32 [−0.02, 0.66]
Carinascincus microlepidotus	Linear	Body	Squeeze	−0.09 [−0.61, 0.43]
Carinascincus microlepidotus	Linear	Body	Squeeze	0.21 [−0.31, 0.74]
Anolis valencienni	Performance	Weapon	Squeeze	0.19 [−0.49, 0.86]
Anolis sagrei	Performance	Weapon	Squeeze	0.47 [−0.11, 1.05]
Anolis lineatopus	Performance	Weapon	Squeeze	0.36 [−0.25, 0.97]
Anolis gundlachi	Performance	Weapon	Squeeze	0.43 [−0.23, 1.10]
Anolis evermanni	Performance	Weapon	Squeeze	0.23 [−0.34, 0.80]
Anolis distichus	Performance	Weapon	Squeeze	−0.35 [−0.95, 0.25]
Anolis cristatellus	Performance	Weapon	Squeeze	0.24 [−0.35, 0.84]
Anolis carolinensis	Performance	Weapon	Squeeze	0.51 [−0.14, 1.16]
Anolis angusticeps	Performance	Weapon	Squeeze	0.44 [−0.19, 1.07]
Neogonodactylus bredini	Performance	Weapon	Impact	−0.04 [−0.51, 0.44]
Riptortus pedestris	Linear	Body	Push, Squeeze & Impact	1.75 [1.18, 2.32]
Narnia femorata	Linear	Weapon	Impact & Squeeze	0.57 [−0.85, 1.99]
Narnia femorata	Linear	Weapon	Impact & Squeeze	0.69 [−0.75, 2.13]
Narnia femorata	Linear	Weapon	Impact & Squeeze	0.03 [−0.85, 0.92]
Narnia femorata	Linear	Weapon	Impact & Squeeze	0.10 [−0.78, 0.99]
Narnia femorata	Linear	Weapon	Impact & Squeeze	0.42 [−0.47, 1.32]
Narnia femorata	Linear	Weapon	Impact & Squeeze	0.43 [−0.46, 1.33]
Narnia femorata	Area	Weapon	Impact & Squeeze	0.78 [−0.67, 2.23]
Narnia femorata	Area	Weapon	Impact & Squeeze	0.07 [−0.81, 0.96]
Narnia femorata	Area	Weapon	Impact & Squeeze	0.48 [−0.42, 1.38]
Anisolabis maritima	Asymmetry	Weapon	Impact & Squeeze	0.26 [−0.74, 1.26]
Anisolabis maritima	Asymmetry	Weapon	Impact & Squeeze	1.31 [0.37, 2.25]
Scopimera globosa	Linear	Weapon	Push	0.85 [0.41, 1.29]
Parastacus pilimanus	Linear	Weapon	Push, Pull & Squeeze	−0.11 [−1.25, 1.03]
Parastacus brasiliensis	Linear	Weapon	Push, Pull & Squeeze	0.04 [−1.02, 1.10]
Faxonius obscurus	Performance	Body	Push & Squeeze	0.76 [0.38, 1.14]
Faxonius obscurus	Performance	Body	Push & Squeeze	−0.17 [−0.55, 0.20]
Faxonius obscurus	Linear	Body	Push & Squeeze	0.87 [0.49, 1.25]
Faxonius obscurus	Linear	Body	Push & Squeeze	0.86 [0.47, 1.25]
Diogenes nitidimanus	Linear	No	Push & Squeeze	0.46 [−0.02, 0.94]
Cherax dispar	Performance	Weapon	Push & Squeeze	0.73 [0.21, 1.25]
Cherax dispar	Linear	Weapon	Push & Squeeze	0.43 [−0.05, 0.91]
Carcinus maenas	Linear	Weapon	Push, Squeeze & Impact	0.45 [−0.22, 1.13]
Carcinus maenas	Linear	Weapon	Push, Squeeze & Impact	0.33 [−0.34, 1.00]
Carcinus maenas	Performance	Weapon	Push, Squeeze & Impact	0.59 [0.04, 1.14]
Carcinus maenas	Mass	Weapon	Push, Squeeze & Impact	0.41 [−0.34, 1.17]
Carcinus maenas	Mass	Weapon	Push, Squeeze & Impact	0.34 [−0.20, 0.88]
Carcinus maenas	Linear	Weapon	Push, Squeeze & Impact	0.61 [0.02, 1.20]
Carcinus maenas	Area	Weapon	Push, Squeeze & Impact	0.50 [−0.26, 1.26]
Cambarus robustus	Performance	Body	Push & Squeeze	1.01 [0.54, 1.47]
Cambarus robustus	Performance	Body	Push & Squeeze	−0.11 [−0.54, 0.33]
Cambarus robustus	Linear	Body	Push & Squeeze	1.81 [1.29, 2.34]
Cambarus robustus	Linear	Body	Push & Squeeze	1.70 [1.18, 2.21]
Cambarus robustus	Linear	Body	Push & Squeeze	0.38 [−0.16, 0.92]
Cambarus robustus	Linear	Body	Push & Squeeze	0.35 [−0.19, 0.89]
Cambarus robustus	Linear	Body	Push & Squeeze	0.37 [−0.17, 0.91]
Cambarus robustus	Linear	Body	Push & Squeeze	0.40 [−0.14, 0.94]
Cambarus carinirostris	Performance	Body	Push & Squeeze	0.80 [0.39, 1.21]
Cambarus carinirostris	Performance	Body	Push & Squeeze	0.25 [−0.15, 0.64]
Cambarus carinirostris	Linear	Body	Push & Squeeze	1.38 [0.95, 1.82]
Cambarus carinirostris	Linear	Body	Push & Squeeze	1.31 [0.88, 1.75]
Austruca annulipes	Linear	Weapon	Squeeze & Lift	0.50 [0.22, 0.78]
Aegla longirostri	Performance	Body	Pull & Squeeze	1.27 [0.47, 2.07]
Aegla longirostri	Linear	Body	Pull & Squeeze	1.22 [0.43, 2.01]
Aegla longirostri	Linear	Body	Pull & Squeeze	1.28 [0.48, 2.08]
Aegla longirostri	Performance	Body	Pull & Squeeze	−0.00 [−0.72, 0.72]
Aegla longirostri	Linear	Body	Pull & Squeeze	0.01 [−0.71, 0.73]
Aegla longirostri	Linear	Body	Pull & Squeeze	0.03 [−0.69, 0.75]
Trypoxylus dichotomus	Linear	No	Push & Lift	0.34 [−0.05, 0.73]
Trypoxylus dichotomus	Linear	No	Push & Lift	1.47 [0.76, 2.18]
Trypoxylus dichotomus	Linear	No	Push & Lift	0.22 [−0.37, 0.81]
Sagra femorata	Linear	No	Squeeze & Lift	0.45 [0.16, 0.73]
Sagra femorata	Linear	No	Squeeze & Lift	0.29 [0.00, 0.57]
Onthophagus taurus	Linear	Weapon	Push & Lift	1.04 [0.49, 1.59]
Onthophagus acuminatus	Linear	Weapon	Push & Lift	0.70 [0.15, 1.25]
Librodor japonicus	Linear	Weapon	Push, Squeeze & Pry	0.81 [0.37, 1.25]
Librodor japonicus	Linear	Weapon	Push, Squeeze & Pry	0.80 [0.04, 1.56]
Librodor japonicus	Linear	Weapon	Push, Squeeze & Pry	0.14 [−0.64, 0.92]
Heterochelus chiragricus	Linear	Weapon	Squeeze & Lift	0.74 [0.02, 1.46]
Heterochelus chiragricus	Linear	Weapon	Squeeze & Lift	0.83 [0.60, 1.06]
Gnatocerus cornutus	Linear	Weapon	Push, Squeeze & Pry	0.56 [0.08, 1.05]
Dicronocephalus wallichii	Linear	Body	Push & Lift	2.53 [1.24, 3.82]
Aegus chelifer	Linear	Body	Push, Squeeze & Pry	0.78 [0.49, 1.08]
Aegus chelifer	Linear	Body	Push, Squeeze & Pry	0.74 [0.44, 1.03]
Aegus chelifer	Linear	Body	Push, Squeeze & Pry	0.73 [−0.19, 1.65]
Ficedula hypoleuca	Linear	Body	Squeeze & Piercing	0.24 [−0.65, 1.12]
Cambridgea foliata	Linear	Body	Push	0.59 [0.24, 0.94]
Tetranychus urticae	Linear	Weapon	Push & Piercing	1.35 [0.22, 2.48]

Hedges g