The hidden links between animal weapons, fighting style and their expression on contest

resolution

Alexandre V. Palaoro[1,2*] & Paulo E. C. Peixoto[3]

[1]LUTA do Departamento de Ecologia e Biologia Evolutiva, Universidade Federal de São Paulo,

Brazil

[2]LAGE do Departamento de Ecologia, Universidade de São Paulo, Brazil

[3]LASEXIA, Departamento de Genética, Ecologia e Evolução, Universidade Federal de Minas

Gerais, Brazil

* Corresponding author: LUTA do Departamento de Ecologia e Evolução, Universidade Federal

de São Paulo, São Paulo, Brazil. Rua Artur Riedel, 275, Eldorado, Diadema, CEP 09972-270. E-

mail: alexandre.palaoro@gmail.com

SUMMARY

1. In many species that fight over resources, individuals use specialized structures to

overpower their rivals (i.e. weapons). Despite their similar roles for contest settlement

(i.e. affecting the winning chances), weapons are highly diverse morphological

structures across species. However, the comprehension on how this diversity evolved is

still open for debate.

2. Unfortunately, most studies on how weapons are used during contests focus on size

asymmetries between winners and losers. Although such information is important to

understand, it does not provide much information regarding the mechanisms promoting

weapon diversification. In addition, in many situations, the effect of weapon size on

contest resolution may be confounded with the effect of body size due to allometric

correlations.

3. One way to decompose the relative importance of weapon and body size on contest

outcome is to focus on the functional role of weapons. That is, investigate the function of

the weapon during contests, and how size should affect its function.

4. We propose two hypotheses that explicitly associate how weapon function may

determine the relative importance of weapon and body size in affecting rivals during

contests. One based on the internal morphology of the weapon – whether muscles are

within the weapon or not – and the other based on fighting style – whether the weapon

is used to injure the rival or not.

5. To test our hypotheses, we developed a meta-analysis spamming 31 genera across the

animal kingdom.

6. We found that internal morphology and fighting style are important to determine the

relative importance of weapon and body size on contest resolution. Weapons that do not

have muscles within them (e.g., horns) affect more the chance of winning than weapons

that have muscles within them (e.g., claws). Similarly, weapons used to throw/push

rivals away (e.g., oversized mandibles) affect more the chance of winning than those

used to make the rival flee (e.g., jaws).

7. Overall, we have shown that weapon usage (i.e. weapon function) may affect weapon

evolution in predictable ways: weapons without muscles and using for pushing may be

selected for size, while muscled weapons used for injuring may be selected for strength.

Keywords. Animal contests, morphological evolution, animal fighting, sexual selection, weapon

biomechanics

INTRODUCTION

Antlers, claws, horns, and enlarged mandibles are just a few of a much larger diversity of

animal weapons we have in the animal kingdom. These weapons are used during contests to

overpower rivals when individuals fight for a resource (McCullough, Miller, & Emlen, 2016).

Consequently, weapons are said to influence the outcome of a contest (Palaoro & Briffa, 2017).

Given that winners secure more, and better, resources than losers, fighting influences fitness of

the individual (Pinto, Palaoro, & Peixoto, 2019; Vieira & Peixoto, 2013). This link between

bearing weapons, winning contests, and fitness is said to explain the wide variety of weapons in

the animal kingdom (Emlen, 2008; Rico-Guevara & Hurme, 2019). However, most studies focus

on how weapon size asymmetries between winners and losers affects contest outcome (Pinto,

Palaoro, & Peixoto, 2019; Vieira & Peixoto, 2013). However, when looking only at the size

asymmetries, we lose some of the information on how the weapon is used during the contest to

affect the rival (i.e. the functional role of the weapon during the contest; Eberhard et al., 2018).

Without information on the functional role of the weapon, we may bias our comprehension on

the magnitude of the effect of weapon size in contest outcome and on how weapons evolve and

diversify across animal taxa (Eberhard et al., 2018; McCullough et al., 2016).

One of the challenges in detecting the effect of weapons on contest resolution stem from

the allometric correlation between weapons and body size (Palaoro & Briffa, 2017, Pinto,

Palaoro, & Peixoto, 2019; Vieira & Peixoto, 2013). Since the allometric correlation between

weapon and body size is strong for most species (Voje, 2016), one may wonder that both

variables should provide similar information about how individual traits affect winning

probability. However, body size also affects contest outcome independent of weapon size,

especially when we consider contests between species that do not bear weapons (e.g., butterflies,

dragonflies, Vieira & Peixoto, 2013). Without accurate estimations of the relative effect of

weapon and body size on contest resolution, we may be unable to identify the selective

pressures acting on each trait. Here, we test the impact of considering the weapon's functional

role to improve our knowledge about the relative effect of weapon and body size on contest

resolution. For this, we use information on weapon internal morphology and on the behaviors

adopted by the rivals during contests. Below, we propose two hypotheses that merge weapon

functional role and the relative effect of weapon and body size for contest resolution. First, we

present the biomechanical hypothesis, in which the morphology of the weapon might reveal

their overall importance in contests. Second, we present the functional hypothesis, in which

how the weapon is used during contests - i.e., the fighting style - matters for their importance in

contest resolution.

Biomechanical hypothesis

We propose that weapons may be separated in two major groups based on their internal

morphology (hereafter referred to as "morphology" for simplicity): 1) weapons that have

specific (generally internal) muscles responsible for the action of the weapon (from now on,

muscled weapons) and 2) weapons that are not attached to a particular set of muscles (from

now on, muscleless weapons). Muscled weapons are comprised by jaws, claws, mandibles, and

legs, and they generally work by one part of the weapon closing on the rival and squeezing it or

by producing an explosive movement to strike the rival (Emlen, 2008). Muscled weapons may

be weakly dependent on individual body size to overpower rivals. For instance, whenever a

crab closes its' claw on a rival, it requires only the action of the closer muscle of the claw; no

other muscle is needed to grab the rival and win the contest (Dennenmoser & Christy, 2013;

Morrell, Backwell, & Metcalfe, 2005). The same pattern is found in the mandibles of stag beetles,

in which the closing muscle closes the jaws on the rival (Goyens, Dirckx, & Aerts, 2015). The

body size, in these instances, will function mainly as a stabilizer of the whole body, working as

an auxiliary trait (Goyens et al., 2015; Okada, Suzaki, Okada, & Miyatake, 2011). Hence, for

species that possesses muscled weapons, the differences between winners and losers should be

greater for weapon size than for body size.

Muscleless weapons are mainly represented by horns and antlers (Emlen, 2008). These

weapons are often used to enter in direct contact with the rival without the need to squeeze or

strike. The entire body, in these cases, is responsible for overpowering the rival - the weapon is

a tool by which the power of the whole body acts upon. For instance, when male deer lock

antlers, the antler is used mainly as a point of contact, a lever, by which the individual transfer

the power generated by their entire body. It is thus the body that is doing all the pushing

(Jennings & Gammell, 2013). For species possessing muscleless weapons, the main trait used to

overpower rivals is the body, while the weapon becomes the auxiliary trait. Hence, in

muscleless weapons, we expect winners and losers to differ more strongly on body size than

weapon size. Therefore, the biomechanical hypothesis posits that the internal morphology of

the weapon (i.e., presence or absence of auxiliary muscles) can shift the relative importance of

weapon and body size to win fights.

Functional hypothesis

We suggest that the importance of the weapon to win contests should be correlated to how the

weapon is used during a fight - the fighting style of the species (Eberhard et al., 2018;

McCullough et al., 2016). Broadly speaking, weapons can are used in contests in two ways: 1)

weapons mainly used to grasp, kick, strike, pierce or squeeze until the loser decides to back

away from the winner (from now on, biting weapons) and 2) weapons used to forcefully

remove the loser from the resource without (necessarily) causing direct injuries (from now on,

pushing weapons). Biting weapons make the rival flee based on the injuries caused by a direct

strike of the weapon (Green & Patek, 2018; Palaoro & Briffa, 2017). For instance, mantis shrimp

strike their rivals using their high-performance raptorial appendages that are able to crack the

shells of molluscs and crabs (Green & Patek, 2015). When individuals are contesting a burrow,

the loser withdraws from a fight after being hit by the rival's powerful strike - typically leaving

the contest before any major injuries are done (Green & Patek, 2018). Leaf-footed bugs follow a

similar pattern, but with a different weapon. The individuals squeeze each other using their

enlarged hind legs until the eventual loser decides to back away from the contested resource

(Nolen, Allen, & Miller, 2017). In both cases, the rival is deciding to retreat after receiving direct

strikes of the weapon. Therefore, for biting weapons, we expect that the weapon should be

more important to settle the contest than the body size of the individual.

Pushing weapons are used, for example, by some beetle species that contest burrows or

tree saps. In such species, the winner uses its horns, enlarged mandibles, or hind legs, to push

the loser away from the burrow or dislodge the loser from the female/tree branch (McCullough

& Simmons, 2016; O'Brien, Katsuki, & Emlen, 2017; Rink, Altwegg, Edwards, Bowie, & Colville,

2019). That is especially true for stag beetles, which use their enlarged mandibles to lift the rival

and throw him away from the resource - they do not use their mandibles to bite the rival into

submission (Goyens et al., 2015). Therefore, although such mandibles have specific muscles

attached to them, they are not used to injury the rival. A similar pattern occurs in cervid fights,

in which individuals are mainly pushed away from the resource (Jennings & Gammell, 2013).

Due to the emphasis in pushing or shoving rivals away, the trait that should enable individuals

to dislodge rivals is the body, rather than the weapon; the weapon is only transmitting the

power generated by the body. Therefore, in injury fights the main factor affecting contest

resolution should be the biting weapon and its' ability to cause injuries, whereas in pushing

fights the main factor determining victory should be the body and its' ability to push the rival

away.

Despite the existence of marked differences in weapon morphology and fighting style,

and on the growing theoretical interest in how these two aspects can influence weapon

evolution (Eberhard et al., 2018; McCullough et al., 2016), there is no evaluation of how weapon

morphology and fighting style might affect the relative importance of body and weapon traits

in animal contests. Thus, by focusing on the weapon functional role during the contest

(according to our categorization of weapons based on morphology and fighting style), we may

be able to provide ways to disentangle how agonistic interactions affect the evolution of body

size and weapon morphology. However, weapons are frequently used during contests

regardless of morphology and fighting style. Because of their frequent use, it might be that the

overall effect of weapons on winning chances is always higher than body size. For this reason,

we used a meta-analytical approach to first test the common prediction that weapons are more

important than body size for contest resolution (Palaoro & Briffa, 2017). Next, we evaluated the

biomechanical hypothesis that posits that muscled weapons (i.e., weapons that have muscles

attached to them) should be more important than body size, whereas muscleless weapons (i.e.,

weapons that do not have muscles attached) should be less (or equally) important than body

size in determining the winner. Lastly, we evaluated the functional hypothesis, which posits

that weapons with the main function of injuring rivals (i.e., biting weapons) should be more

important than body size, whereas weapons that dislodge rivals from the resource (i.e., pushing

weapons) should be less (or equally) important than body size in determining the winner. To

measure the importance of each trait in contest resolution, we used a standardized measure

(Hedges' g) of the mean difference between winners and losers for that trait (Vieira & Peixoto,

2013). Based on this measure, we expect larger differences between winners and losers in traits

that should be more important to resolve fights.

METHODS

Literature search and eligibility criteria

We searched for articles using the Web of Science (https://www.webofknowledge.com) and

Scopus (https://scopus.com) databases. For both searches we used the search strings shown in

the Supplementary file. We obtained a total of 1108 papers through those searches and we also

added 12 relevant papers cited in reviews that we did not find in our primary searches (Emlen,

2008; Pinto et al., 2019; Rico-Guevara & Hurme, 2019; Vieira & Peixoto, 2013). We excluded all

studies in which the species did not bear a weapon (i.e., specialized morphology used to gain

mechanical advantage on the rival, *sensu* (Eberhard et al., 2018)). Since we had to classify

species according to their fighting style, we excluded all species in which the behaviours

adopted during the fights were undescribed. For the selected studies, we searched for

information about mean trait values (and their corresponding variation) for winners and losers

of contests. Therefore, we also excluded any study that reported only the mean trait size

without distinguishing winners and losers. Within each study, we searched for the pairing

method used by researchers. If individuals with similar body or weapon sizes were paired, we

excluded that study from our analysis because pairing influences the relative importance of

fighting ability on fight outcome (Palaoro & Briffa, 2017). Another source of confounding effect

might come from resource value (*sensu* (Arnott & Elwood, 2008)): individuals that value more a

resource often fight harder regardless of their fighting ability (e.g., (Palaoro, Velasque, Santos, &

Briffa, 2017)). To avoid any bias, we only included the treatments in which there was no

evidence that individuals had different motivations to fight. Lastly, we only included replicates

in which there is a clear winner and loser – we excluded fights that ended in a tie.

Categorizing weapon internal morphology and fighting style

We categorized weapon morphology based on the position of the muscle relative to the weapon

(Fig. 1). For each selected species, we considered that a given structure was a weapon if there

was a clear description that the structure was used in the fight to overcome rivals (Eberhard et

al., 2018). We considered claws, jaws, mandibles and legs as muscled weapons because all move

independently of the body and are attached to specific muscles nearby. For example,

individuals with claws need to grasp their rivals and grasping strength depend on the muscles

inside the claw (Dennenmoser & Christy, 2013). In contrast, we considered horns and antlers as

muscleless weapons because they are unable to perform moves independent of the body (or any

movement).

To categorize fighting style, we searched for a description of how the individuals use the

weapon during fights and of how the decision to withdrawal occurs (McCullough et al., 2016;

Rico-Guevara & Hurme, 2019). If individuals used their weapons to push, shove, or dislodge

the rival from the resource and if most fights were terminated by excluding the rival from the

resource, we classified as 'pushing weapons'. If the weapons were used to grab, bite, kick,

pierce or squeeze the rival and if most fights were terminated by the loser retreating from the

resource without being throwed away, we classified them as 'biting weapons'. All

classifications and more detailed justifications can be found in Table S1.

Species	Fighting style	Trait	N	Weapon morphology	Hedges g [95% CI]
Loxoblemmus doenitzi	Pushing	Body	35	Muscled	1.07 [0.31, 1.83]
Gryllus firmus	Pushing	Body	140	Muscled	0.59 [0.35, 0.83]
Gryllus firmus	Pushing	Body	139	Muscled	0.46 [0.22, 0.70]
Pachyrhamma waitomoensis	Pushing	Body	45	Muscled	0.29 [−0.12, 0.70]
Melanotes ornata	Pushing	Body	39	Muscled	0.56 [0.11, 1.02]
Melanotes ornata	Pushing	Body	39	Muscled	0.71 [0.25, 1.17]
Acheta domesticus	Pushing	Body	33	Muscled	0.80 [0.30, 1.31]
Group mean effect					0.56 [0.40, 0.72]
Loxoblemmus doenitzi	Pushing	Weapon	35	Muscled	0.36 [−0.32, 1.04]
Gryllus firmus	Pushing	Weapon	140	Muscled	0.53 [0.29, 0.77]
Pachyrhamma waitomoensis	Pushing	Weapon	45	Muscled	1.00 [0.57, 1.44]
Melanotes ornata	Pushing	Weapon	39	Muscled	0.64 [0.18, 1.09]
Acheta domesticus	Pushing	Weapon	33	Muscled	−0.05 [−0.53, 0.43]
Acheta domesticus	Pushing	Weapon	33	Muscled	0.29 [−0.19, 0.78]
Group mean effect					0.53 [0.23, 0.82]
Kosciuscola tristis	Injury	Body	40	Muscled	0.28 [−1.26, 1.81]
Kosciuscola tristis	Injury	Body	40	Muscled	0.23 [−0.26, 0.71]
Group mean effect					0.23 [−0.23, 0.69]
Kosciuscola tristis	Injury	Weapon	40	Muscled	0.27 [−0.22, 0.75]
Hemideina crassidens	Injury	Weapon	57	Muscled	1.03 [0.44, 1.62]
Group mean effect					0.63 [−0.11, 1.38]
Rangifer tarandus	Pushing	Weapon	15	Muscleless	3.29 [1.25, 5.33]
Rangifer tarandus	Pushing	Weapon	29	Muscleless	1.38 [0.48, 2.28]
Capreolus capreolus	Pushing	Weapon	40	Muscleless	2.34 [1.38, 3.30]
Group mean effect					1.97 [1.34, 2.59]
Niveoscincus microlepidotus	Injury	Body	30	Muscled	0.09 [−0.43, 0.61]
Niveoscincus microlepidotus	Injury	Body	30	Muscled	0.33 [−0.20, 0.85]
Gallotia galloti	Injury	Body	20	Muscled	0.65 [0.01, 1.29]
Gonatodes albogularis	Injury	Body	22	Muscled	0.38 [−0.22, 0.98]
Gonatodes albogularis	Injury	Body	22	Muscled	0.59 [−0.02, 1.20]
Group mean effect					0.38 [0.06, 0.69]
Niveoscincus microlepidotus	Injury	Weapon	30	Muscled	−0.09 [−0.61, 0.43]
Niveoscincus microlepidotus	Injury	Weapon	30	Muscled	0.21 [−0.31, 0.74]
Gallotia galloti	Injury	Weapon	20	Muscled	0.28 [−0.35, 0.90]
Gonatodes albogularis	Injury	Weapon	22	Muscled	0.36 [−0.24, 0.96]
Gonatodes albogularis	Injury	Weapon	22	Muscled	0.08 [−0.51, 0.68]
Group mean effect					0.15 [−0.10, 0.41]
Riptortus pedestris	Pushing	Body	33	Muscled	1.45 [0.89, 2.00]
Riptortus pedestris	Pushing	Weapon	33	Muscled	1.75 [1.18, 2.32]
Group mean effect					−x−
Narnia fermorata	Injury	Body	6	Muscled	0.60 [−0.57, 1.77]
Narnia fermorata	Injury	Body	11	Muscled	0.22 [−0.62, 1.07]
Narnia fermorata	Injury	Body	13	Muscled	0.62 [−0.19, 1.43]
Group mean effect					0.46 [−0.06, 0.99]
Narnia fermorata	Injury	Weapon	4	Muscled	0.78 [−0.67, 2.23]
Narnia fermorata	Injury	Weapon	4	Muscled	0.57 [−0.85, 1.99]
Narnia fermorata	Injury	Weapon	4	Muscled	0.69 [−0.75, 2.13]
Narnia fermorata	Injury	Weapon	10	Muscled	0.07 [−0.81, 0.96]
Narnia fermorata	Injury	Weapon	10	Muscled	0.03 [−0.85, 0.92]
Narnia fermorata	Injury	Weapon	10	Muscled	0.10 [−0.78, 0.99]
Narnia fermorata	Injury	Weapon	10	Muscled	0.48 [−0.42, 1.38]
Narnia fermorata	Injury	Weapon	10	Muscled	0.42 [−0.47, 1.32]
Narnia fermorata	Injury	Weapon	10	Muscled	0.43 [−0.46, 1.33]
Group mean effect					0.46 [−0.06, 0.99]
Scopimera globosa	Pushing	Body	92	Muscleless	0.39 [−0.00, 0.78]
Scopimera globosa	Pushing	Weapon	92	Muscleless	0.85 [0.41, 1.29]
Group mean effect					−x−
Carcinus maenas	Injury	Body	22	Muscled	0.32 [−0.36, 0.99]
Diogenes nitidimanus	Injury	Body	66	Muscled	0.35 [−0.13, 0.83]
Austruca annulipes	Injury	Body	82	Muscled	0.54 [0.20, 0.88]
Aegla longirostri	Injury	Body	15	Muscled	1.21 [0.42, 1.99]
Group mean effect					0.53 [0.28, 0.77]
Carcinus maenas	Injury	Weapon	22	Muscled	0.45 [−0.22, 1.13]
Carcinus maenas	Injury	Weapon	22	Muscled	0.33 [−0.34, 1.00]
Diogenes nitidimanus	Injury	Weapon	66	Muscled	0.46 [−0.02, 0.94]
Cherax dispar	Injury	Weapon	32	Muscled	0.73 [0.21, 1.25]
Cherax dispar	Injury	Weapon	32	Muscled	0.43 [−0.05, 0.91]
Austruca annulipes	Injury	Weapon	82	Muscled	0.50 [0.22, 0.78]
Aegla longirostri	Injury	Weapon	15	Muscled	1.27 [0.47, 2.07]
Aegla longirostri	Injury	Weapon	15	Muscled	1.22 [0.43, 2.01]
Aegla longirostri	Injury	Weapon	15	Muscled	1.28 [0.48, 2.08]
Group mean effect					0.64 [0.34, 0.94]
Dicronocephalus wallichii	Pushing	Body	25	Muscleless	1.57 [0.57, 2.57]
Allomyrina dichotoma	Pushing	Body	101	Muscleless	0.16 [−0.23, 0.55]
Trypoxylus dichotomus	Pushing	Body	47	Muscleless	0.85 [0.23, 1.47]
Group mean effect					0.76 [−0.01, 1.53]
Heterochelus chiragricus	Pushing	Body	164	Muscled	0.78 [0.56, 1.01]
Sagra femorata	Pushing	Body	96	Muscled	0.16 [−0.12, 0.45]
Sagra femorata	Pushing	Body	96	Muscled	0.26 [−0.02, 0.55]
Aegus chelifer	Pushing	Body	94	Muscled	0.58 [−0.34, 1.50]
Aegus chelifer	Pushing	Body	94	Muscled	0.76 [−0.16, 1.68]
Aegus chelifer	Pushing	Body	94	Muscled	0.67 [−0.25, 1.59]
Group mean effect					0.54 [0.15, 0.92]
Dicronocephalus wallichii	Pushing	Weapon	25	Muscleless	2.53 [1.24, 3.82]
Allomyrina dichotoma	Pushing	Weapon	101	Muscleless	0.34 [−0.05, 0.73]
Trypoxylus dichotomus	Pushing	Weapon	47	Muscleless	1.47 [0.76, 2.18]
Trypoxylus dichotomus	Pushing	Weapon	47	Muscleless	0.22 [−0.37, 0.81]
Group mean effect					1.06 [−0.11, 2.23]
Heterochelus chiragricus	Pushing	Weapon	158	Muscled	0.74 [0.02, 1.46]
Heterochelus chiragricus	Pushing	Weapon	158	Muscled	0.83 [0.60, 1.06]
Sagra femorata	Pushing	Weapon	96	Muscled	0.45 [0.16, 0.73]
Sagra femorata	Pushing	Weapon	96	Muscled	0.29 [0.00, 0.57]
Librodor japonicus	Pushing	Weapon	88	Muscled	0.81 [0.37, 1.25]
Gnatocerus cornutus	Pushing	Weapon	34	Muscled	0.56 [0.08, 1.05]
Aegus chelifer	Pushing	Weapon	94	Muscled	0.78 [0.49, 1.08]
Aegus chelifer	Pushing	Weapon	94	Muscled	0.74 [0.44, 1.03]
Aegus chelifer	Pushing	Weapon	94	Muscled	0.73 [−0.19, 1.65]
Librodor japonicus	Pushing	Weapon	30	Muscled	0.80 [0.04, 1.56]
Librodor japonicus	Pushing	Weapon	25	Muscled	0.14 [−0.64, 0.92]
Group mean effect					0.64 [0.46, 0.83]
Ficedula hypoleuca	Injury	Body	10	Muscled	0.39 [−0.51, 1.28]
Ficedula hypoleuca	Injury	Weapon	10	Muscled	0.24 [−0.65, 1.12]
Group mean effect					−x−
Cambridgea foliata	Injury	Body	66	Muscled	0.39 [0.05, 0.73]
Cambridgea foliata	Injury	Weapon	66	Muscled	0.59 [0.24, 0.94]
Group mean effect					−x−
Unionicola formosa	Injury	Body	15	Muscled	0.55 [−0.18, 1.29]
Unionicola formosa	Injury	Body	15	Muscled	0.83 [0.07, 1.58]
Group mean effect					0.68 [0.16, 1.21]
Unionicola formosa	Injury	Weapon	15	Muscled	0.65 [−0.09, 1.39]
Unionicola formosa	Injury	Weapon	15	Muscled	0.73 [−0.02, 1.48]
Group mean effect					0.69 [0.16, 1.21]

Hedges g axis: −2, 0, 2, 4, 6

Fig 1. Summary of Hedges' g (and corresponding confidence intervals) for species, fighting

style, weapon morphology, type of trait, and the number of winners in the study. The grey

polygon behind each Hedges' g is the fitted value of each study on the complete model. The

width of the polygon denotes the credibility interval limits of that study. The darker polygon is

the group mean effect calculated through a meta-analytic model with no moderators. The width

of the polygon denotes the 95% confidence interval of that estimate.

Data extraction and effect sizes

To calculate Hedges' g, we used the mean trait values and their standard deviations for body

size and weapon size measures of winners and losers of each selected study. Hedges' g is a

standardized dimensionless measure and, for this reason, allows comparisons among different

types of measurements and species (Koricheva, Gurevitch, & Mengersen, 2013). Since we

always calculated the Hedges' g as the difference between winner and loser traits, values

greater than zero indicate that winners, on average, have a larger trait than losers, while values

smaller than zero indicate that losers have, on average, a larger trait than winners.

Unfortunately, mean values and standard deviations were not always available on papers. In

those cases, we gathered results from statistical tests comparing winners and losers (e.g., ²

values, F values, Wilcoxon values, degrees of freedom, p-values, and sample sizes) to transform

the statistical results into Hedges' g values using the package "*compute.es*" in R (Del-Re, 2013,

p.). Whenever the mean and standard deviations were displayed on graphs, we used the

webplotdigitizer software to extract the values directly from the figures (Rohatgi, 2019). If none

of those options were available, we contacted the corresponding author to request the data. We

retained the signs of all original statistics before transforming into Hedges' g values.

Since body size and weapon size are allometrically correlated (Voje, 2016), we needed to

ensure that this correlation would not bias our estimates (Weaver, Santos, Tucker, Wilson, &

Hill, 2018). Thus, we gathered the Pearson's correlation coefficient of the body size and weapon

size measures from the papers in which we retrieved the data to calculate the Hedges' g values.

Whenever the study did not provide the correlation coefficient, but the authors sent us the

dataset, we calculated the Pearson correlation coefficient from their data. However, when

neither the study provided the coefficient, nor we had the dataset, we searched the literature for

another study from the same species that had the allometric coefficient. These studies can be

found in Supplementary Table S2.

Data analyses

Our first prediction was that weapons, regardless of morphology and fighting style, would

have higher Hedges' g values than body size. To test this, we used a mixed-effects model in the

R package '*metafor*' (Viechtbauer, 2010). We considered Hedges' g values as the response

variable, the inverse of the variance of Hedges' g as a weight, and the type of trait (i.e., body or

weapon) measured in the original study as our moderator. In this analysis, we included five

different random variables. First, we used study ID because sometimes we extracted more than

one effect size per study (e.g., one effect size for body size and one for weapon size). Given the

allometric correlation between body size and weapon, the effect sizes retrieved for the same

species will have a higher likelihood of being similar than we would expect by chance. To

account for this effect, we built a correlation matrix using the Pearson's correlation coefficient

between body size and the weapon, and used this matrix to account for the effect of allometry

in our analyses. For the three studies that we did not find the correlation coefficients, we used

the mean of all correlation coefficients in our sample (we also substituted the mean values for

0.5 as a correlation coefficient, but the results did not change - data not shown, but present in

github).

For our second random variable, we categorized how each trait was measured. For

instance, some studies use body weight as a proxy for body size, while others use body length;

and whether traits scale according to weight or length might bias our estimates (Houle, Pélabon,

Wagner, & Hansen, 2011). Hence, we used 'type of proxy' as a random variable to account for

any variation that may come from the type of measurement used. Thirdly, we used species ID

to account for effect sizes that came from the same species, but different studies. Fourth, we

used the environment in which the original study was performed (i.e., laboratory or wild).

Finally, we used a matrix containing the phylogenetic relatedness among species. We used the

same random variables in all analyses.

We estimated phylogenetic relatedness based on a phylogeny constructed for the 31

species used in our study (see Results) with the Interactive Tree of Life online tree generator

(iTOL; https://itol.embl.de/, Fig. S1). Since we did not have two species of the same genus in

our analyses, we used the genus as the tips of our tree. After generating the tree, we estimated

branch length using a Brownian motion model of evolution to simulate an ultrametric

phylogenetic tree (Paradis, Claude, & Strimmer, 2004). We then imputed the ultrametric tree as

a variance-covariance matrix of our phylogenetic random variable. We made these analyses

using the packages "*rtol*" (Michonneau, Brown, & Winter, 2016) and "*ape*" in R (Paradis et al.,

2004).

To estimate heterogeneity and biases in our models, we used two approaches. First, we

calculated the ratio of true heterogeneity to the total variation observed across effect estimates

in multivariate studies (I^2, (Borenstein, 2009)). We also partitioned the I^2 into the contribution of

each random variable (Nakagawa & Santos, 2012). Thus, we had estimations of the within-

studies variance (I^2_{study}, similar to most mixed models), the between measurements variance

($I^2_{measurement}$), the between environments variance ($I^2_{environment}$), the phylogenetic variance

($I^2_{phylogeny}$), and the species ID variance ($I^2_{species}$). The sum of these different I^2 are equal to the

total variance observed (I^2_{total}). To estimate the phylogenetic signal in the effect size, we also

calculated the phylogenetic heritability index, H^2, which is similar to the Pagel's λ (Nakagawa &

Santos, 2012). Finally, to test for publication bias, we conducted a modified version of the

Egger's test, in which we used the residuals of our meta-analytic model as the response variable

and the variance of the effect sizes as our predictor variable (Egger, Smith, Schneider, & Minder,

1997). If the intercept of this regression is not different from zero, then there is little evidence for

publication bias (Nakagawa & Santos, 2012).

According to the biomechanical hypothesis, body size will have a higher Hedges' g

value than weapon size for muscleless weapons, while the opposite pattern would occur for

muscled weapons. We also used a mixed-effects model to test this prediction (Viechtbauer,

2010). Again, we considered Hedges' g values as the response variable, the inverse of the

variance of Hedges' g as a weight, but included as moderators: the morphological category (i.e.,

muscled or muscleless), the type of trait (i.e., body or weapon), and their interaction. We used

the same random variables as in the previous analyses: study ID, type of proxy, environment,

phylogeny, and species ID.

Lastly, the functional hypothesis predicts that body size would have a higher Hedges' g

value than weapon size for pushing fights, and the opposite pattern for injury fights. To test the

prediction of this hypothesis, we followed a similar procedure as described before, but our

moderators were: the fighting style (i.e., pushing or injury), the type of trait (i.e., body or

weapon), and their interaction. As in the other analyses, we added study ID, type of proxy,

environment, phylogeny, and species ID as random variables. We performed all analyses in the

R software (R Core Team, 2018).

RESULTS

We gathered 92 effect sizes, of which 55 were related to weapon size and 36 were related to the

body size (Fig. 1). Our search spanned 31 genera, comprising both vertebrate (mainly lizards)

and invertebrate species (mainly arthropods, Fig. S1). Considering the 55 weapon effect sizes, 46

(83.6%) were categorized as muscled weapons and 9 (16.3%) as muscleless weapons, while 29

(52.7%) were included in injury fights and 26 (47.2%) in pushing fights. We had 17 muscled

weapons that were categorized in pushing fights (32.9%) while the remainder 29 were

categorized as injury fights (67.4%). All muscleless weapons were categorized in pushing fights

(Fig. 1, Table S1).

Winners had larger weapons and larger bodies than losers (QM_2 = 48.51, p < 0.0001, Fig.

2), but the Hedges' g estimates for weapon size were similar to body size estimates (QM_1 = 0.5, p

= 0.47, Fig. 2). The dataset showed medium heterogeneity across studies (as measured by I^2,

Table 1), with heterogeneity within studies explaining the largest part of the heterogeneity in

our sample (Table 1). We also found evidence for publication bias (Egger's test; intercept -0.259,

95% CI: -0.39, -0.12, t = -3.87, p <0.0001).

Effect size (Standardized difference between winners and losers)

Fig. 2. Winners had larger bodies and bore larger weapons than losers, but body size and

weapon size have a similar importance in contest resolution. Mean effect size (Hedges' *g*)

expressing trait differences between winners and losers of 31 species. The red dotted line

denotes the threshold in which winners and losers have similar mean trait values. Whiskers

represent the corresponding 95% confidence intervals of the estimated Hedges' *g*.

Biomechanical hypothesis

Although winners had larger bodies and larger weapons than losers, the difference between

winners and losers in body and weapon size estimations were similar for both muscled (QM_1 =

0.28, p = 0.59, Fig. 3) and muscleless (QM_1 = 1.27, p = 0.25) weapons. However, the differences

between winners and losers in weapon size were smaller for muscled muscleless weapons (QM_1

= 5.25, p = 0.021, Fig. 3), while the difference in body sizes between winners and losers did not

differ between species (QM_1 = 1.71, p = 0.19, Fig. 3). The model showed medium heterogeneity

across studies (Table 1). Similar to the results for the overall effect size, the heterogeneity within

studies explained the largest part of the heterogeneity in our sample (Table 1). We also found

evidence for publication bias (Egger's test; intercept -0.225, 95% CI: -0.34, -0.103, t = -3.66, p =

0.0004).

Fig. 3. Winners with muscleless weapons (i.e., no muscles attached to the weapon) have mean

larger weapons than winners with muscled weapons (i.e., muscles attached to the weapon).

Winners are also larger than losers regardless of trait and weapon morphology. Estimated mean

effect (Hedges' g) for trait measures between winners and losers of species that show different

weapon morphologies. For each weapon morphology, results are shown for estimated body

sizes and estimates weapon sizes. The red dotted line denotes the threshold in which winners

and losers have similar trait measures and the whiskers around the estimated mean values

represent the 95% confidence interval.

Functional hypothesis

The difference between winners and losers in weapon size were greater for species involved in

pushing fights than species that exhibit injury fights (QM_1 = 4.71, p = 0.03, Fig. 4). However, the

mean difference in body size between winners and losers did not differ between fighting styles

(QM_1 = 1.65, p = 0.19, Fig. 4). Once again, despite winners being overall larger than losers in all

categories, estimations of weapon size difference were not larger than the body size difference

neither in pushing (QM_1 = 1.02, p = 0.31) nor in injury fights (QM_1 = 0.13, p = 0.71). The model

showed medium heterogeneity across studies (Table 1). Once again, heterogeneity within

studies explained the largest part of the heterogeneity in our sample. We also found evidence

for publication bias (Egger's test; intercept -0.272, 95% CI: -0.39, -0.15, t = -4.46, p <0.001).

Fig. 4. Winners of pushing fights have mean larger weapons than winners of damage fights.

Winners are also larger than losers regardless of trait and fighting style. Estimated mean effect

(Hedges' *g*) for trait differences between winners and losers of species that show different

fighting styles: pushing fights (fights in which weapons are used to push or throw rivals) and

damage fights (fights in which weapons are used to cause injuries on rivals). For each fighting

style, results are shown for estimated body sizes and estimates weapon sizes. The red dotted

line denotes the threshold in which winners and losers have similar traits and the whiskers

around the estimated mean values represent the 95% confidence interval.

DISCUSSION

Here, we evaluated if one aspect of weapon morphology (weapons associated with muscles

responsible for their movement independent of the boy or not) and fighting style (weapons

used to injure rivals or weapons used to push rivals) would impact the relative effect of weapon

and body size on contest resolution. Our results indicated that winners are larger than losers for

both body and weapon size, and that both were similarly important for contest resolution.

Surprisingly, only weapon size differences differed between categories – body size differences

were similar regardless of weapon morphology and fighting style. For weapon size, the

difference between winners and losers is greater in species that bear muscleless weapons and

fight by pushing than in species that bear muscled weapons and fight by injury. We also

covered species belonging to a wide taxonomic range, spanning both vertebrate and

invertebrate groups, and, given the small phylogenetic signal in our sample (Table 1), our

results should be applicable to any animal that bear weapons. Hence, as predicted, weapon

morphology and fighting style are important for predicting the evolution of animal weapons.

However, the differences found were not in the direction we originally expected.

The similar effect of body and weapon sizes on contest resolution indicate that,

independent of weapon morphology and fighting style, both traits are similarly important for

winning. Although the correlation between body and weapon sizes is typically used to explain

this similar effect (Vieira & Peixoto, 2013), we accounted for that correlation in all our models.

Thus, we raise an alternative explanation: weapons might need a certain proportionality to

body size to function properly (Eberhard et al., 2018; Irschick, Meyers, Husak, & Galliard, 2008;

Lailvaux & Irschick, 2006, 2007). Perhaps other traits such as weapon shape and body shape

may have to match weapon characteristics and body size to maximize the winning chances

(McCullough, Tobalske, & Emlen, 2014). For example, males of the stag beetle *Cyclommatus*

metallifer grab rivals using their jaws and lift them up. To perform this movement, the

individual needs a minimum body stability to prevent falling to either side while holding the

rival up in the air. When body and weapon size are mismatched in any direction, body stability

may decrease and even individuals with bigger weapons may have a lower probability of

winning fights (Goyens et al., 2015). Hence, selection may not act solely on the weapon or on the

body, but may also act on how the body functions as a unit - the whole-body performance

(Irschick et al., 2008; Lailvaux & Irschick, 2006, 2007) – and on the covariation between both

traits. Therefore, when several types of weapons are considered simultaneously, whole-body

performance may be the main trait under selection to win fights.

For our surprise, the winners of species with muscleless weapons and pushing fights

presented relatively larger weapons than losers when compared to species with muscled

weapons and injury fights. Two alternative mechanisms might explain this unexpected pattern.

First, given that species with muscled weapons and injury fights face injuries more frequently,

we can expect that each successful hit on the rival increases the amount of injuries accrued and

the likelihood of the rival withdrawing from a fight (Palaoro & Briffa, 2017). If the ability to

cause injuries is sufficiently high, then it is likely that some individuals with smaller weapons

(and body sizes) sometimes might win contests because they were able to hit their rivals first or

more efficiently (Briffa & Lane, 2017; Palaoro & Briffa, 2017). For instance, gladiator frogs

typically use their prepollex spines during contests. These prepollex are a muscleless weapon

that leave deep gashes on rivals (Candaten, Possenti, Mainardi, da Rocha, & Palaoro, 2020) -

some even report frogs blinded or deafened during contests (Dyson, Richert, & Halliday, 2013).

If the smaller rival sometimes wins contests, then the mean difference between winners and

losers will decrease. A similar rationale can be applied to *Anolis* lizards, a group which bear

muscled weapons (their jaws) and enter in injurious contests (Lailvaux & Irschick, 2007). Since

jaw strength is correlated to fighting (and reproductive) success (Husak, Lappin, & Van Den

Bussche, 2009), smaller individuals might win some contests because they bit the larger rival

first or bit in sensible body regions. Therefore, being able to hit first or more efficiently during

the contest may increase the chances to win injury fights. If this is true, it may be that individual

skill or personality traits, such as being highly aggressive or more consistent in striking the rival,

may be more important in species with muscled weapons and injury fights than species with

muscleless weapons and pushing fights (Briffa & Lane, 2017; Camerlink, Turner, Farish, &

Arnott, 2017).

Second, most weapon effect sizes were calculated using weapon size rather than weapon

strength (Fig. 1, Table S1). If weapon strength is more important than weapon size in inflicting

injuries on the rival, then we can expect differences between winners and losers to be more

variable among studies. Ultimately, this would decrease the mean difference between winners

and losers in injury fights and muscled weapons. For example, in species where bluffing occurs

(e.g., crayfish, fiddler crabs), individuals with strong claws frequently engage in physical

contact with rivals to avoid losing to bluffers (Bywater & Wilson, 2012; Wilson, James, Bywater,

& Seebacher, 2009). If individuals pinch one another, winning might be affected more strongly

by the size of the muscles inside the weapon than weapon size *per se*. Any effect of weapon size

on victory might be because of the correlation between weapon size and weapon muscles. On

the other hand, males of flower beetles (*Dicronocephalus wallichii*) needs to get their horned

heads beneath rivals to be able to lift them up and win a fight (Kojima & Lin, 2017). The farther

the individual is from the rival the better - it might win a fight without entering the rival's reach

and without endangering itself from being toppled. In these fights, then, horn length might be a

better predictor of winning than any measure of strength. Therefore, if most studies investigate

only weapon size (as is the case), then we might be missing important information to

understand weapon evolution, when other associated traits (such as the amount of muscles) are

the main responsible for determining individual fighting capacity.

In conclusion, we showed that the effect of weapon size on contest resolution is as

important as the effect of body size. This is surprising, since we expected that in species that

bear weapons, the role of the weapon in distinguishing winners and losers should be more

prevalent than the role of body size. In addition, the similar effect of weapon and body size in

determining the winner occurred independently of weapon morphology and the fighting style.

However, by looking more carefully into weapon morphology and fighting style, we were able

to discover a hidden pattern that indicated greater differences between winners and losers for

weapons that are mainly used to remove rivals from a given resource in comparison to weapons

that are mainly used to cause injuries. This could be caused by the lack of performance

measures of weapons that are primarily selected by their performance. But it also can be that the

selective pressure and the evolutionary pathways associated to the evolution of weapons with

distinct functional roles (such as causing injuries vs expelling rivals) associated to the fighting

behaviours may differ in predictable ways.

Acknowledgments. We thank all the authors who shared the raw data of their paper with us:

Chatchawan Chaisuekul, Christine W. Miller, Devin O'Brien, Kate Umbers, Leilani Walker,

Marcelo M. Dalosto, Murray Fea, Sarah E. Reece, Zach Stahlschmidt, Zachary Emberts. We also

thank Eduardo S.A. Santos for his help with the heterogeneity and bias analyses. A.V.P. thanks

FAPESP for the post-doctoral grant (process no: 2016/22679-3). P.E.C.P. thanks Conselho

Nacional de Desenvolvimento Científico e Tecnológico (CNPq produtividade em pesquisa

311212/2018-2) and Pesquisa e Desenvolvimento of Agência Nacional de Energia Elétrica and

Companhia Energética de Minas Gerais (P&D ANEEL/CEMIG, PROECOS project GT- 599).

REFERENCES

Arnott, G., & Elwood, R. W. (2008). Information gathering and decision making about resource

value in animal contests. *Animal Behaviour*, *76*(3), 529–542. doi:

10.1016/j.anbehav.2008.04.019

Borenstein, M. (2009). Effect sizes for continuous data. In H. Cooper, L. V. Hedges, & J. C.

Valentine (Eds.), *The Handbook of Research Synthesis and Meta-Analysis* (pp. 221–236).

Russell Sage Foundation.

Briffa, M., & Lane, S. M. (2017). The role of skill in animal contests: a neglected component of

fighting ability. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1863),

20171596. doi: 10.1098/rspb.2017.1596

Bywater, C. L., & Wilson, R. S. (2012). Is honesty the best policy? Testing signal reliability in

fiddler crabs when receiver-dependent costs are high. *Functional Ecology*, *26*(4), 804–811.

doi: 10.1111/j.1365-2435.2012.02002.x

Camerlink, I., Turner, S. P., Farish, M., & Arnott, G. (2017). The influence of experience on

contest assessment strategies. *Scientific Reports*, *7*(1), 1–10. doi: 10.1038/s41598-017-

15144-8

Candaten, A., Possenti, A. G., Mainardi, Á. A., da Rocha, M. C., & Palaoro, A. V. (2020). Fighting

scars: heavier gladiator frogs bear more injuries than lighter frogs. *Acta Ethologica*. doi:

10.1007/s10211-019-00333-7

Del-Re, A. C. (2013). compute.es: Compute Effect Size. R Package version 0.2-2. *R Package*.

Retrieved from http://cran.r-project.org/web/packages/compute.es

Dennenmoser, S., & Christy, J. H. (2013). The Design of a Beautiful Weapon: Compensation for

Opposing Sexual Selection on a Trait with Two Functions. *Evolution*, *67*(4), 1181–1188.

doi: 10.1111/evo.12018

Dyson, M., Richert, M., & Halliday, T. (2013). Contest in amphibians. In I. Hardy & M. Briffa

(Eds.), *Animal Contests* (pp. 228–257). Cambridge: Cambridge University Press.

Eberhard, W. G., Rodríguez, R. L., Huber, B. A., Speck, B., Miller, H., Buzatto, B. A., & Machado,

G. (2018). Sexual Selection and Static Allometry: The Importance of Function. *The*

Quarterly Review of Biology, *93*(3), 207–250. doi: 10.1086/699410

Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a

simple, graphical test. *BMJ*, *315*(7109), 629–634. doi: 10.1136/bmj.315.7109.629

Emlen, D. J. (2008). The Evolution of Animal Weapons. *Annual Review of Ecology, Evolution,*

and Systematics, *39*(1), 387–413. doi: 10.1146/annurev.ecolsys.39.110707.173502

Goyens, J., Dirckx, J., & Aerts, P. (2015). Stag Beetle Battle Behavior and its Associated

Anatomical Adaptations. *Journal of Insect Behavior*, *28*(3), 227–244. doi: 10.1007/s10905-

015-9495-3

Green, P. A., & Patek, S. N. (2015). Contests with deadly weapons: telson sparring in mantis

shrimp (Stomatopoda). *Biology Letters, 11*(9), 20150558. doi: 10.1098/rsbl.2015.0558

Green, P. A., & Patek, S. N. (2018). Mutual assessment during ritualized fighting in mantis

shrimp (Stomatopoda). *Proceedings of the Royal Society B: Biological Sciences, 285*(1871),

20172542. doi: 10.1098/rspb.2017.2542

Houle, D., Pélabon, C., Wagner, G. P., & Hansen, T. F. (2011). Measurement and Meaning in

Biology. *The Quarterly Review of Biology, 86*(1), 3–34. doi: 10.1086/658408

Husak, J. F., Lappin, A. K., & Van Den Bussche, R. A. (2009). The fitness advantage of a high-

performance weapon. *Biological Journal of the Linnean Society, 96*(4), 840–845. doi:

10.1111/j.1095-8312.2008.01176.x

Irschick, D. J., Meyers, J. J., Husak, J. F., & Galliard, J. F. L. (2008). How does selection operate on

whole-organism functional performance capacities? A review and synthesis.

Evolutionary Ecology Research, 10(2), 177–196.

Jennings, D. J., & Gammell, M. P. (2013). Contest behaviour in ungulates. In I. C. W. Hardy & M.

Briffa (Eds.), *Animal Contests* (pp. 304–320). Cambridge: Cambridge University Press.

Kojima, W., & Lin, C.-P. (2017). It takes two to tango: functional roles, sexual selection and

allometry of multiple male weapons in the flower beetle Dicronocephalus wallichii

bourgoini. *Biological Journal of the Linnean Society, 121*(3), 514–529. doi:

10.1093/biolinnean/blx018

Koricheva, J., Gurevitch, J., & Mengersen, K. (2013). *Handbook of Meta-analysis in Ecology and*

Evolution. Princeton University Press.

Lailvaux, S. P., & Irschick, D. J. (2006). A functional perspective on sexual selection: insights and

future prospects. *Animal Behaviour, 72*(2), 263–273. doi: 10.1016/j.anbehav.2006.02.003

Lailvaux, S. P., & Irschick, D. J. (2007). The Evolution of Performance-Based Male Fighting

Ability in Caribbean Anolis Lizards. *The American Naturalist, 170*(4), 573–586. doi:

10.1086/521234

McCullough, E. L., Miller, C. W., & Emlen, D. J. (2016). Why Sexually Selected Weapons Are

Not Ornaments. *Trends in Ecology & Evolution, 31*(10), 742–751. doi:

10.1016/j.tree.2016.07.004

McCullough, E. L., & Simmons, L. W. (2016). Selection on male physical performance during

male–male competition and female choice. *Behavioral Ecology, 27*(5), 1288–1295. doi:

10.1093/beheco/arw033

McCullough, E. L., Tobalske, B. W., & Emlen, D. J. (2014). Structural adaptations to diverse

fighting styles in sexually selected weapons. *Proceedings of the National Academy of*

Sciences, 111(40), 14484–14488. doi: 10.1073/pnas.1409585111

Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl: an R package to interact with the

Open Tree of Life data. *Methods in Ecology and Evolution, 7*(12), 1476–1481. doi:

10.1111/2041-210X.12593

Morrell, L. J., Backwell, P. R. Y., & Metcalfe, N. B. (2005). Fighting in fiddler crabs Uca mjoebergi:

what determines duration? *Animal Behaviour, 70*(3), 653–662. doi:

10.1016/j.anbehav.2004.11.014

Nakagawa, S., & Santos, E. S. A. (2012). Methodological issues and advances in biological meta-

analysis. *Evolutionary Ecology, 26*(5), 1253–1274. doi: 10.1007/s10682-012-9555-5

Nolen, Z. J., Allen, P. E., & Miller, C. W. (2017). Seasonal resource value and male size influence

male aggressive interactions in the leaf footed cactus bug, Narnia femorata. *Behavioural*

Processes, 138, 1–6. doi: 10.1016/j.beproc.2017.01.020

O'Brien, D. M., Katsuki, M., & Emlen, D. J. (2017). Selection on an extreme weapon in the frog-

legged leaf beetle (Sagra femorata). *Evolution*, *71*(11), 2584–2598. doi: 10.1111/evo.13336

Okada, K., Suzaki, Y., Okada, Y., & Miyatake, T. (2011). Male Aggressive Behavior and

Exaggerated Hindlegs of the Bean Bug Riptortus pedestris. *Zoological Science*, *28*(9),

659–663. doi: 10.2108/zsj.28.659

Palaoro, A. V., & Briffa, M. (2017). Weaponry and defenses in fighting animals: how allometry

can alter predictions from contest theory. *Behavioral Ecology*, *28*(1), 328–336. doi:

10.1093/beheco/arw163

Palaoro, A. V., Dalosto, M. M., Costa, J. R., & Santos, S. (2014). Freshwater decapod (Aegla

longirostri) uses a mixed assessment strategy to resolve contests. *Animal Behaviour*, *95*,

71–79. doi: 10.1016/j.anbehav.2014.06.014

Palaoro, A. V., Velasque, M., Santos, S., & Briffa, M. (2017). How does environment influence

fighting? The effects of tidal flow on resource value and fighting costs in sea anemones.

Biology Letters, *13*(5), 20170011. doi: 10.1098/rsbl.2017.0011

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in

R language. *Bioinformatics*, *20*(2), 289–290. doi: 10.1093/bioinformatics/btg412

Pinto, N. S., Palaoro, A. V., & Peixoto, P. E. C. (2019). All by myself? Meta-analysis of animal

contests shows stronger support for self than for mutual assessment models. *Biological*

Reviews, *94*(0), 1430–1442. doi: 10.1111/brv.12509

R Core Team. (2018). R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna. *Https://Www.R-Project.Org*.

Rico-Guevara, A., & Hurme, K. J. (2019). Intrasexually selected weapons. *Biological Reviews*,

94(1), 60–101. doi: 10.1111/brv.12436

Rink, A. N., Altwegg, R., Edwards, S., Bowie, R. C. K., & Colville, J. F. (2019). Contest dynamics

and assessment strategies in combatant monkey beetles (Scarabaeidae: Hopliini).

Behavioral Ecology, *30*(3), 713–723. doi: 10.1093/beheco/arz008

Rohatgi, A. (2019). WebPlotDigitizer 4.2: HTML5 based online tool to extract numerical data

from plot images. *Https://Automeris.Io/WebPlotDigitizer*.

Sneddon, L. U., Huntingford, F. A., Taylor, A. C., & Orr, J. F. (2000). Weapon strength and

competitive success in the fights of shore crabs (Carcinus maenas). *Journal of Zoology*,

250(3), 397–403. doi: 10.1111/j.1469-7998.2000.tb00783.x

Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal of*

Statistical Software, *36*(1), 1–48. doi: 10.18637/jss.v036.i03

Vieira, M. C., & Peixoto, P. E. C. (2013). Winners and losers: a meta-analysis of functional

determinants of fighting ability in arthropod contests. *Functional Ecology*, *27*(2), 305–313.

doi: 10.1111/1365-2435.12051

Voje, K. L. (2016). Scaling of Morphological Characters across Trait Type, Sex, and Environment.

The American Naturalist, *187*(1), 89–98. doi: 10.1086/684159

Weaver, R. J., Santos, E. S. A., Tucker, A. M., Wilson, A. E., & Hill, G. E. (2018). Carotenoid

metabolism strengthens the link between feather coloration and individual quality.

Nature Communications, *9*(1), 1–9. doi: 10.1038/s41467-017-02649-z

Wilson, R. S., James, R. S., Bywater, C., & Seebacher, F. (2009). Costs and benefits of increased

weapon size differ between sexes of the slender crayfish, Cherax dispar. *Journal of*

Experimental Biology, *212*(6), 853–858. doi: 10.1242/jeb.024547

Table 1. Heterogeneity estimates obtained from the models.

a) Overall effect of weapon size and body size

	I² (95% CI)	σ² (95% CI)
Study ID	3.86 (1.94, 5.79)	0.0149 (0.00, 0.22)
Measurement	3.75 (1.83, 5.67)	0.014 (0.00, >0.014)
Environment	<0.0001 (0.00, 1.92)	0.000 (0.00, 0.49)
Phylogeny	<0.0001 (0.00, 1.92)	0.000 (0.00, 0.11)
Species ID	21.73 (19.81, 23.65)	0.083 (0.00, 0.22)
Total	29.631 (27.44, 31.28)	-
H^2	<0.00001 (0.00, 1.92)	-

b) Weapon morphology hypothesis

	I² (95% CI)	σ² (95% CI)
Study ID	4.33 (2.41, 6.25)	0.016 (0.00, 0.21)
Measurement	3.71 (1.78, 5.62)	0.014 (0.00, 0.92)
Environment	<0.001 (0.00, 1.92)	0.00 (0.00, 0.55)
Phylogeny	<0.001 (0.00, 1.92)	0.00 (0.00, 0.09)
Species ID	21.44 (19.52, 23.36)	0.082 (0.00, 0.22)
Total	29.48 (27.56, 31.41)	-
H^2	<0.00001 (0.00, 1.92)	-

c) Fighting style hypothesis

	I² (95% CI)	σ² (95% CI)
Study ID	0.51 (0.00, 2.42)	0.002 (0.00, 0.20)
Measurement	5.24 (3.32, 7.16)	0.02 (0.00, 1.61)

Environment	<0.001 (0.00, 1.92)	0.00 (0.00, 0.48)
Phylogeny	<0.001 (0.00, 1.92)	0.00 (0.00, 0.36)
Species ID	23.72 (21.79, 25.64)	0.091 (0.00, 0.21)
Total	29.47 (27.55, 31.39)	-
H^2	<0.00001 (0.00, 1.92)	-